Patterns of Structural Variation Define Prostate Cancer Across Disease States

Meng Zhou[1,2] *, Minjeong Ko[4] *, Anna C. Hoge[4], Kelsey Luu[4], Yuzhen Liu[4], Magdalena L. Russell[4], William W. Hannon[4], Zhenwei Zhang[1,5], Jian Carrot-Zhang[1,2,3], Rameen Beroukhim[1,2], Eliezer M. Van Allen[1,2,6], Atish D. Choudhury[1,3], Peter S. Nelson[4,7], Matthew L. Freedman[1,3,8], Mary-Ellen Taplin[1,3] †, Matthew Meyerson[1,2,3] †, Srinivas R. Viswanathan[1,2,3] †, Gavin Ha[4,7] †

[1] Department of Medical Oncology, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215

[2] Broad Institute of MIT and Harvard, 415 Main Street, Cambridge, MA 02142

[3] Harvard Medical School, 25 Shattuck St, Boston, MA 02115

[4] Public Health Sciences and Human Biology Divisions, Fred Hutchinson Cancer Center, 1100 Fairview Ave. N, Seattle, WA 98109

[5] Department of Pathology, UMass Memorial Medical Center, 1 Innovation Dr. #2, Worcester, MA 01605

[6] Center for Cancer Genomics, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215

[7] Department of Genome Sciences, University of Washington, 1959 Pacific St, Seattle, WA, 98195

[8] Center for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215

* These authors contributed equally to this work.

† These authors jointly supervised this work.

Correspondence:

Gavin Ha
1100 Fairview Ave N
Seattle, WA 98019
1-206-667-2802
gha@fredhutch.org

Srinivas Viswanathan
450 Brookline Ave
1-857-215-0573
Boston MA 02215
srinivas.viswanathan@dfci.harvard.edu

Matthew Meyerson
450 Brookline Ave
Boston MA 02215
Matthew_meyerson@dfci.harvard.edu

Mary-Ellen Taplin
450 Brookline Ave
Boston MA 02215
Mary_Taplin@dfci.harvard.edu

ABSTRACT

The complex genomic landscape of prostate cancer evolves across disease states under therapeutic pressure directed toward inhibiting androgen receptor (*AR*) signaling. While significantly altered genes in prostate cancer have been extensively defined, there have been fewer systematic analyses of how structural variation shapes the genomic landscape of this disease across disease states. We uniformly characterized structural alterations across 278 localized and 143 metastatic prostate cancers profiled by whole genome and transcriptome sequencing. We observed distinct significantly recurrent breakpoints in localized and metastatic castration-resistant prostate cancers (mCRPC), with pervasive alterations in noncoding regions flanking the *AR, MYC*, *FOXA1*, and *LSAMP* genes enriched in mCRPC and *TMPRSS2-ERG* rearrangements enriched in localized prostate cancer. We defined nine subclasses of mCRPC based on signatures of structural variation, each associated with distinct genetic features and clinical outcomes. Our results comprehensively define patterns of structural variation in prostate cancer and identify clinically actionable subgroups based on whole genome profiling.

INTRODUCTION

Over the past decade, genomic sequencing studies have progressively sharpened our view of the genetic landscape of prostate cancer (1). Such studies have defined key driver genes in prostate cancer and have enabled the deployment of therapeutic agents in molecularly-defined disease subsets, including potent androgen receptor (*AR*)-targeted therapies (2, 3), poly (ADP-ribose) polymerase (PARP) inhibitors in *BRCA1/2*-altered prostate cancers, and immune checkpoint inhibitors in cancers with microsatellite instability (4–7).

To date, most cancer genomic studies have employed whole exome sequencing (WES) and have thus been focused on mutations or copy number alterations that occur within the protein-coding regions of genes, which represent only 1-2% of the genome. More recent studies applying whole

genome sequencing (WGS) to prostate and other cancers have identified previously

underappreciated recurrent alterations in regulatory (non-coding) regions of the genome and have

illuminated complex mechanisms of genomic alterations – driven by structural variants (SVs) –

that are difficult to discern by WES; in the case of prostate cancer, most of these studies have

focused on localized disease, the disease state in which tissue is most readily accessible for

profiling (8–22). There remains a need for continued high-resolution genomic discovery efforts in

prostate cancer.

In addition to efforts characterizing entire cancer genomes, recent studies have illustrated the

importance of molecularly profiling prostate cancer across disease states. While many localized

prostate cancers can be cured with surgery or radiotherapy, a substantial portion of higher-risk

cancers recur and progress to metastatic disease, which is incurable. Recurrent prostate cancer

may have a long natural history, during which time a patient may receive several lines of therapy

– with androgen deprivation therapy (ADT) as a backbone – that may shape the cancer's genomic

landscape (23).

Indeed, while hormone-refractory castration-resistant prostate cancer (CRPC) has been less

extensively profiled than localized prostate cancer, several studies have indicated that CRPCs

display genomic landscapes distinct from treatment-naïve disease (24, 25). A cardinal hallmark

of CRPC is the reactivation of *AR* signaling in the face of maximal ADT (22, 26–28). This may

occur via diverse mechanisms, including the production of constitutively active *AR* splice variants

(*AR-V*s) and activating mutations or copy number amplifications of the *AR* gene (29–31) or of

regulatory elements distal to the gene body (13, 15, 32). To date, the relative contribution of each

of these mechanisms in driving *AR* reactivation in CRPC has not been systematically explored.

Also needed is a more global map of significant hotspots of structural variation in prostate cancer

genomes, drawn within a rigorous statistical framework.

In this study, we performed linked-read WGS on 36 mCRPC tumor-normal pairs. We combined

these data with WGS and whole transcriptome sequencing (RNA-Seq) data from previously

described localized and metastatic CRPC cohorts (9, 13, 15, 33). We then established a

harmonized workflow for the integrative genomic analysis of 278 localized and 143 metastatic

CRPC samples, interrogated both hotspots and genome-wide patterns of structural variation, and

evaluated their consequences.

RESULTS

WGS analysis of localized and metastatic prostate cancer cohorts

We performed linked-read whole genome sequencing on 36 biopsy specimens from 33 mCRPC

patients and matched blood normal controls. After quality control, 17 tumor samples were

excluded based on insufficient tumor purity and/or contamination, reflecting the challenge of

obtaining high-purity metastatic biopsies, particularly from bone lesions (34). We included only

samples with tumor purity > 15% in downstream analyses so as to increase confidence in

structural variant calls (**Methods, Figure 1A, Table S1**). We re-analyzed a linked-read WGS

dataset of 23 samples published previously (15), resulting in a total of 42 linked-read WGS

samples from 38 patients with mean coverage of 34X (range 21X - 54X) and 33X (range 25X -

45X) for tumor and normal samples, respectively (**Table S1A**). The mean molecule length was

29 kB and 34 kB in tumor and normal samples, respectively (**Table S1A**).

We further combined these data with 101 mCRPC samples sequenced with standard short-read

sequencing, published previously (13). This resulted in the generation of a final combined cohort

of 143 tumor-normal pairs, which were uniformly analyzed for copy number and structural

alterations via a harmonized pipeline (**Figure 1A**). Fifty-four samples (37.8% of 143 samples)

were collected at castration resistance, prior to receiving treatment of second-generation

androgen receptor signaling inhibitor (ARSi) such as abiraterone and/or enzalutamide ("pre-

treatment"), while the remaining 89 samples (62.2% of 143 samples) were collected at

progression ("post-treatment", **Figure 1B**, **Table S1B**). We analyzed the somatic single nucleotide

variant (SNVs), insertion-deletions (indels), copy number alterations (CNAs), and SVs in the

combined cohort and identified recurrent somatic alterations in each of these classes (**Figure 1A,**

Methods).

A total of 2,315,452 SNVs and indels were called, with a mean tumor mutation burden (TMB) of

2.82 mutations per million bases (Mb). We confirmed that known driver genes of prostate cancer

were enriched for non-synonymous mutations, including *TP53*, *RB1*, *PTEN*, *FOXA1*, *CDK12*, *AR*

and *SPOP* among known COSMIC Cancer Gene Census genes (dndscv, q ≤ 0.1, **Table S1C and**

S1D, Methods). We detected an average of 272 (range 96-833) SV events per sample. Based

on breakpoint orientations, SV events were classified into deletions, inversions, tandem

duplications, inter-chromosomal translocations, and intra-chromosomal translocations, while

intra-chromosomal translocations were further divided into balanced and unbalanced events

based on copy number information (**Methods**). Chromoplexy was detected in 53 samples (37.1%

of 143 samples) while chromothripsis was detected in 37 samples (25.9%); these events were

not mutually exclusive (Fisher's exact test, log-odds=1.417, p-value=0.612). Ten cases (7.0%)

harbored a genome-wide tandem duplicator phenotype (TDP), all of which had *CDK12*

inactivating alterations, as recently reported (15, 35). We found that TDP was mutually exclusive

with ETS rearrangements (Fisher's exact test, log-odds ratio=0.133, p=0.043) and chromothripsis

(log-odds ratio=0.301, p-value=0.007), as previously reported (10, 13, 15, 35).

Analysis of CNA events across the genome revealed amplification and deletion peaks in the

regions of known prostate cancer genes (10, 13, 15, 24). Many oncogenic drivers of mCRPC,

such as *AR* and *MYC*, are within peaks of amplification across the cohort, while tumor

suppressors such as *PTEN, TP53,* and *KMT2C* were found within deletion peaks (**Figure S1C**,

Table S1E and S1F), consistent with prior reports (10, 13, 15, 36).

Recurrent somatic structural variants in prostate cancer-associated genes

Structural variants may either activate or inactivate gene function, depending on the location of the breakpoints and the specific class of SV. We analyzed the potential impact of SVs called across our combined cohort, distinguishing between those with predicted inactivating ("gene transecting events") and activating ("gene flanking events") effects (**Figure 1C, Figure S1C, Table S1G and S1H**). Frequent gene transecting alterations were observed at the *TTC28* (37.1% of 143 samples), *LSAMP* (31.5%), and *PTPRD* (23.8%) loci, which have not been extensively studied in prostate cancer, though they have been reported in callsets for certain cohorts (28). Rearrangements involving *TTC28* were predominantly inter-chromosomal translocations between the gene body and various non-recurrent partner loci (**Figure S2E**). This likely represents retrotransposon activity, given that the *TTC28* locus harbors an active L1 retrotransposon element (37–39). Transecting SVs within the *LSAMP* and *PTPRD* genes were predominantly deletions. Both of these genes are sites of deletion/rearrangement in cancer and have been reported to function as tumor suppressors, though they have not been extensively studied within the context of prostate cancer (40–43) (**Figure 1C**). Of note, although gene transecting events would be predicted to disrupt individual genes, the most frequent transecting events identified via this analysis were deletion events that span the adjacent *TMPRSS2* and *ERG* genes (observed in 37.8%), which actually produces an activating *TMPRSS2-ERG* fusion.

Duplication events that flank an intact gene could activate oncogenes, either by resulting in copy number gain of the gene or by duplicating non-coding regulatory regions (13, 15). In our combined cohort, we observed recurrent tandem duplication events with breakpoints located in the flanking gene regions of several known prostate cancer oncogenes, including *AR* (35.7%), *FOXA1* (16.8%), *MYC* (16.8%), and *CCND1* (14.0%), consistent with frequencies that have been previously reported by us and others (10, 13, 15) (**Figure 1C**).

Certain prostate cancer driver genes were altered by multiple classes of structural alterations in

both the gene body and flanking regions (e.g., *AR, PTEN*), while others were predominantly

altered by a single alteration class (e.g., SNVs for *TP53*, intragenic translocations for *TTC28*, or

flanking tandem duplications for *MYC*) (**Figure 1C, Figure S1C**). Collectively, these results

catalog how diverse classes of rearrangements, both within genes and in intergenic regions, alter

prostate cancer genes across disease states.

Significantly recurrent breakpoint regions in the mCRPC genome are enriched within

enhancer regions and *AR* binding sites

Next, we sought to identify significantly recurrent breakpoint (SRB) regions across our combined

mCRPC cohort of 143 cases in a genome-wide, unbiased manner. We applied a Gamma-Poisson

regression approach to model the occurrences of SV breakpoints within 100 kB windows across

the cohort as previously described (44). Importantly, this model nominates significantly recurrent

breakpoint regions likely to function as cancer drivers by accounting for six different covariates,

including sequence features (e.g., GC-content and transposable elements), fragile sites,

heterochromatin regions, DNase I hypersensitivity sites (DHS), and replication timing (**Methods**),

which may increase specificity over prior studies that have accounted only for SV frequency or

for breakpoint density within a genomic window (10, 13, 15, 45).

We identified a total of 55 significantly recurrent breakpoint regions genome-wide across our

combined mCRPC cohort (Benjamini-Hochberg corrected, q-value ≤ 0.1, **Figure 2A**, **Table S2A**).

Thirty-six (65.5%) SRB regions were located within 1 Mb of 14 known prostate cancer driver

genes, including *AR* and its enhancer, *TMPRSS2/ERG*, *TP53*, *PTEN*, *FOXA1*, and *MYC*. For

these 14 driver genes, we did not observe significant differences in SV alteration frequencies

when comparing between pre-treatment (N=54) and post-progression (N=89) samples, except in

the case of *ERG, for* which the SV frequency was enriched in pre-treatment samples (Fisher's

exact test, p = 0.0395; all other genes had p > 0.05, **Figure S3B**). We also did not identify any

major differences in the alteration frequencies of prostate cancer genes in four patients who had

paired samples collected both before treatment with and after progression on an ARSi. (**Figure**

S3A).

We then sought to compare how SVs drive prostate cancer across disease states. For the

localized disease state, we utilized genome alteration calls from 278 primary localized prostate

cancer tumors from the PCAWG study (9, 33). Using Gamma-Poisson regression, we first

identified 47 SRB regions in localized prostate cancer tumors (**Figure S2B, Table S2B**). Six

prostate cancer genes (*TMPRSS2*, *ERG*, *TP53*, *PTEN*, *IL6ST*, *ELK4*) within mCRPC SRB regions

were also found within or in proximity (less than 1 Mb) to an SRB region in localized disease. By

contrast, four SRBs (three near *SEL1L3* and one near *PRKDC*) were unique to localized disease,

while 27 SRBs were unique to mCRPC with six genes nearby (*LSAMP*, *ETV1*, *MYC*, *PTPRD*,

FOXA1, *AR*). When comparing SV alteration frequencies for the 14 genes located within SRB

regions in either mCRPC or localized tumors, 12 genes were significantly more altered in mCRPC

samples, while *TMPRSS2* and *ERG* were significantly more altered in localized disease (Fisher's

exact test, $p < 0.05$ for all genes, **Figure 2B**). We repeated this comparison using a non-

overlapping cohort of localized prostate cancers profiled by WGS and found similar genes

enriched for SVs in either the localized or metastatic disease states (**Figure S2C-E**) (18). Thus,

localized prostate cancer and mCRPC have significantly different landscapes of recurrent SVs.

To explore the potential functional consequences of SVs in intergenic SRB regions, we

overlapped SV breakpoints with locations of H3K27ac marks specific to mCRPC (46). We

observed that intergenic SVs within SRB regions in the mCRPC cohort included gene flanking

events that were enriched at putative enhancer regions for *AR*, *MYC*, and *FOXA1*, which all had

frequent focal duplication events at sites marked by mCRPC-specific H3K27ac deposition (**Figure**

2C, S2F). Interestingly, an intragenic deletion SRB region was observed near the transcription

start site of *LSAMP*, also overlapping H3K27ac marks. *PTEN* had a high level of both gene

transecting and flanking deletions, leading to SV breakpoints that were spread more broadly

around the gene.

We also observed an enrichment of metastatic-specific *AR* binding sites (ARBS) compared to

localized primary ARBS within the 55 mCRPC SRB regions (**Figure 2D**, one-sided proportion test,

p = 1.05 x 10^{-8}). This enrichment was not observed for localized primary SRB regions (p = 0.22).

These results highlight that SVs within mCRPC SRB regions may be capturing the genome-wide

footprint of activated *AR* signaling that occurs with castration resistance.

Refined landscape of ETS gene fusions from integrated analysis of the genome and

transcriptome

We applied gene fusion analysis by integrating both genome rearrangements and fusion RNA

transcript information from 127 samples with RNA-seq data (**Figure 1A**, **Table S2C**, **Methods**).

For gene fusions involving E26 transformation-specific (ETS) transcription factor gene family

members (*ERG*, *ETV1*, *ETV4* and *ETV5*), we detected 50 events supported by both DNA and

RNA evidence, 15 supported by only DNA evidence, and 10 supported by only RNA evidence

(**Figure 2E**, **Figure S2G**). Overall, 74 samples (51.7% of 143 samples) harbored a fusion event

of the *ETS* gene family, consistent with previous reports (47, 48) (**Figure 1B, Table S2C**).

Among the ETS fusions, *ERG* was most commonly involved with *TMPRSS2* as the fusion partner

(54 out of 57 cases, **Figure 2G**). Other common ETS fusion partners were *SLC45A3* (7 cases)

and lncRNA RP11-356O9.1 downstream of *FOXA1* (3 cases). *ETV1* had eight distinct fusion

partners, which is consistent with previous reports that *ETV1* is a promiscuous ETS fusion

member (49) (**Figure 2F**).

We observed that fusions of the ETS family members *ERG*, *ETV1*, *ETV4* and *ETV5* were mutually

exclusive, except for one sample which harbored fusions of both *ERG* and *ETV1* (**Figure S2G**).

In addition, gene fusion events were correlated with higher expression of the corresponding ETS

genes they involved (Wilcoxon rank-sum tests, p < 0.05 for all genes, **Figure 2E**). In the 38 cases

which did not show any evidence for an ETS fusion, we noted that presence of high-level

expression (z-score > 1) of ETS genes *ERG*, *ETV1*, *ETV4*, and *ETV5* were also mutually

exclusive (Fisher's exact test, p = 0.480 for *ETV4*, p = 0.363 for *ETV5*, **Figure S2G**). These may

represent cases of missed fusion calls, or cases in which ETS family members are

transcriptionally activated through non-genetic mechanisms.

Interestingly, we also observed 20 cases (14.0% of 143 cases) involving fusions between the ETS

family member *ELK4* and its upstream gene *SLC45A3*. While the *ELK4* locus was an SRB in our

analysis (**Figure 2A** and **Figure S2G**), manual inspection of individual samples revealed evidence

for a genomic event capable of producing an *ELK4* fusion in only 1 out of 20 cases (**Figure S2G**

and data not shown). In contrast, 19 other cases showed *ELK4* fusions on RNA-sequencing

alone, consistent with a mechanism of cis-splicing or transcriptional read-through events that may

perhaps be induced by local genomic alterations (50–52) (**Table S2C**). Importantly, although

ELK4 fusions were significantly correlated with higher expression of *ELK4* (Wilcoxon rank-sum

test, p = 7.91×10^{-5}, **Figure S2G**), these events were not mutually exclusive with fusions of other

ETS family members (Fisher's exact test, p = 0.472). Thus, the functional consequences of these

ELK4 fusions and whether they contribute to prostate cancer pathogenesis in a manner similar to

other ETS fusions remains to be determined.

Classes of rearrangements driving *AR* signaling in mCRPC

Genomic alterations involving the *AR* locus play an important role in sustaining *AR* signaling in

mCPRC (13, 15, 26, 53). We sought to catalog the spectrum of diverse structural mechanisms

that underlie *AR* activation in mCRPC, and the relationship between them, in our combined

mCRPC cohort. To understand the relationship between different modes of somatic *AR* activation,

we first determined copy number at the *AR* gene body and its upstream enhancer and categorized

samples into distinct groups of: **(1)** co-amplification (N = 99, 69.2% of 143 cases); **(2)** selective

AR gene body amplification (N = 4, 2.8% of 143 cases); **(3)** selective *AR* enhancer gains (N = 17,

11.9% of 143 cases), and **(4)** lack of amplification for both (N = 23, 16.0% of 143 cases) (**Figure**

3A-C, Table S3). For the 122 samples with expression data available, we observed that *AR* gene

expression was higher in the co-amplification and selective enhancer categories compared to

samples with no amplification, after accounting for tumor purity and ploidy (ANCOVA/TukeyHSD

p-values 5.6×10^{-11} and 4.5×10^{-4}, respectively), but not for selective *AR* status (ANCOVA p = 0.098)

(**Figure 3B, Methods**). Interestingly, we observed that samples with selective enhancer

duplication exhibited similar *AR* expression levels to samples with co-amplification (ANCOVA, p

= 0.31), even though enhancer duplications involved lower-copy gains (mean 2.73, range 1.97 -

5.02) compared to co-amplified samples (mean 12.87, range 1.55 - 150.57) (**Figure 3A**). This is

consistent with previous results (15, 28) and suggests a mechanism whereby *AR* expression

levels are increased through even modest genomic expansion of enhancer elements.

We then systematically and manually curated the diverse mechanisms of rearrangements

activating *AR* signaling by analyzing patterns of SVs at the *AR* locus (**Figure 3C, Table S3,**

Methods). We observed a total of 62 samples (43.4% of 143 samples) with tandem duplication

SV events that spanned the enhancer with breakpoints located within 1 Mb, including 16 cases

(11.2% of 143 samples) with selective enhancer copy number amplification status (**Figure 3D**).

Thirty-two samples (22.4% of 143 samples) harbored intragenic rearrangements within *AR*, which

may have implications for the production of truncated, constitutively-active *AR* splice variants (31).

For example, in case 01115414-TA1, we observed a tandem duplication breakpoint selectively

amplifying exons 1-4 of AR, but not exons 5-8 of *AR*, which includes the ligand binding domain;

such an event could promote selective expression of a constitutively active truncated *AR* variant,

although RNA-Seq data was not available on this sample. (**Figure 3E**). In another case, DTB-

124-BL, our reanalysis confirmed that a focal deletion involving exons 1-4 resulted in the

expression of truncated *AR* variants as previously described (28) (**Figure S3C**). Interestingly, in

the 21 samples with selective *AR* enhancer or selective *AR* gene body copy number gain, none

harbored intragenic SV events in *AR*.

We also examined the landscape of complex rearrangement mechanisms involving *AR*; these

mechanisms involve multiple SV events and copy number patterns, including chromothripsis,

extrachromosomal DNA (ecDNA), chromoplexy, and breakage-fusion-bridge cycle (BFB)

(**Methods**). Chromothripsis of a region or the entire X chromosome involving the *AR* locus was

detected in 5 samples, all of which had co-amplification of *AR* and enhancer, suggesting that

following repair after catastrophic DNA shattering the *AR* locus was retained or further amplified

(**Figure 3F, Figure 3G**). Thirteen samples (9.1% of 143 samples) showed very high levels of *AR*

and enhancer copy number, suggesting the possibility of their presence on extrachromosomal

elements (ecDNA, **Figure 3H**). In 40 samples (28.0% of 143 samples), the most frequent complex

rearrangement mechanism, BFB, led to *AR* locus amplification, including instances following

chromothripsis (14, 54) (**Figure 3G**). Overall, we noted that complex rearrangement events, which

frequently co-occurred, were significantly enriched in samples with co-amplification of *AR* and

enhancer compared to those with selective enhancer copy number gain status (Fisher's exact

test, p = 1.52×10^{-4}).

Distinct signatures of structural rearrangement patterns in mCRPC

To systematically characterize genome-wide structural rearrangement patterns in mCRPC, we

performed rearrangement signature analysis using SV breakpoint features, non-negative matrix

factorization, and known reference signatures (12, 55) (**Methods**). First, we derived signatures

de novo, which identified eight signatures: six that matched reference signatures (RefSigs) also

observed in localized prostate cancer (> 0.91 cosine similarity), one that matched an ovarian

cancer RefSig.R14 associated with large segment (100 kB-10 Mb) TDP (0.96 cosine similarity),

and one that was likely an artifact specific to linked-read sequencing (**Figure S4A-C, Table S4A**

and 4B). Therefore, we excluded the linked-read data and focused on standard WGS data from

101 mCRPC cases for further SV signature analysis. We fit these samples to the nine known

RefSigs from localized prostate cancer (R1-4, R6a-b, R8-9, R15) and the one (R14) from ovarian

cancer (**Figure S4A**, **Table S4C**). Overall, eight of the RefSigs were detected across our cohort

(R1-2, R4, R6a-b, R9, R14-15). Notably absent in mCRPC were RefSig.R8 (short, 1-10 kB

inversions) and RefSig.R3, which is associated with germline *BRCA1* mutations and short (1-100

kB) tandem duplications (11, 12, 55, 56) (**Figure S4D**). By contrast, we observed increased

prevalence of some signatures in mCRPC compared to localized disease, including RefSig.R2

(large SV classes, abundant translocations; 97% vs. 60%), RefSig.R4 (clustered translocation

events; 37% vs. 27%), and RefSig.R15 (large deletions and inversions, 48% vs. 37%) (**Figure**

S4D).

To investigate whether molecular subtypes in mCRPC can be grouped based on SV patterns, we

applied hierarchical clustering on the exposure of the eight fitted signatures and identified nine

distinct SV clusters (**Figure 4, Table S4C**). We observed that samples in SV Cluster 1 were

composed of non-clustered translocation events and were significantly enriched for the presence

of chromoplexy (χ^2 test, FDR corrected, q = 0.12). SV Cluster 3 was characterized by many short

deletions and was significantly enriched for *BRCA2* mutations (q = 5.01x10^{-4}). SV Cluster 5 was

significantly enriched for *SPOP* mutations (q = 0.02), with no instances of ETS gene family fusion

(q=0.06), consistent with previous reports (57). SV Cluster 6 had the highest prevalence of *TP53*

mutation (q = 0.02), while SV Cluster 7 samples harbored the TDP associated with *CDK12*

inactivation (q = 3.52 x 10^{-11}) as well as enrichment for *CCND1* gains (q = 0.02), consistent with

previous reports (35, 58). The remaining clusters did not have enrichment for any alterations in

known driver genes; however, distinct SV patterns were still evident in SV Cluster 4 (non-clustered

tandem duplications), 8, and 9 (increased clustered SV events of various classes).

While SV Clusters 3, 5 and 6 had significant enrichment of mutations in *BRCA2*, *SPOP,* and *TP53*,

respectively, not all samples within each cluster harbored these mutations. Intriguingly, we further

noted that clinical outcomes showed significantly better stratification when using SV Clusters 3,

5, and 6 for outcome stratification compared to using the associated mutation status itself (**Figure**

S4D-E). Specifically, SV Cluster 5 had significantly better overall survival than SV Clusters 3 and

6 (log-rank test, p=0.01), while the sample group with *SPOP* mutations did not have significantly

greater survival compared to the sample groups with *BRCA2* and *TP53* mutations (log-rank test,

p=0.45) in this cohort. Together, these results indicate the analysis of genome-wide patterns of

rearrangements may provide a way to further refine molecular subtypes in mCRPC.

DISCUSSION

We present a large-scale and comprehensive integrative genomic analysis of both localized

prostate cancer and mCRPC, with a focus on how structural variation drives each of these

clinically distinct disease states. The size of our cohort as well as our harmonized analysis pipeline

enable a sharper view of the genetic alterations that drive prostate cancer across its natural history

as compared with prior studies, which have involved either smaller cohorts or been limited to a

single disease state (9, 13, 15, 59).

In contrast to somatic SNVs/indels and CNAs that occur within coding regions, the functional and

clinical significance of alterations within noncoding regions has often been more challenging to

interpret, as localized variations in mutability may result in the nomination of certain recurrently

mutated sites that do not necessarily drive cancer (11, 12, 44). This issue is even more complex

for SVs, in which different classes of SVs spanning the same loci would be predicted to have

distinct functional consequences. Our study addresses the former issue by identifying genomic

hotspots of structural variation with rigorous correction for covariates including nucleotide

composition, replication timing, sensitivity to DNA breaks, repetitive elements, and chromatin

state. We address the latter issue by careful curation of SV classes to distinguish those that are

likely to be activating versus inactivating (**Figures 1B and 3; Methods**).

Our approach has produced several insights into the recurrent rearrangements that drive prostate

cancer. First, several top hotpots of rearrangement genome-wide lie in noncoding regions outside

the boundaries of known prostate cancer genes, as previously reported (10, 13, 15). In many

cases, such as for *AR*, *MYC*, and *FOXA1*, these hotspots overlap with active chromatin marks

and likely represent distal regulatory regions for neighboring prostate cancer genes, as shown by

our analyses overlapping SVs with ChIP-Seq on mCRPC specimens (46) (**Figure 2**). These data

are intriguing in light of the observation that a majority of prostate cancer germline susceptibility

loci are in noncoding regions (60). Second, the loci altered by rearrangements differ across

prostate cancer disease states (**Figure 2B**). For example, *TMPRSS2-ERG* rearrangements are

enriched in localized prostate cancer versus mCRPC, while alterations in *AR, FOXA1, MYC*, and

LSAMP are more frequent in mCRPC than in localized disease. Third, certain driver genes are

enriched for alteration by SVs as compared to other mutagenic processes. For example, *PTEN*

inactivation frequently occurs via gene transecting SV events, while *TP53* inactivation is primarily

caused by SNVs (**Figure 1C and Figure S1**).

Our systematic genomic discovery efforts confirm the primacy of *AR* as a target of somatic

alteration in hormone-refractory mCRPC (13, 15, 26, 53, 61). We have precisely catalogued the

diverse genomic mechanisms leading to *AR* activation across our large aggregate cohort and find

that different alteration mechanisms are associated with differing levels of *AR* amplification.

Whether the precise mechanism by which *AR* is altered in a given patient is associated with

differences in response to *AR* pathway inhibition warrants further investigation in clinically

annotated cohorts. High levels of *AR* signaling in mCRPC may also underlie the patterns of

structural variation seen in this disease state. Strikingly, we found that *AR* binding sites

overlapped several of the top SV hotspots in mCRPC (**Figure 2D**), consistent with the notion that

androgen signaling may induce DNA double-strand breaks that resolve as rearrangements (62).

In addition to alterations in highly validated prostate cancer genes, we identified highly recurrent

rearrangements near or involving genes that have not been extensively studied in prostate cancer

in multiple cohorts, such as *LSAMP, PTPRD*, and *TTC28*. *LSAMP* encodes a cell-surface

glycoprotein and has a possible tumor suppressor role in several cancers (40–42); notably,

deletions near the *LSAMP* locus have been shown in one report to be enriched in African

American men with prostate cancer (63). *PTPRD*, a receptor protein tyrosine kinase, has been

previously identified as a target of inactivating alteration in glioblastoma (43). We observed

frequent SVs near the *TTC28* locus, which encodes an L1 retrotransposon element, specifically

in mCRPC (**Figure 1C**). L1 retrotranspositions originating from *TTC28* have been reported

previously in colorectal cancer (37–39); our results raise the intriguing possibility that they may

also be frequent in prostate cancer, and may be activated by the pressure of hormonal therapy.

Interestingly, we also observed SRBs near *ELK4* along with a relatively high frequency of

SLC45A3-ELK4 chimeric transcripts, although it was not clear how the rearrangements at this

locus produced the chimeric transcripts in most cases. Whether this fusion functions similarly to

or in a distinct mode from other ETS fusions is an exciting area for future study.

Our study also extends beyond the analysis of SVs at individual loci to molecularly subclassify

prostate cancers based on their genome-wide signatures of structural variation. Sample clustering

based on SV signature exposure defines distinct molecular subtypes of prostate cancer and may

find utility alongside signatures of single base substitution and copy number to more precisely

define tumor subtypes (55, 64–67). In the mCRPC cohort, we identified 9 molecular subtypes

based on SV signature, and several clusters had clear associated genomic alterations including

chromoplexy (cluster 1), *BRCA2* alterations (cluster 3), *SPOP* alterations (cluster 5), *TP53*

alterations (cluster 6) and *CDK12*/*CCND1* alterations (cluster 7). Future studies with larger WGS

and RNA-Seq cohorts may identify associated alterations or transcriptional signatures in the

remaining clusters. Notably, unsupervised clustering identified samples with clear SV signatures

but without detectable associated mutations in genes or pathways that plausibly contribute to the

genomic alterations (**Figure 4**). Moreover, clinical outcomes were more separated by SV

signature cluster than by alterations of the mutations associated with those clusters (**Figure S4D-**

E).

In sum, these results highlight the dynamic complexity of rearrangements in prostate cancer

across disease states and provide insights into new mechanisms of oncogenesis that can be

functionally prioritized in future studies. More broadly, our work underscores the key role of large-

scale WGS studies in the derivation of a comprehensive molecular taxonomy of prostate cancer.

METHODS

Data and code availability

- Whole genome sequencing data have been deposited at dbGaP under accession number phs001577 and access is available upon request.
- All original code has been deposited at GitHub and is publicly available as of the date of publication. Links are provided in the key resources table.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Sequence data processing for linked-read genome sequencing data

Data processing of the linked-read genome sequencing data include high molecular weight DNA preparation and sequencing library construction followed protocols as previously described (15). DNA was extracted from tumor samples using the MagAttract HMW DNA Kit (QIAGEN), and then quantified using Quant-it Picogreen assay kit (Thermo Fisher) on a Varioskan Flash Microplate Reader (Thermo Fisher). For germline samples, pre-extracted DNA was size-selected on the PippinHT platform (Sage Science) and then quantified using the Quant-it Picogreen assay kit (Thermo Fisher) on a Varioskan Flash Microplate Reader (Thermo Fisher). Libraries were constructed using the 10X Chromium protocol (10X Genomics), with the fragment sizes determined using the DNA 1000 Kit and 2100 BioAnalyzer (Agilent Technologies) and quantified using qPCR (KAPA Library Quantification Kit, Kapa Biosystems). WGS libraries were sequenced using the Illumina HiSeqX platform. The Long Ranger v2.2.2 pipeline (10X Genomics) was used for aligning sequence reads to the human genome hg38 (GRCh38).

Samples were excluded from the analysis based on having tumor purity less than 15% estimated by TitanCNA or based on cross-individual contamination indicated by SNP fingerprinting. A total of 17 samples with linked-read data was excluded (**Table S1J**).

List of known prostate cancer driver genes

For analyses limited to established prostate cancer driver genes, a curated list of 159 known prostate cancer driver genes was assembled from several prior studies (10, 13, 15, 24). The list of genes are provided in **Table S1**.

Somatic mutation analysis

Somatic mutation detection

Somatic mutation calls for samples based on linked-read sequencing were generated by Mutect2 from the Genome Analysis Toolkit (GATK) (68). Default parameters were used on individual pairs of tumor and normal samples following the standard GATK pipeline. A panel of normals based on all normal samples was used to filter out germline variants. The SNV calls were further processed using the modified version of LoLoPicker (69) as described previously (15). The panel of normals for LoLoPicker was generated from 52 normal samples based on linked-read sequencing. The final SNV call set was composed of the common variants called by both Mutect2 and LoLoPicker. Somatic indels for linked-read samples were called by Strelka (70). All parameters were default except the following modifications: sindelNoise = 0.000001, minTier1Mapq = 20. Somatic mutation calls for the 101 WGS samples based on short-read sequencing including SNV and indels based on Strelka were obtained from a prior study (13). All variants were further annotated using annovar with "table_annovar.pl" to functionally annotate genetic variants. The parameter -neargene was set to 5000 to define the promoter region as 5 kB upstream of the transcription start site of a protein coding gene.

Analysis of significantly mutated genes

R package dndscv (71) was used to identify significantly mutated genes. For driver discovery on GRCh38, a precomputed database corresponding to human genome GRCh38.p12 was downloaded and used as the reference database. A global q-value ≤ 0.1 was applied to identify

statistically significant (novel) driver genes. To reduce false positives and increase the signal to

noise ratio, we only considered mutations in Cancer Gene Census genes (v81) (72).

Copy-number analysis of linked-read WGS and short-read WGS data

Copy-number calls

The ploidy and purity corrected copy-number of all mCRPC samples in this study was analyzed

by TitanCNA (73) and ichorCNA (74), with different pipeline settings. For WCDT samples, the

snakemake workflow for Illumina sequencing was applied with the following parameters modified:

ichorCNA_normal: c(0.25, 0.5, 0.75); ichorCNA_ploidy: c(2,3,4); ichorCNA_includeHOMD: TRUE;

ichorCNA_minMapScore: 0.75;   ichorCNA_maxFracGenomeSubclone: 0.5;

ichorCNA_maxFracCNASubclone: 0.7;   TitanCNA_maxNumClonalClusters: 3;

TitanCNA_maxPloidy: 4.   The   workflow   is   available   at

https://github.com/GavinHaLab/TitanCNA_SV_WGS.

For linked-read data samples, a Snakemake workflow for 10X Genomics whole genome

sequencing   data   was   used   with   the   following   parameters   modified:

TitanCNA_maxNumClonalClusters: 3; TitanCNA_maxPloidy: 4. TitanCNA solutions were

generated for number of clonal clusters from 1 to 3 and ploidy initializations from 2 to 4. Optimal

solutions were selected as described, with manual inspection to confirm tumor ploidy and clonal

cluster selection (15); solutions are provided in **Table S1J**. The workflow can be accessed at

https://github.com/GavinHaLab/TitanCNA_10X_snakemake. The final copy-number call-set is

included in **Table S1I**.

Recurrent somatic copy-number alteration

GISTIC 2.0 was used to detect regions with recurrent CNA in mCRPC samples. For input, all copy

numbers (logR_Copy_Number from TITAN output) were converted to log2 copy ratio using the

median logR copy number from genome-wide (separately for autosomes and X chromosome) as

denominator. We set corrected logR copy number to -1.5 for segments where corrected log R

copy number below -1.5 and set values to 0 if copy neutral. GISTIC2.0 was run with the following

parameters: td 0.5; ta 0.1; genegistic 0; maxseg 5000; js 4; cap 1.5; broad 1; brlen 0.75; conf 0.99;

qvt 0.25; armpeel 1; rx 0; gcm mean; do_gene_gistic 1; savegene 1; scent median. Wide peaks

detected by GISTIC2 were re-annotated based on overlapping genomic coordinates, using

prostate cancer driver genes.

Structural variant analysis

Structural variant detection in linked-read and short-read whole genome sequencing data

For each tumor-normal pair of samples with linked-read genome sequencing data, three variant

callers were used to detect structural variants: SvABA (75), GROC-SVS (76), Long Ranger

version 2.2.2 (https://support.10xgenomics.com/genome-

exome/software/pipelines/latest/using/wgs).

The SvABA analysis was performed using default tumor-normal paired settings. Re-analysis of

low confidence (based on evidence from discordant and split reads) events filtered by SvABA was

performed to 'rescue' SVs using linked-read barcode overlap between pairs of breakpoints within

a given SV event, as previously described (15). Only SV events having span of 1.5 times the

mean molecule length in the library were considered for rescue. We further rescued low

confidence intra-chromosomal SV events with span > 50 kB filtered by SvABA if at least one of

the breakpoint pair was within 100 kB of a CNA boundary or (2) if both breakpoints were each

within 1 Mb of the boundaries for the overlapping CNA event and the length of the SV overlaps

this CNA event by > 75%. Inter-chromosomal translocation SV events filtered by SvABA are

rescued if both breakpoints were within 100 kB of CNA boundaries.

GROC-SVS analysis was performed using two-sample (tumor-normal paired) mode or three-

sample (pre-treatment, post-progression, normal) mode when applicable. SV events were

retained if all following conditions were satisfied: (1) $p < 1\times10^{-10}$, (2) minimum barcode overlap ≥

2 on the same haplotype, (3) no more than 1 barcode overlap between different haplotypes, (4)

FILTER value reported by the software was within this set {"PASS", "NOLONGFRAGS",

"NEARBYSNVS", or "NEARBYSNVS; NOLONGFRAGS"}, and (5) classified as somatic.

Long Ranger analysis generated SV calls for tumor and normal samples, independently. For each

tumor-normal pair, both large SVs ("large_sv_calls.bedpe") and deletions ("dels.vcf") were

combined for individual samples. Somatic tumor SVs were determined as events that were not

found in the matched normal sample based on the left breakpoints in tumor and normal being

within 1 kB and the right breakpoints in tumor and normal samples being within 1 kB. Only SV

events with FILTER values within this set {"PASS", "LOCAL_ASM", "SV", "CNV, SV"} and intra-

chromosomal events with span ≥ 100 kB were considered. SV events were only retained if both

breakpoints of an SV event were within 500 kB the boundaries of an overlapping CNA event and

the length of SV overlaps this CNA event by > 75%.

SV events from these three callers were then combined by taking the union of the filtered events

from. Intersecting events between 2 or more call-sets were determined if both breakpoints of one

event were located within 5 kB from both breakpoints of the event detected by the other tool. Then

the details of this event were retained based the priority ordered by SvABA, GROC-SVS, Long

Ranger. Long Ranger SV events were further filtered out if they were not intersecting events

detected by at least one other tool. SV events with span less than 1 kB were excluded from

downstream analyses.

An SV panel of normals (PoN) was generated using germline events from SvABA and Long

Ranger calls. There are two components to this panel: (1) frequency of germline events at exact

breakpoint locations (SVpon.bkpt) and (2) frequency of germline event breakpoint overlapping

within tiled windows of 1 kB (SVpon.blackListBins). The PoN was used to filter events in the

combined SV call-set when an SV has at least one breakpoint with SVpon.bkpt ≥ 2 and

overlapping bin with SVpon.blackListBins ≥ 100.

The workflow for SV analysis from linked-read sequencing data can be accessed at

https://github.com/GavinHaLab/SV_10X_analysis. Manual curation of filtered SV events in the *AR*

locus was performed and rescued events were labeled "Manual". The final SV call-set is included

in **Table S1K**.

For samples based on short-read WGS, SvABA was used in tumor-normal paired mode for SV

detection with default parameters. Intra-chromosomal SV events with span > 1 kB were retained.

The SvABA workflow can be accessed at https://github.com/GavinHaLab/TitanCNA_SV_WGS

Classification of structural variants in mCRPC

SV types were annotated based on orientations of breakpoints and bin-level copy-number around

breakpoints. The orientation of one breakpoint was defined based on the fragment of DNA

molecule being connected to the altered molecule. If the connected fragment was to the 5'-end of

the breakpoint, *i.e.*, "upstream" or "left" to the breakpoint, then the orientation was annotated as

forward or "+"; on the contrary, if the connected fragment was located to the 3'-end of the

breakpoint, the orientation was annotated as reverse or "-". The copy-number near each

breakpoint was evaluated using 10 kB bins. For one SV event, copy-number values of the bins

located to the upstream and downstream of breakpoint 1 were denoted as c_1^{up} and c_1^{down},

respectively; similarly, the copy-number values for breakpoint 2 were denoted as c_2^{up} and c_2^{down}.

In addition, then mean copy-number c^{mean} of the 10 kB bins between the two breakpoints of one

SV event and the number of bins s were also considered during SV classification. Intra-

chromosomal SV events, *i.e.*, both breakpoints were located on the same chromosome, were

classified to the list of SV types below following the corresponding classification criteria.

•   Deletion. Events having the orientation combination (reverse, forward) and length

between 10 kB and 1 Mb were classified as deletions. The copy-number values of

breakpoints should satisfy $c_1^{up} > c_1^{down}$ or $c_2^{up} < c_2^{down}$, and $c_1^{up} > c^{mean}$ or $c_2^{down} > c^{mean}$, and

$s \leq 5$. In addition, events overlapping copy-number deletion or LOH segments were also

considered as deletions.

•   Tandem duplication. Events having the orientation combination (forward, reverse) and

length between 10 kB and 1 Mb were classified as tandem duplications. The copy-number

values of breakpoints should satisfy $c_1^{up} < c_1^{down}$ or $c_2^{up} > c_2^{down}$, and $c_1^{up} < c^{mean}$ or $c_2^{down} <$

c^{mean}, and $s \leq 5$. In addition, events overlapping copy-number gain or copy neutral LOH

segments were also considered as tandem duplications.

•   Inversion. Events having the orientation combination (forward, forward) or (reverse,

reverse) and length between 10 kB and 5 Mb were classified as inversions. Furthermore,

inversion events shorter than 30 kB with unequal copy-numbers around either breakpoint

were classified as fold-back inversions.

•   Balanced rearrangement (balanced). Events having the orientation combination same to

inversion (forward, forward) or (reverse, reverse), but length larger than 5 Mb were

classified as balanced events. The copy-number values of breakpoints should satisfy c_1^{up}

$= c_1^{down}$ and $c_2^{up} = c_2^{down}$, or $c_1^{up} = c^{mean}$ and $c_2^{down} = c^{mean}$.

•   Unbalanced rearrangement (unbalanced). Intra-chromosomal events which did not fulfill

any of the above criteria and having length larger than 10 kB were classified as

unbalanced events.

•   All SV events with two breakpoints located on different chromosomes were classified as

translocations.

 ICGC/TCGA PCAWG localized prostate cancer structural variants

We obtained localized prostate cancer structural variant calls from ICGC Data Portal release 28

(https://dcc.icgc.org/releases/PCAWG/consensus_sv). In this consensus SV file, each SV event

was predicted by at least two variant callers. Samples that were classified as prostate

adenocarcinoma (PRAD) and early onset prostate cancer (EOPC) were selected. A total of 278

samples successfully lifted over to genome build GRCh38. To maximize consistency with mCRPC

datasets, we used only the PCAWG consensus SVs that included "SNOWMAN" as one of the

tools. Note that "SNOWMAN" was the previous name for SvABA. Intrachromosomal SV events

shorter than 10 kB were excluded.

We obtained ICGC early-onset PC (EOPC) and late-onset (LOPC) structural variant calls from

Gerhauser et al, 2018 (18) (https://data.mendeley.com/datasets/6gtrrxrn2c/1). This dataset

includes a total of 253 samples, consisting of 206 EOPC and 47 LOPC. We used the SV calls

generated in the original study using DELLY2 (Rausch et al., 2012) PCAWG analysis workflow

(https://github.com/ICGC-TCGA-PanCancer/pcawg_delly_workflow). SV coordinates were lifted

over to GRCh38.

Tandem duplicator phenotype

For all samples in the combined cohort, the TDP status was predicted using copy-number and

SV by counting the number of copy-number segments overlapping with tandem duplication SV

events, *i.e.,* gain segments. A sample was considered as TDP if it has more than 300, or 90 gain

segments for samples based on linked-read sequencing and short-read sequencing, respectively.

The number of segments with gain and median length SV are reported in **Table S1L**.

Chromothripsis analysis

Chromothripsis events were detected by ShatterSeek R package (77). Structural variants calls by

SvABA and copy-number calls by TitanCNA were used as input data (excluding Y chromosome).

In the input, consecutive segments were joined as one if they had the same copy-number value

and centromere regions were filtered out.

Manual inspection was performed for reported chromothripsis-like events after adapting criteria

thresholds. For samples based on short-read sequencing, confidence classification criteria were

refined from the ShatterSeek documentation. Following criteria were used for high confidence

calls: total number of intra-chromosomal structural variants events involved in the event ≥ 10; max

number of oscillating CN segments (two states) ≥ 10; satisfying either the chromosomal

enrichment or the exponential distribution of breakpoints test ($p \leq 0.05$). For samples based on

linked-read sequencing, we filtered these calls based on a weighted score that is primarily

determined by the number of SVs in a cluster, with less weight given to CN oscillations. In this

analysis, events with a score over 0.8 were considered as high confidence and all other events

were excluded. The score is defined based on the following terms (ranges from 0 to 1).

• Weight 0.6 if total number of intra-chromosomal structural variants events involved in the
event ≥ 10.

• Weight 0.2 for max number of oscillating CN segments (two states) ≥ 7 or max number of
oscillating CN segments (three states) ≥ 14.

• Weight 0.1 for passing chromosomal enrichment test by ShatterSeek.

• Weight 0.1 for passing exponential distribution of breakpoints test.

Chromoplexy analysis

ChainFinder was used to detect chromoplexy events (8). Ten samples that were considered as

TDP (01115374-TA2, 01115202-TC2, 01115248-TA3, 01115503-TC2, 01115257-TA4,

01115284-TA9, 01115414-TA1, DTB-063-BL, DTB-183-BL, DTB-214-BL) were excluded from

this analysis. In addition, four samples that were found to cause numeric instabilities of

ChainFinder were also excluded (DTB-023-BL, DTB-102-PRO, DTB-111-PRO, DTB-151-BL).

The SV calls of remaining samples were further filtered to exclude those that were located within

5 Mb from chromosomal ends or overlapping chromothripsis regions. For copy-number input,

segments that were determined as copy neutral by TitanCNA were set to have log copy-ratio of

0. Copy-ratio of the other segments were computed from copy-number values generated by

TitanCNA divided by 2 for autosomes or 1 for X chromosome. Log copy-ratio values less than -

1.5 were set to -1.5. The output of ChainFinder was used for determining chromoplexy status of

individual samples. A chromoplexy event was defined as a chain including at least 5

rearrangement events and involving more than 2 different chromosomes. Samples having at least

2 such events were considered positive for chromoplexy status.

ChIP-seq data analysis

ChIP-seq data used in this study were downloaded from Gene Expression Omnibus (GEO) (78,

79) and the Sequence Read Archive (SRA) (1). Short reads were mapped to the human genome

GRCh38 (hg38) using bwa (80). Because read lengths were less than 50bp, the bwa aln

command with default parameters was used for mapping. MACS2 (81) was used to identify peaks

from mapped ChIP-seq data. For histone modification marks, MACS2 callpeak command was

applied with --nomodel --broad –extsize 146. For CTCF data, MACS2 callpeak command was

used with --nomodel --extsize 200. Below is the list of ChIP-seq datasets involved in this analysis.

•   H3K4me3, H3K27me3 and CTCF (GSE38685) (82).

•   H3K36me3 and H3K9me3 (GSE98732) (83).

•   H3K4me1 and H3K27ac (GSE73785) (84).

For *AR* binding site (ARBS), the peak files were downloaded from two different datasets and

converted to hg38 coordinates. For primary prostate cancer, ARBS data were downloaded from

GSE70079 (Pomerantz et al., 2015). The union of all tumor sample peaks was used. For mCRPC,

met-specfic ARBS data were obtained from a previous study (46).

Identification of SRB regions

Masking the human genome based on mappability

The human genome was divided into 100 kB non-overlapping bins for detection of significantly recurrent breakpoint regions (SRB). A low-mappability mask was generated for the hg38 genome to screen out out regions that are difficult for variant calling based on short-read sequencing. We adopted procedures from a previous study (86) to construct a mask corresponding to regions with low mappability in the human genome. Below is a list of masked regions included in the low-mappability mask.

- Composition mask. This set of masked regions includes regions with low sequence complexity detected by mdust, regions with long homopolymers detected by seqtk, satellite regions annotated by RepeatMasker (87), and low complexity regions annotated by RepeatMasker.

- Mappability mask. This mask was based on mappability of k-mers in the human genome hg38. The value k was set to 75 which is half of the read length of WGS data in this study. Each base in the genome was assigned a mappability level, based on the mapping ambiguity of all 75-mers overlapping this specific base. See below for the list of mappability levels. Regions with mappability level 0 and 1 were included in the low-mappability mask.

  - Level 0: all 75-mers overlapping this base could not be mapped to the genome uniquely.

  - Level 1: more than 50% of overlapping 75-mers are not uniquely mapped.

  - Level 2: more than 50% of overlapping 75-mers are uniquely mapped with 1-mismatch hits.

  - Level 3: more than 50% of overlapping 75-mers are uniquely mapped without 1-mismatch hits.

In addition, we used GATK CallableLoci to mark regions with high confidence of variant detection

based on coverage. Together, the intersection of unmasked regions and callable loci were defined

as the eligible territories for SRB detection. The 100 kB bins with less than 75% overlap with

eligible territories were excluded from the analysis.

Generating covariates for regression analysis

To accurately model the genomic features of mCRPC, we incorporated the following covariates.

• Nucleotide composition, including GC content, CpG fraction and TpC fraction per 10 kB
non-overlapping bin in the genome.

• Replication timing of LNCaP (data obtained from ENCODE under accession
ENCFF995YGM, lifted over from hg19 to hg38) (88, 89).

• DNase I hypersensitive sites (data obtained from ENCODE under accession
ENCFF434GSJ, lifted over to hg38).

• Repeats annotated by RepeatMasker, including LINE, SINE, LTR, DNA transposon and
simple repeats.

• Heterochromatin regions inferred by ChromHMM (90) with the 18-state model parameters
from the Roadmap Epigenomics Project (91), based LNCaP ChIP-seq data of H3K4me1,
H3K4me3, H3K4ac H3K27me3, H3K36me3 and H3K9me3.

• Common fragile sites downloaded from HGNC biomart (92).

SRB detection

Structural variants from the final call set were used for statistical enrichment of recurrent

breakpoints within 100 kB bins using a Gamma-Poisson regression implemented in the package,

fish.hook (44). Breakpoints of SVs were treated independently. The Benjamini-Hochberg

procedure was used for multiple testing correction and bins with q-value ≤ 0.1 were determined

to be significant. The distances of individual known driver genes to those significant bins were

663 evaluated based on the shortest genomic distance between the gene and bin boundaries,

664 regardless of gene orientations.

665 **Annotation of gene alteration status**

666 *Gene alteration by copy-number*

667 Copy-number segments were excluded if their cellular fraction was lower than 0.8, except for

668 those which were determined as copy neutral or copy-number greater than 4. The gene

669 annotation was based on known protein coding genes from GenCode release 30 (GRCh38.p12)

670 (93). For each gene, its copy-number was assigned to the copy-number value and LOH status of

671 the segment that has the largest overlap with it. The gene-level copy-number was normalized

672 based on ploidy of the corresponding sample, with autosomal genes normalized by the inferred

673 ploidy rounded to nearest integer, and X-linked genes normalized by half such value. Then the

674 copy-number status of each gene was categorized based on the following criteria.

675 • Amplification. Normalized gene-level copy-number is greater than or equal to 2.5.

676 • Gain. Normalized gene-level copy-number is between 2 and 2.5.

677 • Homozygous deletion. Normalized gene-level copy-number is 0.

678 • Deletion with LOH. Normalized gene-level copy-number is between 0 and 1, and LOH

679   status was found.

680 • Copy neutral LOH. Normalized gene-level copy-number is 1 and LOH status was found.

681 *Gene alteration by structural variant*

682 Gene coordinates were based on ENSEMBL v33 of hg38 (94). Gene body region of one gene

683 was defined as the widest region of all known isoforms collapsed. Gene flanking region was

684 defined as the corresponding two 1 Mb regions next to the gene body region on 5'-end and 3'-

685 end, respectively.

Gene alteration status by genome rearrangements was defined based on the breakpoints and

directions of involving structural variant events. A gene in one WGS sample (gene-sample pair)

was considered having gene transecting events if any breakpoints of SV events were located

within the gene body region. If the gene transecting status did not apply, then this gene-sample

pair was examined for gene flanking status if the breakpoints of any intra-chromosomal SV events,

including tandem duplications, deletions, and inversions, were located within the gene flanking

regions. Additionally, translocation events including intra-chromosomal balanced and unbalanced

events which spanned over 10 Mb, and inter-chromosomal translocation events were considered

altering the gene flanking regions if any of their breakpoints was in the gene flanking region, and

the direction of the SV was going towards the gene body region. The alteration status of

rearrangements for each gene-sample pair was exclusive between gene transecting and gene

flanking, with the former being prioritized in report.

AR alteration analysis

Copy-number of the *AR* gene (chrX:67,544,623-67,730,619) and the *AR* enhancer region

(chrX:66,895,000-66,910,000) were each computed as the mean corrected total copy-number

across the 10 kB bins overlapping each region. The copy-number was further normalized by

sample ploidy as previously described. Amplification status of *AR* was determined by comparing

the log2 fold-change *FC* of enhancer-level over gene-level copy-number. Four distinct groups

were defined based on copy-number and *FC* as below.

• Co-amplification. Ploidy normalized copy-number values of both *AR* gene body and

enhancer are greater than 1.5.

• Selective *AR* amplification. *FC* < -log2(1.5) and enhancer copy-number is less than 1.5.

• Selective enhancer copy gain. *FC* > log2(1.5) and *AR* gene body copy-number is less than

1.5.

• Lack of amplification for both. All other cases were considered as no amplification for both

regions.

ANCOVA test was used to test if different patterns of *AR* amplification have an impact on *AR*

expression. See Statistics section.

Gene expression

TPM values for a subset of the samples based on linked-read sequencing were obtained from

cBioportal (95, 96). For samples based on short-read sequencing the TPM values were obtained

from a previous study (13).

Gene fusion analysis

Fusion status of the main members of the ETS family, including *ERG*, *ETV1*, *ETV4*, *ETV5* and

ELK4 was analyzed. Determination of gene fusion status was based on both DNA and RNA levels.

For DNA, structural variants transecting gene body regions were used. SV events were

considered supporting gene fusion only if they satisfy the following criteria: (1) the breakpoints of

this event must be located within the ETS gene and another protein coding gene, respectively; (2)

the orientation of the breakpoint located within the ETS gene must be pointing towards the coding

sequence of ETS domain. For RNA, arriba was used to detect fusion transcripts from RNA-seq

data (97). The fusion status was only confirmed if all following conditions were satisfied: (1) the

complete ETS domain was included in the fusion product; (2) detection confidence reported by

arriba is "high"; (3) coding sequence in the fusion transcript was in sense orientation and no out-

of-frame shifts.

SV signature analysis

Signature extraction and clustering

De novo signature extraction was performed on all SV events called by SvABA of the combined

cohort using signature.tools.lib (55) with the recommended settings of 20 bootstraps, 200 repeats,

the clustering with matching algorithm, the KLD objective function, and RTOL = 0.001. The

exposure of one signature in one sample is defined as the median activity of the signature within

the sample across all bootstraps. For clustering, the reference signature exposure values for each

sample based on short-read sequencing were normalized such that the sum of exposure values

per sample is 1, and the normalized exposure values for each signature were mean-centered

across all samples. A Euclidean distance matrix was computed and then samples were clustered

with the Ward.D2 algorithm using R's hclust function. We chose the number of clusters to be k =

9 based on dendrogram using cutree function in R.

STATISTICS

Association of AR locus amplification status and AR expression

ANCOVA test was used to test if different patterns of *AR* amplification have an impact on *AR*

expression. Batch corrected log10(TPM+1) values using ComBat from sva R package (v3.34.0)

were used for *AR* expression level. We fit the ANCOVA model using *AR* expression as the

response variable, *AR* amplification status as the predictor variable, and ploidy, purity as

covariates. The function Anova in the car package (v3.0-5) was used with Type III sum of squares

for the model. Post hoc analysis was performed to determine the specific differences among four

different *AR* amplification status. The function glht was used within the multcomp package (v1.4-

11) in R to perform Tukey's Test for multiple comparisons.

 Enrichment of alterations in SV clusters

All 9 identified SV clusters were analyzed for enrichment of alterations. To make the analysis

unbiased by SV signature, we limited our search to alteration types that were orthogonal to

rearrangements, which include SNV, copy-number gain and copy-number loss. We performed

hypothesis testing on each driver-alteration pair, and also on chromoplexy and chromothripsis.

For each SV cluster, a χ^2 test was performed for each driver gene alteration status, with samples

within group being tested against samples belonging to all 8 other SV clusters. Multiple testing

adjustment based on Benjamini-Hochberg FDR was performed to compute q-values. Alteration

categories with q-values less than 0.25 were determined as enriched in the corresponding SV

cluster.

Survival analysis

Survival data was obtained from (98). Survival analyses were conducted using the Kaplan-Meier

method with log-rank testing for significance. The function survfit from survival R package was

used to perform the analysis.

STUDY APPROVAL

For tumor biopsies profiled via linked-read sequencing, samples were collected from individuals

with mCRPC who provided informed consent on institutional IRB-reviewed protocols, as

previously described (15). Uniformly reanalyzed data were generated as described in the

respective studies (9, 13, 18).

AUTHOR CONTRIBUTIONS

Conceptualization: M-E.T, M.M., S.R.V., G.H.

Methodology: M.Z., M.K., S.R.V., G.H.

Software: M.Z., M.K., A.C.H., G.H.

Formal Analysis: M.Z., M.K., A.C.H., K.L., Y.L., M.R., W.H., J.C-Z, S.R.V., G.H.

Data Curation: M.Z., M.K., A.C.H., Z.Z., S.R.V., G.H.

Writing – Original Draft: M.Z., M.M., G.H., S.R.V.

Writing – Review & Editing: M.Z., R.B., E.M.V., A.D.C. P.S.N., M.L.F., M-E.T., M.M., G.H., S.R.V.

Visualization: M.Z., M.K., A.C.H., S.R.V., G.H.

Supervision: M-E.T., M.M., S.R.V., G.H.

Funding Acquisition: S.R.V., G.H., M.M.

CONFLICTS OF INTERESTS

A.D.C.: Honoraria: OncLive, Bayer, Targeted Oncology, Aptitude Health, Journal of Clinical Pathways, Cancer Network; Consulting: Blackstone; Advisory Board: Clovis, Dendreon, Bayer, Eli Lilly, AstraZeneca, Astellas, Blue Earth; Research Funding: Bayer

E.M.V.: Advisory/Consulting: Tango Therapeutics, Genome Medical, Invitae, Enara Bio, Janssen, Manifold Bio, Monte Rosa; Research support: Novartis, BMS; Equity: Tango Therapeutics, Genome Medical, Syapse, Enara Bio, Manifold Bio, Microsoft, Monte Rosa; Travel reimbursement: Roche/Genentech; Patents: Institutional patents filed on chromatin mutations and immunotherapy

response, and methods for clinical interpretation; intermittent legal consulting on patents for

Foaley & Hoag

M-E.T.: Advisory boards: Janssen, Pfizer, Astra Zeneca, Bayer

M.L.F.: Served as a consultant to and has equity in Nuscan Diagnostics. This activity is outside

of the scope of this manuscript.

M.M.: Consultant for Bayer, Interline and Isobl; an inventor of patents licensed to LabCorp and

Bayer; and receives research funding from Bayer, Janssen, and Ono Pharmaceuticals.

P.S.N.: Served as a consultant to Bristol Myers Squibb, Janssen, and Pfizer in work unrelated to

the present study.

S.R.V.: Consulting (current or previous 3 years), MPM Capital and Vida Ventures; spouse is an

employee of and holds equity in Kojin Therapeutics.

All other authors declare no competing interests.

ACKNOWLEDGEMENTS

We thank the many patients and their families for their generosity in contributing to this study. We

also thank the Prostate Cancer Foundation (PCF) and Stand Up 2 Cancer (SU2C) International

Prostate Cancer Dream Team for contributions to specimen acquisition.

This work was supported by the National Institutes of Health (K22 CA237746 to G.H.; P01

CA163227 and R01 CA234715 to P.S.N.; R01 GM107427, R01 CA193910, and R01 CA251555

to M.L.F.; R35 CA197568 to M.M.), Department of Defense Prostate Cancer Research Program

(Physician Research Award W81XWH-17-1-0358 to S.R.V.; W81XWH-19-1-0565 and W81XWH-

21-1-0234 to M.L.F.; PC200262 to P.S.N.), PCF Young Investigator Awards (G.H. and S.R.V.),

PCF-Movember Challenge Award (to E.M.V.), Brotman Baty Institute for Precision Medicine (to

G.H.), the Fund for Innovation in Cancer Informatics Major Grant (to G.H.), the V Foundation

Scholar Grant (to G.H.), Wong Family Award in Translational Oncology and Dana-Farber Cancer

Institute Medical Oncology grant (to A.D.C.), H.L. Snyder Medical Research Foundation and the

Cutler Family Fund for Prevention and Early Detection (to M.L.F.), and the Pan-Mass Challenge

team IMAGINE (to M-E.T.), American Cancer Society Research Professor (M.M.).

This research was also supported in part by the NIH/NCI Cancer Center Support Grant P30

CA015704, Pacific Northwest Prostate Cancer SPORE (P50 CA097186), and Scientific

Computing Infrastructure (ORIP Grant S10OD028685).

REFERENCES

1. Leinonen R, et al. The sequence read archive. *Nucleic Acids Res* 2011;39(Database issue):D19-21.

2. de Bono JS, et al. Abiraterone and Increased Survival in Metastatic Prostate Cancer. *N Engl J Med* 2011;364(21):1995–2005.

3. Scher HI, et al. Increased Survival with Enzalutamide in Prostate Cancer after Chemotherapy. *New England Journal of Medicine* 2012;367(13):1187–1197.

4. Abida W, et al. Analysis of the Prevalence of Microsatellite Instability in Prostate Cancer and Response to Immune Checkpoint Blockade. *JAMA Oncol* 2019;5(4):471–478.

5. Abida W, et al. Rucaparib in Men With Metastatic Castration-Resistant Prostate Cancer Harboring a BRCA1 or BRCA2 Gene Alteration. *J Clin Oncol* 2020;38(32):3763–3772.

6. de Bono J, et al. Olaparib for Metastatic Castration-Resistant Prostate Cancer. *N Engl J Med* 2020;382(22):2091–2102.

7. Pritchard CC, et al. Inherited DNA-Repair Gene Mutations in Men with Metastatic Prostate Cancer. *New England Journal of Medicine* 2016;375(5):443–453.

8. Baca SC, et al. Punctuated evolution of prostate cancer genomes. *Cell* 2013;153(3):666–677.

9. Campbell PJ, et al. Pan-cancer analysis of whole genomes. *Nature* 2020;578(7793):82–93.

10. van Dessel LF, et al. The genomic landscape of metastatic castration-resistant prostate cancers reveals multiple distinct genotypes with potential clinical impact. *Nat Commun* 2019;10(1):5251.

11.   Glodzik D, et al. A somatic-mutational process recurrently duplicates germline susceptibility

loci and tissue-specific super-enhancers in breast cancers. *Nat Genet* 2017;49(3):341–348.

12.   Nik-Zainal S, et al. Landscape of somatic mutations in 560 breast cancer whole-genome

sequences. *Nature* 2016;534(7605):47–54.

13.   Quigley DA, et al. Genomic Hallmarks and Structural Variation in Metastatic Prostate Cancer.

Cell 2018;174(3):758-769.e9.

14.   Stephens PJ, et al. Massive genomic rearrangement acquired in a single catastrophic event

during cancer development. *Cell* 2011;144(1):27–40.

15.   Viswanathan SR, et al. Structural Alterations Driving Castration-Resistant Prostate Cancer

Revealed by Linked-Read Genome Sequencing. *Cell* 2018;174(2):433-447.e19.

16.   Weinhold N, et al. Genome-wide analysis of noncoding regulatory mutations in cancer. *Nat

Genet* 2014;46(11):1160–1165.

17.   Woodcock DJ, et al. Prostate cancer evolution from multilineage primary to single lineage

metastases with implications for liquid biopsy. *Nat Commun* 2020;11(1):5070.

18.   Gerhauser C, et al. Molecular Evolution of Early-Onset Prostate Cancer Identifies Molecular

Risk Markers and Clinical Trajectories. *Cancer Cell* 2018;34(6):996-1011.e8.

19.   Espiritu SMG, et al. The Evolutionary Landscape of Localized Prostate Cancers Drives

Clinical Aggression. *Cell* 2018;173(4):1003-1013.e15.

20.   Wedge DC, et al. Sequencing of prostate cancers identifies new cancer genes, routes of

progression and drug targets. *Nat Genet* 2018;50(5):682–692.

21. Fraser M, et al. Genomic hallmarks of localized, non-indolent prostate cancer. *Nature*
2017;541(7637):359–364.

22. Gundem G, et al. The evolutionary history of lethal metastatic prostate cancer. *Nature*
2015;520(7547):353–357.

23. Mateo J, et al. Accelerating precision medicine in metastatic prostate cancer. *Nat Cancer*
2020;1(11):1041–1053.

24. Armenia J, et al. The long tail of oncogenic drivers in prostate cancer. *Nature Genetics*
2018;50(5):645–651.

25. Grasso CS, et al. The mutational landscape of lethal castration-resistant prostate cancer.
Nature 2012;487(7406):239–243.

26. Chen CD, et al. Molecular determinants of resistance to antiandrogen therapy. *Nature*
Medicine 2004;10(1):33–39.

27. Yuan X, et al. Androgen receptor functions in castration-resistant prostate cancer and
mechanisms of resistance to new agents targeting the androgen axis. *Oncogene*
2014;33(22):2815–2825.

28. Li Y, et al. Diverse AR Gene Rearrangements Mediate Resistance to Androgen Receptor
Inhibitors in Metastatic Prostate Cancer. *Clinical Cancer Research* 2020;26(8):1965–1976.

29. Brand LJ, Dehm SM. Androgen Receptor Gene Rearrangements: New Perspectives on
Prostate Cancer Progression. *Curr Drug Targets* 2013;14(4):441–449.

30. Céraline J, et al. Constitutive activation of the androgen receptor by a point mutation in the

hinge region: a new mechanism for androgen-independent growth in prostate cancer. *Int. J.*

Cancer 2004;108(1):152–157.

31. Henzler C, et al. Truncation and constitutive activation of the androgen receptor by diverse

genomic rearrangements in prostate cancer. *Nat Commun* 2016;7:13668.

32. Takeda DY, et al. A Somatically Acquired Enhancer of the Androgen Receptor Is a

Noncoding Driver in Advanced Prostate Cancer. *Cell* 2018;174(2):422-432.e13.

33. Li Y, et al. Patterns of somatic structural variation in human cancer genomes. *Nature*

2020;578(7793):112–121.

34. Sailer V, et al. Bone Biopsy Protocol for Advanced Prostate Cancer in the Era of Precision

Medicine. *Cancer* 2018;124(5):1008–1015.

35. Wu Y-M, et al. Inactivation of CDK12 Delineates a Distinct Immunogenic Class of Advanced

Prostate Cancer. *Cell* 2018;173(7):1770-1782.e14.

36. Robinson D, et al. Integrative Clinical Genomics of Advanced Prostate Cancer. *Cell*

2015;162(2):454.

37. Pitkänen E, et al. Frequent L1 retrotranspositions originating from TTC28 in colorectal

cancer. *Oncotarget* 2014;5(3):853–859.

38. Pradhan B, et al. Detection of subclonal L1 transductions in colorectal cancer by long-

distance inverse-PCR and Nanopore sequencing. *Sci Rep* 2017;7(1):14521.

39. Tubio JMC, et al. Mobile DNA in cancer. Extensive transduction of nonrepetitive DNA

mediated by L1 retrotransposition in cancer genomes. *Science* 2014;345(6196):1251343.

40. Chen J, et al. The t(1;3) breakpoint-spanning genes LSAMP and NORE1 are involved in

clear cell renal cell carcinomas. *Cancer Cell* 2003;4(5):405–413.

41. Kresse SH, et al. LSAMP, a novel candidate tumor suppressor gene in human

osteosarcomas, identified by array comparative genomic hybridization. *Genes*

Chromosomes Cancer 2009;48(8):679–693.

42. Kühn MWM, et al. High-resolution genomic profiling of adult and pediatric core-binding factor

acute myeloid leukemia reveals new recurrent genomic alterations. *Blood* 2012;119(10):e67.

43. Veeriah S, et al. The tyrosine phosphatase PTPRD is a tumor suppressor that is frequently

inactivated and mutated in glioblastoma and other human cancers. *PNAS*

2009;106(23):9435–9440.

44. Imielinski M, Guo G, Meyerson M. Insertions and Deletions Target Lineage-Defining Genes

in Human Cancers. *Cell* 2017;168(3):460-472.e14.

45. Eteleeb AM, et al. SV-HotSpot: detection and visualization of hotspots targeted by structural

variants associated with gene expression. *Sci Rep* 2020;10(1):15890.

46. Pomerantz MM, et al. Prostate cancer reactivates developmental epigenomic programs

during metastatic progression. *Nat. Genet.* [published online ahead of print: July 20, 2020];

doi:10.1038/s41588-020-0664-8

47. Tomlins SA, et al. Recurrent fusion of TMPRSS2 and ETS transcription factor genes in

prostate cancer. *Science* 2005;310(5748):644–648.

48. Tomlins SA, et al. Distinct classes of chromosomal rearrangements create oncogenic ETS

gene fusions in prostate cancer. *Nature* 2007;448(7153):595–599.

49.   Kumar-Sinha C, Kalyana-Sundaram S, Chinnaiyan AM. Landscape of gene fusions in

930   epithelial cancers: seq and ye shall find. *Genome Medicine* 2015;7(1):129.

931   50.   Qin F, et al. SLC45A3-ELK4 functions as a long non-coding chimeric RNA. *Cancer Lett*

932   2017;404:53–61.

933   51.   Rickman DS, et al. SLC45A3-ELK4 is a novel and frequent erythroblast transformation-

934   specific fusion transcript in prostate cancer. *Cancer Res* 2009;69(7):2734–2738.

935   52.   Zhang Y, et al. Chimeric transcript generated by cis-splicing of adjacent genes regulates

936   prostate cancer cell proliferation. *Cancer Discov* 2012;2(7):598–607.

937   53.   Visakorpi T, et al. In vivo amplification of the androgen receptor gene and progression of

938   human prostate cancer. *Nat. Genet.* 1995;9(4):401–406.

939   54.   Umbreit NT, et al. Mechanisms generating cancer genome complexity from a single cell

940   division error. *Science* 2020;368(6488):eaba0712.

941   55.   Degasperi A, et al. A practical framework and online tool for mutational signature analyses

942   show inter-tissue variation and driver dependencies. *Nat Cancer* 2020;1(2):249–263.

943   56.   Willis NA, et al. Mechanism of tandem duplication formation in BRCA1-mutant cells. *Nature*

944   2017;

945   57.   Barbieri CE, et al. Exome sequencing identifies recurrent SPOP, FOXA1 and MED12

946   mutations in prostate cancer. *Nat Genet* 2012;44(6):685–689.

947   58.   Nguyen B, et al. Pan-cancer Analysis of CDK12 Alterations Identifies a Subset of Prostate

948   Cancers with Distinct Genomic and Clinical Characteristics. *European Urology*

949   2020;78(5):671–679.

59. Cancer Genome Atlas Research Network. The Molecular Taxonomy of Primary Prostate
Cancer. *Cell* 2015;163(4):1011–1025.

60. Giambartolomei C, et al. H3K27ac HiChIP in prostate cell lines identifies risk genes for
prostate cancer susceptibility. *The American Journal of Human Genetics*
2021;108(12):2284–2300.

61. Taplin ME, et al. Mutation of the androgen-receptor gene in metastatic androgen-
independent prostate cancer. *N. Engl. J. Med.* 1995;332(21):1393–1398.

62. Haffner MC, et al. Androgen-induced TOP2B-mediated double-strand breaks and prostate
cancer gene rearrangements. *Nat Genet* 2010;42(8):668–675.

63. Petrovics G, et al. A novel genomic alteration of LSAMP associates with aggressive prostate
cancer in African American men. *EBioMedicine* 2015;2(12):1957.

64. Alexandrov LB, et al. Signatures of mutational processes in human cancer. *Nature*
2013;500(7463):415–421.

65. Alexandrov LB, et al. The repertoire of mutational signatures in human cancer. *Nature*
2020;578(7793):94–101.

66. Macintyre G, et al. Copy number signatures and mutational processes in ovarian carcinoma.
Nat Genet 2018;50(9):1262–1270.

67. Wang S, et al. Copy number signature analysis tool and its application in prostate cancer
reveals distinct mutational processes and clinical outcomes. *PLoS Genet*
2021;17(5):e1009557.

68. Van der Auwera GA, O'Connor B. Genomics in the Cloud2020;https://www.oreilly.com/library/view/genomics-in-the/9781491975183/. cited December 10, 2021

69. Carrot-Zhang J, Majewski J. LoLoPicker: detecting low allelic-fraction variants from low-quality cancer samples. *Oncotarget* 2017;8(23):37032.

70. Saunders CT, et al. Strelka: accurate somatic small-variant calling from sequenced tumor-normal sample pairs. *Bioinformatics* 2012;28(14):1811–1817.

71. Martincorena I, et al. Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* 2017;171(5):1029-1041.e21.

72. Tate JG, et al. COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* 2019;47(D1):D941–D947.

73. Ha G, et al. TITAN: inference of copy number architectures in clonal cell populations from tumor whole-genome sequence data. *Genome Res* 2014;24(11):1881–1893.

74. Adalsteinsson VA, et al. Scalable whole-exome sequencing of cell-free DNA reveals high concordance with metastatic tumors. *Nat. Commun.* 2017;8(1):1324.

75. Wala JA, et al. SvABA: genome-wide detection of structural variants and indels by local assembly. *Genome Res* 2018;28(4):581–591.

76. Spies N, et al. Genome-wide reconstruction of complex structural variants using read clouds. *Nature Methods* 2017;

77. Cortés-Ciriano I, et al. Comprehensive analysis of chromothripsis in 2,658 human cancers using whole-genome sequencing. *Nat Genet* 2020;52(3):331–341.

78.   Barrett T, et al. NCBI GEO: archive for functional genomics data sets--update. *Nucleic Acids*

Res 2013;41(Database issue):D991-995.

79.   Gene Expression Omnibus: NCBI gene expression and hybridization array data repository -

PubMedhttps://pubmed.ncbi.nlm.nih.gov/11752295/. cited January 6, 2022

80.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics 2009;25(14):1754–1760.

81.   Zhang Y, et al. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* 2008;9(9):R137.

82.   Bert SA, et al. Regional activation of the cancer genome by long-range epigenetic

remodeling. *Cancer Cell* 2013;23(1):9–22.

83.   Du Q, et al. Replication timing and epigenome remodelling are associated with the nature of

chromosomal rearrangements in cancer. *Nat Commun* 2019;10(1):416.

84.   Taberlay PC, et al. Three-dimensional disorganization of the cancer genome occurs

coincident with long-range genetic and epigenetic alterations. *Genome Res* 2016;26(6):719–

731.

85.   Pomerantz MM, et al. The androgen receptor cistrome is extensively reprogrammed in

human prostate tumorigenesis. *Nat Genet* 2015;47(11):1346–1351.

86.   Mallick S, et al. The Simons Genome Diversity Project: 300 genomes from 142 diverse

populations. *Nature* 2016;538(7624):201–206.

87.   Smit, AFA, Hubley, R & Green, P. *RepeatMasker Open-4.0..* 2013:

88.   Davis CA, et al. The Encyclopedia of DNA elements (ENCODE): data portal update. *Nucleic*

Acids Res 2018;46(D1):D794–D801.

89.   ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human

1013   genome. *Nature* 2012;489(7414):57–74.

1014   90.   Ernst J, Kellis M. ChromHMM: automating chromatin-state discovery and characterization.

1015   *Nat Methods* 2012;9(3):215–216.

1016   91.   Roadmap Epigenomics Consortium, et al. Integrative analysis of 111 reference human

1017   epigenomes. *Nature* 2015;518(7539):317–330.

1018   92.   Tweedie S, et al. Genenames.org: the HGNC and VGNC resources in 2021. *Nucleic Acids*

1019   *Res* 2021;49(D1):D939–D946.

1020   93.   Frankish A, et al. GENCODE reference annotation for the human and mouse genomes.

1021   *Nucleic Acids Res* 2019;47(D1):D766–D773.

1022   94.   Howe KL, et al. Ensembl 2021. *Nucleic Acids Res* 2021;49(D1):D884–D891.

1023   95.   Cerami E, et al. The cBio Cancer Genomics Portal: An Open Platform for Exploring

1024   Multidimensional Cancer Genomics Data: Figure 1.. *Cancer Discovery* 2012;2(5):401–404.

1025   96.   Gao J, et al. Integrative analysis of complex cancer genomics and clinical profiles using the

1026   cBioPortal. *Sci Signal* 2013;6(269):pl1.

1027   97.   Uhrig S, et al. Accurate and efficient detection of gene fusions from RNA sequencing data.

1028   *Genome Res* 2021;31(3):448–460.

1029   98.   Chen WS, et al. Genomic Drivers of Poor Prognosis and Enzalutamide Resistance in

1030   Metastatic Castration-resistant Prostate Cancer. *Eur Urol* 2019;76(5):562–571.

1031

Figure 1

Figure 1. Study design and genomic landscape of mCRPC.

(A) Workflow of study and data analysis. Tumor specimens (grey) from both primary prostate cancer and mCRPC were included in this study. Linked-read and short-read whole-genome sequencing (WGS) and RNA-sequencing datasets were either generated for this study or reanalyzed from prior studies (13, 15). A pooled dataset of 143 mCRPC samples with WGS data was used in this study after curation (**Methods**). Genomic alteration call-sets for 278 primary localized prostate cancer samples were obtained from ICGC/TCGA Pan-Cancer Analysis of Whole Genomes (PCAWG) (9, 33). For 125 mCRPC samples, RNA-seq was used. The overview of the genomic alteration and characterization analysis is shown.

(B) Clinical annotations and somatic alterations for 143 patient samples in the pooled mCRPC cohort. Samples are ordered by treatment type; the four patients with pre-treatment and post-progression pairs are placed at the right. (Top) Clinical and sample information and genomic pattern classifications. (Middle) Distribution of genomic rearrangement types in individual samples. (Bottom) Mutational burden for SNVs and indels computed as number of mutations per mega-

base pair (Mb). Y-axis shown in logarithmic scale. Threshold lines indicates mutational burden at 2.5 and 5 mutations per Mb.

(C) Genomic rearrangement alteration profiles of key mCRPC genes. (Top) Events were categorized into gene transecting and gene flanking events (**Methods**). Gene transecting: if any of its breakpoints was located within the gene body region. Gene flanking: rearrangements which were not gene transecting and had breakpoints located within 1 Mb of either transcription start site or termination site of the gene. Only 159 genes reported and known to be involved in prostate cancer were considered in this analysis (**Table S1G and S1H**). (Middle) Frequency and distribution of rearrangement types for gene transecting events; genes with ≥ 10% frequency are shown. Gene transecting events were prioritized over flanking events during annotation. The category "Multiple" represents gene-sample pairs carrying more than one type of rearrangement event. (Bottom) Frequency of gene flanking events by tandem duplication; genes with ≥ 10% are shown.

Figure 2

Figure 2. Genome-wide analysis of genomic rearrangements in mCRPC.

(A) Analysis of significantly recurrent breakpoint (SRB) identified regions of rearrangement hotspots, genome-wide, using a Gamma-Poisson regression model. Each dot corresponds to a 100 kB bin (n=26,663 total bins). Statistically significant SRB bins with FDR (Benjamini-Hochberg) q-value ≤ 0.1 (n=55) are colored based on the distance to the nearest known prostate cancer driver gene, within 1 Mb. The driver genes within 1 Mb of the SRB bins are labeled. A square bracket is used for genes spanning multiple bins. Bins with q-value > 0.1 were not significant (grey).

(B) Comparison of SV alteration frequency in mCRPC versus primary localized prostate cancer. The union set of genes (n=14) within 1 Mb of SRB hotspot regions in mCRPC and localized prostate cancer cohorts was included in the comparison. The frequencies represent total gene transecting and flanking SV events. All labeled genes were significantly enriched in either mCRPC or primary localized tumors (Fisher's test, p-value < 0.05).

(C) Patterns of rearrangements at the loci of driver genes identified at SRB regions in mCRPC cohort of 143 tumors. Cumulative counts of intra-chromosomal SV events (tandem duplications "TandemDup", deletions, and inversions) were computed based on the breakpoints and span of the events. Histone H3 lysine 27 acetylation (H3K27ac) and *AR* binding sites (ARBS) specific to mCRPC were obtained from a previous study (46). Inter-chromosomal translocations are not shown. Genome coordinates based on hg38 build.

(D) Overlap of *AR* binding sites (ARBS) within SRB hotspots of mCRPC (55 regions) and primary localized prostate (47 regions) cohorts. Metastatic-specific and primary localized-specific ARBS were obtained from previous studies (46, 85). χ^2 test of independence p-values are shown.

(E) Fusion status and expression of selected genes in ETS transcription factor gene family in the mCRPC cohort with WGS and RNA-seq data. Fusion type was defined as the data evidence that supported the event: DNA-only, corresponds to WGS; RNA-only, corresponds to RNA-seq; DNA+RNA, corresponds to support from both WGS and RNA-seq. Each dot represents a tumor sample and is colored based on fusion type of each sample; grey indicates no evidence of fusion event. Data shown for samples with available expression data for the specific ETS gene. Gene expression values of full-length transcripts are z-score normalized.

(F) Fusion profile of *ETV1*. DNA rearrangement breakpoints supporting the fusion (purple bars) are indicated with the corresponding fusion partners. Exons of the ETS domain (red) are indicated. Genome coordinates based on hg38 build.

(G) Summary of fusion partners for selected genes in ETS transcription factor gene family in mCRPC cohort. Fusion events and partners are indicated by flow connections. Total counts of individual fusion events and partners across the cohort are shown.

Figure 3

Figure 3. Modes of *AR* activation in mCRPC.

(A) Copy number of *AR* gene and its enhancer (~624 kB upstream) for mCRPC cohort samples after adjustment by tumor purity and sample ploidy normalization. Data shown for samples with available *AR* gene expression data. (Left) Copy number of *AR* and its enhancer are shown in $\log_2$ scale, colored based on *AR* gene expression level (transcripts per million, TPM). (Right) Excerpt of figure highlighting *AR* expression for samples with lower copy number values.

(B) *AR* expression for *AR* locus copy number status for 122 samples with available *AR* gene expression data. ANCOVA test was performed to account for tumor purity and ploidy as covariates. TukeyHSD p-values for pair-wise comparisons between groups with *AR* locus amplification status and groups with no amplification.

(C) Patterns of rearrangements involving the *AR* locus in 143 mCRPC samples. Presence of specific alteration events and complex rearrangements (black) were predicted automatically and manually curated. *AR* gene expression shown (blue shades) for same samples in (B); samples with no available expression data are indicated in grey. Representative examples of each category are presented in (D) to (H).

(D-H) Complex and simple rearrangement patterns involving the *AR* locus, including focal duplication events on *AR* enhancer **(D)**, intragenic amplification event leading to a breakpoint within *AR* between exons 4 and 5 **(E)**, chromosomal level chromothripsis events involving *AR* and enhancer **(F)**, arm-level chromothripsis coinciding with *AR* amplification by break-fusion-break cycle **(G)**, extra-chromosomal DNA amplicon including *AR* and enhancer **(H)**. *AR* gene boundary (green) and its enhancer (yellow) are shown; concave arcs, intra-chromosomal SV events; convex arcs, inter-chromosomal SV events. Copy number values represent 10 kB bins and have been tumor purity corrected.

Figure 4. Clustering of mCRPC SV signatures

SV signature analysis and hierarchical clustering identifies nine distinct molecular groups. (Top) Dendrogram of the clustering of SV signature exposure. The prevalence of each signature was computed based on having ≥ 0.05 exposure (proportion of SVs). (Middle) Enrichment of altered prostate cancer drivers. Enriched alterations in Cluster 1, 3, 5, 6, and 7 are shown based on statistical significance by χ^2 test. (Bottom) Composition of SV types and sizes for each SV cluster, separated by non-clustered (nc) and clustered (c) SV events.