Divergent sensory and immune gene evolution in sea turtles with contrasting demographic and life histories

Blair P. Bentley[1,†], Tomás Carrasco-Valenzuela[2,3], Elisa K. S. Ramos[2,3,4], Harvinder Pawar[5], Larissa Souza Arantes[2,3], Alana Alexander[6], Shreya M. Banerjee[1,7], Patrick Masterson[8], Martin Kuhlwilm[5,9,10], Martin Pippel[11,12], Jacquelyn Mountcastle[13,22], Bettina Haase[13], Marcela Uliano-Silva[2,3], Giulio Formenti[13,14], Kerstin Howe[15], William Chow[15], Alan Tracey[15], Ying Sims[15], Sarah Pelan[15], Jonathan Wood[15], Kelsey Yetsko[16], Justin R. Perrault[17], Kelly Stewart[18], Scott R. Benson[19,20], Yaniv Levy[21], Erica V. Todd[22], H. Bradley Shaffer[23,24], Peter Scott[23,25], Brian T. Henen[26], Robert W. Murphy[27], David W. Mohr[28], Alan F. Scott[28], David J. Duffy[16,29], Neil J. Gemmell[30], Alexander Suh[31,32], Sylke Winkler[11,33], Françoise Thibaud-Nissen[8], Mariana F. Nery[4], Tomas Marques-Bonet[5,34,35,36], Agostinho Antunes[37,38], Yaron Tikochinski[39], Peter H. Dutton[17], Olivier Fedrigo[13], Eugene W. Myers[11,12,40], Erich D. Jarvis[13,14,41], Camila J. Mazzoni[2,3, †,*], and Lisa M. Komoroske[1, †,*]

[1]Department of Environmental Conservation, University of Massachusetts, Amherst, MA, USA; [2]Evolutionary Genetics Department, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany; [3]Berlin Center for Genomics in Biodiversity Research, Berlin, Germany; [4]Department of Genetics, Evolution, Microbiology and Immunology, University of Campinas, Campinas, Brazil; [5]Institut de Biologia Evolutiva, (CSIC-Universitat Pompeu Fabra), PRBB, Doctor Aiguader 88, Barcelona, Catalonia, Spain; [6]Department of Anatomy, School of Biomedical Sciences, University of Otago, Dunedin, New Zealand; [7]Department of Biology, Stanford University, Stanford CA, USA; [8]National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD, USA; [9]Department of Evolutionary Anthropology, University of Vienna, Vienna, Austria; [10]Human Evolution and Archaeological Sciences (HEAS), University of Vienna, Vienna, Austria; [11]Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany; [12]Center for Systems Biology, Dresden, Germany; [13]Vertebrate Genome Lab, The Rockefeller University, New York, NY, USA; [14]Laboratory of Neurogenetics of Language, The Rockefeller University, New York, NY, USA; [15]Tree of Life, Wellcome Sanger Institute, Cambridge, CB10 1SA, UK; [16]The Whitney Laboratory for Marine Bioscience and Sea Turtle Hospital, University of Florida, St. Augustine, Florida 32080, USA; [17]Loggerhead Marinelife Center, Juno Beach, FL, USA; [18]Marine Mammal and Turtle Division, Southwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, La Jolla, CA, United States; [19]Marine Mammal and Turtle Division, Southwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Moss Landing, CA, USA; [20]Moss Landing Marine Laboratories, Moss Landing, CA, USA; [21]National Sea Rescue Centre, Israel's Nature and Parks Authority, Mevoot Yam, Michmoret, Israel; [22]School of Life and Environmental Sciences, Deakin University, Queenscliff, VIC, Australia; [23]Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA, USA; [24]La Kretz Center for California Conservation Science, Institute of the Environment and Sustainability, University of California, Los Angeles, CA, USA; [25]Department of Life, Earth, and Environmental Sciences, West Texas A&M University, Canyon, TX, USA; [26]Environmental Affairs, Marine Air Ground Task Force and Training Command, Marine Corps Air Ground Combat Center, Twentynine Palms CA, USA; [27]Centre for Biodiversity, Royal Ontario Museum, Toronto, Ontario, Canada; [28]Genetic Resources Core Facility, McKusick-Nathans Dept of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore MD 21287, USA; [29]Department of Biology, University of Florida, Gainesville, Florida, 32611, USA; [30]Allan Wilson Centre, Department of Anatomy, University of Otago, Dunedin, New Zealand; [31]School of Biological Sciences, University of East Anglia, Norwich NR4 7TU, UK; [32]Department of Organismal Biology, Evolutionary Biology Centre (EBC), Science for Life Laboratory, Uppsala University, Uppsala, Sweden; [33]DRESDEN concept Genome Center, Dresden, Germany; [34]CNAG-CRG, Centre for Genomic Regulation (CRG), Barcelona Institute of Science and Technology (BIST), Baldiri i Reixac 4, 08028 Barcelona, Spain; [35]Institucio Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Catalonia 08010, Spain; [36]Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona; [37]CIIMAR/CIMAR, Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Av. General Norton de Matos, s/n, 4450-208, Porto, Portugal; [38]Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, s/n, 4169-007, Porto, Portugal; [39]Faculty of Marine Sciences, Ruppin Academic Center, Michmoret, Israel; [40]Faculty of Computer Science, Technical University Dresden, Dresden, Germany; [41]Howard Hughes Medical Institute, Chevy Chase, MD, USA.

Classification: Genetics and Evolution

Keywords: marine turtle; gene evolution; conservation genomics; genetic diversity; demographic history

Abbreviations: TE - transposable element; RE - repetitive element; RRC - region of reduced collinearity; FP – Fibropapillomatosis; ROH – runs of homozygosity

[†]Corresponding authors: (1) Blair P. Bentley (bbentley@umass.edu); (2) Camila J. Mazzoni (mazzoni@izw-berlin.de); (3) Lisa M. Komoroske (lkomoroske@umass.edu).
* These authors contributed equally to the work: Camila J. Mazzoni and Lisa M. Komoroske.

Abstract

Sea turtles represent an ancient lineage of marine vertebrates that evolved from terrestrial ancestors over 100 MYA, yet the genomic basis of the unique physiological and ecological traits enabling these species to thrive in diverse marine habitats remains largely unknown. Additionally, many populations have drastically declined due to anthropogenic activities over the past two centuries, and their recovery is a high global conservation priority. We generated and analyzed high-quality reference genomes for the leatherback (*Dermochelys coriacea*) and green (*Chelonia mydas*) turtles, representing the two extant sea turtle families. These genomes are highly syntenic and homologous, but localized regions of non-collinearity were associated with higher copy numbers of immune, zinc-finger, and olfactory receptor (OR) genes in green turtles, with ORs related to waterborne odorants greatly expanded in green turtles. Our findings suggest that divergent evolution of these key gene families may underlie immunological and sensory adaptations assisting navigation, occupancy of neritic versus pelagic environments, and diet specialization. Reduced collinearity was especially prevalent in microchromosomes, with greater gene content, heterozygosity, and genetic distances between species, supporting their critical role in vertebrate evolutionary adaptation. Finally, diversity and demographic histories starkly contrasted between species, indicating that leatherback turtles have had a low yet stable effective population size, exhibit extremely low diversity compared to other reptiles, and harbor a higher genetic load compared to green turtles, reinforcing concern over their persistence under future climate scenarios. These genomes provide invaluable resources for advancing our understanding of evolution and conservation best practices in an imperiled vertebrate lineage.

Statement of significance

Sea turtle populations have undergone recent global declines. We analyzed *de novo* assembled genomes for both extant sea turtle families through the Vertebrate Genomes Project to inform their conservation and evolutionary biology. These highly conserved genomes were differentiated by localized gene-rich regions of divergence, particularly within microchromosomes, suggesting that these genomic elements play key functional roles in the evolution of sea turtles and possibly other vertebrates. We further demonstrate that dissimilar evolutionary histories impact standing genomic diversity and genetic load, and are critical to consider when using these metrics to assess adaptive potential and extinction risk. Our results also demonstrate how reference genome quality impacts inferences of comparative and conservation genomics analyses that need to be considered in their application.

Introduction

Sea turtles recolonized marine environments over 100 MYA (1, 2) and are now one of the most
widely distributed vertebrate groups on the planet (3). Leatherback turtles (*Dermochelys coriacea*)
represent the only remaining species of the family Dermochelyidae, which diverged from the Cheloniidae
(hard-shelled sea turtles) about 60 MYA (4). Unique morphological (Fig. 1a) and physiological traits
allow leatherback turtles to exploit cool, highly productive pelagic habitats (5, 6), while green turtles
(*Chelonia mydas*) and other hard-shelled species largely inhabit warmer nearshore habitats following an
early pelagic life stage. Most previous research in this group has focused on organismal and ecological
adaptations (7), but the genomic basis of traits that differentiate or unite these species is not well
understood.

Anthropogenic pressures have caused substantial population declines in sea turtles, with
contemporary populations currently representing mere fractions of their historical abundances (8, 9).
Although sea turtles spend most of their life in the ocean, they also exhibit long-distance migrations to
natal rookeries for terrestrial reproduction (7, 10, 11). Consequently, they are threatened by human
activities in both terrestrial and marine environments, including direct harvest of meat and eggs (12),
fisheries bycatch (13), coastal development (14, 15), pollution (16), disease (17), and climate change (18,
19), which is exacerbated by their temperature-dependent mechanism of sex determination (TSD) altering
population dynamics (20, 21). The IUCN lists most sea turtle species as vulnerable or endangered, and
while decades of conservation efforts have fueled positive trends for some populations (22), others
continue to decline (23). In particular, leatherback turtles have undergone extensive declines (>95% in
some populations) over the last century (24–27), including the extirpation of the Malaysian nesting
population (28). Leatherback turtle recovery is also impeded by relatively low hatching success compared
to other sea turtle species (29). In contrast, many green turtle populations have recently increased
following conservation actions (22), but their continued recovery remains threatened by anthropogenic
activities and high incidence of the neoplastic disease fibropapillomatosis (FP), a likely viral-mediated
tumor disease that disproportionately impacts this species (30).

Genomic data have been instrumental in advancing understanding of species' evolutionary
histories and ecological adaptations (31–33), and providing critical information for conservation
management (34–37). However, this research has been hampered in taxa where genomic resources remain
limited. In particular, the lack of high-quality reference genomes, which are essential for accurate
comparative evolutionary analyses (38, 39) and robust estimates of a range of metrics to inform
conservation biology such as inbreeding, hybridization, disease susceptibility, genetic load, and
adaptation (36, 40, 41), impede this work in threatened species. An early draft genome for the green turtle
was assembled almost a decade ago (42), and provided important insights into turtle evolution. However,
errors, gaps, mis-assemblies, and fragmentation in draft genomes can lead to spurious inferences,
potentially masking signals of interest (38, 43) and impeding effective management strategies (41). Well-
annotated, chromosomal-level reference genomes can resolve these issues, improving our understanding
of the genomic underpinnings of ecological and evolutionary adaptations (39, 44). For example, high-
quality genomes with accurate annotations have enabled examination of gene changes associated with
recolonization of the marine environment by terrestrial vertebrates, including the loss of olfactory
receptor (OR) gene families (32, 45). Comparative genomic analyses have also demonstrated adaptive
diversity in genes underlying reptilian immunity (46), and high-quality genomes have provided key
insights into mammalian disease susceptibility (33, 47, 48). Such investigations are critical for sea turtles,

with diseases such as FP adversely impacting populations across the globe (30), and information on
immune genes is needed for devising effective conservation strategies (49).
We assembled chromosome-level reference genomes for leatherback and green turtles as part of
the Vertebrate Genomes Project (VGP), and leveraged these resources to address questions centered
around sea turtle evolutionary history and conservation. Specifically, we provide insights into the
genomic underpinnings of phenotypic traits that separate and unite these two species by examining
genome synteny and regions of divergence. Given the contrasting recent population trends of these two
species, we additionally used whole genome re-sequencing (WGR) data of individuals representative of
global populations to compare key conservation-relevant metrics, including patterns of diversity and
deleterious variants across the genomes, and reconstructed demographic histories to inform assessments
of future vulnerability. These genomes represent two of the most contiguous reptilian genomes assembled
to date, and our results provide a foundation for further hypothesis-driven investigations into the
evolutionary adaptation and conservation of this imperiled vertebrate lineage.
143
Results
Genome quality
Reference genomes for the leatherback and green turtles were generated using four genomic
technologies following the VGP pipeline v1.6 (39), with minor modifications (see Methods). A total of
100% of the leatherback and 99.8% of the green turtle assembled sequences were placeable within
chromosomes. The assembled genomes were near full-length (~2.1 GB), with annotations of all 28 known
chromosomes for both species, composed of 11 macrochromosomes (>50 Mb) and 17 microchromosomes
(<50 Mb) (Tables 1 & S1, Fig. S1). These genomes are among the highest quality genomes assembled for
non-avian reptiles to date in terms of both contiguity and completeness (Table S3), with the leatherback
turtle assembly representing the first reptile genome where all scaffolds were assigned to chromosomes.
Scaffold N50s were high for both genomes (Table 1). We annotated 18,775 protein-coding genes in the
leatherback and 19,752 in the green turtle genomes (see below for analysis of these gene differences). For
the leatherback and green turtles, 96.9% and 97.5% of these genes were supported at >95% of their length
from experimental evidence and/or high-quality protein models from related species (see Methods). The
numbers of protein-coding genes are within the range of other reptiles (Table S3) and include 97.7% and
98.2% complete BUSCO copies for leatherback and green turtles based on Sauropsida models (50), which
are similar or higher than all other assembled reptilian genomes to date (Fig. S2).
161
Genome architecture
Despite diverging over 60 MYA (4), leatherback and green turtles show extremely high genome
synteny and collinearity (Figs. 1b,c, S6, S7), with Progressive CACTUS revealing 95% sequence identity
across the length of the genomes (Table S5). After multiple rounds of manual curation to correct artifacts
of mis-assemblies, few large structural rearrangements between the two species remained, including
inversions of up to 7 Mb on chromosomes 12, 13, 24 and 28 (Fig. S6). The high collinearity between
species included near-complete end-to-end contiguous synteny for nine of 28 chromosomes (Fig. S6). The
remaining 19 chromosomes exhibited at least one small region of reduced collinearity (RRC) between the
species, with RRCs representing a total of ~83.4 Mb (~3.9%) and ~110.5 Mb (~5.2%) of the leatherback
and green turtle genome lengths, respectively. Eight chromosomes exhibited small RRCs (0.1–3 Mb), and
11 contained RRCs that were between 3–18 Mb in length (Figs. 2a-d & Table S6). Analyses of coding
regions revealed a similar pattern of strong collinearity between the two species (Figs. 1c, S6),

particularly within the macrochromosomes, which contain more than 80% of the total length of the
genomes. The two genomes also displayed similar percentages of repetitive elements (REs), which were
almost exclusively transposable elements (TEs) and unclassified repeats (Fig. S8). The landscape of TE
superfamily composition over evolutionary time was also similar between the two species, with the
exception of REs with low Kimura values (<5%) which appeared at higher frequency in the leatherback
turtle genome (see Supplementary Appendix 1 for full analyses).
180
181   *Gene families and gene functional analysis*
182   Gene function analysis of localized RRCs revealed that most contained genes with higher copy
numbers in the green turtle compared to the leatherback (Fig. 2a-d, Table S6). Nineteen chromosomes had
RRCs with higher gene copy numbers in the green turtle, and of these, ten contained genes associated
with immune system, olfactory reception and/or zinc-finger protein coding genes. Many of the same gene
families were also detected as high-diversity exonic regions *via* separate, independent analyses (see
Supplemental Appendix 1), reinforcing their importance in the divergent evolution of these species. In
addition to localized RRCs, higher gene copy numbers in the green turtle occurred in many gene
orthologous groups (orthogroups) across the entire genome, and generally in variable multicopy genes
(Fig. 2f, g). Copy number variation accounted for most of the nearly one thousand more genes annotated
in the green turtle genome relative to the leatherback (Fig. 2f, g; Table 1). We detected no evidence of
collapsed multicopy genes in the leatherback turtle assembly across multiple analyses (see Methods;
Table S7), supporting this as a biological signal rather than technical artifact of the assemblies.
194   Olfactory receptors (ORs) represented the largest orthogroups in both genomes, and differences
in copy numbers were connected to many of the identified RRCs. All OR Class I genes were clustered at
the beginning of chromosome 1, and the green turtle had higher copy numbers in this region (Fig. 2a-d).
This area also contained a cluster of OR Class I genes in at least three additional testudinid species (Fig.
S10), and is the only divergent region across the very large chromosome 1 in the turtles analyzed. In
contrast, OR Class II genes were spread across several chromosomes in both sea turtle species, with
higher copy numbers again in the green turtle found within RRCs (Fig. 2b-d). The instability and rapid
evolution of OR gene numbers in turtles is further illustrated in the expansion-contraction analysis of
orthogroups (Fig. 2e, Table S10a-d), which showed that OR Class I genes underwent a modest
contraction in the ancestral sea turtle lineage, followed by an expansion in the green turtle but a further
contraction in the leatherback turtle. Similar trends were detected for OR Class II genes, but with a
greater magnitude of contraction in the ancestral sea turtle lineage followed by a further contraction for
the leatherback turtle and only a small expansion for the green turtle (Fig. 2e).
207   Another important RRC (RRC14) encompassed the major histocompatibility complex (MHC),
which plays a critical role in vertebrate immunity and is particularly relevant to sea turtle conservation
due to the threat of FP and other diseases (32). In addition to the MHC region, this RRC includes several
copies of OR Class II genes, zinc-finger protein coding genes and other genes involved with immunity,
such as butyrophilin subfamily members and killer cell lectin-like receptors (Fig. 2d, Table S6).
Invariably, the green turtle carried higher numbers of all the multicopy genes present in RRC14. RRCs on
other chromosomes similarly showed increased levels of zinc-finger protein genes in the green turtle,
including the RRCs labeled 6A, 11A, 14A, and 28 (Table S6). In particular, zinc-finger protein genes
were highly prevalent on chromosomes 14 and 28 in both sea turtles, representing more than 50% of all
the protein domains present on these chromosomes (Fig. S11). Finally, all but three genes with known
roles in TSD in reptiles (Table S11) were located as single-copy genes within both sea turtle genomes,

with homologous copies located in the same region of the chromosomes in both species (see
Supplementary Appendix I for full analyses).
220
Macro and microchromosomes
Microchromosomes contained significantly higher proportions of genes than macrochromosomes
(Fig. 3a,b; green turtle: $F_{(2,25)}=16.46$, $p < 0.01$; leatherback turtle: $F_{(2,25)}=16.35$, $p < 0.01$), and gene
content was strongly positively correlated with GC content (Fig. S13; green turtle $R^2 = 0.81$, $p < 0.01$;
leatherback turtle $R^2 = 0.87$, $p < 0.01$). These patterns were particularly apparent in small (<20 Mb)
microchromosomes, where GC content reached 50%, compared to the 43-44% genome-wide averages.
Within chromosome groups, larger proportions of multicopy genes were generally associated with higher
total gene counts (green turtle: $R^2=0.84$, $p < 0.01$; leatherback turtle: $R^2=0.92$, $p < 0.01$), and
chromosomes with the highest multicopy genes numbers have increased proportions of RRCs (Fig. 3a,b;
green turtle: $R^2=0.69$, $p < 0.01$; leatherback turtle: $R^2=0.81$, $p < 0.01$).
Mean genetic distances for single-copy regions between the two sea turtles were also higher in
small microchromosomes (0.053) compared to both intermediate (>20 Mb) microchromosomes (0.047),
and macrochromosomes (0.045) (Fig. 3c; $F_{(2,25)}=21.98$, $p < 0.01$). However, examination of intermediate
microchromosome and macrochromosome RRCs revealed elevated genetic distances in these regions that
approached the values observed in small microchromosomes (Table S12). Genetic distances were also
significantly positively correlated with heterozygosity (green turtle: $R^2=0.97$, $p < 0.01$; leatherback turtle
$R^2=0.97$, $p < 0.01$), which was significantly higher in small microchromosomes for both species (Fig. 3d;
green turtle: $F_{(2,25)}=15.72$, $p < 0.01$; leatherback turtle: $F_{(2,25)}=5.09$, $p < 0.05$).
239
Genome diversity
Genome-wide nucleotide diversity was almost a magnitude of order lower in leatherback
compared to green turtles (mean repeat masked $\pi = 2.86\times10^{-4}$ and 2.46×10^{-3}, respectively; $t_{(5.52)} = 36.9$, p
< 0.001; Figs. 4a, S15-17, Table S14). Despite having largely similar gene content identified in the
annotation, this strong pattern was also observed in coding regions (Fig 4a.; $t_{(5.52)} = 37.7$, $p < 0.001$), such
that leatherback turtles possess much less standing functional variation, which may impact their adaptive
capacity to future novel environmental conditions. The strikingly low genomic diversity of leatherback
turtles is also less than almost all other reptile species examined (Fig S19; but see (51)), including
Chelonoidis abingdonii, where low diversity has been considered a contributing factor to their extinction
(52). In contrast, the genomic diversity of the green turtle fell in the mid-range for reptiles, as well as for
mammals examined using similar methods (53, 54). Finally, within both species, heterozygosity was
lower in coding regions (mean $\pi = 2.77\times10^{-4}$ and 2.18×10^{-3} for leatherback and green turtles; Fig. 4a)
relative to non-coding regions (mean $\pi = 3.18\times10^{-4}$ and 2.64×10^{-3}; leatherbacks: [$t_{(4)} = -8.9$, $p < 0.01$] and
greens: [$t_{(5)} = -30.9$, $p < 0.01$]), as expected from selection pressures driving higher sequence conservation
in these functional genomic regions.
255
Runs of homozygosity (ROH)
In addition to lower genome-wide heterozygosity, leatherbacks had a greater total number of
ROHs (>50 Kb) than green turtles (mean $N_{ROH} = 4,510$ and 829, respectively), as well as a greater total
aggregate length of the genome in ROH (range =26.1 – 45.5% in leatherback turtles; 1.8 – 17.7% in green
turtles). The mean length of ROHs was also significantly higher in leatherback ($L_{ROH} = 183.9$ Kb)
compared to green turtles ($L_{ROH} = 154.9$ Kb) ($t_{(7429.4)} = -8.85$, $p < 0.01$). Length distribution breakdown

showed that leatherbacks have a higher aggregate length of all categories of ROHs relative to the green turtles (Figs. 4b, S22). Short ROHs (50-500 Kb) had the highest total aggregate length in leatherbacks, with a mean aggregate length of 597 Mb (Fig. 4b), suggesting long-term low population sizes in the leatherback turtle.

Within species, overall ROH distributions were generally similar between samples representative of different populations for leatherback turtles, although individuals from the Northwest Atlantic and East Pacific populations displayed slightly higher total aggregate lengths of ROHs than those from the West Pacific population, primarily due to greater aggregate lengths of medium and long ROHs (Fig. 4b). Among green turtles, the aggregate length of ROHs in all categories were generally small and similar across individuals, with the clear exception of the genome reference sample that originated from the Mediterranean population. This individual displayed higher numbers and lengths of long ROHs (>1 Mb) compared to all other green turtles (n = 50 compared to <5, and aggregate length = 74 Mb compared to <4 Mb), suggesting higher levels of recent inbreeding relative to the other green turtle populations represented in our dataset. Comparative analyses mapping this individual to the two previous green turtle assemblies failed to detect these long ROHs (Fig. S23), demonstrating the importance of highly contiguous reference genomes for detecting biologically important patterns using this conservation-relevant metric.

Genetic load

Coding region variants with predicted high (e.g., stop-codon gain or loss) or moderate impacts were significantly more common in leatherback compared to green turtles (Fig. 4c; high impact variants: $t_{(4.18)}$ = -65.7, p < 0.001; moderate impact variants: $t_{(4.51)}$ = -29.5, p < 0.001). Conversely, low impact and modifier (i.e. variants predicted to cause negligible impacts) variants were significantly more common in green turtles (Fig. 4c; low impact variants: $t_{(5.88)}$ = 4.0, p < 0.01; modifier variants: $t_{(5.33)}$ = 31.8, p < 0.001). The missense to silent mutation ratio was also higher in leatherbacks than green turtles ($t_{(7.19)}$ = -72.3, p < 0.001; mean = 0.99 and 0.70), further suggesting that genetic load is higher in the leatherback turtles. Within species, there was limited variation between individuals for all variant categories (Fig. 4c).

Demographic history

Pairwise Sequential Markovian Coalescence (PSMC) analyses indicated different historical effective population sizes (N_e) between the two sea turtle species (Fig. 4d). N_e for all leatherback turtle populations represented in our dataset have been relatively small and sustained over time, ranging in size from approximately 2,000 to 21,000 over the last 10 million years, up until the Last Glacial Maximum (LGM) and at the lower end of this range for most of the last 5 million years. This pattern is consistent between all individuals examined, with similar timings and magnitudes of N_e fluctuations until recent history (Fig. 4d). In contrast, green turtles have experienced wider variation and a higher overall N_e in general, fluctuating between approximately 50,000 and 125,000, until the late Pleistocene, with estimates varying by population (Figs. 4d, S24). While N_e for leatherback turtles is relatively low, it modestly increased prior to the Eemian warm period (Fig. 4d [B]), followed by a subsequent decrease during this period until the LGM (Fig. 4d [A]) when all populations exhibit sharp spikes in N_e possibly due to inter-ocean gene flow following warming after the LGM. In contrast, green turtles generally displayed three distinct peaks in N_e (Fig. 4d), associated with ocean connectivity changes following the closure of the Tethys Sea [D], during the Pleistocene period [C], and prior to the Eemian warming period (Fig. 4d [B]).

305   While the patterns of N_e are broadly similar within green turtles, the timing and magnitude of these
306   fluctuations varied between populations (Fig. S24).
307
308   **Discussion**
309
Divergence in localized RRCs and microchromosomes amidst high global genome synteny. The
ancestral lineage leading to leatherback and green turtles diverged over 60 MYA (4), giving rise to
species that are adapted to dissimilar habitats, diets, and modes of life. Despite high overall levels of
genome synteny between the sea turtle families in both macro- and microchromosomes, RRCs and small
microchromosomes were associated with higher concentrations of multicopy gene families, as well as
heightened nucleotide diversity and genetic distances between species, suggesting that these genomic
elements may be important sources of variation underlying phenotypic differentiation. Higher
heterozygosity despite richer gene content in the microchromosomes suggests that these regions are prone
to variation accumulation and therefore may have high adaptation value. Though our results here do not
demonstrate direct causality, we have identified candidate regions and gene families that can be targeted
in further studies quantifying evidence for positive selection and their roles in sea turtle adaptation and
speciation.
322   The high global stability of macro- and microchromosomes between sea turtle families also aligns
with recent work showing similar patterns across reptiles, including birds, emphasizing the important
roles of microchromosomes in vertebrate evolution (55). Higher evolutionary rates for
microchromosomes relative to macrochromosomes has been documented in intraspecific (56) and
interspecific (57) studies with chicken and turkey genomes, respectively, so it is possible that the
characteristics of microchromosomes and RRCs we observed are not unique to sea turtles, but rather, are
prevalent in many vertebrates, which will become clearer as more high-quality assemblies are produced.
The mechanisms driving these patterns are not well-understood, but could be related to higher
recombination rates in micro- compared to macrochromosomes (58) that result in higher nucleotide
diversity and lower haplotype sharing. Once generated, balancing selection may also play a role in
maintaining variation in these gene dense regions, but more work is needed across taxa to determine the
broad support for these hypotheses. Our detailed analyses of RRCs, microchromosomes, and their
associated genes were only possible due to the high-quality of the assembled sea turtle genomes because
these analyses can be sensitive to genome fragmentation and mis-assemblies (39). For example, the RRCs
and many microchromosomes could not be detected using the draft green turtle genome due to
fragmentation and sequence gaps (Figs. S3-4). The prevalence of localized genomic differentiation and
underlying mechanisms among other closely or more distantly related vertebrate groups has yet to be
widely evaluated due to a lack of equivalent quality genomic resources, but this is rapidly changing. As
chromosomal-level genomes across all vertebrate lineages become available, our work provides a
roadmap for identifying genomic regions harboring contrasting expansion/contractions of gene families
and diversity levels. For taxa with highly conserved genomes like sea turtles, analyses of RRCs and
microchromosomes are likely important to understand their divergent evolutionary histories and the
phenotypic connections of the genes within them.
345
Contrasting sensory and immune gene evolution between sea turtle families. Sea turtles have complex
sensory systems and can detect both volatile and water-soluble odorants, which are imperative for
migration, reproduction and identification of prey, conspecifics, and predators (59–63). However,

leatherback and green turtles occupy dissimilar ecological niches and depend on different sensory cues.
While leatherback turtles almost exclusively inhabit the pelagic environment post-hatching, performing
large horizontal and vertical migrations to seek out patches of gelatinous prey (64), green turtles recruit to
neritic coastal and estuarine habitats as juveniles, and can have highly variable diets (65, 66). Sea turtle
nasal cavity morphology also differs between species, with leatherback turtle cavities relatively shorter,
wider, and more voluminous than chelonids (67–69), suggesting reduced requirements for olfactory
reception. OR genes encode proteins used to detect olfactory chemical cues, with the number of OR genes
strongly correlated with the number of detectable odorants (70), and linked to the chemical complexity of
the inhabited environment (71). The two major groups of ORs in amniote vertebrates are separated by
their affinities with hydrophilic molecules (Class I) or hydrophobic molecules (Class II) (72). Class I OR
genes may be particularly important in aquatic adaptation (32), and expansions of Class I ORs in
testudines, including green turtles, have been previously reported. However, the accuracy of these
estimates for complex gene families using short-read assemblies has been uncertain because they may be
prone to mis-assembly (32, 42, 73). We detected an additional 93 Class I OR genes in our green turtle
genome compared to those reported in the draft green turtle genome (42), suggesting they can be
erroneously collapsed in short-read assemblies. Our reconstruction of both Class I and Class II OR gene
evolution throughout the sea turtle lineage revealed that after ancestral contractions, gene copy evolution
diverged in opposite directions between the sea turtle families. The greater loss of Class II compared to
Class I OR genes in the ancestral sea turtle lineage likely reflects relaxed selection for detection of
airborne odorants, as has been observed in other lineages that recolonized marine environments, including
marine mammals (74). However, as sea turtles continue to use terrestrial habitats for reproduction, they
may need to retain some of these capabilities, which could explain why the observed contraction was
weaker than those in exclusively marine species (e.g., the vaquita *Phocaena sinus*; Fig. 2e).
The strong Class I OR expansion in the green turtle may be related to its distribution in complex
neritic habitats and variable diet, requiring detection of a high diversity of waterborne odorants, while the
continued loss of ORs in the leatherback turtle could be a consequence of its more specialized diet and the
lower chemosensory-complexity of pelagic habitats. Although leatherback turtles can detect chemical
cues from their prey, sensory experiments have indicated that visual cues are more important for food
recognition in this species (75, 76). Additionally, while the precise mechanisms underpinning philopatry
in sea turtles remain unclear, green turtles are thought to use olfactory cues to reach specific natal nesting
beaches following long-distance navigation guided by magnetoreception (61, 63). In contrast, leatherback
turtles exhibit more 'straying' from natal rookeries than other species, and such relaxed philopatry may be
related to reduced capability to detect olfactory cues to hone in on specific beaches.
Diversity within the highly-complex MHC region is a key component in the vertebrate immune
response to pathogens, with greater gene copy numbers and heterozygosity linked to lower disease
susceptibility (77). While both sea turtle species contained most of the core MHC-related genes, the green
turtle had more copies of genes involved in adaptive and innate immunity. Pathogen prevalence and
persistence is often greater in neritic habitats than open ocean habitats (78), so green turtles may be
exposed to higher pathogen loads and diversity than leatherback turtles (79). However, reptilian immune
systems are understudied compared to other vertebrates, and very few studies of MHC genes have been
conducted in turtles (80). Thus, it is not yet understood how immune gene diversity translates into disease
susceptibility or ecological adaptation in sea turtles, which is particularly critical for their conservation as
FP continues to threaten the recovery of populations around the globe (30). Although this likely viral-
mediated tumor disease occurs in all sea turtle species, disease prevalence and recovery greatly varies

between and within species, making it plausible that harboring certain genes, copy numbers, or specific
alleles may play important roles in disease dynamics. Despite decades of research on this disease (30)
only one study on the immunogenomic factors governing FP susceptibility or resilience has been
conducted (81), in part due to difficulty in accurately quantifying hypervariable and complex MHC loci
with short-read sequencing technologies (82). Our reference genomes now enable studies to accurately
interrogate MHC and other immune genes to close this critical research gap and advance our fundamental
understanding of immune gene evolution in testudines.
400
Differential genomic diversity and demographic histories. Genomic diversity is a critical metric for
evaluating extinction risk and adaptive potential to environmental perturbation (83–85), with
heterozygosity positively correlated with individual fitness (see reviews by (86, 87). Understanding the
causes and consequences of genomic diversity is imperative for sea turtles, and for leatherback turtles in
particular, where contemporary populations have experienced recent sharp declines due to human
activities (25). Leatherback turtles exhibited exceptionally low genomic diversity relative to the green
turtles and other reptiles and mammals, broadly aligning with previous estimates (88, 89). However,
factors influencing genomic diversity can vary among species (90), and our PSMC and ROH results
indicate that low diversity in the leatherback turtle is likely a consequence of long-term low effective
population sizes and historical bottleneck events, rather than a loss during recent population declines. This
is consistent with mitochondrial analyses suggesting that contemporary populations radiated from a small
number of matriarchal lineages within a single refugium following the Pleistocene (89). The low, yet
relatively evenly spread heterozygosity, is also congruent with sustained low population sizes similar to
that observed in several mammal species (91, 92). In contrast, the higher heterozygosity, limited ROHs
(though see discussion below), and estimated larger, more variable historical N_e in green turtles likely
reflects their radiation from many refugia and frequent admixing of populations (93).
Regardless of the causes of current genomic diversity levels in sea turtles, the amount of standing
variation can have important implications for species' future persistence (94), especially given the
adaptive capacity likely required to keep pace with rapid anthropogenic global change. Although
informative genome-wide diversity estimates can be made without high-quality reference genomes, these
enable deeper examination of diversity patterns that are relevant for conservation. For example, the use of
our reference genomes demonstrated that diversity is very low within coding regions of the leatherback
turtle genomes, indicating limited standing functional variation that may have implications for the
adaptive potential of this species to novel conditions. Additionally, leatherback turtles exhibited higher
genetic load compared to green turtles, and this signal was consistent across all samples regardless of
population source. Leatherback turtles have substantially lower hatching success compared to other sea
turtle species (29), potentially related to the heightened genetic load and low heterozygosity (95, 96), and
may combine with other factors to slow population recoveries despite conservation measures. However,
recent studies have documented low genome diversity in a number of species with wide geographic
distributions and relatively large census population sizes, including some long-lived marine vertebrates
(91, 97–100). Additionally, other species with low diversity have rebounded following population
declines and/or appear to have purged deleterious alleles through long-term low population sizes (98, 101,
102), thereby limiting the impacts on viability (54, 98, 103). Although our results of greater genetic load
despite long-term low N_e suggest this is not the scenario for leatherback turtles, further assessments
including more individuals over greater spatial and temporal scales are needed. Studies enabled by the
reference genomes presented here quantifying diversity and genetic load within and among global

populations will clarify these relationships for leatherback and other sea turtle species to guide
conservation recommendations.
While diversity and genetic load patterns were consistent within species, ROH analyses revealed
variation providing possible insight into different recent population histories, though these results must be
interpreted with caution given our limited sample sizes. Although all leatherback turtles displayed a high
number and aggregate length of short ROHs, consistent with a historical bottleneck generating long
ROHs that are subsequently broken by mutations and recombination events (104), individuals from the
West Pacific population show limited evidence of recent inbreeding (i.e., few long ROHs). In contrast,
individuals from the Northwest Atlantic and Eastern Pacific harbor higher aggregate lengths of medium
and long ROHs, suggesting more recent breeding of related individuals. These populations differ
substantially in their recent census sizes and trends (105) that are generally concordant with these
patterns; for example, the Western Pacific meta-population is relatively larger but declining, while some
Northwest Atlantic populations have undergone rapid increases and others remain small and isolated.
However, there is limited knowledge of abundances for these populations prior to the last several decades,
and the long generation times of sea turtles makes it likely that impacts of very recent demographic
changes may not yet be fully reflected in the genomes. Thus, as conservation efforts continue to mitigate
the ongoing major anthropogenic threats to the survival of this species, genomic monitoring over longer
temporal scales is needed to discern if populations are likely to encounter complications arising through
inbreeding depression during recovery. In green turtles, long ROHs were absent or in very low numbers
in all individuals, with the striking exception of the reference individual from the Mediterranean. This
isolated population that has undergone severe decline over the last century due to human exploitation
(106), and our results indicate that consequent inbreeding is likely occurring, which may impact its
recovery. The specific individual was from the Israel green turtle rookery that is estimated to have only
10-20 nesting females in the last decade (107, 108). However, it is currently unclear if Israel is
demographically isolated from other rookeries in the region (108, 109), so further research is needed to
understand if inbreeding is a concern only for this nesting aggregation, or the Mediterranean population
more broadly. Finally, these findings highlight the utility of ROH even in animals with long generation
times, and the importance of using highly contiguous genomes for accurate ROH assessment to inform
conservation.
The lower, long-term N_e of leatherback turtles detected in our demographic reconstructions may
be associated with this species' greater mass and trophic position, as was found in recent study assessing
relationships between key life-history traits and genomic variation in avian species (110). While it is
widely documented that environmental changes can strongly impact species' abundances and
distributions (111–113), following an initial decrease associated with declining temperatures, N_e of
leatherback turtles remained relatively constant throughout the substantial temperature fluctuations of the
Pleistocene. As ectotherms, reptiles are generally sensitive to climatic thermal fluctuations, however,
leatherback turtles exhibit unique physiological adaptations that produce regional endothermy and
facilitate exploitation of cold-water habitats (6) that potentially led them to being less susceptible to
periods of cooler temperatures. In contrast, wide fluctuations for green turtles appear correlated with
climatic events, beginning with the closure of the Tethys Sea, which altered ocean connectivity and
represented a period of increasing temperatures that may have opened more suitable habitat. As
temperatures subsequently decreased, N_e also decreased, however temperature fluctuations during the
Pleistocene were associated with an additional increases in N_e. While warmer temperatures presumably
allowed for larger population sizes of green turtles, large spikes in N_e around the Eemian warming,

particularly for the Mediterranean individual, are very likely associated with mixing of previously isolated
populations due to warm-water corridors allowing movement between populations and ocean basins
(114). While our overall estimates and trends for both species were broadly concordant with previous
studies (89, 115, 116), a recent study using Multiple Sequentially Markovian Coalescent (MSMC2)
analyses found steep declines in N_e for green turtles >100,000 years before present (116), which was not
detected in our PSMC analyses. Since this decline was also not detected in a prior study using PSMC on
the draft green turtle genome (115), and demographic inferences are generally robust to genome quality
(117, 118), this is likely a consequence of the different methods, with MSMC analyses inferring large N_e
for more ancient time scales (117).
490
Enabling future research and conservation applications. In addition to the insights reported here, the
reference genomes for both extant sea turtle families provide invaluable resources to enable a wide
breadth of previously unattainable fundamental and applied research. Combined with other forthcoming
chromosomal-level vertebrate genomes (39), in-depth comparative genomics analyses can further
investigate ecological adaptation related to immune and sensory gene evolution, as well as the genomic
basis for traits of interest such as adaptation to saltwater, diving capacity, and long-distance natal homing.
Studies leveraging these reference genomes alongside whole-genome sequencing of archival sample
collections can assess how genomic erosion, inbreeding and mutational load are linked to population size,
trajectories, and conservation measures in global sea turtle populations. For instance, the fact that
leatherback turtles have persisted with low diversity and N_e for extended periods offers hope for their
recovery, but given that some populations have now been reduced to only a few hundred individuals
(105), research quantifying purging of deleterious alleles, inbreeding depression, and adaptive capacity
within populations is urgently needed (119). We emphasize that high-quality reference genomes are not
required for all research goals, and combined with other recent studies (117, 118, 120), our findings
provide clear guidance on when they may, or may not, be necessary in order to generate accurate results
to inform conservation. For example, genome-wide diversity estimates are typically robust to assembly
quality, but the ability to detect long ROHs can be strongly affected. As ROH metrics are increasingly
being used to guide species management plans (121–123), it is important for researchers to understand
how genome quality may impact their analyses and inferences. Additionally, many conservation
applications that may not explicitly require whole-genome data can also directly benefit from the utility of
these reference genomes, including the development of amplicon panels and molecular assays to
investigate TSD mechanisms and adaptive capacity under climate change, and assessing linkages between
immune genes and disease risk. Finally, with global distributions and long-distance migratory
connectivity, sea turtle conservation requires international collaboration that has been previously
hampered by difficulty comparing datasets between laboratories. Existing anonymous markers (e.g.
microsatellites and restriction-site based SNP markers) can now be anchored to these genomes, and new
ones can be optimized for conservation-focused questions and shared across the global research
community, facilitating large-scale syntheses and equitable capacity building for genomics research.
While ongoing anthropogenic impacts continue to threaten the viability of sea turtles to persist, combined
with the important work of reducing major threats such as fisheries bycatch and habitat loss, these
genomes will enable research that make critical contributions to recovering imperiled populations.
522

Methods

Reference sample collections, genome assembly and annotation

523
524
Blood was collected from leatherback and green turtles using minimally invasive techniques for
isolation of ultra-high molecular weight DNA, and tissue samples of internal organs for RNA were
collected opportunistically from recently deceased or euthanized animals. Full details of sample
collection, storage, and laboratory processing prior to sequencing can be found in Supplementary
Appendix I. Resulting raw data were deposited into the VGP Genome Ark and NCBI Short-Read Archive
(SRA) (see Data Accessibility Statement). We assembled both genomes using four genomic technologies
following the VGP pipeline v1.6 (39) with a few modifications detailed in Supplementary Appendix I.
Briefly, PacBio Continuous Long Reads were assembled into haplotype phased contigs, with contigs
scaffolded into chromosome-level super scaffolds using a combination of 10X Genomics linked reads,
Bionano Genomics optical maps, and Arima Genomics Hi-C 3D chromosomal interaction linked reads.
Base call errors were corrected to achieve high quality (>Q40). The assemblies were manually curated,
with structural errors corrected according to the Hi-C maps (Fig. S1), and the 28 super scaffolds
(hereinafter referred to as chromosomes) numbered in both species according to sequence lengths in the
leatherback turtle assembly, and synteny between the two species. A manual inspection comparing the
sequence collinearity between the first curated versions of the genomes revealed a small number of
artefactual sequence rearrangements that were corrected in a second round of manual curation (see
Supplementary Appendix I).
To enable accurate, species-specific annotations for each genome, both short and long-read
transcriptome data (RNA-Seq and Iso-Seq) were generated from tissues known for their high transcript
diversity in each species. These data, plus homology-based mapping from other species, were used to
annotate the genomes using the standardized NCBI pipeline (124). We performed annotation as
previously described (39, 125), using the same RNA-Seq, Iso-Seq, and protein input evidence for the
prediction of genes in the leatherback and green turtles. We aligned 3.5 billion RNA-Seq reads from eight
green turtle tissues (blood, brain, gonads, heart, kidney, lung, spleen and thymus) and 427 million reads
from four leatherback turtle tissues (blood, brain, lung and ovary) to both genomes, in addition to 144,000
leatherback turtle and 1.9 million green turtle PacBio IsoSeq reads, and all Sauropsida and *Xenopus*
GenBank proteins, known RefSeq Sauropsida, *Xenopus*, and human RefSeq proteins, and RefSeq model
proteins for *Gopherus evgoodei* and *Mauremys reevesii.*
553
Genome quality analysis
We used the pipeline assembly-stats from https://github.com/sanger-pathogens/assembly-stats to
estimate scaffold N50, size distributions and assembly size. BUSCO analysis (115) and QV value
estimations (116) were conducted to assess the overall completion, duplication, and relative quality of the
assemblies. We used D-GENIES (118) with default parameters to conduct dot plot mapping of the entire
genomes and each individual chromosome to evaluate the synteny between leatherback and green turtle
genomes, and Haibao Tang JCVI utility libraries following the MCScan pipeline (119) to verify the
contiguity of the genomes. Incongruences in gene synteny blocks were manually investigated using
Artemis Comparative Tool (120), identifying possible regions of inversion that could be caused by
artifacts during assembly. These regions were then identified and corrected in the latest version of the
assembly for both species. Only a few structural rearrangements between the two species remained after
two rounds of manual curation with support of sequencing data. The final curated assemblies were

analyzed using the Genome Evaluation Pipeline (https://git.imp.fu-berlin.de/cmazzoni/GEP) to obtain all
final QC plots and summary statistics.
568
Identification and analysis of RRCs and REs
Leatherback and green turtle genomes were mapped to each other using Minimap2 with a dot plot
of the mapping generated using D-GENIES (126). Using windows of 20 Mb, the dot plot was screened
visually with regions larger than 1 Mb showing reduced collinearity (i.e., one or more breaks in the
diagonal indicating homology), as well as smaller regions with obvious signals of genomic
rearrangements (e.g., inversions), cataloged as regions of reduced collinearity (RRCs; Fig S5). Several
genomic features (e.g. GC content, repeat elements) were examined within these regions and compared to
regions of the same length directly up- and down-stream of the RRCs, which are most likely under
relatively similar cellular and molecular influences to the RRCs (Table S9). We identified the functions of
the genes present in RRCs using genome annotations and identified protein domains using Interproscan
(127). The proportion of GO terms in each chromosome was estimated for each species using PANTHER
(128); Fig. S25). To examine if RRCs presented differential patterns of sequence and/or gene duplication
between the species, we aligned the genomes of the sea turtles against each other using Progressive
Cactus (129, 130), and all homologous genes that presented more than one copy for one of the two
species were isolated using an inhouse script (*IdentifyDupsReciprocalBlast.sh*) to retrieve duplicated
genes (see Supplementary Appendix I for further details on Cactus alignments). Repetitive elements
(REs) were identified by creating a *de novo* database of transposable elements using RepeatModeller2
(131), followed by running RepeatMasker (132, 133) to calculate Kimura values for all REs (see full
analysis details in Supplementary Appendix I).
588
Gene families and gene functional analysis
To estimate the timing of gene family evolution for the OR gene families on sea turtles we used
Computational Analysis of gene Family Evolution v5 (134). Briefly, CAFE uses phylogenomics and gene
family sizes to identify gene family expansions and contractions. We used a dataset containing 8 species
of turtle, 4 non-turtle reptiles, 3 mammals and 1 amphibian using OrthoFinder (135, 136). OR
orthogroups were grouped based on subfamily (Class I and Class II; see (73)), and an ultrametric
phylogeny was generated by gathering 1:1 orthologs. We then aligned amino acid sequences for each
orthogroup and generated a phylogenetic tree (see Supplementary Appendix I for details).
To identify genes related to immunity, and the MHC in particular, we searched the genome for
the list of core MHC genes following Gemmell et al. (2020) (46). We conducted initial searches of gene
identifications, followed by a search of protein identifiers. As genes associated with the MHC are diverse,
and vary substantially among species, we did not use a BLAST search for these genes. Locations of the
genes were then compared between species to determine which genes were annotated, and where the core
MHC region is located within the genomes. We conducted a search following similar methods for genes
with known function in TSD in other reptiles (see Supplementary Appendix I for details).
604
Genetic distance, genome diversity, and runs of homozygosity
To estimate the genetic distance between the leatherback and green turtle genomes, we used the
halSnps pipeline (137) to compute interspecific single variants based on genome alignments obtained
with Progressive Cactus (129, 130) using the leatherback turtle genome as the reference. Genetic
distances were calculated for windows across the genome where each window included exactly 10,000

positions presenting single alignments against the green turtle genome in the Cactus output. Positions
with zero, or more than one alignment were ignored, and if this occurred over more than 50% of a given
window, it was skipped entirely (i.e., each window analyzed covered between 10 and 20 Kb of the
genome). Interspecific distances per bp were calculated by dividing the number of variants found within a
window by 10,000. Differences in genetic distance, gene content, GC content, and heterozygosity
between macro-, intermediate micro-, and small microchromosomes were tested using one-way ANOVAs
for each species. Regression analyses were used to test for correlations between these measures across
chromosomes.
For genome diversity, ROH, demographic history, and genetic load analyses, we also included
whole-genome resequencing (WGR) data for additional individuals representing multiple global
populations in each species (Table S13 and Supplemental Appendix I Methods for sample details). We
calculated genome-wide heterozygosity using a method adapted from Robinson et al. (2019) (92), which
used the Genome Analysis Toolkit (GATK) (138) to call genotypes at every site across the genome from
reads mapped to our reference genomes with BWA-mem (139). To avoid any biases arising from
differences in processing between samples, 10X linked-reads from the reference individuals were initially
processed using the proc10xG pipeline (https://github.com/ucdavis-bioinformatics/proc10xG), and then
treated identically to Illumina short-read data from resequenced individuals. Heterozygosity was
calculated within 100 Kb non-overlapping windows, with only sites that had a depth of between $\frac{1}{3}\times$ and
$2\times$ mean coverage retained for genotype scoring. Heterozygosity was calculated for (1) the entire
genome, (2) the genome with repeat-regions masked, (3) exonic regions, (4) and for the non-coding
regions. Statistical comparisons between species were made using T-tests, with paired T-tests used when
comparing between regions within species. We subsequently applied the heterozygosity pipeline to
generate genome-wide heterozygosity for a number of additional reptilian species with sequences sourced
from the NCBI SRA where species-specific reference genomes were available (see details in
Supplementary Appendix I).
ROHs were identified by initially generating a SNP-list using the Analysis of Next Generation
Sequencing Data (ANGSD; (140) pipeline. ANGSD was parameterized to output files that were
configured for use as input for the ROH analysis incorporated in PLINK (141). ROHs were then further
characterized as 'short' (50-500 Kb), 'medium' (500Kb-1 Mb), or 'long' (>1 Mb) , with size class
categories loosely based on (104).
640
Genetic load
Estimates of deleterious allele accumulation were conducted using the snpEff variant annotation
software (142). We estimated the impacts of variants (SNPs and INDELs) from coding regions using the
species-specific genome annotations generated for both species, with a total of 18,775 genes for the
leatherback turtle genome, and 19,752 genes for the green turtle genome used in the analysis. gVCFs
were generated for each individual followed by joint-genotyping using GATK (138), allowing the
reference individuals to include homozygous alleles found in other individuals. Combined VCFs were
then separated for each individual and filtered using based on depth of coverage ($\frac{1}{3}\times$ - $2\times$ mean coverage
for each individual). The snpEff program predicts variant impacts and bins them into 'high', 'moderate',
or 'low' impact categories, and outputs a list of genes that have predicted variant effects. We ran the
snpEff analysis on all individuals for both species, and compared the percentages of each variant between
species using T-tests.
653

Historical demography
Pairwise Sequential Markovian Coalescence (PSMC; (143)) analyses of demographic history
were employed for all individuals for both species. We used SAMtools (144) and BCFtools (145) to call
genotypes with base and mapping quality filters of >Q30, before filtering for insert size (50-5,000bp) and
allele balance (AB) and retaining only biallelic sites with an AB of <0.25 and >0.75. We then ran PSMC
analysis using the first 10 scaffolds, which constituted over 84% of the total length of the genome. We
scaled our outputs using a generation time of 30 years (mid-way between reported generation times for
both species; see Supplementary Appendix I), and a mutation rate of 1.2×10^{-8} (115).
662

Acknowledgments
We are grateful for the assistance with the (1) leatherback turtle sample collection from the St. Croix Sea
Turtle Program and the US Fish and Wildlife Service, the NOAA-SWFSC California in-water leatherback
research team, and the New England Aquarium; (2) green turtle sample collection from the Israel
National Sea Turtle Rescue Centre, the NOAA PIFSC-MTBAP team, and Thierry Work (USGS). We
also thank Estefany Argueta and Jamie Adkins Stoll for assistance with literature searches for TSD and
immune genes and library preparations for leatherback resequenced individuals, and Phillip Morin,
Andrew Foote, Anna Brüniche-Olsen, Annabel Beichman, Morgan McCarthy, David L. Adelson, and
Yuanyuan Cheng for their invaluable discussions surrounding analysis approaches and comments on the
manuscript. Sequencing of the green turtle has been performed by the Long Read Team of the
DRESDEN-concept Genome Center, DFG NGS Competence Center, part of the Center for Molecular and
Cellular Bioengineering (CMCB), Technische Universität Dresden and MPI-CBG. For green turtle
resequenced samples, we also thank Jessica Farrell, Whitney Crowder, Brooke Burkhalter, Nancy
Condron, and the veterinary and rehabilitation staff and volunteers of the University of Florida's (UF) Sea
Turtle Hospital at Whitney Laboratories, and the staff of the South Carolina Aquarium for sampling
support. Thanks also are due to UF's Interdisciplinary Center for Biotechnology Research Core for
sequencing services, and to the Florida Fish and Wildlife Conservation Commission and South Carolina
Department of Natural Resources for valuable assistance with permitting. Leatherback DNA for
additional WGR came from samples archived in NOAA-SWFSC Marine Mammal and Sea Turtle
Research Collection that were originally collected as part of previous population genetics studies, and we
thank Erin LaCasella for her help processing samples from the SWFSC.
684

Funding
Funding was provided by the University of Massachusetts Amherst, NSF-IOS (grant #1904439 to LMK),
NOAA-Fisheries, National Research Council postdoctoral fellowship program (LMK), Vertebrate
Genomes Project, Rockefeller University, to EDJ, HHMI to EDJ, the Sanger Institute, Max-Planck-
Gesellschaft, as well as grant contributions from Tom Gilbert, Paul Flicek, Robert Murphy, Karen A.
Bjorndal, Alan B. Bolten, Ed Braun, Neil Gemmell, Tomas Marques-Bonet, and Alan Scott. We also
acknowledge CONICYT-DAAD for scholarship support to TCV, and EKSR was supported by São Paulo
Research Foundation - FAPESP (grant #2020/10372-6). BeGenDiv is partially funded by the German
Federal Ministry of Education and Research (BMbF, Förderkennzeichen 033W034A). The work of FT-N
and PM was supported by the Intramural Research Program of the National Library of Medicine, National
Institutes of Health. The work of MP was partially funded through the Federal Ministry of Education and
Research (grant 01IS18026C). HP was supported by a Formació de Personal Investigador fellowship
from Generalitat de Catalunya (FI_B100131). MK was supported by "la Caixa" Foundation (ID

100010434), fellowship code LCF/BQ/PR19/11700002 and the Vienna Science and Technology Fund
(WWTF) and the City of Vienna through project VRG20-001. Funding for the green turtle resequencing
was provided by a Welsh Government Sêr Cymru II and the European Union's Horizon 2020 research
and innovation programme under the Marie Skłodowska-Curie grant agreement No. 663830-BU115 and
the Sea Turtle Conservancy, Florida Sea Turtle Grants Program under project number 17-033R.
703

Data Accessibility Statement
Assemblies for both species have been deposited on NCBI GenBank. The NCBI GenBank accession
numbers for the leatherback turtle genome assembly (rDerCor1) are GCF_009764565.3 and
GCA_009762595.2 for the annotated primary and original alternate haplotypes in BioProject
PRJNA561993, and for the green turtle assembly (rCheMyd1) are GCF_015237465.2 and
GCA_015220195.2 for primary and alternate haplotypes respectively in BioProject PRJNA561941. The
raw data used for assemblies are available on the Vertebrate Genome Ark
(https://vgp.github.io/genomeark/). The leatherback turtle RNA-Seq data generated for the purpose of
assembly annotation was deposited in the SRA under accession numbers SRX8787564-SRX8787566
(RNA-Seq) and SRX6360706-SRX6360708 (ISO-Seq). Green turtle RNA-Seq data generated for
annotation were deposited in SRA under accessions SRX10863130-SRX10863133 (RNA-Seq) and as
SRX11164043-SRX11164046 (ISO-Seq). The NovaSeq 6000 DNA-Seq data for the green turtle
resequencing, including raw reads, are deposited in NCBI (https://www.ncbi.nlm.nih.gov/) under
BioProject ID: PRJNA449022. All scripts used for downstream analyses following genome assembly and
annotation have been deposited on GitHub under repository
https://github.com/bpbentley/sea_turtle_genomes.

Tables and Figures

Table 1 | Quality statistics for the genome assemblies and annotations for leatherback (*Dermochelys coriacea*) and green (*Chelonia mydas*) turtles.

	Leatherback turtle (*Dermochelys coriacea*)	Green turtle (*Chelonia mydas*)
Genome ID	rDerCor1	rCheMyd1
Assembly accession	GCA_009764565.3	GCA_015237465.1
Assembly level	Chromosome	Chromosome
Total genome length (bp)	2,164,762,090	2,134,358,617
Contig N50 (bp)	7,029,801	39,415,510
Scaffold N50 (bp)	137,568,771	134,428,053
Number of scaffolds	40	92
Number of chromosomes	28	28
Quality Value (QV)	38.9	47.7
Annotated protein-coding genes	18,775	19,752

BUSCO Assembly and Annotation Completeness Statistics (based on Vertebrate core BUSCOs) and Annotation BUSCO scores

BUSCO category	Assembly	Annotation	Assembly	Annotation
Complete genes	91.6%	97.2	94.2%	97.9%
Complete + fragmented	95.4%	97.7	96.7%	98.2%
Missing	4.10%	1.3%	2.8%	0.7%
Duplicated	0.5%	1.0%	0.5%	1.1%

Fig. 1 | **(a)** Photographs of green turtle (*Chelonia mydas*)*;* photo credit: NOAA NMFS PIFSC under USFWS Permit #TE-72088A-3, and leatherback turtle (*Dermochelys coriacea*); photo credit: Ricardo Tapilatu. **(b)** Dot plot showing regions with an identity greater than 0.5 across the entire genomes of green (red) and leatherback (blue) turtles. **(c)** Gene synteny and collinearity per chromosome among five species of turtles: leatherback turtle (blue), green turtle (red), Chinese pond turtle (*Mauremys reevesii*; green), pond slider turtle (*Trachemys scripta*; purple) and Goode's thornscrub tortoise (*Gopherus evgoodei*; yellow). Each bar represents chromosomes with respective numbers and gray lines represent homolog gene connections among species.

Fig. 2 | (a-d) Dotplots (identity values as color; dark green=1-0.75, green=0.75-0.5, orange=0.5-0.25 and yellow=0.25-0) depicting four of the regions with reduced collinearity (RRC) identified within chromosomes and associated with higher copy numbers of immune system (IS), olfactory receptor (ORs), or zinc finger domain genes (ZFD) in the green turtle (*Chelonia mydas*) relative to leatherback (*Dermochelys coriacea*) turtle (see also Fig. S3, Tables S3 & S5 for full details of all RRCs). Positions of each RRC are marked with gray squares on the dot plots (left; with *D. coriacea* on the X-axes and *C. mydas* on the Y-axes) and gene collinearity maps (right) for each chromosome highlighting the connections among specific gene families in different colors. **(e)** Gene family evolution of olfactory receptors Class I (red) and Class II (black) for amniote phylogeny. Gene numbers are presented on the nodes and gain/loss along each branch are presented below branches. Small scale bar represents substitutions/site and big scale bar represents divergence times (MA). The blue dashed line shows the estimated divergence between the two sea turtle families. **(f)** Number of unique and shared orthogroups and single and multi-copy genes between the two sea turtles (coding genes including genes with rearrangement). The boxes outlined in black denote shared orthogroups, with the higher multi-copy in the green turtle due to greater gene copies within orthogroups. **(g)** Comparison of gene counts between both species per multigenic orthogroup, depicting only those orthogroups where both species have different numbers of genes and a minimum number of five genes for one of the species. Bubbles above the diagonal represent higher counts for the green turtle and below for the leatherback turtle. The size of the bubbles represents the number of orthogroups with the same gene count combination.

Fig. 3 | Number of genes, genetic distance between species and heterozygosity within species in macrochromosomes, small (<20 Mb) and intermediate (>20 Mb) microchromosomes. **(a)** Relation between the number of genes, percentage of reduced collinearity regions (RRCs), and classified TE per chromosome for the green (*Chelonia mydas*) and **(b)** leatherback (*Dermochelys coriacea*) turtles. Dark colors indicate the total number of genes and light colors indicate the number of multicopy genes. **(c)** Average genetic distance between green and leatherback turtles per chromosome. **(d)** Relation between genetic distance and heterozygosity per chromosome for each species.

Fig. 4 | Data is presented for the leatherback (*Dermochelys coriacea*; blue) and green (*Chelonia mydas,* red) turtle genomes, including reference individuals for both species (*), and the individual used to generate the draft genome (⊤ Wang et al. 2013). **(a)** estimates of heterozygosity across the genome calculated with 100 Kb non-overlapping windows for the entire genome, repeat-masked genome, exons and non-exon regions, with outliers removed. **(b)** accumulated lengths of runs of homozygosity (ROH). **(c)** predicted impacts of variants from within coding regions. **(d)** Pairwise sequential Markovian coalescent plot (PSMC) of demographic history of both species using a mutation rate of 1.2×10^{-8} and generation time of 30 years, overlayed with temperature. Letters indicating portions of the PSMC curves (A-D) are geological events referred to in the main text and Supplementary Appendix I.

References

1. R. Hirayama, Oldest known sea turtle. *Nature* **392**, 705–708 (1998).

2. H. B. Shaffer, E. McCartney-Melstad, T. J. Near, G. G. Mount, P. Q. Spinks, Phylogenomic analyses
of 539 highly informative loci dates a fully resolved time tree for the major clades of living turtles
(Testudines). *Mol. Phylogenet. Evol.* **115**, 7–15 (2017).

3. D. A. Pike, Climate influences the global distribution of sea turtle nesting: Sea turtle nesting
distributions. *Glob. Ecol. Biogeogr.* **22**, 555–566 (2013).

4. R. C. Thomson, P. Q. Spinks, H. B. Shaffer, A global phylogeny of turtles reveals a burst of climate-
associated diversification on continental margins. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2012215118
(2021).

5. J. Davenport, Temperature and the life-history strategies of sea turtles. *J. Therm. Biol.* **22**, 479–488
(1997).

6. W. Frair, R. G. Ackman, N. Mrosovsky, Body Temperature of *Dermochelys coriacea*: Warm Turtle
from Cold Water. *Science* **177**, 791–793 (1972).

7. P. C. H. Pritchard, "Evolution, Phylogeny, and Current Status" in *The Biology of Sea Turtles*, P. L.
Lutz, J. A. Musick, Eds. (CRC Press, 1996), pp. 1–28.

8. L. McClenachan, J. B. C. Jackson, M. J. H. Newman, Conservation implications of historic sea turtle
nesting beach loss. *Front. Ecol. Environ.* **4**, 290–296 (2006).

9. J. B. Jackson, *et al.*, Historical overfishing and the recent collapse of coastal ecosystems. *Science*
293, 629–637 (2001).

10. M. A. Grassman, D. W. Owens, J. P. McVey, R. M. M, Olfactory-based orientation in artificially
imprinted sea turtles. *Science* **224**, 83–84 (1984).

11. K. J. Lohmann, C. M. F. Lohmann, There and back again: natal homing by magnetic navigation in
sea turtles and salmon. *J. Exp. Biol.* **222**, jeb184077 (2019).

12. P. S. Tomillo, V. S. Saba, R. Piedra, F. V. Paladino, J. R. Spotila, Effects of illegal harvest of eggs
on the population decline of leatherback turtles in Las Baulas Marine National Park, Costa Rica.
Conserv. Biol. **22**, 1216–1224 (2008).

13. S. Fossette, *et al.*, Pan-atlantic analysis of the overlap of a highly migratory species, the leatherback
turtle, with pelagic longline fisheries. *Proc. Biol. Sci.* **281**, 20133065 (2014).

14. Y. Kaska, *et al.*, Natural and anthropogenic factors affecting the nest-site selection of Loggerhead
Turtles, *Caretta caretta*, on Dalaman-Sarıgerme beach in South-west Turkey: (Reptilia:
Cheloniidae). *Zool. Middle East* **50**, 47–58 (2010).

15. B. Von Holle, *et al.*, Effects of future sea level rise on coastal habitat. *Journal of Wildlife*
Management **83**, 694–704 (2019).

16. N. Mrosovsky, G. D. Ryan, M. C. James, Leatherback turtles: the menace of plastic. *Mar. Pollut.*
Bull. **58**, 287–289 (2009).

798 17. M. Chaloupka, G. H. Balazs, T. M. Work, Rise and fall over 26 years of a marine epizootic in
799    Hawaiian green sea turtles. *J. Wildl. Dis.* **45**, 1138–1142 (2009).

800 18. L. A. Hawkes, A. C. Broderick, M. H. Godfrey, B. J. Godley, Climate change and marine turtles.
801    *Endanger. Species Res.* **7**, 137–154 (2009).

802 19. B. P. Wallace, *et al.*, Global conservation priorities for marine turtles. *PLoS One* **6**, e24510 (2011).

803 20. C. L. Yntema, N. Mrosovsky, Incubation temperature and sex ratio in hatchling loggerhead turtles: a
804    preliminary report. *Mar. Turtle Newsl.* **11**, 9–10 (1979).

805 21. M. P. Jensen, *et al.*, Environmental Warming and Feminization of One of the Largest Sea Turtle
806    Populations in the World. *Curr. Biol.* **28**, 154–159.e4 (2018).

807 22. A. D. Mazaris, G. Schofield, C. Gkazinou, V. Almpanidou, G. C. Hays, Global sea turtle
808    conservation successes. *Sci Adv* **3**, e1600730 (2017).

809 23. IUCN, The IUCN Red List of Threatened Species (2021) (April 16, 2021).

810 24. L. S. Martínez, *et al.*, Conservation and Biology of the Leatherback Turtle in the Mexican Pacific.
811    *Chelonian Conserv. Biol.* **6**, 70–78 (2007).

812 25. P. Santidrián Tomillo, *et al.*, Reassessment of the leatherback turtle (*Dermochelys coriacea*) nesting
813    population at Parque Nacional Marino Las Baulas, Costa Rica: effects of conservation efforts.
814    *Chelonian Conserv. Biol.* **6**, 54–62 (2007).

815 26. J. R. Spotila, R. D. Reina, A. C. Steyermark, P. T. Plotkin, F. V. Paladino, Pacific leatherback turtles
816    face extinction. *Nature* **405**, 529–530 (2000).

817 27. Laúd OPO Network, Enhanced, coordinated conservation efforts required to avoid extinction of
818    critically endangered Eastern Pacific leatherback turtles. *Sci. Rep.* **10**, 4772 (2020).

819 28. E. Chan, H. Liew, Decline of the leatherback population in Terengganu, Malaysia, 1956-1995.
820    *Chelonian Conserv. Biol.* **2**, 196–203 (1996).

821 29. K. L. Eckert, B. P. Wallace, J. G. Frazier, S. A. Eckert, P. C. H. Pritchard, "Synopsis of the
822    biological data on the leatherback sea turtle, *Dermochelys coriacea*" (U.S. Department of Interior,
823    Fish and Wildlife Service, 2012).

824 30. K. Jones, E. Ariel, G. Burgess, M. Read, A review of fibropapillomatosis in Green turtles *(Chelonia
825    mydas)*. *Vet. J.* **212**, 48–57 (2016).

826 31. G. Zhang, *et al.*, Comparative genomics reveals insights into avian genome evolution and adaptation.
827    *Science* **346**, 1311–1320 (2014).

828 32. I. Khan, *et al.*, Olfactory Receptor Subgenomes Linked with Broad Ecological Adaptations in
829    Sauropsida. *Mol. Biol. Evol.* **32**, 2832–2843 (2015).

830 33. D. Jebb, *et al.*, Six reference-quality genomes reveal evolution of bat adaptations. *Nature* **583**, 578–
831    584 (2020).

832 34. B. J. McMahon, E. C. Teeling, J. Höglund, How and why should we implement genomics into
833    conservation? *Evolutionary Applications* **7**, 999–1007 (2014).

35. M. A. Supple, B. Shapiro, Conservation of biodiversity in the genomics era. *Genome Biol.* **19**, 131
(2018).

36. P. Brandies, E. Peel, C. J. Hogg, K. Belov, The Value of Reference Genomes in the Conservation of
Threatened Species. *Genes* **10**, 864 (2019).

37. P. A. Hohenlohe, W. C. Funk, O. P. Rajora, Population genomics for wildlife conservation and
management. *Mol. Ecol.* (2020) https:/doi.org/10.1111/mec.15720.

38. X. Zhang, J. Goodsell, R. B. Norgren Jr, Limitations of the rhesus macaque draft genome assembly
and annotation. *BMC Genomics* **13**, 206 (2012).

39. A. Rhie, *et al.*, Towards complete and error-free genome assemblies of all vertebrate species. *Nature*
592, 737–746 (2021).

40. A. P. Fuentes-Pardo, D. E. Ruzzante, Whole-genome sequencing approaches for conservation
biology: Advantages, limitations and practical recommendations. *Mol. Ecol.* **26**, 5369–5406 (2017).

41. G. Formenti, *et al.*, The era of reference genomes in conservation genomics. *Trends Ecol. Evol.* **37**,
197–202 (2022).

42. Z. Wang, *et al.*, The draft genomes of soft-shell turtle and green sea turtle yield insights into the
development and evolution of the turtle-specific body plan. *Nat. Genet.* **45**, 701–706 (2013).

43. A. Whibley, J. L. Kelley, S. R. Narum, The changing face of genome assemblies: Guidance on
achieving high-quality reference genomes. *Mol. Ecol. Resour.* **21**, 641–652 (2021).

44. V. Peona, *et al.*, Identifying the causes and consequences of assembly gaps using a multiplatform
genome assembly of a bird-of-paradise. *Mol. Ecol. Resour.* **21**, 263–286 (2021).

45. Y. Yuan, *et al.*, Comparative genomics provides insights into the aquatic adaptations of mammals.
Proc. Natl. Acad. Sci. U. S. A. **118** (2021).

46. N. J. Gemmell, *et al.*, The tuatara genome reveals ancient features of amniote evolution. *Nature* **584**,
403–409 (2020).

47. R. N. Johnson, *et al.*, Adaptation and conservation insights from the koala genome. *Nat. Genet.* **50**,
1102–1111 (2018).

48. J.-N. Hubert, T. Zerjal, F. Hospital, Cancer- and behavior-related genes are targeted by selection in
the Tasmanian devil (*Sarcophilus harrisii*). *PLoS One* **13**, e0201838 (2018).

49. L. M. Zimmerman, The reptilian perspective on vertebrate immunity: 10 years of progress. *J. Exp.*
Biol. **223**, jeb214171 (2020).

50. M. Seppey, M. Manni, E. M. Zdobnov, BUSCO: Assessing Genome Assembly and Annotation
Completeness. *Methods Mol. Biol.* **1962**, 227–245 (2019).

51. Q.-H. Wan, *et al.*, Genome analysis and signature discovery for diving and sensory properties of the
endangered Chinese alligator. *Cell Res.* **23**, 1091–1105 (2013).

52. V. Quesada, *et al.*, Giant tortoise genomes provide insights into longevity and age-related disease.
Nat Ecol Evol **3**, 87–95 (2019).

53. P. A. Morin, *et al.*, Reference genome and demographic history of the most endangered marine
mammal, the vaquita. *Mol. Ecol. Resour.* **21**, 1008–1020 (2021).

54. J. A. Robinson, *et al.*, Genomic Flatlining in the Endangered Island Fox. *Curr. Biol.* **26**, 1183–1189
(2016).

55. P. D. Waters, *et al.*, Microchromosomes are building blocks of bird, reptile, and mammal
chromosomes. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

56. H.-J. Megens, *et al.*, Comparison of linkage disequilibrium and haplotype diversity on macro- and
microchromosomes in chicken. *BMC Genet.* **10**, 86 (2009).

57. E. Axelsson, M. T. Webster, N. G. C. Smith, D. W. Burt, H. Ellegren, Comparison of the chicken
and turkey genomes reveals a higher rate of nucleotide divergence on microchromosomes than
macrochromosomes. *Genome Res.* **15**, 120–125 (2005).

58. A. V. Rodionov, Micro vs. macro: structural-functional organization of avian micro- and
macrochromosomes. *Genetika* **32**, 597–608 (1996).

59. C. S. Endres, K. J. Lohmann, Detection of coastal mud odors by loggerhead sea turtles: a possible
mechanism for sensing nearby land. *Mar. Biol.* **160**, 2951–2956 (2013).

60. C. S. Endres, N. F. Putman, K. J. Lohmann, Perception of airborne odors by loggerhead sea turtles.
J. Exp. Biol. **212**, 3823–3827 (2009).

61. M. Manton, A. Karr, D. W. Ehrenfeld, Chemoreception in the migratory sea turtle, *Chelonia mydas*.
Biol. Bull. **143**, 184–195 (1972).

62. C. Kitayama, *et al.*, Behavioral effects of scents from male mature Rathke glands on juvenile green
sea turtles (*Chelonia mydas*). *J. Vet. Med. Sci.* **82**, 1312–1315 (2020).

63. C. S. Endres, *et al.*, Multi-Modal Homing in Sea Turtles: Modeling Dual Use of Geomagnetic and
Chemical Cues in Island-Finding. *Front. Behav. Neurosci.* **10**, 19 (2016).

64. K. L. Dodge, J. M. Logan, M. E. Lutcavage, Foraging ecology of leatherback sea turtles in the
Western North Atlantic determined through multi-tissue stable isotope analyses. *Mar. Biol.* **158**,
2813–2824 (2011).

65. K. E. Arthur, M. C. Boyle, C. J. Limpus, Ontogenetic changes in diet and habitat use in green sea
turtle (*Chelonia mydas*) life history. *Mar. Ecol. Prog. Ser.* **362**, 303–311 (2008).

66. J. A. Seminoff, *et al.*, Large-scale patterns of green turtle trophic ecology in the eastern Pacific
Ocean. *Ecosphere* **12**, e03479 (2021).

67. C. Kitayama, *et al.*, Morphological features of the nasal cavities of hawksbill, olive ridley, and black
sea turtles: Comparative studies with green, loggerhead and leatherback sea turtles. *PLoS One* **16**,
e0250873 (2021).

68. D. Kondoh, C. Kitayama, Y. K. Kawai, The nasal cavity in sea turtles: adaptation to olfaction and
seawater flow. *Cell Tissue Res.* **383**, 347–352 (2021).

69. Y. Yamaguchi, *et al.*, Computed tomographic analysis of internal structures within the nasal cavities
of green, loggerhead and leatherback sea turtles. *Anat. Rec.* **304**, 584–590 (2021).

70.   Y. Niimura, M. Nei, Evolutionary dynamics of olfactory and other chemosensory receptor genes in
vertebrates. *J. Hum. Genet.* **51**, 505–517 (2006).

71.   L. R. Yohe, M. Fabbri, M. Hanson, B.-A. S. Bhullar, Olfactory receptor gene evolution is unusually
rapid across Tetrapoda and outpaces chemosensory phenotypic change. *Curr. Zool.* **66**, 505–514
(2020).

72.   H. Saito, Q. Chi, H. Zhuang, H. Matsunami, J. D. Mainland, Odor coding by a Mammalian receptor
repertoire. *Sci. Signal.* **2**, ra9 (2009).

73.   M. W. Vandewege, *et al.*, Contrasting Patterns of Evolutionary Diversification in the Olfactory
Repertoires of Reptile and Bird Genomes. *Genome Biol. Evol.* **8**, 470–480 (2016).

74.   A. Liu, *et al.*, Convergent degeneration of olfactory receptor gene repertoires in marine mammals.
BMC Genomics **20**, 977 (2019).

75.   M. A. Constantino, M. Salmon, Role of chemical and visual cues in food recognition by leatherback
posthatchlings (*Dermochelys coriacea* L). *Zoology* **106**, 173–181 (2003).

76.   N. Warraich, J. Wyneken, N. Blume, Feeding behavior and visual field differences in loggerhead and
leatherback sea turtles may explain differences in longline fisheries interactions. *Endanger. Species*
Res. **41**, 67–77 (2020).

77.   H. V. Siddle, J. Marzec, Y. Cheng, M. Jones, K. Belov, MHC gene copy number variation in
Tasmanian devils: implications for the spread of a contagious cancer. *Proc. Biol. Sci.* **277**, 2001–
2006 (2010).

78.   L. E. Escobar, *et al.*, A global map of suitability for coastal Vibrio cholerae under current and future
climate conditions. *Acta Trop.* **149**, 202–211 (2015).

79.   L. Zhang, *et al.*, Massive expansion and functional divergence of innate immune genes in a
protostome. *Sci. Rep.* **5**, 8693 (2015).

80.   J. P. Elbers, S. S. Taylor, Others, Major histocompatibility complex polymorphism in reptile
conservation. *Herpetol. Conserv. Biol.* **11**, 1–12 (2016).

81.   K. R. Martin, K. L. Mansfield, A. E. Savage, Adaptive evolution of major histocompatibility
complex class I immune genes and disease associations in coastal juvenile sea turtles. *R Soc Open*
Sci **9**, 211190 (2022).

82.   X. Vekemans, *et al.*, Whole-genome sequencing and genome regions of special interest: Lessons
from major histocompatibility complex, sex determination, and plant self-incompatibility. *Mol. Ecol.*
30, 6072–6086 (2021).

83.   C. Moritz, Strategies to protect biological diversity and the evolutionary processes that sustain it.
Syst. Biol. **51**, 238–254 (2002).

84.   P. Fernandez-Fournier, J. M. M. Lewthwaite, A. Ø. Mooers, Do We Need to Identify Adaptive
Genetic Variation When Prioritizing Populations for Conservation? *Conserv. Genet.* **22**, 205–216
(2021).

85.   Y. J. Borrell, *et al.*, Heterozygosity-fitness correlations in the gilthead sea bream *Sparus aurata*
using microsatellite loci from unknown and gene-rich genomic locations. *J. Fish Biol.* **79**, 1111–

1129 (2011).

86. J. A. DeWoody, A. M. Harder, S. Mathur, J. R. Willoughby, The long-standing significance of
genetic diversity in conservation. *Mol. Ecol.* **30**, 4147–4154 (2021).

87. Y. Willi, *et al.*, Conservation genetics as a management tool: The five best-supported paradigms to
assist the management of threatened species. *Proc. Natl. Acad. Sci. U. S. A.* **119** (2022).

88. L. M. Komoroske, *et al.*, A versatile Rapture (RAD-Capture) platform for genotyping marine turtles.
Mol. Ecol. Resour. **19**, 497–511 (2019).

89. P. H. Dutton, B. W. Bowen, D. W. Owens, A. Barragan, S. K. Davis, Global phylogeography of the
leatherback turtle (*Dermochelys coriacea*). *J. Zool.* **248**, 397–409 (1999).

90. J. Romiguier, *et al.*, Comparative population genomics in animals uncovers the determinants of
genetic diversity. *Nature* **515**, 261–263 (2014).

91. M. V. Westbury, B. Petersen, E. Garde, M. P. Heide-Jørgensen, E. D. Lorenzen, Narwhal Genome
Reveals Long-Term Low Genetic Diversity despite Current Large Abundance Size. *iScience* **15**,
592–599 (2019).

92. J. A. Robinson, *et al.*, Genomic signatures of extensive inbreeding in Isle Royale wolves, a
population on the threshold of extinction. *Sci Adv* **5**, eaau0757 (2019).

93. M. P. Jensen, *et al.*, The evolutionary history and global phylogeography of the green turtle
(*Chelonia mydas*). *J. Biogeogr.* **46**, 860–870 (2019).

94. M. Kardos, *et al.*, The crucial role of genome-wide genetic variation in conservation. *Proc. Natl.*
Acad. Sci. U. S. A. **118**, e2104642118 (2021).

95. P. Dobrynin, *et al.*, Genomic legacy of the African cheetah, *Acinonyx jubatus*. *Genome Biol.* **16**, 277
(2015).

96. A. L. K. Mattila, *et al.*, High genetic load in an old isolated butterfly population. *Proc. Natl. Acad.*
Sci. U. S. A. **109**, E2496–505 (2012).

97. M. V. Westbury, *et al.*, Extended and Continuous Decline in Effective Population Size Results in
Low Genomic Diversity in the World's Rarest Hyena Species, the Brown Hyena. *Molecular Biology*
and Evolution **35**, 1225–1237 (2018).

98. J. A. Robinson, C. Brown, B. Y. Kim, K. E. Lohmueller, R. K. Wayne, Purging of Strongly
Deleterious Mutations Explains Long-Term Persistence and Absence of Inbreeding Depression in
Island Foxes. *Curr. Biol.* **28**, 3487–3494.e4 (2018).

99. Y. Xue, *et al.*, Mountain gorilla genomes reveal the impact of long-term population decline and
inbreeding. *Science* **348**, 242–245 (2015).

100. A. D. Foote, *et al.*, Killer whale genomes reveal a complex history of recurrent admixture and
vicariance. *Mol. Ecol.* **28**, 3427–3444 (2019).

101. N. Dussex, *et al.*, Population genomics of the critically endangered kākāpō. *Cell Genomics* **1**,
100002 (2021).

102. C. C. Kyriazis, R. K. Wayne, K. E. Lohmueller, Strongly deleterious mutations are a primary
determinant of extinction risk due to inbreeding depression. *Evol Lett* **5**, 33–47 (2021).

103. Y. D. Dewoody, J. A. Dewoody, On the estimation of genome-wide heterozygosity using molecular
markers. *J. Hered.* **96**, 85–88 (2005).

104. F. C. Ceballos, P. K. Joshi, D. W. Clark, M. Ramsay, J. F. Wilson, Runs of homozygosity: windows
into population history and trait architecture. *Nat. Rev. Genet.* **19**, 220–234 (2018).

105. United States National Marine Fisheries Service, U.S. Fish and Wildlife Service, "Endangered
Species Act status review of the leatherback turtle (*Dermochelys coriacea*)" (United States National
Marine Fisheries Service, 2020).

106. P. Casale, *et al.*, Mediterranean sea turtles: current knowledge and priorities for conservation and
research. *Endanger. Species Res.* **36**, 229–267 (2018).

107. Y. Tikochinski, *et al.*, Mitochondrial DNA STR analysis as a tool for studying the green sea turtle
(*Chelonia mydas*) populations: the Mediterranean Sea case study. *Mar. Genomics* **6**, 17–24 (2012).

108. Y. Tikochinski, *et al.*, Mitochondrial DNA short tandem repeats unveil hidden population structuring
and migration routes of an endangered marine turtle. *Aquat. Conserv.* (2018)
https:/doi.org/10.1002/aqc.2908.

109. S. Karaman, *et al.*, Population genetic diversity of green turtles, *Chelonia mydas*, in the
Mediterranean revisited. *Mar. Biol.* **169** (2022).

110. A. Brüniche-Olsen, K. F. Kellner, J. L. Belant, J. A. DeWoody, Life-history traits and habitat
availability shape genomic diversity in birds: implications for conservation. *Proc. Biol. Sci.* **288**,
20211441 (2021).

111. G. M. Hewitt, Genetic consequences of climatic oscillations in the Quaternary. *Philos. Trans. R. Soc.*
Lond. B Biol. Sci. **359**, 183–95; discussion 195 (2004).

112. F. V. Seersholm, *et al.*, Rapid range shifts and megafaunal extinctions associated with late
Pleistocene climate change. *Nat. Commun.* **11**, 2770 (2020).

113. I.-C. Chen, J. K. Hill, R. Ohlemüller, D. B. Roy, C. D. Thomas, Rapid range shifts of species
associated with high levels of climate warming. *Science* **333**, 1024–1026 (2011).

114. J. P. van der Zee, *et al.*, The population genomic structure of green turtles (*Chelonia mydas*) suggests
a warm-water corridor for tropical marine fauna between the Atlantic and Indian oceans during the
last interglacial. *Heredity* **127**, 510–521 (2021).

115. R. R. Fitak, S. Johnsen, Green sea turtle (*Chelonia mydas*) population history indicates important
demographic changes near the mid-Pleistocene transition. *Mar. Biol.* **165**, 110 (2018).

116. S. T. Vilaça, *et al.*, Divergence and hybridization in sea turtles: Inferences from genome data show
evidence of ancient gene flow between species. *Mol. Ecol.* (2021) https:/doi.org/10.1111/mec.16113.

117. A. H. Patton, *et al.*, Contemporary Demographic Reconstruction Methods Are Robust to Genome
Assembly Quality: A Case Study in Tasmanian Devils. *Mol. Biol. Evol.* **36**, 2906–2921 (2019).

118. K. Nadachowska-Brzyska, R. Burri, L. Smeds, H. Ellegren, PSMC analysis of effective population

sizes in molecular ecology and its application to black-and-white Ficedula flycatchers. *Mol. Ecol.* **25**,
1058–1072 (2016).

119. A. Khan, *et al.*, Genomic evidence for inbreeding depression and purging of deleterious genetic
variation in Indian tigers. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

120. A. Prasad, E. D. Lorenzen, M. V. Westbury, Evaluating the role of reference-genome phylogenetic
distance on evolutionary inference. *Mol. Ecol. Resour.* **22**, 45–55 (2022).

121. A. Brüniche-Olsen, K. F. Kellner, C. J. Anderson, J. A. DeWoody, Runs of homozygosity have
utility in mammalian conservation and evolutionary studies. *Conserv. Genet.* **19**, 1295–1307 (2018).

122. A. D. Foote, *et al.*, Runs of homozygosity in killer whale genomes provide a global record of
demographic histories. *Mol. Ecol.* (2021) https:/doi.org/10.1111/mec.16137.

123. J. A. Robinson, *et al.*, The critically endangered vaquita is not doomed to extinction by inbreeding
depression. *Science* **376**, 635–639 (2022).

124. N. A. O'Leary, *et al.*, Reference sequence (RefSeq) database at NCBI: current status, taxonomic
expansion, and functional annotation. *Nucleic Acids Res.* **44**, D733–45 (2016).

125. K. D. Pruitt, *et al.*, RefSeq: an update on mammalian reference sequences. *Nucleic Acids Res.* **42**,
D756–63 (2014).

126. F. Cabanettes, C. Klopp, D-GENIES: dot plot large genomes in an interactive, efficient and simple
way. *PeerJ* **6**, e4958 (2018).

127. M. Blum, *et al.*, The InterPro protein families and domains database: 20 years on. *Nucleic Acids Res.*
49, D344–D354 (2021).

128. H. Mi, *et al.*, PANTHER version 16: a revised family classification, tree-based classification tool,
enhancer regions and extensive API. *Nucleic Acids Res.* **49**, D394–D403 (2021).

129. B. Paten, *et al.*, Cactus graphs for genome comparisons. *J. Comput. Biol.* **18**, 469–481 (2011).

130. J. Armstrong, *et al.*, Progressive Cactus is a multiple-genome aligner for the thousand-genome era.
Nature **587**, 246–251 (2020).

131. J. M. Flynn, *et al.*, RepeatModeler2: automated genomic discovery of transposable element families.
Proceedings of the National Academy of Sciences **117**, 9451–9457 (2020).

132. M. Tarailo-Graovac, N. Chen, Using RepeatMasker to identify repetitive elements in genomic
sequences. *Curr. Protoc. Bioinformatics* **Chapter 4**, Unit 4.10 (2009).

133. A. Smit, R. Hubley, P. Green, RepeatMasker Open-4.0. 2013-2015http. *repeatmasker. org* (2015).

134. F. K. Mendes, D. Vanderpool, B. Fulton, M. W. Hahn, CAFE 5 models variation in evolutionary
rates among gene families. *Bioinformatics* (2020) https:/doi.org/10.1093/bioinformatics/btaa1022.

135. D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157 (2015).

136. D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative genomics.

Genome Biol. **20**, 238 (2019).

137. G. Hickey, B. Paten, D. Earl, D. Zerbino, D. Haussler, HAL: a hierarchical format for storing and
analyzing multiple genome alignments. *Bioinformatics* **29**, 1341–1342 (2013).

138. A. McKenna, *et al.*, The Genome Analysis Toolkit: A MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

139. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-
bio.GN]* (2013).

140. T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next Generation Sequencing
Data. *BMC Bioinformatics* **15**, 356 (2014).

141. S. Purcell, *et al.*, PLINK: a tool set for whole-genome association and population-based linkage
analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

142. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-
3. *Fly* **6**, 80–92 (2012).

143. H. Li, R. Durbin, Inference of human population history from individual whole-genome sequences.
Nature **475**, 493–496 (2011).

144. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
(2009).

145. H. Li, A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987–2993
(2011).

(a) Chromosome 1

(c) Chromosome 13

(b) Chromosome 6

(d) Chromosome 14

zinc-finger gene   olfactory receptor I   olfactory receptor II   MHC I   MHC II   other MHC genes   immune gene (not MHC)

(e)

Olfactory Receptors Class I Class II

(f)

(g)