# Cell surface receptor kinase FERONIA linked to nutrient sensor TORC signaling controls root hair growth at low temperature linked to low nitrate in *Arabidopsis thaliana*

Javier Martínez Pacheco[1,*], Limei Song[2,3,*], Lenka Kubĕnová[4], Miroslav Ovečka[4], Victoria Berdion Gabarain[1], Juan Manuel Peralta[1], Tomás Urzúa Lehuedé[5,6], Miguel Angel Ibeas[5,6], Martiniano M. Ricardi[7], Sirui Zhu[2], Yanan Shen[2], Mikhail Schepetilnikov[8], Lyubov A Ryabova[8], José M. Alvarez[5,10], Rodrigo A. Gutierrez[9,10], Guido Grossman[11], Jozef Šamaj[3], Feng Yu[2,†] & José M. Estevez[1,5,6,10,†]

[1]Fundación Instituto Leloir and IIBBA-CONICET. Av. Patricias Argentinas 435, Buenos Aires C1405BWE, Argentina.

[2]State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Biology, and Hunan Key Laboratory of Plant Functional Genomics and Developmental Regulation, Hunan University, Changsha 410082, P.R. China.

[3]Laborarory of Species Interaction and Biological Invasion, School of Life Science, Hebei University, Baoding 071002, P.R. China

[4]Department of Biotechnology, Faculty of Science, Palacký University Olomouc, Šlechtitelů 27, 783 71 Olomouc, Czech Republic.

[5]Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad Andres Bello.

[6]ANID - Millennium Nucleus for the Development of Super Adaptable Plants (MN-SAP), Santiago, Chile.

[7]Instituto de Fisiología, Biología Molecular y Neurociencias (IFIBYNE-UBA-CONICET) and Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Buenos Aires, Argentina.

[8]Institut de Biologie Moléculaire des Plantes, CNRS, UPR 2357, Université de Strasbourg, Strasbourg, France.

[9]Centre FONDAP Center for Genome Regulation, Santiago, Chile; Departamento de Genética Molecular y Microbiología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile.

[10]ANID - Millennium Institute for Integrative Biology (iBio), Santiago, Chile.

[11]Institute of Cell and Interaction Biology, Heinrich-Heine-University Düsseldorf, Germany; Cluster of Excellence in Plant Sciences, Heinrich-Heine-University Düsseldorf, Germany.

*Co-first authors.

†Correspondence should be addressed. Email: feng_yu@hnu.edu.cn (F.Y) and jestevez@leloir.org.ar (J.M.E)

**Abstract**

Root hairs (RH) are excellent model systems for studying cell size and polarity since they elongate
several hundred-fold their original size. Their tip growth is determined both by intrinsic and
environmental signals. Although nutrient availability and temperature are key factors for a
sustained plant growth, the molecular mechanisms underlying their sensing and downstream
signaling pathways remain unclear. Here, we identified that low temperature (10°C) triggers a
strong RH elongation response involving the cell surface receptor kinase FERONIA (FER) and the
nutrient sensing TOR Complex 1 (TORC). In this study, we found that FER is required to perceive
limited nutrient availability caused by low temperature. FER interacts with and activates TORC-
downstream components to trigger RH growth. In addition, the small GTPase Rho-related protein
from plants 2 (ROP2) is also involved in this RH growth response linking FER and TOR. We also
found that limited nitrogen nutrient availability can mimic the RH growth response at 10°C in a
NRT1.1-dependent manner. These results uncover a molecular mechanism by which a central
hub composed by FER-ROP2-TORC is involved in the control of RH elongation under low
temperature and nitrogen deficiency.

Abstract Word counts 186

Text Word counts 3,885

Figures 1-7

Passwords: Arabidopsis, cell surface, FERONIA, nitrogen, low temperature, root hairs, ROP2, TOR

kinase.

**Introduction**

Root hairs (RH) are cell outgrowths that develop as cylindrical protrusions from the root epidermis in a developmentally regulated manner[1]. RHs are able to expand in a polar manner several hundred times their original size in a couple of hours to reach water-soluble nutrients in the soil, to promote interactions with the local microbiome, and to support the anchoring of the plant[2]. RH growth is controlled by the coordination of a plethora of environmental and endogenous factors[3,4]. Recently, an autocrine mechanism of RH growth was described where RALF1-FER complex recruits and phosphorylates the early translation initiation factor 4E1 (eIF4E1) to enhance protein synthesis of specific mRNAs, including the RH growth master regulator *ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4)*[5]. The observation that RH cells can respond to their local environment in a cell-autonomous manner within minutes[6], points towards mechanisms that directly modulate the growth machinery at the growing apex.

It is known that macronutrient availability is a key factor that promotes rapid RH growth[3,7,8] Recently, we showed that under low temperature conditions (10°C), nutrient availability in the media is reduced and triggers an enhanced RH growth that is suppressed if nutrients are increased[9]. We specifically showed that RH growth of WT Col-0 plants is highly responsive to increasing nutrient concentrations (from 0.5X Murashige and Skoog (MS) to 2.0X MS) with regular concentration gelling agents (0.8%). High nutrient concentration impairs RH growth even if they are exposed to low temperatures. In a similar manner, an increase in agar concentration (from 0.8% to 2.5%) in the MS medium, which likely restrains nutrient mobility and nutrient uptake[10–13] blocked low temperature-induced RH elongation[9]. These observations suggested that low temperature restrict nutrient mobility and availability in the culture medium, leading to promotion of polar RH growth. It is still unclear which specific nutrients are affected and the signaling pathways involved in the temperature effect that triggers this RH growth response. Diminished nutrient availability is known to activate RH expansion through a transcriptional reprogramming governed by the transcription factors (TF) RHD6-RSL4[9]. Specifically, a novel ribonucleoprotein complex composed of lncRNA AUXIN-REGULATED PROMOTER LOOP (APOLO) and the TF WRKY42 forms a regulatory hub to activate RHD6 by shaping its epigenetic environment and integrate low temperature signals governing RH growth[9,14]. In addition, we recently identified new molecular components involved in low temperature RH growth. We found that cell wall-apoplastic related peroxidases, PRX62 and PRX69, are important for low temperature triggered RH growth by inducing changes in ROS homeostasis and cell wall EXTENSIN insolubilization[15]. Although relevant advances have been achieved in our understanding on how RH growth occurs at low temperature, it is still unclear how nutrient availability caused by low temperature is perceived in the RH cells and the extent of molecular responses that control RH growth.

TARGET OF RAPAMYCIN (TOR) is an evolutionarily conserved Ser/Thr protein kinase in all
eukaryotic organisms that acts as a central growth regulator controlling metabolism and protein
synthesis[16–18]. TOR is found in at least two distinct multiprotein complexes called TOR Complex 1
and 2 (TORC1 and TORC2) in animal cells, although only the equivalent for TORC1 has been
experimentally validated in plants[19,20]. The *Arabidopsis* TOR complex is encoded by one *TOR* gene
(*AtTOR*)[21], two *REGULATORY-ASSOCIATED PROTEIN OF TOR (RAPTOR 1A and 1B)* genes[22–27], and
two *LETHAL WITH SEC THIRTEEN 8 (LST8)* genes[28]. In contrast to the embryo-lethal *tor*-null
mutant lines, *raptor1b* and *lst8-1* mutants are viable but show significant growth defects and
developmental phenotypes[29]. Some canonical downstream targets of TOR are conserved in
plants, such as the 40S ribosomal S6 kinase (S6K) which stimulates protein synthesis[24,30–33]. In
plants, TOR complex plays a key role during many stages of the plant life cycle by controlling both
anabolic and catabolic downstream processes. In addition, TOR is activated by nutrient
availability and inactivated by stresses that alter cellular homeostasis[18,34–38]. The TOR complex
senses and integrates signals from the environment to coordinate developmental and metabolic
processes including hormones (e.g., auxin), several nutrients (e.g., nitrogen and sulfur), amino
acids and glucose[16,27,39–43]. However, the underlying molecular mechanism by which TOR
operates at a single plant cell level has yet to be elucidated.

To date, few upstream regulators of TOR have been described in plants. Among them, there is a
subfamily of small GTPase Rho-related protein from plants (ROPs) involved in the spatial control
of cellular processes by signaling to the cytoskeleton and vesicular trafficking. Particularly, ROP2
activates TOR in response to auxin and nitrogen signals[43–45]. ROP2 is described as a monomeric
GTP-binding protein that participates in many cellular signaling processes, including the polar
growth of pollen tubes and root hairs[46–49]. ROP activation is regulated by ROP guanine nucleotide
exchange factors (ROP-GEFs) which interact with several receptor-like kinases (RLKs) including
the *Cr*RLK family member FERONIA (FER)[47]. During the elongation of RH, ROP2 is activated by the
ROPGEF1-FER interaction. This process is also regulated by ROP-GEF4 and ROP-GEF10[47,50]. FER
was also linked to carbon/nitrogen balance during plant growth[51]. Even more, it was shown that
the cytoplasmic kinase domain of FER and its partner RIPK (RPM1-induced protein kinase)
interact with TOR and RAPTOR-1B forming a complex to positively modulate the TOR pathway
under low nitrogen nutrient conditions[52]. Currently, it is clear that plant growth can be regulated
via the FER-TOR pathway as it links amino acid and/or nutrients signaling with true leaves
development[51]. This evidence suggests that nutrient-mediated sensing at the cell surface level
triggers a response using the TOR signaling pathway. However, it is still unknown which are the
environmental signals activated in RH by the FER-TOR pathway in RH. Considering that low
temperature stress can result in enhanced RH growth[9,14], this stress condition was used as a
proxy to investigate environmental signals that activate RH cell elongation. Our research reveals a

novel mechanism in which FER, ROP2 and TOR are necessary to regulate RH growth in
*Arabidopsis* in the context of the nitrogen nutrient availability as caused by low temperature
stress.

**Results**

**FER is required to trigger a strong RH growth response at low temperature.** We asked which
might be the RH surface protein involved in perceiving or transducing the low temperature
stimulus. Since CrRLK1L FER was shown as an important hub between the cell surface signaling
and downstream processes during RH growth[5,47], we tested whether *fer-4* null mutant or *fer-5* (a
truncated version with shortened kinase domain (KD)) can respond to low temperature stimulus
(**Figure S1A**). Both *fer* mutants failed to trigger RH growth at low temperature (**Figure 1A**). We
considered plant genotype (Geno), treatment (Temp), and the interaction between genotype and
treatment (Geno:Temp) as the factors for two-way analysis of variance (ANOVA) to define
significant changes in RH growth in response to cold. We determined that the genotype factor
has a substantial and significant effect on RH growth in *fer-4* and *fer-5* mutants. Notably, the
Geno:Temp interaction term is also significant for *fer-4* and *fer-5*. These results suggest that FER
contributes to general RH growth and RH response to cold (**Figure S1B**). In contrast, the mutant
*eru*, impaired in the related CrRLK1L ERULUS (ERU), previously linked to RH growth and cell wall
integrity processes with very short RH phenotype at 22°C[53,54], was able to react to low
temperature which triggered RH growth response although to a lower extent than wild-type Col-0
plants (**Figure 1A**). This indicates that FER, but not ERU, might be involved specifically in this
growth response despite being phylogenetically related[54]. Then, a fluorescent translational
reporter line of FER (*p*FER:FER-GFP), was used to study *FER* expression in roots and RH. FER
protein levels are increased clearly activated in root cells under low temperature stimulus (**Figure
1B**), and during RH growth (**Figure 1C**). Distribution of FER-GFP in the clear zone and punctate
compartments in the cytoplasm was not affected by the temperature change from 22°C to 10°C.
However, accumulation of FER-GFP at the plasma membrane was increased at low temperature
(at 10°C) (**Figure 1C**). Finally, phosphorylated, and non-phosphorylated levels of FER (FER-p/FER)
were quantified at both temperatures. Under low temperature, almost half of the FER protein
levels were present as a non-phosphorylated form and after 3 days of growth, almost all FER
protein was present as a phosphorylated version (**Figure 1D**). This indicates that low temperature
not only increases FER protein levels at the plasma membrane of growing RH, but also promotes
a complete phosphorylation of FER protein. This is possibly enhancing the interaction with
putative partners and triggering the activation of downstream signaling components of RH polar
growth.

**TOR pathway is involved in low temperature induced RH growth.** Recently, it was shown that FER-RIPK interacts with TOR-RAPTOR1B and phosphorylates it leading to TOR pathway activation in the context of nutrient perception and regulation of global metabolism[52]. We then asked whether TOR might be also involved in the RH growth process under low temperature. We used a β-estradiol (es)-induced TOR knockdown (*tor-es,* es-induced RNAi silencing of TOR) line[55] which showed short RH as reported previously[32]. Notably, induction of TOR silencing with estradiol completely blocked the RH response to low temperature (**Figure 2A**). Similarly, inhibition of TOR kinase activity with AZD-8055[56], an ATP-competitive inhibitor of mTOR kinase activity, abolished the RH growth at both 22°C and low temperature (**Figure 2B**). Overexpression of TOR enhanced RH growth at both temperatures assayed (**Figure 2C**). Using two-way ANOVA analysis, we found that alteration of TOR levels impacts general RH growth (Geno factor, **Figure S2A**). However, the significant interaction between genotype and treatment indicates TOR is also involved in RH responses to cold (Geno:Temp, **Figure S2A**). These results suggest that the TOR pathway is involved in the RH growth and specifically in RH low temperature growth response and might operate downstream of FER. At the transcriptional level, TOR root expression level is upregulated up to three-folds at 10°C vs 22°C (**Figure S2B**). We tested whether the growth response at low temperature in RH is also affected in mutants of the TOR pathway and some downstream components. All the plant mutants tested *(raptor1b, rps6b,* and *lst8-1)* were unable to respond to the low temperature treatment. They exhibited a broad spectrum of RH phenotypes as compared to Wt Col-0 plants (**Figure 2C**). For instance, *raptor1b* behaved similarly to *tor-es* mutant, while *rps6b* showed an intermediate phenotype and *lst8* was comparable to Wt Col-0 plants (at 22°C) (**Figure 2C**). In addition, overexpression of TOR and the S6 KINASE 1 (S6K1), a direct downstream target of TOR, showed enhanced RH growth at 22°C. Since previous reports showed phosphorylation of the S6K1 can be used to monitor TOR protein kinase activity in plants[32,55], we measured ratios of S6K-p/S6K. Strong induction of the S6K-p/S6K ratio was observed in Wt Col-0 under low temperature (**Figure 2D**). These results suggest TOR and some downstream components (e.g. S6K and RPS6b) are required to promote RH growth under low temperature.

**FER facilitates TOR polar localization in RH and activation of TOR pathway at low temperature.** Since TOR activation under a plethora of stimuli usually triggers phosphorylation of the downstream factor S6K (S6K-p), we asked whether low temperature responses regulated by FER might control this output. We investigated the molecular mechanisms by which TOR-S6K are induced at 10°C and we tested whether this might be mediated by FER. Since *fer-4* and *tor-es* showed similar RH phenotypes at low temperature (**Figure 2A-B**), possibly indicating that they may act in the same pathway, we first tested whether FER and TOR interaction is enhanced under 10°C vs 22°C. By performing a co-immunoprecipitation (Co-IP) analysis, we found that FER-TOR interaction is enhanced under low temperature (**Figure 3A**). This result suggests an interaction between FER and TOR kinase is implicated in RH growth under low temperature. This

corroborates previous findings that showed direct interaction of the FER kinase domain and the N-terminal domain of TOR (aa 1-1449 of TOR, NTOR)[52]. Next, we evaluated TOR levels in growing RH by immunolocalization. We found that FER is required for the apical accumulation of TOR since TOR lost the polar pattern in *fer-4* root hairs. However, the level of RH tip-localized TOR detected was similar at both, low temperature, and control conditions (22°C) (**Figure 3B-C and Figure S3**). This result indicates that localization of TOR at RH tip is dependent on FER. Then, we tested whether FER is involved in S6K activation by TOR. We measured S6K-p/S6K protein ratio under both growth conditions in Wt Col-0, *fer-4*, *fer-5*, and FER[K565R] (which has been shown to abolish FER autophosphorylation and transphosphorylation in an *in vitro* kinase activity assay[57,58], although some activity might remain[59]). Interestingly, low temperature enhanced S6K-p levels after 3 days of growth in Wt Col-0 this was partially suppressed in all three *fer* mutants tested (**Figure 3D**). This suggests that FER controls the level of TOR activation under low temperature in RH. Collectively, these results indicate that TOR localization in the RH tip and TOR activation is dependent on FER and enhanced at low temperature.

**ROP2 is required for low temperature RH growth.** Previous studies have shown a key role of ROP2 in the regulation of RH polarity and elongation[48,60,61], as an important molecular link between FER and downstream components involved in RH growth[47]. In addition, ROP2 promotes the activation of TOR and its relocation to the cell periphery and induces the downstream signal transduction pathway[44]. More importantly, ROP2 was shown to integrate diverse nitrogen and hormone signals for TOR activation[43]. We asked whether low temperature (10°C) is able to change ROP2 targeting (as ROP2p:ROP2-mCitrine) to the plasma membrane in roots and RH. Low temperature (10°C) led to noticeable increase of ROP2 fluorescence intensity in root cells (**Figure 4A**), in the apical and subapical cytoplasm, and at the apical plasma membrane of RH (**Figure 4B**). Then, we observed that *rop2* and *rop2 rop4* partially abolishes the differential growth responses at 10°C and produces a very short RH phenotype (**Figure 4C**). On the contrary, ROP2 OE triggers a constitutive RH growth at both temperatures (**Figure S4**) similarly to *S6K1 OE* and *TOR OE* lines (**Figure 2C**). In addition, when ROP2 OE is expressed in the *fer* mutant background *fer-8* (ROP2 OE/*fer-8*; **Figure S4**), this strong growth effect disappears highlighting the role of FER on the ROP2 function. On the other hand, while high levels of TOR enhance RH growth regardless of the temperature treatments, the constitutively active ROP2 version (CA-ROP2) is able to abolish TOR activation of RH growth (**Figure 4C**). The levels of S6K-p/S6K in TOR OE, with constitutive long RH at both temperatures, was reduced by the presence of CA-ROP2 (in TOR OE/CA-ROP2), which correlates with a diminished RH growth (**Figure 4D**). Taken together, we suggest a direct function of ROP2 (and possibly ROP4) at the RH tip. ROP2 function is dependent on the presence of FER and it modulates TOR activity in the RH growth responses to low temperature.

**Nitrate perception and transport mediated by NRT1.1 controls RH growth at low temperature.**
Nitrate is a key nitrogen nutrient required for plant growth and development[62–64]. The CHLORINA1/NITRATE TRANSPORTER1.1 (CHL1/NRT1.1) is the only known nitrate receptor (transporter and receptor) [65–67] as well as for auxin transport[68]. NRT1.1 belong to the low affinity nitrate transporter family. However, when NRT1.1 is phosphorylated at threonine 101, it behaves as a high-affinity $NO_3^-$ (nitrate) transporter, giving this protein a dual-affinity capability[69–72]. Previous studies have shown that nitrogen and specifically nitrate is important for TOR signaling[43,73]. Since low temperature growth conditions reduce nutrient availability in the media and trigger a strong response in RH growth[9,14], we hypothesized that the nitrate signaling pathway was involved. We directly test this hypothesis, and we measured using a qualitative method the temperature effect of $NO_3^-$ diffusion in agar plates at 22°C and at 10°C (**Figure S5**). As expected, the $NO_3^-$ diffusion at 22°C was twice as high as at 10°C, confirming our previous results that the temperature effect affects the amount of nitrates available for the roots and RHs[9,14]. In agreement, we found that high levels of nitrate (18.8 and 37.6 mM) supplied in the M407 media without nitrogen (see methods) were able to partially repress low temperature-mediated RH growth while low levels of nitrate (0.5 mM) did not affect growth under this temperature (**Figure 5A**). As a reference, previous experiments were carried out with 0.5X MS media containing 9.3 mM of nitrate and several other compounds (see **Table S1**). In addition, an increase in agar concentration in the 407 media (from 0.8% to 1.5%) even with high $NO_3^-$ (18.8 mM), which likely restrains $NO_3^-$ diffusion mobility induced RH expansion independently of the temperature (**Figure S5**). Three-way ANOVA analysis across all conditions (**Figure S6A and S6B**) reveals that temperature (T), the interaction of agar with $NO_3^-$ (agar:N); and the interaction of agar, temperature, and $NO_3^-$ (agar:T:N) largely explain the variance of RH elongation; while $NO_3^-$ and agar factors on their own are less influential (**Figure S5C**). Altogether, these observations suggest that low temperatures restrict $NO_3^-$ diffusion and availability in the culture medium, leading to the promotion of polar RH growth. This result indicated nitrate can impact RH cell elongation at low temperature, particularly under low nutrient mobility environment. Next, we tested whether NRT1.1 was involved in this low temperature RH growth response. The NRT1.1 null mutants *chl1-5, chl1.9* (NRT1.1 harbors a substitution P492L that suppresses its root nitrate uptake activity[8]), as well as the *CHL1^T101D* (which mimics phosphorylated version of NRT1.1) showed strong RH growth response regardless of temperature conditions (**Figure 5B**), similarly to *S6K1 OE* and *TOR OE* lines (**Figure 2C**), but to a lower extent in terms of RH elongation. Only the *CHL1^T101A* (dephosphorylated version of NRT1.1) was able to slightly respond to the change in temperature. As expected, similar levels of S6K-p/S6K were detected in the *chl1-9* line at both temperatures (**Figure 5C**). When these NRT1.1 mutants were grown under high (18.8 mM) or low (0.5mM) nitrate concentrations (high N/low N) at both temperatures (at 22°C and at 10°C), a similar RH phenotype was detected between both nitrate conditions either at 22°C or 10°C (**Figure 5D**). In contrast, Wt Col-0 RH was sensitive to both low temperature and nitrate levels. Taken together,

291    these results suggest low temperature may restrict nitrate accessibility to the RH linked to a
292    lower mobility in the media affecting NRT1.1-mediated signaling upstream of TOR-S6K activation.
293
294

**FER-ROP2-TOR hub is required for nitrate starvation-mediated RH growth response**. Recently it
was reported that RALF1-FERONIA complex is able to interact with TOR kinase under low N
conditions modulating the TOR downstream signaling pathway[52]. As described above, we found
that Col-0 grown at 22°C under high N condition have a shorter RH phenotype as compared to
low N condition. This growth response is consistent with previous studies in which the RH length
in *Arabidopsis* and other species decrease as the nitrate concentration increases although the
molecular mechanism remained unknown[60,74]. To gain a better insight into the role of FER and
TOR in this process, we decided to test the *fer* mutants, the TOR pathway mutants and selected
overexpressing lines (TOR and S6K) in low (0.5 mM) and high nitrate (18.8 mM) media conditions
(**Figure 6A** and **Figure S7**). Under low nitrate conditions, the RH response was suppressed in both
*fer* mutants and in the *tor-es* inducible mutant compared to Col-0. This result supports the idea
that FER and TOR are necessary for the RH growth mediated by low nitrate conditions. The *TOR*
*OE* line showed a RH growth phenotype similar to Col-0 under low N and a higher RH growth in
excess of N. It is well known that nitrogen starvation causes enhanced root and RH growth,
whereas an excess of nitrate inhibits primary root growth[75], because of osmotic stress[71,].
Furthermore, plants overexpressing TOR in a nitrate excess medium have a longer primary root
phenotype[71]. According to our data that *TOR OE* line showed longer RH than Col-0 under normal
conditions (**Figure 6A**). The constitutive overexpression of TOR can lead not only to a longer
primary root but also to longer RH when plants are grown on a high nitrate media to alleviate the
nutrient stress. The constitutive active mutant of ROP2 suppressed the TOR OE RH phenotype
while *rop2* mutant was unable to respond to contrasting levels of nitrates. A model was proposed
where ROP GTPases and the cytoplasmic and apoplastic pH fluctuations can regulate RH tip
growth in a nitrogen supply dependent manner thus maintaining the unidirectional growth during
the RH elongation[77]. Additionally, nitrate and ammonium levels restore TOR activation under
nitrogen-starvation condition via ROP2 activation[43]. We then tested if other components on the
TOR complex are also involved in the low nitrate perception linked to RH growth (**Figure 6B**).
Similarly to their low temperature phenotype, *raptor1b* and *lst8-1* do not respond with
differential RH growth under low nitrate. Mutation of RAPTOR1B resulted in a strong reduction of
TOR kinase activity, leading to massive changes in carbon and nitrogen metabolism[78]. The RH
phenotype derived from the overexpression of S6K resembled that of TOR OE line (**Figure 6A, 6B**).
Altogether, our results show that low-nitrate (low N) RH responses are similar to those
determined for low temperature. This is consistent with the idea that nitrate is one of the main
signals that trigger RH expansion upon perception by NRT1.1, and subsequently transduction by
FER, ROP2 and TOR pathway. We then tested how changed are the levels of S6K-p/S6K as a

readout of TOR activation in low N and high N conditions in FER, TOR silenced line (*tor-es*) and
NRT1.1 mutants compared to Wt Col-0 (**Figure 6C**). First, Wt Col-0 showed an increased in S6K-
p/S6K ratio in low nitrogen similarly to the low temperature effect. On the contrary, and as
expected, *fer-4*, *fer-5* and *tor-es* showed much reduced levels of S6K-p/S6K in both nitrate
conditions but much lower in low nitrogen. Importantly, the effect of low N on increasing S6K-p
levels is abolished in NRT1.1 mutants (*chl1-9*, *chl1-5* and CHLT101D), which is similar to the
results observed under low temperature conditions. These results indicate that nitrate effect on
S6K-p/S6K ratio depends on NRT1.1 and the FER-TOR complex.
337
Since an integrative gene regulatory network analysis of TF–target interactions positioned TGA1
and its homolog TG4 as the most influential TFs in the nitrate response[79,80], we decide to test if
they play a role in the response to low temperature in RHs and low/high levels of Nitrogen
(**Figure S8**). As expected *tga1 tga4* double mutant failed to respond to low temperature and low
nitrogen while TGA1 OE showed a constitutive growth response regardless the temperature and
nitrogen levels. This confirms that nitrate, NRT1.1 and its downstream signaling pathway
including the transcriptional responses controlled by TGA1-TG4 act an important pathway in RH
growth at low temperature.
346
347 **Discussion**
348
Growth and development of plants and animals are based on nutrient and hormonal signaling
that constitute as the main regulatory networks in eukaryotes. Unraveling the functions of the
regulatory hubs and their detailed molecular mechanisms are critical for deeper understanding of
these signaling pathways. In all cells, there is a central hub composed by the evolutionarily
conserved TOR protein kinase that integrates nutrient and energy levels coupled to stress
signaling networks to further modulate cell growth[81–84]. In addition, the cell surface receptor FER
acts as a versatile sensor of signals coming from the environment[51,85,86]. Specifically, our study
uncovers a new molecular mechanism by which plants use FER-ROP2-TOR signaling pathway to
control RH elongation under low nutrient conditions, specifically nitrate, induced by low
temperature stress (**Figure 7**).
359
In this work, we discovered that the cell surface FER receptor regulates TOR apical localization
and further downstream activation, both controlling S6K phosphorylation linked to RH growth.
Based on the results obtained and previous evidences, here we propose a model in which the
FER-ROP2-TOR axis might be able to regulate the RH growth under low temperature with
remarked attention to variable nitrate conditions (**Figure 7**). In agreement with this, it was
previously shown that ROP2 in response to auxin is an upstream effector of TOR activation[44] and
by inorganic and organic nitrogen inputs[43]. In addition, FER was shown to directly activate the

TOR/RAPTOR1B signaling pathway under low inorganic and organic nitrogen conditions[52] and is
an upstream regulator of the ROPGEF4-ROP2 signaling pathway that controls ROS-mediated RH
development[47,87]. In addition, ROP2 promotes TOR accumulation close to the cell periphery[44],
suggesting that plasma membrane-localized FER together with ROP2 function in regulating the
relocation of TOR, close to the RH tip. Interestingly, such a local, non-genomic regulatory circuit
may also explain the rapid, cell-autonomous growth regulation in RH cells observed under
asymmetric conditions in the dual-flow-RootChip[6]. All these findings are in line with our results
and support the idea of the existence of the FER-ROP2-TOR pathway being highly relevant for RH
growth in a changing nutritional environment. Interestingly, we recently showed that FER
regulates localized protein synthesis during polar RH cell growth[5,88], suggesting that FER-TOR may
also regulates polar protein synthesis in RHs. All the data together clearly highlight a key role of
the FER-ROP2-TOR pathway as a core complex in sensing nutrients (e.g. nitrates) to direct RH
growth.

Nitrogen is an essential nutrient for plants that when scarce limits plant growth. Although
inorganic nitrogen (e.g. nitrate) is a major source of nitrogen for plants in soil, organic forms of
nitrogen, like amino acids, can also be assimilated by plants from soil[89,90]. Changes in nitrate
concentrations in the media mimicked this response at low temperature and NRT1.1 mutants are
insensitive at RH growth level regardless the temperature or nitrate levels. We hypothesize that
low nitrate in the media might rapidly induce the levels of mature-active RALF1 (and possibly
other RALF peptides), which might then activate the FER-ROP2-TOR pathway and the
downstream fast cell elongation effect to search for further nutrients sources. This autocrine
mechanism dependent on RALF1-FER growth activation has been demonstrated for RH[88], but not
in a context that involves low temperature/low nitrate and TOR pathway. It is expected that low
nitrates levels will impact on amino acid homeostasis. In line with our results, branched-chain
amino acids have been shown to serve as upstream activators of TOR in plants[40]. TOR is also
involved in mediating amino acid-derived metabolic regulatory signals that influence respiratory
activity and plant metabolic rate[91]. Specifically, a total of 15 proteinogenic amino acids can
reactivate TOR under inorganic-N starvation conditions[43]. In addition, the FER-TOR pathway
responds to Gln, Asp, and Gly, reinforcing that amino acids serve as conserved upstream
regulators of the TOR signaling pathway in animals and plants[52]. It is tempting to speculate that
under low nitrates condition (and low temperature), the RH cell compensates with an altered
amino acid homeostasis to trigger TOR activation linked fast cell growth.

Finally, several questions in our proposed model remain to be answered in future studies. How
does low nitrate concentration (low temperature) triggers FER-TOR activation?. Does RALF1 and
other RALFs respond to low nitrate to activate FER?. Does FER activate NRT1.1 directly?. Recently,
it was shown that FER-regulated ROP2 triggers a new mechanism of negative regulation in the

case of rhizosphere microorganisms such as *Pseudomonas*[76]. Although not tested here, FER-
ROP2-TOR signaling coupled to enhanced RH growth under low temperature (low-nutrients)
might be linked to the root growth in specific soil conditions to select favorable microbiota in the
soil. This pathway may integrate complex signals from soil nutrients and microbiota in the
rhizosphere to plant root cell growth mechanism.

**Figure 1. High levels of FER in its phosphorylated form are required to trigger the low**
**temperature RH growth.**

(**A**) Scatterplot of RH length of Col-0, of *fer-4*, *fer-5*, and *eru* mutants grown at 22°C or at 10°C. RH
growth is enhanced at low temperature in Wt Col-0 and *eru* mutant but not in the *fer-4* and *fer-5*
mutants. Each point is the mean of the length of the 10 longest RHs identified in the maturation
zone of a single root. Data are the mean ± SD (N=10 roots), two-way ANOVA followed by a
Tukey–Kramer test; (\*\*\*) *p* <0.001, NS= non-significant. Results are representative of three
independent experiments. Asterisks indicate significant differences between Col-0 and the
corresponding genotype at the same temperature or between the same genotype at different
temperatures. Representative images of each genotype are shown on the right. Scale bars= 300
μm. <span style="color:red">Numbers under the plots represents RH growth ratio 10°C/22°C.</span>
(**B-C**) Increased level of FER in root cells after its activation by low temperature. (**B**) Confocal
images of root apex of a fluorescent translational reporter line of FER (*pFER:FER-GFP*) at 22°C and
after transfer from ambient to low temperature (22°C→10°C). On the Bottom: semi-quantitative
evaluation of the GFP fluorescence intensity across the root tip at 22°C and after transfer from
ambient to low temperature at the transition zone, indicated by a interrupted white line.
Fluorescence intensity is expressed in arbitrary units (AU), N=4-6 roots. Results are representative
of three independent experiments. Scale bars: 50 μm.
(**C**) Confocal images of short root hairs of a fluorescent translational reporter line of FER
(*pFER:FER-GFP*) at 22°C and after transfer from ambient to low temperature (22°C→10°C) On the
bottom: semi-quantitative evaluation of the GFP fluorescence intensity across the subapical root
hair zone at 22°C and after transfer from ambient to low temperature, indicated by a interrupted
white line. Fluorescence intensity is expressed in arbitrary units (A.U.), N=6-8 roots, 8-11 root
hairs. Results are representative of three independent experiments. Scale bars=5 μm.
(**D**) Phosphorylation levels on FER (FER-p in *pFER:FER-GFP*) increases after 3 days at low
temperature in roots. Protein loading control (Coomasie Blue) is indicated below. FER-p/FER
ratios were analyzed by ImageJ.

**Figure 2. TORC signaling pathway is required for low temperature triggered RH growth.**

(**A**) Scatterplot of RH length of Col-0 and *tor-es* line grown at 22°C or at 10°C. Differential growth of RH at low temperature is suppressed in the estradiol inducible *RNAi TOR* line. Each point is the mean of the length of the 10 longest RHs identified in the maturation zone of a single root. Data are the mean ± SD (N=7-10 roots), two-way ANOVA followed by a Tukey–Kramer test; (***) $p<0.001$, NS=non-significant. Results are representative of three independent experiments. Asterisks indicate significant differences between Col-0 and the corresponding genotype at the same temperature or between the same genotype at different temperatures. Representative images of each line are shown below. Scale bars=300 µm. Numbers under the plots represents RH growth ratio 10°C/22°C.

(**B**) Differential growth of RH at low temperature is abolished in the Col-0 treated with 250 nM of TOR inhibitor, AZD-8055. Each point is the mean of the length of the 10 longest RHs identified in the maturation zone of a single root. Data are the mean ± SD (N=7 roots), two-way ANOVA followed by a Tukey–Kramer test; (***) $p<0.001$, NS=non-significant. Results are representative of three independent experiments. Asterisks indicate significant differences between Col-0 and the corresponding genotype at the same temperature or between the same genotype at different temperatures. Representative images of each line are shown below. Scale bars=300 µm. Numbers under the plots represents RH growth ratio 10°C/22°C.

(**C**) RH elongation of TOR and downstream TOR signaling pathway mutants under low temperature. Each point is the mean of the length of the 10 longest RHs identified in the maturation zone of a single root. Data are the mean ± SD (N=7-12 roots), two-way ANOVA followed by a Tukey–Kramer test; (*) $p<0.05$, (**) $p<0.01$, (***) $p<0.001$, NS=non-significant. Results are representative of three independent experiments. Asterisks indicate significant differences between Col-0 and the corresponding genotype at the same temperature or between the same genotype at different temperatures. Representative images of each line are shown on the right. Scale bars=300 µm. Numbers under the plots represents RH growth ratio 10°C/22°C.

(**D**) Analysis of the phosphorylation state of S6K (S6K-p/S6K ratio) in Col-0. A representative immunoblot is shown of two biological replicates (see **Supplementary Material 1**). S6K-p/S6K ratio was analyzed by ImageJ.

Figure 3. FER controls polar TOR localization and subsequent activation thus increasing phosphorylation of S6K under low temperature.

(**A**) Enhanced FER-TOR interaction at low temperature in roots by Immunoprecipitation (IP). A
representative experiment of three replicates is shown (see **Supplementary Material 1**).
(**B**)  Representative images showing TOR immunolocalization in RHs is FER-dependent. Scale
bar=10 μm. Control images with the pre-immune serum as primary antibody are included in
**Figure S3**.
(**C**) Apical and total TOR signal quantification in RHs showed in (B). A ROI at the RH tip of the
fluorescent signal or at the entire RH TOR-derived fluorescent signal was measured. Fluorescence
AU data are the mean ± SD (N=10-15 root hairs), two-way ANOVA followed by a Tukey–Kramer
test; (***) $p<0.001$, NS= non-significant. Results are representative of two independent
experiments. Asterisks on the graph indicate significant differences between genotypes.
(**D**) Analysis of the phosphorylation state of S6K in Wt Col-0, *fer-4, fer-5*, and *FER*$^{K565R}$ mutants
Arabidopsis roots. Phosphorylation of S6K (S6K-p) is enhanced at low temperature and requires
FER active kinase. A representative immunoblot is shown of three biological replicates (see
**Supplementary Material 1**). S6K-p/S6K ratio was analyzed by Image J. Wt Col-0 immublot is the
same of the Figure 2D.

**Figure 4.** ROP2 is required to triggers RH growth at low temperature.

(**A-B**) Increased level of ROP2 after its activation by low temperature.

(**A**) Confocal images of root apex of a fluorescent translational reporter line of ROP2

(*p*ROP2:ROP2-mCitrine) at 22°C (A) and after transfer from ambient to low temperature (22°C→

10°C). On the bottom: semi-quantitative evaluation of the mCitrine fluorescence intensity across

the root tip at 22°C and after transfer from ambient to low temperature at the transition zone,

indicated by a interrupted white line. Fluorescence intensity is expressed in arbitrary units (A.U.),

N=4-5 roots. Results are representative of three independent experiments. Scale bars=50 μm.

(**B**) Increased level of ROP2 in the plasma membrane of RHs after its activation by low

temperature. Confocal images of short RHs of a fluorescent translational reporter line of ROP2

(ROP2p:ROP2-mCitrine) at 22°C and after transfer from ambient to low temperature (22°C→10°

C). On the bottom: semi-quantitative evaluation of the mCitrine fluorescence intensity across the

RH apex at 22°C and after transfer from ambient to low temperature, indicated by a interrupted

white line. Fluorescence intensity is expressed in arbitrary units (A.U.), N=4-5 roots, 6-8 root

hairs. Results are representative of three independent experiments. Scale bars=5 μm.

(**C**) ROP2 is required for RH at low temperature meanwhile *rop2, rop2 rop4* and *CA-ROP2/TOR OE*

blocks RH growth. Each point is the mean of the length of the 10 longest RHs identified in a single

root. Data are the mean ± SD (N=10-15 roots), two-way ANOVA followed by a Tukey–Kramer test;

(**\*\***) $p<0.01$, (**\*\*\***) $p<0.001$. Results are representative of three independent experiments.

Asterisks indicate significant differences. Representative images of each line are shown below.

Scale bars=300 μm. <span style="color:red">Numbers under the plots represents RH growth ratio 10°C/22°C.</span>

(**D**) Analysis of the phosphorylation state of S6K in *TOR OE* and *TOR OE /CA-ROP2* Arabidopsis

roots. Phosphorylation of S6K (S6K-p) is enhanced in *TOR OE* and suppressed in *TOR OE*/ROP2-CA.

A representative immunoblot is shown of <span style="color:red">three biological replicates (see **Supplementary**</span>

<span style="color:red">**Material 1**)</span>. S6K-p/S6K ratio was analyzed by Image J.

**Figure 5**. Nitrate acts as a RH growth signal perceived by NRT1.1 and triggered by TOR pathway
at low temperature growth.

(**A**) High nitrate supplemented in M407 media (18.8 and 37.6 mM) partially suppresses low
temperature enhanced RH growth in Wt Col-0 plants. Each point is the mean of the length of the
10 longest RHs identified in the maturation zone of a single root. Data are the mean ± SD (N=10
roots), two-way ANOVA followed by a Tukey–Kramer test; (**\*\*\***) $p<0.001$. Results are
representative of three independent experiments. Asterisks indicate significant differences
between 0.5mM N concentration and every other N concentration at the same temperature or
between the same N concentrations at different temperatures. Representative images of Col-0
under the different nitrate concentrations are shown below. Scale bars=300 μm. Numbers under
the plots represents RH growth ratio $10^{o}C/22^{o}C$.
(**B**) Low temperature RH growth is regulated by the nitrate sensor NRT1.1 (CHL1).  Scatter-plot of
RH length of Col-0 and NRT1.1 mutants in 0.5X MS (contains 9.4 mM Nitrates) grown at 22°C or
10°C. Each point is the mean of the length of the 10 longest RHs identified in the maturation zone
of a single root. Data are the mean ± SD (N=10-15 roots), two-way ANOVA followed by a Tukey–
Kramer test; (**\*\*\***) $p<0.001$, NS=non-significant. Results are representative of three independent
experiments. Asterisks indicate significant differences between Col-0 and the corresponding
genotype at the same temperature or between the same genotype at different temperatures.
Representative images of each line are shown below. Scale bars=300 μm. Numbers under the
plots represents RH growth ratio $10^{o}C/22^{o}C$.
(**C**) Analysis of the phosphorylation state of S6K in Wt Col-0 and *chl1-9* mutant roots. A
representative immunoblot is shown of three biological replicates (see **Supplementary Material**
**1**). S6K-p/S6K ratio was analyzed by ImageJ. Wt Col-0 immunoblot is the same of the Figure 2D
and Figure3D.
(**D**) RH growth responses under high and low nitrate (18.8 mM and 0.5 mM, respectively) in Wt
Col-0, *chl1-5*, *chl1-9* mutants, and *CHL1$^{T101D}$* grown at 22°C or 10°C. Each point is the mean of the
length of the 10 longest RHs identified in the maturation zone of a single root. Data are the mean
± SD (N=10-15 roots), two-way ANOVA followed by a Tukey–Kramer test; (**\*\*\***) $p<0.001$, NS= non-
significant. Results are representative of three independent experiments. Asterisks indicate
significant differences between Col-0 and the corresponding genotype at the same temperature
or between the same genotype at different temperatures. Numbers under the plots represents
RH growth ratio $10^{o}C/22^{o}C$.

**Figure 6. Low nitrate perception relays in FER, ROP2 and TOR to trigger RH growth.**

(**A**) Scatterplot of RH length of Col-0, *fer-4*, *fer-5*, *rop2*, TOR OE/CA-ROP2 and TOR OE grown in
low (0.5 mM) and high nitrate (18.8 mM) conditions in M407 media, both at 22°C. Each point is
the mean of the length of the 10 longest RHs identified in the maturation zone of a single root.
Data are the mean ± SD (N=10 roots), two-way ANOVA followed by a Tukey–Kramer test; (**)
*p*<0.01, (***) *p*<0.001, NS=non-significant. Results are representative of three independent
experiments. Asterisks indicate significant differences between the same genotype at different N
concentration or between different genotypes at the same N concentration. Representative
images of each line are shown on the right. Scale bars=300 μm. Numbers under the plots
represents RH growth ratio low N/high N.
(**B**) Scatterplot of RH length of Col-0, *raptor1b*, *lst8-1*, *rps6b* and S6K OE mutants grown in low
and high N conditions. Each point is the mean of the length of the 10 longest RHs identified in the
maturation zone of a single root. Data are the mean ± SD (N=10 roots), two-way ANOVA followed
by a Tukey–Kramer test; (***) *p*<0.001, NS=non-significant. Results are representative of three
independent experiments. Asterisks indicate significant differences between the same genotype
at different N concentration or between different genotypes at the same N concentration.
Numbers under the plots represents RH growth ratio low N/high N.
(**C**) Analysis of the phosphorylation state of S6K in Wt Col-0, in NRT1.1 mutants (*chl1-9, chl1-5,*
CHLT101D), and in FER mutants (*fer-4, fer-5,* and *FER$^{K565R}$*). A representative immunoblot is shown
of a two biological replicates (see **Supplementary Material 1**). S6K-p/S6K ratio was analyzed by
Image J.

**Figure 7. Model of FER-ROP2-TOR function in RH elongation at limited nutrient availability caused by low temperature**. The model is based on the current evidence shown in this work and together with previous works. We have demonstrated that low temperature and nitrate conditions trigger RH elongation response through activation of the receptor kinase FERONIA. FERONIA triggers further activation of TOR either via direct binding to TOR (Left) or induction of the GTPase ROP2-TOR-S6K1 signaling axis via activation of GTPase ROP2 (Right). However, high levels of active ROP2 are able to negatively repress TORC activity. FERONIA can also activate NRT1.1 to promote RH elongation. Several questions (?) remain to be answered in future studies such as how low nitrate (triggered by low temperature) modulates FER-TOR activation? Does FER activate NRT1.1 directly? Do RALF1 and other RALFs respond to low nitrate to activate FER and downstream FER-TOR signaling?

585 **Experimental Procedures**
586

**Plant Material and Growth Conditions.** *A. thaliana* Col-0 ecotype was used as a wild-type plant.
To test the low and high nitrate response a 0.5X M407 medium without N, P nor K (M407,
PhytoTechnology Laboratories, https://www.phytotechlab.com/) supplemented with 0.8% plant
agar (Duchefa, Netherlands); 1.17mM MES, 0.625 mM $KH_2PO_4$ monobasic and 0.5mM $KNO_3$ (low
nitrate medium) 9.4mM, 18.8 mM and 37.6 mM $KNO_3$ (high nitrate medium) was used. Media
composition is detailed in **Table S1**. Plants were grown in the above media in continuous light
(120 µmol.seg$^{-1}$.m$^{-2}$) either: 8 days at 22°C in the case of *tor-es* line assay, 8 days at 22°C for
nitrate response assay or 5 days at 22°C + 3 days at 10°C for the low temperature and nitrate
response combination assay. For the rest of the low temperature experiments plants were grown
on regular 0.5X MS agar plates. For the imaging of fluorescence intensity distribution in root tips
and RH, seedlings were grown for 3 days at 22°C + 3-4 days at 10°C. All mutants and transgenic
lines are listed in **Table S2**. *chl1-5*, *chl1-9* and T101D mutants were kindly donated by Dr Yi-Fang
Tsay. *tor-es* was kindly donated by Dr Ezequiel Petrillo, *lst8-1*, *rps6b*, *raptor1b* and overexpressing
line S6K1 (S6K OE) kindly donated by Dr Elina Welchen and TOR OE and TOR OE/CA-ROP2 kindly
donated by Dr. Lyubov Ryabova.
602

**Pharmacological Treatments.** For all experiments, plants were grown first on solid 0.5X MS
medium at 22°C for 5 days in continuous light. According to the specific treatment plants were
transferred to plates containing regular solid 0.5X MS supplemented with 100nM IAA (auxin
treatment), 250 nM AZD-8055 (TOR inhibition) or 10 µM β-estradiol (*tor-es* line low temperature
treatment) and grown at 22°C for 5 days + 3 days at 10°C in continuous light. RH phenotype was
quantified after each 3 days span. For nitrate response treatment plants were transferred to
plates containing low or high nitrate solid media supplemented with 10 µM β-estradiol (*tor-es*
line), grown at 22°C for 3 days in continuous light and RH phenotype quantified.
611

**Root hair phenotype.** Seeds were surface sterilized and stratified in darkness for 3 days at 4°C.
Then grown *in vitro* on a specific condition and medium in a plant growth chamber in continuous
light (120 µmol.sec$^{-1}$.m$^{-2}$) at 22°C and/or 10°C. The quantitative analyses of RH phenotypes of Col-
0 and transgenic lines were made the last day of the growth conditions described in the two
previous sections. In the case of low temperature treatment, the measurements were done after
5 days at 22°C and after 3 days at 10°C. For that purpose, 10 fully elongated RH from the
maturation zone were measured per root under the same conditions from each treatment and
control. Images were captured using an Olympus SZX7 Zoom Stereo Microscope (Olympus, Japan)
equipped with a Q-Colors digital camera and Q Capture Pro 7 software (Olympus, Japan). Results
were expressed as the mean ± SD using the GraphPad Prism 8.0.1 (USA) statistical analysis

software. Results are representative of three independent experiments, each involving 7–20
roots.

**Confocal Microscopy.** For measurements of fluorescence intensity distributions after cold stress
(22°C→10°C) in root tips and root hairs of $p$FER::FER-GFP and $p$ROP2::ROP2-mCitrine lines
confocal laser scanning microscopy Zeiss LSM 710 (Carl Zeiss, Germany) was used. For image
acquisition, 10x/0.3 NA EC Plan-Neofluar objective for root tips, and 40x/1.4 Oil DIC Plan-
Apochromat objective for root hairs were used. GFP signal was excited with 488 nm argon laser
at 4% laser power intensity and emission band of 493-549 nm. mCytrine signal was excited with
514 nm argon laser at 4% laser power intensity and emission band 519-583 nm. Fluorescence
intensity measurements (in A.U.) were generated using Zen Black 2011 software (Carl Zeiss,
Germany) and graphically edited in Microsoft Excel. Results are representative of three
independent experiments, each involving 1–2 roots and 1 to 4 hairs per root.

**Co-IP assay.** For Co-IP assays using A/G agarose and an anti-TOR antibody[32], 30 µL of A/G beads
(Thermo Fisher Scientific Inc., 20421) was resuspended and washed three times using NEB buffer
(20 mM HEPES [pH 7.5], 40 mM KCl, 5 mM MgCl$_2$) before adding 8 µL of anti-TOR antibody or
preimmune serum as a negative control[32] in a total volume of 500 µL of NEB buffer, followed by
incubation for 4 h at 4 °C. Col-0 seedlings were first grown at 22 °C for 5 days then transferred to
22 °C or 10 °C for 2 days or 3 days. For protein extraction from plants, the collected materials
were ground to a fine powder in liquid nitrogen and solubilized with NEB-T buffer (20 mM HEPES
[pH 7.5], 40 mM KCl, 5 mM MgCl$_2$, 0.5% Triton X-100) containing 1 × protease inhibitor cocktail
(Thermo Fisher Scientific Inc., 78430) and 1 × phosphatase inhibitor (Thermo Fisher Scientific Inc.,
78420) and incubated for 1 h on the ice. The extracts were centrifuged at 16,000 g at 4 °C for 15
min, and the resultant supernatant was incubated with prepared antibody-beads from the above
step. After overnight incubation at 4°C with rotation, the agarose beads were washed five times
with the NEB buffer and eluted with elution buffer (0.2 M glycine, 0.5% Triton X-100, pH 7.5).
Anti-FER and anti-TOR antibodies[30] were used for immunoblotting to detect the
immunoprecipitates.

**TOR immunolocalization**. Col-0 and *fer-4* seedlings were first grown at 22 °C for 5 days before
transferred to 22 °C or 10 °C for 3 days. Then seedlings were collected and incubated for 10 min
under vacuum (0.05 MPa) in phosphate-buffered saline (PBS) containing 4% paraformaldehyde
and 0.1% Triton X-100. Seedlings were washed gently three times (10 min for each wash) in PBS
and then the cell wall was digested in 2% Driselase (Sigma, D8037) in PBS for 18 min at 37 °C and
washed five times with PBS. The permeability of the seedlings was increased by incubating them
in 3% IGEPAL CA-630 (Sigma, 18896) and 10% DMSO in PBS for 18 min, followed by washed three

times with PBS. Seedlings were incubated in 2% bovine serum albumin (BSA) (Ameresco, 0332) in
PBS for 1.5 h and then incubated with primary TOR antibody[32] (antibody diluted 1:600 in 2% BSA)
for overnight at 4°C. The seedlings were washed with PBS for five times. Fluorophore-labeled
secondary antibody (goat–mouse secondary antibody, diluted 1:600 in 2% BSA) was incubated
with the samples at 37°C for 5 h in the dark. Seedlings were washed five times with PBS before
observation. Fluorescent signal detection and documentation were performed using a Nikon
confocal laser scanning microscope with a 560-nm band-pass filter for IF555 detection.
666
**S6K-p/S6K immunoblotting detection.** For immunological detection of S6K-p and S6K1/2, total
soluble proteins were extracted from 50 mg of plant materials grown as indicated previously with
100 μL 2 × Laemmli buffer supplemented with 1% Phosphatase Inhibitor Cocktail 2 (Thermo
Fisher Scientific Inc., 78430). Proteins were denatured for 10 min at 95°C and separated on 10%
or 8% SDS-PAGE. Rabbit *At*TOR polyclonal antibodies (Abiocode, R2854-2), rabbit polyclonal
S6K1/2 antibodies (Agrisera, AS121855), and S6K1-p (phospho T449) antibody (Abcam,
ab207399) and ACTIN antibody (Abmart, M20009) were used for immunoblotting.
674
**Quantitative PCR (qPCR).** Total root RNA was extracted from plantlets grown *in vitro* at 22 °C and
10 °C using the RNeasy®Plant Mini Kit (QIAGEN, Germany). One microgram of total RNA was
reverse transcribed using an oligo(dT)$_{20}$ primer and the Super Script™ IV RT (Invitrogen,USA)
according to the manufacturer's instructions. cDNA was diluted 20-fold before PCR. *q*PCR was
performed on a LightCycler®480 Instrument II (Roche, USA) using 2 μL of 5X HOT
FIREPol®EvaGreen® qPCR Mix Plus(no ROX) (Solis BioDyne, Estonia), 2 μL of cDNA, and 0.25 μM of
each primer in a total volume of 10 μL per reaction. ACT2 (AT3G18780) gene was used as
reference for normalization of gene expression levels (ACT2 primers, F:
GGTAACATTGTGCTCAGTGGTGG R: CTCGGCCTTGGAGATCCACATC; TOR primers, F:
GAAGATGAAGATCCCGCTGA R: GCATCTCCAAGCATATTTACAGC[44]). The cycling conditions were: 95
°C for 12 min., 35 cycles of 95°C for 15 sec., 60°C for 1 min and finally a melting curve from 60°C
to 95 °C (0.05°/sec). Data were analyzed using the ΔΔC$_t$ method[92] and LightCycler®480 Software,
version 1.5 (Roche). Two independent experiments with three biological and three technical
replicates per experiment, were performed.
689
**Qualitative determination of KNO$_3$ diffusion at 10°C and 22°C.** This method was adapted from a
previous protocol[93]. First, 25 ml of 0.8% agar-water media was prepared into plastic Petri dishes.
After gelation it was removed the center of the agar plate by using the rear part of a p1000 tip as
a puncher (removing a volume of approx. 200 μl of gel). After equilibration at either 10°C or 22°C
it was added 200 μl of liquid 0.8% Agar (at 40°C) containing 0,94M KNO$_3$ to fill up the well and
started the diffusion time. After 3 h of incubation, 20 μl agar disks were taken at different
distances from the border of the KNO$_3$ zone (from 5mm to 25mm in 5mm steps). For this

purpose, we used a Pasteur pipette as a puncher. The agar disks were incubated with 200 µl of
2mM diphenylamine (($C_6H_5$)2NH) colorless solution in 14,4M $H_2SO_4$ with eventual shaking to
homogenize the reaction media. After 30 min we transferred 100ul of the bright blue resultant
reaction to a 96 well plate and measured OD at 595 nm. Serial dilutions showed deviations from
the expected linear behavior preventing the possibility of a robust quantitative assay. We used
high $KNO_3$ concentrations to prevent excessive dilution of the $NO_3^-$ below detection levels and
speed up the diffusion process.

**Acknowledgements**

We would like to thank Elina Welchen, Ezequiel Petrillo, Yi-Fang Tsay, Kriss Vissenberg for the seed lines. We thank NASC (Ohio State University) for providing T-DNA lines seed lines. J.M.E. is investigator of the National Research Council (CONICET) from Argentina. M.I. is supported by ANID FONDECYT POSTDOCTORADO [grant 3220138]. This work was supported by grants Natural Science Foundation of China (NSFC-32001974 and NSFC-31871396) to F.Y. and (NSFC-31900232) to L.S. and from ANPCyT (PICT2017-0066, and PICT2019-0015), by ANID – Programa Iniciativa Científica Milenio ICN17_022, NCN2021_010 and Fondo Nacional de Desarrollo Científico y Tecnológico [1200010] to J.M.E., by the Deutsche Forschungsgemeinschaft (DFG; GR4559_4-1, GR4559_5-1, EXC-2048/1 project ID 390686111) to G.G., and by the Czech Science Foundation GAČR (project Nr. 19-18675S) to J.Š.

**Author Contribution**

J.M.P performed most of the experiments, analysed the data and helped in the writing process of the manuscript. L.S. performed the S6K-p determinations, IP of TOR-FER and immunolocalization assay of TOR. L.K., M.O. and J.Š. provided live cell imaging data on FER and ROP2. V.B.G., J.M.P., T.U.I., M.A.I., M.M.R., S-Z., Y.S., R.A.G., M.S., L.A.R., J.M.A., G.G., J.S. helped in the data analysis and writing process of the manuscript. F.Y. designed research and analysed part of the data and J.M.E. designed research, analysed the data, supervised the project, and wrote the paper. All authors commented on the results and the manuscript. This manuscript has not been published and is not under consideration for publication elsewhere. All the authors have read the manuscript and have approved this submission.

**Competing financial interest**

The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to F.Y. (feng_yu@hnu.edu.cn) and J.M.E. (Email: jestevez@leloir.org.ar).

**REFERENCES**

1.    Šamaj, J., Müller, J., Beck, M., Böhm, N., and Menzel, D. (2006). Vesicular trafficking, cytoskeleton and signalling in root hairs and pollen tubes. Trends Plant Sci. *11*, 594–600.

2.    Pascale, A., Proietti, S., Pantelides, I.S., and Stringlis, I.A. (2019). Modulation of the Root Microbiome by Plant Molecules: The Basis for Targeted Disease Suppression and Plant Growth Promotion. Front. Plant Sci. *10*.

3.    Vijayakumar, P., Datta, S., and Dolan, L. (2016). ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4) promotes root hair elongation by transcriptionally regulating the expression of genes required for cell growth. New Phytol. *212*, 944–953.

4.    Kuběnová, L., Tichá, M., Šamaj, J., and Ovečka, M. (2022). ROOT HAIR DEFECTIVE 2 vesicular delivery to the apical plasma membrane domain during Arabidopsis root hair development. Plant Physiol., kiab595.

5.    Zhu, S., Estévez, J.M., Liao, H., Zhu, Y., Yang, T., Li, C., Wang, Y., Li, L., Liu, X., Pacheco, J.M., et al. (2020). The RALF1–FERONIA Complex Phosphorylates eIF4E1 to Promote Protein Synthesis and Polar Root Hair Growth. Mol. Plant *13*, 698–716.

6.    Stanley, C.E., Shrivastava, J., Brugman, R., Heinzelmann, E., van Swaay, D., and Grossmann, G. (2018). Dual-flow-RootChip reveals local adaptations of roots towards environmental asymmetry at the physiological and genetic levels. New Phytol. *217*, 1357–1369.

7.    Yi, K., Menand, B., Bell, E., and Dolan, L. (2010). A basic helix-loop-helix transcription factor controls cell growth and size in root hairs. Nat. Genet. *42*, 264–267.

8.    Datta, S., Prescott, H., and Dolan, L. (2015). Intensity of a pulse of RSL4 transcription factor synthesis determines Arabidopsis root hair cell size. Nat. Plants *1*, 1–6.

9.    Moison, M., Pacheco, J.M., Lucero, L., Fonouni-Farde, C., Rodríguez-Melo, J., Mansilla, N., Christ, A., Bazin, J., Benhamed, M., Ibañez, F., et al. (2021). The lncRNA APOLO interacts with the transcription factor WRKY42 to trigger root hair cell expansion in response to cold. Mol. Plant *14*, 937–948.

10.   Singha, S., Townsend, E.C. and Oberly, G.H. (1985). Mineral nutrient status of crabapple and pear shoots cultured in vitro on varying commercial agars. J. Am. Soc. Hortic. Sci. *110*, 407-411.

11.   Nonami, H., and Boyer, J.S. (1989). Turgor and Growth at Low Water Potentials. Plant Physiol. *89*, 798–804.

12.   Ghashghaie, J., Brenckmann, F., and Saugier, B. (1991). Effects of agar concentration on water status and growth of rose plants cultured in vitro. Physiol. Plant. *82*, 73–78.

13.   Buah, J.N., Kawamitsu, Y., Sato, S., and Murayama, S. (1999). Effects of Different Types and Concentrations of Gelling Agents on the Physical and Chemical Properties of Media and the Growth of Banana ( Musa spp.) in Vitro. Plant Prod. Sci. *2*, 138–145.

14.   Pacheco, J.M., Mansilla, N., Moison, M., Lucero, L., Gabarain, V.B., Ariel, F., and Estevez, J.M. (2021). The lncRNA APOLO and the transcription factor WRKY42 target common cell wall EXTENSIN encoding genes to trigger root hair cell elongation. Plant Signal. Behav. *16*, 1920191.

15.   Pacheco, J.M., Ranocha, P., Kasulin, L., Fusari, C.M., Servi, L., Aptekmann, A.A., Gabarain, V.B., Peralta, J.M., Borassi, C., Marzol, E., et al. (2022). Apoplastic class III peroxidases

775     PRX62 and PRX69 promote Arabidopsis root hair growth at low temperature. Nat.
776     Commun. *13*, 1310.

777 16. Xiong, Y., and Sheen, J. (2014). The role of target of rapamycin signaling networks in plant
778     growth and metabolism. Plant Physiol. *164*, 499–512.

779 17. Dobrenel, T., Mancera-Martínez, E., Forzani, C., Azzopardi, M., Davanture, M., Moreau, M.,
780     Schepetilnikov, M., Chicher, J., Langella, O., Zivy, M., et al. (2016). The arabidopsis TOR
781     kinase specifically regulates the expression of nuclear genes coding for plastidic ribosomal
782     proteins and the phosphorylation of the cytosolic ribosomal protein S6. Front. Plant Sci. *7*,
783     1–16.

784 18. Saxton, R.A., and Sabatini, D.M. (2017). mTOR Signaling in Growth, Metabolism, and
785     Disease. Cell *168*, 960–976.

786 19. Tatebe, H., and Shiozaki, K. (2017). Evolutionary Conservation of the Components in the
787     TOR Signaling Pathways. Biomolecules *7*, 1–17.

788 20. Van Leene, J., Han, C., Gadeyne, A., Eeckhout, D., Matthijs, C., Cannoot, B., De Winne, N.,
789     Persiau, G., Van De Slijke, E., Van de Cotte, B., et al. (2019). Capturing the phosphorylation
790     and protein interaction landscape of the plant TOR kinase. Nat. Plants *5*, 316–327.

791 21. Menand, B., Desnos, T., Nussaume, L., Bergert, F., Bouchez, D., Meyer, C., and Robaglia, C.
792     (2002). Expression and disruption of the Arabidopsis TOR (target of rapamycin) gene. Proc.
793     Natl. Acad. Sci. U. S. A. *99*, 6422–6427.

794 22. Anderson, G.H., Veit, B., and Hanson, M.R. (2005). The Arabidopsis AtRaptor genes are
795     essential for post-embryonic plant growth. BMC Biol. *3*, 1–12.

796 23. Deprost, D., Truong, H.N., Robaglia, C., and Meyer, C. (2005). An Arabidopsis homolog of
797     RAPTOR/KOG1 is essential for early embryo development. Biochem. Biophys. Res.
798     Commun. *326*, 844–850.

799 24. Mahfouz, M.M., Kim, S., Delauney, A.J., and Verma, D.P.S. (2006). Arabidopsis TARGET of
800     RAPAMYCIN interacts with RAPTOR, which regulates the activity of S6 kinase in response to
801     osmotic stress signals. Plant Cell *18*, 477–490.

802 25. Yang, H., Rudge, D.G., Koos, J.D., Vaidialingam, B., Yang, H.J., and Pavletich, N.P. (2013).
803     MTOR kinase structure, mechanism and regulation. Nature *497*, 217–223.

804 26. Rexin, D., Meyer, C., Robaglia, C., and Veit, B. (2015). TOR signalling in plants. Biochem. J.
805     *470*, 1–14.

806 27. Salem, M.A., Li, Y., Wiszniewski, A., and Giavalisco, P. (2017). Regulatory-associated protein
807     of TOR (RAPTOR) alters the hormonal and metabolic composition of Arabidopsis seeds,
808     controlling seed morphology, viability and germination potential. Plant J. *92*, 525–545.

809 28. Moreau, M., Azzopardi, M., Clé ment, G., Dobrenel, T., Marchive, C., Renne, C., Martin-
810     Magniette, M.L., Taconnat, L., Renou, J.P., Robaglia, C., et al. (2012). Mutations in the
811     Arabidopsis homolog of LST8/GβL, a partner of the target of Rapamycin kinase, impair
812     plant growth, flowering, and metabolic adaptation to long days. Plant Cell *24*, 463–481.

813 29. McCready, K., Spencer, V., and Kim, M. (2020). The Importance of TOR Kinase in Plant
814     Development. Front. Plant Sci. *11*, 1–7.

815 30. Henriques, R., Magyar, Z., Monardes, A., Khan, S., Zalejski, C., Orellana, J., Szabados, L., De
816     La Torre, C., Koncz, C., and BÖgre, L. (2010). Arabidopsis S6 kinase mutants display
817     chromosome instability and altered RBR1-E2F pathway activity. EMBO J. *29*, 2979–2993.

31.  Ahn, C.S., Han, J.A., Lee, H.S., Lee, S., and Pai, H.S. (2011). The PP2A regulatory subunit
Tap46, a component of the TOR signaling pathway, modulates growth and metabolism in
plants. Plant Cell *23*, 185–209.
32.  Xiong, Y., and Sheen, J. (2012). Rapamycin and glucose-target of rapamycin (TOR) protein
signaling in plants. J. Biol. Chem. *287*, 2836–2842.
33.  Patrick, R.M., Lee, J.C.H., Teetsel, J.R.J., Yang, S.H., Choy, G.S., and Browning, K.S. (2018).
Discovery and characterization of conserved binding of eIF4E 1 (CBE1), a eukaryotic
translation initiation factor 4E– binding plant protein. J. Biol. Chem. *293*, 17240–17247.
34.  Xiong, Y., and Sheen, J. (2015). Novel links in the plant TOR kinase signaling network. Curr.
Opin. Plant Biol. *28*, 83–91.
35.  Dobrenel, T., Caldana, C., Hanson, J., Robaglia, C., Vincentz, M., Veit, B., and Meyer, C.
(2016). TOR Signaling and Nutrient Sensing. Annu. Rev. Plant Biol. *67*, 261–285.
36.  González, A., and Hall, M.N. (2017).  Nutrient sensing and TOR signaling in yeast and
mammals . EMBO J. *36*, 397–408.
37.  Bakshi, A., Moin, M., Madhav, M.S., and Kirti, P.B. (2019). Target of rapamycin, a master
regulator of multiple signalling pathways and a potential candidate gene for crop
improvement. Plant Biol. *21*, 190–205.
38.  Pacheco, J.M., Canal, M.V., Pereyra, C.M., Welchen, E., Martínez-Noël, G.M.A., and
Estevez, J.M. (2021). The tip of the iceberg: emerging roles of TOR, and its regulatory
functions in plant cells. J. Exp. Bot. *72*, 4085–4101.
39.  Dong, Y., Silbermann, M., Speiser, A., Forieri, I., Linster, E., Poschet, G., Allboje Samami, A.,
Wanatabe, M., Sticht, C., Teleman, A.A., et al. (2017). Sulfur availability regulates plant
growth via glucose-TOR signaling. Nat. Commun. *8*, 1–10.
40.  Cao, P., Kim, S.J., Xing, A., Schenck, C.A., Liu, L., Jiang, N., Wang, J., Last, R.L., and Brandizzi,
F. (2019). Homeostasis of branched-chain amino acids is critical for the activity of TOR
signaling in Arabidopsis. Elife *8*, 1–24.
41.  Schaufelberger, M., Galbier, F., Herger, A., de Brito Francisco, R., Roffler, S., Clement, G.,
Diet, A., Hörtensteiner, S., Wicker, T., and Ringli, C. (2019). Mutations in the Arabidopsis
ROL17/isopropylmalate synthase 1 locus alter amino acid content, modify the TOR
network, and suppress the root hair cell development mutant lrx1. J. Exp. Bot. *70*, 2313–
2323.
42.  Sun, X., Chen, H., Wang, P., Chen, F., Yuan, L., and Mi, G. (2020). Low nitrogen induces root
elongation via auxin-induced acid growth and auxin-regulated target of rapamycin (TOR)
pathway in maize. J. Plant Physiol. *254*, 153281.
43.  Liu, Y., Duan, X., Zhao, X., Ding, W., Wang, Y., and Xiong, Y. (2021). Diverse nitrogen signals
activate convergent ROP2-TOR signaling in Arabidopsis. Dev. Cell *56*, 1283-1295.e5.
44.  Schepetilnikov, M., Makarian, J., Srour, O., Geldreich, A., Yang, Z., Chicher, J., Hammann, P.,
and Ryabova, L.A. (2017).  GTP ase ROP 2 binds and promotes activation of target of
rapamycin, TOR , in response to auxin . EMBO J. *36*, 886–903.
45.  Li, X., Cai, W., Liu, Y., Li, H., Fu, L., Liu, Z., Xu, L., Liu, H., Xu, T., and Xiong, Y. (2017).
Differential TOR activation and cell proliferation in Arabidopsis root and shoot apexes.
Proc. Natl. Acad. Sci. U. S. A. *114*, 2765–2770.
46.  Carol, R.J., Takeda, S., Linstead, P., Durrant, M.C., Kakesova, H., Derbyshire, P., Drea, S.,

861            Zarsky, V., and Dolan, L. (2005). A RhoGDP dissociation inhibitor spatially regulates growth
862            in root hair cells. Nature *438*, 1013–1016.

863    47.    Duan, Q., Kita, D., Li, C., Cheung, A.Y., and Wu, H.M. (2010). FERONIA receptor-like kinase
864            regulates RHO GTPase signaling of root hair development. Proc. Natl. Acad. Sci. U. S. A.
865            *107*, 17821–17826.

866    48.    Kang, E., Zheng, M., Zhang, Y., Yuan, M., Yalovsky, S., Zhu, L., and Fu, Y. (2017). The
867            microtubule-associated protein MAP18 affects ROP2 GTPase activity during root hair
868            growth. Plant Physiol. *174*, 202–222.

869    49.    Denninger, P., Reichelt, A., Schmidt, V.A.F., Mehlhorn, D.G., Asseck, L.Y., Stanley, C.E.,
870            Keinath, N.F., Evers, J.F., Grefen, C., and Grossmann, G. (2019). Distinct RopGEFs
871            Successively Drive Polarization and Outgrowth of Root Hairs. Curr. Biol. *29*, 1854-1865.e5.

872    50.    Huang, G.Q., Li, E., Ge, F.R., Li, S., Wang, Q., Zhang, C.Q., and Zhang, Y. (2013). Arabidopsis
873            RopGEF4 and RopGEF10 are important for FERONIA-mediated developmental but not
874            environmental regulation of root hair growth. New Phytol. *200*, 1089–1101.

875    51.    Xu, G., Chen, W., Song, L., Chen, Q., Zhang, H., Liao, H., Zhao, G., Lin, F., Zhou, H., and Yu, F.
876            (2019). FERONIA phosphorylates E3 ubiquitin ligase ATL6 to modulate the stability of 14-3-
877            3 proteins in response to the carbon/nitrogen ratio. J. Exp. Bot. *70*, 6375–6388.

878    52.    Song, L., Xu, G., Li, T., Zhou, H., Lin, Q., Chen, J., Wang, L., Wu, D., Li, X., Wang, L., et al.
879            (2022). The RALF1-FERONIA Complex Interacts with and Activates TOR Signaling in
880            Response to Low Nutrients. Mol. Plant *0*.

881    53.    Kwon, T., Alan Sparks, J., Liao, F., and Blancaflor, E.B. (2018). Erulus is a plasma membrane-
882            localized receptor-like kinase that specifies root hair growth by maintaining tip-focused
883            cytoplasmic calcium oscillations[OPEN]. Plant Cell *30*, 1173–1177.

884    54.    Schoenaers, S., Balcerowicz, D., Breen, G., Hill, K., Zdanio, M., Mouille, G., Holman, T.J., Oh,
885            J., Wilson, M.H., Nikonorova, N., et al. (2018). The Auxin-Regulated CrRLK1L Kinase ERULUS
886            Controls Cell Wall Composition during Root Hair Tip Growth. Curr. Biol. *28*, 722-732.e6.

887    55.    Xiong, Y., McCormack, M., Li, L., Hall, Q., Xiang, C., and Sheen, J. (2013). Glucose-TOR
888            signalling reprograms the transcriptome and activates meristems. Nature *496*, 181–186.

889    56.    Montané, M.H., and Menand, B. (2013). ATP-competitive mTOR kinase inhibitors delay
890            plant growth by triggering early differentiation of meristematic cells but no developmental
891            patterning change. J. Exp. Bot. *64*, 4361–4374.

892    57.    Escobar-Restrepo, J.M., Huck, N., Kessler, S., Gagliardini, V., Gheyselinck, J., Yang, W.C.,
893            and Grossniklaus, U. (2007). The Feronia receptor-like kinase mediates male-female
894            interactions during pollen tube reception. Science (80-. ). *317*, 656–660.

895    58.    Haruta, M., Sabat, G., Stecker, K., Minkoff, B.B., and Sussman, M.R. (2014). A peptide
896            hormone and its receptor protein kinase regulate plant cell expansion. Science (80-. ). *343*,
897            408–411.

898    59.    Chakravorty, D., Yu, Y., and Assmann, S.M. (2018). A kinase-dead version of FERONIA
899            receptor-like kinase has dose-dependent impacts on rosette morphology and RALF1-
900            mediated stomatal movements. FEBS Lett. *592*, 3429–3437.

901    60.    Jones, M.A., Shen, J.J., Fu, Y., Li, H., Yang, Z., and Grierson, C.S. (2002). The arabidopsis
902            Rop2 GTPase is a positive regulator of both root hair initiation and tip growth. Plant Cell
903            *14*, 763–776.

61.    Gendre, D., Baral, A., Dang, X., Esnay, N., Boutte, Y., Stanislas, T., Vain, T., Claverol, S.,
Gustavsson, A., Lin, D., et al. (2019). Rho-of-plant activated root hair formation requires
arabidopsis YIP4a/b gene function. Dev. *146*.
62.    Vatter, T., Neuhäuser, B., Stetter, M., and Ludewig, U. (2015). Regulation of length and
density of Arabidopsis root hairs by ammonium and nitrate. J. Plant Res. *128*, 839–848.
63.    Canales, J., Contreras-López, O., Álvarez, J.M., and Gutiérrez, R.A. (2017). Nitrate induction
of root hair density is mediated by TGA1/TGA4 and CPC transcription factors in Arabidopsis
thaliana. Plant J. *92*, 305–316.
64.    Liu, B., Wu, J., Yang, S., Schiefelbein, J., and Gan, Y. (2020). Nitrate regulation of lateral root
and root hair development in plants. J. Exp. Bot. *71*, 4405–4414.
65.    Ho, C.H., Lin, S.H., Hu, H.C., and Tsay, Y.F. (2009). CHL1 Functions as a Nitrate Sensor in
Plants. Cell *138*, 1184–1194.
66.    Gojon, A., Krouk, G., Perrine-Walker, F., and Laugier, E. (2011). Nitrate transceptor(s) in
plants. J. Exp. Bot. *62*, 2299–2308.
67.    Bouguyon, E., Brun, F., Meynard, D., Kubeš, M., Pervent, M., Leran, S., Lacombe, B., Krouk,
G., Guiderdoni, E., Zazímalová, E., et al. (2015). Multiple mechanisms of nitrate sensing by
Arabidopsis nitrate transceptor NRT1.1. Nat. Plants *1*, 15015.
68.    Krouk, G., Lacombe, B., Bielach, A., Perrine-Walker, F., Malinska, K., Mounier, E., Hoyerova,
K., Tillard, P., Leon, S., Ljung, K., et al. (2010). Nitrate-regulated auxin transport by NRT1.1
defines a mechanism for nutrient sensing in plants. Dev. Cell *18*, 927–937.
69.    Liu, K.-H., and Tsay, Y.-F. (2003). Switching between the two action modes of the dual-
affinity nitrate transporter CHL1 by phosphorylation. EMBO J. *22*, 1005–13.
70.    Forde, B.G., and Walch-Liu, P. (2009). Nitrate and glutamate as environmental cues for
behavioural responses in plant roots. Plant, Cell Environ. *32*, 682–693.
71.    Léran, S., Varala, K., Boyer, J.C., Chiurazzi, M., Crawford, N., Daniel-Vedele, F., David, L.,
Dickstein, R., Fernandez, E., Forde, B., et al. (2014). A unified nomenclature of nitrate
transporter 1/peptide transporter family members in plants. Trends Plant Sci. *19*, 5–9.
72.    Wang, W., Hu, B., Li, A., and Chu, C. (2020). NRT1.1s in plants: Functions beyond nitrate
transport. J. Exp. Bot. *71*, 4373–4379.
73.    Deprost, D., Yao, L., Sormani, R., Moreau, M., Leterreux, G., Bedu, M., Robaglia, C., and
Meyer, C. (2007). The Arabidopsis TOR kinase links plant growth, yield, stress resistance
and mRNA translation. EMBO Rep. *8*, 864–870.
74.    ROBINSON, D., and RORISON, I.H. (1987). ROOT HAIRS AND PLANT GROWTH ATLOW
NITROGEN AVAILABILITIES. New Phytol. *107*, 681–693.
75.    Deak, K.I., and Malamy, J. (2005). Osmotic regulation of root system architecture. Plant J.
*43*, 17–28.
76.    Song, Y., Wilson, A.J., Zhang, X.C., Thoms, D., Sohrabi, R., Song, S., Geissmann, Q., Liu, Y.,
Walgren, L., He, S.Y., et al. (2021). FERONIA restricts Pseudomonas in the rhizosphere
microbiome via regulation of reactive oxygen species. Nat. Plants *7*, 644–654.
77.    Bloch, D., Monshausen, G., Gilroy, S., and Yalovsky, S. (2011). Co-regulation of root hair tip
growth by ROP GTPases and nitrogen source modulated pH fluctuations. Plant Signal.
Behav. *6*, 426–429.
78.    Salem, M.A., Li, Y., Bajdzienko, K., Fisahn, J., Watanabe, M., Hoefgen, R., Schöttler, M.A.,

and Giavalisco, P. (2018). RAPTOR controls developmental growth transitions by altering
the hormonal and metabolic balance. Plant Physiol. *177*, 565–593.

79.  Vidal, E.A., Alvarez, J.M., Araus, V., Riveras, E., Brooks, M.D., Krouk, G., Ruffel, S., Lejay, L.,
Crawford, N.M., Coruzzi, G.M., et al. (2020). Nitrate in 2020: Thirty years from transport to
signaling networks. Plant Cell *32*, 2094–2119.

80.  Alvarez, J.M., Riveras, E., Vidal, E.A., Gras, D.E., Contreras-López, O., Tamayo, K.P.,
Aceituno, F., Gómez, I., Ruffel, S., Lejay, L., et al. (2014). Systems approach identifies TGA1
and TGA4 transcription factors as important regulatory components of the nitrate
response of Arabidopsis thaliana roots. Plant J. *80*, 1–13.

81.  Wullschleger, S., Loewith, R., and Hall, M.N. (2006). TOR signaling in growth and
metabolism. Cell *124*, 471–484.

82.  Wang, P., Zhao, Y., Li, Z., Hsu, C.C., Liu, X., Fu, L., Hou, Y.J., Du, Y., Xie, S., Zhang, C., et al.
(2018). Reciprocal Regulation of the TOR Kinase and ABA Receptor Balances Plant Growth
and Stress Response. Mol. Cell *69*, 100-112.e6.

83.  Wu, Y., Shi, L., Li, L., Fu, L., Liu, Y., Xiong, Y., and Sheen, J. (2019). Integration of nutrient,
energy, light, and hormone signalling via TOR in plants. J. Exp. Bot. *70*, 2227–2238.

84.  Fu, L., Wang, P., and Xiong, Y. (2020). Target of Rapamycin Signaling in Plant Stress
Responses. Plant Physiol. *182*, 1613–1623.

85.  Yang, T., Wang, L., Li, C., Liu, Y., Zhu, S., Qi, Y., Liu, X., Lin, Q., Luan, S., and Yu, F. (2015).
Receptor protein kinase FERONIA controls leaf starch accumulation by interacting with
glyceraldehyde-3-phosphate dehydrogenase. Biochem. Biophys. Res. Commun. *465*, 77–
82.

86.  Yeats, T.H., Sorek, H., Wemmer, D.E., and Somerville, C.R. (2016). Cellulose Deficiency Is
Enhanced on Hyper Accumulation of Sucrose by a H+-Coupled Sucrose Symporter. Plant
Physiol. *171*, 110–124.

87.  Yu, F., Qian, L., Nibau, C., Duan, Q., Kita, D., Levasseur, K., Li, X., Lu, C., Li, H., Hou, C., et al.
(2012). FERONIA receptor kinase pathway suppresses abscisic acid signaling in Arabidopsis
by activating ABI2 phosphatase. Proc. Natl. Acad. Sci. U. S. A. *109*, 14693–14698.

88.  Zhu, S., Martínez Pacheco, J., Estevez, J.M., and Yu, F. (2020). Autocrine regulation of root
hair size by the RALF-FERONIA-RSL4 signaling pathway. New Phytol. *227*, 45–49.

89.  Jones, D.L., Healey, J.R., Willett, V.B., Farrar, J.F., and Hodge, A. (2005). Dissolved organic
nitrogen uptake by plants—an important N uptake pathway? Soil Biol. Biochem. *37*, 413–
423.

90.  Forsum, O., Svennerstam, H., Ganeteg, U., and Näsholm, T. (2008). Capacities and
constraints of amino acid utilization in Arabidopsis. New Phytol. *179*, 1058–1069.

91.  O'Leary, B.M., Khim Oh, G.G., Lee, C.P., and Harvey Millar, A. (2020). Metabolite regulatory
interactions control plant respiratory metabolism via target of rapamycin (TOR) kinase
activation. Plant Cell *32*, 666–682.

92.  Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-
time quantitative PCR and the 2-ΔΔCT method. Methods *25*, 402–408.

93.  Grebber, K., and Karabinos, J.V. (1952). A study of the diphenylamine test for aliphatic
nitrocompounds. Journal of Research of the National Bureau of Standards *49*, 163-166.