Cell surface receptor kinase FERONIA linked to nutrient sensor TORC1 signaling controls root hair growth at low temperature in *Arabidopsis thaliana*

Javier Martínez Pacheco[1,*], Limei Song[2,*], Victoria Berdion Gabarain[1], Juan Manuel Peralta[1], Tomás Urzúa Lehuedé[3,4], Miguel Angel Ibeas[3,4], Sirui Zhu[2], Yanan Shen[2], Feng Yu[2,†] & José M. Estevez[1,3,4,†]

[1]Fundación Instituto Leloir and IIBBA-CONICET. Av. Patricias Argentinas 435, Buenos Aires C1405BWE, Argentina.

[2]State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Biology, and Hunan Key Laboratory of Plant Functional Genomics and Developmental Regulation, Hunan University, Changsha 410082, P.R. China.

[3]Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad Andres Bello.

[4]ANID - Millennium Science Initiative Program - Millennium Institute for Integrative Biology (iBio) and Millennium Nucleus for the Development of Super Adaptable Plants (MN-SAP), Santiago, Chile.

*Co-first authors.

[†]Correspondence should be addressed. Email: feng_yu@hnu.edu.cn (F.Y) and jestevez@leloir.org.ar (J.M.E).

Abstract

Root hairs (RH) are excellent model systems for studying cell size regulation since they elongate
several hundred-fold their original size. Their growth is determined both by intrinsic and
environmental signals. Although nutrients availability in the soil are key factors for a sustained
plant growth, the molecular mechanisms underlying their sensing and downstream signaling
pathways remains unclear. Here, we identified the low temperature (10°C) triggers a strong RH cell
elongation response involving the cell surface receptor kinase FERONIA (FER) and the nutrient
sensing role of TOR Complex 1(TORC1). We found that FER is required to perceive limited nutrients
availability caused by low temperature, to interacts with and activate TORC1-downstream
components to trigger RH growth. We also found that nitrate availability could mimic the RH
growth response at 10°C via an unknown specific mechanism involving the NRT1.1 transceptor.
Our findings reveal a new molecular mechanism by which a central hub composed by FER-TORC1
might be involved in the control of RH cell elongation under low temperature.

Abstract Word counts 164

Text Word counts 3,377

Figures 1-5

Figure S1-S3

Introduction

43
44

Root hairs (RHs) are single cells that develop as a cylindrical protrusion from the root epidermis in
a developmentally regulated manner. RHs are able to expand in a polar manner several hundred
times their original size in a couple of hours to reach water-soluble nutrients in the soil, to promote
interactions with the local microbiome, and to support the anchoring of the plant. RH growth is
controlled by the coordination of a plethora of environmental and endogenous factors[1]. Recently,
an autocrine mechanism of RH growth was described where RALF1-FER complex recruits and
phosphorylates the early translation factor eIF4E1 to enhance protein synthesis of specific mRNAs,
including the RH growth master regulator: *ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4)*[2]. It is known that
macronutrients availability are key factors that promotes rapid RH growth[1,3,4]. Although nitrate is
detrimental to RH growth, their mechanisms of action remain unknown[5–7] . The
CHLORINA1/NITRATE TRANSPORTER1.1 (CHL1/NRT1.1) was proposed as a transceptor (transporter
and receptor) for nitrates[8–10] as well as for auxin transport[11]. When the position T101 of NRT1.1 is
phosphorylated, NRT1.1 becomes a high-affinity NO_3^- (nitrate) transporter, giving this protein a
dual-affinity capability[12–15] . Recently, we have shown that at low temperature conditions (at 10
°C), nutrients availability in the media is reduced and triggers an exacerbated RH growth that is
suppressed if nutrients are increased[16]. Although great advances were achieved in our
understanding on how RH growth occurs at low temperature, it is still unclear how nutrient
availability derived from low temperature is perceived in the RH cells and which are the non-
transcriptional responses that control RH growth.

TARGET OF RAPAMYCIN (TOR) is an evolutionarily conserved Ser/Thr protein kinase in all
eukaryotic organisms that acts as a central growth regulator controlling metabolism and protein
synthesis[17–19] .TOR is found in at least two distinct multiprotein complexes called TOR Complex 1
and 2 (TORC1 and TORC2), although only TORC1 was experimentally validated in plants[20,21] . The
Arabidopsis TORC1 complex is encoded by one *TOR* gene (*At*TOR)[22] ,two *REGULATORY-ASSOCIATED*
PROTEIN OF TOR (RAPTOR A and B) genes[23–28], and two *LETHAL WITH SEC THIRTEEN 8 (LST8)*
genes[29]. In contrast to the embryo-lethal *tor*-null mutant lines, *raptor1b* and *lst8-1* mutants are
viable but show significant growth issues and developmental phenotypes[30]. Some canonical
downstream targets of TOR are conserved in plants, such as the S6 kinase (S6K) which stimulates
protein translation[25,31–34]. In plants, TORC1 complex plays a key role during many stages of the plant
life cycle by controlling both anabolic and catabolic downstream processes. In addition, TORC1 is
activated by nutrient availability and inactivated by stresses that alter cellular homeostasis[19,35–39].
The TORC1 complex senses and integrates signals from the environment to coordinate
developmental and metabolic processes including hormones (e.g., auxin), several nutrients (e.g.,
Nitrogen and Sulfur), amino acids and glucose[17,28,40–44]. However, the underlying molecular
mechanisms by which TORC1 operates at a single plant cell level has yet to be elucidated. Although

the sensing and uptake of nutrients by RHs is crucial for the plant productivity, the molecular
mechanisms underlying these processes remain unknown. Until now, few upstream regulators of
TOR have been described in plants. Among them, there is a single subfamily of Rho GTPases (ROPs)
involved in the spatial control of cellular processes by signaling to the cytoskeleton and vesicular
trafficking. Particularly, ROP2 GTPase activates TOR in response to auxin and nitrogen signals[44–46].
In addition, ROP2 is described as a monomeric GTP-binding protein that participates in many
cellular signaling processes, including the polar growth of pollen tubes and root hairs[47–50]. ROPs
activation is regulated by ROP guanine nucleotide exchange factors (ROP-GEFs) which interact with
several receptor-like kinases (RLKs) including the *Cr*RLK family member: FERONIA (FER)[48]. During
the cell elongation of RH, ROP2 is activated by the ROPGEF1-FER interaction. This process is also
regulated by ROP-GEF4 and ROP-GEF10[48,51] .Therefore, ROP2 (among other ROPs) may bridge the
membrane receptor FER with TORC1 activation, although this was not yet demonstrated.
93
FERONIA was also linked to carbon/nitrogen balance during the plant growth[52]. Even more, it was
shown that the cytoplasmic kinase domain of FER and its partner RIPK (RPM1-induced protein
kinase) interact with TOR and RAPTOR-A forming a complex to positively modulate TOR pathway
under nutrient starvation. Currently, it is clear that plant growth can be affected via FER-TOR axis
since they act early by linking amino acid and/or nutrients signaling with global protein synthesis[52].
Existing evidences suggest that nutrient mediated sensing at the cell surface level triggers a
response thought the TOR signaling pathway. However, it is still unknown which are the
environmental signals activated in RH by the FER-TORC1 pathway. Taking into account that
previously we described that a low temperature stress is able to trigger an exacerbated RH
growth[16,53], this stress condition was used as a proxy to investigate which are the environmental
signals that activate RH cell elongation. Our research reveals a novel mechanism in which FER and
TORC1 are necessary to regulate RH growth in *Arabidopsis* in the context of the nutrient availability
(nitrate) as cause of low temperature stress.
107
108
109   **Results**
110
FER is required to trigger a strong RH growth response at low temperature. Here, we asked which
might be the cell surface protein involved in perceiving and/or transducing the low temperature
stimulus. Since FER was shown as an important hub between the cell surface signaling with
downstream processes during RH growth[2,48] ,we tested if the *fer-4* null mutant and *fer-5* (a
truncated version with shortened kinase domain (KD)) are able to respond to low temperature
stimulus. Both *fer* mutants failed to trigger RH growth regardless the temperature (**Figure 1A**). On
the contrary, the related CrRLK1L ERULUS (ERU), previously linked to RH growth and cell wall
integrity processes with a severe short cell elongated RH phenotype at 22°C[54,55], was able to react

to low temperature and triggered a RH growth response although to a lower extent than Wt Col-
0. This indicates that FER, but not ERU, might be involved specifically in this growth response
despite being phylogenetically related[55] .Then, a fluorescent translational reporter line of FER
(FER*p*:FER-GFP), was used to study *FER* expression in RHs in two growth stages, an initial and a later
stage. FER is clearly activated under low temperature stimulus at the initiation stage of RH growth
and to a lower extent at a later stage (**Figure 1B**). Finally, phosphorylated and non-phosphorylated
levels of FER (FER-p/FER) were quantified at both temperatures. Under low temperature, almost
half of the FER protein levels was present as a non-phosphorylated form and after 3 days of growth,
almost all FER protein was present as a phosphorylated version (**Figure 1C**). This indicates that low
temperature not only increases FER protein levels in growing RHs at the plasma membrane but
also promotes a complete phosphorylation of FER protein, possibly enhancing the interaction with
putative partners and triggering the activation of downstream signaling components of RH polar
growth.
132
TORC1 pathway is involved in low temperature induced RH growth. Recently, it was shown that
FER-RIPK interacts with TOR-RAPTOR and phosphorylates it leading to the TOR pathway activation
in the context of nutrient perception and global metabolism regulation. We then asked if TOR might
be also involved in the RH growth process under low temperature. In the first place, a β-estradiol
(es)-induced TOR knockdown (*tor-es,* es-induced RNAi silencing of TOR) line[56] showed short RHs as
previously has been reported[33] and notably, a lack of response to low temperature stimulus (**Figure**
2A). Secondly, in a similar manner, inhibition of TOR kinase activity with AZD-8055[57], a novel ATP-
competitive inhibitor of mTOR kinase activity, abolished the RH growth at both 22°C and low
temperature (**Figure 2B**). On the other hand, overexpression of TOR enhanced RH growth at both
temperatures assayed (**Figure 2C**). These results lead us to suggest that the TOR pathway is
involved in the RH low temperature growth response and might operates downstream of FER. At
the transcriptional level, TOR expression levels are upregulated in roots up to three-folds at 10 °C
vs 22 °C (**Figure S1**). To deepen in this study, we tested if the growth response at low temperature
in RHs is also affected in mutants of the TORC1 pathway and some downstream components. All
the mutants *(raptor1b, rps6b,* and *lst8-1)* were all unable to respond to the low temperature
treatment but their RHs phenotypes showed a broad spectrum of responses in terms of cell length
with respect to the Wt Col-0 (**Figure 2C**). For instance, *raptor1b* behaved similarly to *tor-es* mutant,
while *rps6b* had an intermediate phenotype and *lst8* was comparable to Wt Col-0 (at 22°C) (**Figure**
2C). In addition, the overexpression of TOR and the S6 kinase 1 (S6K1), as a direct downstream
target of TOR, showed enhanced RH growth at 22 °C. Since previous reports showed that
phosphorylation of the ribosomal S6K1 can be used to monitor TOR protein kinase activity in
plants[33,56] ,we measure the ratios of S6K-p/S6K. Lower levels of S6K-p/S6K were detected for those
lines with strong short RH phenotype (mutants in components of the TORC1 complex and in FER
protein) while those with longer RHs (overexpression of S6K1 and TOR) showed higher ratios

(**Figure 2D, 3D**). These results clearly indicate that TORC1 and some downstream components (e.g.
S6K) promote RH growth under low temperature.

FER regulates TOR polar localization in RH and activation of TORC1 pathway at low temperature.
Since TORC1 activation under a plethora of stimuli usually triggers the phosphorylation of the
downstream factor S6K (S6K-p), we asked if low temperature responses regulated by FER might
control this output. With this in mind, we investigated the molecular mechanisms by which TORC1-
S6K are induced at 10 °C and we tested whether this might be mediated by FER. Since *fer-4* and
tor-es showed a similar RH phenotypes at low temperature (**Figure 2A-B**), possibly indicating that
they may act in the same pathway, we first tested if FER and TOR interaction is enhanced under
10°C vs 22°C. By performing a co-immunoprecipitation (Co-IP) analysis, we found that FER-TOR
protein-protein interaction is enhanced under low temperature (**Figure 3A**). This confirms a direct
interaction between FER and TOR kinase in low temperature RH mediated growth. Then, by
evaluating the immunolocalization of TOR in the growing RHs, we found that FER is required for
the apical accumulation of TOR since TOR lost the polar pattern in *fer-4* root hairs. In addition, the
level of RH tip-localized TOR detected was similar at both, low temperature and control conditions
(22°C) (**Figure 3B-C**). This indicates that in root hairs, TOR is able to localize into the tip and this is
dependent on FER. Then, it was tested if FER is involved in the S6K activation by TOR by measuring
the ratio of S6K-p/S6K proteins under both growth conditions in Wt Col-0, *fer-4*, *fer-5*, and FER^{K565R},
which has been shown to abolish FER autophosphorylation and transphosphorylation in an *in vitro*
kinase activity assay[58,59] ,although some activity might remain[60]. Interestingly, low temperature
enhanced S6K-p levels after 3 days of growth in Wt Col-0 at low temperature but this was partially
suppressed in all *fer* mutants tested (**Figure 3D**). This suggests that FER controls the level of TORC1
activation under low temperature in RH. Collectively, these results indicate that TOR localization in
the RH tip and TORC1 activation is dependent on FER and enhanced at low temperature.

Nitrate perception and transport mediated by NRT1.1 controls RH growth at low temperature.
Previous studies have been shown that nitrogen and specifically nitrates are important nutrient for
TOR signaling[44,61] . Since low temperature growth conditions reduce the availability of nutrients in
the media and trigger a strong response in RH growth[16,53] , we decided to test if nitrate signaling
pathway might be involved. We found that high levels of nitrate (18.8 mM) were able to repress
low temperature mediated RH growth while low levels of nitrate (0.5 mM) did not affect growth
under this temperature (**Figure 4A**). This clearly indicated that nitrate is one of the major growth
signals that activate RH cell elongation at low temperature where there is a low nutrient mobility
environment. Then, we tested if the dual affinity nitrate transporter and sensor NRT1.1 is involved
in this low temperature RH growth response. The NRT1.1 null mutants *chl1-5*, *chl1.9* (NRT1.1
harbors a substitution P492L that suppresses its root nitrate uptake activity[8] ,as well as the
catalytically inactive *CHL1^{T101D}* version showed a strong RH growth response despite the

temperature conditions (**Figure 4B**), similarly to *S6K1-OE* and *TOR-OE* lines (**Figure 2C**), but to a
lower extent in terms of RH cell elongation. As expected, similar levels of S6K-p/S6K were detected
in the *chl1-9* line at both temperatures (**Figure 4C**). When these NRT1.1 mutants were grown under
high (18.8 mM) or low (0.5mM) nitrate concentrations at both temperatures (at 22°C and at 10°C),
a similar RH phenotype was detected between both nitrate conditions either at 22 °C or 10 °C
(**Figure 4D**). On the contrary, as expected Wt Col-0 RH were sensitive to both, low temperature and
nitrate levels. Overall, this suggested a key role of NRT1.1 in the perception of changes in nitrate
levels and in the regulation of the downstream TORC1-S6K activation. At this point, we
hypothesized that FER might be a good candidate to controls NRT1.1 activity under low
temperature (and low nitrate), but this remains to be tested.
205

ROP2 is required for low temperature RH growth. Previous studies have shown a key role of ROP2
in the regulation of RH polarity and elongation[49,62,63], as an important molecular link between FER
and downstream components involved in RH growth[48]. In addition, ROP2 promotes the activation
of TOR and its relocation to the cell periphery and induces the downstream translation pathway[45].
More importantly, ROP2 was shown to integrate diverse nitrogen and hormone signals for TOR
activation[44]. In the first place, *rop2* abolishes the differential growth responses at 10°C and
produces a very short RH phenotype (**Figure 5A**). While high levels of TOR are able to enhance RH
growth regardless of the temperature treatments (**Figure 2C**), the constitutively active ROP2
version (CA-ROP2) is able to repress TOR activation (**Figure 5A**). As expected, 35S-TOR-GFP (TOR
OE) with constitutive long RHs at both temperatures showed much higher levels of S6K-p/S6K than
Wt Col-0 while the presence of CA-ROP2 (in TOR OE/CA-ROP2) was able to reduce the levels of S6K-
p/S6K linked to a diminished RH growth (**Figure 5B**). Then, we asked if TOR partition between
plasma membrane/cell surface-cytoplasmic localization might be changed by the effect of low
temperature (equal to reduced nutrient environment) and if this is changed by the presence of CA-
ROP2 (**Figure 5C**). Low temperature triggers a higher cell surface TOR localization while the
presence of CA-ROP2 abolishes this effect. Taken together, we observed a direct effect of CA-ROP2
version in the TOR localization at the RH tip, RH growth responses and TORC1 activation. Then, we
tested if a treatment with exogenous auxin (100 nM IAA) that is able to trigger a strong RH growth
response and ROP2 activation, might be able to bypass silenced TOR growth inhibition effect in RH
at both, room and low temperatures (**Figure S2**). In this case, auxin treatment was able to partially
revert RH growth inhibition under TOR silenced line. Because of ROP2 activity and other possible
unknown effectors, auxin is able to partially rescue RH growth when TOR is inhibited, highlighting
that other pathway might be still being active downstream auxin independently of TORC1 to trigger
RH cell elongation.
230
231

Discussion

Growth and development of plants and animals are based on nutrient and hormonal signaling that
constitute as the main regulatory networks in eukaryotes. Unraveling the functions of the
regulatory hubs and their detailed molecular mechanisms are critical for understanding the
deepened modes of regulation in these signaling pathways. Our study uncovers a new molecular
mechanism by which plants use FER-ROP2-TORC1 signaling pathway to control cell elongation in
RH under nutrient starvation induced by low temperature stress. Here we showed that the cell
surface FER receptor is able to regulate TOR apical localization and further downstream activation,
both controlling S6K phosphorylation linked to RH growth. Based on the results obtained, here we
propose a model in which the FER-ROP2-TORC1 axis might be able to regulate the RH growth under
low temperature with remarked attention to variable nitrate conditions (**Figure S3**). It is based on
the mutants tested in this signaling pathway, FER-TOR enhanced interactions, TOR polar
localization in RHs and S6K-p/S6K as a TORC1 activation response. Changes in nitrate
concentrations in the media mimicked this response at low temperature and mutants on the
NRT1.1 transceptor are insensitive at RH growth level regardless the temperature or nitrate levels.
We hypothesize that low nitrate in the media might rapidly induce the levels of mature-active
RALF1 (and possibly other RALF peptides), which might then activate FER-ROP2-TORC1 pathway
and the downstream fast cell elongation effect to search for further nutrients sources. This
autocrine mechanism dependent on RALF1-FER growth activation has been recently demonstrated
for RH[2,64] ,but not in a context that involves low temperature/low nitrate and TORC1 pathway.
Finally, several questions in our proposed model (**Figure S3**) remain to be answered in future
studies. How does low nitrate concentration (low temperature) triggers FER-TORC1 activation?
Does RALF1 and other RALFs respond to low nitrate to activate FER? Does FER activate NRT1.1
directly? Recently, it was shown that FER-regulated ROP2 triggers a new mechanism for the
negative regulation of rhizosphere microorganisms such as *Pseudomonas*[65] .Although not tested
here, FER-ROP2-TORC1 signaling coupled to enhanced RH growth under low temperature (low-
nutrients) might be linked in root growth in specific soil conditions to select favorable microbiota
in the soil. This pathway may integrate complex signals from soil nutrients and microbiota in the
rhizosphere to plant root cell growth mechanism.

Figure 1. High levels of FER in its phosphorylated form is required to trigger the low temperature RH growth.

(A) Scatter-plot of RH length of Col-0, of *fer-4*, *fer-5*, and *eru* mutants grown at 22°C or at 10°C. RH growth is enhanced at low temperature in Wt Col-0 and *eru* mutant but not in the *fer-4* and *fer-5* mutants. Each point is the mean of the length of the 10 longest RHs identified in a single root. Data are the mean ± SD (N=10-20 roots), two-way ANOVA followed by a Tukey–Kramer test; (*) $p < 0.05$, (***) $p < 0.001$, NS=non-significant. Results are representative of three independent experiments. Asterisks indicate significant differences between Col-0 and the corresponding genotype at the same temperature. Representative images of each genotype are shown below. Scale bars= 500 µm.

(B) Confocal images of the reporter line of FER in two growth stages of RH cells, short RHs (on top) and long RHs (on the bottom) at 22°C *vs* low temperature (10°C). Below, quantification of the GFP fluorescence intensity in RH tip. Fluorescence AU data are the mean ± SD (N=11-20 root hairs), two-way ANOVA followed by a Tukey–Kramer test; (*) $p < 0.05$, (***) $p < 0.001$. Results are representative of two independent experiments. Asterisks on the graph indicate significant differences. Scale bars: 100 µm.

278   (C) Phosphorylation levels on FER (FER-p in *p*FER:FER-GFP) increases after 3 days at low
279   temperature in roots. Phosphorylated levels of FER were analyzed by ImageJ.

280
281
Figure 2. TORC1 signaling pathway is required for low temperature triggered RH growth.

(A) Scatter-plot of RH length of Col-0 and *tor-es* line grown at 22°C or at 10°C. Differential growth
of RH at low temperature is suppressed in the estradiol inducible *RNAi TOR* line. Each point is the
mean of the length of the 10 longest RHs identified in a single root. Data are the mean ± SD (N=7-
10 roots), two-way ANOVA followed by a Tukey–Kramer test; (***) p <0.001, NS=non-significant.
Results are representative of three independent experiments. Asterisks indicate significant
differences. Representative images of each line are shown below. Scale bars= 500 μm.
(B) Differential growth of RH at low temperature is abolished in the Col-0 treated with 250nM of
TOR inhibitor, AZD-8055. Each point is the mean of the length of the 10 longest RHs identified in a
single root. Data are the mean ± SD (N=7 roots), two-way ANOVA followed by a Tukey–Kramer test;
(***) p <0.001, NS=non-significant. Results are representative of three independent experiments.
Asterisks indicate significant differences. Representative images of each line are shown below.
Scale bars= 500 μm.
(C) RH cell elongation of TORC1 and downstream TORC1 signaling pathway mutants under low
temperature. Each point is the mean of the length of the 10 longest RHs identified in a single root.
Data are the mean ± SD (N=7-12 roots), two-way ANOVA followed by a Tukey–Kramer test; (*) p
<0.05, (**) p <0.01, (***) p <0.001, NS=non-significant. Results are representative of three
independent experiments. Asterisks indicate significant differences. Representative images of each
line are shown on the right. Scale bars= 500 μm.
(D) Analysis of the phosphorylation state of S6K (S6K-p/S6K ratio) in Col-0 and *S6K-OE* line. A
representative immunoblot is shown of a three biological replicates. S6K-p/S6K ratio was analyzed
by ImageJ.

Figure 3. **FER controls apical TOR localization and subsequent activation thus increasing phosphorylation of S6K under low temperature.** (A) Enhanced FER-TOR interaction at low temperature in roots by Immunoprecipitation (IP). A representative experiment of three replicates is shown.

(B) On the top, TOR immunolocalization in RHs is FER-dependent. Scale bar=20 μm

(C) Apical and total TOR signal quantification in RHs showed in (B). A ROI at the RH tip of the fluorescent signal or all the RH TOR-derived fluorescent signal was measured. Fluorescence AU

data are the mean ± SD (N=10-15 root hairs), two-way ANOVA followed by a Tukey–Kramer test;

(***) p < 0.001, NS= non significant. Results are representative of two independent experiments.

Asterisks on the graph indicate significant differences.

(D) Analysis of the phosphorylation state of S6K in Wt Col-o, *fer-4, fer-5*, and *FER^K565R* mutants

Arabidopsis roots. Phosphorylation of S6K (S6K-p) is enhanced at low temperature and requires

FER active kinase. A representative immunoblot is shown of a three biological replicates. S6K-p/S6K

ratio was analyzed by ImageJ.

320
321

Figure 4. Nitrate acts as a RH growth signal perceived by NRT1.1 and triggered by TORC1 pathway at low temperature growth.

(A) High nitrate (18.8 mM) suppresses low temperature enhanced RH growth in Wt Col-0 plants. . Each point is the mean of the length of the 10 longest RHs identified in a single root. Data are the mean ± SD (N=10 roots), two-way ANOVA followed by a Tukey–Kramer test; (*) p <0.05, (***) p

<0.001. Results are representative of three independent experiments. Asterisks indicate significant
differences. Representative images of each line are shown below. Scale bars= 500 µm.
(B) Low temperature RH growth is regulated by the nitrate sensor NRT1.1 (CHL1). Scatter-plot of
RH length of Col-0 in low or high nitrate conditions(right); RH length of Col-0 and NRT1.1 mutants,
(left) grown at 22°C or 10°C. Each point is the mean of the length of the 10 longest RHs identified
in a single root (N=15-20). Each point is the mean of the length of the 10 longest RHs identified in
a single root. Data are the mean ± SD (N= 10-15 roots), two-way ANOVA followed by a Tukey–
Kramer test; (***) p <0.001, NS= non-significant. Results are representative of three independent
experiments. Asterisks indicate significant differences. Representative images of each line are
shown below. Scale bars= 500 µm.
(C) Analysis of the phosphorylation state of S6K in Wt Col-0 and *chl1-9* mutant roots. A
representative immunoblot is shown of a three biological replicates. S6K-p/S6K ratio was analyzed
by ImageJ.
(D) RH growth responses under high and low nitrate (18.8 mM and 0.5mM, respectively) in Wt Col-
0, *chl1-5*, *chl1-9* mutants, and $CHL1^{T101D}$ grown at 22°C or 10°C. Each point is the mean of the length
of the 10 longest RHs identified in a single root. Data are the mean ± SD (N= 10-15 roots), two-way
ANOVA followed by a Tukey–Kramer test; (***) p <0.001, NS= non-significant. Results are
representative of three independent experiments. Asterisks indicate significant differences.

Figure 5. ROP2 is required to triggers RH growth at low temperature. (A) ROP2 is required for RH at low temperature meanwhile *rop2* and CA-ROP2/35S-TOR block RH growth. Each point is the mean of the length of the 10 longest RHs identified in a single root. Data are the mean ± SD (N= 10-15 roots), two-way ANOVA followed by a Tukey–Kramer test; (**) p <0.01, (***) p <0.001. Results

are representative of three independent experiments. Asterisks indicate significant differences.

Representative images of each line are shown below. Scale bars= 500 µm.

(B) Analysis of the phosphorylation state of S6K in *TOR-OE* and *TOR-OE / ROP2-CA Arabidopsis*

roots. Phosphorylation of S6K (S6K-p) is enhanced in *TOR-OE* and suppressed in *TOR-OE*/ROP2-CA.

A representative immunoblot is shown of a three biological replicates. S6K-p/S6K ratio was

analyzed by ImageJ.

(C) Quantification of TOR-GFP localization in the cell surface versus cytoplasmic localization in the

RH tip in Wt Col-0 and in CA-ROP2 background. Fluorescence AU data are the mean ± SD (N=12-15

root hairs), two-way ANOVA followed by a Tukey–Kramer test; (*) $p < 0.05$, (**) $p < 0.01$, NS= non-

significant. Results are representative of two independent experiments. Asterisks on the graph

indicate significant differences. Representative images of each line are shown on the right. Scale

bars= 1 µm.

TOR expression in Col-0 roots

364
365
Figure S1. Quantitative PCR of *TOR* expression levels in Col-0 roots grown at 22°C and 10°C. *ACT2*
expression was used for normalization of gene expression. Three biological replicates and three
technical replicates per experiment were performed.

Figure S2. TOR-pathway on RH growth is partially dependent on auxin.

RH length of non-treated and treated Col-0 and TOR-RNA*i* line (*tor-es*) with 100nM IAA at 22°C *vs*

10°C. Each point is the mean of the length of the 10 longest RHs identified in a single root. Data are

the mean ± SD (N= 10-15 roots), two-way ANOVA followed by a Tukey–Kramer test; (***) p <0.001,

NS=non-significant. Results are representative of three independent experiments. Asterisks

indicate significant differences.

378
379
Figure S3. **Proposed model of FER-ROP2-TORC1 pathway at low temperature-nitrate that triggers**
cell elongation in RHs. On the left, several mutants characterized in this work. On the right, the
model based on the current evidence shown in this work and together with previous works. Low
temperature triggers a low nitrate environment that leads to FER activation, enhanced FER-TOR
interaction and TORC1 activation. This promotes cell elongation of RHs. This response also requires
ROP2 that might bridge FER with TORC1. Several questions (?) remain to be answered in future
studies such as how low nitrate (low temperature) triggers FER-TORC1 activation? Does FER
activate NRT1.1 directly? Do RALF1 and other RALFs respond to low nitrate to activate FER?.

Experimental Procedures
389
Root hair phenotype. Seeds were surface sterilized and stratified in darkness for 3 days at 4°C.
Then grown on ½ strength MS agar plates supplemented with MES (Duchefa, Netherlands), in a
plant growth chamber at 22°C in continuous light (120 µmol.sec-1.m-2) for 5 days at 22°C plus 3
days at 10°C (moderate-low temperature treatment) or for 8 days at 22°C as control. For
quantitative analyses of RH phenotypes, 10 fully elongated RH from the maturation zone were
measured per root under the same conditions from treatment and control. Measurements were
made after 8 days. Images were captured using an Olympus SZX7 Zoom Stereo Microscope
(Olympus, Japan) equipped with a Q-Colors digital camera and Q Capture Pro 7 software (Olympus,
Japan). Results were expressed as the mean ± SD using the GraphPad Prism 8.0.1 (USA) statistical
analysis software. Results are representative of three independent experiments, each involving 7–
20 roots.
401
Plant Material and Growth Conditions. Plants were grown on agar plates in a plant chamber at
22 °C for 7 days at 24 hs light (120 µmol.seg-1.m-2). *A. thaliana* Col-0 ecotype was used as a wild-
type plant. A Murashige & Skoog-based media without N, P nor K (M407, PhytoTechnology
Laboratories, https://phytotechlab.com/) was used as the growth base medium under sterile
conditions. To test the low and high nitrate response, 0.5X M407 media was supplemented with
0.8% plant agar (Duchefa, Netherlands); 1.17mM MES,0.625 mM KH_2PO_4 monobasic and 0.5mM
or 18.8 mM KNO_3, respectively. Plants were treated for the indicated period in accordance with
prior conditions. *chl1-5*, *chl1-9 and T101D* mutants were kindly donated by Dr Yi-Fang Tsay. *tor-es*
was kindly donated by Dr Ezequiel Petrillo, *lst8-1*, *rps6b*, *raptor1b* and overexpressing line S6K1
kindly donated by Dr Elina Welchen and *35S::GFP-TOR* and *35S::GFP-TOR/CA-ROP2* kindly donated
by Dr. Lyubov Ryabova.
413
Pharmacological Treatments. Plants were grown on solid 0.5x MS medium plates at 22°C for 5 days
at 24 hs light (110 µmol.seg-1.m-2). Then transferred to plates containing solid 0.5x MS, according
with the specific treatment, supplemented with 100nM IAA (auxin treatment), 250nM AZD-8055
(TOR inhibition) or 10 µM β-estradiol (TOR RNA*i* estradiol inducible line). Treatments were
incubated at 22 °C for 3 days at 24 hs light (110 µmol.seg^{-1}.m^{-2}) and then at 10 °C for 3 days at 24
hs light (110 µmol.seg^{-1}.m^{-2}).
420
Confocal Microscopy. Confocal laser scanning microscopy for the lines *pFER::FER-GFP, 35S::GFP-*
TOR and 35S::GFP-TOR/CA-ROP2, was performed using a Zeiss LSM 710 NLO microscope (Zeiss,
Germany) (Excitation: 488 nm argon laser; Emission: 490-525 nm, EC Plan Neofluar 40X/1.3 Oil or
a Zeiss Plan-Apochromat 63X/1.4 Oil objectives). GFP signal at the cell surface and cytoplasm of the
RH tip were quantified using ROIs with the ImageJ software. Fluorescence AU were expressed as

the mean ± SD using the GraphPad Prism 8.0.1 (USA) statistical analysis software. Results are
representative of two independent experiments, each involving 10–15 roots and approximately,
between 10 to 20 hairs per root were observed.

Co-IP assay. For Co-IP assays using A/G agarose and an anti-TOR antibody[33], 30 µL of A/G beads
(Thermo Fisher Scientific Inc., 20421) was resuspended and washed three times using NEB buffer
(20 mM HEPES [pH 7.5], 40 mM KCl, 5 mM MgCl$_2$) before adding 8 µL of anti-TOR antibody or
preimmune serum as a negative control [33] in a total volume of 500 µL of NEB buffer, followed by
incubation for 4 h at 4 ˚C. Col-0 seedlings were first grown at 22 ˚C for 5 days then transferred to
22 ˚C or 10 ˚C for 2 days or 3 days. For protein extraction from plants, the collected materials were
ground to a fine powder in liquid nitrogen and solubilized with NEB-T buffer (20 mM HEPES [pH
7.5], 40 mM KCl, 5 mM MgCl$_2$, 0.5% Triton X-100) containing 1 × protease inhibitor cocktail (Thermo
Fisher Scientific Inc., 78430) and 1 × phosphatase inhibitor (Thermo Fisher Scientific Inc., 78420)
and incubated for 1 h on the ice. The extracts were centrifuged at 16,000 g at 4 ˚C for 15 min, and
the resultant supernatant was incubated with prepared antibody-beads from the above step. After
overnight incubation at 4 ˚C with rotation, the agarose beads were washed five times with the NEB
buffer and eluted with elution buffer (0.2 M glycine, 0.5% Triton X-100, pH 7.5). Anti-FER and anti-
TOR antibodies[33] were used for immunoblotting to detect the immunoprecipitates.

TOR immunolocalization. Col-0 and *fer-4* seedlings were first grown at 22 ˚C for 4 days before
transferred to 22 ˚C or 10 ˚C for 2 days and treatment with or without 1 µM RALF1 for 30min. Then
seedlings were collected and incubated for 10 min under vacuum (0.05 MPa) in phosphate-
buffered saline (PBS) containing 4% paraformaldehyde and 0.1% Triton X-100. Seedlings were
washed gently three times (10 min for each wash) in PBS and then the cell wall was digested in 2%
Driselase (Sigma, D8037) in PBS for 18 min at 37 ˚C and washed five times with PBS. The
permeability of the seedlings was increased by incubating them in 3% IGEPAL CA-630 (Sigma,
18896) and 10% DMSO in PBS for 18 min, followed by washed three times with PBS. Seedlings were
incubated in 2% bovine serum albumin (BSA) (Ameresco, 0332) in PBS for 1.5 h and then incubated
with primary TOR antibody[33] (antibody diluted 1:600 in 2% BSA) for overnight at 4 ˚C. The seedlings
were washed with PBS for five times. Fluorophore-labeled secondary antibody (goat–mouse
secondary antibody, diluted 1:600 in 2% BSA) was incubated with the samples at 37 ˚C for 5 h in
the dark. Seedlings were washed five times with PBS before observation. Fluorescent signal
detection and documentation were performed using a Nikon confocal laser scanning microscope
with a 560-nm band-pass filter for IF555 detection.

S6K-P/S6K immunoblotting detection. For immunological detection of S6K-P and S6K1/2, total
soluble proteins were extracted from 50 mg of plant materials grown as indicated previously with
100 µL 2 × Laemmli buffer supplemented with 1% Phosphatase Inhibitor Cocktail 2 (Thermo Fisher

Scientific Inc., 78430). Proteins were denatured for 10 min at 95 ˚C and separated on 10% or 8%

SDS-PAGE. Rabbit *At*TOR polyclonal antibodies (Abiocode, R2854-2), rabbit polyclonal S6K1/2

antibodies (Agrisera, AS121855), and S6K1-p (phospho T449) antibody (Abcam, ab207399) were

used for immunoblotting.

Quantitative PCR (qPCR). Total root RNA was extracted from plantlets grown *in vitro* at 22 ˚C and

10 ˚C using the RNeasy®Plant Mini Kit (QIAGEN, Germany). One microgram of total RNA was

reverse transcribed using an oligo(dT)$_{20}$ primer and the Super Script™ IV RT (Invitrogen,USA)

according to the manufacturer's instructions. cDNA was diluted 20-fold before PCR. *q*PCR was

performed on a LightCycler®480 Instrument II (Roche, USA) using 2 μL of 5X HOT

FIREPol®EvaGreen® qPCR Mix Plus(no ROX) (Solis BioDyne, Estonia), 2 μL of cDNA, and 0.25 μM of

each primer in a total volume of 10 μL per reaction. ACT2 (AT3G18780) gene was used as reference

for normalization of gene expression levels (ACT2 primers, F: GGTAACATTGTGCTCAGTGGTGG R:

CTCGGCCTTGGAGATCCACATC; TOR primers, F: GAAGATGAAGATCCCGCTGA R:

GCATCTCCAAGCATATTTACAGC[45]). The cycling conditions were: 95 °C for 12 min., 35 cycles of 95 °C

for 15 sec., 60 °C for 1 min. and finally a melting curve from 60 °C to 95 °C (0.05°/sec). Data were

analyzed using the $\Delta\Delta C_t$[66] method and LightCycler®480 Software, version 1.5 (Roche). Two

independent experiments with three biological and three technical replicates per experiment,

were performed.

Acknowledgements

We would like to thank Elina Welchen, Ezequiel Petrillo, Lyubov A. Ryabova, Yi-Fang Tsay, Kriss

Vissenberg for the seed lines. We thank NASC (Ohio State University) for providing T-DNA lines

seed lines. J.M.E. is investigator of the National Research Council (CONICET) from Argentina. M.I.

is supported by ANID FONDECYT POSTDOCTORADO [grant 3220138]. This work was supported by

grants Natural Science Foundation of China (NSFC-32001974 and NSFC-31871396) to F.Y. and from

ANPCyT (PICT2017-0066, and PICT2019-0015), by ANID – Programa Iniciativa Científica Milenio

ICN17_022, NCN2021_010 and Fondo Nacional de Desarrollo Científico y Tecnológico [1200010] to

J.M.E.

Author Contribution

J.M.P performed most of the experiments, analysed the data and helped in the writing process of

the manuscript. L.S. performed the S6K-p determinations, IP of TOR-FER and immunolocalization

assay of TOR. V.B.G., J.M.P., T.U.I., M.A.I., S-Z., Y.S., helped in the data analysis and writing process

of the manuscript. F.Y. designed research and analysed part of the data and J.M.E. designed

research, analysed the data, supervised the project, and wrote the paper. All authors commented

on the results and the manuscript. This manuscript has not been published and is not under

consideration for publication elsewhere. All the authors have read the manuscript and have

approved this submission.

Competing financial interest

504 The authors declare no competing financial interests. Correspondence and requests for materials

505 should be addressed to F.Y. (feng_yu@hnu.edu.cn) and J.M.E. (Email: jestevez@leloir.org.ar).

REFERENCES

1. Vijayakumar, P., Datta, S. & Dolan, L. ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4) promotes root hair elongation by transcriptionally regulating the expression of genes required for cell growth. *New Phytol.* **212**, 944–953 (2016).

2. Zhu, S. *et al.* The RALF1–FERONIA Complex Phosphorylates eIF4E1 to Promote Protein Synthesis and Polar Root Hair Growth. *Mol. Plant* (2020) doi:10.1016/j.molp.2019.12.014.

3. Yi, K., Menand, B., Bell, E. & Dolan, L. A basic helix-loop-helix transcription factor controls cell growth and size in root hairs. *Nat. Genet.* **42**, 264–267 (2010).

4. Datta, S., Prescott, H. & Dolan, L. Intensity of a pulse of RSL4 transcription factor synthesis determines Arabidopsis root hair cell size. *Nat. Plants* **1**, 1–6 (2015).

5. Vatter, T., Neuhäuser, B., Stetter, M. & Ludewig, U. Regulation of length and density of Arabidopsis root hairs by ammonium and nitrate. *J. Plant Res.* **128**, 839–848 (2015).

6. Canales, J., Contreras-López, O., Álvarez, J. M. & Gutiérrez, R. A. Nitrate induction of root hair density is mediated by TGA1/TGA4 and CPC transcription factors in Arabidopsis thaliana. *Plant J.* **92**, 305–316 (2017).

7. Liu, B., Wu, J., Yang, S., Schiefelbein, J. & Gan, Y. Nitrate regulation of lateral root and root hair development in plants. *Journal of Experimental Botany* vol. 71 4405–4414 (2020).

8. Ho, C. H., Lin, S. H., Hu, H. C. & Tsay, Y. F. CHL1 Functions as a Nitrate Sensor in Plants. *Cell* **138**, 1184–1194 (2009).

9. Gojon, A., Krouk, G., Perrine-Walker, F. & Laugier, E. Nitrate transceptor(s) in plants. *J. Exp. Bot.* **62**, 2299–2308 (2011).

10. Bouguyon, E. *et al.* Multiple mechanisms of nitrate sensing by Arabidopsis nitrate transceptor NRT1.1. *Nat. Plants* **1**, 15015 (2015).

11. Krouk, G. *et al.* Nitrate-regulated auxin transport by NRT1.1 defines a mechanism for nutrient sensing in plants. *Dev. Cell* **18**, 927–937 (2010).

12. Liu, K.-H. & Tsay, Y.-F. Switching between the two action modes of the dual-affinity nitrate transporter CHL1 by phosphorylation. *EMBO J.* **22**, 1005–13 (2003).

13. Forde, B. G. & Walch-Liu, P. Nitrate and glutamate as environmental cues for behavioural responses in plant roots. *Plant, Cell Environ.* **32**, 682–693 (2009).

14. Léran, S. *et al.* A unified nomenclature of nitrate transporter 1/peptide transporter family members in plants. *Trends in Plant Science* vol. 19 5–9 (2014).

15. Wang, W., Hu, B., Li, A. & Chu, C. NRT1.1s in plants: Functions beyond nitrate transport. *J. Exp. Bot.* **71**, 4373–4379 (2020).

16. Moison, M. *et al.* The lncRNA APOLO interacts with the transcription factor WRKY42 to trigger root hair cell expansion in response to cold. *Mol. Plant* **14**, 937–948 (2021).

17. Xiong, Y. & Sheen, J. The role of target of rapamycin signaling networks in plant growth and metabolism. *Plant Physiol.* **164**, 499–512 (2014).

18. Dobrenel, T. *et al.* The arabidopsis TOR kinase specifically regulates the expression of nuclear genes coding for plastidic ribosomal proteins and the phosphorylation of the cytosolic ribosomal protein S6. *Front. Plant Sci.* **7**, 1–16 (2016).

19. Saxton, R. A. & Sabatini, D. M. mTOR Signaling in Growth, Metabolism, and Disease. *Cell* **168**, 960–976 (2017).

20. Tatebe, H. & Shiozaki, K. Evolutionary Conservation of the Components in the TOR

Signaling Pathways. *Biomolecules* **7**, 1–17 (2017).

21.   Van Leene, J. *et al.* Capturing the phosphorylation and protein interaction landscape of the
plant TOR kinase. *Nat. Plants* **5**, 316–327 (2019).

22.   Menand, B. *et al.* Expression and disruption of the Arabidopsis TOR (target of rapamycin)
gene. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 6422–6427 (2002).

23.   Anderson, G. H., Veit, B. & Hanson, M. R. The Arabidopsis AtRaptor genes are essential for
post-embryonic plant growth. *BMC Biol.* **3**, 1–12 (2005).

24.   Deprost, D., Truong, H. N., Robaglia, C. & Meyer, C. An Arabidopsis homolog of
RAPTOR/KOG1 is essential for early embryo development. *Biochem. Biophys. Res.*
Commun. **326**, 844–850 (2005).

25.   Mahfouz, M. M., Kim, S., Delauney, A. J. & Verma, D. P. S. Arabidopsis TARGET of
RAPAMYCIN interacts with RAPTOR, which regulates the activity of S6 kinase in response to
osmotic stress signals. *Plant Cell* **18**, 477–490 (2006).

26.   Yang, H. *et al.* MTOR kinase structure, mechanism and regulation. *Nature* **497**, 217–223
(2013).

27.   Rexin, D., Meyer, C., Robaglia, C. & Veit, B. TOR signalling in plants. *Biochem. J.* **470**, 1–14
(2015).

28.   Salem, M. A., Li, Y., Wiszniewski, A. & Giavalisco, P. Regulatory-associated protein of TOR
(RAPTOR) alters the hormonal and metabolic composition of Arabidopsis seeds, controlling
seed morphology, viability and germination potential. *Plant J.* **92**, 525–545 (2017).

29.   Moreau, M. *et al.* Mutations in the Arabidopsis homolog of LST8/GβL, a partner of the
target of Rapamycin kinase, impair plant growth, flowering, and metabolic adaptation to
long days. *Plant Cell* **24**, 463–481 (2012).

30.   McCready, K., Spencer, V. & Kim, M. The Importance of TOR Kinase in Plant Development.
Front. Plant Sci. **11**, 1–7 (2020).

31.   Henriques, R. *et al.* Arabidopsis S6 kinase mutants display chromosome instability and
altered RBR1-E2F pathway activity. *EMBO J.* **29**, 2979–2993 (2010).

32.   Ahn, C. S., Han, J. A., Lee, H. S., Lee, S. & Pai, H. S. The PP2A regulatory subunit Tap46, a
component of the TOR signaling pathway, modulates growth and metabolism in plants.
Plant Cell **23**, 185–209 (2011).

33.   Xiong, Y. & Sheen, J. Rapamycin and glucose-target of rapamycin (TOR) protein signaling in
plants. *J. Biol. Chem.* **287**, 2836–2842 (2012).

34.   Patrick, R. M. *et al.* Discovery and characterization of conserved binding of eIF4E 1 (CBE1),
a eukaryotic translation initiation factor 4E– binding plant protein. *J. Biol. Chem.* **293**,
17240–17247 (2018).

35.   Xiong, Y. & Sheen, J. Novel links in the plant TOR kinase signaling network. *Curr. Opin. Plant*
Biol. **28**, 83–91 (2015).

36.   Dobrenel, T. *et al.* TOR Signaling and Nutrient Sensing. *Annu. Rev. Plant Biol.* **67**, 261–285
(2016).

37.   González, A. & Hall, M. N. Nutrient sensing and TOR signaling in yeast and mammals .
EMBO J. **36**, 397–408 (2017).

38.   Bakshi, A., Moin, M., Madhav, M. S. & Kirti, P. B. Target of rapamycin, a master regulator of
multiple signalling pathways and a potential candidate gene for crop improvement. *Plant*

592		*Biology* vol. 21 190–205 (2019).
593	39.	Pacheco, J. M. *et al.* The tip of the iceberg: emerging roles of TORC1, and its regulatory
594		functions in plant cells. *J. Exp. Bot.* **72**, 4085–4101 (2021).
595	40.	Dong, Y. *et al.* Sulfur availability regulates plant growth via glucose-TOR signaling. *Nat.*
596		*Commun.* **8**, 1–10 (2017).
597	41.	Cao, P. *et al.* Homeostasis of branched-chain amino acids is critical for the activity of TOR
598		signaling in Arabidopsis. *Elife* **8**, 1–24 (2019).
599	42.	Schaufelberger, M. *et al.* Mutations in the Arabidopsis ROL17/isopropylmalate synthase 1
600		locus alter amino acid content, modify the TOR network, and suppress the root hair cell
601		development mutant lrx1. *J. Exp. Bot.* **70**, 2313–2323 (2019).
602	43.	Sun, X. *et al.* Low nitrogen induces root elongation via auxin-induced acid growth and
603		auxin-regulated target of rapamycin (TOR) pathway in maize. *J. Plant Physiol.* **254**, 153281
604		(2020).
605	44.	Liu, Y. *et al.* Diverse nitrogen signals activate convergent ROP2-TOR signaling in
606		Arabidopsis. *Dev. Cell* **56**, 1283-1295.e5 (2021).
607	45.	Schepetilnikov, M. *et al.* GTP ase ROP 2 binds and promotes activation of target of
608		rapamycin, TOR , in response to auxin . *EMBO J.* **36**, 886–903 (2017).
609	46.	Li, X. *et al.* Differential TOR activation and cell proliferation in Arabidopsis root and shoot
610		apexes. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 2765–2770 (2017).
611	47.	Carol, R. J. *et al.* A RhoGDP dissociation inhibitor spatially regulates growth in root hair
612		cells. *Nature* **438**, 1013–1016 (2005).
613	48.	Duan, Q., Kita, D., Li, C., Cheung, A. Y. & Wu, H. M. FERONIA receptor-like kinase regulates
614		RHO GTPase signaling of root hair development. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 17821–
615		17826 (2010).
616	49.	Kang, E. *et al.* The microtubule-associated protein MAP18 affects ROP2 GTPase activity
617		during root hair growth. *Plant Physiol.* **174**, 202–222 (2017).
618	50.	Denninger, P. *et al.* Distinct RopGEFs Successively Drive Polarization and Outgrowth of
619		Root Hairs. *Curr. Biol.* **29**, 1854-1865.e5 (2019).
620	51.	Huang, G. Q. *et al.* Arabidopsis RopGEF4 and RopGEF10 are important for FERONIA-
621		mediated developmental but not environmental regulation of root hair growth. *New*
622		*Phytol.* **200**, 1089–1101 (2013).
623	52.	Xu, G. *et al.* FERONIA phosphorylates E3 ubiquitin ligase ATL6 to modulate the stability of
624		14-3-3 proteins in response to the carbon/nitrogen ratio. *J. Exp. Bot.* **70**, 6375–6388
625		(2019).
626	53.	Pacheco, J. M. *et al.* The lncRNA APOLO and the transcription factor WRKY42 target
627		common cell wall EXTENSIN encoding genes to trigger root hair cell elongation. *Plant*
628		*Signal. Behav.* **16**, 1920191 (2021).
629	54.	Kwon, T., Alan Sparks, J., Liao, F. & Blancaflor, E. B. Erulus is a plasma membrane-localized
630		receptor-like kinase that specifies root hair growth by maintaining tip-focused cytoplasmic
631		calcium oscillations[OPEN]. *Plant Cell* vol. 30 1173–1177 (2018).
632	55.	Schoenaers, S. *et al.* The Auxin-Regulated CrRLK1L Kinase ERULUS Controls Cell Wall
633		Composition during Root Hair Tip Growth. *Curr. Biol.* **28**, 722-732.e6 (2018).
634	56.	Xiong, Y. *et al.* Glucose-TOR signalling reprograms the transcriptome and activates

meristems. *Nature* **496**, 181–186 (2013).

57.   Montané, M. H. & Menand, B. ATP-competitive mTOR kinase inhibitors delay plant growth
by triggering early differentiation of meristematic cells but no developmental patterning
change. *J. Exp. Bot.* **64**, 4361–4374 (2013).

58.   Escobar-Restrepo, J. M. *et al.* The Feronia receptor-like kinase mediates male-female
interactions during pollen tube reception. *Science (80-.).* **317**, 656–660 (2007).

59.   Haruta, M., Sabat, G., Stecker, K., Minkoff, B. B. & Sussman, M. R. A peptide hormone and
its receptor protein kinase regulate plant cell expansion. *Science (80-.).* **343**, 408–411
(2014).

60.   Chakravorty, D., Yu, Y. & Assmann, S. M. A kinase-dead version of FERONIA receptor-like
kinase has dose-dependent impacts on rosette morphology and RALF1-mediated stomatal
movements. *FEBS Lett.* **592**, 3429–3437 (2018).

61.   Deprost, D. *et al.* The Arabidopsis TOR kinase links plant growth, yield, stress resistance
and mRNA translation. *EMBO Rep.* **8**, 864–870 (2007).

62.   Jones, M. A. *et al.* The arabidopsis Rop2 GTPase is a positive regulator of both root hair
initiation and tip growth. *Plant Cell* **14**, 763–776 (2002).

63.   Gendre, D. *et al.* Rho-of-plant activated root hair formation requires arabidopsis YIP4a/b
gene function. *Dev.* **146**, (2019).

64.   Zhu, S., Martínez Pacheco, J., Estevez, J. M. & Yu, F. Autocrine regulation of root hair size
by the RALF-FERONIA-RSL4 signaling pathway. *New Phytol.* **227**, 45–49 (2020).

65.   Song, Y. *et al.* FERONIA restricts Pseudomonas in the rhizosphere microbiome via
regulation of reactive oxygen species. *Nat. Plants* **7**, 644–654 (2021).

66.   Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time
quantitative PCR and the 2-ΔΔCT method. *Methods* **25**, 402–408 (2001).