Pervasive conditional selection of driver mutations and modular epistasis networks in cancer

Jaime Iranzo[1,2]*, George Gruenhagen[3], Jorge Calle-Espinosa[1], Eugene V. Koonin[4]*

[1]Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid (UPM) - Instituto
Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA), Madrid, Spain.

[2]Institute for Biocomputation and Physics of Complex Systems (BIFI), University of Zaragoza,
Zaragoza, Spain.

[3]Institute of Bioengineering and Biosciences, School of Biological Sciences, Georgia Institute of
Technology, Atlanta GA, USA.

[4]National Center for Biotechnology Information, National Library of Medicine, National Institutes of
Health, Bethesda MD, USA.

*Corresponding authors: koonin@ncbi.nlm.nih.gov, jaime.iranzo@upm.es

Summary

Cancer driver mutations often display mutual exclusion or co-occurrence, underscoring the key role
of epistasis in carcinogenesis. However, estimating the magnitude of epistasis and quantifying its
effect on tumor evolution remains a challenge. We developed a method to quantify *COnditional*
SELection on the Excess of Nonsynonymous Substitutions (*Coselens*) in cancer genes. *Coselens* infers
the number of drivers per gene in different partitions of a cancer genomics dataset using covariance-
based mutation models and determines whether coding mutations in a gene affect selection for
drivers in any other gene. Using *Coselens*, we identified 296 conditionally selected gene pairs across
16 cancer types in the TCGA dataset. Conditional selection affects 25-50% of driver substitutions in
tumors with >2 drivers. Conditionally co-selected genes form modular networks, whose structures
challenge the traditional interpretation of within-pathway mutual exclusivity and across-pathway
synergy, suggesting a more complex scenario, where gene-specific across-pathway epistasis shapes
differentiated cancer subtypes.

Keywords: cancer genomics, conditional selection, epistasis, mutational landscape, cancer subtypes,
co-occurrence, mutual exclusivity.

1   Introduction

Cancer is a result of somatic evolution, during which cells accumulate driver mutations (drivers for
short) that enable them to proliferate, bypassing cell cycle checkpoints and avoiding senescence,
invade the surrounding tissues, escape the immune system, and ultimately disseminate to distant
body locations(Hanahan and Weinberg, 2000; Stratton et al., 2009). By virtue of increasing the
growth and survival rates of (pre)cancer cells, drivers are positively selected and can reach high
prevalence within the tumor although not necessarily fixed (Garraway and Lander, 2013; Greaves
and Maley, 2012; Nik-Zainal et al., 2012; Vogelstein et al., 2013; Williams et al., 2018). Concomitantly,
(pre)cancer cells acquire passenger mutations which, despite not contributing to cancer initiation or
progression, can spread by hitchhiking on drivers and typically are the dominant component of
cancer mutational landscapes(Martincorena et al., 2017; Weghorn and Sunyaev, 2017).

The precise set of drivers acquired by a tumor determines some of its fundamental properties, such
as metastatic potential and sensitivity to targeted therapies(Senft et al., 2017). Some of these
properties can be related to the presence or absence of a single driver, whereas others result from
the combination of multiple drivers that interact synergistically or interfere with each other, critically
affecting tumor initiation, progression, and clinical outcome(Ashworth et al., 2011; Mina et al., 2017;
Zhang et al., 2017). Two well-studied examples involve the synergy between activating mutations in
oncogene *RAS* and inactivation of the tumor suppressor *ATM*, and the synthetic lethality of *EGFR* and
KRAS double mutants in lung and colon cancers. In the first case, inactivation of ATM allows tumor
cells to overcome oncogene-induced senescence associated with RAS activation, and therefore
exacerbates cell proliferation(Di Micco et al., 2006). In the second case, oncogenic mutations in *KRAS*
make EGFR-mutant tumors resistant to therapies that target EGFR, but the synergistic deleterious
effect of *KRAS* and *EGFR* mutations makes such escape clones inviable in the absence of the
drug(Misale et al., 2014).

Epistatic interactions, such as those mentioned above, have the potential to substantially affect
cancer somatic evolution by making selection on a driver conditional on the presence or absence of
another driver (Wilkins et al., 2018). Epistasis results in patterns of co-occurrence when synergy
between drivers enhances positive selection or mutual exclusivity when redundancy or antagonism
diminish selection on the double mutant. Epistasis-driven conditional selection implies that past
driver mutations affect the fitness of subsequent driver mutations and, with that, the evolutionary
trajectory of the tumor.

Searching for mutually exclusive mutations is a main focus of research in cancer genomics because
such findings can lead to identification of synthetic lethal pairs and cancer-associated pathways,
which are potential therapeutic targets(Babur et al., 2015; Ciriello et al., 2012; Jerby-Arnon et al.,
2014; Kim et al., 2017; Leiserson et al., 2015; Matlak and Szczurek, 2017; Mina et al., 2017; Srihari et
al., 2015; Wappett et al., 2016). However, because co-occurrence and mutual exclusivity can also
result from systematic biases in tumor mutational load, their presence cannot be taken as
unequivocal evidence of epistasis(van de Haar et al., 2019). Furthermore, the statistical frameworks
commonly used to test for mutation co-occurrence and mutual exclusivity fail to provide quantitative
estimates of biologically meaningful effect sizes, such as the extent to which epistasis modifies the
fitness of a mutant clone and the subsequent probability that a mutation becomes a driver(Cannataro
et al., 2018).

Obtaining quantitative estimates of epistatic effects in a systematic and direct manner is extremely challenging, although a recent study has approached this goal by conducting combinatorial RNAi screening in cell culture followed by image-based phenotyping of growth-related traits(Wang et al., 2014). An alternative approach is to quantify the contribution of epistasis to cancer development indirectly, through its effect on the conditional selection of driver mutations. This approach, which takes advantage of the recent surge in cancer genomic data, has the fundamental advantage of not relying on ad-hoc (*in vitro*) proxies for *in vivo* tumor fitness. A major difficulty, however, is that quantification of conditional selection requires disentangling the contributions of mutation and selection to cancer genomic landscapes, a challenging task that has only recently begun to be addressed by adapting tools from evolutionary genetics for cancer genomics(Cannataro et al., 2018; Martincorena et al., 2017; Persi et al., 2018; Weghorn and Sunyaev, 2017).

Along these lines, we developed a method, Coselens (COnditional SELection on the Excess of Nonsynonymous Substitutions), that compares the mean number of drivers per tumor in a gene depending on whether a second gene harbors non-synonymous mutations, returning an estimate for the magnitude of conditional selection, while controlling for differences in mutational load and spectrum. The principal advantages of Coselens over previously developed methods for analysis of epistasis in cancer is that this method rigorously controls for the biases introduced by cancer subtypes and explicitly estimates the size of epistatic effects. Application of Coselens to somatic mutation data from more than 10,000 cancer exomes led to the identification of a complex network of cancer type-specific epistasis, shedding light on the underlying genetic constraints that shape cancer development.

Results and Discussion

<u>Detection of pairwise conditional selection among cancer genes</u>

Driver mutations have been the subject of intense research in recent years, leading to multiple definitions and strategies for driver detection. In this work, we adopted an evolutionary genetics approach, defining drivers as mutations that provide (pre)cancer cells with a selective advantage, and therefore are subject to positive selection. Because purifying selection is typically negligible in non-hypermutated tumors (Martincorena et al., 2017; Persi et al., 2018), the number of positively selected mutations (that is, drivers) in a locus can be obtained as the difference between the number of mutations observed in that locus and the number of mutations that would be expected in the absence of selection. The main challenge of this approach is to infer the null mutation model under the assumption of neutrality, which we did by combining information from synonymous mutations and genomic covariates. This method accounts for the possibility that the same mutation does not necessarily act as driver in different types of tumors and provides a quantitative estimate of the probability that a particular mutation, if observed, is indeed a driver. Although this definition applies, in principle, to any kind of mutation, we focused on nonsynonymous and truncating substitutions as driver candidates because those are the mutations for which the most reliable null models can be built.

To investigate the role of conditional selection in cancer progression, we quantified the dependency of the number of driver substitutions in a cancer gene (the "query" gene) on the presence or absence

of coding mutations in a second gene (the "split" gene). For each cancer type in The Cancer Genome Atlas (TCGA), we first identified the set of query genes by pooling all somatic mutations from the same cancer type and searching for genes for which the dN/dS indicated significant levels of positive selection (Supplementary Table 1). Query genes with coding mutations (nonsynonymous substitutions, truncations, and small indels) in >25 samples were additionally considered as split genes. Then, for all split-query gene pairs, we classified tumor samples into those that did and did not harbor coding mutations in the split gene and applied our evolutionary genetics-inspired method *Coselens* to estimate and compare the number of driver substitutions in the query gene in each of the groups (Figure 1a). In the first step, *Coselens* fits a mutational model to each group of samples and calculates the mean number of nonsynonymous and truncating substitutions that would be expected in the query gene in the absence of selection. Then, it calculates the number of drivers as the difference between the number of observed and expected mutations (see Methods for details). Finally, *Coselens* implements a likelihood ratio test to assess whether the number of driver events significantly differs between samples that do and do not harbor coding mutations in the split gene.

Our analysis identified 296 gene pairs subject to significant conditional selection (FDR $\leq$ 5%) in at least one cancer type (Supplementary Table 2). Of these, 262 were deemed significant by the standard workflow and the rest were identified through restricted hypothesis testing of reciprocal conditional selection (that is, by testing the reverse pairs, in which the split and query genes were swapped). The effect size distribution of significant pairs identified through restricted hypothesis testing closely resembles that of significant pairs detected through the more stringent, standard workflow (Figure 1b), suggesting that the former represent genuine cases of conditional selection. Overall, conditional selection affects 112 genes across 16 cancer types. In 70% of the conditionally selected gene pairs, coding mutations in the split gene increase the probability of finding driver mutations in the query gene; henceforth, we refer to these conditional pairs as positive. In the remaining 30%, coding mutations in the split gene decrease the probability to find driver mutations in the query gene, and henceforth, we refer to these conditional pairs as negative (Figure 1c).

The magnitude of conditional selection varies among gene pairs, with coding mutations in the split gene typically associated with an increase or decrease of at least 0.15 driver substitutions per query gene per tumor (Figure 1b). In some extreme cases, conditional selection results in almost complete presence or absence of a driver mutation in the query gene. The pair of closely related oncogenes *GNAQ* and *GNA11* is a good example of such extreme conditionality: driver substitutions in either GNAQ or GNA11 are found in 80-90% of uveal melanomas. However, mutations in these genes are mutually exclusive(Decatur et al., 2016; Van Raamsdonk et al., 2010), so that the absence of mutations in one implies, with high probability, the presence of driver mutations in the other. On the other extreme, the pair formed by *TP53* and *ATRX* shows strong, mutual positive conditionality in low-grade gliomas, such that driver substitutions in *ATRX* appear almost exclusively in mutated *TP53* backgrounds(Liu et al., 2012).

We compared the results of *Coselens* with those obtained using a mutual information approach (*SELECT*), which was also developed to identify conditional selection in cancer(Mina et al., 2017). The comparison indicates that *Coselens* detects many more conditionally selected gene pairs than *SELECT*, but the two approaches seem to capture complementary aspects of conditional selection in cancer genomes (Figure 1d). By leveraging information from copy number alterations (CNA), *SELECT* can detect conditional selection in genes whose role in cancer depends on amplification and/or

deletion, albeit at the cost of not providing easily interpretable effect sizes. In contrast, *Coselens*
proves substantially more powerful at detecting and quantifying conditional selection among cancer
genes that harbor driver substitutions, with 7- and 4-fold increases in the number of positive and
negative conditional pairs compared to *SELECT* (Supplementary Figure 1). Such higher detection
power can be partly explained by *Coselens*' ability to analyze genes that fail some of the pre-filtering
criteria applied by *SELECT* (this is the case for 79 of the 250 *Coselens*-exclusive pairs). In turn, 17 of
the 23 non-CNA, *SELECT*-exclusive pairs involved genes that did not show evidence of positive
selection as per their *dN/dS* values.

<u>Mutational load does not substantially confound *Coselens*' estimates of conditional selection</u>

Previous studies have noted that detection of mutual exclusivity can be confounded by intrinsic
differences in the frequency and spectrum of mutations across cancer subtypes(Kim et al., 2017; van
de Haar et al., 2019). To minimize this risk, *Coselens* independently fits two baseline mutation models,
one for samples with coding mutations in the split gene and another for samples without such
mutations. The use of subset-specific models for the expected number of substitutions helps prevent
spurious correlations resulting from across-subset variations in mutational signatures. However,
differences in mutational load can also affect the detection of conditional selection indirectly, in cases
when mutation availability is a limiting factor in tumor evolution. If differences in mutational load
significantly affect the availability of mutations on which selection can act, then, associations
between rare cancer genes could reflect patterns of differential mutation availability rather than
conditional selection. To account for that possibility, we adopted two alternative correction methods:
one applies sampling theory to assess the likelihood that each significant gene pair can be explained
by differences in mutation availability, and the other uses regression to identify biases in the number
of significant pairs that correlate with mutation load (see Methods). The results obtained with both
methods were in close agreement, leading to the conclusion that >80% of conditional gene pairs
could not be explained by differential mutation availability (Figure 1c).

We also investigated whether nonsynonymous passenger mutations in split genes quantitatively
affected the inference of conditional selection. Because conditional selection is inferred by
partitioning tumors into those with and without coding mutations in one of the genes, the inclusion
of non-driver mutants in the former set could lead to underestimating the real magnitude of the
association between driver mutations. To test whether estimates of conditional selection are indeed
affected by passenger coding mutations in the split gene, we used Bayes theorem to derive an
equivalence expression that relates the standardized effect size of conditional selection upon
swapping the split and query genes (see Methods). This equivalence should hold exactly in the case
when conditional selection results from epistasis between driver mutations, but deviations are
expected to appear if the inference of conditional selection is biased by passenger coding mutations
in the split gene. Following this rationale, we compared the standardized effect sizes of conditional
selection in swapped gene pairs (Figure 1e). The close agreement between empirical estimates and
theoretical predictions (intraclass correlation = 0.88, N = 78, p<1e-4) confirms that estimates of
coding-to-driver conditional selection are good proxies for the magnitude of driver-to-driver
epistasis.

Functional association and classification of epistasis in conditionally selected gene pairs

To better characterize the mechanisms underlying conditional selection, we searched the STRING
database for functional and physical interactions between the split- and query-gene products. Based
on that search, we calculated an interaction enrichment ratio that quantifies, for any subset of gene
pairs, the enrichment in predicted interactions of different types with respect to the set of all tested
gene pairs. Given that STRING compiles extensive information on various types of association
between genes and the encoded proteins, including functional data extracted from the literature,
experimental evidence of physical interactions, and coexpression across a broad diversity of
organisms (Szklarczyk et al., 2021), we considered it the best available source of background
knowledge of gene-gene interactions. We found that enrichment in interactions increases with the
increase of the effect size of conditional selection (Figure 2a). Pairs subject to significant conditional
selection encompass more interactions than expected by chance, with a 1.5-fold enrichment (95% CI
1.2-2.0) in the case of positive conditional pairs and a 4.2-fold enrichment (95% CI 2.7-6.6) in the case
of negative conditional pairs (Figure 2b). Regarding the specific nature of interactions, no significant
differences were found between functional association and physical binding, although the latter
seem to be slightly more prevalent among the negative conditional pairs.

Epistasis between cancer genes can be classified attending to the sign and magnitude of the
conditional selection on driver substitutions. We identified 4 major classes of epistasis (Figure 2c):
strict dependence, when driver (that is, positively selected) substitutions in the query gene are only
observed in the presence of coding mutations in the split gene; *facilitation*, when driver substitutions
are more likely to occur when the split gene is mutated; *interference*, when driver substitutions are
less likely to occur when the split gene is mutated; and *masking*, when mutations in the split gene
completely abolish the effect of driver substitutions in the query gene. Although strict dependence
and masking appear as the extremes in the continuum that goes from facilitation to independence
to interference, they can be distinguished by testing the null hypothesis that there is no significant
selection on nonsynonymous mutations. By far, the most prevalent form of conditional selection is
facilitation, which was identified in more than half of the significant gene pairs (Figure 2d). Focusing
on the extreme classes, masking represents almost 40% of negative epistatic interactions (11% of the
total), whereas strict dependence only represents 20% of positive interactions (14% of the total).

Positive and negative epistasis differ in the degree of mutual interdependency between the two
genes of the same pair (Figure 2e). Negative epistasis tends to be symmetrical and often involves
mutual masking. This is consistent with negative epistasis arising from complete functional
redundancy although it could also reflect the existence of multiple, incompatible mutational paths
to cancer (see below). On the other side, the spectrum of positive epistasis is broader and combines
cases of strict interdependence and asymmetric facilitation. A biologically plausible interpretation of
asymmetric facilitation is that driver mutations in a "master" cancer gene open new pathways of
tumor development that involve the dependent gene. As a result, mutations in the dependent gene
only act as drivers if the master gene carries drivers as well , but not vice versa. Supporting this
interpretation, we found that in pairs with moderate and high asymmetry, mutations in the master
gene tend to show a higher degree of clonality (mean relative difference in variant allele frequency
= 0.13, p = 0.009, 1-sample Student's t-test, n = 20) and therefore are likely to precede mutations in
the dependent gene (Figure 2f). In contrast, no systematic differences in clonality were found
between genes that participate in symmetric epistasis. We observed that 6 of the 15 most

asymmetric pairs identified in this study involved *PTEN* mutations in endometrial cancers
(Supplementary Table 3), which underscores the fundamental role of *PTEN* inactivation as an early
driver event in this cancer type(Levine et al., 1998; Zhang and Yu, 2010). Another highly asymmetric
pair, observed in bladder and colorectal cancer, consists of *TP53* as the master and *RB1* as the
dependent gene. Considering the tumor suppressor role of these genes, the asymmetry of
conditional selection implies that, for *RB1* loss to act as a driver, *TP53* must also harbor inactivating
driver mutations, whereas *TP53* inactivation can act as a driver on its own. Thus, an intact *TP53* could
partially compensate for the loss of *RB1*, but not vice versa.

<u>Conditional selection across cancer types</u>

Different cancer types broadly vary in the number of gene pairs subject to conditional selection, with
the highest numbers found in endometrial and colorectal cancers, followed by stomach and bladder
cancers (Figure 3a). To quantify the contribution of conditional selection to cancer somatic evolution,
we compared the numbers of driver substitutions per tumor that were affected by conditional
selection across cancer types (Figure 3b). We found that positive conditional selection is responsible
for the spread of a sizable fraction (25-50%) of all driver substitutions. The relative contributions of
interference and masking are generally smaller although in some tumors, such as uveal melanoma,
lower glade gliomas, and endometrial cancers, negative epistasis could lead to lack of selection for
(and therefore hinder fixation of) >40% potential driver mutations.

Both the number of gene pairs subject to conditional selection and the average number of
conditionally-selected driver substitutions strongly correlate with the total number of driver
substitutions per tumor (Figure 3c-d). These associations persist when controlling for several possible
confounding factors, including the number of samples and mutations per cancer type, median
mutational load, and number of gene pairs tested, that might unequally affect our ability to detect
conditional selection in different cancer types (Spearman partial correlation coefficients rho = 0.95
($p<10^{-8}$) for the number of gene pairs and rho = 0.80 ($p<10^{-4}$) for the number of conditional drivers,
n = 22). A strong correlation was observed also with the number of non-conditionally-selected driver
substitutions (Spearman partial correlation coefficients rho = 0.96 ($p<10^{-8}$) for the number of gene
pairs and rho = 0.75 (p<0.001) for the number of conditional drivers, n = 22). Taken together, these
observations imply that conditional selection is intrinsic to cancer somatic evolution, its contribution
becoming increasingly important as the number of driver mutations increases.

Most gene pairs subject to conditional selection are highly specific to a single cancer type, with only
3% positive and 12% negative conditional pairs found in >1 cancer type (Figure 3d). Such higher
promiscuity of negative conditional gene pairs, though small in magnitude, is statistically significant
(chi-square = 6.24, p = 0.03, permutation test). Not surprisingly, multi-tissue gene pairs include well-
known examples of genes that display mutually exclusive or concurrent mutation patterns in pan-
cancer analyses(Kandoth et al., 2013). Among the negative conditional pairs found in multiple cancer
types, those involving oncogenes *KRAS*/*NRAS* and *BRAF* are easily ascribed to the functional
redundancy in activating the Ras signaling pathway. The mutual exclusivity between *ARID1A* and
TP53 possibly results from functional codependency in suppressing tumor development through
transcriptional regulation(Guan et al., 2011; Wu et al., 2014). However, the mechanisms underlying
the negative conditionality between mutations in *TP53* and *KRAS* (and also *TP53* and *PIK3CA*) remain
unclear and could reflect fundamental, but poorly understood interactions among major pathways

involved in cancer development. Positive conditional pairs spanning multiple cancer types include associations among the genes *ATRX*, *IDH1*, and *TP53*, which are known to harbor concurrent mutations in brain tumors (glioblastoma and lower grade gliomas)(Brennan et al., 2013; Cancer Genome Atlas Research et al., 2015; Liu et al., 2012). Other multi-tissue pairs are *TP53-RB1* (detected in bladder cancer and glioblastoma), *ARID1A-ERBB3* (bladder and stomach cancer), and *PGM5-RNF43* (colorectal and stomach cancer). Interestingly, the gene *PGM5* (encoding phosphoglucomutase-like protein 5) has not been previously reported as a cancer gene, although low *PGM5* expression has been linked to poor survival in liver and colorectal cancers(Jiao et al., 2019; Sun et al., 2019). Almost 75% (35 out of 47) coding mutations in *PGM5* detected in stomach and colorectal cancers correspond to a single nonsynonymous substitution with predicted deleterious effect (SIFT: 0.01 - deleterious; PolyPhen: 0.573 - possibly damaging) affecting position 98 (Ile>Val) of the encoded protein, without loss of the second copy of the gene. In contrast, this substitution only represents 7% (7 out of 96) *PGM5* coding mutations in other cancer types (Figure 3f). Thus, the PGM5 p.I98V mutation is likely to be a specific driver of gastric and colorectal cancers that possibly operates through a gain-of-function mechanism.

Apart from the broadly spread gene pairs described above, the specificity of most conditional pairs for a single cancer type suggests that epistasis is often context-dependent and is conditioned on the physiology of the tumor(Park and Lehner, 2015; Yeang et al., 2008). This context dependency is manifest at two levels: first, cancer genes involved in conditional selection are often specific to one or a small number of cancer (sub)types; second, when the same gene appears in more than one cancer type, it is often involved in different conditional pairs. The first situation is illustrated by *HLA-C*, which we identified as a conditional cancer gene in *BRAF*-mutant colorectal cancer (Figure 4). Conditional selection on *HLA-C* inactivating mutations is probably mediated by the highly immunogenic nature of tumors of the consensus molecular subtype 1 (CMS-1), whereby the loss of MHC-I-mediated antigen presentation confers a selective advantage(Jhunjhunwala et al., 2021). Supporting this interpretation, multiple studies have described driver mutations in *B2M* and *HLA* genes in CMS-1 colorectal cancer and other microsatellite unstable tumors characterized by high levels of CD8+ T cell infiltration(Castro et al., 2019; Kloor et al., 2007; Shukla et al., 2015). The second situation is exemplified by *RNF43* and its different conditionality patterns observed in colorectal, gastric, and endometrial cancers (Figure 4 and Supplementary Figure 2). In agreement with these differences, two distinct mechanisms have been proposed to explain how the loss of *RNF43* contributes to carcinogenesis, with deregulation of Wnt signaling being dominant in the colon and modulation of the DNA damage response playing the principal role in the stomach(Neumeyer et al., 2021; Neumeyer et al., 2019). An even more striking example is presented by conditionally selected mutations in *KRAS* and *TP53*, which are antagonistic in lung and endometrial tumors, but synergistic in pancreatic cancer. For the latter cancer type, recent research has shown that mutant KRAS and p53 interact through the CREB1 protein to promote metastasis(Kim et al., 2021). The pronounced cancer type-specificity of conditional selection among widespread cancer genes highlights the need to integrate functional pleiotropy, cellular environment, and tumor physiology in models of cancer evolution(DeGregori, 2017).

Epistasis networks

To further elucidate how epistasis among somatic mutations affects tumor evolution, we built epistasis networks for each cancer type by connecting the genes that are involved in conditional

selection. The most salient feature of such networks is their modular organization (Figure 4). Modularity implies that cancer genes organize into clearly differentiated groups (hereafter modules), such that genes within a module connect to each other through positive epistasis, whereas genes from different modules display negative or no epistasis. Modularity is most evident in cancer types where conditional selection affects a large fraction of drivers, such as low-grade glioma, colorectal, and endometrial cancers.

Inspection of the genes that belong to different modules suggests an association between the modular structure of the epistasis network and the existence of well-defined molecular cancer subtypes. This association is especially clear in low-grade gliomas, where the three main modules encompassing (i) *IDH1*, *ATRX*, and *TP53*; (ii) *CIC* and *FUBP1*; and (iii) *EGFR* and *PTEN*, correspond, respectively, to three recently proposed molecular subtypes based on the *IDH1* mutation and 1p/19q codeletion status(Cancer Genome Atlas Research et al., 2015; Ceccarelli et al., 2016), whereas negative conditional selection between the genes of the first module and the *NF1* gene is reminiscent of the molecular landscape found in gliomas from patients with neurofibromatosis type 1 and the LGm6 subgroup of sporadic glioma(Ceccarelli et al., 2016; D'Angelo et al., 2019; Verhaak et al., 2010). In glioblastoma, positive conditional selection among *IDH1*, *ATRX*, and *TP53* defines a module associated with G-CIMP (glioma-CpG island methylator phenotype) tumors from the proneural subtype(Brennan et al., 2013; Noushmehr et al., 2010). To a lesser extent, the high prevalence of mutations in *BRAF* suggests a connection between consensus molecular subtype 1 of colorectal cancer and a module involving *BRAF*, *BMPR2*, *RNF43* and *HLA-C* (Guinney et al., 2015). Similarly, in endometrial cancer, the detection of positive conditional selection between *TP53* and *PPP2R1A* and the antagonism between these two genes and *PTEN* are consistent with the mutational landscape of serous endometrial tumors of the "copy number-high" molecular subtype(Cancer Genome Atlas Research et al., 2013; McConechy et al., 2011).

A possible explanation for the association between conditionally selected gene sets and cancer subtypes could be that the latter are a confounding factor for the inference of conditional selection. More precisely, because all samples from the same cancer type are pooled for the analysis, groups of genes that are preferentially mutated in a single subtype but not in the rest could appear as being jointly selected(van de Haar et al., 2019). To rule out this possibility, we carried out a reanalysis of conditional selection in cancer types with sufficient sample size and well-defined molecular subtypes, stratifying the data according to subtype classification. Although the reduction in sample size affected the ability to detect conditional pairs as statistically significant, we recovered some of the key interactions that characterized the modules of the epistasis network (Supplementary Table 4). More generally, we found that the estimated effect sizes for conditional selection in individual subtypes were highly correlated with the values obtained in the whole dataset (Figure 5a). Altogether, the results of the stratified analysis show that conditional selection also occurs within molecular subtypes and, therefore, is not an artifact of the joint analysis of samples from different groups.

Mutual exclusivity and co-occurrence among cancer somatic mutations have traditionally been interpreted in terms of within-pathway redundancy and across-pathway synergy(Yeang et al., 2008). In the simplest scenario, the synergy involves activation of an oncogenic pathway that promotes proliferation coupled with inactivation of a tumor suppressor pathway that regulates a cell cycle checkpoint(Cui et al., 2007). A relevant question concerning across-pathway synergy of driver mutations is whether it occurs in a combinatorial (or gene-agnostic) manner, that is, regardless of

the specific genes that are mutated in each pathway(Oikonomou et al., 2014). To address this question, we investigated the distribution of triangular motifs in the epistasis networks. If across-pathway synergy was gene-agnostic, positive epistasis would manifest as an excess of triangular motifs containing two positive and one negative connection (Figure 5b). Contrary to this expectation, we found a significant excess of triangular motifs containing one positive and two negative connections in the epistasis networks (Figure 5c). Such configuration suggests a more complex scenario, in which across-pathway synergy is gene-specific, that is, depends on which genes are mutated in each pathway (Figure 5d). In that scenario, mutual exclusivity among functionally redundant mutations in the same pathway can still be expected; however, only some combinations of mutations affecting different pathways can act synergistically driving the progression of specific cancer subtypes. Consistent with that scenario, visual inspection of epistasis networks reveals a highly gene-specific pattern of across-pathway interconnections, whereby genes from the same pathway (for example, *RNF43* and *APC*, or *BRAF* and *RAS*) rarely display positive epistasis with the same partners. This finding calls for a reappraisal of the adequacy of pathway-centric approaches to cancer molecular biology. Although focusing on cancer pathways rather than individual genes can simplify the overall picture and boost the sensitivity of genomic analyses, it comes at the cost of overlooking the possibility that driver mutations in different genes from the same pathway can have distinct effects on tumor progression due to their involvement in different interactions with other pathways.

To better understand the effect of epistasis on cancer mutation landscapes, we simulated tumor evolution in the presence of conditional selection (Figure 6a). The simulation results suggest that the structure of epistasis networks can play a major role in determining the set of driver mutations observed in evolved tumors. In particular, highly modular facilitation networks with across-module interference, such as those observed in low grade gliomas, colorectal, and endometrial cancers, spontaneously give rise to clearly differentiated mutation-based subtypes (Figure 6b, column 4). This phenomenon is less pronounced in star-like epistasis networks (that is, those characterized by a single master driver that enhances the fitness of several secondary drivers; Figure 6b, columns 2-3) and completely absent in negative-modular networks (those corresponding to an idealized scenario of full within-pathway redundancy and across-pathway synergy; Figure 6b, column 5).

Based on these results, we hypothesize that tumor evolution is constrained by the structure of epistasis networks. In cancer types with strong modular epistasis, early drivers could be major determinants of the evolutionary path that a precancerous clone will take by modifying the strength of selection on subsequent mutations. Eventually, such evolutionary path can lead to an advanced tumor in which the mutational landscape is characterized by drivers in genes from the same epistasis module as the original one. Accordingly, differences in the pervasiveness of conditional selection could underlie the existence of clearly delineated (sub)type-specific gene sets in some cancer types, such as gliomas, colorectal, and endometrial cancers, but not in others, such as lung and liver cancers (Iranzo et al., 2018). The hypothesis that subtypes are shaped by conditional selection is consistent with previous findings(Zhang et al., 2017) and could open a new avenue for forecasting cancer evolution, encouraging further research as new somatic mutation datasets for precancer states and early cancers become available.

Conditional selection affects survival rates

To assess the clinical relevance of conditional selection among cancer genes, we investigated the
association between the modules of the epistasis network and patient survival times (Figure 7). To
that end, we assigned tumor samples to epistatic modules by selecting, for each sample, the module
that harbored the highest number of coding mutations. Module assignment was significantly
associated with differential survival in 5 cancer types (glioblastoma, lower grade glioma, head and
neck squamous carcinomas, pancreas adenocarcinoma, and endometrial cancer). An association
between conditional selection and differential survival was also observed at the gene pair level,
although the statistical significance was lower due to the small sample sizes (Suppl Figure 3). Of the
80 gene pairs with sufficient number of samples (at least 10 double mutants and 10 single mutants
for each gene), we identified 9 instances (11% of the pairs), in which patient survival differed between
tumors carrying coding mutations in both genes and those mutated in each gene alone (Cox
regression and log-rank tests with p<0.1 for the omnibus test and p<0.15 for double- vs single-mutant
comparisons). For comparison, we ran the same analysis on non-conditionally selected gene pairs,
finding that only 4.6% (27 out of 580) showed differences in patient survival between double and
single mutants. Compared to single-gene mutants, the survival of patients with double mutations can
be improved, worsened, or unaffected. Notably, worse patient survival is associated with gene pairs
subject to positive conditional selection (*ARID1A-ERBB3* in bladder cancer, *CASP8-FAT1* in head and
neck cancers, *FGFR2-TP53* in endometrial cancer, and *MUC17-NF1* in melanoma), whereas improved
patient survival is associated with negative conditional selection (*PPP2R1A-PTEN* and *ARID1A-*
PPP2R1A in endometrial cancer). The association with better patient survival in *PPP2R1A-PTEN* and
ARID1A-PPP2R1A double mutants holds when controlling for mutational load, and suggests that
mutual exclusivity in these gene pairs could result from synthetic lethality. Likewise, synthetic
lethality might underlie other instances of negative conditional selection that could not be tested
due to a lack of double mutants, as has been shown for *KRAS* and *EGFR* in lung cancer(Unni et al.,
2015).

These results are of particular interest in the context of personalized medicine, as they show that
epistasis-driven mutational profiles could be better predictors of cancer prognosis than single
mutations. Conceivably, epistasis profiles, such as those described here, could be leveraged to aid
the selection of optimal therapeutic strategies. In particular, the set of epistatic interactions, in which
a driver mutation is involved, could be indicative of the mechanism, through which that driver
contributes to the fitness of the tumor. Thus, epistasis profiles could be key for understanding the
variability in the effect of gene-targeting therapies across patients and cancer subtypes.

Limitations of the study

An obvious limitation of the approach developed in this work is that we only sought to identify
pairwise interactions between driver mutations. Extending *Coselens* to search for higher-order
interactions is straightforward. However, because quantifying conditional selection while accounting
for heterogeneity in mutational mechanisms requires partitioning the data in as many groups as
conditions, and the number of conditions grows exponentially with the interaction order, the
detailed study of high-order interactions can only be performed in large datasets and for frequently
mutated cancer genes.

Another limitation is that we only analyzed point mutations that lead to amino acid replacement. Extending the approach to other types of mutations, in particular Copy Number Alterations (CNA), will require developing the corresponding mutational models that, in the case of CNA, will have to take into account correlations among adjacent genes. We anticipate that, by including CNA in future studies, additional instances of conditional selection will be identified for tumor suppressor genes that are typically affected by homozygous deletions (Ciriello et al., 2012; Mina et al., 2017). It also seems plausible that positive selection for CNA in tumors with chromosome instability is associated with inactivating mutations in the p53 pathway given that the latter facilitates the survival of cells with genomic aberrations (Hanel and Moll, 2012; Motoyama and Naka, 2004).

A third aspect that deserves further investigation concerns the temporal dynamics of epistasis. Changes in the mutation landscape and the selective pressures experienced by the tumor likely modify the structure of epistasis networks in different stages of cancer progression (Persi et al., 2021). However, due to the transversal nature of the TCGA, the dynamic nature of epistasis was not captured in this study. Future analyses of somatic mutations in healthy tissues and precancerous lesions would be of great interest, as they could lead to the discovery of new epistasis modules associated with protection against cancer progression.

Conclusions

In this work, we developed a method, Coselens (COnditional SELection on the Excess of Nonsynonymous Substitutions), for analysis of epistasis in cancers. Coselens includes two major features that, to the best of our knowledge, are missing in methods previously developed for the same purpose, namely, rigorous control for biases introduced by cancer subtypes and explicit estimation of the size of epistatic effects. The results reported here expand the catalogue of conditionally selected pairs of cancer genes by several folds and demonstrate wide spread and importance of conditional selection in cancer evolution. Indeed, in some cancer types, up to 50% of driver mutations are involved in such epistatic interactions. Furthermore, we observed association between conditional selection and patient survival, attesting to the clinical relevance of epistasis. Perhaps, most notably, the findings of this work emphasize the striking complexity of the epistasis among driver mutations that is manifest at several levels. First, we identified multiple cases of all theoretically conceivable forms of epistasis, both positive and negative, including strict dependence, facilitation, interference, and masking. Second, many of the conditionally selected gene pairs are cancer subtype-specific and organize into highly modular epistasis networks. Third, we found that synergistic epistasis between signaling pathways is highly gene-specific. Conceivably, epistasis shapes differentiated cancer subtypes by constraining the evolutionary path that follows early driver mutations. This evolutionary path towards advanced tumors involves mutations in a specific set of additional genes that belong to the same module of the epistasis network and determine the key properties of a molecular cancer subtype. Finally, the finding that epistatic interactions involving widespread cancer genes are cancer type-dependent raises the possibility that the same gene might act through different mechanisms in different cancer types. Thus, epistasis profiles potentially could guide the application of gene-targeting therapies across cancer types.

Acknowledgements

We thank Iñigo Martincorena and Koonin group members for helpful discussions. This work was
initiated during an internship of G.G. at Koonin group, which was made possible thanks to the NIH
Summer Internship Program in Biomedical Research. This work utilized the computational resources
of the NIH HPC Biowulf cluster (http://hpc.nih.gov).

J.I. is supported by the Ramón y Cajal Programme of the Spanish Ministry of Science (Grant No. RYC-
2017-22524), the Agencia Estatal de Investigación of Spain (Grant No. PID2019-106618GA-I00), and
the Severo Ochoa Programme for Centres of Excellence in R&D of the Agencia Estatal de Investigación
of Spain (Grant No. SEV-2016-0672 (2017–2021) to the CBGP). G.G. is supported by the
DHHS/PHS/National Institutes of Health (Grant No. 2R01DE019637-10, "Patterning the vertebrate
dentition through replacement and repair). J.C.-E. is supported by the Youth Employment Initiative
of the European Social Fund through a junior postdoctoral contract from Comunidad de Madrid
(Grant No. PEJD-2019-POST/BIO-16377). E.V.K. is supported by intramural research program funds
of the National Institutes of Health (National Library of Medicine).

Author contributions

J.I. and E.V.K. designed and supervised the study; J.I. and G.G. developed the *coselens* software and
generated the primary results; J.I., G.G., and E.V.K. analyzed the results; J.C.-E. developed and
simulated the computational model of tumor evolution; J.I., G.G., J.C-E., and E.V.K. wrote the paper.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1: Inference of pairwise conditional selection in cancer genes. a) Overview of the *Coselens*
method. The effect size ΔN_d is the difference in the frequency of driver substitutions in a given gene
(the query gene) between tumors that do and do not harbor coding mutations in a second gene (the
split gene). b) Effect size distributions for statistically significant and not-significant conditional gene
pairs. Negative effect sizes are represented by their absolute values. RHT, statistically significant pairs
under Restricted Hypothesis Testing of reciprocal interactions. c) Number of gene pairs subject to
significant conditional selection (FDR ≤ 0.05), classified by the sign of their interaction. The shaded
green bar (*) indicates a small subset of significant positive pairs whose conditional mutation
patterns can be explained by differences in mutation availability. d) Comparison of *Coselens* and
SELECT(Mina et al., 2017) methods for the detection of conditionally selected gene pairs. CNA: copy
number alterations. e) Comparison of the standardized effect sizes that result from exchanging the
query gene by the split gene (an outlier at [18.4, 8.7] is not shown). The symmetry of standardized
effect sizes implies that passenger mutations in the split gene do not substantially affect the
estimates of conditional selection acting on the query gene.

Figure 2: Functional association and epistasis classification of conditionally selected gene pairs. a)
Protein-protein interactions between gene products of the same conditional pair. The STRING
enrichment ratio (y axis) was calculated using the formula $R(x) = \frac{\sum_{G(x)} s \sum_{G(0)} (1-s)}{\sum_{G(x)} (1-s) \sum_{G(0)} s}$, where s is the
STRING score (rescaled in the range 0-1) and $G(x)$ is the set of conditional gene pairs such that $\Delta N_d \geq$
x (x axis). $R(x)$, which is formally analogous to a risk ratio for continuous response variables,
compares gene pairs above a given intensity of conditional selection with a reference base value
calculated using all possible gene pairs. The small arrow at $\Delta N_d = 0.15$ indicates the effect size of
conditional selection at which most interactions become statistically significant. Shaded regions
correspond to 95% confidence intervals. The noisy values at $\Delta N_d > 0.4$ are due to the small number
of gene pairs that reach such high levels of conditional selection. b) Enrichment in different types of
protein-protein interactions, calculated for the 296 gene pairs subject to significant conditional
selection (whiskers indicate 95% confidence intervals). c) Epistasis is quantified attending to the
frequency of driver substitutions in the query gene when the split gene does not harbor coding
mutations (x axis) and when it does (y axis). In the absence of epistasis, the frequency of drivers in
the query gene is not affected by the mutation status of the split gene, resulting in a 1:1 trend.
Facilitation and interference manifest as positive and negative deviations from that trend and their
magnitude is quantified by the angle θ (normalized by a factor $\pi/4$). Strict dependence and masking
correspond to the extreme cases $x = 0$ ($\theta = 1$) and $y = 0$ ($\theta = -1$), respectively. None of the
negative data points in the figure significatively deviates from 0. d) Relative abundances of different
classes of epistasis among 296 significantly conditional gene pairs. e) Asymmetry of epistasis. Each
data point corresponds to a statistically significant conditional gene pair whose reversed pair (with
split and query gene roles exchanged) is also significant. The diagonal represents purely symmetric
epistasis, whereby both genes equally affect selection on each other. f) Mean relative difference in
the variant allele frequencies for coding mutations in genes subject to reciprocal positive epistasis.
Gene pairs were grouped based on the asymmetry of epistasis (low asymmetry if $|\theta_{AB} - \theta_{BA}| < 0.2$;
high asymmetry if $|\theta_{AB} - \theta_{BA}| \geq 0.2$). Boxes comprise the 25th-75th percentiles; thick lines and
notches represent the median and its 95% confidence interval, respectively; whiskers extend to the

last data point within 1.5 times the interquartile range; outliers beyond that range are represented as with the symbol +.

Figure 3: Distribution of conditionally selected gene pairs across cancer types. a) Number of gene pairs affected by significant conditional selection in each cancer type. Gene pairs with reversed split and query genes were counted only once. No significant gene pairs were detected for sarcoma, thymoma, esophageal, renal (clear cell), liver, ovarian, and prostate cancers. b) Number of conditionally selected driver substitutions and total driver substitutions per tumor. Negative values can be interpreted as substitutions that could have been drivers in the absence of epistasis, but did not become established due to lack of selection. c, d) Association between the total number of driver substitutions per tumor (x axis) and (c) the number of significant gene pairs, (d) the number of conditionally selected driver substitutions (y axis). Each data point corresponds to a cancer type (n = 22). These associations remain highly significant after controlling for sample size (number of tumors and total mutations), median mutational load, and number of pairs tested. e) Conditional gene pairs found in multiple cancer types. f) Distribution of mutations along the PGM5 protein in stomach, colorectal, and other cancer types. The cartoon at the bottom indicates the location of alpha-D-phosphohexomutase, alpha/beta/alpha domains I, II, and III, with the predicted active site in red.

Figure 4: Epistasis networks based on conditional selection for some representative cancer types (see Suppl. Fig. 2 for the remaining cancer types). Nodes represent cancer genes involved in conditional selection; edges denote conditionally selected gene pairs. Cancer genes that are not involved in conditional selection are not represented, although they sometimes harbor a sizeable fraction of driver substitutions. Node size is proportional to the frequency of driver substitutions in a gene; edge width is proportional to the effect size; edge color represents the sign of epistasis; and arrows indicate the degree of symmetry (fully unidirectional arrows indicate lack of statistical significance and/or insufficient sample size for testing the reverse interaction).

Figure 5: Conditional selection and structure of cancer molecular subtypes. a) Effect sizes of conditional selection at the cancer type (x axis) and subtype (y axis) levels. Each data point corresponds to a gene pair tested in at least one subtype. Gray lines represent the orthogonal least squares fit for each cancer type. b) Simple scenario of cancer progression through the alteration of two independent pathways (each represented in a gray rectangle), generating a pattern of within-pathway mutual exclusivity and across-pathway co-occurrence. c) Number of signed triangular motifs in the epistasis networks of colon cancer (COAD), endometrial cancer (UCEC), and all cancer types. Gray bars correspond to randomized networks with the same size and signed degree distributions obtained by pairwise rewiring of the network edges, and serve as a null model for motif counts (error bars represent 95% confidence intervals based on 2000 randomizations). BRCA: breast cancer; LGG: low-grade glioma. d) A more complex scenario of cancer progression, compatible with the overrepresentation of (+,-,-) triangular motifs. In this scenario, gene-specific, across-pathway synergistic interactions can lead to the development of distinct cancer subtypes.

Figure 6: Modular epistasis in simulated networks leads to differentiation of cancer genomes into mutational subtypes. a) Tumor evolution was simulated in the presence of conditional selection, with epistatic interactions determined by networks with different degrees of modularity. For each network, the mutation composition of 500 simulated tumors was explored using principal component analysis to search for mutational differentiation into subtypes. b) Conditional selection

can generate clearly delineated mutational subtypes if the underlying epistasis network (top) is characterized by intra-module facilitation and inter-module interference. The bottom panels show the results of a principal component analysis for each network, with each point representing a simulated tumor.

Figure 7: Differences in patient survival associated to the modules of the epistasis network. Patients were assigned to the module with which they shared the greatest number of mutated genes. Patients without mutations in genes from the epistasis network were assigned to a null module (e.g., GBM-0). P-values correspond to the log-rank test for the null hypothesis that there are no differences among the three curves. GBM: glioblastoma; LGG: lower grade glioma; PAAD: pancreatic cancer; UCEC: endometrial cancer; HNSC: head and neck squamous carcinoma. Genes included in each module GBM-1: *ATRX, IDH1, RB1, TP53*; GBM-2: *PTEN*; LGG-1: *ATRX, IDH1, TP53*; LGG-2: *CIC, FUBP1*; LGG-3: *EGFR, PTEN*; LGG-4: *IDH2*; LGG-5: *NF1*; PAAD-1: *KRAS, TP53*; UCEC-1: *ARID1A, CTCF, CTNNB1, INPPL1, JAK1, KANSL1, KMT2B, KMT2D, KRAS, NFE2L2, PIK3R1, PTEN, RB1, RNF43, TAF1, ZFHX3, ATM, CCND1, FGFR2, UPF3A*; UCEC-5: *PIK3CA*; UCEC-6: *PPP2R1A, TP53*; HNSC-1: *CASP8, CTCF, EP300, EPHA2, FAT1, FBXW7, HLA-A, HLA-B, HRAS, RAC1, TGFBR2*; HNSC-2: *CDKN2A*.p14arf, *CDKN2A*.p16INK4a, *TP53*.

STAR Methods

Dataset and data preprocessing

Somatic mutation calls for The Cancer Genome Atlas (TCGA) dataset, generated by the Multi-Center
Mutation Calling in Multiple Cancers project(Ellrott et al., 2018), were downloaded from the Genomic
Data Commons repository of the National Institutes of Health (https://gdc.cancer.gov/about-
data/publications/mc3-2017). Clinical and survival data for TCGA were also downloaded from the
Genomic   Data   Commons   repository   of   the   National   Institutes   of   Health
(https://gdc.cancer.gov/node/905/; file TCGA-CDR-SupplementalTableS1.xlsx)(Liu et al., 2018). To
avoid the confounding effect of purifying selection, which is non- negligible in hypermutated
tumors(Persi et al., 2018), samples with >3000 coding mutations were classified as hypermutators
and removed from subsequent analyses.

Inference of conditional selection: the Coselens method

To quantify conditional selection at the gene level, we developed *Coselens* (*COnditional SELection on
the Excess of Nonsynonymous Substitutions*), a tool that makes extensive use of the *dndscv* R package
(dndSCV, RRID:SCR_017093) to quantify selection in cancer and normal tissues(Martincorena et al.,
2017). We briefly describe the fundamentals of the *dndscv* method, as long as it is needed to
understand the modifications introduced by *Coselens*; we refer the reader to the original publication
for further details. In short, *dndscv* estimates the number of missense and truncating (nonsense and
essential splice site) substitutions per gene that would be expected in the absence of selection by
combining a nucleotide substitution model with 192 rates (12 possible substitutions in each of the
16 possible trinucleotide contexts) and a set of genomic covariates, which greatly improve the
performance of the method at low mutation loads. The model parameters are estimated through
negative binomial regression on the number of synonymous mutations to maximize the following
likelihood function:

$$\mathcal{L}\left(\boldsymbol{r}, \boldsymbol{t}, \alpha, \beta; \boldsymbol{n}_s^{obs}\right) = \prod_{g,j} P_{Poisson}\left(n_{s,g,j}^{obs}; \lambda = t_g\, r_j\, L_{s,g,j}\right) P_{Gamma}(t_g; \alpha, \beta)$$

where $n_{s,g,j}^{obs}$ is the number of synonymous substitutions observed in the gene g and the index j
corresponds to the substitution classes (192 by default although substitution models with fewer or
more classes can be easily accommodated), r_j is the mutation rate for each substitution class, $L_{s,g,j}$
is the number of sites in gene g that are synonymous under the substitution class j, and t_g is a factor
accounting for variable mutation rates across genes, which is assumed to follow a gamma distribution
with parameters α and β. Once those parameters have been inferred, the expected number of
nonsynonymous substitutions in the absence of selection is obtained by replacing $L_{s,g,j}$ by the
number of sites in gene g that are susceptible to missense and truncating substitutions ($L_{m,g,j}$ and
$L_{t,g,j}$, respectively). The selection parameter $\omega_{k,g}$ is then calculated for each gene as the ratio
between the observed number of mutations and their neutral expectation ($\omega_{k,g} = \sum_j n_{k,g,j}^{obs}$ /
$\sum_j n_{k,g,j}^{exp}$, where the subscript $k \in \{m, t\}$ refers to missense and truncating substitutions). From the
perspective of the likelihood function, the selection parameter is a multiplicative factor in λ that
modifies the observed substitution frequency with respect to the neutral expectation, with $\omega_{k,g} > 1$
and $\omega_{k,g} < 1$ implying positive and purifying selection, respectively. The significance of $\omega_{k,g}$ is

computed through a likelihood ratio test, where the null and alternative hypotheses correspond to $\omega_{k,g} = 1$ and $\omega_{k,g} \neq 1$, respectively.

To develop *Coselens*, we modified the *dndscv* method in two substantial ways. First, we quantified the differences (rather than the ratios) between the observed number of missense and truncating substitutions and their neutral expectations:

$$\Delta n_{m,g} = \frac{\sum_j n_{m,g,j}^{obs} - \sum_j n_{m,g,j}^{exp}}{T} \quad ; \quad \Delta n_{t,g} = \frac{\sum_j n_{t,g,j}^{obs} - \sum_j n_{t,g,j}^{exp}}{T}$$

where T is the number of samples used in the analysis. Missense and truncating substitution excesses result from mutations that have reached detectable frequencies through positive selection. Because the effect of purifying selection in non-hypermutator tumors is negligible(Martincorena et al., 2017; Persi et al., 2018), the total excess of nonsynonymous mutations in a gene, $\Delta n_g = \Delta n_{m,g} + \Delta n_{t,g}$, is a good estimate of the number of driver substitutions N_d in that gene (formally, $N_d = \left((\omega_{m,g} - 1)n_{m,g}^{exp} + (\omega_{t,g} - 1)n_{t,g}^{exp}\right)/T = \Delta n_g$; note that we omitted the subscript g in N_d for simplicity although it is a gene-specific quantity). Second, we modified the likelihood ratio test to allow for comparison of Δn_g between two sets of samples. To this end, *Coselens* separately fits the *dndscv* substitution model to each sample group (hereafter *X* and *Y*), obtaining two sets of parameters for the neutral scenario ($\boldsymbol{\theta}(X)$ and $\boldsymbol{\theta}(Y)$) and two sets of nonsynonymous mutation excesses ($\Delta n_{k,g}(X)$ and $\Delta n_{k,g}(Y)$, with $k \in \{m,t\}$). The likelihood ratio test is built according to the following null and alternative hypotheses (2 degrees of freedom):

$$H_0: \Delta n_{m,g}(X) = \Delta n_{m,g}(Y) \; ; \; \Delta n_{t,g}(X) = \Delta n_{t,g}(Y)$$

$$H_1: \Delta n_{m,g}(X) \neq \Delta n_{m,g}(Y) \; ; \; \Delta n_{t,g}(X) \neq \Delta n_{t,g}(Y)$$

To restrict the hypothesis test to differences in the number of drivers, but not in the underlying mutation rates, the null hypothesis does not introduce any constraint on the parameters $\boldsymbol{\theta}(X)$ and $\boldsymbol{\theta}(Y)$. Accordingly, the likelihood of the null hypothesis is evaluated using group-specific parameters for each observation and joint nonsynonymous mutation excesses $\Delta n_{m,g}(X + Y)$ and $\Delta n_{t,g}(X + Y)$), for which the general expression is

$$\Delta n_{k,g}(X + Y) = \frac{\Delta n_{k,g}(X)\, T(X) + \Delta n_{k,g}(Y)\, T(Y)}{T(X) + T(Y)} \quad ; \quad k \in \{m,t\}$$

Here, $T(X)$ and $T(Y)$ are the numbers of samples in each group. The joint nonsynonymous excesses can be incorporated in the likelihood function by leveraging the relationship between the mutation excess $\Delta n_{k,g}$ and the mutation ratio $\omega_{k,g}$, that is, $\omega_{k,g} = 1 + \Delta n_{k,g} T/n_{k,g}^{exp}$. Under the null hypothesis, this relationship becomes

$$\omega_{k,g}(X|H_0) = 1 + \frac{\Delta n_{k,g}(X + Y)\, T(X)}{n_{k,g}^{exp}(X)} \quad ; \quad \omega_{k,g}(Y|H_0) = 1 + \frac{\Delta n_{k,g}(X + Y)\, T(Y)}{n_{k,g}^{exp}(Y)}$$

Thus, the likelihood of the null hypothesis becomes

$$\mathcal{L}(H_0) = \mathcal{L}\left(H_0, \boldsymbol{\theta}(X); \boldsymbol{n}_{k,g}^{obs}(X)\right) \mathcal{L}\left(H_0, \boldsymbol{\theta}(Y); \boldsymbol{n}_{k,g}^{obs}(Y)\right)$$

$$1 \quad \mathcal{L}\big(\mathrm{H}_0, \boldsymbol{\theta}; \boldsymbol{n}_{k,g}^{obs}\big) = \prod_{k,j} P_{Poisson}(n_{k,g,j}^{obs}; \lambda = t_g \, r_j \, L_{k,g,j} \omega_{k,g}(\cdot \,|\mathrm{H}_0)) \, P_{Gamma}(t_g; \alpha, \beta)$$

2   The expression for the likelihood of the alternative hypothesis is analogous, except that the $\omega_{k,g}$
3   parameters are calculated using sample-specific mutation excesses, that is,

$$4 \quad \omega_{k,g}(X|\mathrm{H}_1) = 1 + \frac{\Delta n_{k,g}(X)\, T(X)}{n_{k,g}^{exp}(X)} \; ; \quad \omega_{k,g}(Y|\mathrm{H}_1) = 1 + \frac{\Delta n_{k,g}(Y)\, T(Y)}{n_{k,g}^{exp}(Y)}$$

5   The output of the likelihood ratio test is the probability that the excess of nonsynonymous
6   substitutions (missense, truncating, or both) in a given gene differs between the two sets of samples.
7   As mentioned above, in the context of nearly absent purifying selection, such as in cancer evolution,
8   the effect size can be interpreted as the difference in the number of driver mutations per sample in
9   the gene of interest:

$$10 \quad \Delta N_d = N_d(X) - N_d(Y) = \; \Delta n_{m,g}(X) + \Delta n_{t,g}(X) - \Delta n_{m,g}(Y) - \Delta n_{t,g}(Y)$$

In our case, the sets *X* and *Y* correspond to samples with and without coding mutations in a second
gene; however, the *Coselens* method is general and can be applied to compare any pair of sample
sets (e.g. to study differential selection across tissues or clinical conditions). Although we focused on
genes under positive selection, *Coselens* can also detect differences in purifying selection if the excess
of substitutions is negative. To achieve the highest accuracy, *Coselens* should be run on whole exome
data or targeted sequencing data including as many genes as possible, as that boosts the
performance of parameter inference. The output can be subsequently filtered to restrict false
discovery rate correction and downstream analysis to a set of genes of interest.

We provide *Coselens* as an open-source R package that can be downloaded from
https://github.com/ggruenhagen3/coselens. Besides the main method described above, the package
includes some extensions to study the excess of small indels and test for differences in missense or
truncating substitutions only.

Conditional selection within cancer types

To investigate pairwise conditional selection among cancer genes, we compiled a set of candidate
gene pairs, to which we would apply the *Coselens* method. Each candidate gene pair consisted of a
split gene (the gene inducing conditional selection) and a *query gene* (the gene that experiences
conditional selection). Candidate gene pairs were separately identified in each cancer type; in that
sense, one can think of gene pairs as triplets (cancer type – split gene – query gene). Given a split
gene, conditional selection was assessed by dividing the samples into those that carried coding
mutations (nonsynonymous substitutions and small indels) in the split gene and those that do not,
comparing both groups with *Coselens*, and filtering the results according to the list of query genes
associated to that split gene.

A preliminary set of query genes was built by finding genes evolving under significant positive
selection in each cancer type with the R package *dndscv* (default parameters). From this preliminary
set, we built the set of split genes that harbored coding mutations in at least 25 samples, while being
free of coding mutations in at least another 25 samples. Then, we expanded the set of query genes
by running *Coselens* and searching for conditional selection on known cancer genes not included in

the preliminary set of query genes. For this exploratory step, we considered a list of 369 cancer genes from previous studies(Iranzo et al., 2018; Martincorena et al., 2017) and applied the Benjamini-Hochberg false discovery rate correction to all possible pairs consisting of a split gene and one of those known cancer genes. This approach identified 20 additional query genes, for which positive selection is only detectable in a restricted subset of samples (e.g., *HLA-C* and *PGM5* in *BRAF*-mutant colorectal cancer). We finally studied conditional selection on all possible split - query gene pairs within each cancer type, applying a false discovery rate correction to the pooled results (that is, jointly considering all cancer types). Additionally, to capture cases of reciprocal conditionality among gene pairs that could have been missed due to insufficient statistical power, we applied restricted hypothesis testing to split-query gene combinations whose reciprocal had been found significant under the standard approach.

Negative control

A control dataset was built by randomly assigning mock "mutated split gene" and "wild-type split gene" labels to each tumor. Relabeling was done such that the total number of mutations in the control subsets remained approximately the same as in the original ones. Specifically, for each cancer type, tumor samples were randomly chosen and assigned to the "mutated split gene" subset until the total number of mutations in the group reached or surpassed the number of mutations in the original subset of samples with mutations in the split gene. The reason to keep the number of mutations (rather than the number of samples) per group fixed is that *Coselens* pools together all mutations from all samples within a group. By keeping that number approximately unchanged, we ensured that the sensitivity of *Coselens* was similar in the negative control and in the original dataset. Furthermore, it is important to perform the randomization at the sample (rather than mutation) level to conserve the structure of within-sample correlations, so that the model covariates remain informative. Note that, because this randomization procedure removes systematic biases in mutational load and subtype composition between subsets, it is not suitable to assess the rate of false positives induced by such confounding factors. The entire workflow described in the previous section was repeated for 40 randomized control datasets, finding an average of 6.5 significant pairs per run (0.1% of the tested pairs). The comparison between this number and the value found in the original dataset (296 significant pairs) yields an empirical false discovery rate of ~2%.

Conditional selection within cancer subtypes

Samples from colorectal cancer were classified into molecular subtypes, and each subtype was separately analyzed as described above. The consensus molecular subtypes of colorectal cancer samples were obtained from the data repository of the colorectal cancer subtyping consortium (https://www.synapse.org/#!Synapse:syn2623706/files/)(Guinney et al., 2015). After removing hypermutated tumors (>3000 coding mutations), only 49 samples remained in the CMS1 group. Therefore, the sample size threshold to consider a candidate gene as a split gene in CMS1 was lowered to at least 18 samples with and 18 samples without coding mutations in that gene. A similar subtype-based analysis was applied to endometrial cancer, low-grade glioma, and breast cancer. The molecular subtype classification for those samples was obtained from the original publications(Cancer Genome Atlas Research et al., 2013; Ceccarelli et al., 2016) and the PanCancer Atlas dataset(Berger et al., 2018) of the cBioPortal repository(Cerami et al., 2012).

Classification of conditional pairs by into modes of epistasis

Gene pairs were classified by epistasis modes attending to the variation in the excess of nonsynonymous mutations in the first gene (A) when the second gene (B) does or does not harbor coding mutations ($N_d(A|B)$ and $N_d(A|\bar{B})$, respectively). First, we tested whether $N_d(A|B)$ or $N_d(A|\bar{B})$ were significantly distinct from zero. Within the *dndscv* framework, testing $N_d \neq 0$ is equivalent to testing whether the ratio $dn/ds \neq 0$. To account for multiple comparisons, we applied the Benjamini-Hochberg correction to all pairs tested for conditional selection and set the significance threshold to a false discovery rate of 5%. Next, we classified conditionally selected gene pairs in one of the following categories: (i) masking epistasis, if $N_d(A|\bar{B}) > N_d(A|B) = 0$; (ii) interference, if $N_d(A|\bar{B}) > N_d(A|B) > 0$; (iii) facilitation, if $N_d(A|B) > N_d(A|\bar{B}) > 0$; (iv) strict dependence, if $N_d(A|B) > N_d(A|\bar{B}) = 0$; (v) strict dependence with sign change, if $N_d(A|B) > 0 > N_d(A|\bar{B})$; (vi) compensation, if $N_d(A|\bar{B}) < N_d(A|B) = 0$; (vii) conditional deleterious, if $N_d(A|B) > N_d(A|\bar{B}) = 0$; and (viii) incompatible, if $N_d(A|\bar{B}) > 0 > N_d(A|B)$. Because the test for conditional selection can be more sensitive than the test for $N_d \neq 0$, there were some cases where $N_d(A|B) \neq N_d(A|\bar{B})$ but neither $N_d(A|B)$ nor $N_d(A|\bar{B})$ were significantly different from zero. Those cases were left unclassified.

Estimation of the number of driver substitutions affected by conditional selection

For a given cancer type and a split gene B, the average number of driver events per tumor that involve conditional selection on the query gene A was obtained as $\Delta N_d(A, B) \, X_B$, where $\Delta N_d(A, B) = N_d(A|B) - N_d(A|\bar{B})$ is the effect size of conditional selection and X_B is the fraction of tumors with coding mutations in the split gene B. To estimate the total number of driver events affected by conditional selection, we summed over all directed pairs (A, B) that were significant with FDR ≤ 5%, including reciprocal pairs that were significant under restricted hypothesis testing.

Estimation of the number of driver mutations per patient

Although the number of driver mutations can be estimated within the *dndscv* (and *Coselens*) framework, we chose an independent approach to minimize the risk that correlations with the number of conditional driver substitutions were affected by the presence of uncontrolled confounding factors. Therefore, we implemented a previously described simple regression-based approach(Iranzo et al., 2018). The number of driver mutations per patient was estimated as the intercept of a linear regression model that had, as independent variables, the cancer type, the number of coding mutations in passenger genes, and the interaction of both; and as the dependent variable, the number of coding mutations in 441 significantly mutated cancer genes (the union set of those previously detected(Martincorena et al., 2017) and those found in this study).

Functional association between genes in conditionally selected pairs

Each split-query gene pair was searched against the STRING database of protein interactions (v11.0). If an interaction was found in the database, the pair was assigned a score s equal to the STRING score divided by 1000 (this normalization accounted for the fact that STRING scores range from 0 to 1000, with higher values representing higher confidence in the predicted interaction). Pairs not found in the STRING database were assigned a score of zero. Given a subset G of gene pairs, the STRING interaction ratio was calculated as $r_G = \sum_G s / \sum_G (1 - s)$. In the special case when s takes a binary value (1 or 0), r_G becomes the ratio between the number of gene pairs that are known to interact and those for which there is no evidence of interaction. The STRING interaction ratio was computed

for increasing magnitudes of conditional selection by defining a nested collection of subsets G_x, with
$0 < x < \max(\Delta N_d)$, such that each subset includes all gene pairs whose effect size ΔN_d is greater
than x (accordingly, G_0 corresponds to the set of all gene pairs included in the analysis). The
interaction enrichment ratio for a given effect size was then calculated as $R(\Delta N_d) = r_{G_{\Delta N_d}}/r_{G_0}$.
Conceptually, the enrichment ratio is analogous to a risk ratio, with the difference that the response
variable is continuous rather than binary. Following Pagano and Gauvreau (2000), the 95%
confidence interval was calculated as $CI_{95}(\Delta N_d) = \exp\left(\ln(R) \pm 1.96 \times SE(\ln(R))\right)$, where
$SE(\ln(R)) = \sqrt{\dfrac{1}{\Sigma_{G_{\Delta N_d}}s} + \dfrac{1}{\Sigma_{G_{\Delta N_d}}(1-s)} - \dfrac{1}{\Sigma_{G_0}s} - \dfrac{1}{\Sigma_{G_0}(1-s)}}$.

Reciprocity of conditional selection

Let $P(X|\bar{Y})$ be the conditional probability that query gene X harbors a driver mutation provided that
there are no driver mutations in the split gene Y. Conversely, let $P(X|Y)$ be the conditional
probability that the query gene X harbors a driver mutation provided that there are driver mutations
in the split gene Y. Then, the magnitude of epistasis-driven conditional selection can be quantified
as the difference $\Delta P(X,Y) = P(X|Y) - P(X|\bar{Y})$. For any pair of genes A and B, we are interested in
finding an expression that relates $\Delta P(A,B)$ and $\Delta P(B,A)$, that is, the magnitude of conditional
selection when A is the query gene and B is the split gene, and when the query and split roles are
reversed. Combining Bayes theorem and the identity $P(X) + P(\bar{X}) = 1$, the difference $\Delta P(A,B)$ can
be expressed as

$$\Delta P(A,B) = P(A|B) - P(A|\bar{B}) = P(A)\left(\frac{P(B|A)}{P(B)} - \frac{1 - P(B|A)}{1 - P(B)}\right) = \frac{P(A)\left(P(B|A) - P(B)\right)}{P(B)\left(1 - P(B)\right)}$$

where $P(A)$ and $P(B)$ are the marginal probabilities for finding driver mutations in genes A and B,
respectively. In a similar way, $\Delta P(B,A)$ can be written as

$$\Delta P(B,A) = P(B|A) - P(B|\bar{A}) = P(B|A) - \frac{P(B) - P(B|A)P(A)}{1 - P(A)} = \frac{P(B|A) - P(B)}{1 - P(A)}$$

Extracting $P(B|A) - P(B)$ from both expressions, and after some algebraic manipulation, we obtain

$$\frac{\Delta P(A,B)}{P(A)(1 - P(A))} = \frac{\Delta P(B,A)}{P(B)(1 - P(B))} \qquad (1)$$

If presence or absence of driver mutations in genes A and B are modeled as independent Bernoulli
random variables with success probabilities $P(A)$ and $P(B)$, the denominators in eq. (1) correspond
to the variances of such random variables across tumors. Accordingly, the left and right terms of eq.
(1) can be interpreted as the standardized effect sizes of conditional selection for the query genes A
and B, respectively. Eq. (1) implies that, if conditional selection results from epistasis between driver
mutations, the standardized effect sizes should be equal for reciprocal query-split gene pairs.

Qualitatively, reciprocity implies that, if conditional selection is detected for a given split-query gene
pair, it should be also detected for the reverse pair, in which the split and query genes are exchanged.
Attending to this qualitative criterion, we found that 68% of the pairs that are subject to significant

conditional selection are reciprocal (this fraction does not include reverse pairs that could not be
tested due to insufficient number of coding mutations in one of the genes).

To carry out a more quantitative comparison between the theoretical prediction with the empirical
estimates, we first established a correspondence between the marginal and conditional probabilities
from eq. (1) and the nonsynonymous mutation excesses calculated from cancer genomic data. To
that end, we assumed that only the first driver substitution in each cancer gene is positively selected,
and therefore the probability to find a driver mutation in a given gene can be approximated by the
mean number of driver substitutions per tumor in that gene. Although this assumption is strictly valid
only for single-hit oncogenes, it also holds for tumor suppressor genes, as long as the inactivation of
the second copy of the gene can occur via large deletions. Thus, to validate eq. (1) with empirical
genomic data, we replaced $\Delta P(A, B) \approx \Delta N_d(A, B)$ and $P(A) \approx N_d(A|B) \times X_B + N_d(A|\bar{B}) \times (1 -$
$X_B)$, where X_B is the fraction of tumors with coding mutations in the split gene. This second
approximation potentially could become a major source of error because the empirical split of tumor
samples is based on the presence of coding mutations, whereas conditionality in the theoretical
expression specifically involves driver mutations. Therefore, calculating the goodness of fit of the
empirical data to the theoretical expectation in eq. (1) is a straightforward way to assess whether the
empirical estimates are robust with respect to the use of coding mutations rather than driver
mutations in the split gene to quantify conditional selection.

Assessment of the effect of differential mutation availability on conditional selection estimates

To investigate whether conditional selection can be explained by differences in mutation availability,
two alternative approaches were adopted. The first approach consisted of inferring, given the
expected number of mutations per tumor in the query gene, the probability that at least one driver
mutation becomes available for selection to act on. We considered a Poisson distribution for the
number of substitutions per gene per tumor, such that given a local mutation rate μ, the probability
that at least one driver mutation becomes available in the gene of interest is $\pi(\mu) = 1 - \exp(b\mu)$.
Given a set of *T* independent tumors, the number of tumors in which the driver mutation becomes
available (*k*) follows a binomial distribution with parameters *T* and $\pi(\mu)$. Following the central limit
theorem, for large enough *T*, the fraction of tumors in which the driver mutation becomes available
(*f* = *k/T*), can be approximated by a normal distribution with mean $\pi(\mu)$ and variance
$\pi(\mu)\big(1 - \pi(\mu)\big)/T$. Therefore, given two sets of values for *f*, μ, and *T* (corresponding to tumors with
and without mutations in the split gene), it is possible to apply a maximum likelihood approach to
infer the only unknown parameter of the model (*b*) and then obtain the probability that differences
in *f* can be explained by changes in μ. To perform these calculations, we took the expected number
of non-synonymous and non-sense mutations per sample per gene inferred with the dndscv method
as a gene-wise measure of μ; and the average number of drivers per gene per tumor as a gene-wise
measure of *f* (i.e., $f \approx N_d$, following the same rationale described above for the analysis of
reciprocity). A continuity correction was included when assessing the likelihood, such that *L*(*b*; *f*, μ,
T) = NormCDF(*f* + 0.5/*T*) − NormCDF(*f* - 0.5/*T*), where "NormCDF" is the cumulative probability
function for a normal distribution with mean $\pi(\mu)$ and variance $\pi(\mu)\big(1 - \pi(\mu)\big)/T$. For each
conditional gene pair, the probability that it can be explained by differential mutation availability was
calculated as the product $p = p_1 p_2$, where $p_1(f_1, \mu_1, T_1)$ is the probability that f_1 (the average
number of drivers per tumor in samples with a mutated split gene) comes from a normal distribution

with mean $\pi(\mu_1)$ and variance $\pi(\mu_1)\big(1 - \pi(\mu_1)\big)/T_1$ and $p_2(f_2, \mu_2, T_2)$ represents the same probability for samples without coding mutations in the split gene. In the case of reciprocal interactions, we combined the probabilities of the direct and reverse pairs (p^{AB} and p^{BA}, respectively) as $p = p^{AB}p^{BA}$. The resulting probabilities were corrected for multiple comparisons with FDR = 5%.

The second approach involved a regression analysis to assess how the excess of gene pairs displaying positive epistasis (compared to those that display negative epistasis) changes with the difference in the median mutational load of tumors with and without mutations in the split gene. By focusing on the difference between positive and negative epistatic pairs, rather than on the total numbers, we accounted for the fact that mutational load has a directed effect on mutation availability, that is, higher (respectively lower) mutational loads increase (decrease) mutation availability, resulting in an apparent excess of positive (negative) epistatic pairs. We found a weak, but significant association between the difference in mutational load and the excess of positive epistatic pairs in which a split gene is involved (0.16 pairs per every 100 mutations/exome difference in samples with and without coding mutations in the split gene; ANCOVA stratified by cancer types, $R^2 = 0.15$; p = 1e-9). To assess how this association could translate into apparent (mutation availability-driven) conditional selection, we adopted a Bayesian approach to infer the probability that a pair involving a given split gene can be attributed to differences in mutational load. First, we modeled the prior for the number of such pairs as a Poisson distribution with mean $\lambda = a \times \Delta\mu_g$, where a is the slope of the regression model and $\Delta\mu_g$ is the difference in the median mutational load of tumors that do and do not harbor mutations in the split gene of interest. To obtain the posterior distribution, we took into account the total number of conditional interactions observed for that split gene, k_g, by truncating the Poisson distribution between 0 and k_g. The expected contribution of differential mutation availability to the number of conditional pairs was then calculated as the mean of the truncated distribution.

Detection of modules in the epistasis networks

Positive conditional modules in the epistasis networks were identified with SiMap, a community detection algorithm that takes into account the sign and weight of interactions(Esmailian and Jalili, 2015). We used the effect sizes $\Delta N_d(A, B)$ as interaction weights and preprocessed the networks to extract their connected components before running the community detection algorithm with default parameters. Overlapping modules were identified by adding, for each gene, the weights of its interactions with genes from other communities. A secondary module assignation was made for those genes for which the total weight of interactions with the original community was <1.5 times the total weight of interactions with the secondary module. Samples were assigned to the module with which they shared the greatest number of mutated genes. Samples without mutations in genes from the epistasis network were assigned to a null module. Ambiguous cases (samples with the same number of genes mutated in >1 module) were resolved only if they met the following two conditions: (i) there were only 2 candidate modules, to which the sample could be assigned, and (ii) the fraction of mutated genes belonging to each module was different. If both conditions were met, the sample was assigned to the module with the largest fraction of mutated genes; otherwise, the module assignation was considered ambiguous.

Survival analysis

We conducted a survival analysis to investigate if the combination of mutations in two conditionally

selected genes affected patient survival, compared to mutations in a single gene. For each gene pair

(A, B) in a given cancer type, tumor samples were split in 3 groups: (i) those with coding mutations in

gene A but not in gene B, (ii) those with coding mutations in gene B but not in A, and (iii) those with

coding mutations in both genes (samples without mutations in A or B were not included). Gene pairs

that did not reach a minimum of 10 samples in each group and 100 samples overall were no further

considered. For those gene pairs that reached sufficient sample size, we adopted two approaches to

assess the statistical significance of differences in survival between the double-mutation group and

each of the single-mutation groups: (a) Kaplan-Meier estimation followed by non-parametrical log-

rank tests(Clark et al., 2003), and (b) Cox regression analysis with the double-mutation group as the

reference category(Bradburn et al., 2003). In both cases, we assessed the overall significance

(omnibus test for the null hypothesis that survival is the same in the three groups) and the

significance of the differences between each single-gene mutant and the double mutant. We

reported the cases with $p<0.1$ for the omnibus test and $p<0.15$ for both single vs double mutant

comparisons. Statistical analyses were carried out with the R packages *survminer* and

survival(Therneau and Grambsch, 2000).

The association between patient survival and positive epistatic modules with >15 samples was

assessed using the same approaches, and the cancer types reported in the text are those with $p<0.05$

for the omnibus tests.

Comparison of Coselens and SELECT methods

The list of conditionally selected events in the TGCA identified using the *SELECT* algorithm was

collected from the Supplementary Information of the original publication(Mina et al., 2017). To

reconcile the fact that *SELECT* operates at the level of "selected functional events" (SFE) and *Coselens*

operates at the gene level, each SFE was mapped to the gene or genes affected by that SFE. Pairs of

conditional SFE that mapped to the same gene (typically associated with double inactivation of tumor

suppressor genes) were not considered. Directionality was ignored and the significance threshold of

both methods was adjusted to ensure a FDR <10%, which was the value originally used by the

developers of *SELECT* when applying it to the TCGA dataset.

Simulation of tumor evolution

The evolution of a population of (pre)cancer cells was modeled as a discrete time birth-and-death

process, where death also accounted for differentiation into non-proliferative lineages. At each time

step, each cell could either die (with probability P_d) or divide (with probability $1 - P_d$). Following

previous work by McFarland et al(McFarland et al., 2013), the death probability was set to

34
$$P_d = \frac{D(N)}{B(d_i, d_e, p) + D(N)}$$

35   where $D(N) = N/K$ represents density-dependent death rate (N is the current population size and

36   parameter K determines the equilibrium population size in the absence of mutations). The birth term

37   was modeled as

38
$$B(d_i, d_e, p) = \frac{(1 + s_d)^{d_i}(1 + \epsilon s_d)^{d_e}}{(1 + s_p)^p}$$

where s_p is the fitness effect of a passenger mutation, s_d is the fitness effect of a driver mutation in the absence of epistasis, and ϵ is an epistasis coefficient that accounts for the increase in the fitness effect of a driver mutation due to positive epistasis. The variable p represents the number of passenger mutations in the genome of a given cell, d_i is the number of driver mutations that are not affected by epistasis, and d_e is the number of driver mutations affected by positive epistasis. We assumed a scenario, in which negative epistasis fully masks the effect of driver mutations, and therefore, driver mutations affected by negative epistasis do not contribute to the birth term.

Upon division, one of the daughter cells can acquire driver and passenger mutations. The number of new driver and passenger mutations was drawn from Poisson distributions with the means $T_d\mu$ (for drivers) and $T_p\mu$ (for passengers), where T_d and T_p are the mutation target sizes for drivers and passengers, respectively. New driver mutations were randomly assigned to the cancer genes from the epistasis network; if a gene had already been hit by a previous driver mutation, all subsequent driver mutations in that gene were treated as neutral.

To study the effect of epistasis in cancer genome evolution, we simulated 5 scenarios represented by the following epistasis networks: (i) no epistasis, represented as a "network" of 10 disconnected genes; (ii) 2 x 5-node star networks, with positive epistasis between the central and peripheral genes of each star; (iii) 2 x 5-node star networks as in (ii), adding negative epistasis between the central genes of each star; (iv) 2 x 5-node clique networks, with positive epistasis among all gene pairs from the same clique, and negative epistasis affecting half of all possible gene pairs from different cliques; and (v) 3 x 5-node "negative clique" network, with negative epistasis among all gene pairs from the same clique, and positive epistasis affecting 10% of all possible gene pairs from different cliques (this percentage was chosen to achieve a positive-to-negative interaction ratio that inversely mirrored the synergistic clique scenario).

Simulations started from a clonal population of size K with one driver mutation in a randomly selected gene and no passengers. We used the following parameter values: $K = 1000$, $\mu = 10^{-8}$, $T_d = 700$, $T_p = 5 \times 10^6$, $s_p = 10^{-3}$, $\epsilon = 5$. To facilitate comparisons, the parameter s_d was separately set in each scenario, such that the average fitness effect when considering epistasis was the same ($\langle s_d \rangle = 0.1$). Thus, in the disconnected and star networks (scenarios i-iii), we set $s_d = \langle s_d \rangle = 0.1$, in the clique network (scenario iv) $s_d = 2\langle s_d \rangle/(1 + \epsilon) \approx 0.03$, and in the negative clique network (scenario v) $s_d = 3\langle s_d \rangle/\epsilon = 0.06$. Simulations ended when the cell population doubled its initial size, became extinct, or after 30,000 generations. The final composition of those populations that doubled the initial size (representing tumor progression) was used to investigate differentiation into mutation-based subtypes by running a principal component analysis on the mutation composition matrix that represented the most frequent genotypes of 500 independently evolving populations.

Supplemental Information

Supplementary Figure 1 (related to Figure 1d): Comparison of *Coselens* and SELECT methods for the
detection of conditionally selected gene pairs, separately showing the overlap of both methods for
positive and negative epistasis. CNA: copy number alterations.

Supplementary Figure 2 (related to Figure 4): Epistasis networks for cancer types not shown in Figure
4. Nodes represent cancer genes involved in conditional selection; edges denote conditionally
selected gene pairs. Cancer genes that are not involved in conditional selection are not represented,
although they sometimes harbor a sizeable fraction of driver substitutions. Node size is proportional
to the frequency of driver substitutions in a gene; edge width is proportional to the effect size; edge
color represents the sign of epistasis; and arrows indicate the degree of symmetry (fully
unidirectional arrows indicate lack of statistical significance and/or insufficient sample size for testing
the reverse interaction).

Supplementary Figure 3: Kaplan-Meier curves showing association between conditional selection at
the gene pair level and differential survival. The p-value indicated by p_{all} corresponds to the log-rank
test for the null hypothesis that there are no differences among the three survival curves; p_1 and p_2
correspond to pairwise comparisons between the double mutant and each of the single mutants (log-
rank test).

Supplementary Table 1: Genes under significant levels of positive selection in at least one cancer
type, used as "query genes" in this study. For each gene, the selection parameters for missense,
nonsense, and indels, and the estimated number of driver mutations per tumor are shown. The table
also indicates if a query gene was additionally considered as a "split gene".

Supplementary Table 2: Gene pairs under significant conditional selection in at least one cancer type,
sorted by effect size. For each pair, the table shows the mean number of driver substitutions per
tumor in the query gene in the presence (Nd.with) and absence (Nd.without) of coding mutations in
the split gene, the magnitude of conditional selection (DeltaNd), the dependence index (theta), the
epistasis class, and the q-values for the null hypotheses Nd.with = 0, Nd.without = 0, and DeltaNd =
0.

Supplementary Table 3: Gene pairs that display mutual conditional selection, sorted by asymmetry.
For each pair, the table shows the dependence indices in both directions (theta_AB and theta_BA),
the asymmetry, the sign of epistasis, and the mean relative difference in variant allele frequency.

Supplementary Table 4: Analysis of conditional selection within cancer subtypes for breast cancer
(BRCA), colorectal cancer (COAD), endometrial cancer (UCEC), and lower grade glioma (LGG).
Statistical significance levels are coded as n.s. (not significant), * (p<0.05),** (p<0.01), and ***
(p<0.01).

References

Ashworth, A., Lord, C. J., and Reis-Filho, J. S. (2011). Genetic interactions in cancer progression and treatment. Cell *145*, 30-38.

Babur, O., Gonen, M., Aksoy, B. A., Schultz, N., Ciriello, G., Sander, C., and Demir, E. (2015). Systematic identification of cancer driving signaling pathways based on mutual exclusivity of genomic alterations. Genome biology *16*, 45.

Berger, A. C., Korkut, A., Kanchi, R. S., Hegde, A. M., Lenoir, W., Liu, W., Liu, Y., Fan, H., Shen, H., Ravikumar, V.*, et al.* (2018). A Comprehensive Pan-Cancer Molecular Study of Gynecologic and Breast Cancers. Cancer cell *33*, 690-705 e699.

Bradburn, M. J., Clark, T. G., Love, S. B., and Altman, D. G. (2003). Survival analysis part II: multivariate data analysis--an introduction to concepts and methods. British journal of cancer *89*, 431-436.

Brennan, C. W., Verhaak, R. G., McKenna, A., Campos, B., Noushmehr, H., Salama, S. R., Zheng, S., Chakravarty, D., Sanborn, J. Z., Berman, S. H.*, et al.* (2013). The somatic genomic landscape of glioblastoma. Cell *155*, 462-477.

Cancer Genome Atlas Research, N., Brat, D. J., Verhaak, R. G., Aldape, K. D., Yung, W. K., Salama, S. R., Cooper, L. A., Rheinbay, E., Miller, C. R., Vitucci, M.*, et al.* (2015). Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. The New England journal of medicine *372*, 2481-2498.

Cancer Genome Atlas Research, N., Kandoth, C., Schultz, N., Cherniack, A. D., Akbani, R., Liu, Y., Shen, H., Robertson, A. G., Pashtan, I., Shen, R.*, et al.* (2013). Integrated genomic characterization of endometrial carcinoma. Nature *497*, 67-73.

Cannataro, V. L., Gaffney, S. G., and Townsend, J. P. (2018). Effect Sizes of Somatic Mutations in Cancer. J Natl Cancer Inst *110*, 1171-1177.

Castro, A., Ozturk, K., Pyke, R. M., Xian, S., Zanetti, M., and Carter, H. (2019). Elevated neoantigen levels in tumors with somatic mutations in the HLA-A, HLA-B, HLA-C and B2M genes. BMC Med Genomics *12*, 107.

Ceccarelli, M., Barthel, F. P., Malta, T. M., Sabedot, T. S., Salama, S. R., Murray, B. A., Morozova, O., Newton, Y., Radenbaugh, A., Pagnotta, S. M.*, et al.* (2016). Molecular Profiling Reveals Biologically Discrete Subsets and Pathways of Progression in Diffuse Glioma. Cell *164*, 550-563.

Cerami, E., Gao, J., Dogrusoz, U., Gross, B. E., Sumer, S. O., Aksoy, B. A., Jacobsen, A., Byrne, C. J., Heuer, M. L., Larsson, E.*, et al.* (2012). The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov *2*, 401-404.

Ciriello, G., Cerami, E., Sander, C., and Schultz, N. (2012). Mutual exclusivity analysis identifies oncogenic network modules. Genome research *22*, 398-406.

Clark, T. G., Bradburn, M. J., Love, S. B., and Altman, D. G. (2003). Survival analysis part I: basic concepts and first analyses. British journal of cancer *89*, 232-238.

Cui, Q., Ma, Y., Jaramillo, M., Bari, H., Awan, A., Yang, S., Zhang, S., Liu, L., Lu, M., O'Connor-McCourt, M.*, et al.* (2007). A map of human cancer signaling. Mol Syst Biol *3*, 152.

D'Angelo, F., Ceccarelli, M., Tala, Garofano, L., Zhang, J., Frattini, V., Caruso, F. P., Lewis, G., Alfaro, K. D., Bauchet, L.*, et al.* (2019). The molecular landscape of glioma in patients with Neurofibromatosis 1. Nature medicine *25*, 176-187.

Decatur, C. L., Ong, E., Garg, N., Anbunathan, H., Bowcock, A. M., Field, M. G., and Harbour, J. W. (2016). Driver Mutations in Uveal Melanoma: Associations With Gene Expression Profile and Patient Outcomes. JAMA Ophthalmol *134*, 728-733.

DeGregori, J. (2017). Connecting Cancer to Its Causes Requires Incorporation of Effects on Tissue Microenvironments. Cancer research *77*, 6065-6068.

Di Micco, R., Fumagalli, M., Cicalese, A., Piccinin, S., Gasparini, P., Luise, C., Schurra, C., Garre, M., Nuciforo, P. G., Bensimon, A.*, et al.* (2006). Oncogene-induced senescence is a DNA damage response triggered by DNA hyper-replication. Nature *444*, 638-642.

Ellrott, K., Bailey, M. H., Saksena, G., Covington, K. R., Kandoth, C., Stewart, C., Hess, J., Ma, S., Chiotti, K. E., McLellan, M.*, et al.* (2018). Scalable Open Science Approach for Mutation Calling of Tumor Exomes Using Multiple Genomic Pipelines. Cell Syst *6*, 271-281 e277.

Esmailian, P., and Jalili, M. (2015). Community Detection in Signed Networks: the Role of Negative ties in Different Scales. Scientific reports *5*, 14339.

Garraway, L. A., and Lander, E. S. (2013). Lessons from the cancer genome. Cell *153*, 17-37.

Greaves, M., and Maley, C. C. (2012). Clonal evolution in cancer. Nature *481*, 306-313.

Guan, B., Wang, T. L., and Shih Ie, M. (2011). ARID1A, a factor that promotes formation of SWI/SNF-mediated chromatin remodeling, is a tumor suppressor in gynecologic cancers. Cancer research *71*, 6718-6727.

Guinney, J., Dienstmann, R., Wang, X., de Reynies, A., Schlicker, A., Soneson, C., Marisa, L., Roepman, P., Nyamundanda, G., Angelino, P.*, et al.* (2015). The consensus molecular subtypes of colorectal cancer. Nature medicine *21*, 1350-1356.

Hanahan, D., and Weinberg, R. A. (2000). The hallmarks of cancer. Cell *100*, 57-70.

Hanel, W., and Moll, U. M. (2012). Links between mutant p53 and genomic instability. J Cell Biochem *113*, 433-439.

Iranzo, J., Martincorena, I., and Koonin, E. V. (2018). Cancer-mutation network and the number and specificity of driver mutations. Proceedings of the National Academy of Sciences of the United States of America *115*, E6010-E6019.

Jerby-Arnon, L., Pfetzer, N., Waldman, Y. Y., McGarry, L., James, D., Shanks, E., Seashore-Ludlow, B., Weinstock, A., Geiger, T., Clemons, P. A.*, et al.* (2014). Predicting cancer-specific vulnerability via data-driven detection of synthetic lethality. Cell *158*, 1199-1209.

Jhunjhunwala, S., Hammer, C., and Delamarre, L. (2021). Antigen presentation in cancer: insights into tumour immunogenicity and immune evasion. Nature reviews Cancer *21*, 298-312.

Jiao, Y., Li, Y., Jiang, P., Han, W., and Liu, Y. (2019). PGM5: a novel diagnostic and prognostic biomarker for liver cancer. PeerJ *7*, e7070.

Kandoth, C., McLellan, M. D., Vandin, F., Ye, K., Niu, B., Lu, C., Xie, M., Zhang, Q., McMichael, J. F., Wyczalkowski, M. A.*, et al.* (2013). Mutational landscape and significance across 12 major cancer types. Nature *502*, 333-339.

Kim, M. P., Li, X., Deng, J., Zhang, Y., Dai, B., Allton, K. L., Hughes, T. G., Siangco, C., Augustine, J. J., Kang, Y. a.*, et al.* (2021). Oncogenic KRAS recruits an expansive transcriptional network through mutant p53 to drive pancreatic cancer metastasis. Cancer Discovery, candisc.1228.2020.

Kim, Y. A., Madan, S., and Przytycka, T. M. (2017). WeSME: uncovering mutual exclusivity of cancer drivers and beyond. Bioinformatics *33*, 814-821.

Kloor, M., Michel, S., Buckowitz, B., Ruschoff, J., Buttner, R., Holinski-Feder, E., Dippold, W., Wagner, R., Tariverdian, M., Benner, A.*, et al.* (2007). Beta2-microglobulin mutations in microsatellite unstable colorectal tumors. Int J Cancer *121*, 454-458.

Leiserson, M. D., Vandin, F., Wu, H. T., Dobson, J. R., Eldridge, J. V., Thomas, J. L., Papoutsaki, A., Kim, Y., Niu, B., McLellan, M.*, et al.* (2015). Pan-cancer network analysis identifies combinations of rare somatic mutations across pathways and protein complexes. Nat Genet *47*, 106-114.

Levine, R. L., Cargile, C. B., Blazes, M. S., van Rees, B., Kurman, R. J., and Ellenson, L. H. (1998). PTEN mutations and microsatellite instability in complex atypical hyperplasia, a precursor lesion to uterine endometrioid carcinoma. Cancer research *58*, 3254-3258.

Liu, J., Lichtenberg, T., Hoadley, K. A., Poisson, L. M., Lazar, A. J., Cherniack, A. D., Kovatich, A. J.,
Benz, C. C., Levine, D. A., Lee, A. V.*, et al.* (2018). An Integrated TCGA Pan-Cancer Clinical Data
Resource to Drive High-Quality Survival Outcome Analytics. Cell *173*, 400-416 e411.
Liu, X. Y., Gerges, N., Korshunov, A., Sabha, N., Khuong-Quang, D. A., Fontebasso, A. M., Fleming, A.,
Hadjadj, D., Schwartzentruber, J., Majewski, J.*, et al.* (2012). Frequent ATRX mutations and loss of
expression in adult diffuse astrocytic tumors carrying IDH1/IDH2 and TP53 mutations. Acta
Neuropathol *124*, 615-625.
Martincorena, I., Raine, K. M., Gerstung, M., Dawson, K. J., Haase, K., Van Loo, P., Davies, H.,
Stratton, M. R., and Campbell, P. J. (2017). Universal Patterns of Selection in Cancer and Somatic
Tissues. Cell.
Matlak, D., and Szczurek, E. (2017). Epistasis in genomic and survival data of cancer patients. PLoS
computational biology *13*, e1005626.
McConechy, M. K., Anglesio, M. S., Kalloger, S. E., Yang, W., Senz, J., Chow, C., Heravi-Moussavi, A.,
Morin, G. B., Mes-Masson, A. M., Australian Ovarian Cancer Study, G.*, et al.* (2011). Subtype-
specific mutation of PPP2R1A in endometrial and ovarian carcinomas. J Pathol *223*, 567-573.
McFarland, C. D., Korolev, K. S., Kryukov, G. V., Sunyaev, S. R., and Mirny, L. A. (2013). Impact of
deleterious passenger mutations on cancer progression. Proceedings of the National Academy of
Sciences of the United States of America *110*, 2910-2915.
Mina, M., Raynaud, F., Tavernari, D., Battistello, E., Sungalee, S., Saghafinia, S., Laessle, T., Sanchez-
Vega, F., Schultz, N., Oricchio, E., and Ciriello, G. (2017). Conditional Selection of Genomic
Alterations Dictates Cancer Evolution and Oncogenic Dependencies. Cancer cell *32*, 155-168 e156.
Misale, S., Di Nicolantonio, F., Sartore-Bianchi, A., Siena, S., and Bardelli, A. (2014). Resistance to
anti-EGFR therapy in colorectal cancer: from heterogeneity to convergent evolution. Cancer Discov
4, 1269-1280.
Motoyama, N., and Naka, K. (2004). DNA damage tumor suppressor genes and genomic instability.
Current opinion in genetics & development *14*, 11-16.
Neumeyer, V., Brutau-Abia, A., Allgauer, M., Pfarr, N., Weichert, W., Falkeis-Veits, C., Kremmer, E.,
Vieth, M., Gerhard, M., and Mejias-Luque, R. (2021). Loss of RNF43 Function Contributes to Gastric
Carcinogenesis by Impairing DNA Damage Response. Cell Mol Gastroenterol Hepatol *11*, 1071-1094.
Neumeyer, V., Grandl, M., Dietl, A., Brutau-Abia, A., Allgauer, M., Kalali, B., Zhang, Y., Pan, K. F.,
Steiger, K., Vieth, M.*, et al.* (2019). Loss of endogenous RNF43 function enhances proliferation and
tumour growth of intestinal and gastric cells. Carcinogenesis *40*, 551-559.
Nik-Zainal, S., Van Loo, P., Wedge, D. C., Alexandrov, L. B., Greenman, C. D., Lau, K. W., Raine, K.,
Jones, D., Marshall, J., Ramakrishna, M.*, et al.* (2012). The life history of 21 breast cancers. Cell *149*,
994-1007.
Noushmehr, H., Weisenberger, D. J., Diefes, K., Phillips, H. S., Pujara, K., Berman, B. P., Pan, F.,
Pelloski, C. E., Sulman, E. P., Bhat, K. P.*, et al.* (2010). Identification of a CpG island methylator
phenotype that defines a distinct subgroup of glioma. Cancer cell *17*, 510-522.
Oikonomou, E., Koustas, E., Goulielmaki, M., and Pintzas, A. (2014). BRAF vs RAS oncogenes: are
mutations of the same pathway equal? Differential signalling and therapeutic implications.
Oncotarget *5*, 11752-11777.
Pagano, M., and Gauvreau, K. (2000). Principles of biostatistics, 2nd edn (Pacific Grove, CA:
Duxbury).
Park, S., and Lehner, B. (2015). Cancer type-dependent genetic interactions between cancer driver
alterations indicate plasticity of epistasis across cell types. Mol Syst Biol *11*, 824.
Persi, E., Wolf, Y. I., Horn, D., Ruppin, E., Demichelis, F., Gatenby, R. A., Gillies, R. J., and Koonin, E.
V. (2021). Mutation-selection balance and compensatory mechanisms in tumour evolution. Nature
reviews Genetics *22*, 251-262.

Persi, E., Wolf, Y. I., Leiserson, M. D. M., Koonin, E. V., and Ruppin, E. (2018). Criticality in tumor evolution and clinical outcome. Proceedings of the National Academy of Sciences of the United States of America *115*, E11101-E11110.

Senft, D., Leiserson, M. D. M., Ruppin, E., and Ronai, Z. A. (2017). Precision Oncology: The Road Ahead. Trends Mol Med *23*, 874-898.

Shukla, S. A., Rooney, M. S., Rajasagi, M., Tiao, G., Dixon, P. M., Lawrence, M. S., Stevens, J., Lane, W. J., Dellagatta, J. L., Steelman, S.*, et al.* (2015). Comprehensive analysis of cancer-associated somatic mutations in class I HLA genes. Nature biotechnology *33*, 1152-1158.

Srihari, S., Singla, J., Wong, L., and Ragan, M. A. (2015). Inferring synthetic lethal interactions from mutual exclusivity of genetic events in cancer. Biol Direct *10*, 57.

Stratton, M. R., Campbell, P. J., and Futreal, P. A. (2009). The cancer genome. Nature *458*, 719-724.

Sun, Y., Long, H., Sun, L., Sun, X., Pang, L., Chen, J., Yi, Q., Liang, T., and Shen, Y. (2019). PGM5 is a promising biomarker and may predict the prognosis of colorectal cancer patients. Cancer Cell Int *19*, 253.

Szklarczyk, D., Gable, A. L., Nastou, K. C., Lyon, D., Kirsch, R., Pyysalo, S., Doncheva, N. T., Legeay, M., Fang, T., Bork, P.*, et al.* (2021). The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. Nucleic acids research *49*, D605-D612.

Therneau, T. M., and Grambsch, P. M. (2000). Modeling survival data : extending the Cox model, (New York: Springer).

Unni, A. M., Lockwood, W. W., Zejnullahu, K., Lee-Lin, S. Q., and Varmus, H. (2015). Evidence that synthetic lethality underlies the mutual exclusivity of oncogenic KRAS and EGFR mutations in lung adenocarcinoma. Elife *4*, e06907.

van de Haar, J., Canisius, S., Yu, M. K., Voest, E. E., Wessels, L. F. A., and Ideker, T. (2019). Identifying Epistasis in Cancer Genomes: A Delicate Affair. Cell *177*, 1375-1383.

Van Raamsdonk, C. D., Griewank, K. G., Crosby, M. B., Garrido, M. C., Vemula, S., Wiesner, T., Obenauf, A. C., Wackernagel, W., Green, G., Bouvier, N.*, et al.* (2010). Mutations in GNA11 in uveal melanoma. The New England journal of medicine *363*, 2191-2199.

Verhaak, R. G., Hoadley, K. A., Purdom, E., Wang, V., Qi, Y., Wilkerson, M. D., Miller, C. R., Ding, L., Golub, T., Mesirov, J. P.*, et al.* (2010). Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1. Cancer cell *17*, 98-110.

Vogelstein, B., Papadopoulos, N., Velculescu, V. E., Zhou, S., Diaz, L. A., Jr., and Kinzler, K. W. (2013). Cancer genome landscapes. Science *339*, 1546-1558.

Wang, X., Fu, A. Q., McNerney, M. E., and White, K. P. (2014). Widespread genetic epistasis among cancer genes. Nature communications *5*, 4828.

Wappett, M., Dulak, A., Yang, Z. R., Al-Watban, A., Bradford, J. R., and Dry, J. R. (2016). Multi-omic measurement of mutually exclusive loss-of-function enriches for candidate synthetic lethal gene pairs. BMC genomics *17*, 65.

Weghorn, D., and Sunyaev, S. (2017). Bayesian inference of negative and positive selection in human cancers. Nat Genet *49*, 1785-1788.

Wilkins, J. F., Cannataro, V. L., Shuch, B., and Townsend, J. P. (2018). Analysis of mutation, selection, and epistasis: an informed approach to cancer clinical trials. Oncotarget *9*, 22243-22253.

Williams, M. J., Werner, B., Heide, T., Curtis, C., Barnes, C. P., Sottoriva, A., and Graham, T. A. (2018). Quantification of subclonal selection in cancer from bulk sequencing data. Nat Genet *50*, 895-903.

Wu, R. C., Wang, T. L., and Shih Ie, M. (2014). The emerging roles of ARID1A in tumor suppression. Cancer Biol Ther *15*, 655-664.

Yeang, C. H., McCormick, F., and Levine, A. (2008). Combinatorial patterns of somatic gene
mutations in cancer. FASEB J *22*, 2605-2622.
Zhang, H., Deng, Y., Zhang, Y., Ping, Y., Zhao, H., Pang, L., Zhang, X., Wang, L., Xu, C., Xiao, Y., and Li,
X. (2017). Cooperative genomic alteration network reveals molecular classification across 12 major
cancer types. Nucleic acids research *45*, 567-582.
Zhang, S., and Yu, D. (2010). PI(3)king apart PTEN's role in cancer. Clin Cancer Res *16*, 4325-4330.

a all tested pairs

b 296 significant pairs

c strict dependence
facilitation
$\theta_{AB} > 0$
1:1
$\theta_{AB} < 0$
interference
masking

d facilitation (53%)
strict dependence (14%)
other (4%)
masking (11%)
interference (18%)

e masking
interference
independence
facilitation
strict depend.
strict dependence
facilitation
independence
interference
masking
dependence θ_{BA}
dependence θ_{AB}

f mean rel. diff. VAF
low (< 0.2)
high (≥ 0.2)
asymmetry $|\theta_{AB} - \theta_{BA}|$

a

epistasis network

cond. +
cond. -

500 simulations (mutation + selection)

...

mutation matrix

	gene A	gene B	gene C	gene D
tumor 1	■			
tumor 2	■	■		
tumor 3			■	■
...				
tumor 500			■	■

identification of subtypes (PCA)

b

epistasis network

mutation-based subtypes

PC2

PC1

14%   27%   28%   54%   11%

% mutational variance explained by PC1