The genomic landscape of contemporary western Remote Oceanians

Lara R. Arauna[a,1], Jacob Bergstedt[a,b,c], Jeremy Choin[a,d,e], Javier Mendoza-Revilla[a], Christine Harmant[a], Maguelonne Roux[a,f], Alex Mas-Sandoval[g], Laure Lémée[h], Heidi Colleran[i], Alexandre François[j], Frédérique Valentin[k], Olivier Cassar[l], Antoine Gessain[l], Lluis Quintana-Murci[a,e,1,2], Etienne Patin[a,1,2]

[a]Human Evolutionary Genetics Unit, Institut Pasteur, UMR 2000, CNRS, Paris 75015, France
[b]Institute of Environmental Medicine, Karolinska Institutet, Stockholm 171 77, Sweden
[c]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm 171 77, Sweden
[d]Université de Paris, Paris 75015, France
[e]Chair Human Genomics and Evolution, Collège de France, Paris 75005, France
[f]Bioinformatics and Biostatistics Hub, Institut Pasteur, Université de Paris, Paris 75015, France
[g]Department of Life Sciences, Imperial College London, Ascot SL5 7PY, United Kingdom
[h]Biomics Platform, Institut Pasteur, Paris 75015, France
[i]BirthRites Independent Max Planck Research Group, Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Leipzig 04103, Germany
[j]Langues, Textes, Traitements Informatiques, Cognition (LaTTiCe), UMR 8094, CNRS, Paris 75015, France
[k]MSH Mondes, UMR 7041, CNRS, Nanterre 92023, France
[l]Oncogenic Virus Epidemiology and Pathophysiology Unit, Institut Pasteur, UMR 3569, CNRS, Paris 75015, France

[1]To whom correspondence may be addressed. Email: lrubioar@pasteur.fr (L.R.A.); quintana@pasteur.fr (L.Q.-M.); epatin@pasteur.fr (E.P.)

[2]L.Q.-M. and E.P. contributed equally to this work.

Classification: Biological Sciences, Anthropology

Keywords: Pacific, populations, migrations, admixture, genetics, cultural practices

This PDF file includes:
    Main Text
    Figures 1 to 4

Abstract

The Vanuatu archipelago served as a gateway to Remote Oceania during one of the most extensive human migrations to uninhabited lands ~3,200 years ago. Ancient DNA studies suggest an initial settlement by East Asian-related peoples that was quickly followed by the arrival of Papuan-related populations, leading to a major population turnover. Yet, there is uncertainty over the population processes and the sociocultural factors that have shaped the genomic diversity of ni-Vanuatu, who present nowadays among the world's highest linguistic and cultural diversity. Here, we report genome-wide data for 1,433 contemporary ni-Vanuatu from 29 different islands, including 287 couples. We find that ni-Vanuatu derive their East Asian- and Papuan-related ancestry from the same source populations and descend from relatively synchronous admixture events that occurred ~1,700-2,300 years ago, indicating a peopling history common to all the archipelago. However, our analyses reveal that the Papuan population turnover was geographically uneven, and that the genetic contribution of Papuan-related peoples was male-biased. Furthermore, we detect Polynesian ancestry arriving ~600-1,000 years ago to South Vanuatu, and map its distribution to both Polynesian- and non-Polynesian-speaking islands. Lastly, we provide evidence for a tendency of spouses to carry similar genetic ancestry, when accounting for relatedness avoidance. The signal is not driven by strong genetic effects of specific loci or trait-associated variants, suggesting that it results instead from social assortative mating. Altogether, our findings provide insight into both the genetic history of ni-Vanuatu populations and how sociocultural processes have shaped the diversity of their genomes.

Significance statement

Vanuatu served as a gateway for the peopling of Remote Oceania but how demography and culture have shaped the genomic diversity of its peoples remains partially understood. Here, we report genomic data for 1,433 contemporary ni-Vanuatu covering 29 islands. We demonstrate that their genomic diversity derives from relatively synchronous, geographically uneven and sex-biased admixture events between the Austronesian-related first settlers and the Papuan-related subsequent incomers. We show that the more recent arrival of Polynesian ancestry did not always trigger shifts to Polynesian languages. Lastly, we find evidence for non-random mating, probably owing to social assortative mating. Our findings reveal the impact of multiple migrations, admixture and sociocultural practices on the genomic landscape of a key region of the South Pacific.

Introduction

Vanuatu, an archipelago located in western Remote Oceania, is a key region for understanding the peopling history of the Pacific. The cultural, anthropological and genetic diversity of Vanuatu reflects three distinct phases of population movements. The first, which started in present-day Taiwan ~5,000 ya, was associated with the spread of Austronesian languages to Near and Remote Oceania (1-3). The so-called Austronesian expansion led to the emergence of the well-characterized Lapita cultural complex, which emerged in the Bismarck Archipelago and reached Vanuatu 3,200 ya (4, 5). Morphometric and ancient DNA (aDNA) studies indicate that the Lapita people in Vanuatu carried East Asian ancestry, supporting a connection with the Austronesian expansion (6, 7). The second migration occurred at the end of the Lapita period, ~2,500 ya, and involved the arrival of Papuan-related peoples who shared ancestry with contemporary Bismarck Archipelago islanders. aDNA studies have shown that these migrations triggered a dramatic shift in genetic ancestry, from the East Asian-related ancestry associated with the Lapita culture to the Papuan-related ancestry that has remained predominant since then (8, 9). Finally, it has been postulated that "Polynesian outliers" from Vanuatu (i.e., Polynesian speakers living outside the Polynesia triangle) are the descendants of migrations from Polynesia into western Remote Oceania (10, 11), a model that has received recent support (12).

Although aDNA studies have revealed that ni-Vanuatu are descended from at least three ancestral populations (8, 9, 12), whether the settlement process was uniform across the multiple islands of the archipelago remains an open question. aDNA data indicate that individuals dated to ~2,500 ya from South and Central Vanuatu carried largely different proportions of East Asian-related ancestry (8, 9, 12), in line with either a unique population turnover that was geographically heterogeneous or separate admixture events between different populations across islands. Furthermore, Vanuatu is the country with the world's largest number of languages per capita (13), which supports the view that languages have rapidly diversified since the initial settlement and/or that the arrival of new groups to the archipelago further increased linguistic diversity. Nevertheless, these questions have been difficult to resolve because, to date, available ancient and modern DNA data from the region have remained sparse (8, 9).

Here, we generated genome-wide genotype data for 1,433 contemporary ni-Vanuatu and assessed their fine-scale genetic structure, in order to address central questions about the settlement of western Remote Oceania: do all contemporary ni-Vanuatu derive their ancestry from three populations only? Was the contribution of these three ancestral populations

different across the archipelago? Was the late-Lapita shift to Papuan-related ancestry

heterogeneous across islands? In addition, the recent settlement of Vanuatu allows for the

study of how socio-cultural practices have shaped genetic diversity in humans over the past

~3,000 years, in the form of sex-biased admixture, relatedness avoidance and non-random

mating (14-16). We thus used the comprehensive set of population genomics data presented

here, which includes 287 male-female couples, to answer important questions relating to the

genetic history of a population that descends from diverse ancestral populations: Was

admixture sex-biased? Was admixture accompanied by language shifts? Has socio-cultural

structure influenced mating? Can residence rules and urbanization affect human genetic

structure?

Results

The Genetic Variation of Ni-Vanuatu is Spatially Structured. To shed light on the genetic make-up of ni-Vanuatu, we collected >4,000 blood samples from contemporary individuals between 2003 and 2005, and genotyped a selection of 1,433 of these sampled individuals at 2,358,955 single nucleotide polymorphisms (SNPs). After quality filters and merging with high-coverage whole genome sequencing data (17), 1,439 ni-Vanuatu samples were kept, including those from 522 males and 917 females living in 29 islands and 179 different villages (Fig. 1*A* and Table S1).

Principal Component (PCA) and ADMIXTURE analyses indicate that ni-Vanuatu fall on a genetic gradient between East Asian-related and Papuan-related populations (Fig. 1*B* and Fig. S1), supporting the view that their ancestry derives from these two population groups (6, 8, 9, 12). Furthermore, ni-Vanuatu show high genetic similarities with individuals from the Bismarck Archipelago, in line with the hypothesis that their Papuan-related ancestors originated from these islands (6, 8, 9, 12, 17). However, the extensive geographic coverage of the dataset presented here enabled us to reveal substantial genetic substructure among ni-Vanuatu islanders. PCA showed that genetic variation in ni-Vanuatu is explained by two contiguous but distinct groups that reflect geography (Fig. 1*C* and Fig. S2). Accordingly, when considering the number of ancestral populations that is most supported by ADMIXTURE ($K_{ADM} = 10$), populations from different islands also show different ancestral components (Fig. S1). These results reveal that contemporary ni-Vanuatu show substantial genetic differentiation (Fig. S3), which could result either from different admixture histories during the settlement period and/or from the existence of barriers to gene flow that formed after the settlement of the archipelago.

To gain further insights into the genetic history of ni-Vanuatu, we next assessed the fine-scale genetic structure of the 1,439 sampled individuals, using ChromoPainter and fineSTRUCTURE (18-20). Haplotype-based clustering revealed a first separation between ni-Vanuatu living north and south of the strait separating Epi and Tongoa islands ($K_{FS} = 2$; Fig. S4). At $K_{FS} = 4$, populations separate into four clusters, here referred to as North, South, Central East and Central West Vanuatu clusters (Fig. 2*A* and Fig. S4 and Table S2), in general agreement with the classification of Oceanic languages spoken in the archipelago (21-23). Remarkably, individuals from the north and south of Pentecost island cluster with individuals from different neighboring islands, indicating that the sea does not necessarily act as a barrier to gene flow in the region, in line with linguistic and ethnographic data that reveal cultural networks between islands (21, 22, 24, 25). At $K_{FS} = 20$, the highest K_{FS} value for which

statistical robustness remains maximal (Material and Methods) (19), we found that genetic

clusters are often island-specific (Fig. 2*A*, Figs. S4-S7 and Table S1). These observations

suggest that clusters inferred by fineSTRUCTURE are reliable, as they reflect expected

geographic and linguistic barriers to gene flow.

The Late-Lapita Shift to Papuan-related Ancestry was Geographically Uneven. We

detected important differences among fineSTRUCTURE clusters in the proportions of East

Asian-related ancestry, that is, ancestry related to contemporary populations from either

Taiwan and the Philippines or Polynesia (range of SOURCEFIND estimates: 0.15 to 0.55;

Fig. 2*C*, Figs. S8-S11 and Table S3). East Asian-related ancestry was found to be lowest for

the Central Vanuatu clusters (e.g., Malekula, Ambrym, Epi; median = 0.230, SD = 0.056) and

highest in the South Vanuatu cluster (i.e., Efate, Tongoa, Ifira and Emae islands; median =

0.323, SD = 0.077), where "Polynesian outlier" communities live today (10). Nevertheless,

East Asian-related ancestry is also high in islands where Polynesian ancestry is low, such as

Ambae (median = 0.405, SD = 0.042; Fig. 2*D*), suggesting that differences in East Asian-

related ancestry are not due solely to differences in Polynesian ancestry (Figs. S10 and S11

and Table S3).

To assess whether differences in East Asian-related ancestry originate from a

geographically uneven ancestry turnover or separate admixture events between distinct

populations, we dated admixture in each genetic cluster separately. Besides more recent

events relating to Polynesian migrations (see next section), all estimates overlap the same

time period that ranges from 1,700 to 2,300 ya (Fig. 2*B*, Fig. S12 and Table S4), suggesting

that all ni-Vanuatu share the same admixture history. To test this hypothesis more formally,

we evaluated if the Papuan-related ancestry carried by present-day ni-Vanuatu derives from a

single source. PCA and f_4-statistics of the form f_4(*X*, New Guinean highlanders; Solomon or

Bismarck Archipelago islanders, East Asians) indicate that all ni-Vanuatu show similar

genetic relatedness with populations from New Guinea, the Bismarck Archipelago or the

Solomon islands (Figs. S13 and S14), in agreement with aDNA data (12). Furthermore, the

haplotype-based SOURCEFIND method detected that the same cluster of Bismarck

Archipelago islanders is the source that contributed the most to all ni-Vanuatu (Fig. S15 and

Table S5). Of note, SOURCEFIND also detected a small contribution from New Guinean

highlanders and Santa Cruz islanders in all ni-Vanuatu clusters, which negatively correlates

with Polynesian ancestry (r = -0.378, *P*-value < 2.22×10^{-16}), probably because Polynesian

source populations capture some Papuan-related ancestry. Collectively, these findings

indicate that ni-Vanuatu are descended from relatively synchronous admixture events between

the same sources of Papuan- and East Asian-related ancestry; yet, their proportions differ

markedly across islands, indicating that the dramatic Papuan-related ancestry shift that started

~2,500 ya was geographically uneven.

Polynesian Migrations Did Not Necessarily Trigger Language Shifts. Admixture

proportions and date estimates indicate that ni-Vanuatu ancestry partly derives from a third

and more recent migration originating from Polynesia (Fig. 2*B-D* and Tables S3 and S4). We

dated admixture events between 600 and 1,000 ya for genetic clusters predominant in Efate,

Ifira, Emae and Futuna islands (Fig. 2*B*, Fig. S12 and Table S4), where Polynesian languages

are spoken today (26). These clusters show higher Polynesian ancestry and, therefore, higher

East Asian-related ancestry proportions, relative to North and Central Vanuatu clusters

(SOURCEFIND estimates; Fig. 2*C-D*, Figs. S8-S11 and Table S3). Ni-Vanuatu Polynesian

speakers show a higher ratio of Polynesian-to-East Asian ancestry, when compared to non-

Polynesian speakers (ratio = 0.647 *vs.* 0.479; Wilcoxon test *P*-value < 2.22×10^{-16}; Fig. 2*D* and

Fig. S10). Furthermore, f_4-statistics of the form f_4(*X*, East Asians; Tongans, New Guinean

highlanders) suggest that ni-Vanuatu from Futuna, Ifira, Efate, Emae and Tongariki share

more alleles with Polynesians than other ni-Vanuatu do (Fig. S16).

Interestingly, our analyses also revealed that Polynesian ancestry is not restricted to

Vanuatu islands where Polynesian languages are spoken. Non-Polynesian-speaking

populations assigned to South Vanuatu (e.g., Tongoa, Tongariki and Tanna) and North

Vanuatu (i.e., Banks and Torres islands) clusters also show a higher Polynesian-to-East Asian

ancestry ratio than Central East and West ni-Vanuatu (ratio = 0.557 and 0.510, *vs.* 0.461;

Wilcoxon test *P*-value < 2.22×10^{-16}; Fig. 2*D*, Figs. S10 and S11 and Table S3). Furthermore,

estimated admixture dates are similar among South Vanuatu clusters where both Polynesian

and non-Polynesian languages are spoken (Fig. 2*B*, Fig. S12 and Table S4). These results

support the view that Polynesian migrations, as well as subsequent contacts between Vanuatu

islands (26, 27) or with "Polynesian outliers" from the Solomon islands (28), also introduced

Polynesian ancestry among non-Polynesian-speaking groups.

Conversely, we found no evidence of admixture with Polynesians in individuals assigned

to Central Vanuatu clusters, including Epi islanders, despite the geographic proximity

between Epi and Tongoa (Fig. 2*D*, Figs. S10 and S11 and Table S3). Of note, ADMIXTURE,

PCA and fineSTRUCTURE analyses separate ni-Vanuatu into northern and southern

populations, the frontier between the two being located between Epi and Tongoa (Figs. 1*C*

and 2*A* and Fig. S1). The strait that separates the two islands today is the location of the

Kuwae caldera (29, 30), which may have been a barrier to gene flow. Together, these results

reveal that, since 1,000 ya onward, Polynesians migrated to Vanuatu where they admixed

with local populations, and that such interactions did not necessarily result in a shift to

Polynesian languages.

A Limited Genetic Impact of European Colonization. Despite the relatively long history of

European colonization of Vanuatu, which started in 1606 CE (31, 32), our genetic data

indicate that admixture between ni-Vanuatu and Europeans has been rare or has left few

descendants in Vanuatu. In total, only 28 individuals out of 1,439 (1.95%) show a genetic

contribution from Europeans higher than 1%, according to SOURCEFIND analyses (range:

0.03% to 35.3%; Fig. S8 and Table S3). The two fineSTRUCTURE clusters with the highest

European ancestry (median = 0.093 and 0.107, *vs.* 0.002 for other clusters) show an admixture

event in the last 120 years (mean = 78 ya, 95%CI: [74–81] and mean = 111 ya, 95%CI: [104-

118]) (Fig. 2*B* and Table S4). Similarly, the three other clusters that include individuals

carrying more than 1% of European ancestry show a pulse of admixture occurring in the last

200 years. These results are consistent with historical records, which indicate that contacts

between Europeans and ni-Vanuatu were initially rare but became more common when

Christian missionaries and colonists settled in the archipelago during the 19[th] century, when

the first intermarriages were reported (31, 32).

Genetic Admixture in Vanuatu was Sex-Biased. Genetic studies have suggested that

Papuan-related migrations from the Bismarck Archipelago into Remote Oceania were male-

biased, because contemporary Polynesians and a few ancient individuals from present-day

Vanuatu show lower Papuan-related ancestry on the X chromosome, relative to autosomes (6,

12). To confirm that admixture between the ancestors of present-day ni-Vanuatu was sex-

biased, we estimated Papuan- and East Asian-related ancestry in ni-Vanuatu on each

chromosome separately, using local ancestry inference (33). We found that Papuan-related

ancestry is indeed significantly lower on the X chromosome, relative to autosomes (α_X =

75.2% *vs.* α_{auto} = 79.8%, Wilcoxon test *P*-value $< 1.36 \times 10^{-5}$; Fig. 3*A*), confirming aDNA

results (12). These values were similar between Polynesian and non-Polynesian speakers (Fig.

S17*A*, Wilcoxon test *P*-value = 0.13), which indicates that it is not explained by recent

migrations from Polynesia. We replicated the results using high-coverage genome sequences

from a subset of 179 ni-Vanuatu (17), indicating that our results are not biased due to SNP

ascertainment (Fig. S17B). Assuming that admixture proportions have reached equilibrium

values (34), we estimated that the genetic contribution of Papuan-related males to ni-Vanuatu

was 27% higher than that of Papuan-related females (α_m = 93.5% *vs.* α_f = 66.1%).

Accordingly, Y chromosomes of ni-Vanuatu are dominated by haplogroups found at high

frequency in Near Oceanians (e.g., M1a3b2, S1), whereas mitochondrial DNAs (mtDNA)

show a high proportion of haplogroups typically found in East Asia (e.g., B4a1a1, E1a2a4)

(Fig. 3B and Table S1). Collectively, these results indicate that ni-Vanuatu ancestry

predominantly results from admixture between Papuan-related males and East Asian-related

females.

Recent Migrations are Influenced by Residence Rules and Urbanization. The genetic

structure of contemporary ni-Vanuatu is also expected to reflect socio-cultural practices (e.g.,

social networks, exchange and marriage rules) that have culturally evolved since the

settlement of the archipelago. We leveraged the high-resolution genetic data to infer recent

migrations between Vanuatu islands – indicated by individuals who inhabit an island but

belong to a genetic cluster that is prevalent in another island – and determine if these

migrations have involved mainly females or males, in line with virilocal or uxorilocal

postmarital residence rules, respectively. We found that 5.70% of the sampled individuals (54

individuals) migrated at a large geographical scale (K_{FS} = 4), while 11.81% (112 individuals)

migrated at a local scale (K_{FS} = 20; Table S6), suggesting more genetic connections between

closer islands. We estimated that local mobility among females is higher than among males

(K_{FS} = 20; odds ratio = 2.20, Fisher's exact test P-value = 9.16×10^{-4}; Fig. 3C), as expected

under virilocal residence and/or female exogamy (35). The same trend was observed at a

larger geographical scale, when considering migrations between four broader regions (K_{FS} =

4; odds ratio = 1.86, P-value = 0.075) and when restricting the analyses to the reported birth

place of male-female couples included in the dataset (Fig. S18) (odds ratio = 1.96, P-value =

6.89×10^{-3}). Notably, comparisons of the places of birth and residence of sampled individuals

did not support female-biased migrations (odds ratio = 0.807, P-value = 0.18; Fig. S19 and

Table S6), possibly reflecting a bias in the self-reported birth places. This might occur if

women are inclined to report their husband's or children's place rather than their father or

mother's place (36).

We also explored the direction of the inferred migrations, and found that both female and

male migrations mainly occurred from the northernmost and southernmost islands towards the

center of the archipelago (Fig. 3C), where Port Vila – the largest city – has developed (37).

Migrations are also common among Central Vanuatu islands, consistent with a long-term

network of cultural and material exchanges in the region (28). Thus, our genomic data reflect

the migration patterns that characterize the recent history of ni-Vanuatu, including residence

rules and urbanization.

Evidence for Relatedness Avoidance. Human kinship systems vary tremendously,

regulating marriage, exchange, endogamy or exogamy according to indigenous relational

concepts (38). Given the small census size of ni-Vanuatu populations, it is interesting to

consider whether and how sociocultural features like marriage and exchange rules influence

local levels of genetic relatedness. We found that genetic relatedness was higher within

islands than between islands (Wilcoxon test P-value $< 2.22 \times 10^{-16}$; Figs. S20-S22), consistent

with an isolation by distance model (Mantel test P-value $= 0.001$; 1,000 permutations).

Accordingly, out of the 287 male-female couples included in the dataset, 78.4% were born on

the same island (Fig. S23). Genetic relatedness is also higher among inhabitants of the same

village, relative to individuals living on the same island (Wilcoxon test P-value $< 2.22 \times 10^{-16}$)

(Fig. S20), suggesting that the local community is often the source of marriage partners.

However, despite the generally high levels of genetic relatedness observed between ni-

Vanuatu, we did not observe an excess of genetic relatedness among couples (Figs. S23 and

S24). Specifically, spouses tend to show slightly lower kinship coefficients than random pairs

of individuals from the same village, after excluding first-degree related individuals (logistic

regression model P-value $= 0.057$; resampling P-value $= 0.079$) (Figs. S24 and S25). These

results indicate that the practice of exogamy has affected the genetic diversity of ni-Vanuatu,

and provide a vivid example that human populations can show high levels of genetic

relatedness in the absence of endogamous marriage practices.

Ni-Vanuatu Spouses Tend to Share Similar Genetic Ancestry. Research on admixed

populations from other parts of the world has shown that, in addition to low genetic

relatedness, partners tend to show similar genetic ancestry (15), because mating often occurs

within socio-cultural groups, which can correlate, in turn, with genetic ancestry. To verify

whether this phenomenon is observed in Vanuatu, we implemented a logistic regression

model that jointly estimates the effects of geography, genetic relatedness and genetic ancestry

on the probability to be partners (Material and Methods). These analyses confirmed that

spouses tend to originate from the same island, while showing lower kinship coefficients than

non-spouses (Fig. 4*A*). Importantly, we found that spouses show lower differences in genetic

ancestry than non-spouses, when considering Papuan-related (β = -4.233, *P*-value = 0.018),

East Asian-related (β = -4.221 *P*-value = 0.018) or Polynesian (β = -5.197, *P*-value = 0.024)

ancestry. These results suggest that ni-Vanuatu tend to mate with a partner who carries similar

genetic ancestry.

Two hypotheses have been proposed to explain these observations. First, social assortative

mating may underlie the observed signal, as genetic ancestry can correlate with socio-cultural

structure. Second, spouses may choose their partner because they share biological traits, such

as physical appearance (39, 40). To test these hypotheses, we searched for genomic loci that

could play a role in partner choice, by including, in the logistic regression model, a term that

measures dissimilarity between individuals at each SNP (Material and Methods). No SNP

showed statistical evidence for a significantly lower or higher genotype similarity between

spouses, when accounting for multiple testing (Fig. S26). To test for assortative mating

according to polygenic traits, we then evaluated if genotype similarity between spouses is

significantly higher or lower at SNPs associated with candidate traits relating to physical

appearance, when compared to non-associated SNPs. When we did not account for genetic

structure and ancestry-associated assortative mating (i.e., effect sizes are estimated from a

model where these confounders are not included, see Material and Methods), we found

evidence for assortative mating according to body mass index (BMI, Fig. 4*B*). However,

when accounting for such possible confounders, we found no statistical evidence that

genotypes at trait-associated variants are more similar or dissimilar between spouses than

expected. Collectively, our results do not support a marked tendency for partner choice

according to genetic or phenotypic features, and suggest instead the occurrence of assortative

mating driven by social structure as the cause for ancestry-based assortments among ni-

Vanuatu.

Discussion

By leveraging an extensive genomic dataset of 1,433 contemporary individuals, we show here that all ni-Vanuatu initially descend from admixture between the same ancestral populations: an East Asian-related population, which shares genetic affinities with groups living today in Taiwan and the Philippines, and a Papuan-related population, which shares genetic affinities with groups living today in the Bismarck Archipelago. We also find that admixture was relatively synchronous across islands, in agreement with a peopling history common to all the archipelago. Thus, our results suggest that the exceptionally high cultural diversity of ni-Vanuatu results from a rapid cultural diversification that developed *in situ*, as suggested by linguistic, archaeological and archaeogenetic studies (12, 13, 28). Nevertheless, our results cannot exclude a scenario where the islands were settled by multiple, closely related groups carrying East Asian-related or Papuan-related ancestry. Furthermore, we caution that admixture date estimates from modern DNA data are uncertain and can be biased downward when admixture was gradual (41), which was likely the case in Vanuatu (9). Additional aDNA time transects from multiple islands will be required to provide a definitive picture of the genetic history of ni-Vanuatu.

Our analyses reveal substantial differences in East Asian-related ancestry proportions between islands. We show that these differences do not result solely from Polynesian migrations, as our haplotype-based analyses could differentiate ancestry attributed to the Austronesian expansion from that introduced by Polynesians. A compelling example is Ambae, where East Asian-related ancestry is 1.8-times higher than in surrounding islands but where Polynesian ancestry is low. These findings indicate that the major population turnover following the arrival of Papuan-related peoples was geographically uneven, possibly because, at the time of admixture, the two ancestral populations of ni-Vanuatu were of different sizes across islands, some of which being preferentially settled by Austronesian-related groups and others by Papuan-related groups. Lastly, we find evidence that admixture was sex-biased; either Papuan-related migrants were predominantly males or both males and females migrated, but admixture was more common between Papuan-related males and East Asian-related females.

A recent archaeogenetic study has reported that ancient ni-Vanuatu from Efate and Eretok show genetic similarities with Polynesians (12), supporting the occurrence of migrations from Polynesia, which were previously postulated by linguistic studies (10, 11). We confirm that "Polynesian outlier" communities in Vanuatu are descended from admixture events between Polynesians and local populations. We dated these admixture events to 600-1,000 ya, in line

with archaeological records (42). Furthermore, we extend previous findings by mapping the genetic impact of Polynesian migrations to some Vanuatu islands where Polynesian languages are not spoken today (e.g., Makura, Tongoa, Tongariki and Tanna) (26). These results indicate that genetic interactions between ni-Vanuatu and Polynesian incomers did not systematically trigger shifts to Polynesian languages. Intriguingly, Polynesian ancestry is not detected north of the Kuwae caldera, a large submarine volcano that separates Tongoa and Epi islands. Geological data have shown that the Kuwae volcano erupted in ca. 1452, producing among the largest volumes of magma and aerosol ever recorded (29, 30). We suggest that this caldera, or other active volcanoes in the region, may have acted as a barrier to Polynesian gene flow. Nowadays, ni-Vanuatu living north and south of Kuwae form two genetic groups with distinct socio-cultural practices (e.g., grade-taking and chiefly title political systems) (43), indicating that the area has remained a genetic and cultural frontier. Besides these two groups, we observed subtle genetic differences among ni-Vanuatu, which recapitulate geography and known language subgroups. Yet, migration between islands is common, probably owing to long-term networks of exchange. An illustrative example is Pentecost island, where northern and southern groups show fewer genetic similarities than either do with groups from nearby islands.

By building upon the well-defined genetic history of Vanuatu, we also explored how genetic diversity has been shaped by cultural practices. While levels of genetic relatedness are high among ni-Vanuatu, we do not find evidence for generalized endogamy, which challenges the frequent association geneticists make between the two processes (16, 44-46). Nonetheless, even if ni-Vanuatu spouses are generally less related than non-spouses, we show that their genetic ancestries are more similar than expected, indicating that mating in Vanuatu is not random. Importantly, ancestry similarity between partners is not stronger at trait-associated SNPs, suggesting that ancestry-associated assortments are due to social structure, which may, in turn, be correlated with levels of East Asian-related and/or Polynesian ancestry. Other studies have suggested non-random mating according to ancestry in regions of the world where socio-cultural structure is highly correlated with ancestry (15, 47). Our findings extend the occurrence of such socio-cultural assortments to Oceanians, raising questions of how common this phenomenon is in human societies and whether non-random mating should systematically be accounted for in human genetic studies. Collectively, our study emphasizes the need to include diverse populations in genetic studies, not only to address key anthropological and evolutionary questions that are important for specific geographic regions, but also to identify factors shaping the genetic diversity of human populations as a whole.

Material and Methods

Sample collection and approvals

The sampling survey was conducted in the Republic of Vanuatu between April 2003 and August 2005. The purpose of the study was the estimation of the seroprevalence of HTLV-1 viral infection and the assessment of human genetic diversity in ni-Vanuatu. The recruitment of participants was carried out after the agreement of the Ministry of Health of Vanuatu, the head of each Directorate from the sampled province, and the chief of the sampled village. The data collectors, Olivier Cassar, Helene Walter, Woreka Mera and Antoine Gessain, were accompanied by the village chief and/or the head of the local dispensary. The nature and the scope of the study were explained in detail by Olivier Cassar in English, and by Helene Walter in Bislama (i.e., an English-based Creole that is the main *lingua franca* of Vanuatu), during information meetings organized in each village. Participants could ask any questions after the information meeting. After several hours of reflection, each volunteer participant of at least 18 years of age was asked to sign a written informed consent form, including consent for research on human genetic diversity. Sex, age and birth place, as well as the date and place of blood collection, were collected through a structured questionnaire. Couples were identified through interviews in English or Bislama, and were preferentially sampled. Blood samples were collected either at the local dispensary, a gymnasium or a hut provided by the village chiefs.

The study received approval from the Institutional Review Board of Institut Pasteur (n°2016-02/IRB/5) and the Ministry of Health of the Republic of Vanuatu. It was conducted in full respect of the legal and ethical requirements and guidelines for good clinical practice, in accordance with national and international rules. Namely, research was conducted in accordance with: (i) ethical principles set forth in the Declaration of Helsinki (Version: Fortaleza October 2013), (ii) European directives 2001/20/CE and 2005/28/CE, (iii) principles promulgated in the UNESCO International Declaration on Human Genetic Data, (iv) principles promulgated in the Universal Declaration on the Human Genome and Human Rights, (v) the principle of respect for human dignity and the principles of non-exploitation, non-discrimination and non-instrumentalisation, (vi) the principle of individual autonomy, (vii) the principle of justice, namely with regard to the improvement and protection of health and (viii) the principle of proportionality. The rights and welfare of the subjects have been respected, and the hazards did not outweigh the benefits of the study.

SNP genotyping and quality filters

Five millilitres of blood were obtained by venepuncture from each volunteer participant and

transferred to the Institut Pasteur of New Caledonia, where plasma and buffy coats were

isolated, frozen, and stored at –80°C. Samples were then transferred to the Institut Pasteur in

Paris (France). Out of the 4,428 collected samples, 1,433 samples were selected for genetic

analyses. These samples were selected in order to (i) cover the largest number of islands and

villages, (ii) cover locations where "Polynesian outliers" live nowadays and (iii) include as

many couples as possible. After sample selection, a total of 179 different villages were

covered, located on 29 islands (Table S1). DNA was purified from frozen buffy coats at the

Institut Pasteur of Paris (France), using QIAamp DNA Blood Mini Kit protocol, and eluted in

AE buffer. DNA concentration was quantified with the Invitrogen Qubit 3 Fluorometer using

the Qubit dsDNA broad-range assay. Prior to SNP array genotyping, DNA integrity was

checked on agarose gels.

The 1,433 selected samples were genotyped on the Illumina Infinium Omni 2.5-8 array

(San Diego, California). Genotype calling was performed using the Illumina GenomeStudio

software. We excluded 2,491 SNPs with missing annotations, 9,661 duplicated SNPs, 1,772

SNPs with a GenTrain score < 0.4, SNPs with a missingness > 0.05 and SNPs that deviate

from Hardy-Weinberg equilibrium (i.e., P-value < 0.01 in more than one Vanuatu island).

Only autosomal SNPs were kept for the analyses, unless otherwise stated. After filters, a total

of 2,269,868 SNPs were kept. When the analyses required minor allele frequency (MAF)

filters, a MAF > 0.01 threshold was applied. When a linkage disequilibrium pruning was

required, we pruned the data using a window size of 50 Kb, a step size of 5 SNPs and a r^2

threshold of 0.5. After all these filtering steps were applied, the remaining number of SNPs

was 294,806. All filters were applied using PLINK v.1.9b (48).

Sample quality filters

The highest genotype missingness per sample was 0.018. We removed 21 samples with

outlier values for heterozygosity (mean ± 3 SD), suggestive of DNA contamination, leaving

1,412 samples. Cryptic relatedness between samples was detected using KING v.2.1 (49). We

excluded 456 samples with a kinship coefficient > 0.08 with another sample, whenever

analyses required unrelated samples. For 12 samples, the reported sex did not match the

genetic sex inferred by the Y- and X-chromosome call rates. These include one couple, for

which the reported sex is the opposite to the genetic sex. We thus exchanged the sex of the

two samples, as it is most likely an error on the sample annotation. We did not include the 10

remaining samples, when performing analyses relating to mating practices and sex-specific

migrations. After quality filters, a total of 287 couples was present in the dataset.

Demographic information

Demographic data include the island of birth and the village of residence. Birth place was

missing for 73 samples and the residence for 49 samples. From this information, we retrieved,

for each sample, the geographical coordinates of the birth place (around the center of the

island) and of the place of residence (around the village of residence), using MapAction

reference maps (www.mapaction.org). We considered as the locations of "Polynesian

outliers" all the villages within Futuna, the villages of Mele and Imere in Efate, Ifira island

and the villages of Makatea, Tongamea and Vaitini in Emae (26). Among the 287 couples,

207 reported the same island of birth, while 57 reported a different island. This information

was missing for 23 couples.

Merging with reference datasets

We merged the new SNP array data with whole genome sequences of different populations

across the Pacific (17) and worldwide populations from the SGDP project (50). We also

merged the dataset with SNP array data from Oceanian populations (41), to perform some

population genetics analyses. Datasets were merged using PLINK v.1.9b (48). Transversions

were excluded, to avoid allele strand inconsistences.

Haplotype phasing

We phased the dataset using SHAPEIT v.2 (51, 52) using 500 conditioning states, 10 burn-in

steps, 50 Markov chain Monte Carlo (MCMC) main steps, a window length of 1 cM and an

effective population size of 15,000. We used 1000 Genomes data (53) as a reference panel

and therefore removed all array SNPs that did not align with this data, prior to haplotype

phasing.

Genetic structure: PCA and ADMIXTURE

We performed Principal Component Analyses (PCA) with the 'SmartPCA' algorithm

implemented in EIGENSOFT v. 6.1.4 (54). aDNA samples were projected on PCs by using

the "lsqproject" and "shrinkmode" options. We inferred population structure with

ADMIXTURE v. 1.3.0 (55), using 10 different seeds and assuming that K_{ADM} varies from 2 to

17, and visualized ADMIXTURE results using pong v1.4.7 (56). We estimated pair-wise F_{ST}

values using the StAMPP R package (57), and assessed their significance with 100 bootstraps.

We computed f_4-statistics with ADMIXTOOLS v.6.0 and estimated standard errors by block

jackknife.

Genetic structure: ChromoPainter and fineSTRUCTURE

We ran ChromoPainter (18) to infer haplotype sharing among individuals. This algorithm is

based on a Hidden Markov Model in which each sample is treated as a "recipient", i.e., a

mosaic of haplotypes from a set of "donor" samples. We performed different analyses with

ChromoPainter. In Analysis 1, we inferred the genetic structure of ni-Vanuatu running

ChromoPainter only for the unrelated ni-Vanuatu samples. We set each sample both as a

recipient and as a donor for all the samples (-a mode). We first estimated the global mutation

probability and the switch rate with 10 expectation-maximization (EM) iterations, by using

the '-in -iM' options and by averaging estimates obtained for chromosomes 1, 7, 13 and 19.

The estimated values were 1.542×10^{-4} and 81.415, respectively. Then, we ran ChromoPainter

in the same mode for all chromosomes but fixing the parameters to estimated values. In

Analysis 2, we repeated the same process independently for the reference populations. In this

case, we ran ChromoPainter only for the samples outside Vanuatu (17, 50) and repeated the

two-step process described above. In this case, the global mutation probability was estimated

to 4.53×10^{-4} and the switch rate was 255.42.

We then produced a coancestry matrix by summing the coancestry matrices for all

chromosomes and estimated the C parameter with ChromoCombine. The C factor was

estimated at 0.452 for the Vanuatu dataset (Analysis 1), and 0.498 for the reference

population dataset (Analysis 2). We applied the model-based Bayesian clustering method

fineSTRUCTURE v.2.0.7 (18) on ChromoPainter coancestry matrices. We ran 1 million

burn-in iterations (-y option) and 1 million MCMC iterations (-x option) and sampled values

every 10,000 iterations. We then inferred the fineSTRUCTURE tree (-m T option). We

repeated this process 5 times, with different random seeds.

Following a previous study (19), we estimated for each individual the robustness of the

clustering assignation, by comparing the final state (i.e., the state with the highest posterior

probability) with the 100 MCMC samples. For each individual i, we estimated $x_i^{(m)}$, the

number of individuals clustering with i for each of the $m = 1,…,100$ MCMC samples. We

also estimated $y_{ik}^{(m)}$, the number of individuals that cluster with i in the final state and in each

of the MCMC samples, for each inferred cluster k. Finally, we computed

$P_i = \sum_m [y_{ik}^{(m)} / x_i^{(m)}]$. Therefore, we ended up with a matrix of this sum P_i, for all individuals

and for each genetic cluster, that shows the robustness in the assignation of each individual to

its corresponding cluster in the final state. We repeated this process for $k = 2,…,K$, K being

the maximum number of clusters identified by fineSTRUCTURE. We also repeated the

process for the 5 random seeds and compared the performance of each seed. Fig. S6 shows

the number of ni-Vanuatu individuals with $P_i < 0.9$ for each seed s and for each $k = 2,…K$.

These estimations allowed to test the robustness in the cluster assignation of each individual,

to compare the performance across seeds and to determine the robustness at each k value.

To compare robustness among seeds, we followed a similar approach. We estimated $x_i^{(s)}$,

where $s = 1,…,5$, which corresponds to the number of individuals clustering with individual i

for each seed. We then estimated $y_{ik}^{(s)}$ and $\sum_s [y_{ik}^{(s)} / x_i^{(s)}]$, which shows the concordance in the

assignation of each individual to a cluster across the different seeds. To summarize this

information, we analyzed all the individual sums, for each seed and each cluster k (Fig. S7*A*

and *C*), as well as the number of individuals who were not assigned to the same cluster,

between one seed and all the other seeds (Fig. S7*B* and *D*).

Based on these analyses, we chose the seed that showed the highest consensus among

seeds (Fig. S7) and in which the largest number of samples are robustly assigned (Fig. S6)

(seed number 289 for Analysis 1 and 161 for Analysis 2). Then, we defined the maximum k

value based on the number of individuals who are robustly assigned to a cluster for each k

value and we focused our analyses on this value. Finally, once we chose a seed and a k value,

we removed samples that showed some uncertainty in the cluster assignation ($P_i \leq 0.9$).

Following this process, we observed that the robustness decreases from $K_{FS} = 20$ and we

removed 3 samples from Analysis 1 (Vanuatu). For the reference dataset (Analysis 2), all

samples showed a $P_i > 0.9$ across all k values, therefore we used genetic clusters from the k

value that best suited the hypotheses to test ($K_{FS} = 25$) (Table S5).

For the subsequent analyses, we redefined populations based on fineSTRUCTURE results

and therefore on the genetic data itself. This approach is rather trivial when applied to

continental populations, like the reference populations presented here, because

fineSTRUCTURE clusters coincide broadly with the geographical or cultural categories

assigned to the populations. Yet, this strategy has proven useful when complex admixture

shapes the genetic structure (58, 59), or when the studied populations are sampled from a

small geographic region (19), in which case, defining populations based on geography or

culture may lead to a biased view of the genetic structure of the populations.

Ancestry estimation

We estimated admixture proportions for each ni-Vanuatu individual using SOURCEFIND

(60). We first ran ChromoPainter using the "donors" mode: we considered each ni-Vanuatu

individual as a recipient only and defined donor populations by using the genetic clusters

defined above (Analysis 2). We ran SOURCEFIND by considering all surrogates ("all

surrogates" analysis; Fig. S15 and Table S5) or by limiting as much as possible the number of

surrogates ("limited surrogates" analysis). In the latter case, we considered as surrogates

West_Eurasia1, West_Eurasia2, West_Eurasia3, EastAsia1, EastAsia2, EastAsia3, Atayal,

Paiwan, Southeast_Asia, Cebuano, RenBell, Tikopia, PNG1, PNG2 and PNG_SGDP. Then,

we summed the estimated ancestry proportions for these surrogates and grouped them as

follows: Taiwanese.Philippines (Atayal, Paiwan, Cebuano), EastAsian_mainland (EastAsia1,

EastAsia2, EastAsia3, Southeast_Asia), Polynesian (RenBell, Tikopia), Papuan-related

(PNG1, PNG2 and PNG_SGDP) and European (West_Eurasia1, West_Eurasia2,

West_Eurasia3). The East Asian-related ancestry was estimated as the sum of

Taiwanese.Philippines group and Polynesian group (the EastAsian_mainland was negligible,

with a maximum proportion = 0.007). To facilitate visualization, we excluded an individual

born in Futuna and living in Malekula, who shows a Polynesian ancestry of 0.745 in figures

showing ancestry proportions (Fig. 2, Fig. S8-11). The next individual carrying the highest

Polynesian ancestry shows 0.383. For both the "all surrogates" and "limited surrogates"

analyses, we ran 200,000 MCMC iterations with a burn-in of 50,000 iterations, sampling

every 5,000 iterations and not allowing for self-copy. We estimated the final ancestry of each

ni-Vanuatu sample as the mean of the 30 sampled MCMC runs.

Admixture date estimation

We estimated admixture dates using GLOBETROTTER (61). For this analysis, we ran

ChromoPainter using the "donors" mode, by considering the ni-Vanuatu genetic clusters

defined in (Analysis 1) as recipients and all the reference populations (Analysis 2) as both

recipients and donors. Surrogates were the same as those considered for the "limited

surrogates" SOURCEFIND analysis. For each of the 20 recipient ni-Vanuatu clusters, we

performed 100 bootstraps, which are implemented as resamplings of chromosomes among the

available samples. We set the "Null.ind" option to 1. We assumed a generation time of 28

years, to estimate dates from the number of generations. We also dated admixture in each ni-

Vanuatu sample following the same approach.

Sex-biased admixture

We studied sex-biased admixture by comparing ancestry proportions estimated for the

autosomes and the X chromosome. We estimated ancestry proportions using RFMix v.1.5.4

(33), considering as ancestral populations New Guinean highlanders (approximating Papuan-

related ancestry) and Taiwanese Indigenous peoples and the Cebuano from the Philippines

(approximating East Asian-related ancestry). We ran the "TrioPhased" algorithm, allowing

for phase correction and 3 EM iterations. The window size was set at 0.03 cM (17) and the

admixture date was set at 50 generations. The same parameters were used for the X

chromosome and the autosomes. We combined the X chromosome haploid data of males from

the same island to obtain diploid individuals. We filtered out SNPs with an RFMix posterior

probability < 0.9, those within centromeres and within 2 Mb from the telomeres. We then

estimated East Asian-related and Papuan-related ancestry in the autosomes and the X

chromosome separately. We estimated α_f and α_m, that is, the proportion of female and male

ancestors of ni-Vanuatu who carried Papuan-related ancestry, respectively, by assuming that

admixture proportions have reached equilibrium values (34). In such a case, $\alpha_f = 3\alpha_X -$

$2\alpha_{auto}$ and $\alpha_m = 4\alpha_{auto} - 3\alpha_X$, where α_{auto} and α_X are the average Papuan-related

ancestry proportions estimated on the autosomes and the X chromosome, respectively.

To avoid potential biases due to SNP ascertainment, we also studied sex-biased admixture

by comparing the autosomes and the sex chromosomes of 179 ni-Vanuatu sequenced at 30×

coverage (17). Because genotypes on the X chromosome were not called in the previous

study, we mapped *fastq* files on the X chromosome of the human reference genome (version

hs37d5) and performed genotype calling for X-linked variants, as previously described (17),

setting the ploidy parameter to 1 for males and 2 for females. We kept only biallelic SNPs and

filtered *vcf* files following GATK best practices (62): QualByDepth < 2; FisherStrand value $>$

60; StrandOddsRatio > 3; RMSMappingQuality < 40; MappingQualityRankSumTest < -12.5;

and ReadPosRankSum < -8. We also removed genotypes with a DP < 10 for females and < 3

for males, and those with a GQ < 30. We removed PAR1, PAR2 and XTR regions according

to the annotations provided by the UCSC browser, and amplicon regions as reported in (63).

We filtered out SNPs that were not in Hardy-Weinberg equilibrium in females (*P*-value $<$

0.0001), those with a missingness > 0.05 and those with a MAF < 0.01. We estimated mtDNA

haplogroups with MToolBox (64) from the *bam* and *fastq* files, and Y chromosome

haplogroups were estimated with Yleaf (65).

Inferring migrations within Vanuatu

We used the genetic structure inferred by fineSTRUCTURE to study migration patterns among Vanuatu islands. First, we assigned each genetic cluster to one or more islands, each time more than 25% of the sampled individuals inhabiting the island were assigned to that cluster. We then identified "outlier" individuals who inhabit an island but are assigned to a genetic cluster that is prevalent in another island. These individuals, either themselves or their close ancestors, likely migrated from the island/s where their genetic cluster is predominant to their current place of residence. We removed from these analyses Vanuatu_8 and Vanuatu_9 clusters ($K_{FS} = 20$) because those clusters are driven by recent European gene flow, as well as Vanuatu_15 because no island could be assigned to this cluster when using the 25% rule. To study sex-specific migrations, we calculated the proportion of female (male) migrants by dividing the number of "outlier" females (males) by the number of females (males) in each cluster. We did not include clusters where the number of males or females was < 2.

Tests for exogamy

We tested if spouses show lower genetic relatedness than non-spouses by testing if the average kinship coefficient between spouses is higher or lower than expected by chance, using either permutations or a logistic regression model. When using permutations, we accounted for isolation by distance by sampling random pairs of males and females among individuals born in the same island (Fig. S24*A*) or living in the same village (Fig. S24*B*). Because marriages between first-degree related individuals are very unlikely (they are indeed not observed in the data), we excluded from this analysis individuals who are first-degree relatives, but also tested how the inclusion of these individuals affect the results (Fig. S24*C*). We performed 10,000 permutations for each island. In each permutation, we sampled equally many random pairs as there were observed pairs in the island. The null distribution was built by estimating the average kinship coefficient among randomly sampled pairs, using KING v.2.1. We calculated *P*-values by comparing the average kinship coefficient among the observed couples to the null distribution.

Based on the sample scheme described above, we designed a logistic regression model that could be generalized to multiple explanatory variables, such as ancestry. Let the variable i index all possible pairs of males and females in the dataset, except pairs of individuals who are first-degree relatives, and define the dependent binary variable Y_i with $Y_i = 1$ if i indexes an observed pair of spouses and $Y_i = 0$ otherwise. Define the probability that a pair is an

684   observed couple $p_i(x) = P(Y_i = 1)$, and introduce ϕ_i as the kinship coefficient between

685   individuals in the i:th pair. We estimated the effect of kinship on mate choice by the

686   parameter β^ϕ in the logistic regression model

$$\log\left[\frac{p_i(x)}{1 - p_i(x)}\right] = \mu + \phi_i\beta^\phi + I(village_i)\beta^v + I(island_i)\beta^I, \qquad \textit{Eq. 1}$$

where $I(village_i) = 1$ if individuals of pair i are from the same village and $I(village_i) = 0$

otherwise, and $I(island_i) = 1$ if individuals of pair i are from the same island and

$I(island_i) = 0$ otherwise. As a sensitivity analysis, we also considered a model in which

pairs of first-degree related individuals were included (Fig. S25).

Tests for ancestry-based assortative mating

We extended the logistic regression model shown in *Eq. 1* to test for ancestry-based

assortative mating, by testing if the genetic ancestry of spouses is more similar than that of

non-spouses, accounting for population structure and relatedness avoidance. Let q_{i1}^{EA} and q_{i2}^{EA}

be the proportion of East Asian-related ancestry for the individuals 1 and 2 of pair i and

introduce the variable $q_i^{EA} = \left|q_{i1}^{EA} - q_{i2}^{EA}\right|$. We estimated the effect of having similar

proportions of East Asian-related ancestry between two individuals on the probability that a

pair is an observed couple $p_i(x)$ by the parameter β^q in the logistic regression model,

$$\log\left[\frac{p_i(x)}{1 - p_i(x)}\right] = \mu + q_i^{EA}\beta^q + \phi_i\beta^\phi + I(village_i)\beta^v + I(island_i)\beta^I, \qquad \textit{Eq. 2}$$

where we use the same notations as in *Eq. 1*. A negative (positive) effect size β^q is interpreted

as evidence for assortative (disassortative) mating according to ancestry. Effect sizes for other

ancestries were calculated similarly. The same model was also tested in ni-Vanuatu

originating or not from islands where Polynesian languages are spoken. Note that in standard

regression analyses, such as those used in GWAS, population stratification is usually

corrected by principal components of the genetic relatedness matrix; here we have taken a

more general approach by decomposing the genetic structure in two variables: the kinship and

the ancestry, and have studied how both variables affect mate choice independently.

Tests for SNP-based assortative mating

We extended the logistic regression model shown in *Eq. 1* to test for SNP-based assortative

mating, by testing if the genotypes at a given SNP are more similar between spouses than

non-spouses, accounting for population structure, ancestry-based associated mating and

relatedness avoidance. Define, for each SNP s and each pair i of individuals, the allele

sharing distance (ASD) $d_{i,s}$ as $d_{i,s} = 0$ if both alleles are identical between individuals,

$d_{i,s} = 1$ if only one allele is identical and $d_{i,s} = 2$ if none of the alleles are identical. We

estimated the association between SNP s and mate choice by the parameter β_s^{ASD} in the model

$$\log\left[\frac{p_i(x)}{1 - p_i(x)}\right] = \mu + d_{i,s}\,\beta_s^{ASD} + q_i^{EA}\beta^q + \phi_i\beta^\phi + I(village_i)\beta^v + I(island_i)\beta^I, \qquad Eq.\ 3$$

721   where we use the same notations as in *Eq. 2*. A negative (positive) effect size β_s^{ASD} indicates

722   higher (lower) similarity at SNP s between the members of the observed couples than

723   between non-couples, in line with SNP-based assortative (disassortative) mating.

Tests for trait-based assortative mating

We tested for trait-based assortative mating by testing if genotypes of spouses are more

similar or dissimilar at trait-associated SNPs, relative to non-associated SNPs. We obtained

GWAS summary statistics for 8 candidate traits from the UK Biobank database

(http://www.nealelab.is/uk-biobank). Candidate traits include traits relating to morphology

and physical appearance. Let $y_s = \beta_s^{ASD}$, where β_s^{ASD} is the effect size of SNP s on mate

choice estimated by *Eq. 1*. We estimated if trait-associated SNPs are more similar or

dissimilar between spouses by the parameter β^{trait} in the model

$$y_s = \mu + I(associated_s)\beta^{trait} + MAF_s\beta^{MAF} + GERP_s\beta^{GERP} + rec_s\beta^{rec}, \qquad Eq.\ 4$$

where $I(associated_s) = 1$ if SNP s is significantly associated with the candidate trait in

GWAS (with GWAS *P*-value $< 5\times10^{-8}$) and $I(associated_s) = 0$ otherwise, MAF_s is the

minor allele frequency of SNP s in ni-Vanuatu, $GERP_s$ is the Genomic Evolutionary Rate

Profiling (GERP) score of SNP s and rec_s is the interpolated recombination rate between

SNP s and SNP $s - 1$, estimated in cM/Mb from the 1000 Genomes Phase 3 combined

recombination map. We considered that there is trait-based assortative (disassortative) mating

if β^{trait} is significantly negative (positive). We adjusted *P*-values with the Bonferroni

correction for multiple testing, to account for the number of traits tested.

Acknowledgments

We thank all volunteers in Vanuatu who kindly provided samples and participated in this research. We thank the Ministry of Health of the Republic of Vanuatu (especially Myriam Abel, Rose Bahor, Yvannah Taga, Maturine Tary and the former Minister Jerôme Ludvaune), the Vanuatu Family Health Association (especially Blandine Boulekone and Marie Nickllum Wala), the WHO Office in Port Vila (especially Corinne Capuano) and the Institut Pasteur of New Caledonia at Nouméa (especially Paul M. V. Martin and Eliane Chungue), for their continuous support and interest in this work. We are particularly thankful to Helene Walter, Woreka Mera, Françoise Charavay and Sonia Treptow for their assistance in collecting samples. We thank S. Créno and the HPC Core Facility of Institut Pasteur (Paris) for the management of computational resources. L.R.A. was funded by a Pasteur-Roux-Cantarini fellowship from the Institut Pasteur. The laboratory of Human Evolutionary Genetics is supported by the Institut Pasteur, the Collège de France, the CNRS, the Fondation Allianz-Institut de France, the French Government's Investissement d'Avenir programme, Laboratoires d'Excellence 'Integrative Biology of Emerging Infectious Diseases' (ANR-10-LABX-62-IBEID) and 'Milieu Intérieur' (ANR-10-LABX-69-01), the Fondation de France (n°00106080), and the Fondation pour la Recherche Médicale (Equipe FRM DEQ20180339214) and the French National Research Agency (ANR-19-CE35-0005).

Authors contributions: L.R.A, E.P. and L.Q.-M. designed research; L.R.A analyzed data, with input from J.B., J.C., J.M.R., M.R. and E.P. C.H. and L.L. performed DNA extractions and genotyping. A.M.S., H.C., A.F., F.V., O.C., and A.G. provided resources, expertise and feedback. L.Q.-M and E.P. supervised the study. L.Q.M secured funding. L.R.A., L.Q.M. and E.P. wrote the paper, with input from all authors.

Competing Interest Statement: The authors declare no competing interests.

References

References

1.   Bellwood P (2004) *First Farmers: The Origins of Agricultural Societies* (Wiley-Blackwell).

2.   Gray RD, Drummond AJ, & Greenhill SJ (2009) Language phylogenies reveal expansion pulses
and pauses in Pacific settlement. *Science* 323(5913):479-483.

3.   Hung H-C & Carson MT (2015) Foragers, fishers and farmers: origins of the Taiwanese Neolithic.
Antiquity 88(342):1115-1131.

4.   Kirch PV (2017) *On the road of the winds: An archeological history of the Pacific islands before*
European contact (University of California Press).

5.   Sheppard PJ (2011) Lapita Colonization across the Near/Remote Oceania Boundary. *Curr*
Anthropol 52(6):799-840.

6.   Skoglund P, *et al.* (2016) Genomic insights into the peopling of the Southwest Pacific. *Nature*
538(7626):510-513.

7.   Valentin F, Detroit F, Spriggs MJ, & Bedford S (2016) Early Lapita skeletons from Vanuatu show
Polynesian craniofacial shape: Implications for Remote Oceanic settlement and Lapita origins.
Proc Natl Acad Sci U S A 113(2):292-297.

8.   Lipson M, *et al.* (2018) Population Turnover in Remote Oceania Shortly after Initial Settlement.
Curr Biol 28(7):1157-1165 e1157.

9.   Posth C, *et al.* (2018) Language continuity despite population replacement in Remote Oceania.
Nat Ecol Evol 2(4):731-740.

10. Kirch PV (1984) The Polynesian outliers: Continuity, change, and replacement. *J Pac Hist*
19:224-238.

11. Feinberg RS (2012) *Polynesian Outliers: The State of the Art* (University of Pittsburgh).

12. Lipson M, *et al.* (2020) Three Phases of Ancient Migration Shaped the Ancestry of Human
Populations in Vanuatu. *Curr Biol* 30(24):4846-4856 e4846.

13. François A, Lacrampe S, Franjieh M, & Schnell S (2015) *The Languages of Vanuatu: Unity and*
Diversity (Canberra: Asia–Pacific Linguistics).

14. Heyer E, Chaix R, Pavard S, & Austerlitz F (2012) Sex-specific demographic behaviours that
shape human genomic variation. *Mol Ecol* 21(3):597-612.

15. Zou JY, *et al.* (2015) Genetic and socioeconomic study of mate choice in Latinos reveals novel
assortment patterns. *Proc Natl Acad Sci U S A* 112(44):13621-13626.

16. Ceballos FC, Joshi PK, Clark DW, Ramsay M, & Wilson JF (2018) Runs of homozygosity:
windows into population history and trait architecture. *Nat Rev Genet* 19(4):220-234.

17. Choin J, *et al.* (2021) Genomic insights into population history and biological adaptation in
Oceania. *Nature* 592(7855):583-589.

18. Lawson DJ, Hellenthal G, Myers S, & Falush D (2012) Inference of population structure using
dense haplotype data. *PLoS Genet* 8(1):e1002453.

19. Leslie S*, et al.* (2015) The fine-scale genetic structure of the British population. *Nature*
519(7543):309-314.

20. van Dorp L*, et al.* (2015) Evidence for a Common Origin of Blacksmiths and Cultivators in the
Ethiopian Ari within the Last 4500 Years: Lessons for Clustering-Based Inference. *PLoS Genet*
11(8):e1005397.

21. Tryon DT (1976) *New Hebrides Languages: An Internal Classification* (Australian National
University, Canberra) p 545.

22. François A (2011) Where *R they all? The Geography and History of *R-loss in Southern
Oceanic Languages. *Oceanic Linguistics* 50(1):140-197.

23. Lynch JD, Ross MT, & Crowley T (2011) *The Oceanic Languages* (Routledge).

24. Crowley T (2000) The Language Situation in Vanuatu. *Current Issues in Language Planning* 1(1).

25. Hau'ofa E (1994) Our Sea of Islands. *The Contemporary Pacific* 6(1):148-161.

26. Hermann A & Walworth M (2020) Approche interdisciplinaire des échanges interculturels et de
l'intégration des communautés polynésiennes dans le centre du Vanuatu. *Journal de la société des*
océanistes 151:239-262.

27. Lynch JD & Fakamuria K (1994) Borrowed Moieties, Borrowed Names: Sociolinguistic Contact
between Tanna and Futuna-Aniwa, Vanuatu. *Pacific Studies* 17(1):79-91.

28. Bedford S & Spriggs M (2008) Northern Vanuatu as a Pacific Crossroads: The Archaeology of
Discovery, Interaction, and the Emergence of the "Ethnographic Present". *Asian Perspectives*
47(1):95-120.

29. Monzier M, Robin C, & Eissen J-P (1994) Kuwae (≈ 1425 A.D.): the forgotten caldera. *J Volcanol*
Geoth Res 59(3):207-218.

30. Gao C*, et al.* (2006) The 1452 or 1453 A.D. Kuwae eruption signal derived from multiple ice core
records : Greatest volcanic sulfate event of the past 700 years. *J Geophysical Res* 111(D12107):1-
11.

31. Flexner J, Spriggs M, Bedford S, & Abong M (2016) Beginning Historical Archaeology in
Vanuatu: Recent Projects on the Archaeology of Spanish, French, and Anglophone Colonialism.
Archaeologies of Early Modern Spanish Colonialism, (Springer International Publishing), pp 205-
227.

32. Jolly M (2009) The Sediment of Voyages: Re-membering Quirós, Bougainville and Cook in
Vanuatu. *Oceanic Encounters: Exchange, Desire, Violence*, (ANU Press), pp 57-112.

33. Maples BK, Gravel S, Kenny EE, & Bustamante CD (2013) RFMix: a discriminative modeling
approach for rapid and robust local-ancestry inference. *Am J Hum Genet* 93(2):278-288.

34. Goldberg A & Rosenberg NA (2015) Beyond 2/3 and 1/3: The Complex Signatures of Sex-Biased
Admixture on the X Chromosome. *Genetics* 201(1):263-279.

35. Vienne B (1984) Gens de Motlav. Idéologie et pratique sociale en Mélanésie. *Publication de la*
Société des Océanistes 42:232-240.

36. Bolton L (1999) Women, Place and Practice in Vanuatu: a View from Ambae. *Oceania* 70:43-55.

37. Petrou K & Connell J (2017) Rural-urban migrants, translocal communities and the myth of return

migration in Vanuatu: the case of Paama. *Journal de la société des Océanistes* 144-145(1-2):51-

62.

38. Godelier M (2004) *Métamorphoses de la parenté* (Fayard, Paris).

39. Parra FC*, et al.* (2003) Color and genomic ancestry in Brazilians. *Proc Natl Acad Sci U S A*

100(1):177-182.

40. Alvarez L & Jaffe K (2004) Narcissism guides mate selection: Humans mate assortatively, as

revealed by facial resemblance, following an algorithm of "self seeking like". *Evol Psychol* 2:177–

194.

41. Pugach I*, et al.* (2018) The gateway from Near into Remote Oceania: new insights from genome-

wide data. *Mol Biol Evol* 35(4):871–886.

42. Flexner JL, Bedford S, & Valentin F (2019) Who was Polynesian? Who was Melanesian?

Hybridity and ethnogenesis in the South Vanuatu Outliers. *J Social Archaeol* 19:403–426.

43. Bonnemaison J (1996) *Arts of Vanuatu* (University of Hawaii Press).

44. Kirin M*, et al.* (2010) Genomic runs of homozygosity record population history and

consanguinity. *PLoS One* 5(11):e13996.

45. McQuillan R*, et al.* (2008) Runs of homozygosity in European populations. *Am J Hum Genet*

83(3):359-372.

46. Bianco E*, et al.* (2020) Recent Common Origin, Reduced Population Size, and Marked Admixture

Have Shaped European Roma Genomes. *Mol Biol Evol* 37(11):3175-3187.

47. Ruiz-Linares A*, et al.* (2014) Admixture in Latin America: geographic structure, phenotypic

diversity and self-perception of ancestry based on 7,342 individuals. *PLoS Genet* 10(9):e1004572.

48. Chang CC*, et al.* (2015) Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* 4:7.

49. Manichaikul A*, et al.* (2010) Robust relationship inference in genome-wide association studies.

Bioinformatics 26(22):2867-2873.

50. Mallick S*, et al.* (2016) The Simons Genome Diversity Project: 300 genomes from 142 diverse

populations. *Nature* 538(7624):201-206.

51. Delaneau O, Marchini J, & Zagury JF (2011) A linear complexity phasing method for thousands

of genomes. *Nat Methods* 9(2):179-181.

52. Delaneau O, Zagury JF, & Marchini J (2013) Improved whole-chromosome phasing for disease

and population genetic studies. *Nat Methods* 10(1):5-6.

53. Auton A*, et al.* (2015) A global reference for human genetic variation. *Nature* 526(7571):68-74.

54. Patterson N, Price AL, & Reich D (2006) Population structure and eigenanalysis. *PLoS Genet*

2(12):e190.

55. Alexander DH, Novembre J, & Lange K (2009) Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res* 19(9):1655-1664.
56. Behr AA, Liu KZ, Liu-Fang G, Nakka P, & Ramachandran S (2016) pong: fast analysis and
visualization of latent clusters in population genetic data. *Bioinformatics* 32(18):2817-2823.
57. Pembleton LW, Cogan NO, & Forster JW (2013) StAMPP: an R package for calculation of
genetic differentiation and structure of mixed-ploidy level populations. *Mol Ecol Resour*
13(5):946-952.
58. Arauna LR, *et al.* (2017) Recent Historical Migrations Have Shaped the Gene Pool of Arabs and
Berbers in North Africa. *Mol Biol Evol* 34(2):318-329.
59. Mas-Sandoval A, *et al.* (2019) Reconstructed Lost Native American Populations from Eastern
Brazil Are Shaped by Differential Je/Tupi Ancestry. *Genome Biol Evol* 11(9):2593-2604.
60. Chacon-Duque JC, *et al.* (2018) Latin Americans show wide-spread Converso ancestry and
imprint of local Native ancestry on physical appearance. *Nat Commun* 9(1):5388.
61. Hellenthal G, *et al.* (2014) A genetic atlas of human admixture history. *Science* 343(6172):747-
751.
62. Van der Auwera GA, *et al.* (2013) From FastQ data to high confidence variant calls: the Genome
Analysis Toolkit best practices pipeline. *Curr Protoc Bioinformatics* 43:11 10 11-11 10 33.
63. Lucotte EA, *et al.* (2018) Dynamic Copy Number Evolution of X- and Y-Linked Ampliconic
Genes in Human Populations. *Genetics* 209(3):907-920.
64. Calabrese C, *et al.* (2014) MToolBox: a highly automated pipeline for heteroplasmy annotation
and prioritization analysis of human mitochondrial variants in high-throughput sequencing.
Bioinformatics 30(21):3115-3117.
65. Ralf A, Montiel Gonzalez D, Zhong K, & Kayser M (2018) Yleaf: Software for Human Y-
Chromosomal Haplogroup Inference from Next-Generation Sequencing Data. *Mol Biol Evol*
35(5):1291-1294.
906
907

Figure legends

Fig. 1. Sampling locations and genetic structure in Vanuatu. (*A*) Map showing the location of the 1,439 sampled ni-Vanuatu individuals. The inset at the bottom left shows the location of Vanuatu in the Asia-Pacific region. (*B,C*) Principal Component Analysis (PCA) of genotypes of ni-Vanuatu at 301,774 SNPs, in the context of (*B*) the broad Asia-Pacific region and (*C*) the Vanuatu archipelago only. (*A-C*) Each point indicates an individual, colored according to the latitude of their island of residence.

Fig. 2. Fine-scale genetic structure and admixture in Vanuatu. (*A*) Map showing the clustering of 979 ni-Vanuatu into 20 genetic clusters, according to fineSTRUCTURE ($K_{FS} = 20$). Each point indicates an individual, located according to their village of residence. Colors indicate genetic clusters, so that the closer the colors, the closer the clusters. Noise was added to sampling locations to facilitate visualization. (*B*) Admixture date estimates for each genetic cluster, based on 100 bootstrap replicates. Colors are the same as those used in (*A*). The x-axis shows the admixture date in years before present, assuming a generation time of 28 years. The y-axis indicates the latitude of the islands assigned to each genetic cluster. The heights of the density curves are proportional to the sample size of each cluster. (*C*) Proportion of East Asian-related ancestry estimated by SOURCEFIND. (*D*) Ratio of the Polynesian and East Asian-related ancestry proportions, estimated by SOURCEFIND. The inset at the bottom left shows the location of Polynesian-speaking individuals, indicated by black points. (*C-D*) Each point indicates an individual, colored according to their ancestry proportions.

Fig 3. Sex-biased admixture and migration patterns in Vanuatu. (*A*) Papuan-related ancestry proportions in ni-Vanuatu, estimated for the 22 autosomes and the X chromosome separately by RFMix. (*B*) Frequencies of Y chromosome and mtDNA haplogroups, colored according to their assumed origins (i.e., shades of blue or red indicate Papuan- or East Asian-related origins, respectively). Haplogroups were inferred from high-coverage genome sequencing data obtained for a subset of 179 ni-Vanuatu (17). (*C*) Recent migrations among Vanuatu islands, inferred based on fineSTRUCTURE clusters. The arrows connect the location of the genetic cluster to which individuals were assigned to their actual place of residence. The colors indicate the predominant genetic cluster in the island of origin. The width is proportional to the number of inferred migrant individuals, relative to the number of females or males in the genetic cluster.

Fig 4. Ancestry- and trait-based assortative mating in Vanuatu. (*A*) Effects of kinship and

ancestry differences on partner choice. Effect sizes were estimated with a logistic regression

model, while accounting for population structure (island of birth and village of residence).

The effect size and *P*-value were estimated using different ancestries as predictors,

independently. (*B*) Increased (purple) or decreased (orange) genetic similarity among spouses

(as measured by β_S^{ASD}) at trait-associated SNPs, relative to non-associated SNPs. Results on

the left are based on a logistic model that includes only the genotype dissimilarity among

spouses at each SNP, whereas results on the right are based on the logistic model that also

controls for possible confounders (i.e. geography, kinship and ancestry differences).