The genomic landscape of contemporary western Remote Oceanians

Lara R. Arauna[1,*], Jacob Bergstedt[1,2,3,13], Jeremy Choin[1,4,13], Javier Mendoza-Revilla[1,5,13], Christine Harmant[1], Maguelonne Roux[1,6], Alex Mas-Sandoval[7], Laure Lémée[8], Heidi Colleran[9], Alexandre François[10], Frédérique Valentin[11], Olivier Cassar[12], Antoine Gessain[12], Lluis Quintana-Murci[1,4,14,*], Etienne Patin[1,14,15,*]

[1]Institut Pasteur, Université Paris Cité, CNRS UMR2000, Human Evolutionary Genetics Unit, Paris 75015, France

[2]Institute of Environmental Medicine, Karolinska Institutet, Stockholm 171 77, Sweden

[3]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm 171 77, Sweden

[4]Chair Human Genomics and Evolution, Collège de France, Paris 75005, France

[5]Laboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima, Perú

[6]Bioinformatics and Biostatistics Hub, Institut Pasteur, Université Paris Cité, Paris 75015, France

[7]Department of Life Sciences, Imperial College London, Ascot SL5 7PY, United Kingdom

[8]Biomics Platform, Institut Pasteur, Paris 75015, France

[9]BirthRites Independent Max Planck Research Group, Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Leipzig 04103, Germany

[10]Langues, Textes, Traitements Informatiques, Cognition (LaTTiCe), UMR 8094, CNRS, Paris 75015, France

[11]MSH Mondes, UMR 8068, CNRS, Nanterre 92023, France

[12]Oncogenic Virus Epidemiology and Pathophysiology Unit, Institut Pasteur, UMR 3569, CNRS, Paris 75015, France

[13]These authors contributed equally

[14]Senior authors

[15]Lead contact

[*]Correspondence: lrubioar@pasteur.fr (L.R.A.); quintana@pasteur.fr (L.Q.-M.); epatin@pasteur.fr (E.P.)

SUMMARY

The Vanuatu archipelago served as a gateway to Remote Oceania during one of the most extensive human migrations to uninhabited lands, ~3,000 years ago. Ancient DNA studies suggest an initial settlement by East Asian-related peoples that was quickly followed by the arrival of Papuan-related populations, leading to a major population turnover. Yet, there is uncertainty over the population processes and the sociocultural factors that have shaped the genomic diversity of ni-Vanuatu, who present nowadays among the world's highest linguistic and cultural diversity. Here, we report new genome-wide data for 1,433 contemporary ni-Vanuatu from 29 different islands, including 287 couples. We find that ni-Vanuatu derive their East Asian- and Papuan-related ancestry from the same source populations and descend from relatively synchronous, sex-biased admixture events that occurred ~1,700-2,300 years ago, indicating a peopling history common to all the archipelago. However, East Asian-related ancestry proportions differ markedly across islands, suggesting that the Papuan-related population turnover was geographically uneven. Furthermore, we detect Polynesian ancestry arriving ~600-1,000 years ago to South Vanuatu in both Polynesian- and non-Polynesian-speaking populations. Lastly, we provide evidence for a tendency of spouses to carry similar genetic ancestry, when accounting for relatedness avoidance. The signal is not driven by strong genetic effects of specific loci or trait-associated variants, suggesting that it results instead from social assortative mating. Altogether, our findings provide insight into both the genetic history of ni-Vanuatu populations and how sociocultural processes have shaped the diversity of their genomes.

Keywords: Pacific, populations, migrations, admixture, genetics, cultural practices

INTRODUCTION

Vanuatu, an archipelago located in western Remote Oceania, is a key region for understanding the peopling history of the Pacific. The cultural, anthropological and genetic diversity of Vanuatu reflects three distinct phases of population movements. The first, which started in present-day Taiwan ~5,000 ya, was associated with the spread of Austronesian languages to Near and Remote Oceania [1-3]. The so-called Austronesian expansion led to the emergence of the well-characterized Lapita cultural complex, which emerged in the Bismarck Archipelago and reached Vanuatu ~3,000 ya [4-6]. Morphometric and ancient DNA (aDNA) studies indicate that the Lapita people in Vanuatu carried East Asian-related ancestry, supporting a connection with the Austronesian expansion [7, 8]. The second migration occurred after the Lapita period, ~2,500 ya, and involved the arrival of Papuan-related peoples who shared ancestry with contemporary Bismarck Archipelago islanders. aDNA studies have shown that these migrations triggered a dramatic shift in genetic ancestry, from the East Asian-related ancestry observed in first Remote Oceanians to the Papuan-related ancestry that has remained predominant since then [9, 10]. Finally, it has been postulated that "Polynesian outliers" from Vanuatu (i.e., Polynesian speakers living outside the Polynesia triangle) are the descendants of migrations from Polynesia into western Remote Oceania [11, 12], a model that has received recent genetic support [13].

Although aDNA studies have revealed that ni-Vanuatu are descended from at least three ancestral populations [9, 10, 13], whether the settlement process was uniform across the multiple islands of the archipelago remains an open question. aDNA data indicate that individuals dated to ~2,500 ya from Central and South Vanuatu carried largely different proportions of East Asian-related ancestry [9, 10, 13], in line with either a unique population turnover that was geographically heterogeneous, or separate admixture events between different populations across islands. Furthermore, Vanuatu is the country with the world's largest number of languages per capita [14], which supports the view that languages have rapidly diversified since the initial settlement and/or that the arrival of new groups to the archipelago further increased linguistic diversity. Nevertheless, these questions have been difficult to resolve because, to date, available ancient and modern DNA data from the region have remained sparse [9, 10].

Here, we generated genome-wide genotype data for 1,433 contemporary ni-Vanuatu and assessed their fine-scale genetic structure, in order to address central questions about the settlement of western Remote Oceania: do all contemporary ni-Vanuatu derive their ancestry from three populations only? Was the contribution of these three ancestral populations

different across the archipelago? Was the Post-Lapita shift to Papuan-related ancestry heterogeneous across islands? In addition, the recent settlement of Vanuatu allows for the study of how socio-cultural practices have shaped genetic diversity in humans over the past ~3,000 years, in the form of sex-biased admixture, relatedness avoidance and non-random mating [15-17]. We thus used the comprehensive set of population genomics data presented here, which includes 287 male-female couples, to answer important questions relating to the genetic history of a population that descends from diverse ancestral populations: Was admixture sex-biased? Was admixture accompanied by language shifts? Has socio-cultural structure influenced mating? Can residence rules and urbanization affect human genetic structure?

RESULTS

The Genetic Variation of Ni-Vanuatu is Spatially Structured. To shed light on the genetic make-up of ni-Vanuatu, we collected >4,000 blood samples from contemporary individuals between 2003 and 2005, and genotyped a selection of 1,433 of these sampled individuals at 2,358,955 single nucleotide polymorphisms (SNPs). After merging with 179 high-coverage whole genome sequencing data [18] and excluding 173 low-quality, duplicated or related samples, a total of 1,439 ni-Vanuatu samples was included in subsequent analyses, including those from 522 males and 917 females living in 29 islands and 179 different villages (Fig. 1A and Table S1).

   Principal Component (PCA) and ADMIXTURE analyses indicate that contemporary ni-Vanuatu fall on a genetic gradient between East Asian-related and Papuan-related populations (Fig. 1B and Fig. S1), supporting the view that their ancestry derives from these two population groups. When projecting ancient samples from Vanuatu, we found that Lapita individuals show higher affinity with present-day East Asian-related populations, whereas Post-Lapita individuals are closer to Papuan-related populations, in line with a Papuan-related population turnover occurring after the Lapita period [8-10, 13]. Furthermore, contemporary ni-Vanuatu show high genetic similarities with individuals from the Bismarck Archipelago, in line with the hypothesis that their Papuan-related ancestors originated from these islands [8-10, 13, 18]. However, the extensive geographic coverage of the dataset presented here enabled us to reveal substantial genetic substructure among ni-Vanuatu islanders. PCA showed that genetic variation in ancient and contemporary individuals from Vanuatu is explained by two contiguous but distinct groups that broadly reflect geography (Fig. 1C and Fig. S2). When considering the number of ancestral components that is most supported by ADMIXTURE ($K_{ADM} = 9$), populations from different islands also show different ancestral components (Fig. S1). These results reveal that contemporary ni-Vanuatu show substantial genetic differentiation (Fig. S3), which could result either from different admixture histories during the settlement period and/or from the existence of barriers to gene flow that formed after the settlement of the archipelago.

   To gain further insights into the genetic history of ni-Vanuatu, we next assessed the fine-scale genetic structure of the 1,439 sampled individuals, using ChromoPainter and fineSTRUCTURE [19-21]. Haplotype-based clustering revealed a first separation between ni-Vanuatu living north and south of the strait separating Epi and Tongoa islands ($K_{FS} = 2$; Fig. S4). At $K_{FS} = 4$, populations separate into clusters, here referred to as the *Banks and Torres Islands* cluster and the *North East*, *North West* and *Central-South Vanuatu* clusters (Fig. 2A

and Fig. S4 and Table S2), that are in general agreement with the classification of Oceanic

languages spoken in the archipelago [22-24]. Of note, individuals from the north and south of

Pentecost island cluster with individuals from different neighboring islands, indicating that

the sea does not necessarily act as a barrier to gene flow in the region, in line with linguistic

and ethnographic data that reveal cultural networks between islands [22, 23, 25]. Likewise, in

the Ambae island, northwest and south inhabitants cluster separately (Fig. 2A), suggesting

that the periodic volcanic activity of the Ambae caldera has affected gene flow in the region.

At $K_{FS} = 20$, i.e., the highest K_{FS} value for which statistical robustness remains maximal

(Methods) [20], we found that genetic clusters are often island-specific (Fig. 2A, Figs. S4-S7

and Table S1). These observations suggest that clusters inferred by fineSTRUCTURE are

reliable, as they reflect expected geographic and linguistic barriers to gene flow.

The Post-Lapita Shift to Papuan-related Ancestry was Geographically Uneven. We

detected important differences among fineSTRUCTURE clusters in the proportions of East

Asian-related ancestry, that is, ancestry related to contemporary populations from either

Taiwan and the Philippines or Polynesia (range of SOURCEFIND estimates: 0.15 to 0.55;

Fig. 2C, Figs. S8-S11 and Table S3). East Asian-related ancestry was found to be lowest for

the *North Vanuatu* clusters (e.g., Malekula, Ambrym, Epi; median = 0.230, SD = 0.056) and

highest in the *Central-South Vanuatu* cluster (i.e., Mele and Imere in Efate island, Makatea,

Tongamea and Vaitini in Emae island, and Tongoa and Ifira islands; median = 0.323, SD =

0.077), where "Polynesian outlier" communities live today [11, 12]. Nevertheless, East Asian-

related ancestry is also high in islands where Polynesian ancestry is low, such as Ambae

(median = 0.405, SD = 0.042; Fig. 2D), suggesting that differences in East Asian-related

ancestry are not due solely to differences in Polynesian ancestry (Figs. S10 and S11 and Table

S3).

To assess whether differences in East Asian-related ancestry originate from a

geographically uneven ancestry turnover or separate admixture events between distinct

populations, we dated admixture in each genetic cluster separately. Besides more recent

events relating to Polynesian migrations (see next section), all estimates overlap the same

time period that ranges from 1,700 to 2,300 ya (Fig. 2B, Fig. S12 and Table S4), suggesting

that all ni-Vanuatu share the same admixture history. To test this hypothesis more formally,

we evaluated if the Papuan-related ancestry carried by present-day ni-Vanuatu derives from a

single source. PCA and f_4-statistics of the form f_4(X, New Guinean highlanders; Solomon or

Bismarck Archipelago islanders, East Asians) indicate that all ni-Vanuatu show similar

genetic relatedness with populations from the Bismarck Archipelago, New Guinea or the Solomon islands (Figs. S13 and S14), in agreement with aDNA data [13]. Furthermore, the haplotype-based SOURCEFIND method detected that the same cluster of Bismarck Archipelago islanders is the source that contributed the most to all ni-Vanuatu (Fig. S15 and Table S5). Of note, SOURCEFIND also detected a small contribution from New Guinean highlanders and Santa Cruz islanders in all ni-Vanuatu clusters, which negatively correlates with Polynesian ancestry ($r = -0.378$, P-value $< 2.22 \times 10^{-16}$), probably because Polynesian source populations capture some Papuan-related ancestry. Collectively, these findings indicate that ni-Vanuatu are descended from relatively synchronous admixture events between the same sources of Papuan- and East Asian-related ancestry; yet, their proportions differ markedly across islands, suggesting that the dramatic Papuan-related ancestry shift that started ~2,500 ya was geographically uneven.

Polynesian Migrations Did Not Necessarily Trigger Language Shifts. Admixture proportions and date estimates indicate that ni-Vanuatu ancestry partly derives from a third and more recent migration originating from Polynesia (Fig. 2B-D and Tables S3 and S4). We dated admixture events between 600 and 1,000 ya for genetic clusters predominant in Mele and Imere (Efate island), Makatea, Tongamea and Vaitini (Emae island), as well as in Ifiraand Futuna islands (Fig. 2B, Fig. S12 and Table S4), where Polynesian languages are spoken today [12, 26]. These clusters show higher Polynesian ancestry and, therefore, higher East Asian-related ancestry proportions, relative to *Banks and Torres Islands* and *North Vanuatu* clusters (SOURCEFIND estimates; Fig. 2C-D, Figs. S8-S11 and Table S3). Ni-Vanuatu Polynesian speakers show a higher ratio of Polynesian-to-East Asian ancestry, when compared to non-Polynesian speakers (ratio = 0.647 *vs.* 0.479; Wilcoxon test *P*-value $< 2.22 \times 10^{-16}$; Fig. 2D and Fig. S10). Furthermore, f_4-statistics of the form f_4(X, East Asians; Tongans, New Guinean highlanders) suggest that ni-Vanuatu from Efate, Ifira and Emae share more alleles with Polynesians than other ni-Vanuatu do (Fig. S16).

Interestingly, our analyses also revealed that Polynesian ancestry is not restricted to Vanuatu islands where Polynesian languages are spoken. Non-Polynesian-speaking populations assigned to the *Central-South Vanuatu* (e.g., Tongoa, Tongariki and Tanna) and the *Banks and Torres islands* clusters also show a higher Polynesian-to-East Asian ancestry ratio, relative to *North East* and *North West* clusters (ratio = 0.557 and 0.510, *vs.* 0.461; Wilcoxon test *P*-value $< 2.22 \times 10^{-16}$; Fig. 2D, Figs. S10 and S11 and Table S3). Furthermore, estimated admixture dates are similar among groups from the *Central-South Vanuatu* cluster

that speak or not Polynesian languages (Fig. 2B, Fig. S12 and Table S4). These results support the view that Polynesian migrations, as well as subsequent contacts between Vanuatu islands [12, 27] or with "Polynesian outliers" from the Solomon islands [28], also introduced Polynesian ancestry among non-Polynesian-speaking groups.

Conversely, we found no evidence of admixture with Polynesians in individuals assigned to *North Vanuatu* clusters, including Epi islanders, despite the geographic proximity between Epi and Tongoa (Fig. 2D, Figs. S10 and S11 and Table S3). Of note, ADMIXTURE, PCA and fineSTRUCTURE analyses separate ni-Vanuatu into northern and southern populations, the frontier between the two being located between Epi and Tongoa (Figs. 1C and 2A and Fig. S1). The strait that separates the two islands today is the location of the Kuwae caldera [29, 30], whose volcanic activity may have been a barrier to gene flow and/or may have triggered large-scale population movements that disrupted the isolation by distance patterns. Together, these results reveal that, since 1,000 ya onward, Polynesians migrated to Vanuatu where they admixed with local populations, and that such interactions did not necessarily result in a shift to Polynesian languages.

A Limited Genetic Impact of European Colonization. Our genetic data indicate that admixture between ni-Vanuatu and Europeans has been rare or has left few descendants in Vanuatu. In total, only 28 individuals out of 1,439 (1.95%) show a genetic contribution from Europeans higher than 1%, according to SOURCEFIND analyses (range: 0.03% to 35.3%; Fig. S8 and Table S3). The two fineSTRUCTURE clusters with the highest European ancestry (median = 0.093 and 0.107, *vs*. 0.002 for other clusters) show an admixture event in the last 120 years (mean = 78 ya, 95%CI: [74–81] and mean = 111 ya, 95%CI: [104-118]) (Fig. 2B and Table S4). Similarly, the three other clusters that include individuals carrying more than 1% of European ancestry show a pulse of admixture occurring in the last 200 years. These results are consistent with historical records; early, fleeting European contacts with ni-Vanuatu began in 1606, with subsequent documented exploratory voyages in 1768, 1774 and 1809; yet, it was only from around 1829 onwards that contacts became more common, when Christian missionaries and European colonists settled in the archipelago and the first intermarriages were reported [31, 32].

Genetic Admixture in Vanuatu was Sex-Biased. Genetic studies have suggested that Papuan-related migrations from the Bismarck Archipelago into Remote Oceania were male-biased, because contemporary Polynesians and ancient individuals from present-day Vanuatu

show lower Papuan-related ancestry on the X chromosome, relative to autosomes [8, 13]. To

confirm that admixture between the ancestors of present-day ni-Vanuatu was sex-biased, we

estimated Papuan- and East Asian-related ancestry in ni-Vanuatu on each chromosome

separately, using local ancestry inference [33]. We found that Papuan-related ancestry is

indeed significantly lower on the X chromosome, relative to autosomes (α_X = 75.2% *vs.* α_{auto}

= 79.8%, Wilcoxon test *P*-value < 1.36×10^{-5}; Fig. 3A), in agreement with aDNA results [13].

These values were similar between Polynesian and non-Polynesian speakers (Fig. S17A,

Wilcoxon test *P*-value = 0.13), which indicates that it is not explained by recent migrations

from Polynesia. We replicated the results using high-coverage genome sequences from a

subset of 179 ni-Vanuatu [18], implying that our results are not biased due to SNP

ascertainment (Fig. S17B). Assuming that admixture proportions have reached equilibrium

values [34], we estimated that the genetic contribution of Papuan-related males to ni-Vanuatu

was 27% higher than that of Papuan-related females (α_m = 93.5% *vs.* α_f = 66.1%).

Accordingly, Y chromosomes of ni-Vanuatu are dominated by haplogroups found at high

frequency in Near Oceanians (e.g., M1a3b2, S1), whereas mitochondrial DNAs (mtDNA)

show a high proportion of haplogroups typically found in East Asia (e.g., B4a1a1, E1a2a4)

(Fig. 3B and Table S1). Collectively, these results support the notion that ni-Vanuatu ancestry

predominantly results from admixture between Papuan-related males and East Asian-related

females.

Recent Migrations are Influenced by Residence Rules and Urbanization. The genetic

structure of contemporary ni-Vanuatu is also expected to reflect socio-cultural practices (e.g.,

social networks, exchange and marriage rules) that have culturally evolved since the

settlement of the archipelago. We leveraged the high-resolution genetic data to infer recent

migrations between Vanuatu islands – indicated by individuals who inhabit an island but

belong to a genetic cluster that is prevalent in another island – and determine if these

migrations have involved mainly females or males, in line with virilocal or uxorilocal

postmarital residence rules, respectively. We found that 5.70% of the sampled individuals (54

individuals) migrated at a large geographical scale (K_{FS} = 4), while 11.81% (112 individuals)

migrated at a local scale (K_{FS} = 20; Table S6), suggesting more genetic connections between

closer islands. We estimated that local mobility among females is higher than among males

(K_{FS} = 20; odds ratio = 2.20, Fisher's exact test *P*-value = 9.16×10^{-4}; Fig. 3C), as expected

under virilocal residence and/or female exogamy [35]. The same trend was observed at a

larger geographical scale, when considering migrations between four broader regions (K_{FS} =

4; odds ratio = 1.86, *P*-value = 0.075) and when restricting the analyses to the reported birth

place of male-female couples included in the dataset (Fig. S18) (odds ratio = 1.96, *P*-value =

6.89×10^{-3}). Notably, comparisons of the places of birth and residence of sampled individuals

did not support female-biased migrations (odds ratio = 0.807, *P*-value = 0.18; Fig. S19 and

Table S6), possibly reflecting a bias in the self-reported birth places. This might occur if

women are inclined to report their husband's or children's place rather than their father or

mother's place [36] or under other, more complex patterns.

We also explored the direction of the inferred migrations, and found that both female and

male migrations mainly occurred from the northernmost and southernmost islands towards the

center of the archipelago (Fig. 3C), where Port Vila – the largest city – has developed [37].

Migrations are also common among North Vanuatu islands, consistent with a long-term

network of cultural and material exchanges in the region [25, 28]. Thus, our genomic data

reflect the migration patterns that characterize the recent history of ni-Vanuatu, including

residence rules and urbanization.

No Evidence for Endogamous Practices among Ni-Vanuatu Spouses. Human kinship

systems vary tremendously, regulating marriage, exchange, endogamy or exogamy according

to relational concepts [38]. Given the small census size of ni-Vanuatu populations, it is

interesting to consider whether and how sociocultural features like marriage and exchange

rules influence local levels of genetic relatedness. We found that genetic relatedness was

higher within islands than between islands (Wilcoxon test *P*-value < 2.22×10^{-16}; Figs. S20-

S22 and Table S7), consistent with an isolation by distance model (Mantel test *P*-value =

0.001; 1,000 permutations). Accordingly, out of the 287 male-female couples included in the

dataset, 78.4% were born on the same island (Fig. S23). Genetic relatedness is also higher

among inhabitants of the same village, relative to individuals living on the same island

(Wilcoxon test *P*-value < 2.22×10^{-16}) (Fig. S20 and Table S7), suggesting that the local

community is often the source of marriage partners. However, despite the generally high

levels of genetic relatedness observed between ni-Vanuatu, we did not observe an excess of

genetic relatedness among couples (Figs. 4A, S23 and S24). Specifically, spouses tend to

show slightly lower kinship coefficients than random pairs of individuals from the same

village, when excluding first-degree related individuals from all possible pairs (logistic

regression model *P*-value = 0.057; resampling *P*-value = 0.079) (Figs. S24 and S25) and

when adjusting on differences in ancestry (*P*-value < 0.05; Figs. 4A). Assuming that our total

sample is not biased towards related individuals, these results suggest that endogamy is not a

common practice among contemporary ni-Vanuatu and more generally illustrate that

populations can show high levels of genetic relatedness in the absence of endogamous

marriage practices.

Ni-Vanuatu Spouses Tend to Share Similar Genetic Ancestry. Research on admixed

populations from other parts of the world has shown that, in addition to low genetic

relatedness, partners tend to show similar genetic ancestry [16], because mating often occurs

within socio-cultural groups, which can correlate, in turn, with genetic ancestry. To verify

whether this phenomenon is observed in Vanuatu, we implemented a logistic regression

model that jointly estimates the effects of geography, genetic relatedness and genetic ancestry

on the probability to be partners (Methods). These analyses confirmed that spouses tend to

originate from the same island, while showing lower kinship coefficients than non-spouses

(Fig. 4A). Importantly, we found that spouses show lower differences in genetic ancestry than

non-spouses, when considering Papuan-related (β = -4.233, P-value = 0.018), East Asian-

related (β = -4.221 P-value = 0.018) or Polynesian (β = -5.197, P-value = 0.024) ancestry.

These results suggest that ni-Vanuatu tend to mate with a partner who carries similar genetic

ancestry.

Two hypotheses have been proposed to explain these observations. First, social assortative

mating may underlie the observed signal, as genetic ancestry can correlate with socio-cultural

structure. Second, spouses may choose their partner because they share biological traits, such

as physical appearance [39, 40]. To test these hypotheses, we searched for genomic loci that

could play a role in partner choice, by including, in the logistic regression model, a term that

measures dissimilarity between individuals at each SNP (Methods). No SNP showed

statistical evidence for a significantly lower or higher genotype similarity between spouses,

when accounting for multiple testing (Fig. S26). To test for assortative mating according to

polygenic traits, we then evaluated if genotype similarity between spouses is significantly

higher or lower at SNPs associated with candidate traits relating to physical appearance, when

compared to non-associated SNPs. When we did not account for genetic structure and

ancestry-associated assortative mating (i.e., effect sizes are estimated from a model where

these confounders are not included, see Methods), we found evidence for assortative mating

according to body mass index (BMI, Fig. 4B). However, when accounting for such possible

confounders, we found no statistical evidence that genotypes at trait-associated variants are

more similar or dissimilar between spouses than expected. Collectively, our results do not

support a marked tendency for partner choice according to genetic or phenotypic features, and

342   suggest instead the occurrence of assortative mating driven by social structure as the cause for

343   ancestry-based assortments among ni-Vanuatu.

DISCUSSION

By leveraging an extensive genomic dataset of 1,439 contemporary individuals, we show here that ni-Vanuatu initially descend from admixture between the same ancestral populations: an East Asian-related population, which shares genetic affinities with groups living today in Taiwan and the Philippines, and a Papuan-related population, which shares genetic affinities with groups living today in the Bismarck Archipelago. We also estimate that admixture occurred after the Lapita period, ~1,700-2,300 years ago, and was relatively synchronous across islands, in agreement with a peopling history common to all the archipelago. Thus, our results suggest that the high cultural diversity of ni-Vanuatu results from a rapid cultural diversification that developed *in situ*, as suggested by linguistic, archaeological and archaeogenetic studies [13, 14, 28]. Nevertheless, our analyses cannot definitely rule out that, after the Lapita period, Vanuatu islands were settled by different, already admixed groups carrying varying levels of East Asian-related and Papuan-related ancestry, as recently suggested [18]. Furthermore, we caution that admixture date estimates from modern DNA data are uncertain and can be biased downward when admixture was gradual [41], which was likely the case in Vanuatu [10]. Additional aDNA time transects from multiple islands will be required to provide a definitive picture of the admixture history of ni-Vanuatu.

Our analyses reveal substantial differences in East Asian-related ancestry proportions between islands. We show that these differences do not result solely from Polynesian migrations, as our haplotype-based analyses could differentiate ancestry attributed to the Austronesian expansion from that introduced by Polynesians. A compelling example is Ambae, where East Asian-related ancestry is 1.8-times higher than in surrounding islands but where Polynesian ancestry is low. Assuming a simple admixture model, these findings imply that the major population turnover following the arrival of Papuan-related peoples was geographically uneven, possibly because, at the time of admixture, the two ancestral populations of ni-Vanuatu were of different sizes across islands, some of which being preferentially settled by East Asian-related groups and others by Papuan-related groups. Lastly, we confirm that admixture was sex-biased [8, 13]; either Papuan-related migrants were predominantly males or both males and females migrated, but admixture was more common between Papuan-related males and East Asian-related females.

A recent archaeogenetic study has reported that ancient ni-Vanuatu from the Chief Roi Mata's Domain in Efate show genetic similarities with Polynesians [13], supporting the occurrence of migrations from Polynesia, which were previously postulated by linguistic studies [11, 12]. We confirm that "Polynesian outlier" communities in Vanuatu are descended

from admixture events between Polynesians and local populations. We dated these admixture

events to 600-1,000 ya, in line with archaeological records [11]. Furthermore, we extend

previous findings by mapping the genetic impact of Polynesian migrations to some Vanuatu

islands where Polynesian languages are not spoken today (e.g., Makura, Tongoa, Tongariki

and Tanna) [12]. These results indicate that genetic interactions between ni-Vanuatu and

Polynesian incomers did not systematically trigger shifts to Polynesian languages.

Intriguingly, Polynesian ancestry is not detected north of the Kuwae caldera, a large

submarine volcano that separates Tongoa and Epi islands. Geological data have shown that

the Kuwae volcano erupted in ca. 1452, producing among the largest volumes of magma and

aerosol ever recorded [29, 30], and oral traditions and linguistic evidence suggest that, after

this eruption, Tongoa and Epi were repopulated by distant populations [42, 43]. Our genetic

results support this view; present-day Tongoa and Emae islanders, as well as Epi and

southwest Malekula islanders, show close genetic affinities (Fig. S4), at odds with the gradual

genetic differences expected under isolation by distance. Nowadays, ni-Vanuatu living north

and south of Kuwae form two genetic groups with distinct socio-cultural practices (e.g.,

grade-taking and chiefly title political systems) [44], indicating that the area has remained a

genetic and cultural frontier.

By building upon the well-defined genetic history of Vanuatu, we also explored how

genetic diversity has been shaped by cultural practices. While levels of genetic relatedness are

high among ni-Vanuatu, we do not find evidence for generalized endogamy, which challenges

the frequent association geneticists make between the two processes [17, 45-47]. Nonetheless,

even if ni-Vanuatu spouses are generally less related than non-spouses, we show that their

genetic ancestries are more similar than expected, indicating that mating in Vanuatu is not

random. Importantly, ancestry similarity between partners is not stronger at trait-associated

SNPs, suggesting that ancestry-associated assortments are due to social structure, which may,

in turn, be correlated with levels of East Asian-related and/or Polynesian ancestry. Other

studies have suggested non-random mating according to ancestry in regions of the world

where socio-cultural structure is highly correlated with ancestry [16, 48]. Our findings extend

the occurrence of such socio-cultural assortments to Oceanians, raising questions of how

common this phenomenon is in human societies and whether non-random mating should

systematically be accounted for in human genetic studies. Collectively, our study emphasizes

the need to include diverse populations in genetic studies, not only to address key

anthropological and evolutionary questions that are important for specific geographic regions,

but also to identify factors shaping the genetic diversity of human populations as a whole.

METHODS

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Etienne Patin (epatin@pasteur.fr).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

The SNP array data generated in this study has been deposited to the European Genome-

Phenome Archive (EGA) under accession number EGAS00001005910.

EXPERIMENTAL MODEL AND PARTICIPANT DETAILS

The sampling survey was conducted in the Republic of Vanuatu between April 2003 and

August 2005. The purpose of the study was the estimation of the seroprevalence of HTLV-1

viral infection and the assessment of human genetic diversity in ni-Vanuatu. The recruitment

of participants was carried out after the agreement of the Ministry of Health of Vanuatu, the

head of each Directorate from the sampled province, and the chief of the sampled village. The

data collectors, Olivier Cassar, Helene Walter, Woreka Mera and Antoine Gessain, were

accompanied by the village chief and/or the head of the local dispensary. The nature and the

scope of the study were explained in detail by Olivier Cassar in English, and by Helene

Walter in Bislama (i.e., an English-based pidgin-creole [14] that is the main *lingua franca* of

Vanuatu), during information meetings organized in each village. Participants could ask any

questions after the information meeting. After several hours of reflection, each volunteer

participant of at least 18 years of age was asked to sign a written informed consent form,

including consent for research on human genetic diversity. Sex, age and birth place, as well as

the date and place of blood collection, were collected through a structured questionnaire.

Couples were identified through interviews in English or Bislama, and were preferentially

sampled. Blood samples were collected either at the local dispensary, a gymnasium or a hut

provided by the village chiefs.

The study received approval from the Institutional Review Board of Institut Pasteur

(n°2016-02/IRB/5) and the Ministry of Health of the Republic of Vanuatu. It was conducted

in full respect of the legal and ethical requirements and guidelines for good clinical practice,

in accordance with national and international rules. Namely, research was conducted in

accordance with: (i) ethical principles set forth in the Declaration of Helsinki (Version:

Fortaleza October 2013), (ii) European directives 2001/20/CE and 2005/28/CE, (iii) principles

promulgated in the UNESCO International Declaration on Human Genetic Data, (iv)

principles promulgated in the Universal Declaration on the Human Genome and Human

Rights, (v) the principle of respect for human dignity and the principles of non-exploitation,

non-discrimination and non-instrumentalisation, (vi) the principle of individual autonomy,

(vii) the principle of justice, namely with regard to the improvement and protection of health

and (viii) the principle of proportionality. The rights and welfare of the participants have been

respected, and the hazards did not outweigh the benefits of the study. Feedback to local

communities and partners in Vanuatu is planned for 2023, under the guidance of H.C. The

results of this study will be presented to key stakeholders, including the Ministry of Health

and the Vanuatu Cultural Center (Port Vila). Written and recorded resources will be provided

in Bislama and distributed to interested individuals and communities.

METHOD DETAILS

SNP genotyping and quality filters

Five millilitres of blood were obtained by venepuncture from each volunteer participant and

transferred to the Institut Pasteur of New Caledonia, where plasma and buffy coats were

isolated, frozen, and stored at –80°C. Samples were then transferred to the Institut Pasteur in

Paris (France). Out of the 4,428 collected samples, 1,433 samples were selected for genetic

analyses. These samples were selected in order to (i) cover the largest number of islands and

villages, (ii) cover locations where "Polynesian outliers" exist nowadays and (iii) include as

many couples as possible. After sample selection, a total of 179 different villages were

covered, located on 29 islands (Table S1). DNA was purified from frozen buffy coats at the

Institut Pasteur of Paris (France), using QIAamp DNA Blood Mini Kit protocol, and eluted in

AE buffer. DNA concentration was quantified with the Invitrogen Qubit 3 Fluorometer using

the Qubit dsDNA broad-range assay. Prior to SNP array genotyping, DNA integrity was

checked on agarose gels.

The 1,433 selected samples were genotyped on the Illumina Infinium Omni 2.5-8 array

(San Diego, California). Genotype calling was performed using the Illumina GenomeStudio

software. We excluded 2,491 SNPs with missing annotations, 9,661 duplicated SNPs, 1,772

SNPs with a GenTrain score < 0.4, SNPs with a missingness > 0.05 and SNPs that deviate

from Hardy-Weinberg equilibrium (i.e., P-value < 0.01 in more than one Vanuatu island).

Only autosomal SNPs were kept for the analyses, unless otherwise stated. After filters, a total

of 2,269,868 SNPs were kept. When the analyses required minor allele frequency (MAF)

filters, a MAF > 0.01 threshold was applied. When a linkage disequilibrium pruning was

required, we pruned the data using a window size of 50 Kb, a step size of 5 SNPs and a r^2

threshold of 0.5. After all these filtering steps were applied, the remaining number of SNPs

was 294,806. All filters were applied using PLINK v.1.9b [49].

Sample quality filters

The highest genotype missingness per sample was 0.018. We removed 21 samples with

outlier values for heterozygosity (mean ± 3 SD), suggestive of DNA contamination, leaving

1,412 samples. Cryptic relatedness between samples was detected using KING v.2.1 [50]. We

excluded 456 samples with a kinship coefficient > 0.08 with another sample, whenever

analyses required unrelated samples. For 12 samples, the reported sex did not match the

genetic sex inferred by the Y- and X-chromosome call rates. These include one couple, for

which the reported sex is the opposite to the genetic sex. We thus exchanged the sex of the

two samples, as it is most likely an error on the sample annotation. We did not include the 10

remaining samples, when performing analyses relating to mating practices and sex-specific

migrations. After quality filters, a total of 287 couples was present in the dataset.

Demographic information

Demographic data include the island of birth and the village of residence. Birth place was

missing for 73 samples and the residence for 49 samples. From this information, we retrieved,

for each sample, the geographical coordinates of the birth place (around the center of the

island) and of the place of residence (around the village of residence), using MapAction

reference maps (www.mapaction.org). We considered as the locations of "Polynesian

outliers" all the villages within Futuna, the villages of Mele and Imere in Efate, Ifira island

and the villages of Makatea, Tongamea and Vaitini in Emae [12]. Among the 287 couples,

207 reported the same island of birth, while 57 reported a different island. This information

was missing for 23 couples.

Merging with reference datasets

We merged the new SNP array data with whole genome sequences of different populations

across the Pacific [18] and worldwide populations from the SGDP project [51]. We also

merged the dataset with SNP array data from Oceanian populations [41], to perform some

population genetics analyses. Datasets were merged using PLINK v.1.9b [49]. Transversions

were excluded, to avoid allele strand inconsistences.

Haplotype phasing

We phased the dataset using SHAPEIT v.2 [52, 53] using 500 conditioning states, 10 burn-in

steps, 50 Markov chain Monte Carlo (MCMC) main steps, a window length of 1 cM and an

effective population size of 15,000. We used 1000 Genomes data [54] as a reference panel

and therefore removed all array SNPs that did not align with this data, prior to haplotype

phasing.

Genetic structure: PCA and ADMIXTURE

We performed Principal Component Analyses (PCA) with the 'SmartPCA' algorithm

implemented in EIGENSOFT v. 6.1.4 [55]. aDNA samples were projected on PCs by using

the "lsqproject" and "shrinkmode" options. We inferred population structure with

ADMIXTURE v. 1.3.0 [56], using 10 different seeds and assuming that K_{ADM} varies from 2 to

17, and visualized ADMIXTURE results using pong v1.4.7 [57]. We estimated pair-wise F_{ST}

values using the StAMPP R package [58], and assessed their significance with 100 bootstraps.

We computed f_4-statistics with ADMIXTOOLS v.6.0 and estimated standard errors by block

jackknife.

Genetic structure: ChromoPainter and fineSTRUCTURE

We ran ChromoPainter [19] to infer haplotype sharing among individuals. This algorithm is

based on a Hidden Markov Model in which each sample is treated as a "recipient", i.e., a

mosaic of haplotypes from a set of "donor" samples. We performed different analyses with

ChromoPainter. In Analysis 1, we inferred the genetic structure of ni-Vanuatu running

ChromoPainter only for the unrelated ni-Vanuatu samples. We set each sample both as a

recipient and as a donor for all the samples (-a mode). We first estimated the global mutation

probability and the switch rate with 10 expectation-maximization (EM) iterations, by using

the '-in -iM' options and by averaging estimates obtained for chromosomes 1, 7, 13 and 19.

The estimated values were $1.542{\times}10^{-4}$ and 81.415, respectively. Then, we ran ChromoPainter

in the same mode for all chromosomes but fixing the parameters to estimated values. In

Analysis 2, we repeated the same process independently for the reference populations. In this

case, we ran ChromoPainter only for the samples outside Vanuatu [18, 51] and repeated the

two-step process described above. In this case, the global mutation probability was estimated

to 4.53×10^{-4} and the switch rate was 255.42.

We then produced a coancestry matrix by summing the coancestry matrices for all

chromosomes and estimated the *C* parameter with ChromoCombine. The *C* factor was

estimated at 0.452 for the Vanuatu dataset (Analysis 1), and 0.498 for the reference

population dataset (Analysis 2). We applied the model-based Bayesian clustering method

fineSTRUCTURE v.2.0.7 [19] on ChromoPainter coancestry matrices. We ran 1 million

burn-in iterations (-y option) and 1 million MCMC iterations (-x option) and sampled values

every 10,000 iterations. We then inferred the fineSTRUCTURE tree (-m T option). We

repeated this process 5 times, with different random seeds.

Following a previous study [20], we estimated for each individual the robustness of the

clustering assignation, by comparing the final state (i.e., the state with the highest posterior

probability) with the 100 MCMC samples. For each individual *i*, we estimated $x_i^{(m)}$, the

number of individuals clustering with *i* for each of the $m = 1,…,100$ MCMC samples. We

also estimated $y_{ik}^{(m)}$, the number of individuals that cluster with *i* in the final state and in each

of the MCMC samples, for each inferred cluster *k*. Finally, we computed

$P_i = \sum_m [y_{ik}^{(m)} / x_i^{(m)}]$. Therefore, we ended up with a matrix of this sum P_i, for all individuals

and for each genetic cluster, that shows the robustness in the assignation of each individual to

its corresponding cluster in the final state. We repeated this process for $k = 2,…,K$, *K* being

the maximum number of clusters identified by fineSTRUCTURE. We also repeated the

process for the 5 random seeds and compared the performance of each seed. Fig. S6 shows

the number of ni-Vanuatu individuals with $P_i < 0.9$ for each seed *s* and for each $k = 2,…K$.

These estimations allowed to test the robustness in the cluster assignation of each individual,

to compare the performance across seeds and to determine the robustness at each *k* value.

To compare robustness among seeds, we followed a similar approach. We estimated $x_i^{(s)}$,

where $s = 1,…,5$, which corresponds to the number of individuals clustering with individual *i*

for each seed. We then estimated $y_{ik}^{(s)}$ and $\sum_s [y_{ik}^{(s)} / x_i^{(s)}]$, which shows the concordance in the

assignation of each individual to a cluster across the different seeds. To summarize this

information, we analyzed all the individual sums, for each seed and each cluster *k* (Fig. S7A

and C), as well as the number of individuals who were not assigned to the same cluster,

between one seed and all the other seeds (Fig. S7B and D).

Based on these analyses, we chose the seed that showed the highest consensus among

seeds (Fig. S7) and in which the largest number of samples are robustly assigned (Fig. S6)

(seed number 289 for Analysis 1 and 161 for Analysis 2). Then, we defined the maximum k

value based on the number of individuals who are robustly assigned to a cluster for each k

value and we focused our analyses on this value. Finally, once we chose a seed and a k value,

we removed samples that showed some uncertainty in the cluster assignation ($P_i \leq 0.9$).

Following this process, we observed that the robustness decreases from $K_{FS} = 20$ and we

removed 3 samples from Analysis 1 (Vanuatu). For the reference dataset (Analysis 2), all

samples showed a $P_i > 0.9$ across all k values, therefore we used genetic clusters from the k

value that best suited the hypotheses to test ($K_{FS} = 25$) (Table S5).

For the subsequent analyses, we redefined populations based on fineSTRUCTURE results

and therefore on the genetic data itself. This approach is rather trivial when applied to

continental populations, like the reference populations presented here, because

fineSTRUCTURE clusters coincide broadly with the geographical or cultural categories

assigned to the populations. Yet, this strategy has proven useful when complex admixture

shapes the genetic structure [59, 60], or when the studied populations are sampled from a

small geographic region [20], in which case, defining populations based on geography or

culture may lead to a biased view of the genetic structure of the populations.

Ancestry estimation

We estimated admixture proportions for each ni-Vanuatu individual using SOURCEFIND

[61]. We first ran ChromoPainter using the "donors" mode: we considered each ni-Vanuatu

individual as a recipient only and defined donor populations by using the genetic clusters

defined above (Analysis 2). We ran SOURCEFIND by considering all surrogates ("all

surrogates" analysis; Fig. S15 and Table S5) or by limiting as much as possible the number of

surrogates ("limited surrogates" analysis). In the latter case, we considered as surrogates

West_Eurasia1, West_Eurasia2, West_Eurasia3, EastAsia1, EastAsia2, EastAsia3, Atayal,

Paiwan, Southeast_Asia, Cebuano, RenBell, Tikopia, PNG1, PNG2 and PNG_SGDP. Then,

we summed the estimated ancestry proportions for these surrogates and grouped them as

follows: Taiwanese.Philippines (Atayal, Paiwan, Cebuano), EastAsian_mainland (EastAsia1,

EastAsia2, EastAsia3, Southeast_Asia), Polynesian (RenBell, Tikopia), Papuan-related

(PNG1, PNG2 and PNG_SGDP) and European (West_Eurasia1, West_Eurasia2,

West_Eurasia3). The East Asian-related ancestry was estimated as the sum of

Taiwanese.Philippines group and Polynesian group (the EastAsian_mainland was negligible,

with a maximum proportion = 0.007). To facilitate visualization, we excluded an individual

born in Futuna and living in Malekula, who shows a Polynesian ancestry of 0.745 in figures

showing ancestry proportions (Fig. 2, Fig. S8-11). The next individual carrying the highest

Polynesian ancestry shows 0.383. For both the "all surrogates" and "limited surrogates"

analyses, we ran 200,000 MCMC iterations with a burn-in of 50,000 iterations, sampling

every 5,000 iterations and not allowing for self-copy. We estimated the final ancestry of each

ni-Vanuatu sample as the mean of the 30 sampled MCMC runs.

Admixture date estimation

We estimated admixture dates using GLOBETROTTER [62]. For this analysis, we ran

ChromoPainter using the "donors" mode, by considering the ni-Vanuatu genetic clusters

defined in (Analysis 1) as recipients and all the reference populations (Analysis 2) as both

recipients and donors. Surrogates were the same as those considered for the "limited

surrogates" SOURCEFIND analysis. For each of the 20 recipient ni-Vanuatu clusters, we

performed 100 bootstraps, which are implemented as resamplings of chromosomes among the

available samples. We set the "Null.ind" option to 1. We assumed a generation time of 28

years, to estimate dates from the number of generations. We also dated admixture in each ni-

Vanuatu sample following the same approach.

Sex-biased admixture

We studied sex-biased admixture by comparing ancestry proportions estimated for the

autosomes and the X chromosome. We estimated ancestry proportions using RFMix v.1.5.4

[33], considering as ancestral populations New Guinean highlanders (approximating Papuan-

related ancestry) and Taiwanese Indigenous peoples and the Cebuano from the Philippines

(approximating East Asian-related ancestry). We ran the "TrioPhased" algorithm, allowing

for phase correction and 3 EM iterations. The window size was set at 0.03 cM [18] and the

admixture date was set at 50 generations. The same parameters were used for the X

chromosome and the autosomes. We combined the X chromosome haploid data of males from

the same island to obtain diploid individuals. We filtered out SNPs with an RFMix posterior

probability < 0.9, those within centromeres and within 2 Mb from the telomeres. We then

estimated East Asian-related and Papuan-related ancestry in the autosomes and the X

chromosome, separately. We estimated α_f and α_m, that is, the proportion of female and male

ancestors of ni-Vanuatu who carried Papuan-related ancestry, respectively, by assuming that

admixture proportions have reached equilibrium values [34]. In such a case, $\alpha_f = 3\alpha_X -$

$2\alpha_{auto}$ and $\alpha_m = 4\alpha_{auto} - 3\alpha_X$, where α_{auto} and α_X are the average Papuan-related

ancestry proportions estimated on the autosomes and the X chromosome, respectively.

To avoid potential biases due to SNP ascertainment, we also studied sex-biased admixture

by comparing the autosomes and the sex chromosomes of 179 ni-Vanuatu sequenced at 30×

coverage [18]. Because genotypes on the X chromosome were not called in the previous

study, we mapped *fastq* files on the X chromosome of the human reference genome (version

hs37d5) and performed genotype calling for X-linked variants, as previously described [18],

setting the ploidy parameter to 1 for males and 2 for females. We kept only biallelic SNPs and

filtered *vcf* files following GATK best practices [63]: QualByDepth < 2; FisherStrand value >

60; StrandOddsRatio > 3; RMSMappingQuality < 40; MappingQualityRankSumTest < -12.5;

and ReadPosRankSum < -8. We also removed genotypes with a DP < 10 for females and < 3

for males, and those with a GQ < 30. We removed PAR1, PAR2 and XTR regions according

to the annotations provided by the UCSC browser, and amplicon regions as reported in [64].

We filtered out SNPs that were not in Hardy-Weinberg equilibrium in females (*P*-value <

0.0001), those with a missingness > 0.05 and those with a MAF < 0.01. We estimated mtDNA

haplogroups with MToolBox [65] from the *bam* and *fastq* files, and Y chromosome

haplogroups were estimated with Yleaf [66].

665 Inferring migrations within Vanuatu

We used the genetic structure inferred by fineSTRUCTURE to study migration patterns

among Vanuatu islands. First, we assigned each genetic cluster to one or more islands, each

time more than 25% of the sampled individuals inhabiting the island were assigned to that

cluster. We then identified "outlier" individuals who inhabit an island but are assigned to a

genetic cluster that is prevalent in another island. These individuals, either themselves or their

close ancestors, likely migrated from the island/s where their genetic cluster is predominant to

their current place of residence. We removed from these analyses Vanuatu_8 and Vanuatu_9

clusters ($K_{FS} = 20$) because those clusters are driven by recent European gene flow, as well as

Vanuatu_15 because no island could be assigned to this cluster when using the 25% rule. To

study sex-specific migrations, we calculated the proportion of female (male) migrants by

dividing the number of "outlier" females (males) by the number of females (males) in each

cluster. We did not include clusters where the number of males or females was < 2.

679 Tests for exogamy

We tested if spouses show lower genetic relatedness than non-spouses by testing if the

average kinship coefficient between 287 observed couples is higher or lower than expected by

chance, using either permutations or a logistic regression model. Kinship coefficients were

computed between all possible pairs of individuals using KING v.2.1 (Table S7). When using

permutations, we accounted for isolation by distance by sampling random pairs of males and

females among individuals born in the same island (Fig. S24A) or living in the same village

(Fig. S24B). Because marriages between first-degree related individuals are very unlikely

(they are indeed not observed in the data), we excluded from this analysis individuals who are

first-degree relatives, but also tested how the inclusion of these individuals affect the results

(Fig. S24C). We performed 10,000 permutations for each island. In each permutation, we

sampled equally many random pairs as there were observed pairs in the island. We calculated

P-values by comparing the average kinship coefficient among the observed couples to the null

distribution. The null distribution was made of 10,000 average kinship coefficients between

randomly sampled pairs of individuals from the same island or village.

Based on the sample scheme described above, we designed a logistic regression model

that could be generalized to multiple explanatory variables, such as ancestry. Let the variable

i index all possible pairs of males and females in the dataset, except pairs of individuals who

are first-degree relatives, and define the dependent binary variable Y_i with $Y_i = 1$ if i indexes

an observed pair of spouses and $Y_i = 0$ otherwise. Define the probability that a pair is an

observed couple $p_i(x) = P(Y_i = 1)$, and introduce ϕ_i as the kinship coefficient between

individuals in the i:th pair. We estimated the effect of kinship on mate choice by the

parameter β^ϕ in the logistic regression model

$$\log\left[\frac{p_i(x)}{1 - p_i(x)}\right] = \mu + \phi_i\beta^\phi + I(village_i)\beta^v + I(island_i)\beta^I, \qquad Eq.\ 1$$

where $I(village_i) = 1$ if individuals of pair i are from the same village and $I(village_i) = 0$

otherwise, and $I(island_i) = 1$ if individuals of pair i are from the same island and

$I(island_i) = 0$ otherwise. As a sensitivity analysis, we also considered a model in which

pairs of first-degree related individuals were included (Fig. S25).

Tests for ancestry-based assortative mating

We extended the logistic regression model shown in $Eq.\ 1$ to test for ancestry-based

assortative mating, by testing if the genetic ancestry of spouses is more similar than that of

non-spouses, accounting for population structure and relatedness avoidance. Let q_{i1}^{EA} and q_{i2}^{EA}

be the proportion of East Asian-related ancestry for the individuals 1 and 2 of pair i and

introduce the variable $q_i^{EA} = |q_{i1}^{EA} - q_{i2}^{EA}|$. We estimated the effect of having similar

proportions of East Asian-related ancestry between two individuals on the probability that a

pair is an observed couple $p_i(x)$ by the parameter β^q in the logistic regression model,

$$\log\left[\frac{p_i(x)}{1 - p_i(x)}\right] = \mu + q_i^{EA}\beta^q + \phi_i\beta^\phi + I(village_i)\beta^v + I(island_i)\beta^I, \qquad Eq.\ 2$$

where we use the same notations as in *Eq. 1*. A negative (positive) effect size β^q is interpreted

as evidence for assortative (disassortative) mating according to ancestry. Effect sizes for other

ancestries were calculated similarly. The same model was also tested in ni-Vanuatu

originating or not from islands where Polynesian languages are spoken. Note that in standard

regression analyses, such as those used in GWAS, population stratification is usually

corrected by principal components of the genetic relatedness matrix; here we have taken a

more general approach by decomposing the genetic structure in two variables: the kinship and

the ancestry, and have studied how both variables affect mate choice independently.

Tests for SNP-based assortative mating

We extended the logistic regression model shown in *Eq. 1* to test for SNP-based assortative

mating, by testing if the genotypes at a given SNP are more similar between spouses than

non-spouses, accounting for population structure, ancestry-based associated mating and

relatedness avoidance. Define, for each SNP s and each pair i of individuals, the allele

sharing distance (ASD) $d_{i,s}$ as $d_{i,s} = 0$ if both alleles are identical between individuals,

$d_{i,s} = 1$ if only one allele is identical and $d_{i,s} = 2$ if none of the alleles are identical. We

estimated the association between SNP s and mate choice by the parameter β_s^{ASD} in the model

$$\log\left[\frac{p_i(x)}{1 - p_i(x)}\right] = \mu + d_{i,s}\beta_s^{ASD} + q_i^{EA}\beta^q + \phi_i\beta^\phi + I(village_i)\beta^v + I(island_i)\beta^I, \qquad Eq.\ 3$$

736 where we use the same notations as in *Eq. 2*. A negative (positive) effect size β_s^{ASD} indicates

737 higher (lower) similarity at SNP s between the members of the observed couples than

738 between non-couples, in line with SNP-based assortative (disassortative) mating.

Tests for trait-based assortative mating

We tested for trait-based assortative mating by testing if genotypes of spouses are more similar or dissimilar at trait-associated SNPs, relative to non-associated SNPs. We obtained GWAS summary statistics for 8 candidate traits from the UK Biobank database (http://www.nealelab.is/uk-biobank). Candidate traits include traits relating to morphology and physical appearance. Let $y_s = \beta_s^{ASD}$, where β_s^{ASD} is the effect size of SNP s on mate choice estimated by *Eq. 1*. We estimated if trait-associated SNPs are more similar or dissimilar between spouses by the parameter β^{trait} in the model

$$y_s = \mu + I(associated_s)\beta^{trait} + MAF_s\beta^{MAF} + GERP_s\beta^{GERP} + rec_s\beta^{rec}, \qquad \textit{Eq. 4}$$

where $I(associated_s) = 1$ if SNP s is significantly associated with the candidate trait in GWAS (with GWAS P-value $< 5{\times}10^{-8}$) and $I(associated_s) = 0$ otherwise, MAF_s is the minor allele frequency of SNP s in ni-Vanuatu, $GERP_s$ is the Genomic Evolutionary Rate Profiling (GERP) score of SNP s and rec_s is the interpolated recombination rate between SNP s and SNP $s-1$, estimated in cM/Mb from the 1000 Genomes Phase 3 combined recombination map. We considered that there is trait-based assortative (disassortative) mating if β^{trait} is significantly negative (positive). We adjusted P-values with the Bonferroni correction for multiple testing, to account for the number of traits tested.

ACKNOWLEDGMENTS

We thank all volunteers in Vanuatu who kindly provided samples and participated in this research. We thank the Ministry of Health of the Republic of Vanuatu (especially Myriam Abel, Rose Bahor, Yvannah Taga, Maturine Tary and the former Minister Jerôme Ludvaune), the Vanuatu Family Health Association (especially Blandine Boulekone and Marie Nickllum Wala), the WHO Office in Port Vila (especially Corinne Capuano) and the Institut Pasteur of New Caledonia at Nouméa (especially Paul M. V. Martin and Eliane Chungue), for their continuous support and interest in this work. We are particularly thankful to Helene Walter, Woreka Mera, Françoise Charavay and Sonia Treptow for their assistance in collecting samples. We thank S. Créno and the HPC Core Facility of Institut Pasteur (Paris) for the management of computational resources. L.R.A. was funded by a Pasteur-Roux-Cantarini fellowship from the Institut Pasteur. The laboratory of Human Evolutionary Genetics is supported by the Institut Pasteur, the Collège de France, the CNRS, the Fondation Allianz-Institut de France, the French Government's Investissement d'Avenir programme, Laboratoires d'Excellence 'Integrative Biology of Emerging Infectious Diseases' (ANR-10-LABX-62-IBEID) and 'Milieu Intérieur' (ANR-10-LABX-69-01), the Fondation de France (n°00106080), and the Fondation pour la Recherche Médicale (Equipe FRM DEQ20180339214) and the French National Research Agency (ANR-19-CE35-0005).

AUTHOR CONTRIBUTIONS

Conceptualization: L.R.A., E.P.; L.Q.-M.; Methodology: L.R.A., J.B., A.M.S., E.P.; Software: L.R.A., M.R.; Formal Analysis: L.R.A.; Data Generation: C.H., L.L.; Resources: O.C., A.G.; Writing – Original Draft: L.R.A., L.Q.M., E.P.; Writing – Review & Editing: L.R.A., J.B., J.C., J.M.R., M.R., A.M.S., H.C., A.F., F.V., L.Q.-M., E.P.; Visualization: L.R.A.; Supervision: L.Q.-M., E.P.; Funding Acquisition: L.Q.M.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. Sampling locations and genetic structure in Vanuatu.

(A) Map showing the sampling location of the 1,439 contemporary ni-Vanuatu individuals. The inset at the bottom left shows the location of Vanuatu in the Asia-Pacific region. Noise was added to sampling locations to facilitate visualization.

(B,C) Principal Component Analysis (PCA) of genotypes of ni-Vanuatu at 301,774 SNPs, in the context of (B) the broad Asia-Pacific region and (C) the Vanuatu archipelago only.

(A-C) Each point indicates an individual, colored according to the latitude of their island of residence. (B,C) Black crosses and points indicate projected ancient samples from Vanuatu dated to the Lapita and Post-Lapita periods, respectively.

Figure 2. Fine-scale genetic structure and admixture in Vanuatu.

(A) Map showing the clustering of 979 ni-Vanuatu into 20 genetic clusters, according to fineSTRUCTURE (K_{FS} = 20). Each point indicates an individual, located according to their village of residence. Colors indicate genetic clusters, so that the closer the colors, the closer the clusters. Noise was added to sampling locations to facilitate visualization.

(B) Admixture date estimates for each genetic cluster, based on 100 bootstrap replicates. Colors are the same as those used in (A). The x-axis shows the admixture date in years before present, assuming a generation time of 28 years. The y-axis indicates the latitude of the islands assigned to each genetic cluster. The heights of the density curves are proportional to the sample size of each cluster.

(C) Proportion of East Asian-related ancestry estimated by SOURCEFIND.

(D) Ratio of the Polynesian and East Asian-related ancestry proportions, estimated by SOURCEFIND. The inset at the bottom left shows the location of Polynesian-speaking individuals, indicated by black points.

(C-D) Each point indicates an individual, colored according to their ancestry proportions.

Figure 3. Sex-biased admixture and migration patterns in Vanuatu.

(A) Papuan-related ancestry proportions in ni-Vanuatu, estimated for the 22 autosomes and the X chromosome separately by RFMix.

(B) Frequencies of Y chromosome and mtDNA haplogroups, colored according to their assumed origins (i.e., shades of blue or red indicate Papuan- or East Asian-related origins, respectively). Haplogroups were inferred from high-coverage genome sequencing data obtained for a subset of 179 ni-Vanuatu [18].

(C) Recent migrations among Vanuatu islands, inferred based on fineSTRUCTURE clusters. The arrows connect the location of the genetic cluster to which individuals were assigned to their actual place of residence. The colors indicate the predominant genetic cluster in the island of origin. The width is proportional to the number of inferred migrant individuals, relative to the number of females or males in the genetic cluster.

Figure 4. Ancestry- and trait-based assortative mating in Vanuatu.

(A) Effects of kinship and ancestry differences on partner choice. Effect sizes were estimated with a logistic regression model, while accounting for population structure (island of birth and village of residence). The effect size and *P*-value were estimated using different ancestries as predictors, independently.

(B) Increased (purple) or decreased (orange) genetic similarity among spouses (as measured by β_S^{ASD}) at trait-associated SNPs, relative to non-associated SNPs. Results on the left are based on a logistic model that includes only the genotype dissimilarity among spouses at each SNP, whereas results on the right are based on the logistic model that also controls for possible confounders (i.e. geography, kinship and ancestry differences).

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Genetic structure of ni-Vanuatu and a selection of worldwide populations.

(A) ADMIXTURE analyses of the SNP array data for 1,439 ni-Vanuatu, together with whole

genome sequences of other populations from the Pacific region [18]. (B) ADMIXTURE

cross-validation errors for different K_{ADM} values and 10 iterations with different seeds,

indicated by colors.

Figure S2. Principal Component Analyses of the SNP array dataset for ni-Vanuatu.

Each point is an individual, colored according to the geographical latitude coordinates of their

island of residence, as in Fig. 1A. Percentages indicate the proportion of variance explained.

PC1 and PC2 are shown in Fig. 1. Black crosses and points indicate projected ancient samples

from Vanuatu dated to the Lapita and Post-Lapita periods, respectively.

Figure S3. F_{ST} matrix between ni-Vanuatu and other worldwide populations. F_{ST} values

and their significance were obtained with the StAMPP R package [58]. The lower triangle of

the matrix indicates F_{ST} values, whereas the upper triangle indicates P-values. All values were

significant (P-value < 0.01, using 100 bootstraps).

Figure S4. Map showing the clustering of ni-Vanuatu into $K_{FS} = 2$ to 18 genetic clusters,

according to fineSTRUCTURE. Each point indicates an individual, located according to

their village of residence. Colors indicate clusters, so that the closer the colors, the closer the

clusters. Noise was added to sampling locations to facilitate visualization.

Figure S5. Sample sizes of each genetic cluster, according to fineSTRUCTURE at $K_{FS} =$

20. Filled bars indicate the number of individuals who live in a village where Polynesian

languages are spoken.

Figure S6. Robustness of fineSTRUCTURE analyses. (A) Number of individuals below a

0.9 threshold of confidence in the assignation to a specific cluster (y-axis), as a function of a

number of clusters K_{FS} that varies between 1 and 125 (x-axis). Colors indicate the 5 different

seeds used for the MCMC algorithm. (B) Zoom-in of the plot in (A), for K_{FS} between 1 and

50. As expected, the robustness on the assignation of each individual to a genetic cluster

decreases with increasing K_{FS} values (i.e., more samples are below the 0.9 threshold). (C)

Robustness as measured by the average of $\Sigma_m(Y_i / X_i)$ (Methods), as a function of a number of

clusters K that varies between 1 and 125 (x-axis). (D) Zoom-in of the plot in (C), for K_{FS}

between 1 and 50. The shaded areas indicate the standard deviation of $\Sigma_m(Y_i / X_i)$ for each K_{FS}

value.

Figure S7. Robustness of fineSTRUCTURE analyses across seeds, for the chosen K_{FS}

values. (A,C) Heatmaps showing the average of $\Sigma_S(Y_i / X_i)$ across clusters for (A) the ni-

Vanuatu (Analysis 1) (K_{FS} = 20) and (C) the reference populations (Analysis 2) (K_{FS} = 25).

Each row corresponds to one seed (the number of which is circled in black), and each column

corresponds to a genetic cluster. Colors indicate the mean $\Sigma_S(Y_i / X_i)$ for each cluster and seed.

(B,D) Heatmaps showing the mismatch between each seed and the four other seeds for (B) the

ni-Vanuatu (Analysis 1) (K_{FS} = 20) and (D) the reference populations (Analysis 2) (K_{FS} = 25).

Each row corresponds to the seed used as reference to estimate $\Sigma_S(Y_i / X_i)$. Each column

corresponds to a genetic cluster. Colors indicate the proportion of individuals who are

assigned to a different cluster in the other seeds.

Figure S8. Ancestry proportions in ni-Vanuatu, estimated by SOURCEFIND. Each point

indicates an individual, colored according to their ancestry proportions. The East Asian-

related ancestry is the sum of the Taiwan.Philippines and Polynesian ancestry. Noise was

added to sampling locations to facilitate visualization.

Figure S9. Correlations between East Asian-related and Polynesian ancestry inferred by

SOURCEFIND. (A) Each individual is colored according to their island of residence. (B)

The black points indicate the individuals living in villages where Polynesian languages are

spoken, the rest of the samples are colored in gray. The blue line indicates the regression line

considering all the individuals from Vanuatu. The gray area indicates the 95% confidence

level.

Figure S10. Distributions of ancestry proportions in ni-Vanuatu, estimated by

SOURCEFIND, for each island and ancestry. The line, box, whiskers and points,

respectively, indicate the median, interquartile range (IQR), 1.5*IQR and outliers. Colors

indicate the geographical latitude coordinates of each island, as in Fig. 1A.

Figure S11. Distributions of ancestry proportions of ni-Vanuatu among genetic clusters,

according to fineSTRUCTURE at K_{FS} = 4. The line, box and whiskers, respectively,

indicate the median, interquartile range (IQR) and 1.5*IQR. We tested significant differences

in ancestry between each cluster using a Wilcoxon test with Bonferroni correction. $*P < 0.05$,

$**P < 0.01$, $***P < 0.001$, $****P < 0.0001$, ns: $P > 0.05$.

Figure S12. Map showing admixture dates estimated per individual with

GLOBETROTTER.

Figure S13. Allele sharing of ni-Vanuatu with Near Oceanians, according to f_4-statistics.

The x-axis indicates f_4(X, New Guinean Highlanders; EastAsian, Australian), which measures

allele sharing of ni-Vanuatu and Near Oceanians (X) with East Asians. The y-axis indicates

f_4(X, New Guinean Highlanders; Near Oceanian, EastAsian), which measures allele sharing

of ni-Vanuatu and Near Oceanians (X) to each Near Oceanian population tested (Nakanai,

Nasioi, Baining), relative to their allele sharing with New Guinean Highlanders. The

regression line is estimated considering only modern individuals from Vanuatu. The bars

show two standard errors. Deviations upper from the regression line indicate a higher affinity

with the Near Oceanian population tested.

Figure S14. Principal Component Analysis (PCA) of ni-Vanuatu, together with other

populations from Near Oceania. The modern and ancient samples from Vanuatu were

projected into a PCA of a previous SNP array data set [41].

Figure S15. Ancestry proportions in ni-Vanuatu, estimated by SOURCEFIND,

considering all the reference populations as possible surrogates (Table S4). The bar plot

indicates the mean value across individuals, for each ancestry in each island of residence.

Figure S16. Allele sharing of ni-Vanuatu with Polynesians, according to f_4-statistics. The

x-axis indicates f_4(X, New Guinean Highlanders; EastAsia, Australian), which measures allele

sharing of ni-Vanuatu and Near Oceanians (X) with East Asians. The y-axis indicates f_4(X,

EastAsian; Polynesian, Tolai), which measures allele sharing of ni-Vanuatu and Near

Oceanians (X) with Polynesians (Rennell and Bellona [RenBel], Tikopia, Tonga), relative to

their East Asian ancestry. The regression line is estimated considering only modern

individuals from Vanuatu. The bars show two standard errors. Deviations upper from the

regression line indicate a higher affinity with the Polynesian population tested.

Figure S17. Sex-biased admixture in Vanuatu. (A) X-to-autosome ratio of Papuan-related ancestry proportions in ni-Vanuatu who speak non-Polynesian or Polynesian languages, estimated using the SNP array data (Wilcoxon test P-value $< 1.36 \times 10^{-5}$, after Bonferroni correction). (B) Papuan-related ancestry proportions in ni-Vanuatu, estimated for the 22 autosomes and the X chromosome separately, using the high-coverage genome sequencing data obtained for a subset of 179 ni-Vanuatu [18].

Figure S18. Self-reported migrations among ni-Vanuatu spouses. (A) Map of self-reported migrations for females (blue) and males (green), separately. The arrows connect the place of birth of individuals to their place of residence. (B) Barplot showing the number of females (blue) and males (green) among ni-Vanuatu spouses who reported to migrate during their lifetime.

Figure S19. Proportions of female and male ni-Vanuatu migrants, based on genetic clusters or self-reported information.

Figure S20. Comparison of kinship levels among ni-Vanuatu living in the same village, the same island or the entire archipelago. Significance of the differences was tested by a Wilcoxon test.

Figure S21. Genetic relatedness and islands of residence of related ni-Vanuatu. Each line connects two individuals who are related up to third degree. Islands are order from north to south (clockwise).

Figure S22. **Degrees of genetic relatedness of ni-Vanuatu from the different Vanuatu islands.** Bar plots indicate the proportion of pairs of related individuals according to the inferred degree of genetic relatedness, in each island. The black points and indicate the mean kinship per island. Significance was tested by a Fisher's exact test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure S23. Genetic relatedness and islands of residence of ni-Vanuatu spouses. Each line connects the two spouses, and the color indicates the degree of genetic relatedness. Islands are order from north to south (clockwise).

Figure S24. **Kinship among spouses and random pairs of individuals.** Kinship coefficients
among spouses and random pairs of individuals from the same (A) island (excluding first-
degree related pairs), (B) village (excluding first-degree related pairs) and (C) village
(including first-degree related pairs). The orange vertical line indicates the average kinship
coefficient among spouses. The blue density curve indicates the null distribution of kinship
coefficients estimated from the sampling of random pairs of individuals. Each data point of
the null distribution is estimated as the mean kinship coefficient in randomly sampled pairs of
individuals. Of note, negative kinship coefficients are expected when the two compared
individuals are from different populations.

Figure S25. **Effects of geographical location and kinship on mate choice.**. (A) Effect sizes
estimated when not including the village of residence as a predictor. (B) Effect sizes
estimated when including the village of residence as a predictor and excluding first-degree
related pairs from the analysis. (C) Effect sizes estimated when including the village of
residence as a predictor and including first-degree related pairs from the analysis. (A-C)
Effect sizes of the logistic regression modelling the probability of mating as a function of
three predictors: the island of birth, the village of residence and kinship coefficients.

1005

1006 **Figure S26. Q-Q plots of observed and expected *P*-values of a regression model testing**
1007 **for SNP-based assortative mating**. (A) *P*-values are from a model that does not control for
1008 population structure. (B) *P*-values are from a model that controls for population structure
1009 (island of birth, kinship and ancestry). The shaded area indicates the 95% confidence interval.
1010

REFERENCES

1. Bellwood, P. (2004). First Farmers: The Origins of Agricultural Societies, (Wiley-Blackwell).

2. Gray, R.D., Drummond, A.J., and Greenhill, S.J. (2009). Language phylogenies reveal expansion pulses and pauses in Pacific settlement. Science *323*, 479-483.

3. Hung, H.-C., and Carson, M.T. (2015). Foragers, fishers and farmers: origins of the Taiwanese Neolithic. Antiquity *88*, 1115-1131.

4. Bedford, S., and Spriggs, M. (2014). The archaeology of Vanuatu: 3000 years of history across islands of ash and coral, (Oxford University Press).

5. Petchey, F., Spriggs, M., Bedford, S., and Valentin, F. (2015). The chronology of occupation at Teouma, Vanuatu: Use of a modified chronometric hygiene protocol and Bayesian modeling to evaluate midden remains. Journal of Archaeological Science: Reports *4*, 95-105.

6. Petchey, F., Spriggs, M., Bedford, S., Valentin, F., and Buckley, H. (2014). Radiocarbon dating of burials from the Teouma Lapita cemetery, Efate, Vanuatu. J Archaeol Sci, 227-242.

7. Valentin, F., Detroit, F., Spriggs, M.J., and Bedford, S. (2016). Early Lapita skeletons from Vanuatu show Polynesian craniofacial shape: Implications for Remote Oceanic settlement and Lapita origins. Proc Natl Acad Sci U S A *113*, 292-297.

8. Skoglund, P., Posth, C., Sirak, K., Spriggs, M., Valentin, F., Bedford, S., Clark, G.R., Reepmeyer, C., Petchey, F., Fernandes, D., et al. (2016). Genomic insights into the peopling of the Southwest Pacific. Nature *538*, 510-513.

9. Lipson, M., Skoglund, P., Spriggs, M., Valentin, F., Bedford, S., Shing, R., Buckley, H., Phillip, I., Ward, G.K., Mallick, S., et al. (2018). Population Turnover in Remote Oceania Shortly after Initial Settlement. Current biology *28*, 1157-1165 e1157.

10. Posth, C., Nagele, K., Colleran, H., Valentin, F., Bedford, S., Kami, K.W., Shing, R., Buckley, H., Kinaston, R., Walworth, M., et al. (2018). Language continuity despite population replacement in Remote Oceania. Nat Ecol Evol *2*, 731-740.

11. Flexner, J.L., Bedford, S., and Valentin, F. (2019). Who was Polynesian? Who was Melanesian? Hybridity and ethnogenesis in the South Vanuatu Outliers. Journal of Social Archaeology *19*, 403–426.

12. Hermann, A., and Walworth, M. (2020). Approche interdisciplinaire des échanges interculturels et de l'intégration des communautés polynésiennes dans le centre du Vanuatu. Journal de la société des océanistes *151*, 239-262.

13. Lipson, M., Spriggs, M., Valentin, F., Bedford, S., Shing, R., Zinger, W., Buckley, H.,
Petchey, F., Matanik, R., Cheronet, O., et al. (2020). Three Phases of Ancient Migration
Shaped the Ancestry of Human Populations in Vanuatu. Curr Biol *30*, 4846-4856 e4846.

14. François, A., Lacrampe, S., Franjieh, M., and Schnell, S. (2015). The Languages of
Vanuatu: Unity and Diversity, Volume 5, (Canberra: Asia–Pacific Linguistics).

15. Heyer, E., Chaix, R., Pavard, S., and Austerlitz, F. (2012). Sex-specific demographic
behaviours that shape human genomic variation. Mol Ecol *21*, 597-612.

16. Zou, J.Y., Park, D.S., Burchard, E.G., Torgerson, D.G., Pino-Yanes, M., Song, Y.S.,
Sankararaman, S., Halperin, E., and Zaitlen, N. (2015). Genetic and socioeconomic study
of mate choice in Latinos reveals novel assortment patterns. Proc Natl Acad Sci U S A
112, 13621-13626.

17. Ceballos, F.C., Joshi, P.K., Clark, D.W., Ramsay, M., and Wilson, J.F. (2018). Runs of
homozygosity: windows into population history and trait architecture. Nat Rev Genet *19*,
220-234.

18. Choin, J., Mendoza-Revilla, J., Arauna, L.R., Cuadros-Espinoza, S., Cassar, O., Larena,
M., Ko, A.M., Harmant, C., Laurent, R., Verdu, P., et al. (2021). Genomic insights into
population history and biological adaptation in Oceania. Nature *592*, 583-589.

19. Lawson, D.J., Hellenthal, G., Myers, S., and Falush, D. (2012). Inference of population
structure using dense haplotype data. PLoS Genet *8*, e1002453.

20. Leslie, S., Winney, B., Hellenthal, G., Davison, D., Boumertit, A., Day, T., Hutnik, K.,
Royrvik, E.C., Cunliffe, B., Lawson, D.J., et al. (2015). The fine-scale genetic structure of
the British population. Nature *519*, 309-314.

21. van Dorp, L., Balding, D., Myers, S., Pagani, L., Tyler-Smith, C., Bekele, E., Tarekegn,
A., Thomas, M.G., Bradman, N., and Hellenthal, G. (2015). Evidence for a Common
Origin of Blacksmiths and Cultivators in the Ethiopian Ari within the Last 4500 Years:
Lessons for Clustering-Based Inference. PLoS Genet *11*, e1005397.

22. Tryon, D.T. (1976). New Hebrides Languages: An Internal Classification, (Canberra:
Australian National University).

23. François, A. (2011). Where *R they all? The Geography and History of *R-loss in
Southern Oceanic Languages. Oceanic Linguistics *50*, 140-197.

24. Lynch, J.D., Ross, M.T., and Crowley, T. (2011). The Oceanic Languages, (Routledge).

25. Huffman, K. (1996). Trading, cultural exchange and copyright: Important aspects of
Vanuatu art. In Arts of Vanuatu, J. Bonnemaison, K. Huffman, C. Kaufmann and D.
Tryon, eds. (Bathurst: Crawford House Press), pp. 182-194.

26. Walworth, M., Dewar, A., Ennever, T., Takau, L., and I., R. (2021). Multilingualism in Vanuatu: Four case studies. International Journal of Bilingualism *25*, 1120-1141.

27. Lynch, J.D., and Fakamuria, K. (1994). Borrowed Moieties, Borrowed Names: Sociolinguistic Contact between Tanna and Futuna-Aniwa, Vanuatu. Pacific Studies *17*, 79-91.

28. Bedford, S., and Spriggs, M. (2008). Northern Vanuatu as a Pacific Crossroads: The Archaeology of Discovery, Interaction, and the Emergence of the "Ethnographic Present". Asian Perspectives *47*, 95-120.

29. Monzier, M., Robin, C., and Eissen, J.-P. (1994). Kuwae (≈ 1425 A.D.): the forgotten caldera. Journal of Volcanology and Geothermal Research *59*, 207-218.

30. Gao, C., Robock, A., Self, S., Witter, J., Steffenson, J., Clausen, H., Siggaard-Andersen, M., Johnsen, S., Mayewski, P., and Ammann, C. (2006). The 1452 or 1453 A.D. Kuwae eruption signal derived from multiple ice core records : Greatest volcanic sulfate event of the past 700 years. Journal of Geophysical Research *111*, 1-11.

31. Flexner, J., Spriggs, M., Bedford, S., and Abong, M. (2016). Beginning Historical Archaeology in Vanuatu: Recent Projects on the Archaeology of Spanish, French, and Anglophone Colonialism. In Archaeologies of Early Modern Spanish Colonialism. (Springer International Publishing), pp. 205-227.

32. Jolly, M. (2009). The Sediment of Voyages: Re-membering Quirós, Bougainville and Cook in Vanuatu. In Oceanic Encounters: Exchange, Desire, Violence. (ANU Press), pp. 57-112.

33. Maples, B.K., Gravel, S., Kenny, E.E., and Bustamante, C.D. (2013). RFMix: a discriminative modeling approach for rapid and robust local-ancestry inference. Am J Hum Genet *93*, 278-288.

34. Goldberg, A., and Rosenberg, N.A. (2015). Beyond 2/3 and 1/3: The Complex Signatures of Sex-Biased Admixture on the X Chromosome. Genetics *201*, 263-279.

35. Vienne, B. (1984). Gens de Motlav. Idéologie et pratique sociale en Mélanésie. Publication de la Société des Océanistes *42*, 232-240.

36. Bolton, L. (1999). Women, Place and Practice in Vanuatu: a View from Ambae. Oceania *70*, 43-55.

37. Petrou, K., and Connell, J. (2017). Rural-urban migrants, translocal communities and the myth of return migration in Vanuatu: the case of Paama. Journal de la société des Océanistes *144-145*, 51-62.

38. Godelier, M. (2004). Métamorphoses de la parenté, (Paris: Fayard).

39. Parra, F.C., Amado, R.C., Lambertucci, J.R., Rocha, J., Antunes, C.M., and Pena, S.D.
(2003). Color and genomic ancestry in Brazilians. Proc Natl Acad Sci U S A *100*, 177-
182.

40. Alvarez, L., and Jaffe, K. (2004). Narcissism guides mate selection: Humans mate
assortatively, as revealed by facial resemblance, following an algorithm of "self seeking
like". Evolutionary Psychology *2*, 177–194.

41. Pugach, I., Duggan, A.T., Merriwether, D.A., Friedlaender, F.R., Friedlaender, J.S., and
Stoneking, M. (2018). The gateway from Near into Remote Oceania: new insights from
genome-wide data. Mol Biol Evol *35*, 871–886.

42. Clark, R. (1996). Linguistic Consequences of the Kuwae Eruption. In Oceanic Culture
History: Essays in Honour of Roger Green, J. Davidson, G. Irwin, F. Leach, A. Pawley
and D. Brown, eds. (New Zealand Journal of Archaeology Special Publication), pp. 275–
285.

43. Ballard, C. (2021). Transmission's end? Cataclysm and chronology in Indigenous oral
tradition. In The Routledge Companion to Global Indigenous History, A. McGrath and L.
Russell, eds. (Taylor & Francis Group), pp. 571-602.

44. Bonnemaison, J. (1996). Le tissu de nexus. In Arts of Vanuatu, J. Bonnemaison, K.
Huffman, C. Kaufmann and D. Tryon, eds. (Bathurst: Crawford House Press), pp. 176-
177.

45. Kirin, M., McQuillan, R., Franklin, C.S., Campbell, H., McKeigue, P.M., and Wilson, J.F.
(2010). Genomic runs of homozygosity record population history and consanguinity.
PLoS One *5*, e13996.

46. McQuillan, R., Leutenegger, A.L., Abdel-Rahman, R., Franklin, C.S., Pericic, M., Barac-
Lauc, L., Smolej-Narancic, N., Janicijevic, B., Polasek, O., Tenesa, A., et al. (2008). Runs
of homozygosity in European populations. Am J Hum Genet *83*, 359-372.

47. Bianco, E., Laval, G., Font-Porterias, N., Garcia-Fernandez, C., Dobon, B., Sabido-Vera,
R., Sukarova Stefanovska, E., Kucinskas, V., Makukh, H., Pamjav, H., et al. (2020).
Recent Common Origin, Reduced Population Size, and Marked Admixture Have Shaped
European Roma Genomes. Mol Biol Evol *37*, 3175-3187.

48. Ruiz-Linares, A., Adhikari, K., Acuna-Alonzo, V., Quinto-Sanchez, M., Jaramillo, C.,
Arias, W., Fuentes, M., Pizarro, M., Everardo, P., de Avila, F., et al. (2014). Admixture in
Latin America: geographic structure, phenotypic diversity and self-perception of ancestry
based on 7,342 individuals. PLoS Genet *10*, e1004572.

49. Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., and Lee, J.J. (2015).
Second-generation PLINK: rising to the challenge of larger and richer datasets.
Gigascience *4*, 7.

50. Manichaikul, A., Mychaleckyj, J.C., Rich, S.S., Daly, K., Sale, M., and Chen, W.M.
(2010). Robust relationship inference in genome-wide association studies. Bioinformatics
26, 2867-2873.

51. Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., Zhao, M.,
Chennagiri, N., Nordenfelt, S., Tandon, A., et al. (2016). The Simons Genome Diversity
Project: 300 genomes from 142 diverse populations. Nature *538*, 201-206.

52. Delaneau, O., Marchini, J., and Zagury, J.F. (2011). A linear complexity phasing method
for thousands of genomes. Nat Methods *9*, 179-181.

53. Delaneau, O., Zagury, J.F., and Marchini, J. (2013). Improved whole-chromosome
phasing for disease and population genetic studies. Nat Methods *10*, 5-6.

54. Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M., Korbel, J.O.,
Marchini, J.L., McCarthy, S., McVean, G.A., and Abecasis, G.R. (2015). A global
reference for human genetic variation. Nature *526*, 68-74.

55. Patterson, N., Price, A.L., and Reich, D. (2006). Population structure and eigenanalysis.
PLoS Genet *2*, e190.

56. Alexander, D.H., Novembre, J., and Lange, K. (2009). Fast model-based estimation of
ancestry in unrelated individuals. Genome Res *19*, 1655-1664.

57. Behr, A.A., Liu, K.Z., Liu-Fang, G., Nakka, P., and Ramachandran, S. (2016). pong: fast
analysis and visualization of latent clusters in population genetic data. Bioinformatics *32*,
2817-2823.

58. Pembleton, L.W., Cogan, N.O., and Forster, J.W. (2013). StAMPP: an R package for
calculation of genetic differentiation and structure of mixed-ploidy level populations. Mol
Ecol Resour *13*, 946-952.

59. Arauna, L.R., Mendoza-Revilla, J., Mas-Sandoval, A., Izaabel, H., Bekada, A.,
Benhamamouch, S., Fadhlaoui-Zid, K., Zalloua, P., Hellenthal, G., and Comas, D. (2017).
Recent Historical Migrations Have Shaped the Gene Pool of Arabs and Berbers in North
Africa. Mol Biol Evol *34*, 318-329.

60. Mas-Sandoval, A., Arauna, L.R., Gouveia, M.H., Barreto, M.L., Horta, B.L., Lima-Costa,
M.F., Pereira, A.C., Salzano, F.M., Hunemeier, T., Tarazona-Santos, E., et al. (2019).
Reconstructed Lost Native American Populations from Eastern Brazil Are Shaped by
Differential Je/Tupi Ancestry. Genome Biol Evol *11*, 2593-2604.

61. Chacon-Duque, J.C., Adhikari, K., Fuentes-Guajardo, M., Mendoza-Revilla, J., Acuna-
Alonzo, V., Barquera, R., Quinto-Sanchez, M., Gomez-Valdes, J., Everardo Martinez, P.,
Villamil-Ramirez, H., et al. (2018). Latin Americans show wide-spread Converso ancestry
and imprint of local Native ancestry on physical appearance. Nat Commun *9*, 5388.
62. Hellenthal, G., Busby, G.B., Band, G., Wilson, J.F., Capelli, C., Falush, D., and Myers, S.
(2014). A genetic atlas of human admixture history. Science *343*, 747-751.
63. Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Del Angel, G., Levy-
Moonshine, A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., et al. (2013). From FastQ
data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline.
Curr Protoc Bioinformatics *43*, 11 10 11-11 10 33.
64. Lucotte, E.A., Skov, L., Jensen, J.M., Macia, M.C., Munch, K., and Schierup, M.H.
(2018). Dynamic Copy Number Evolution of X- and Y-Linked Ampliconic Genes in
Human Populations. Genetics *209*, 907-920.
65. Calabrese, C., Simone, D., Diroma, M.A., Santorsola, M., Gutta, C., Gasparre, G., Picardi,
E., Pesole, G., and Attimonelli, M. (2014). MToolBox: a highly automated pipeline for
heteroplasmy annotation and prioritization analysis of human mitochondrial variants in
high-throughput sequencing. Bioinformatics *30*, 3115-3117.
66. Ralf, A., Montiel Gonzalez, D., Zhong, K., and Kayser, M. (2018). Yleaf: Software for
Human Y-Chromosomal Haplogroup Inference from Next-Generation Sequencing Data.
Mol Biol Evol *35*, 1291-1294.

1200