Enhancement of central norepinephrinergic neurotransmission contributes to

inhibition of seizure-induced respiratory arrest by targeting beta-1 adrenergic receptor

(β1-AR) locating in the cardiomyocytes in the DBA/1 mouse SUDEP model

ˏXiTing Lian[1], Qian Yu[2], HaiXiang Ma[2], LeYuan Gu, Qing Xu[1,2] HongHai Zhang*[1,2]

[1]Department of Anesthesiology, Affiliated Hangzhou First People's Hospital, Zhejiang

University School of Medicine, Hangzhou, 310006, China

[2]Department of Anesthesiology, the Fourth Clinical School of Medicine, Zhejiang

Chinese Medical University, Hangzhou, 310006, China

Corresponding authors: HongHai Zhang

Email address: zhanghonghai_0902@163.com

Abstract

Sudden unexpected death of epilepsy (SUDEP) is the leading cause of of death

in patients with epilepsy. Due to the complicated pathogenesis of SUDEP, however,

the exact mechanism of SUDEP remains elusive. Currently, although it is recognized

that the seizure-induced respiratory arrest (S-IRA) may be a main cause for SUDEP,

other factors resulting in SUDEP can't be excluded e.g arrhythmias. Our previous

findings indicated that the incidence of seizure-induced respiratory arrest S-IRA and

SUDEP evoked by acoustic stimulation or pentetrazol (PTZ) injection was

significantly reduced by atomoxetine, a norepinephrine reuptake inhibitor (NRI),

suggesting that noradrenergic neurotransmission modulates S-IRA and SUDEP. Given

that norepinephrine acts on the target to modulate respiratory and circulation function

by targeting adrenergic receptor α and beta (a-AR and β-AR) and the arrhythmias can

be contributed to SUDEP. Meanwhile, to further test whether cardiac factors are

implicated in S-IRA and SUDEP, we choose esmolol hydrochloride, a selective

antagonist of beta-1 adrenergic receptor (β1-AR) to test it in our models. Our findings

demonstrated that the lower incidence of S-IRA and SUDEP evoked by acoustic

stimulation in DBA/1 mice by administration with atomoxetine was significantly

reversed by intraperitoneal (IP) of esmolol hydrochloride. What's more, the data of

electrocardiogram (ECG) showed that the cardiac arrhythmia evoked by acoustic

stimulation including the ventricular tachycardia, ventricular premature beat and

atrioventricular block and administration of atomoxetine significantly reduced theses arrhythmias and the incidence of S-IRA and SUDEP in our models. Thus, the dysfunction of respiratory and circulation may be implicated in the pathogenesis of S-IRA and SUDEP hand in hand and enhancing central norepinephrinergic neurotransmission in the brain contributes to inhibition of seizure-induced respiratory arrest by targeting β1-AR locating in the cardiomyocytes. Our findings will show a new light on decoding the pathogenesis of SUDEP.

Keywords: sudden unexpected death in epilepsy (SUDEP); seizure-induced respiratory arrest S-IRA); esmolol hydrochloride (**Esmolol**); pentylenetetrazole (PTZ); Electrocardiogram (ECG); locus coeruleus (LC); cardiac arrhythmia

1. INTRODUCTION

It had been recognized that sudden unexpected death in epilepsy (SUDEP) is the key cause leading to the morality of epilepsy patients, especially those who had undergone antiepileptic drug resistance [1-4]. Although some important advancements concerning the pathogenesis of SUDEP had been obtained, to our certain knowledge, decoding SUDEP seems to be a huge challenge in views of the complicated mechanism of SUDEP. Indeed, seizure-induced respiratory arrest (S-IRA) is an important cause resulting in the death of epilepsy including our previous findings[2-6]. However, more and more evidence hinted that the mechanisms of SUDEP are likely to be heterogeneous. Some data showed that the dysfunction in circulation may be a contributor to SUDEP. In fact, both of the dysfunction of respiratory and circulation was in implicated in S-IRA and SUDEP. Previous findings showed that the cardiac and respiratory function were simultaneously suppressed to induce the S-IRA and SUDEP evoked by acoustic stimulation in DBA/1 mice, suggesting that the autonomic nerve including the sympathetic and parasympathetic nerve were involved in the course of S-IRA and SUDEP.

Our previous data unveiled that the incidence of S-IRA and SUDEP evoked by acoustic or PTZ can be significantly reduced by IP with atomoxetine, a norepinephrine reuptake inhibitor (NRI) [6]. Considering that the central and peripheral norepinephrine transmission was associated closely with the activity of the

sympathetic nerve. It was well known that norepinephrine transmission modulates the respiratory and circulation function by targeting the adrenergic α1, α2, β1, and β2 receptors. The findings from our and other laboratories showed α1 and α2 receptors were implicated in the S-IRA and SUDEP medicated by atomoxetine in the DBA/1 mice models [7-8]. However, whether both of β1 and β2 receptors, especially β1 receptor located in the cardiomyocytes that play a key role in modulating the cardiac function seems to essentially be explored based on our and other findings. Therefore, we choose esmolol hydrochloride, a selective antagonist of beta-1 adrenergic receptor (β1-AR) to test whether the β1-AR was implicated in the atomoxetine mediated S-IRA and SUDEP. Our findings showed that the lower incidence of S-IRA and SUDEP evoked by acoustic stimulation medicated by atomoxetine was significantly reversed by Esmolol. Thus, our data further demonstrated that the cardiac factors implicated in the reduction of the incidence of S-IRA and SUDEP by atomoxetine can't be excluded and further contribute to explaining the possibility of heterogeneous factors leading to SUDEP.

2. MATERIALS AND METHODS

2.1 Animals

All experimental procedures were the agreement with the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and approved by the Animal Advisory Committee of Zhejiang University. DBA/1 mice were housed and bred in the Animal Center of Zhejiang University School of Medicine and provided with rodent food and water ad libitum. DBA/1 mice of either sex were used in this experiment, as previous studies have shown, gender is not a variable that affects the S-IRA of DBA/1 mice. Aiming at acoustic stimulation murine model, DBA/1 mice were "primed" starting from postnatal day 26-28 to establish the consistent susceptibility to audiogenic seizures and S-IRA as previously described.

2.2 Seizure induction and resuscitation

S-IRA was established by acoustic stimulation as we previously described [5-12]. For the acoustic stimulation model, each mouse was placed in a cylindrical plexiglass chamber in a sound-isolated room, and audiogenic seizures (AGSZs) were evoked by

using an electric bell (96 dB SPL, Zhejiang People's Electronics, China). Acoustic stimulation was given for a maximum duration of 60 s or until the onset of tonic seizures and S-IRA in most mice in each group. Mice with S-IRA have resuscitated within 5 s post the final respiratory gasp by a rodent respirator (YuYan Instruments, Shanghai, China).

2.3 Effects of intraperitoneal (IP) administration of Esmolol on the atomoxetine-mediated suppression of S-IRA evoked by acoustic stimulation

The susceptibility of DBA/1 mice to S-IRA was confirmed 24 hours before atomoxetine or vehicle administration. To IP administration, atomoxetine and esmolol were dissolved in saline. Different dosages of esmolol were applied to a vehicle control group and treatment groups, in which DBA/1 mice would be acoustically stimulated later. Atomoxetine (15 mg/kg, Ca # Y0001586; Sigma-Aldrich) or vehicle was intraperitoneal (IP) injection in DBA/1 mice 120min prior to acoustic stimulation, 115min after infusion of atomoxetine (or vehicle), IP injection of esmolol (25mg/kg, 50mg/kg, H19991058; Qilu Pharmaceutical Co., Ltd.) or vehicle 5min before acoustic stimulation (n=5-7/per group). The incidence of S-IRA, latency to AGSZs, duration of wild running, clonic seizures, tonic-clonic seizures, and seizure scores were videotaped for offline analysis. DBA/1 mice from the different pre-treatment groups, the monitoring of ECG was performed before acoustic stimulation and post-S-IRA. To exclude the effects of esmolol on the incidence of the morality of DBA/1 mice, the group, subjected to confirmation of S-IRA before 24 hours to experiment, was pre-treated with the esmolol (50mg/kg, i.p) without administration of atomoxetine and acoustic stimulation in the same manner to observe the incidence of the morality of DBA/1 mice(n= 6).

2.4 Effects of IP administration of Esmolol on the atomoxetine-mediated suppression of S-IRA evoked by PTZ

The experiments were performed as follows: 1) For the control group, saline (i.p) was administered 120 min prior to and 10 min post PTZ (75 mg/kg, i.p), respectively (n=8); 2) For the Esmolol group, saline (i.p) was administered 120 min and prior to and Esmolol (50 mg/kg, i.p) was administered 10 min post PTZ (75 mg/kg, i.p),

respectively (n=7); 3) For the atomoxetine group, atomoxetine (15 mg/kg, i.p) was administered 120 min and prior to and saline (i.p) was administered 10 min post PTZ (75 mg/kg, i.p), respectively (n=5); 4) For the atomoxetine with Esmolol group, atomoxetine (15 mg/kg, i.p) was administered 120 min prior to and Esmolol (50 mg/kg, i.p) was administered 10 min post PTZ (75 mg/kg, i.p), respectively (n=5). The S-IRA latency to AGSz, duration of wild running and clonic seizures, duration tonic-clonic seizures, and seizure scores were determined by offline analysis of video recordings (n= 6-7/per group).

2.5 Data analysis

Statistical analyses were performed using SPSS 23(SPSS Software Inc., USA). The incidence of S-IRA was compared in different groups using Wilcoxon Signed Rank test, as these data are nonparametric. The data on seizure scores and the latency to AGSZs, the duration of wild running, clonic seizures, and tonic-clonic seizures were evaluated using the one-way ANOVA tests, which were presented as the mean ± SEM. Statistical significance was inferred if $p < 0.05$.

3. Results

3.1 Administration of Esmolol significantly reversed the atomoxetine-mediated suppression of S-IRA evoked by acoustic stimulation

To investigate the effects of Esmolol on the incidence of S-IRA and SUDEP medicated by atomoxetine based on our previous findings. The following experiments were performed. The incidence of S-IRA evoked by acoustic stimulation in primed DBA/1 mice was obviously reduced by atomoxetine(15 mg/kg,i.p)compared with the vehicle group(n=6, n=7, $p < 0.001$). There was no significant difference between the vehicle group and the group pre-treated with esmolol (50 mg/kg,i.p) and with vehicle (n=7,n=6, $p > 0.05$). However, the incidence of -S-IRA in the group pre-treated with atomoxetine(15 mg/kg,i.p) and esmolol(50 mg/kg,i.p) was significantly increased than the group pre-treated with atomoxetine(15 mg/kg,i.p) and the vehicle(n=6, n=7, $p < 0.05$). The incidence of -S-IRA in the group pre-treated with atomoxetine(15 mg/kg,i.p) and the vehicle was significantly decreased than in the group pre-treated with the vehicle and with Esmolol(50 mg/kg,i.p) (n=6, n=6, $p < 0.001$). The

difference of latencies to AGSZs, durations of wild running, clonic seizures, tonic-clonic seizures, and seizure scores in the treatment group showed administration of esmolol significantly reversed the atomoxetine-mediated suppression of S-IRA evoked by acoustic stimulation without sensitivity to seizures. These data showed the atomoxetine-mediated suppression of S-IRA evoked by acoustic stimulation can be significantly reversed by esmolol, which means that administration of atomoxetine reduced the incidence of S-IRA by targeting the β1-AR locating in the cardiomyocytes in our models. **(FIG 1)**

3.2 Administration of Esmolol specifically reversed the atomoxetine-mediated suppression of S-IRA by targeting the β1-AR locating in the cardiomyocytes

To exclude whether the dose of Esmolol (50 mg/kg,i.p) produces the morality of DBA/1 mice to affect the specificity to reverse the reversed atomoxetine-mediated suppression of S-IRA in our models. The following experiments were performed in different groups. There was no significant difference in the mortality of DBA/1 mice between the group pre-treated with the vehicle and the group pre-treated with Esmolol (50 mg/kg,i.p), suggesting that the atomoxetine-mediated suppression of S-IRA can be specifically reversed by esmolol by targeting the β1-AR locating in the cardiomyocytes.   **(FIG 2)**

3.3 Obvious cardiac arrhythmia immediately occurred with S-IRA evoked by acoustic stimulation

To observe the changing of ECG in the different treatment groups to analyze the role of atomoxetine and/or esmolol in cardiac electrophysiology. We observed that the ventricular tachycardia, ventricular premature beat, and atrioventricular block frequently occurred immediately following S-IRA evoked by acoustic stimulation in DBA/1 mice in a group pre-treated with the vehicle and with the vehicle. However, Compared with the vehicle group, the arrhythmia was significantly decreased in the group pre-treated with atomoxetine (15 mg/kg,i.p) and vehicle. We also observed the arrhythmia frequently occurred in the group pre-treated with esmolol and the vehicle and the group pre-treated with atomoxetine (15 mg/kg,i.p) and esmolol, suggesting that Esmolol can specifically reverse the atomoxetine-mediated suppression of S-IRA

by targeting the β1-AR locating in the cardiomyocytes without resulting in extra

arrhythmia.   **(FIG 3-4)**

3.4 Administration of Esmolol significantly reversed the atomoxetine-mediated

suppression of S-IRA evoked by PTZ

To further investigate whether administration of Esmolol significantly reversed the

atomoxetine-mediated suppression of S-IRA depends on S-IRA and SUDEP models

or not. The PTZ-induced S-IRA was accepted to test in the present study. Compared

with the vehicle group, the lower incidence of S-IRA by PTZ was reversed by

atomoxetine (p<0.01). No obvious difference in the group of vehicle and in the group

of Esmolol (p>0.05), meaning that administration of Esmolol (50 mg/kg,i.p) produced

no effects on the incidence of mortality caused by PTZ itself. However, Compared

with the group treated with atomoxetine and vehicle, the incidence of S-IRA and

SUDEP significantly increased in the group treated with atomoxetine and Esmolol

(p<0.01), suggesting that administration of Esmolol can specifically reverse the

atomoxetine-mediated suppression of S-IRA evoked by PTZ. Thus, it can be

speculated from our findings that enhancing the norepinephrinergic neurotransmission

contributes to inhibition of S-IRA via targeting the (β 1-AR) localized in the

cardiomyocytes in the DBA/1 mouse SUDEP model. **(FIG 5)**

4.DISCUSSION

There are about 70 million people in the world who suffer from epilepsy, and 2.4

million have been diagnosed with epilepsy every year, nearly 90% occurred in

developing countries[1]. The latest epidemiological studies of the American Epilepsy

Society show that the incidence of SUDEP in western developed countries is about

1.11 per 1000 people per year[2], while the results of the multi-center epidemiological

survey in China show that the incidence of SUDEP in China is about 2.3 per 1000

people per year[3], and the number of domestic epilepsy patients dying from SUDEP is

more than 20,000 per year. More importantly, SUDEP is the main factor that leads to

the death in young patients with epilepsy with an incidence two to three times that of

other diseases that can lead to sudden death in young adults, and children's SUDEP

incidence is also close to adults[4]. Japan's latest epidemiological survey shows that

the incidence of SUDEP in pregnant women is significantly higher than the incidence of non-pregnant women[5]. However, due to the complexity of its pathogenesis and the difficulty to obtain the real-time monitoring data of SUDEP in patients with epilepsy, the understanding of the SUDEP mechanism is still limited, and it is difficult to develop a prevention strategy. Many previous studies have shown that the pathophysiology of SUDEP is complex, including the disfunction of heart, autonomic nerves, respiratory, brain structural obstacles, and polygenesis.

In the current study, we firstly reported that the suppressant effects of S-IRA and SUDEP in DBA/1 mice by atomoxetine can be significantly reversed by administration of esmolol, a selective β1 receptor blocker. Our previous findings showed that administration of atomoxetine inhibited S-IRA and SUDEP in DBA/1 mice independently of seizure induction methods, and its inhibition of S-IRA induced by acoustic stimulation was dose-dependent[6]. Given the current clinical reports that atomoxetine can produce the effects of peripheral circulation. For example, after oral administration of atomoxetine, ADHD (Attention-Deficit/ Hyperactivity Disorder) patients showed symptoms such as increased heart rate and other symptoms[7], considering that atomoxetine may play a role in SUDEP passing through the central - peripheral - cardiac sympathetic mechanisms by affecting β1 receptors as dominate receptors in the heart. In the present study, we used the β1 receptor blocker esmolol to block the β1 receptor which locates preferentially in the cardiomyocytes of the heart to test whether the lower incidence of S-IRA by atomoxetine can be reversed by Esmolol. It turned out that the suppressant effects of S-IRA by atomoxetine were markedly reversed by esmolol, suggesting that administration of atomoxetine significantly reduced the incidence of S-IRA by enhancing the concentration of norepinephrine of the cleft between the sympathetic synapse and cardiomyocytes of the heart to combine with β1 receptors of cardiomyocytes. Of course, atomoxetine medicating the reduction of S-IRA by enhancing the concentration of norepinephrine to target the adrenergic receptor α1 and α2 in the brain was the main pathway as well. To be exact, atomoxetine medicates the reduction of S-IRA by interacting between α1, α2, and β1 receptors located in central and peripheral sympathetic nerves. In fact,

although S-IRA was the key cause leading to SUDEP, other factors are involved in the course can't be excluded as well. A previous study showed that S-IRA evoked by acoustic stimulation in DBA/1mice was characterized by the simultaneous suppression of respiration and circulation though the respiratory arrest preceded the cardiac arrest. What's more, the recent study showed that the tonic phase apnea evoked by acoustic stimulation is not sufficient for seizure-induced death in a W/+[Emx1-Cre] mouse. Therefore, the factors resulting in the S-IRA and SUDEP including respiration and circulation may be heterogeneous.

The previous research has shown that the main risk factors of SUDEP are generalized tonic-clonus and nocturnal seizures. The neuronal network of generalized tonic-clonic seizures may lead to hypoventilation, apnea, and cardiovascular failure by inhibiting brainstem respiration or autonomic nerve control centers, leading to death [10, 11]. Since the most sudden death cases occur at night or in sleep, with no witnesses, leaving many questions unanswered. In recent years, more and more researches have shown that apnea and heart failure are important potential mechanisms in SUDEP. Studies in animal models and SUDEP patients have shown that seizure-induced respiratory arrest (S-IRA) is one of the main induced factors that lead to death in many cases[12-14]. However, a respiratory arrest is not the only lethal cause, tonic-clonic seizures caused sudden, simultaneous respiratory and cardiac depression in mice, and the time of recovery of respiration and heart rate during resuscitation is also closely matched[15]. Therefore, the heart also plays an important role in the mechanism of SUDEP.

Seizure-associated arrhythmias are common and have been considered as the potential pathogenesis of SUDEP, such as tachycardia, bradycardia, and cardiac arrest after the seizures, resulting in cardiac repolarization, changes in electrolyte levels, blood pH, and catecholamine release in patients with epilepsy, which latter three factors may promote arrhythmias and lead to death by affecting cardiac excitability[16]. Tachycardia and several factors that increase susceptibility to tachyarrhythmias, known as pathological cardiac repolarization, are considered as risk factors for sudden cardiac death (SCD) in healthy individuals and patients with epilepsy which, in this

particular setting, are important factors in triggering SUDEP events[17, 18]. However, the exact mechanism of SUDEP and cardiac arrest induced by seizures remains unclear. Seizures may interfere with the normal functioning of the heart by causing autonomic disturbances [19], leading to fatal arrhythmias. Studies have found that cardiac electrical instability and autonomic dystonia are caused by cumulative cardiac damage during repeated attacks[20]. Studies also have shown that catecholamine (norepinephrine and adrenaline) released from seizures is related to a wide range of calcium ($Ca2+$)-mediated physiological changes, which can lead to damage to the heart structure. Our previous studies found that repeated S-IRA in DBA / 1 mice will result in necrotic damage in the ventricle which is caused by the local $Ca2+$ homeostasis disturbances, and the incidence and severity of injury depend on the total number of S-IRA[21]. Relevant clinical data have found that paroxysmal arrhythmias (including arrest, atrioventricular block, and rare atrial and ventricular fibrillation) usually occur after convulsive episodes and are often associated with (near)SUDEP[22]. Studies have shown that chronic epilepsy can lead to hypoxemia and the increase of catecholamine, causing damage of the heart and vascular, resulting in cardiac electrical, and mechanical dysfunction[23,24]. Some fatal incentives may be the acidosis caused by persistent hypoxemia and hypercapnia, leading to bradycardia or cardiac arrest, while in others it may be the seizures due to malignant arrhythmias[25]. This suggests that SUDEP is inextricably linked to arrhythmia.

In the present study, the ECG of DBA/ 1 mice with administration of esmolol (50 mg/kg, i.p) that can significantly reverse the lower incidence of S-IRA evoked by acoustic stimulation by administration of atomoxetine showed the mixture of sinus bradycardia, atrioventricular block, ventricular premature beat, and ventricular tachycardia in order. In the meantime, the ECG of DBA/1 mice suffered from the S-IRA evoked by acoustic stimulation characterized by the mixture of sinus bradycardia, atrioventricular block, ventricular premature beat, and ventricular tachycardia as well in our model. However, no obvious arrhythmia appeared in the DBA/1 mice without suffering from S-IRA other than the sinus bradycardia and no apparent mortality occurred in the group of DBA/1 mice with pre-treatment with the

dose of esmolol 50 mg/kg (i.p). Furthermore, the arrhythmias can be significantly reduced by atomoxetine in our models. Thus, the lower incidence of S-IRA by acoustic stimulation can be significantly as well as especially reversed by esmolol by blocking the β1-AR located in cardiomyocytes to combine the norepinephrine released by cardiac sympathetic nerve synaptic terminal in our models. Additionally, it is possible that the locus coeruleus (LC), as the largest nuclei to synthesize and release norepinephrine in the brain, may be implicated in the course of administration of esmolol to reverse the lower incidence of S-IRA and SUDEP and plays a key role in modulating S-IRA and SUDEP by regulating the respiratory and circulation function in our models. In future experiments, we will further explore it.

4. Conclusions

The current finding in our models suggested that enhancing central norepinephrinergic neurotransmission in the brain contributes to inhibition of seizure-induced respiratory arrest by targeting β1-AR locating in the cardiomyocytes and the norepinephrinergic neurotransmission-β1-AR between the pathway brain and heart may be a potential and specific target to prevent SUDEP.

5. Funding

The work was supported by the National Natural Science Foundation of China (Grant.NO: 81771403 and 81974205); by the Natural Science Foundation of Zhejiang Province (LZ20H090001); by the Program of New Century 131 outstanding young talent plan top-level of Hang Zhou to HHZ

6. DISCLOSURE

All authors declare no competing interests. We had confirmed that we have read the Journal's position on issues involved in ethical publication and affirm that this study was in accordance with those guidelines.

7. Figure legends

Figure 1

Figure 1 Administration of Esmolol significantly reversed the atomoxetine-mediated suppression of S-IRA evoked by acoustic stimulation

A. The protocol to explore the influence of administration of esmolol on atomoxetine-mediated reduction of S-IRA evoked by acoustic stimulation. **B.** Compared to the group treated with vehicle (n = 7) or vehicle and esmolol (n = 6), S-IRA evoked by acoustic stimulation was markedly lower in groups treated with i.p. atomoxetine (n = 6, **p < 0.01) at 15mg/Kg. However, the protective effect of atomoxetine in primed DBA/1 mice was significantly reversed by esmolol doses of 50mg/kg (n = 7, *p < 0.05). **C.** Furthermore, compared with the group treated with atomoxetine and vehicle (n = 6), the latencies to AGSZs in the control group (n = 7)

or the group treated with atomoxetine and esmolol (n = 7) was significantly increased (p < 0.05). **D-E.** There were no intergroup differences in durations of wild running plus clonic seizures (W+C), tonic-clonic seizures (p > 0.05). **F.** Compared to the control group (n = 7) , the seizure score was lower in the group treated with atomoxetine and vehicle (n = 6, *p < 0.05).

Figure 2

Figure 2 The dose of Esmolol (50 mg/kg, i.p.) does not produce the mortality of DBA/1 mice.

A. The protocol to explore the dose of esmolol (50 mg/kg,i.p.) to varify whether produce the morality of DBA/1 mice. **B.** There was no significant difference in the

mortality of DBA/1 mice between the group pre-treated with the vehicle and the

group pre-treated with esmolol(50 mg/kg, i.p, n=7) (p > 0.05). **C.** Compared with the

vehicle group, the heart rate of DBA/1 in the group pre-treated with esmolol significantly

increased (p < 0.001). However, no ventricular tachycardia, atrioventricular block to be

found in the DBA/1 mice from the vehicle group and the group pre-treated with

esmolol .

Figure 3

Figure 3 After administration of atomoxetine, obvious cardiac arrhythmia immediately occurred with S-IRA evoked by acoustic stimulation

A. The protocol to explore the changes of electrocardiogram on atomoxetine-mediated reduction of S-IRA evoked by acoustic stimulation. **B.** The ventricular tachycardia and atrioventricular block occurred immediately following S-IRA evoked by acoustic stimulation in DBA/1 mice in the control group. **C.** What's more, the group pre-treated with the atomoxetine(15mg/Kg) was observed the ventricular tachycardia, atrioventricular block, and ventricular premature beat after acoustic stimulation, and the DBA/1 mice were then recovered.

Figure 4

A

-10 min   0 min   115 min   120 min   130 min

ECG Recording   Vehicle / Atomoxetine i.p.   Esmolol i.p.   Acoustic Stimulation   ECG Recording

10 min ECG Recording   10 min ECG Recording

B   **Vehicle+Esmolol**   **C**   **Atomoxetine+Esmolol**

Normal   Normal

Acoustic Stimulation

Post-S-IRA

Ventricular Tachycardia (VT)   **Ventricular Tachycardia (VT)**

I Atrioventricular Block (I AVB)   **I Atrioventricular Block (I AVB)**   Post-S-IRA

II Atrioventricular Block (II AVB)   **II Atrioventricular Block (II AVB)**

III Atrioventricular Block (III AVB)   **III Atrioventricular Block (III AVB)**

Death   10mm/mv   10mm/mv   Death

25mm/s   25mm/s

3 **Figure 4 After administration of esmolol, obvious cardiac arrhythmia**
4 **immediately occurred with S-IRA evoked by acoustic stimulation**

5 A. The protocol to explore the changes of electrocardiogram on the administration of

6 esmolol reversed the atomoxetine-mediated reduction of S-IRA evoked by acoustic

7 stimulation. **B.** The ventricular tachycardia and atrioventricular block occurred

immediately following S-IRA evoked by acoustic stimulation in DBA/1 mice in the group treated with vehicle and esmolol(50mg/Kg). **C.** The group pre-treated with the atomoxetine(15mg/Kg) and treated with esmolol(50mg/Kg) was observed the ventricular tachycardia and atrioventricular block after acoustic stimulation.

Figure 5

Fig 5 Administration of Esmolo significantly reversed the atomoxetine-mediated suppression of S-IRA evoked by PTZ

B. Compared with the group treated with vehicle (n = 5) or vehicle and esmolol (n =5), S-IRA evoked by acoustic stimulation was markedly lower in groups treated with i.p. atomoxetine (n = 7, **p < 0.01). However, Compared with the group treated with

i.p. atomoxetine and vehicle, the incidence of S-IRA significantly increased in the group treated with i.p. atomoxetine and esmolol (n = 7, **p < 0.05). C.No significant difference of GSZ latency(s) and W+C duration (s) in the different treatment group (n = 6-7/group, p > 0.05). However, compared with the vehicle and other groups, duration of Tonic-clonics (s) was significantly decreased in the group with the pre-treatment (atomoxetine 15mg/kg + vehicle). Compared with the vehicle and other groups, seizure score was significantly decreased in the group with the pre-treatment (atomoxetine 15mg/kg + vehicle).

8. References

[1]   SINGH A, TREVICK S. The Epidemiology of Global Epilepsy [J]. Neurol Clin, 2016, 34(4): 837-47.

[2]   LHATOO S, NOEBELS J, WHITTEMORE V, et al. Sudden unexpected death in epilepsy: Identifying risk and preventing mortality [J]. Epilepsia, 2015, 56(11): 1700-6.

[3]   GE Y, DING D, ZHANG Q, et al. Incidence of sudden unexpected death in epilepsy in community-based cohort in China [J]. Epilepsy Behav, 2017, 76: 76-83.

[4]   KELLER A E, WHITNEY R, LI S A, et al. Incidence of sudden unexpected death in epilepsy in children is similar to adults [J]. Neurology, 2018, 91(2): e107-e11.

[5]   TANAKA H, KATSURAGI S, HASEGAWA J, et al. Maternal Death Related to Sudden Unexpected Death in Epilepsy: A Nationwide Survey in Japan [J]. Brain Sci, 2021, 11(8).

[6]   ZHANG H, ZHAO H, FENG H J. Atomoxetine, a norepinephrine reuptake inhibitor, reduces seizure-induced respiratory arrest [J]. Epilepsy Behav, 2017, 73: 6-9.

[7]   Yue Shen , Hai Xiang Ma , Han Lu 3, Hai Ting Zhao , Jian Liang Sun, Yuan Cheng , Hong Hai Zhang. Central deficiency of norepinephrine synthesis and norepinephrinergic neurotransmission contributes to seizure-induced respiratory arrest.Biomed Pharmacother. 2021:111024. doi:

10.1016/j.biopha.2020.111024.

[8]Rui Zhang, Zheren Tan, Jianguo Niu Hua-Jun Feng. Adrenergic α2 receptors are implicated in seizure-induced respiratory arrest in DBA/1 mice. Life Sci. 2021;284:119912. doi: 10.1016/j.lfs.2021.119912

[9]   HENNISSEN L, BAKKER M J, BANASCHEWSKI T, et al. Cardiovascular Effects of Stimulant and Non-Stimulant Medication for Children and Adolescents with ADHD: A Systematic Review and Meta-Analysis of Trials of Methylphenidate, Amphetamines and Atomoxetine [J]. CNS Drugs, 2017, 31(3): 199-215.

[10]   ALLEN L A, HARPER R M, KUMAR R, et al. Dysfunctional Brain Networking among Autonomic Regulatory Structures in Temporal Lobe Epilepsy Patients at High Risk of Sudden Unexpected Death in Epilepsy [J]. Front Neurol, 2017, 8: 544.

[11]   MUELLER S G, NEI M, BATEMAN L M, et al. Brainstem network disruption: A pathway to sudden unexplained death in epilepsy? [J]. Hum Brain Mapp, 2018, 39(12): 4820-30.

[12]   FAINGOLD C L, RANDALL M, TUPAL S. DBA/1 mice exhibit chronic susceptibility to audiogenic seizures followed by sudden death associated with respiratory arrest [J]. Epilepsy Behav, 2010, 17(4): 436-40.

[13]   RYVLIN P, NASHEF L, LHATOO S D, et al. Incidence and mechanisms of cardiorespiratory arrests in epilepsy monitoring units (MORTEMUS): a retrospective study [J]. Lancet Neurol, 2013, 12(10): 966-77.

[14]   BUCHANAN G F, MURRAY N M, HAJEK M A, et al. Serotonin neurones have anti-convulsant effects and reduce seizure-induced mortality [J]. J Physiol, 2014, 592(19): 4395-410.

[15]   SCHILLING W P, MCGRATH M K, YANG T, et al. Simultaneous cardiac and respiratory inhibition during seizure precedes death in the DBA/1 audiogenic mouse model of SUDEP [J]. PLoS One, 2019, 14(10): e0223468.

[16]   DEGIORGIO C M, CURTIS A, HERTLING D, et al. Sudden unexpected death in epilepsy: Risk factors, biomarkers, and prevention [J]. Acta Neurol

Scand, 2019, 139(3): 220-30.

[17] SURGES R, TAGGART P, SANDER J W, et al. Too long or too short? New insights into abnormal cardiac repolarization in people with chronic epilepsy and its potential role in sudden unexpected death [J]. Epilepsia, 2010, 51(5): 738-44.

[18] CONE E J, DICKERSON S L, DARWIN W D, et al. Elevated drug saliva levels suggest a "depot-like" effect in subjects treated with sublingual buprenorphine [J]. NIDA Res Monogr, 1990, 105: 569.

[19] SEVCENCU C, STRUIJK J J. Autonomic alterations and cardiac changes in epilepsy [J]. Epilepsia, 2010, 51(5): 725-37.

[20] PANG T D, NEARING B D, KRISHNAMURTHY K B, et al. Cardiac electrical instability in newly diagnosed/chronic epilepsy tracked by Holter and ECG patch [J]. Neurology, 2019, 93(10): 450-8.

[21] ZHAO H, ZHANG H, SCHOEN F J, et al. Repeated generalized seizures can produce calcified cardiac lesions in DBA/1 mice [J]. Epilepsy Behav, 2019, 95: 169-74.

[22] VAN DER LENDE M, SURGES R, SANDER J W, et al. Cardiac arrhythmias during or after epileptic seizures [J]. J Neurol Neurosurg Psychiatry, 2016, 87(1): 69-74.

[23] POWELL K L, LIU Z, CURL C L, et al. Altered cardiac structure and function is related to seizure frequency in a rat model of chronic acquired temporal lobe epilepsy [J]. Neurobiol Dis, 2021, 159: 105505.

[24] VERRIER R L, PANG T D, NEARING B D, et al. Epileptic heart: A clinical syndromic approach [J]. Epilepsia, 2021, 62(8): 1780-9.

[25] GOLDMAN A M. Mechanisms of sudden unexplained death in epilepsy [J]. Curr Opin Neurol, 2015, 28(2): 166-74.