A non-canonical RNA-binding domain of the Fragile X protein, FMRP, elicits translational repression independent of mRNA G-quadruplexes

MaKenzie R. Scarpitti[1,2,3], Julia E. Warrick[1,2,3], Evelyn L. Yoder[2,3], and Michael G. Kearse[1,2,3*]

[1]The Biomedical Sciences Graduate Program, [2]Department of Biological Chemistry and Pharmacology, [3]Center for RNA Biology, The Ohio State University, Columbus, OH 43210 USA

*To whom correspondence should be addressed. Tel: +1 614 685 3709; Email: michael.kearse@osumc.edu

ABSTRACT

Loss of functional fragile X mental retardation protein (FMRP) causes fragile X syndrome, the leading form of inherited intellectual disability and the most common monogenic cause of autism spectrum disorders. FMRP is an RNA-binding protein that controls neuronal mRNA localization and translation. FMRP is thought to inhibit translation elongation after being recruited to target transcripts via binding RNA G-quadruplexes (G4s) within the coding sequence. Here, we directly test this model and report that FMRP inhibits translation independent of mRNA G4s. Furthermore, we found that the RGG box motif together with its natural C-terminal domain forms a non-canonical RNA-binding domain (ncRBD) that is essential for translational repression. The ncRBD elicits broad RNA binding ability and binds to multiple reporter mRNAs and all four homopolymeric RNAs. Serial deletion analysis of the ncRBD identified that the regions required for mRNA-binding and translational repression overlap but are not identical. Consistent with FMRP stalling elongating ribosomes and causing the accumulation of slowed 80S ribosomes, transcripts bound by FMRP via the ncRBD co-sediment with heavier polysomes and were present in puromycin-resistant ribosome complexes. Together, this work identifies a ncRBD and translational repression domain that shifts our understanding of how FMRP inhibits translation independent of mRNA G4s.

INTRODUCTION

Loss of functional fragile X mental retardation protein (FMRP) causes fragile X syndrome (FXS) (1-4), the leading form of inherited intellectual disability and the most common monogenic cause of autism spectrum disorders (ASD) (5). FXS affects around one in 4,000 males and one in 8,000 females (5). Approximately one third of FXS patients are also diagnosed with ASD (6). The vast majority of FXS cases is caused by a CGG trinucleotide repeat expansion in the 5□ untranslated region (UTR) of the *FMR1* gene. The expanded repeat is subsequently hypermethylated which causes transcriptional silencing of the locus (1-4). As a result, most FXS patients express little to no functional FMRP.

FMRP is an RNA-binding protein (RBP) with multiple RNA-binding domains (RBDs) including three K homology (KH) domains (KH0, KH1, and KH2) and a positively-charged RGG box motif (7-9). In general, KH domains canonically bind to short stretches of RNA to provide binding specificity to particular transcripts (10-14). However, the KH domains of FMRP are unable to bind strongly to any combination of five ribonucleotides (15). *In vitro* selection experiments have found that the KH2 domain of FMRP does have high affinity for an artificial RNA pseudoknot (e.g., Δ kissing complex 2; Δkc2) (16). Reports using *in vitro* selection, filter binding, and fluorescence anisotropy have concluded that the RGG box motif of FMRP has preference for RNA G-quadruplexes (G4s), and in particular an artificial RNA molecule called Sc1 that harbors a G4 (15,17). Using these multiple binding domains and motifs, FMRP is thought to bind mRNAs with higher order RNA structure (e.g., pseudoknots and G4s). The importance of the KH domains and RGG box motif in FMRP function is highlighted by independent point mutations found in rare FXS cases. Enigmatically, a G266E mutation in the KH1 domain (18), an I304N mutation in the KH2 domain (19), and a guanine insertion that causes a frameshift early within the RGG box motif (20) all cause FXS. This suggests that FMRP may have multiple functions within neurons dictated by interactions between specific RBDs and mRNA sequences or structures.

FMRP is known to regulate translation and, in alignment with this idea, it is found to primarily bind the coding sequence (CDS) of mRNAs in mouse brain tissue (21). Most previous reports (21-25), but not all (26-28), suggest that FMRP represses translation of its target mRNAs by stalling bound elongating ribosomes. Thus, at least one facet of the FXS phenotype is thought to result from aberrant and unregulated protein synthesis of dendritic mRNAs at synapses. Because FMRP mutants that lack the RGG box motif do not repress translation (22), it is commonly postulated that FMRP binds to target mRNAs at RNA G4s and subsequently blocks translation elongation. In this model, the G4-bound FMRP acts as a physical roadblock and directly contacts the elongating ribosome near the A site to sterically hinder delivery of aminoacyl-tRNAs to the ribosome.

Here we directly test the ability of FMRP to inhibit translation of G4-lacking and G4-containing mRNAs. Our data show that FMRP represses translation independent of mRNA G4s and that the RGG box works synergistically with the positively-charged C-terminal domain (CTD) to form a non-canonical RNA-binding domain (ncRBD) that is critical for repression. The RGG+CTD ncRBD is able to bind all four homopolymeric RNA sequences with slightly less preference for homopolymeric A RNA, providing FMRP the ability to target multiple mRNAs. Through scanning deletion analysis of the ncRBD, we found that the residues required for mRNA binding and translational repression overlap but are not identical. Consistent with inhibiting translation post-initiation, our data show that FMRP harboring the ncRBD, but not a mutant that is missing the ncRBD, promotes accumulation of inhibited reporter mRNA on heavy polysomes and forms puromycin-resistant ribosome complexes. Taken together, our data indicate that FMRP harbors a small RNA-binding element that extends from the annotated RGG

box motif that overlaps with a translational repression domain to stall and slow elongating ribosomes independent of mRNA G4s.

MATERIALS AND METHODS
Plasmids

The nLuc coding sequence from pNL1.1 (Promega) was analyzed by Quadruplex forming G-Rich Sequences (QGRS) Mapper (https://bioinformatics.ramapo.edu/QGRS/index.php) and manually codon optimized to eliminate predicted G4 motifs. The final nLuc coding sequence harboring the P2A ribosome skipping motif and human β-globin 5□ UTR was synthesized by Integrated DNA Technologies and cloned into pcDNA3.1(+). The G4 sequence and larger 5□ UTR (three human β-globin 5□ UTR sequences in tandem) were inserted using the Q5 Site-Directed Mutagenesis Kit (NEB # E0552S). pcDNA3.1(+)/mEGFP was a kind gift from Jeremy Wilusz (Baylor College of Medicine). pCRII/FFLuc, which contains the FFLuc coding sequence from pGL4.13 (Promega) downstream from the T7 RNA polymerase promoter, was previously described (29).

An *E. coli* optimized coding sequence for human FMRP (isoform 1) was designed and synthesized by Genscript, and then subcloned into pET His6 MBP TEV LIC cloning vector (1M), which was a gift from Scott Gradia (Addgene plasmid # 29656), through ligation-independent cloning (LIC) using Novagen's LIC-qualified T4 DNA polymerase (Sigma # 70099-M) as described by Q3 Macrolab (http://qb3.berkeley.edu/macrolab/). The His6-tag was deleted from the N-terminus and inserted at the C-terminus. The NT-hFMRP sequence included a P451S mutation to prevent ribosome stalling at a poly-proline stretch and formation of truncated recombinant protein, as previously described (30). Point mutations and deletions were achieved using the Q5 Site-Directed Mutagenesis Kit. To be as consistent as possible across the previous literature, we refer to the RGG box motif as to the minimal region identified by Darnell and colleagues that bound Sc1 RNA with highest affinity (17).

All plasmids were propagated in TOP10 *E. coli* (Thermo Fisher # C404006), purified using the PureYield Plasmid Miniprep or Midiprep Systems (Promega # A1222 and A2495), and validated by Sanger sequencing at The Ohio State University Comprehensive Cancer Center Genomics Shared Resource (OSUCCC GSR). Nucleotide sequences of the reporters and recombinant proteins are provided in the Supplemental Material.

Reporter mRNA *in vitro* transcription

All nLuc plasmids were linearized with XbaI and purified using a Zymo DNA Clean & Concentrator 25 (Zymo Research # D4065). pcDNA3.1(+)/mEGFP was linearized with PspOMI. pCRII/FFLuc was linearized with HindIII. DNA was transcribed into mRNA which was co-transcriptionally capped with the Anti-Reverse Cap Analog (ARCA) 3′-O-Me-m7G(5′)ppp(5′)G (NEB # S1411) using the HiScribe T7 High Yield RNA Synthesis Kit (NEB # E2040). Our standard 10 μL reactions used 0.5 μg of linear plasmid template and an 8:1 ARCA:GTP ratio. Reactions were incubated at 30°C for 2 hrs, then incubated with 20 U of DNaseI (NEB # M0303S) at 37°C for 15 min, and then purified using a Zymo RNA Clean & Concentrator 25 (Zymo Research # R1018). Reporter mRNA was eluted in 75 μL RNase-free water, aliquoted in single use volumes, and stored at -80°C. Reporter mRNA integrity was confirmed by denaturing formaldehyde agarose gel electrophoresis and ethidium bromide visualization. We routinely found the 30°C incubation resulted in less observable truncated products than incubation at 37°C and did not significantly affect yield for our purposes.

Recombinant protein expression and purification

All recombinant proteins were expressed in Rosetta 2(DE3) *E. coli* (Sigma # 71397-4) using MagicMedia *E. coli* Expression Medium (Thermo Fisher # K6803) supplemented with 50 µg/mL kanamycin and 35 µg/mL chloramphenicol for auto-induction. A 5 mL starter culture in LB media supplemented with 50 µg/mL kanamycin, 35 µg/mL chloramphenicol, and 1% glucose (w/v) was inoculated with a single colony and grown overnight at 37°C, 250 rpm. 1 mL of a fresh overnight starter culture was then used to inoculate 50 mL of room temperature MagicMedia and incubated for 48 hrs at 18°C, 160 rpm in a 250 mL baffled flask. After auto-induction, cultures were pelleted and stored at -20°C for purification later. Recombinant proteins were purified using a dual affinity approach, first using the C-terminal His6-tag, then the N-terminal MBP-tag. Cell pellets were resuspended and lysed with BugBuster Master Mix (Sigma # 71456) using the recommended 5 mL per 1 g wet cell pellet ratio for 10 min at room temperature with gentle end-over-end rotation (10-15 rpm). Lysates were placed on ice and kept cold moving forward. Lysates were cleared by centrifugation for 20 min at 18,000 rcf in a chilled centrifuge (4°C). Lysates were then incubated with HisPur Cobalt Resin (Thermo Fisher # 89965) in a Peirce centrifugation column (Thermo # 89897) for 30 min at 4°C with gentle end-over-end rotation. Columns were centrifuged in a pre-chilled (4°C) Eppendorf 5810R for 2 min at 700 rcf to eliminate the flow through and then were washed 5X with two resin-bed volumes of ice-cold Cobalt IMAC Wash Buffer (50 mM Na_3PO_4, 300 mM NaCl, 10 mM imidazole; pH 7.4) in a pre-chilled (4°C) Eppendorf 5810R for 2 min at 700 rcf. His-tagged proteins were then eluted in a single elution step with two resin-bed volumes of ice-cold Cobalt IMAC Elution Buffer (50 mM Na_3PO_4, 300 mM NaCl, 150 mM imidazole; pH 7.4) by gravity flow. Eluates were then incubated with Amylose resin (NEB # E8021) in a centrifugation column for 2 hrs at 4°C with gentle end-over-end rotation. Columns were washed 5X with at least two bed-volumes of ice-cold MBP Wash Buffer (20 mM Tris-HCl, 200 mM NaCl, 1 mM EDTA; pH 7.4) by gravity flow. MBP-tagged proteins were then eluted by a single elution step with two resin-bed volumes of ice-cold MBP Elution Buffer (20 mM Tris-HCl, 200 mM NaCl, 1 mM EDTA, 10 mM maltose; pH 7.4) by gravity flow. Recombinant proteins were then desalted and buffer exchanged into Protein Storage Buffer (25 mM Tris-HCl, 125 mM KCl, 10% glycerol; pH 7.4) using a 7K MWCO Zeba Spin Desalting Column (Thermo Fisher # 89892) and, if needed, concentrated using 10K MWCO Amicon Ultra-4 (EMD Millipore # UFC803024). Recombinant protein concentration was determined by Pierce Detergent Compatible Bradford Assay Kit (Thermo Fisher # 23246) with BSA standards diluted in Protein Storage Buffer before aliquoting in single use volumes, snap freezing in liquid nitrogen, and storage at -80°C.

mRNA folding, mRNP formation, and *in vitro* translation

In vitro translation was performed in the dynamic linear range as previously described but adapted to translate mRNPs (29). 28 nM *in vitro* transcribed reporter mRNA in RNA folding buffer (10 mM Tris-HCl, 100 mM KCl; pH 7.4) was heated for 5 min at 70°C then gradually cooled for 30 min at room temperature on the bench. 5 mM $Mg(OAc)_2$ (final) was then added, gently mixed, and allowed to cool for an additional 30 min on the bench. In a total of 4 µL, 28 fmol of folded reporter mRNA was mixed with 0-10 picomol of recombinant protein and 100 picomol of UltraPure BSA (Thermo Fisher # AM2618) on ice for 1 hr. UltraPure BSA stock was diluted in protein storage buffer and its addition was necessary to prevent non-specific binding of the reporter mRNA to the tube. For *in vitro* translation reactions, 6 µL of a Rabbit Reticulocyte Lysate (RRL) master mix was added to each 4µL mRNP complex. 10 µL *in vitro* translation reactions were performed in the linear range using 2.8 nM mRNA in the Flexi RRL System (Promega # L4540) with final concentrations of reagents at 30% RRL, 10 mM amino acid mix minus leucine, 10 mM amino acid mix minus Methionine, 0.5 mM $Mg(OAc)_2$, 100 mM KCl, 8 U murine RNase inhibitor (NEB # M0314), 0-1 µM recombinant protein, and 10 µM UltraPure BSA. Reactions were incubated for 30 min at 30°C, terminated by incubation on ice and diluted 1:5 in Glo Lysis Buffer (Promega # E2661). 25 µL of prepared Nano-Glo reagent (Promega # N1120)

was mixed with 25 µL of diluted reaction and incubated at room temperature for 5 min in the dark (with gentle shaking during the first minute), and then read on a Promega GloMax Discover Multimode Microplate Reader.

IC$_{50}$ measurements were calculated using non-liner regression analysis following the [inhibitor] versus normalized response analysis in GraphPad Prism 9.1.2

Denaturing PAGE, native PAGE, and nucleic acid staining
Denaturing TBE-Urea 6% PAGE gels (Thermo Fisher # EC68652BOX) were run with 1X TBE-Urea Sample Buffer (Thermo Fisher # LC6876) and 1X TBE Running Buffer (Thermo Fisher # LC6675). Gels were pre-run at 180 volts (constant) for 20 minutes, then samples were loaded and run at 180 volts (constant) for 3 hrs. Native TBE 6% PAGE gels (Thermo Fisher # EC62652BOX) were run with 1X Hi-Density TBE Sample Buffer (Thermo Fisher # LC6678) and 1X TBE Running Buffer. Gels were pre-run at 180 volts (constant) for 1 hr, then samples were loaded and run at 180 volts (constant) for 3 hrs. Total RNA was stained with 1X SYBR Green II RNA Gel Stain (Thermo Fisher # S7568) diluted in milliQ water for 10 min in the dark. G4s were selectively stained with 0.1 mg/mL N-methyl mesoporphyrin IX (NMM) (Frontier Scientific # NMM58025MG) in milliQ water for 10 min in the dark. Stained gels were imaged on a Bio-Rad GelDoc Go Gel Imaging System using the SYBR Green setting. NMM stock was made at 10 mg/mL in DMF and stored in single use aliquots at -20°C. For Native TBE PAGE, the mRNA was folded as described above. Due to the different sensitivity, 100 ng and 2,000 ng were loaded for staining with SYBR Green II and NMM, respectively.

Electrophoretic mobility shift assays
For EMSAs with mRNAs to assess mRNP formation, 4% PAGE gels made with 0.5X TBM (45 mM Tris, 45 mM borate, 2.5 mM MgCl$_2$) and acrylamide/bis-acrylamide, 37.5:1 (2.7% crosslinker) were poured between glass plates and allowed to polymerize for at least 1 hr. Gels were pre-run for 20 min at 100 volts (constant) with 0.5X TBM as the running buffer. In 18 µL, 1 picomol of folded reporter mRNA was mixed with 20 picomol of recombinant protein and 200 picomol of UltraPure BSA (Thermo Fisher # AM2618) in binding buffer (10 mM Tris-HCl, 100 mM KCl, 5 mM Mg(OAc)$_2$; pH 7.4) on ice for 1 hr. 2 µL of 20% Ficoll 400 (Sigma # F5415-25ML) was then added and the entire sample was loaded immediately. After loading, gels were run for 45 min at 100 volts (constant) at room temperature, stained with 1X SYBR Green II RNA Gel Stain, and visualized as described above.

For EMSAs with 5' FAM-labeled RNA oligos, 4% PAGE gels were made as described above with 0.5X TBM and acrylamide/bis-acrylamide, 37.5:1. 5' FAM-labeled RNA olgios were first diluted in nuclease-free water and not heated. In 18 µL, 4.8 picomol of 5' FAM-labeled RNA oligo was mixed with 54 picomol of recombinant protein and 542 picomol of UltraPure BSA in binding buffer (10 mM Tris-HCl, 100 mM KCl, 5 mM Mg(OAc)$_2$; pH 7.4) on ice for 1 hr in the dark. 2 µL of 20% Ficoll 400 was then added, and the entire sample was loaded immediately. For the FAM-U(G)17 RNA oligo, 24 picomol per reaction was used due to FAM being slightly quenched by proximal guanosines (a known caution provided by the RNA oligo manufacture). Gels were pre-run for 20 min at 100 volts (constant) with 0.5X TBM as the running buffer. Samples were loaded and gels were run for 40 min at 100 volts (constant) at room temperature in the dark. Gels in glass plates were then directly imaged using an Azure Sapphire Biomolecular Imager. The 5☐ FAM-labeled RNA oligo sequences are provided in the Supplemental Data. A single uridine was added as a spacer between the 5' FAM label and the polymeric guanosines to avoid quenching and was kept for consistency in the other labeled polymeric RNA oligos. We found flexible linkers did not help further avoid quenching by proximal guanosines.

Sucrose gradient ultracentrifugation and RT-qPCR

In vitro translation reactions were scaled up 10-fold to 100 µL but were performed identically as described above. After 30 min at 30°C, reactions were kept on ice, diluted two-fold with ice-cold Polysome Dilution Buffer (10 mM Tris-HCl, 140 mM KCl, 10 mM MgCl$_2$, 1 mM DTT, 100 µg/mL cycloheximide; pH 7.4), and layered on top of a linear 10-50% (w/v) buffered sucrose gradient (10 mM Tris-HCl, 140 mM KCl, 10 mM MgCl$_2$, 1 mM DTT, 100 µg/mL cycloheximide; pH 7.4) in a 14 mm × 89 mm thin-wall Ultra-Clear tube (Beckman # 344059) that was formed using a Biocomp Gradient Master. Gradients were centrifuged at 35K rpm for 120 min at 4°C in a SW-41Ti rotor (Beckman) with maximum acceleration and no brake using a Beckman Optima L-90 Ultracentrifuge. Gradients were subsequently fractionated into 0.9 mL volumes using a Biocomp piston fractionator with a TRIAX flow cell (Biocomp) recording a continuous A$_{260\ nm}$ trace. Total RNA was extracted from 400 µL of each fraction (spiked with 0.2 ng exogenous control FFLuc mRNA; Promega # L4561) by adding 600 µL TRIzol (Thermo Fisher # 15596018) and following the manufacturer's protocol. Glycogen (Thermo Fisher # R0561) was added at the isopropanol precipitation step. The resulting RNA pellet was resuspended in 30 µL nuclease-free water. 16 µL of extracted RNA was converted to cDNA using iScript Reverse Transcription Supermix for RT-qPCR (Bio-Rad # 1708841). cDNA reactions were then diluted 10-fold with nuclease-free water and stored at -20°C or used immediately. RT-qPCR was performed in 15 µL reactions using iTaq Universal SYBR Green Supermix (Bio-Rad # 1725124) in a Bio-Rad CFX Connect Real-Time PCR Detection System with 1.5 µL diluted cDNA and 250 nM (final concentration) primers. For each fraction, nLuc reporter mRNA abundance was normalized to the spiked-in control FFLuc mRNA using the Bio-Rad CFX Maestro software (ΔΔCt method). Abundance of total signal in each fraction was calculated using $Q_n = 2^{\Delta\Delta Ct}$ and $P = 100 \times Q_n/Q_{total}$ as previously described (31). Primers for RT-qPCR can be found in **Supplemental Table S1**.

Puromycin treatment and low-speed sucrose cushions

nLuc reporter mRNA translation was performed as described above except that translation was limited to 15 min at 30°C. Samples were then placed on ice for 3 min before the addition of 0.1 mM puromycin (final) and further incubation at 30°C for 30 min. Control samples lacking puromycin (water added instead) were kept on ice. Cycloheximide (final concentration of 1.43 mg/mL) was then added to all samples to preserve ribosome complexes on mRNAs and halt puromycin incorporation. In a separate tube, Firefly luciferase (FFLuc) reporter mRNA was translated as described above (3 nM mRNA conditions) for 15 min at 30°C and was terminated by the addition of 1.43 mg/mL cycloheximide (final) and incubation on ice.

nLuc and FFLuc translation reactions were combined on ice and then mixed with an equal volume of ice-cold 2X dilution buffer (40 mM Tris-HCl, 280 mM KCl, 20 mM MgCl$_2$, 200 µg/ml cycloheximide, 2 mM DTT; pH 7.4). The entire 40 µl volume was then layered on top of 130 µl of ice-cold 35% (w/v) buffered sucrose (20 mM Tris-HCl, 140 mM KCl, 10 mM MgCl$_2$, 100 µg/mL cycloheximide, 1 mM DTT; pH 7.4) in a pre-chilled 7 mm x 20 mm thick-walled polycarbonate ultracentrifuge tubes (Thermo Scientific # 45233) and centrifuged in a S100AT3 rotor at 4°C for 60 min at 50,000 x g (43,000 rpm) in a Sorvall Discovery M120 SE Micro-Ultracentrifuge. The supernatant was then discarded and each pellet was resuspended in 0.5 mL of TRIzol (Thermo Fisher # 15596018). Total RNA was extracted from each pellet following the manufacturer's protocol with glycogen (Thermo Fisher # R0561) added at the isopropanol precipitation step. The resulting RNA pellet was resuspended in 30 µL nuclease-free water. 16 µL of extracted RNA was converted to cDNA and analyzed by RT-qPCR as described in above. For each sample, nLuc reporter mRNA abundance was normalized to FFLuc mRNA (coding

sequence from pGL4.13) using the Bio-Rad CFX Maestro software. See **Supplemental Figure S9** for flowchart. Primers for RT-qPCR can be found in **Supplemental Table S1**.

RESULTS
FMRP inhibits translation independent of mRNA G4s

Previous reports have shown that deletion of the RGG box motif from FMRP abolishes translational repression and ribosome binding *in vitro* (22). Dependence of the RGG box motif provided support to others that FMRP must target mRNAs and/or ribosomes by binding to intramolecular RNA G4s (22,24). When tested in isolation, the FMRP RGG box motif has high affinity for G-rich sequences that can form RNA G4 structures (15,17,32,33). Together, these data shaped the leading model that FMRP binds to RNA G4s in the CDS via the RGG box motif and then sterically blocks the A site of an elongating ribosome (22,24). This model has yet to be directly tested as most reporters used by the field harbor predicted G4 sequences and altering the sequence would mutate the reporter protein. To our knowledge, it has yet to be shown experimentally that RNA G4s are present and required in coding sequences for FMRP to inhibit translation.

To directly test this model, we first dual-affinity purified the N-terminally truncated human FMRP (NT-hFMRP) (**Figure 1A, B**) because it is more stable than the full-length isoform and it retains translational repression activity (22,30). We then generated specialized nLuc reporters that either lacked or harbored one of two RNA G4 structures within the CDS (**Figure 1C**). Importantly, this nLuc nucleotide sequence was customized to lack predicted G4 structure without altering the amino acid sequence of the reporter protein. We also included a P2A ribosome skipping motif, which releases the nascent peptide but allows the ribosome to stay bound and continue elongation, directly upstream of the nLuc CDS. This allows uniform luciferase detection across all reporters. To experimentally confirm this reporter design, we took advantage of the selective G4 staining properties of N-methyl mesoporphyrin IX (NMM) (34). As expected, the total RNA stain SYBR Green II detected both control and G4-containing reporter mRNAs (**Figure 1D, left panel**). However, NMM staining only detected the G4 reporter mRNAs (**Figure 1D, right panel**). This same selective staining pattern of NMM for the G4 reporter mRNAs was also seen in native PAGE (**Supplemental Figure S1**). These data support that only the G4 reporter mRNAs form an intramolecular RNA G4 structure (as depicted in **Figure 1C**).

We next tested to what extent recombinant NT-hFMRP represses translation of control and G4 reporter mRNAs. If FMRP did in fact require RNA G4s on target transcripts to inhibit translation, we would expect enhanced repression on both the G_{15} and $(GGGU)_4$ reporters. However, FMRP repressed translation of the G4 reporter mRNAs marginally less than the control reporter mRNA (**Figure 1E**). Identical results were seen with a reporter harboring the Sc1 RNA (**Supplemental Figure S2**). These data indicate that human FMRP represses translation independent of mRNA G4s in the CDS.

For FMRP to inhibit translation elongation on select mRNAs, it is logical that it must bind target mRNAs before acting on a translating ribosome. High throughput sequencing with crosslinking immunoprecipitation of FMRP from mouse brain tissue revealed that FMRP binds to mRNA predominately in the CDS (21). However, these data are derived from multiple RBDs within FMRP, and it is not yet known which binding sites represent true FMRP translational repression targets (as opposed to mRNA transport or localization). Recombinant FMRP can also bind purified 80S ribosomes near the A site and purified 60S ribosomal subunits alone

(22,24,35), raising the possibility that FMRP can directly inhibit the ribosome independent of the mRNA sequence.

To determine if FMRP requires binding to target mRNA first to inhibit translation in our assay, we performed *in vitro* translation assays using different pre-incubation protocols (**Figure 1F**). As a negative control, we programed *in vitro* translation assays with reporter mRNA and protein storage buffer. FMRP was either added directly to the translation reaction immediately before the *in vitro* translation reaction began or was allowed to first form an mRNP with reporter mRNA (which was used in **Figure 1E**). Translational repression was only observed when FMRP was pre-incubated with the reporter mRNA (**Figure 1F**), demonstrating that FMRP must bind a target first to inhibit translation. To further dissect this mechanism, we solely used the mRNA•FMRP pre-incubation strategy and the control nLuc mRNA reporter that lacks an RNA G4 for all remaining experiments.

The RGG box motif and CTD of hFMRP together, but not independently, inhibit translation

We next sought to identify the critical RNA-binding element in FMRP required for translational repression. FMRP contains at least three canonical RBDs (KH0, KH1, KH2) and a single RGG box motif. FXS patient mutations suggest that multiple regions of FMRP are critical for RNA binding-dependent function and contribute to pathology if mutated. The I304N patient mutation in the KH2 domain abolishes FMRP binding to polysomes in human cells (16). A guanosine insertion (ΔRGG+CTD) within the sequence that encodes the RGG box motif causes a frameshift and results in a truncated FMRP that lacks most of the RGG box motif and the entire C-terminal domain (CTD) (20).

To further define the domain(s) that are critical for translational repression by FMRP, we purified recombinant NT-hFMRP harboring I304N and ΔRGG+CTD mutations (**Figure 2A, B**) and tested their ability to inhibit translation. Multiple attempts were made to purify a KH1 domain G266E mutant (18), but we were unable to recover soluble protein. We then determined the half-maximal inhibitory concentration (IC_{50}) of WT and each mutant NT-hFMRP (**Figure 2C-F**). As expected, the His6-MBP tag alone did not inhibit translation (**Figure 2C**). WT and I304N NT-hFMRP both inhibited translation in our assay, with the I304N mutant having a ~2-fold more potent IC_{50} than that of the WT isoform (**Figure 2D, E**). This suggests that although the I304N mutation alters FMRP binding to an optimal RNA pseudoknot substrate (i.e., Δkc2) and causes FMRP to dissociate from polysomes in cells (16), the mutation does not interfere with translational repression.

The ΔRGG+CTD mutant that contains the FXS-patient guanosine insertion did not repress translation (**Figure 2F**). This insertion mutation creates a frameshift in the RGG box motif and results in the encoding of a short novel peptide upstream of a premature termination codon. To validate the loss of translational repression was due to the truncation of the RGG box motif and complete deletion of the CTD rather than the addition of the short novel peptide, we purified a mutant of the NT-hFMRP that completely lacks both the RGG box motif as well as the CTD, which we termed NT-hFMRP ΔRGG+CTD complete (**Supplemental Figure S3A, B**). Full deletion of the RGG+CTD resulted in loss of translational repression (**Supplemental Figure S3C**), which is in alignment with previous reports (22,24). Taken together, the RGG+CTD is essential for translational repression.

To determine if the RGG+CTD was not only essential but also sufficient for translational repression, we purified the RGG box motif and CTD regions both together and separately (**Figure 2A, B**). Robust translational repression was observed with the isolated RGG+CTD

region (**Figure 2G**). However, neither the RGG box alone nor the CTD alone effectively inhibited translation (**Figure 2H, I**). Together, these data suggest that the RGG box must be appended to the CTD to inhibit translation and that the RGG+CTD is sufficient for translational repression by FMRP (22,24).

The RGG+CTD region forms a ncRBD that has broad RNA binding ability

We next sought to determine the RNA binding capability of the RGG+CTD region. We used electrophoretic mobility shift assays (EMSAs) to test the ability of the RGG box motif and CTD, both together and separately, to bind FAM-labeled homopolymeric RNAs. The first identified RGG box motif, belonging to heterogenous nuclear ribonucleoprotein (RNP) U, was found to bind both homopolymeric G and U RNA sequences *in vitro*, with a higher preference for polymeric G RNA (36). In agreement with the FMRP RGG box motif favoring G-rich sequences and G4s (15,37), the RGG box motif alone had some observable binding to $U(G)_{17}$ RNA, but little to no binding to $U(A)_{17}$, $U(C)_{17}$, or $(U)_{18}$ RNA (**Figure 3A-E**). Similar results were seen with the positively charged CTD (**Figure 3A-E**). Conversely, the RGG+CTD robustly bound all four homopolymeric RNAs, with slightly less preference for $U(A)_{17}$ RNA (**Figure 3A-E**). These data suggest that the RGG box motif and CTD must be together to form a non-canonical RNA-binding domain (ncRBD) that can bind to a wide range of RNAs.

It should be noted that we routinely did not see the RNP enter the gel for most of the homopolymeric RNAs tested, suggesting the high positive charge of the RGG+CTD neutralized the negative charge of the RNA oligo and/or the FAM label was quenched. The RGG box motif and CTD have theoretical isoelectric points of 12.1 and 10.0, respectively. Nevertheless, to be as consistent as possible across all the samples, unbound homopolymeric RNA was quantified (**Figure 3E**). It is possible that the $U(G)_{17}$ formed intermolecular G4s, providing a high net negative charge to the complex that facilitated its entry into the gel if RGG+CTD was sub-stoichiometric.

To further dissect how the ncRBD formed by the RGG+CTD elicits FMRP to inhibit translation, we tested whether recombinant NT-hFMRP WT and each mutant can bind reporter mRNA by EMSA (**Figure 3F**). Remarkably, the ability of WT and each mutant NT-hFMRP to bind reporter mRNA (**Figure 3F**) mirrors the observed translational repression with each mutant (**Figure 2**), except for ΔRGG+CTD. WT and I304N NT-hFMRP, as well as the RGG+CTD fusion, caused a complete gel shift (**Figure 3F**) and were translationally repressive (**Figure 2D, E, G**). RGG box alone and CTD alone did not cause a gel shift and were not robust translational repressors. Unique, ΔRGG+CTD did cause a complete gel shift (**Figure 3F**) but was not translationally repressive (**Figure 2F**). We rationalized that the functional KH domains in ΔRGG+CTD allowed mRNA binding since the ncRBD was absent in this mutant. Indeed, adding the I304N mutation in the KH2 domain to ΔRGG+CTD blocked reporter mRNA binding (**Figure 3F, Supplemental Figure S4**). Identical results were seen with mEGFP mRNA (**Supplemental Figure S5**). The decreased binding observed with ΔRGG+CTD when the I304N mutation was added suggests that the KH2 does compete with the ncRBD. This is also observed in our translational repression assays with the I304N mutant having an ~2-fold lower IC_{50} compared to WT **(Figure 2D,E)**. Together, these data support that the RGG box motif and CTD forms a ncRBD that elicits broad RNA binding.

Discrete regions of the ncRBD are required for translational repression and mRNA binding

We next asked if mRNA-binding and translational repression are elicited by the same or different regions of the ncRBD. Our pre-incubation data (**Figure 1F**) suggest that mRNA-binding is a prerequisite for FMRP translational repression. Thus, we postulated that if the region of the

ncRBD that elicited mRNA-binding and translational repression were the same, we would see a mirrored loss in mRNA-binding and translational repression when mutated. The ncRBD of human FMRP is highly conserved among most vertebrates and is predicted to be largely flexible and disordered (**Supplemental Figure S6**), which hindered our ability to make refined mutations *a priori* as typically is done for canonical RBDs. We generated a series of large and more refined serial deletions of the CTD from the C-terminal end (**Figure 4A**) and identified a sharp decline in translational repression between the RGG+CTD Δ54 and RGG+CTD Δ55 mutants (**Figure 4B**). This transition point between deleting 54 or 55 amino acids from the C-terminus of the ncRBD was also present when tested in the NT-hFMRP isoform that harbors the other canonical RBDs (**Supplemental Figure S7**). NT-hFMRP Δ54 had an IC$_{50}$ of ~0.88 µM (**Figure 4C**) and NT-hFMRP Δ55 had an IC$_{50}$ of ~2.29 µM (**Figure 4D**). We attempted to introduce the I304N KH2 domain mutation to offset any altered binding of the truncated C-terminus but these mutants were insoluble. These data demonstrated that RGG+CTD Δ54 region of the ncRBD represents the minimal repressive element of human FMRP.

If the change in translational repression between RGG+CTD Δ54 and RGG+CTD Δ55 was due to altered mRNA-binding ability, we would predict a similar drastic change in mRNA binding. However, both RGG+CTD Δ54 and RGG+CTD Δ55 had nearly identical ability to cause a robust gel shift of reporter mRNA by EMSA (**Figure 5A, B**). Further deletional analysis and serial single amino acid truncations (RGG+CTD Δ60, Δ61, Δ62, and Δ63) identified RGG+CTD Δ62 as the key region of the ncRBD for robust mRNA binding (**Figure 5A, B**). Together, these data suggest that the regions of the ncRBD that are responsible for mRNA binding and translational repression overlap but are not identical (**Figure 5C**).

FMRP inhibits translation post-initiation when bound to mRNA via the ncRBD
FMRP bound to mRNA could directly inhibit translation by either blocking the scanning pre-initiation complex (PIC) in the 5 UTR or stalling the elongating ribosome in the CDS. In general, scanning PICs are more susceptible and sensitive to obstacles in their path (i.e., RNA structure or bound RBPs) than elongating ribosomes. Most mRNA-bound FMRP is mapped to the CDS of mRNAs *in vivo*, not to the 5 UTR (21). However, in our assays, it is possible that a portion of recombinant FMRP is bound to the 5 UTR and is simply blocking the scanning PIC. To confirm that FMRP inhibits translation post-initiation, consistent with previous reports that FMRP slows or stalls elongating ribosomes (21-25), we used the following three distinct strategies.

First, we rationalized that if 5 UTR-bound recombinant FMRP was blocking scanning PICs, extending the 5 UTR length would enhance repression. In this case, extending the 5 UTR would provide FMRP increased opportunity to bind the 5 UTR instead of the CDS. To achieve this, we mutated the control reporter that harbors the 50 nt human β-globin 5 UTR and extended the 5 UTR three-fold by inserting two additional β-globin 5 UTR sequences (resulting in a 150 nt 5 UTR). Nevertheless, FMRP inhibited translation of the control and long 5 UTR reporter mRNAs to similar extents (**Figure 6A**). These data suggest that the predominant mechanism by which FMRP inhibits translation in our assays is not by inhibiting scanning PICs.

Second, we used sucrose density gradient ultracentrifugation to confirm the expected distribution of mRNAs with stalled elongating ribosomes. For example, elongation inhibitors (e.g., cycloheximide or emetine) cause an increase in polysomes as they do not prevent initiation, but slow down and stabilize elongating ribosomes on mRNAs (38). This is typically seen as an increase in polysome signal in the heavier fractions of sucrose gradients. If FMRP inhibits translation post-initiation when bound to mRNA via its ncRBD, we would predict that

polysomes accumulate on inhibited transcripts only when FMRP harbored the RGG+CTD ncRBD. To test this prediction, we used the ΔRGG+CTD mutant as a negative control as it does not inhibit translation (**Figure 2F**) but still binds the reporter mRNA (**Figure 3F**) in our assays. After *in vitro* translation and ultracentrifugation, we quantified nLuc reporter mRNA abundance in each fraction. The monosome peak (which is primarily inactive 80S ribosomes native to RRL (39)) was routinely found in fractions 4 and 5 (**Supplemental Figure S8A**), indicating that polysomes would sediment in fractions 6 through 12. Compared to the negative control that lacks the translationally repressive ncRBD, WT NT-hFMRP increased the abundance of nLuc mRNA in the heavy polysomes (**Figure 6B-D**, **Supplemental Figure S8B-E**). Specifically, nLuc mRNA abundance in the heavy polysomes in fractions 10-12 increased ~2-fold (**Figure 6C, D, Supplemental Figure S8C, E**). This increase of reporter mRNA at the bottom of the gradient with the heavy polysomes is consistent with accumulation of slowed and stalled ribosomes.

Lastly, we tested the ability of NT-hFMRP to generate ribosome complexes on reporter mRNPs that are resistant to puromycin. FMRP was previously identified by immunogold labeling and electron microscopy to be bound to ribosomes within puromycin-resistant polysomes from mouse brain (21). Puromycin is an aminonucleoside antibiotic and acyl tRNA analog that is incorporated into nascent polypeptides and results in ribosomes releasing both the nascent polypeptide and bound mRNA. Puromycin sensitivity is specific for elongating 80S ribosomes that are in an unrotated, pre-translocation state (e.g., during the decoding step of elongation or when inhibited by emetine). Ribosomes that are stalled in the rotated state and inhibited during translocation (e.g., when inhibited by cycloheximide) are resistant to puromycin and stay bound to mRNA. We predicted that NT-hFMRP would stall and slow elongating ribosomes and elicit puromycin-resistance only when harboring the ncRBD. We optimized a low-speed sucrose cushion protocol to be selective for mRNAs only bound by ribosomes to be recovered, allowing us to assay puromycin-resistant ribosomes on nLuc•FMRP mRNPs (**Supplemental Figure S9**). A series of controls demonstrates that nLuc mRNA is detected in the ribosome pellet only after being translated and this detection is strongly prevented when translation reactions are treated with 0.1 mM puromycin prior to low-speed centrifugation over a 35% sucrose cushion (**Figure 6E, Lanes 1-4**). Only minor amounts of nLuc mRNA were recovered from untranslated reactions with nLuc•NT-FMRP mRNPs (ΔRGG+CTD or WT) in RRL that were kept on ice (**Figure 6E, Lanes 5 & 6**). In agreement with our prediction that only NT-hFMRP harboring the ncRBD would elicit puromycin-resistance, ribosome-bound nLuc•NT-hFMRP ΔRGG+CTD mRNP, but not nLuc•NT-hFMRP WT mRNP, was sensitive to puromycin (**Figure 6F, Lanes 1 & 2**). We were unable to detect a change in pelleted nLuc from the translated nLuc•NT-hFMRP WT mRNP when treated with puromycin (**Figure 6F, Lanes 3 & 4**). In total, these data support that FMRP uses a ncRBD formed by the RGG box motif and CTD to bind mRNA independent of mRNA G4s resulting in ribosome stalling and translational repression (**Figure 7**).

DISCUSSION

The complexity of molecular and cellular phenotypes seen in FMRP-null neurons and model systems has led to much debate surrounding how FMRP targets transcripts and what functions it may possess once bound to mRNA (40). Most early data support FMRP as a translational repressor (21-25), but recent work in *Drosophila* and mouse models has provided compelling evidence that FMRP may in fact also act a translational activator (26-28). It remains largely unclear which RNA motifs or structures FMRP targets as few common enriched sequences were found across multiple transcriptome-wide FMRP binding studies (reviewed in 40). A single consensus sequence for FMRP may be an oversimplification since FMRP has multiple RBDs and motifs.

Here we refine the RNA-binding element that contains the RGG box motif and provide evidence that FMRP inhibits translation after binding mRNA via the ncRBD comprised of the RGG box motif and the CTD. The ncRBD is essential and sufficient for repression (24,41) (**Figure 2**) and required for FMRP to bind mRNA to inhibit translation (**Figures 2 & 3**). Scanning deletion analysis of the ncRBD identified that the residues required for mRNA binding reside within the region critical for translational repression (**Figures 4 & 5**). We and others (24,41) have found the KH1 and KH2 domains dispensable for translational repression, yet cryo-EM determination of *Drosophila* FMRP•80S ribosome complexes show that KH1 and KH2 domains partially overlap with where the anticodon stem of a bound P-site tRNA would be located. In this structure, the RGG+CTD region of FMRP is near the A site and leading edge of the ribosome (22). Chen *et al.* proposed that the RGG box motif binds to the mRNA via G4s within the CDS and that the KH domains bind directly to the ribosome near the A site to sterically hinder delivery of charged tRNAs (22). Early studies identified FMRP-bound polysomes from mouse brain lysate as puromycin-resistant, a characteristic we demonstrate is reproduced in our translation assays (**Figure 6E, F**). Puromycin is a substrate for peptidyl transferase only when ribosomes are in an unrotated state during the decoding step of elongation. Puromycin-resistant FMRP-stalled ribosomes are thus thought to be inhibited during translocation (21). However, the fly FMRP•80S ribosome structure was solved using empty 80S ribosomes in a unrotated state unbound to mRNA. To our knowledge, our report is the first to provide *in vitro* biochemical data that demonstrate FMRP creates puromycin-resistant ribosome complexes on mRNA, but in agreement with what was previously shown in brain tissue lysates (21). The abundance of inhibited transcripts on heavier polysomes confirms translational repression is not due to decreased initiation (**Figure 6B-D; Supplemental Figure S8**). Our approach reported here provides valuable samples and tools to enrich for these complexes, which is not easily achievable from brain tissue lysates. Future structural determination of FMRP-inhibited ribosomes on mRNA will be critical for fully understanding FMRP-mediated translational repression.

Although the RGG box motif within the FMRP ncRBD has high affinity to the G4-containing Sc1 RNA (17), this interaction is not required for translational repression, nor does it provide enhanced translational repression (**Figure 1, Supplemental Figure S2**). In fact, structural determinations of the RGG box motif and Sc1 RNA complex show that the RGG box motif actually binds to the major groove of the duplexed RNA region of the Sc1 RNA and not to the G4 itself (32,33). Moreover, G4s are enriched in the UTRs of mRNAs but not in the CDS (42), while FMRP predominately binds to the CDS but not to the UTRs (21). If FMRP is recruited to the CDS via RGG box-G4 interaction, one would predict the distributions of FMRP binding events and G4s to be positively correlated, but they are instead negatively correlated. Recent ribosome profiling studies that incorporated RNA G4 prediction analyses show that transcripts that are derepressed in FMRP knockout cell lines are not enriched for RNA G4s (43). Goering *et al.* conclude that RNA G4s were instead correlated with FMRP-mediated mRNA localization, not translational repression. A more recent study from Darnell and colleagues found that G4s were not enriched in dendritic FMRP targets in CA1 pyramidal neurons (44).

Consistent with our finding that repression occurs independent of RNA G4s in the CDS (**Figure 1**), the ncRBD has a broad ability to bind RNA (**Figure 3, Supplemental Figure S5**). In further support of this G4-independent mode of translational repression by FMRP, two other reports have identified that FMRP binds to a stem loop-containing RNA (devoid of a G4) to inhibit translation. First, Maurin *et al.* identified mammalian FMRP inhibited expression of a reporter mRNA that contained theSL1 and SL2 stem-loops when inserted into the open reading frame but not either UTR. Which RNA-binding domain (KH1, KH2, or ncRBD) was responsible for this inhibition was not reported (45). Second, while this report was in preparation, Edwards *et*

al. reported that the RGG+CTD region bound to a stem-loop in a short peptide reporter mRNA to inhibit translation (41). For both reports, further studies are warranted to determine if the stem-loops in the open reading frame cause FMRP to inhibit initiation and/or elongation. Nevertheless, the stem-loop sequences identified by Maurin *et al.* and Edwards *et al.* are absent in our nLuc reporter mRNA. Our finding that the ncRBD can bind all four homopolymeric RNAs and multiple G4-less reporter mRNAs suggests that FMRP binding is more promiscuous than previously appreciated. Future iCLIP studies in cells expressing the ncRBD alone will be important to determine, if any, sequence or RNA secondary structure preferences for binding and subsequent translational repression.

The structure of the ncRBD in FMRP bound or unbound to RNA is not known, but is predicted to be disordered and flexible (**Supplemental Figure S6**). A small fragment of the RGG box motif was used in determining the RGG box•Sc1 RNA complex and was found to be highly flexible. Recent reports have also shown that small repetitive regions (46) (e.g., poly-K patches, poly-R patches, and YGG box motifs) and other larger flexible regions have previously undefined RNA-binding abilities. For example, the disordered and flexible N-terminal half of *C. elegans* MEG-3 elicits RNA-binding (yet, does not cause phase transition that is commonly reported for such disordered regions) (47). Additionally, Xu *et al.* recently characterized human nuclear hormone receptor estrogen receptor α (ERα) as an RNA-binding protein that contains a functional arginine-rich RBD with binding preference to A-U rich sequences in 3□ UTRs (48) to control post-transcriptional regulation. Further assessment in AlphaFold and IUPred predicts the arginine-rich RBD in ERα is centered in a large highly flexible region (data not shown), similar to the FMRP ncRBD we define in this report. Our understanding of how flexible RNA-binding domains mechanistically associate with RNA is increasing but whether they contain sequence specificity is still unclear.

FMRP also facilitates the transport of dendritic mRNAs to synapses *in vivo* and this transport can be dependent on its ability to bind mRNA (43,49). Given that the translationally repressive ncRBD binds mRNA more promiscuously than previously appreciated (**Figure 3, Supplemental Figure S5**) and the lack of a shared consensus sequence that has been found throughout multiple transcriptome-wide FMRP binding screens (40), the data supports a model where FMRP could regulate translation of dendritic mRNAs primarily based on local concentration and not mRNA sequence. Since FMRP and dendritic mRNAs are transported to the synapses, often together, both are found at relatively high concentrations compared to other translationally active sites of neurons. Future studies of directly shuttling FMRP from one region of neuron to another and monitoring its ability to repress translation of proximal transcripts would prove beneficial in further defining a concentration-dependent model.

FUNDING

MRS was supported by The Ohio State University Fellowship and The Ohio State University Center for RNA Biology Graduate Fellowship. This work was supported by NIH grant R00GM126064 to MGK. The Ohio State University Comprehensive Cancer Center Genomics Shared (OSUCCC GSR) is supported by NIH grant P30CA016058.

ACKNOWLEDGEMENTS

Experiments were conceived by MRS and MGK and were performed by MRS, JEW, ELY, and MGK. The manuscript was written by MRS and MGK with input from JEW and ELY. We thank members of the Kearse lab and Aaron Goldstrohm for critically reading the

manuscript. We also thank Brian Scarpitti for his help with image analyses and Christine Daugherty at the OSUCCC GSR.

REFERENCES

1.   Kremer, E. J., Pritchard, M., Lynch, M., Yu, S., Holman, K., Baker, E., Warren, S. T., Schlessinger, D., Sutherland, G. R., and Richards, R. I. (1991) Mapping of DNA instability at the fragile X to a trinucleotide repeat sequence p(CCG)n. *Science* **252**, 1711-1714

2.   Oberle, I., Rousseau, F., Heitz, D., Kretz, C., Devys, D., Hanauer, A., Boue, J., Bertheas, M. F., and Mandel, J. L. (1991) Instability of a 550-base pair DNA segment and abnormal methylation in fragile X syndrome. *Science* **252**, 1097-1102

3.   Verkerk, A. J., Pieretti, M., Sutcliffe, J. S., Fu, Y. H., Kuhl, D. P., Pizzuti, A., Reiner, O., Richards, S., Victoria, M. F., Zhang, F. P., and et al. (1991) Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* **65**, 905-914

4.   Yu, S., Pritchard, M., Kremer, E., Lynch, M., Nancarrow, J., Baker, E., Holman, K., Mulley, J. C., Warren, S. T., Schlessinger, D., and et al. (1991) Fragile X genotype characterized by an unstable region of DNA. *Science* **252**, 1179-1181

5.   Niu, M., Han, Y., Dy, A. B. C., Du, J., Jin, H., Qin, J., Zhang, J., Li, Q., and Hagerman, R. J. (2017) Autism Symptoms in Fragile X Syndrome. *J Child Neurol* **32**, 903-909

6.   Kelleher, R. J., 3rd, and Bear, M. F. (2008) The autistic neuron: troubled translation? *Cell* **135**, 401-406

7.   Ashley, C. T., Jr., Wilkinson, K. D., Reines, D., and Warren, S. T. (1993) FMR1 protein: conserved RNP family domains and selective RNA binding. *Science* **262**, 563-566

8.   Siomi, H., Siomi, M. C., Nussbaum, R. L., and Dreyfuss, G. (1993) The protein product of the fragile X gene, FMR1, has characteristics of an RNA-binding protein. *Cell* **74**, 291-298

9.   Myrick, L. K., Hashimoto, H., Cheng, X., and Warren, S. T. (2015) Human FMRP contains an integral tandem Agenet (Tudor) and KH motif in the amino terminal domain. *Hum Mol Genet* **24**, 1733-1740

10.   Jensen, K. B., Musunuru, K., Lewis, H. A., Burley, S. K., and Darnell, R. B. (2000) The tetranucleotide UCAY directs the specific recognition of RNA by the Nova K-homology 3 domain. *Proc Natl Acad Sci U S A* **97**, 5740-5745

11.   Lewis, H. A., Musunuru, K., Jensen, K. B., Edo, C., Chen, H., Darnell, R. B., and Burley, S. K. (2000) Sequence-specific RNA binding by a Nova KH domain: implications for paraneoplastic disease and the fragile X syndrome. *Cell* **100**, 323-332

12.   Hollingworth, D., Candel, A. M., Nicastro, G., Martin, S. R., Briata, P., Gherzi, R., and Ramos, A. (2012) KH domains with impaired nucleic acid binding as a tool for functional analysis. *Nucleic Acids Res* **40**, 6873-6886

13.   Nicastro, G., Taylor, I. A., and Ramos, A. (2015) KH-RNA interactions: back in the groove. *Curr Opin Struct Biol* **30**, 63-70

14.   Nicastro, G., Candel, A. M., Uhl, M., Oregioni, A., Hollingworth, D., Backofen, R., Martin, S. R., and Ramos, A. (2017) Mechanism of beta-actin mRNA Recognition by ZBP1. *Cell Rep* **18**, 1187-1199

15.   Athar, Y. M., and Joseph, S. (2020) RNA-Binding Specificity of the Human Fragile X Mental Retardation Protein. *J Mol Biol* **432**, 3851-3868

16.   Darnell, J. C., Fraser, C. E., Mostovetsky, O., Stefani, G., Jones, T. A., Eddy, S. R., and Darnell, R. B. (2005) Kissing complex RNAs mediate interaction between the Fragile-X

mental retardation protein KH2 domain and brain polyribosomes. *Genes Dev* **19**, 903-918

17.   Darnell, J. C., Jensen, K. B., Jin, P., Brown, V., Warren, S. T., and Darnell, R. B. (2001) Fragile X mental retardation protein targets G quartet mRNAs important for neuronal function. *Cell* **107**, 489-499

18.   Myrick, L. K., Nakamoto-Kinoshita, M., Lindor, N. M., Kirmani, S., Cheng, X., and Warren, S. T. (2014) Fragile X syndrome due to a missense mutation. *Eur J Hum Genet* **22**, 1185-1189

19.   De Boulle, K., Verkerk, A. J., Reyniers, E., Vits, L., Hendrickx, J., Van Roy, B., Van den Bos, F., de Graaff, E., Oostra, B. A., and Willems, P. J. (1993) A point mutation in the FMR-1 gene associated with fragile X mental retardation. *Nat Genet* **3**, 31-35

20.   Okray, Z., de Esch, C. E., Van Esch, H., Devriendt, K., Claeys, A., Yan, J., Verbeeck, J., Froyen, G., Willemsen, R., de Vrij, F. M., and Hassan, B. A. (2015) A novel fragile X syndrome mutation reveals a conserved role for the carboxy-terminus in FMRP localization and function. *EMBO Mol Med* **7**, 423-437

21.   Darnell, J. C., Van Driesche, S. J., Zhang, C., Hung, K. Y., Mele, A., Fraser, C. E., Stone, E. F., Chen, C., Fak, J. J., Chi, S. W., Licatalosi, D. D., Richter, J. D., and Darnell, R. B. (2011) FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* **146**, 247-261

22.   Chen, E., Sharma, M. R., Shi, X., Agrawal, R. K., and Joseph, S. (2014) Fragile X mental retardation protein regulates translation by binding directly to the ribosome. *Mol Cell* **54**, 407-417

23.   Liu, B., Li, Y., Stackpole, E. E., Novak, A., Gao, Y., Zhao, Y., Zhao, X., and Richter, J. D. (2018) Regulatory discrimination of mRNAs by FMRP controls mouse adult neural stem cell differentiation. *Proc Natl Acad Sci U S A* **115**, E11397-E11405

24.   Athar, Y. M., and Joseph, S. (2020) The Human Fragile X Mental Retardation Protein Inhibits the Elongation Step of Translation through Its RGG and C-Terminal Domains. *Biochemistry* **59**, 3813-3822

25.   Shah, S., Molinaro, G., Liu, B., Wang, R., Huber, K. M., and Richter, J. D. (2020) FMRP Control of Ribosome Translocation Promotes Chromatin Modifications and Alternative Splicing of Neuronal Genes Linked to Autism. *Cell Rep* **30**, 4459-4472 e4456

26.   Greenblatt, E. J., and Spradling, A. C. (2018) Fragile X mental retardation 1 gene enhances the translation of large autism-related proteins. *Science* **361**, 709-712

27.   Shu, H., Donnard, E., Liu, B., Jung, S., Wang, R., and Richter, J. D. (2020) FMRP links optimal codons to mRNA stability in neurons. *Proc Natl Acad Sci U S A* **117**, 30400-30411

28.   Flanagan, K., Baradaran-Heravi, A., Yin, Q., Dao Duc, K., Spradling, A. C., and Greenblatt, E. J. (2022) FMRP-dependent production of large dosage-sensitive proteins is highly conserved. *Genetics* **221**

29.   Kearse, M. G., Green, K. M., Krans, A., Rodriguez, C. M., Linsalata, A. E., Goldstrohm, A. C., and Todd, P. K. (2016) CGG Repeat-Associated Non-AUG Translation Utilizes a Cap-Dependent Scanning Mechanism of Initiation to Produce Toxic Proteins. *Mol Cell* **62**, 314-322

30.   Edwards, M., Xu, M., and Joseph, S. (2020) A simple procedure for bacterial expression and purification of the fragile X protein family. *Sci Rep* **10**, 15858

31.   Pringle, E. S., McCormick, C., and Cheng, Z. (2019) Polysome Profiling Analysis of mRNA and Associated Proteins Engaged in Translation. *Curr Protoc Mol Biol* **125**, e79

32.   Phan, A. T., Kuryavyi, V., Darnell, J. C., Serganov, A., Majumdar, A., Ilin, S., Raslin, T., Polonskaia, A., Chen, C., Clain, D., Darnell, R. B., and Patel, D. J. (2011) Structure-function studies of FMRP RGG peptide recognition of an RNA duplex-quadruplex junction. *Nat Struct Mol Biol* **18**, 796-804

33. Vasilyev, N., Polonskaia, A., Darnell, J. C., Darnell, R. B., Patel, D. J., and Serganov, A. (2015) Crystal structure reveals specific recognition of a G-quadruplex RNA by a beta-turn in the RGG motif of FMRP. *Proc Natl Acad Sci U S A* **112**, E5391-5400

34. Yett, A., Lin, L. Y., Beseiso, D., Miao, J., and Yatsunyk, L. A. (2019) N-methyl mesoporphyrin IX as a highly selective light-up probe for G-quadruplex DNA. *J Porphyr Phthalocyanines* **23**, 1195-1215

35. Siomi, M. C., Zhang, Y., Siomi, H., and Dreyfuss, G. (1996) Specific sequences in the fragile X syndrome protein FMR1 and the FXR proteins mediate their binding to 60S ribosomal subunits and the interactions among them. *Mol Cell Biol* **16**, 3825-3832

36. Kiledjian, M., and Dreyfuss, G. (1992) Primary structure and binding activity of the hnRNP U protein: binding RNA through RGG box. *EMBO J* **11**, 2655-2664

37. Brown, V., Small, K., Lakkis, L., Feng, Y., Gunter, C., Wilkinson, K. D., and Warren, S. T. (1998) Purified recombinant Fmrp exhibits selective RNA binding as an intrinsic property of the fragile X mental retardation protein. *J Biol Chem* **273**, 15521-15527

38. Juszkiewicz, S., Slodkowicz, G., Lin, Z., Freire-Pritchett, P., Peak-Chew, S. Y., and Hegde, R. S. (2020) Ribosome collisions trigger cis-acting feedback inhibition of translation initiation. *Elife* **9**

39. Shao, S., Murray, J., Brown, A., Taunton, J., Ramakrishnan, V., and Hegde, R. S. (2016) Decoding Mammalian Ribosome-mRNA States by Translational GTPase Complexes. *Cell* **167**, 1229-1240.e1215

40. Richter, J. D., and Zhao, X. (2021) The molecular biology of FMRP: new insights into fragile X syndrome. *Nat Rev Neurosci* **22**, 209-222

41. Edwards, M., Huang, M., and Joseph, S. (2022) The Fragile X Protein Disordered Regions Bind a Novel RNA Target. *Biochemistry* **61**, 1199-1212

42. Kwok, C. K., Marsico, G., Sahakyan, A. B., Chambers, V. S., and Balasubramanian, S. (2016) rG4-seq reveals widespread formation of G-quadruplex structures in the human transcriptome. *Nat Methods* **13**, 841-844

43. Goering, R., Hudish, L. I., Guzman, B. B., Raj, N., Bassell, G. J., Russ, H. A., Dominguez, D., and Taliaferro, J. M. (2020) FMRP promotes RNA localization to neuronal projections through interactions between its RGG domain and G-quadruplex RNA sequences. *Elife* **9**

44. Hale, C. R., Sawicka, K., Mora, K., Fak, J. J., Kang, J. J., Cutrim, P., Cialowicz, K., Carroll, T. S., and Darnell, R. B. (2021) FMRP regulates mRNAs encoding distinct functions in the cell body and dendrites of CA1 pyramidal neurons. *Elife* **10**

45. Maurin, T., Melko, M., Abekhoukh, S., Khalfallah, O., Davidovic, L., Jarjat, M., D'Antoni, S., Catania, M. V., Moine, H., Bechara, E., and Bardoni, B. (2015) The FMRP/GRK4 mRNA interaction uncovers a new mode of binding of the Fragile X mental retardation protein in cerebellum. *Nucleic Acids Res* **43**, 8540-8550

46. Castello, A., Fischer, B., Eichelbaum, K., Horos, R., Beckmann, B. M., Strein, C., Davey, N. E., Humphreys, D. T., Preiss, T., Steinmetz, L. M., Krijgsveld, J., and Hentze, M. W. (2012) Insights into RNA biology from an atlas of mammalian mRNA-binding proteins. *Cell* **149**, 1393-1406

47. Schmidt, H., Putnam, A., Rasoloson, D., and Seydoux, G. (2021) Protein-based condensation mechanisms drive the assembly of RNA-rich P granules. *Elife* **10**

48. Xu, Y., Huangyang, P., Wang, Y., Xue, L., Devericks, E., Nguyen, H. G., Yu, X., Oses-Prieto, J. A., Burlingame, A. L., Miglani, S., Goodarzi, H., and Ruggero, D. (2021) ERα is an RNA-binding protein sustaining tumor cell survival and drug resistance. *Cell* **184**, 5215-5229.e5217

49. Ramos, A., Hollingworth, D., and Pastore, A. (2003) G-quartet-dependent recognition between the FMRP RGG box and RNA. *RNA* **9**, 1198-1207

Figure 1. FMRP inhibits translation independent of mRNA G-quadruplexes in the CDS. A) Schematic of MBP- and His6-tagged WT N-terminally truncated human FMRP (NT-hFMRP). B)

Coomassie stain of recombinant WT NT-hFMRP. C) Schematic of custom nLuc reporters either lacking a G4 (control reporter) or harboring a G4 in the coding sequence (G_{15} and $(GGGU)_4$ reporters). A P2A ribosome skipping motif was included immediately upstream of the nLuc coding sequence to ensure equal nLuc function between reporters. D) Denaturing PAGE of control, G_{15}, and $(GGGU)_4$ reporters stained for total RNA with SYBR Green II or for G4 structures with NMM. E) *In vitro* translation of control, G_{15}, and $(GGGU)_4$ reporter mRNA with protein buffer or WT NT-hFMRP. Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using a two-tailed unpaired t test with Welch's correction. F) *In vitro* translation of control nLuc reporter mRNA with protein storage buffer as a negative control, with WT NT-hFMRP and nLuc mRNA pre-incubated together, and with WT NT-hFMRP without a pre-incubation step. Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using a two-tailed unpaired t test with Welch's correction.

Figure 2. The RGG box motif and CTD together are essential and sufficient to inhibit translation. A) Schematic of recombinant WT and mutant NT-hFMRP. Mutated/truncated

domains are highlighted in red. B) Coomassie stain of recombinant proteins. C-I) *In vitro* translation of nLuc mRNA with a titration of recombinant wildtype and mutant NT-hFMRP isoforms. IC_{50} values were determined for the His6-MBP tag (negative control; 633.50 $\pm$ infinity µM) (C), WT NT-hFMRP (D), I304N KH2 domain patient-derived mutant NT-hFMRP (E), ΔRGG+CTD mutant NT-hFMRP (25.72 $\pm$ infinity µM) (F), the RGG box motif + CTD fusion (G), the RGG box motif alone (H), and the CTD alone ($5.75 \times 10^{33} \pm$ infinity µM) (I). n=3 biological replicates. A non-linear regression was used to calculate the IC_{50} and is shown as the line with the 95% confidence interval (CI) included as a watermark. The IC_{50} is reported $\pm$ 95% CI.

Figure 3. The RGG+CTD non-canonical RBD binds all four homopolymeric RNAs and mRNA. A-D) EMSAs of 5□ FAM-labeled homopolymeric RNA oligonucleotides with the indicated recombinant proteins. Due to the suspected net neutral charge of the RNA oligo•FMRP complex causing the RNP to not enter the gel, the intensity of unbound RNA was quantified and set relative to the protein storage buffer only sample (E). F) EMSA of control reporter mRNA incubated with the indicated recombinant protein and stained with SYBR Green II.

Figure 4. The RGG+CTD Δ54 is required for translational repression. A) Coomassie stain of recombinant proteins with serial truncations to the CTD. The number of amino acids truncated from the CTD is indicated. B) *In vitro* translation of nLuc mRNA with CTD truncated recombinant RGG+CTD proteins and protein storage buffer as a negative control. Data are shown as mean ± SD. n = 3 biological replicates. All recombinant protein tested was statistically significant compared to buffer negative control using a two-tailed unpaired t test with Welch's correction (p < 0.01 for all). C, D) *In vitro* translation of nLuc mRNA with a titration of recombinant NT-hFMRP Δ54 (C) and NT-hFMRP Δ55 (D). n=3 biological replicates. A non-linear regression was used to calculate the IC_{50} value for each truncation and is shown as the line with the 95% confidence interval (CI) included as a watermark. The IC_{50} is reported ± 95% CI.

Figure 5. The RGG+CTD Δ62 is the key region of the ncRBD required for robust mRNA-binding. A) Coomassie stain of recombinant proteins with C-terminal truncations to the ncRBD (RGG+CTD). The number of amino acids truncated from the CTD is indicated. B) EMSA of control reporter mRNA incubated with recombinant protein and stained with SYBR Green II. C) Amino acid composition of overlapping regions of the ncRBD that are required for mRNA-binding (RGG+CTD Δ62) and translational repression (RGG+CTD Δ54).

Figure 6. Human FMRP inhibits translation post-initiation when binding mRNA via the ncRBD. A) *In vitro* translation of the control reporter mRNA that harbors the 50 nt human β-globin 5' UTR and a long 5' UTR reporter mRNA that harbors three β-globin 5' UTRs in

tandem (150 nt total). mRNPs were formed with protein storage buffer or WT NT-hFMRP. Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using a two-tailed unpaired t test with Welch's correction. B) Distribution of nLuc reporter mRNA across sucrose gradients to assess polysome formation when pre-incubated with ΔRGG+CTD (Control) or WT NT-hFMRP. Abundance of reporter mRNA in each gradient fraction was determined by RT-qPCR. C) Cumulative nLuc abundance in heavy polysomes in fractions 10-12 from ΔRGG+CTD (Control) and WT NT-hFMRP samples. D) Fold change of nLuc mRNA abundance in heavy polysomes. Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using a two-tailed unpaired t test with Welch's correction. E) Relative quantification of nLuc reporter mRNA pelleted through a 35% (w/v) sucrose cushion after a low-speed centrifugation (see **Supplemental Figure S9**). Lane 1 is a negative control lacking nLuc mRNA. Lane 2 is a negative control containing mRNA in RRL but not incubated at 30°C to start translation. Lane 3 is nLuc mRNA in RRL translated for 15 min at 30°C. Lane 4 is nLuc mRNA in RRL translated for 15 min at 30°C and then incubated with 0.1 mM puromycin (final) for 30 min at 30°C. Lanes 5 and 6 are negative control untranslated reactions of nLuc•FMRP mRNPs formed with ΔRGG+CTD (Control) and WT NT-hFMRP, respectively, in RRL kept on ice demonstrating poor pelleting without active translation through the low-speed sucrose cushion as described in the Materials and Methods and outlined in **Supplemental Figure S9**). Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using a two-tailed unpaired t test with Welch's correction. F) Relative quantification of nLuc reporter mRNA pelleted through a 35% (w/v) sucrose cushion after a low-speed centrifugation. nLuc•ΔRGG+CTD (Control) and nLuc•WT NT-hFMRP mRNPs were translated and treated with 0.1 mM puromycin (final) before being overlayed on the cushion and low-speed centrifugation. Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using a two-tailed unpaired t test with Welch's correction.

Figure 7. Model of FMRP-mediated translational repression via a ncRBD. FMRP can bind mRNA through canonical and non-canonical RBDs. When bound to mRNA through its canonical KH domain(s), FMRP does not robustly inhibit translation. The RGG box motif and CTD of FMRP together form a ncRBD that allows FMRP to bind multiple mRNAs and subsequently inhibit translation elongation.

SUPPLEMENTAL DATA

Supplemental Table S1. Oligonucleotides used in this study.

Name	Sequence (5′-3′)
F_nLuc RT-qPCR	ATCAAGTGCTCGAACAAGGC
R_nLuc RT-qPCR	CAATTTTCAGGCCGTTTTCGC
F_Control FFLuc RT-qPCR	AACGCTTCCATCTTCCAGGG
R_Control FFLuc RT-qPCR	CCAGATCCACAACCTTCGCT
F_pGL4.13 RT-qPCR	GCAGTACCGGATTGCCCAAG
R_pGL4.13 RT-qPCR	GTCGGGGATGATCTGGTTGC

Supplemental Figure S1. G_{15} and $(GGGU)_4$ reporter mRNA, but not control reporter mRNA, harbor G-quadruplexes. Native PAGE of control, G_{15}, and $(GGGU)_4$ reporters stained for total RNA with SYBR Green II or for G4 structures with NMM.

Supplemental Figure S2. The Sc1 G4 does not enhance translational repression by FMRP. A) Schematic of custom nLuc reporter harboring the Sc1 G4 in the coding sequence. A P2A ribosome skipping motif was included immediately upstream of the nLuc coding sequence to ensure equal nLuc function between reporters. B) *In vitro* translation of Sc1 reporter mRNA with protein storage buffer or WT NT-hFMRP. Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using an unpaired t test with Welch's correction.

Supplemental Figure S3. NT-hFMRP ΔRGG+CTD complete does not inhibit translation. A) Schematic of MBP- and His6-tagged NT-hFMRP ΔRGG+CTD complete. B) Coomassie stain of recombinant NT-hFMRP ΔRGG+CTD complete. C) *In vitro* translation of nLuc control mRNA with protein storage buffer or NT-hFMRP ΔRGG+CTD complete. Data are shown as mean ± SD. n = 3 biological replicates. Comparisons were made using an unpaired t test with Welch's correction.

Supplemental Figure S4. NT-hFMRP I304N & ΔRGG+CTD double mutant. A) Schematic of MBP- and His6-tagged NT-hFMRP I304N & ΔRGG+CTD. Mutated/truncated domains are highlighted in red. B) Coomassie stain of recombinant NT-hFMRP I304N & ΔRGG+CTD.

Supplemental Figure S5. The RGG+CTD ncRBD binds mEGFP mRNA. A) EMSA of mEGFP mRNA with the indicated recombinant protein resolved on a native PAGE and subsequently stained with SYBR Green II. B) Denaturing PAGE of G_{15} reporter and mEGFP mRNAs stained for total RNA with SYBR Green II or for G4 structures with NMM.

Supplemental Figure S6. The FMRP ncRBD is predicted to be flexible and is conserved among most vertebrates. A) IUPred3 analysis (https://iupred.elte.hu/) of the full-length human FMRP. The ncRBD is bracketed. Scores above 0.5 (solid horizontal line) signal predicted

unstructured flexible regions. B) AlphaFold structural prediction of full-length human FMRP with the ncRBD highlighted in magenta. C) Clustal Omega (1.2.4) alignment of vertebrate FMRP orthologs. The human FMRP ncRBD is highlighted in magenta. The residue numbering corresponds to the full-length protein sequence. D) Clustal Omega (1.2.4) alignment of human, zebrafish, and fruit fly FMRP. The human FMRP ncRBD is highlighted in magenta. The residue numbering corresponds to the full-length protein sequence.

Supplemental Figure S7. NT-hFMRP Δ54 and Δ55. A) Schematic of MBP- and His6-tagged NT-hFMRP Δ54 and Δ55. The truncated CTD is highlighted in red. B) Coomassie stain of recombinant NT-hFMRP Δ54 and Δ55.

Supplemental Figure S8. WT NT-hFMRP causes heavy polysomes to accumulate. A) A representative $A_{260\,nm}$ trace of sucrose gradient ultracentrifugation (polysome analysis) of *in vitro* translation reactions. The 80S monosome sediments in fractions 4 and 5. Thus, fractions 6-12 contain polysomes. Robust polysome curves are not visualized as mRNA input amounts were limited to ensure responses were in the linear dynamic range (see Materials and Methods). B-E)

Separate biological replicates of experiments shown in **Figure 6B-D**. B & D) Distribution of reporter mRNA across sucrose gradients to assess polysome formation with ΔRGG+CTD (Control) or WT NT-hFMRP. Abundance of reporter mRNA in each gradient fraction was determined by RT-qPCR. C & E) Cumulative nLuc abundance in heavy polysomes in fractions 10-12 from ΔRGG+CTD (Control) and WT NT-hFMRP samples.

Supplemental Figure S9. Workflow of puromycin-mediated ribosome dissociation and low-speed sucrose cushion assay. WT or mutant NT-hFMRP was allowed to form an mRNP with nLuc mRNA as described in the Material and Methods. FFLuc (pGL4.13) mRNA was translated and used as an internal spike-in control for ribosome-bound mRNA pelleting and RT-qPCR. See details in the Material and Methods.

Reporter Sequences
T7 promoter
Human beta globin 5▢ UTR
Heterologous 5▢ UTR
Start codon
P2A ribosome skipping motif
nLuc
3xFLAG
Stop codon
G4 insert

Control nLuc reporter
*TAATACGACTCACTATAG*GGACATTTGCTTCTGACACAACTGTGTTCACTAGCAACCTCAAA
CAGACACCATGGGAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAAGCTGGAGACGTCG
AAGAAAACCCTGGACCTGTGTTTACCCTTGAAGATTTTGTCGGCGATTGGCGCCAGACCGC
GCGCTATAACCTCGATCAAGTGCTCGAACAAGGCGGCGTGAGCAGCCTGTTTCAGAACCT
GGGCGTGAGCGTGACCCCGATTCAGCGCATTGTGCTGAGCGGCGAAAACGGCCTGAAAA
TTGATATTCATGTGATTATTCCGTATGAAGGCCTGAGCGGCGATCAGATGGGCCAGATTGA
AAAAATTTTTAAAGTCGTATATCCTGTCGATGATCATCATTTTAAAGTGATTCTGCATTATGG
CACCCTGGTGATTGATGGCGTGACCCCGAACATGATTGATTATTTTGGCCGCCCGTATGAA
GGCATTGCTGTTTTTGATGGCAAGAAGATTACCGTGACCGGCACCCTGTGGAACGGCAAC
AAAATTATTGATGAACGCCTGATTAACCCAGATGGCAGCCTGCTGTTTCGCGTGACCATTA
ACGGCGTGACCGGCTGGCGCCTGTGCGAACGCATTCTCGCAGACTACAAAGACCATGAC
GGTGATTATAAAGATCATGACATCGATTACAAGGATGACGATGACAAGTAAGGCCGCGACT
CTAGAG

G₁₅ nLuc reporter
*TAATACGACTCACTATAG*GGACATTTGCTTCTGACACAACTGTGTTCACTAGCAACCTCAAA
CAGACACCATGCAACAACAACAACAAGGGGGGGGGGGGGGGCAACAACAACAACAAGGA
AGCGGAGCTACTAACTTCAGCCTGCTGAAGCAAGCTGGAGACGTCGAAGAAAACCCTGGA
CCTGTGTTTACCCTTGAAGATTTTGTCGGCGATTGGCGCCAGACCGCGCGCTATAACCTCG
ATCAAGTGCTCGAACAAGGCGGCGTGAGCAGCCTGTTTCAGAACCTGGGCGTGAGCGTGA
CCCCGATTCAGCGCATTGTGCTGAGCGGCGAAAACGGCCTGAAAATTGATATTCATGTGAT
TATTCCGTATGAAGGCCTGAGCGGCGATCAGATGGGCCAGATTGAAAAAATTTTTAAAGTC
GTATATCCTGTCGATGATCATCATTTTAAAGTGATTCTGCATTATGGCACCCTGGTGATTGA
TGGCGTGACCCCGAACATGATTGATTATTTTGGCCGCCCGTATGAAGGCATTGCTGTTTTT
GATGGCAAGAAGATTACCGTGACCGGCACCCTGTGGAACGGCAACAAAATTATTGATGAA
CGCCTGATTAACCCAGATGGCAGCCTGCTGTTTCGCGTGACCATTAACGGCGTGACCGGC
TGGCGCCTGTGCGAACGCATTCTCGCAGACTACAAAGACCATGACGGTGATTATAAAGATC
ATGACATCGATTACAAGGATGACGATGACAAGTAAGGCCGCGACTCTAGAG

(GGGU)₄ nLuc reporter
*TAATACGACTCACTATAG*GGACATTTGCTTCTGACACAACTGTGTTCACTAGCAACCTCAAA
CAGACACCATGCAACAACAACAACAAGGGTGGGTGGGTGGGTTTCAACAACAACAACAAG
GAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAAGCTGGAGACGTCGAAGAAAACCCTG
GACCTGTGTTTACCCTTGAAGATTTTGTCGGCGATTGGCGCCAGACCGCGCGCTATAACCT
CGATCAAGTGCTCGAACAAGGCGGCGTGAGCAGCCTGTTTCAGAACCTGGGCGTGAGCG
TGACCCCGATTCAGCGCATTGTGCTGAGCGGCGAAAACGGCCTGAAAATTGATATTCATGT
GATTATTCCGTATGAAGGCCTGAGCGGCGATCAGATGGGCCAGATTGAAAAAATTTTTAAA
GTCGTATATCCTGTCGATGATCATCATTTTAAAGTGATTCTGCATTATGGCACCCTGGTGAT
TGATGGCGTGACCCCGAACATGATTGATTATTTTGGCCGCCCGTATGAAGGCATTGCTGTT

TTTGATGGCAAGAAGATTACCGTGACCGGCACCCTGTGGAACGGCAACAAAATTATTGATG
AACGCCTGATTAACCCAGATGGCAGCCTGCTGTTTCGCGTGACCATTAACGGCGTGACCG
GCTGGCGCCTGTGCGAACGCATTCTCGCAGACTACAAAGACCATGACGGTGATTATAAAG
ATCATGACATCGATTACAAGGATGACGATGACAAGTAAGGCCGCGACTCTAGAG

Sc1 nLuc reporter

*TAATACGACTCACTATAG*GGACATTTGCTTCTGACACAACTGTGTTCACTAGCAACCTCAAA
CAGACACCATGCAACAACAACAACAAGCTGCGGTGTGGAAGGAGTGGTCGGGGTTGCGCA
GCGCAACAACAACAACAAGGAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAAGCTGGA
GACGTCGAAGAAAACCCTGGACCTGTGTTTACCCTTGAAGATTTTGTCGGCGATTGGCGCC
AGACCGCGCGCTATAACCTCGATCAAGTGCTCGAACAAGGCGGCGTGAGCAGCCTGTTTC
AGAACCTGGGCGTGAGCGTGACCCCGATTCAGCGCATTGTGCTGAGCGGCGAAAACGGC
CTGAAAATTGATATTCATGTGATTATTCCGTATGAAGGCCTGAGCGGCGATCAGATGGGCC
AGATTGAAAAAATTTTTAAAGTCGTATATCCTGTCGATGATCATCATTTTAAAGTGATTCTGC
ATTATGGCACCCTGGTGATTGATGGCGTGACCCCGAACATGATTGATTATTTTGGCCGCCC
GTATGAAGGCATTGCTGTTTTTGATGGCAAGAAGATTACCGTGACCGGCACCCTGTGGAAC
GGCAACAAAATTATTGATGAACGCCTGATTAACCCAGATGGCAGCCTGCTGTTTCGCGTGA
CCATTAACGGCGTGACCGGCTGGCGCCTGTGCGAACGCATTCTCGCAGACTACAAAGACC
ATGACGGTGATTATAAAGATCATGACATCGATTACAAGGATGACGATGACAAGTAAGGCCG
CGACTCTAGAG

Long 5▯ UTR nLuc reporter

*TAATACGACTCACTATAG*GGACATTTGCTTCTGACACAACTGTGTTCACTAGCAACCTCAAA
CAGACACCACATTTGCTTCTGACACAACTGTGTTCACTAGCAACCTCAAACAGACACCACA
TTTGCTTCTGACACAACTGTGTTCACTAGCAACCTCAAACAGACACCATGGGAAGCGGAGC
TACTAACTTCAGCCTGCTGAAGCAAGCTGGAGACGTCGAAGAAAACCCTGGACCTGTGTTT
ACCCTTGAAGATTTTGTCGGCGATTGGCGCCAGACCGCGCGCTATAACCTCGATCAAGTG
CTCGAACAAGGCGGCGTGAGCAGCCTGTTTCAGAACCTGGGCGTGAGCGTGACCCCGAT
TCAGCGCATTGTGCTGAGCGGCGAAAACGGCCTGAAAATTGATATTCATGTGATTATTCCG
TATGAAGGCCTGAGCGGCGATCAGATGGGCCAGATTGAAAAAATTTTTAAAGTCGTATATC
CTGTCGATGATCATCATTTTAAAGTGATTCTGCATTATGGCACCCTGGTGATTGATGGCGTG
ACCCCGAACATGATTGATTATTTTGGCCGCCCGTATGAAGGCATTGCTGTTTTTGATGGCA
AGAAGATTACCGTGACCGGCACCCTGTGGAACGGCAACAAAATTATTGATGAACGCCTGAT
TAACCCAGATGGCAGCCTGCTGTTTCGCGTGACCATTAACGGCGTGACCGGCTGGCGCCT
GTGCGAACGCATTCTCGCAGACTACAAAGACCATGACGGTGATTATAAAGATCATGACATC
GATTACAAGGATGACGATGACAAGTAAGGCCGCGACTCTAGAG

mEGFP reporter

*TAATACGACTCACTATAG*GGAGACCCAAGCTGGCTAGCGTTTAAACTTAAGCTTGGCAATC
CGGTACTGTTGGTAAATAAGCCACCATGGTGAGCAAGGGCGAGGAGCTGTTCACCGGGGT
GGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCG
GCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACC
GGCAAGCTGCCCGTGCCCTGGCCCACCCTCGTGACCACCCTGACCTACGGCGTGCAGTG
CTTCAGCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTCCGCCATGCCCGA
AGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGC
CGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACT
TCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACTACAACAGCCACAACG
TCTATATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAAGATCCGCCACAA
CATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCG
ACGGCCCCGTGCTGCTGCCCGACAACCACTACCTGAGCACCCAGTCCAAGCTGAGCAAAG
ACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGGGATC

ACTCTCGGCATGGACGAGCTGTACAAG`TAA`GGCCGCGACTCTAGAG

Recombinant Protein Coding Sequences
`Start codon`
`MBP`
`His6`
`NT-hFMRP (WT and mutant)`
`Stop codon`

His6-MBP
`ATG`GGTTCTTCT`CACCATCACCATCACCAT`GGTTCTTCTATGAAAATCGAAGAAGGTAAACT
GGTAATCTGGATTAACGGCGATAAAGGCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTC
GAGAAAGATACCGGAATTAAAGTCACCGTTGAGCATCCGGATAAACTGGAAGAGAAATTCC
CACAGGTTGCGGCAACTGGCGATGGCCCTGACATTATCTTCTGGGCACACGACCGCTTTG
GTGGCTACGCTCAATCTGGCCTGTTGGCTGAAATCACCCCGGACAAAGCGTTCCAGGACA
AGCTGTATCCGTTTACCTGGGATGCCGTACGTTACAACGGCAAGCTGATTGCTTACCCGAT
CGCTGTTGAAGCGTTATCGCTGATTTATAACAAAGATCTGCTGCCGAACCCGCCAAAAACC
TGGGAAGAGATCCCGGCGCTGGATAAAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGAT
GTTCAACCTGCAAGAACCGTACTTCACCTGGCCGCTGATTGCTGCTGACGGGGGTTATGC
GTTCAAGTATGAAAACGGCAAGTACGACATTAAAGACGTGGGCGTGGATAACGCTGGCGC
GAAAGCGGGTCTGACCTTCCTGGTTGACCTGATTAAAAACAAACACATGAATGCAGACACC
GATTACTCCATCGCAGAAGCTGCCTTTAATAAAGGCGAAACAGCGATGACCATCAACGGCC
CGTGGGCATGGTCCAACATCGACACCAGCAAAGTGAATTATGGTGTAACGGTACTGCCGA
CCTTCAAGGGTCAACCATCCAAACCGTTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCG
CCAGTCCGAACAAAGAGCTGGCAAAAGAGTTCCTCGAAAACTATCTGCTGACTGATGAAGG
TCTGGAAGCGGTTAATAAAGACAAACCGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGA
AGAGTTGGCGAAAGATCCACGTATTGCCGCCACTATGGAAAACGCCCAGAAAGGTGAAAT
CATGCCGAACATCCCGCAGATGTCCGCTTTCTGGTATGCCGTGCGTACTGCGGTGATCAA
CGCCGCCAGCGGTCGTCAGACTGTCGATGAAGCCCTGAAAGACGCGCAGACTAATGGGA
TCGAGGAAAACCTGTACTTCCAATCCAATATTGGAAGTGGA`TAA`

MBP-(WT NT-hFMRP)-His6
`ATG`GGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG

CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGCTTCCACGAACAATTTATTGTTCGCGAGGACCTGATGGGCTTGGCGATTGGCAC
CCACGGCGCGAACATTCAGCAGGCGCGTAAAGTCCCAGGTGTTACCGCTATTGATCTGGA
CGAAGACACGTGCACCTTTCATATTTATGGTGAAGACCAGGATGCTGTGAAAAAGGCGCGT
TCCTTCCTGGAATTTGCGGAAGACGTGATCCAGGTTCCGCGTAACCTGGTTGGTAAAGTCA
TCGGCAAAAACGGTAAGTTAATCCAAGAGATCGTGGACAAAAGCGGGGGTAGTTCGCGTTC
GGATTGAAGCGGAAAATGAGAAGAACGTTCCGCAGGAGGAAGAAATTATGCCGCCAAATA
GCCTGCCGAGCAACAACTCACGTGTCGGTCCGAACGCTCCGGAAGAGAAGAAGCACCTG
GATATTAAAGAGAACAGCACCCATTTCAGCCAACCAAACTCCACTAAGGTGCAGCGTGTTT
TGGTAGCCAGCTCCGTTGTTGCCGGTGAGTCGCAAAAGCCGGAACTGAAAGCGTGGCAG
GGTATGGTGCCGTTCGTCTTTGTGGGCACCAAGGACAGCATCGCCAACGCAACGGTTCTG
CTGGACTACCATCTGAATTACCTGAAAGAAGTCGATCAGCTTCGTTTGGAACGCTTGCAAA
TCGATGAGCAACTGCGCCAGATCGGTGCGAGCTCTCGCCCGTCTCCGAACCGTACCGACA
AAGAGAAGAGCTACGTGACCGACGACGGTCAAGGTATGGGCCGTGGCAGCCGTCCGTAT
CGTAATAGGGGACATGGCCGCCGTGGTCCGGGTTACACCTCAGGTACGAACTCTGAGGC
GTCTAACGCCTCCGAAACCGAGTCGGATCATCGTGATGAGCTGAGCGACTGGTCACTGGC
GCCGACCGAAGAGGAGCGCGAGAGCTTTCTGCGTCGCGGTGATGGTCGCCGCAGAGGC
GGTGGCGGTCGTGGCCAGGGCGGCCGTGGCAGAGGCGGCGGTTTTAAAGGTAATGATGA
TCACAGCCGCACTGACAACCGTCCGCGTAATCCGCGTGAGGCGAAGGGCAGAACTACAG
ATGGTAGCTTGCAAATCCGTGTGGACTGTAATAACGAACGCAGCGTGCATACGAAAACCCT
ACAAAACACCAGTAGCGAGGGTAGCCGCCTTCGTACCGGTAAAGACCGCAACCAGAAGAA
AGAGAAACCGGACAGCGTTGATGGCCAACAACCGTTGGTGAATGGTGTTCCGGGTTCTTC
TCACCATCACCATCACCATTAA

MBP-(I304N NT-hFMRP)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGCTTCCACGAACAATTTATTGTTCGCGAGGACCTGATGGGCTTGGCGATTGGCAC
CCACGGCGCGAACATTCAGCAGGCGCGTAAAGTCCCAGGTGTTACCGCTATTGATCTGGA
CGAAGACACGTGCACCTTTCATATTTATGGTGAAGACCAGGATGCTGTGAAAAAGGCGCGT
TCCTTCCTGGAATTTGCGGAAGACGTGATCCAGGTTCCGCGTAACCTGGTTGGTAAAGTCA
TCGGCAAAAACGGTAAGTTAAACCAAGAGATCGTGGACAAAAGCGGGGGTAGTTCGCGTTC

MBP-(ΔRGG+CTD NT-hFMRP)-His6

CGTAATAGGGGACATGGCCGCCGTGGTCCGGGTTACACCTCAGGTACGAACTCTGAGGC
GTCTAACGCCTCCGAAACCGAGTCGGATCATCGTGATGAGCTGAGCGACTGGTCACTGGC
GCCGACCGAAGAGGAGCGCGAGAGCTTTCTGCGTCGCGGTGATGGTCGCCGCAGAGGC
GGTGGCGGGTCGTGGCCAGGGCGGCCGTGGCAGAGGCGGCGGTTTGGTTCTTCTCACCA
TCACCATCACCATTAA

MBP-(NT-hFMRP ΔRGG+CTD complete)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGCTTCCACGAACAATTTATTGTTCGCGAGGACCTGATGGGCTTGGCGATTGGCAC
CCACGGCGCGAACATTCAGCAGGCGCGTAAAGTCCCAGGTGTTACCGCTATTGATCTGGA
CGAAGACACGTGCACCTTTCATATTTATGGTGAAGACCAGGATGCTGTGAAAAAGGCGCGT
TCCTTCCTGGAATTTGCGGAAGACGTGATCCAGGTTCCGCGTAACCTGGTTGGTAAAGTCA
TCGGCAAAAACGGTAAGTTAATCCAAGAGATCGTGGACAAAAGCGGGGTAGTTCGCGTTC
GGATTGAAGCGGAAAATGAGAAGAACGTTCCGCAGGAGGAAGAAATTATGCCGCCAAATA
GCCTGCCGAGCAACAACTCACGTGTCGGTCCGAACGCTCCGGAAGAGAAGAAGCACCTG
GATATTAAAGAGAACAGCACCCATTTCAGCCAACCAAACTCCACTAAGGTGCAGCGTGTTT
TGGTAGCCAGCTCCGTTGTTGCCGGTGAGTCGCAAAAGCCGGAACTGAAAGCGTGGCAG
GGTATGGTGCCGTTCGTCTTTGTGGGCACCAAGGACAGCATCGCCAACGCAACGGTTCTG
CTGGACTACCATCTGAATTACCTGAAAGAAGTCGATCAGCTTCGTTTGGAACGCTTGCAAA
TCGATGAGCAACTGCGCCAGATCGGTGCGAGCTCTCGCCCGTCTCCGAACCGTACCGACA
AAGAGAAGAGCTACGTGACCGACGACGGTCAAGGTATGGGCCGTGGCAGCCGTCCGTAT
CGTAATAGGGGACATGGCCGCCGTGGTCCGGGTTACACCTCAGGTACGAACTCTGAGGC
GTCTAACGCCTCCGAAACCGAGTCGGATCATCGTGATGAGCTGAGCGACTGGTCACTGGC
GCCGACCGAAGAGGAGCGCGAGAGCTTTCTGGGTTCTTCTCACCATCACCATCACCATTA
A

MBP-(I304N & ΔRGG+CTD NT-hFMRP)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT

ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCAGCTTCCACGAACAATTTATTGTTCGCGAGGACCTGATGGGCTTGGCGATTGGCACC
CACGGCGCGAACATTCAGCAGGCGCGTAAAGTCCCAGGTGTTACCGCTATTGATCTGGAC
GAAGACACGTGCACCTTTCATATTTATGGTGAAGACCAGGATGCTGTGAAAAAGGCGCGTT
CCTTCCTGGAATTTGCGGAAGACGTGATCCAGGTTCCGCGTAACCTGGTTGGTAAAGTCAT
CGGCAAAAACGGTAAGTTAAACCAAGAGATCGTGGACAAAGCGGGGTAGTTCGCGTTCG
GATTGAAGCGGAAAATGAGAAGAACGTTCCGCAGGAGGAAGAAATTATGCCGCCAAATAG
CCTGCCGAGCAACAACTCACGTGTCGGTCCGAACGCTCCGGAAGAGAAGAAGCACCTGG
ATATTAAAGAGAACAGCACCCATTTCAGCCAACCAAACTCCACTAAGGTGCAGCGTGTTTT
GGTAGCCAGCTCCGTTGTTGCCGGTGAGTCGCAAAAGCCGGAACTGAAAGCGTGGCAGG
GTATGGTGCCGTTCGTCTTTGTGGGCACCAAGGACAGCATCGCCAACGCAACGGTTCTGC
TGGACTACCATCTGAATTACCTGAAAGAAGTCGATCAGCTTCGTTTGGAACGCTTGCAAAT
CGATGAGCAACTGCGCCAGATCGGTGCGAGCTCTCGCCCGTCTCCGAACCGTACCGACAA
AGAGAAGAGCTACGTGACCGACGACGGTCAAGGTATGGGCCGTGGCAGCCGTCCGTATC
GTAATAGGGGACATGGCCGCCGTGGTCCGGGTTACACCTCAGGTACGAACTCTGAGGCGT
CTAACGCCTCCGAAACCGAGTCGGATCATCGTGATGAGCTGAGCGACTGGTCACTGGCGC
CGACCGAAGAGGAGCGCGCGAGAGCTTTCTGCGTCGCGGTGATGGTCGCCGCAGAGGCGGT
GGCGGGTCGTGGCCAGGGCGGCCGTGGCAGAGGCGGCGGTTTGGTTCTTCTCACCATCA
CCATCACCATTAA

MBP-(NT-hFMRP Δ54)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG

ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGCTTCCACGAACAATTTATTGTTCGCGAGGACCTGATGGGCTTGGCGATTGGCAC
CCACGGCGCGAACATTCAGCAGGCGCGTAAAGTCCCAGGTGTTACCGCTATTGATCTGGA
CGAAGACACGTGCACCTTTCATATTTATGGTGAAGACCAGGATGCTGTGAAAAAGGCGCGT
TCCTTCCTGGAATTTGCGGAAGACGTGATCCAGGTTCCGCGTAACCTGGTTGGTAAAGTCA
TCGGCAAAAACGGTAAGTTAATCCAAGAGATCGTGGACAAAAGCGGGGGTAGTTCGCGTTC
GGATTGAAGCGGAAAATGAGAAGAACGTTCCGCAGGAGGAAGAAATTATGCCGCCAAATA
GCCTGCCGAGCAACAACTCACGTGTCGGTCCGAACGCTCCGGAAGAGAAGAAGCACCTG
GATATTAAAGAGAACAGCACCCATTTCAGCCAACCAAACTCCACTAAGGTGCAGCGTGTTT
TGGTAGCCAGCTCCGTTGTTGCCGGTGAGTCGCAAAAGCCGGAACTGAAAGCGTGGCAG
GGTATGGTGCCGTTCGTCTTTGTGGGCACCAAGGACAGCATCGCCAACGCAACGGTTCTG
CTGGACTACCATCTGAATTACCTGAAAGAAGTCGATCAGCTTCGTTTGGAACGCTTGCAAA
TCGATGAGCAACTGCGCCAGATCGGTGCGAGCTCTCGCCCGTCTCCGAACCGTACCGACA
AAGAGAAGAGCTACGTGACCGACGACGGTCAAGGTATGGGCCGTGGCAGCCGTCCGTAT
CGTAATAGGGGACATGGCCGCCGTGGTCCGGGTTACACCTCAGGTACGAACTCTGAGGC
GTCTAACGCCTCCGAAACCGAGTCGGATCATCGTGATGAGCTGAGCGACTGGTCACTGGC
GCCGACCGAAGAGGAGCGCGAGAGCTTTCTGCGTCGCGGTGATGGTCGCCGCAGAGGC
GGTGGCGGTCGTGGCCAGGGCGGCCGTGGCAGAGGCGGCGGTTTTAAAGGTAATGATGA
TCACAGCCGCACTGACAACCGTCCGCGTAATCCGCGTGAGGCGAAGGGCAGAACTACAG
ATGGTAGCTTGGGTTCTTCTCACCATCACCATCACCATTAA

MBP-(NT-hFMRP Δ55)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGCTTCCACGAACAATTTATTGTTCGCGAGGACCTGATGGGCTTGGCGATTGGCAC
CCACGGCGCGAACATTCAGCAGGCGCGTAAAGTCCCAGGTGTTACCGCTATTGATCTGGA
CGAAGACACGTGCACCTTTCATATTTATGGTGAAGACCAGGATGCTGTGAAAAAGGCGCGT
TCCTTCCTGGAATTTGCGGAAGACGTGATCCAGGTTCCGCGTAACCTGGTTGGTAAAGTCA
TCGGCAAAAACGGTAAGTTAATCCAAGAGATCGTGGACAAAAGCGGGGGTAGTTCGCGTTC
GGATTGAAGCGGAAAATGAGAAGAACGTTCCGCAGGAGGAAGAAATTATGCCGCCAAATA
GCCTGCCGAGCAACAACTCACGTGTCGGTCCGAACGCTCCGGAAGAGAAGAAGCACCTG
GATATTAAAGAGAACAGCACCCATTTCAGCCAACCAAACTCCACTAAGGTGCAGCGTGTTT
TGGTAGCCAGCTCCGTTGTTGCCGGTGAGTCGCAAAAGCCGGAACTGAAAGCGTGGCAG

GGTATGGTGCCGTTCGTCTTTGTGGGCACCAAGGACAGCATCGCCAACGCAACGGTTCTG
CTGGACTACCATCTGAATTACCTGAAAGAAGTCGATCAGCTTCGTTTGGAACGCTTGCAAA
TCGATGAGCAACTGCGCCAGATCGGTGCGAGCTCTCGCCCGTCTCCGAACCGTACCGACA
AAGAGAAGAGCTACGTGACCGACGACGGTCAAGGTATGGGCCGTGGCAGCCGTCCGTAT
CGTAATAGGGGACATGGCCGCCGTGGTCCGGGTTACACCTCAGGTACGAACTCTGAGGC
GTCTAACGCCTCCGAAACCGAGTCGGATCATCGTGATGAGCTGAGCGACTGGTCACTGGC
GCCGACCGAAGAGGAGCGCGAGAGCTTTCTGCGTCGCGGTGATGGTCGCCGCAGAGGC
GGTGGCGGTCGTGGCCAGGGCGGCCGTGGCAGAGGCGGCGGTTTTAAAGGTAATGATGA
TCACAGCCGCACTGACAACCGTCCGCGTAATCCGCGTGAGGCGAAGGGCAGAACTACAG
ATGGTAGCGGTTCTTCTCACCATCACCATCACCATTAA

MBP-(RGG+CTD)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGTCGCGGTGATGGTCGCCGCAGAGGCGGTGGCGGTCGTGGCCAGGGCGGCCGT
GGCAGAGGCGGCGGTTTTAAAGGTAATGATGATCACAGCCGCACTGACAACCGTCCGCGT
AATCCGCGTGAGGCGAAGGGCAGAACTACAGATGGTAGCTTGCAAATCCGTGTGGACTGT
AATAACGAACGCAGCGTGCATACGAAAACCCTACAAAACACCAGTAGCGAGGGTAGCCGC
CTTCGTACCGGTAAAGACCGCAACCAGAAGAAAGAGAAACCGGACAGCGTTGATGGCCAA
CAACCGTTGGTGAATGGTGTTCCGGGTTCTTCTCACCATCACCATCACCATTAA

MBP-(RGG)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT

TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGTCGCGGTGATGGTCGCCGCAGAGGCGGTGGCGGTCGTGGCCAGGGCGGCCGT
GGCAGAGGCGGCGGTTTTTAAAGGTAATGATGATCACAGCCGCGGTTCTTCTCACCATCAC
CATCACCATTAA

MBP-(CTD)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCAACTGACAACCGTCCGCGTAATCCGCGTGAGGCGAAGGGCAGAACTACAGATGGTAG
CTTGCAAATCCGTGTGGACTGTAATAACGAACGCAGCGTGCATACGAAAACCCTACAAAAC
ACCAGTAGCGAGGGTAGCCGCCTTCGTACCGGTAAAGACCGCAACCAGAAGAAAGAGAAA
CCGGACAGCGTTGATGGCCAACAACCGTTGGTGAATGGTGTTCCGGGTTCTTCTCACCAT
CACCATCACCATTAA

MBP-(RGG+CTD Δ19)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC

CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGTCGCGGTGATGGTCGCCGCAGAGGCGGTGGCGGTCGTGGCCAGGGCGGCCGT
GGCAGAGGCGGCGGTTTTAAAGGTAATGATGATCACAGCCGCACTGACAACCGTCCGCGT
AATCCGCGTGAGGCGAAGGGCAGAACTACAGATGGTAGCTTGCAAATCCGTGTGGACTGT
AATAACGAACGCAGCGTGCATACGAAAACCCTACAAAACACCAGTAGCGAGGGTAGCCGC
CTTCGTACCGGTAAAGACCGCAACCAGGGTTCTTCTCACCATCACCATCACCATTAA

MBP-(RGG+CTD Δ54)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGTCGCGGTGATGGTCGCCGCAGAGGCGGTGGCGGTCGTGGCCAGGGCGGCCGT
GGCAGAGGCGGCGGTTTTAAAGGTAATGATGATCACAGCCGCACTGACAACCGTCCGCGT
AATCCGCGTGAGGCGAAGGGCAGAACTACAGATGGTAGCTTGGGTTCTTCTCACCATCAC
CATCACCATTAA

MBP-(RGG+CTD Δ55)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT

TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGTCGCGGTGATGGTCGCCGCAGAGGCGGTGGCGGTCGTGGCCAGGGCGGCCGT
GGCAGAGGCGGCGGTTTTTAAAGGTAATGATGATCACAGCCGCACTGACAACCGTCCGCGT
AATCCGCGTGAGGCGAAGGGGCAGAACTACAGATGGTAGCGGTTCTTCTCACCATCACCAT
CACCATTAA

MBP-(RGG+CTD Δ62)-His6

ATGGGTTCTTCTATGAAAATCGAAGAAGGTAAACTGGTAATCTGGATTAACGGCGATAAAG
GCTATAACGGTCTCGCTGAAGTCGGTAAGAAATTCGAGAAAGATACCGGAATTAAAGTCAC
CGTTGAGCATCCGGATAAACTGGAAGAGAAATTCCCACAGGTTGCGGCAACTGGCGATGG
CCCTGACATTATCTTCTGGGCACACGACCGCTTTGGTGGCTACGCTCAATCTGGCCTGTTG
GCTGAAATCACCCCGGACAAAGCGTTCCAGGACAAGCTGTATCCGTTTACCTGGGATGCC
GTACGTTACAACGGCAAGCTGATTGCTTACCCGATCGCTGTTGAAGCGTTATCGCTGATTT
ATAACAAAGATCTGCTGCCGAACCCGCCAAAAACCTGGGAAGAGATCCCGGCGCTGGATA
AAGAACTGAAAGCGAAAGGTAAGAGCGCGCTGATGTTCAACCTGCAAGAACCGTACTTCA
CCTGGCCGCTGATTGCTGCTGACGGGGGTTATGCGTTCAAGTATGAAAACGGCAAGTACG
ACATTAAAGACGTGGGCGTGGATAACGCTGGCGCGAAAGCGGGTCTGACCTTCCTGGTTG
ACCTGATTAAAAACAAACACATGAATGCAGACACCGATTACTCCATCGCAGAAGCTGCCTT
TAATAAAGGCGAAACAGCGATGACCATCAACGGCCCGTGGGCATGGTCCAACATCGACAC
CAGCAAAGTGAATTATGGTGTAACGGTACTGCCGACCTTCAAGGGTCAACCATCCAAACCG
TTCGTTGGCGTGCTGAGCGCAGGTATTAACGCCGCCAGTCCGAACAAAGAGCTGGCAAAA
GAGTTCCTCGAAAACTATCTGCTGACTGATGAAGGTCTGGAAGCGGTTAATAAAGACAAAC
CGCTGGGTGCCGTAGCGCTGAAGTCTTACGAGGAAGAGTTGGCGAAAGATCCACGTATTG
CCGCCACTATGGAAAACGCCCAGAAAGGTGAAATCATGCCGAACATCCCGCAGATGTCCG
CTTTCTGGTATGCCGTGCGTACTGCGGTGATCAACGCCGCCAGCGGTCGTCAGACTGTCG
ATGAAGCCCTGAAAGACGCGCAGACTAATGGGATCGAGGAAAACCTGTACTTCCAATCCAA
TGCACGTCGCGGTGATGGTCGCCGCAGAGGCGGTGGCGGTCGTGGCCAGGGCGGCCGT
GGCAGAGGCGGCGGTTTTTAAAGGTAATGATGATCACAGCCGCACTGACAACCGTCCGCGT
AATCCGCGTGAGGCGAAGGGTTCTTCTCACCATCACCATCACCATTAA