1

cdon and _boc_ affect trunk neural crest cell migration non-cell autonomously through a

reduction of hedgehog signaling in zebrafish slow-twitch muscle

Ezra Lencer PhD[1,2,3]

Rytis Prekeris PhD[2,4]

Kristin Artinger PhD[3,4]

1. Current Address: Department of Biology, Lafayette College, United States

2. Department of Cell and Developmental Biology, University of Colorado, United States

3. Department of Craniofacial Biology, University of Colorado, United States

4. Co-corresponding authors

Abstract

The transmembrane proteins *cdon* and *boc* are implicated in regulating hedgehog signaling during vertebrate development. Recent work showing roles for these genes in axon guidance and neural crest cell migration suggest that *cdon*/*boc* may play additional functions in regulating directed cell movements. We use novel and existing mutants to investigate a role for *cdon* and *boc* in zebrafish neural crest cell migration. We find that single mutant embryos exhibit normal neural crest phenotypes, but that neural crest migration is strikingly disrupted in double *cdon*;*boc* mutant embryos. We further show that this migration phenotype is associated with defects to the differentiation of slow-twitch muscle cells, and the loss of a Col1a1a containing extracellular matrix, suggesting that neural crest defects are a secondary consequence to defects in mesoderm development. Combined, our data add to a growing literature showing that *cdon* and *boc* act synergistically to promote hedgehog signaling during vertebrate development, and provide a foundation for using zebrafish to study the function of hedgehog receptor paralogs.

Introduction

During vertebrate development, neural crest cells (NCCs) migrate ventrally from the dorsal neural tube along conserved pathways in the embryo to contribute to numerous traits such as the craniofacial skeleton, peripheral nervous system, and pigment (Theveneau and Mayor, 2012). Defects in NCC migration result in multiple human hereditary disorders (Tobin et al., 2008; Vega-Lopez et al., 2018), and modifications to NCC migration is associated with the evolution of morphological diversity among some vertebrate taxa (Powder et al., 2014). Further, diseases, such as metastatic melanoma, have been shown to progress through a reactivation of a NCC-like migratory genetic program (Diener and Sommer, 2021; Kaufman et al., 2016). Thus, understanding the genetic regulation of NCC migration is fundamental to understanding the

etiology to multiple human diseases as well as a major developmental event that occurs in

every vertebrate embryo.

NCC migration is highly coordinated along conserved routes in the embryo (Mayor and

Etienne-Manneville, 2016), and is regulated by both NCC intrinsic factors, such as genes

regulating cell adhesion and actin cytoskeleton (Clay and Halloran, 2011, 2013, 2014; Williams

et al., 2018), as well as cell extrinsic factors such as proteins in the extracellular matrix (Perris

and Perissinott, 2000; Szabo et al., 2016), tissue stiffness (Shellard and Mayor, 2021), and

chemical gradients (Barriga et al., 2013; Olesnicky Killian et al., 2009; Shellard et al., 2018) that

funnel NCCs along the migratory route. Mediating cell intrinsic and extrinsic regulation of NCC

migration are cell signaling interactions (Carmona-Fontaine et al., 2008; Mayor and Etienne-

Manneville, 2016; Scarpa et al., 2015; Szabo et al., 2016; Theveneau and Mayor, 2012). These

include the chemical gradients that attract and inhibit migrating NCCs, as well as adhesion

molecules that adhere NCCs to each other and mediate cell-cell interactions such as contact

inhibition of locomotion (Carmona-Fontaine et al., 2008; Mayor and Etienne-Manneville, 2016;

Scarpa et al., 2015; Szabo et al., 2016; Theveneau and Mayor, 2012; Vega-Lopez et al., 2017).

Work has implicated a role for the immunoglobin superfamily member protein *cell*

adhesion associated, oncogene regulated (*cdon*) as a cell surface protein regulating trunk NCC

(tNCC) migration in zebrafish (Powell et al., 2015). Morpholino knockdown of *cdon* results in

arrested (tNCC) migration, with tNCCs mislocalizing N-cadherin (Cdh2). *cdon* and its paralog

boc, brother of cell adhesion associated, oncogene regulated (*boc*), are vertebrate members of

a larger gene family with identifiable orthologs in most metazoan taxa including the *Drosophila*

paralogs *ihog* and *boi*. Recent work has established a role for these proteins in activating

hedgehog signaling. *cdon* and *boc* act as co-receptors binding to hedgehog ligand and patched

receptor to promote smoothened activity. Other work suggests that *cdon*/*boc* may also interact

with Wnt and Nodal signaling pathways (Hong et al., 2020; Jeong et al., 2017). *cdon* has been

shown to form a complex with N-cadherin/Cdc42/Bnip-2 in a hedgehog independent manner to

regulate actin cytoskeletal dynamics during myogenic differentiation in cell culture (Kang et al.,

2008; Lu and Krauss, 2010). *boc* has been shown to play a role in axon guidance neuron

growth (Connor et al., 2005; Okada et al., 2006). Both proteins localize to the ends of long

membrane extensions (e.g. cytonemes), suggesting that these receptors help cells sense the

external environment and communicate over long distances (Bilioni et al., 2013; Ferent et al.,

2019; Hall et al., 2021). We hypothesize that these transmembrane proteins are reiteratively

used as cell surface proteins to integrate multiple signals during processes such as cell

migration and axon guidance. While a role for *cdon* in tNCC migration has been suggested

(Powell et al., 2015), the mechanism by which *cdon*/*boc* regulates NCC movements is not fully

understood.

In zebrafish, hedgehog signaling is thought to affect tNCC migration in a non-cell

autonomous manner by regulating the differentiation of adaxial mesoderm (Honjo and Eisen,

2005). Adaxial cells form at the border of the notochord during gastrulation and differentiate into

slow-twitch muscle cells that migrate laterally through the somites (Elworthy et al., 2008; Wolff

et al., 2003). A non-migratory population of adaxial cells, called muscle pioneers, remain at the

body midline. In hedgehog mutant embryos, adaxial cells fail to differentiate into slow-twitch

muscle/muscle pioneer derivatives (Elworthy et al., 2008; Roy et al., 2001; Yin et al., 2018).

Zebrafish embryos with slow-twitch muscle defects also exhibit tNCC migration defects. Though

the mechanisms of how these two tissues interact is not fully understood, it is suggested to

depend on extracellular matrix deposition/modification (Banerjee et al., 2011; Banerjee et al.,

2013; Guillon et al., 2016; Honjo and Eisen, 2005). Whether *cdon*/*boc* affect slow-twitch muscle

differentiation is unknown (though see (Bergeron et al., 2011)), and it is possible that *cdon*/*boc*

affect NCC development through autonomous mechanism (Powell et al., 2015).

We use novel CRISPR/Cas9 generated mutations and existing mutants to investigate

the role of *cdon* and *boc* in tNCC migration and slow-twitch muscle differentiation. We show that

cdon and *boc* are expressed by tNCCs and adaxial mesoderm. Loss of either *cdon* or *boc*

individually has no observable effect on tNCC migration and minimal effects on slow-twitch

muscle differentiation. However, embryos that are double mutant for *cdon* and *boc* exhibit both

tNCC migratory defects and loss of slow-twitch muscle. We show that tNCC mobility is not

impaired by loss of *cdon*/*boc* suggesting that the tNCC migration defect is a non-cell

autonomous consequence of loss of the migratory path. We further provide data suggesting that

cdon/*boc* are modifying hedgehog signal reception during mesoderm differentiation. These data

contribute to a growing literature on the critical role of *cdon*/*boc* in modulating hedgehog

signaling across multiple tissues during embryogenesis.

Results

cdon/boc paralogs exhibit shared sequence conservation in extracellular domains

105   *cdon* and *boc*, share a similar genomic domain architecture that includes 4

106   immunoglobin (IG) domains, 3 fibronectin type 3 repeat (FNIII) domains, and a cytosolic C-

107   terminal tail (**Figure 1A**). A fifth IG domain is variable present in *cdon* orthologs of tetrapod taxa

108   (grey in **Figure 1A**). The FNIII domains have been implicated in binding to hedgehog ligand and

109   patched receptor (Izzi et al., 2011; Song et al., 2015), and evidence suggests that the c-terminal

110   region is needed for protein localization in the cell (Okumura et al., 2021).

111   As functional conservation is likely to be reflected by sequence conservation across

evolutionary time, we investigated the molecular evolution of *cdon*/*boc* orthologs across

vertebrate taxa. An alignment of *cdon* and *boc* amino acid sequences from 11 select jawed

vertebrate taxa shows that extracellular domains exhibit high conservation across taxa and

paralogs, suggesting that these extracellular domains are critical for gene function (**Figure 1A**).

In contrast, the C-terminal cytosolic tail exhibits relatively low sequence conservation (Kang et

al., 2002) suggesting that cytosolic region is evolutionarily labile.

118   Identifiable orthologs to *cdon* and *boc* are present in non-vertebrate bilaterian taxa.

These include *Drosophila* paralogs *ihog* and *boi*, whose function in hedgehog signaling has

been well characterized (Bilioni et al., 2013; Camp et al., 2010; Yao et al., 2006). However, the

orthologous relationships between *cdon*/*boc* and non-vertebrate orthologs such as *ihog* and *boi*

are less well known. To explore the molecular evolution of this gene family we built a maximum

likelihood tree (phyml) using 44 amino acid sequences from taxa representative for major

bilaterian clades (**Figure 1B**). These data identify *cdon*/*boc* as shared vertebrate specific

paralogs that emerge from a duplication event at either the base of the vertebrate phylogeny or

along the gnathostome lineage following splitting from cyclostomes (hagfish and lamprey).

Importantly, these data are consistent with a hypothesis that *cdon*/*boc* originated during one of

the two vertebrate specific genome duplications (Dehal and Boore, 2005). Analyses suggest

that an independent gene duplication event led to the origins of *boi* and *ihog* paralogs in

drosophilids (**Figure 1B**)(Allen et al., 2011; Izzi et al., 2011; Sanchez-Arrones et al., 2012), a

finding that is concordant with gene trees made publicly available by the TreeFam consortium

(Li et al., 2006; Ruan et al., 2008).

To explore the function of *cdon*/*boc* we used mutations that target the conserved

functional extracellular domains (**Figure 1C**, see methods). Two mutants for each gene were

initially used to confirm an association between phenotypes and gene mutations. As we

observed similar phenotypes for both *cdon* mutations and both *boc* mutations, we focused our

analyses on two mutations in particular, cdon^{co1019} and boc^{co25}, and refer to these mutants

simply as *cdon* and *boc* except where stated.

cdon and boc transcripts are expressed in tNCCs and slow-twitch muscle cells

To determine where endogenous *cdon* and *boc* transcripts are expressed we used

multiplexed hybrid chain reaction *in situ* hybridization (HCR).

We found *cdon* and *boc* transcripts to be abundant in the dorsal central nervous system

(CNS) of 24hpf-48hpf zebrafish embryos (**Figure 2A-D**), consistent with similar data from prior

studies (Bergeron et al., 2011; Cardozo et al., 2014; Kearns et al., 2021). This pattern is

established early in development where *cdon* and *boc* are expressed along the border of the

neural plate in cells that contribute to the dorsal neural tube/NCC (**Figure 2A**). While *boc*

transcripts in the CNS form a broad dorsal-ventral gradient (Kearns et al., 2021), *cdon*

transcripts are restricted to the dorsal-most roofplate cells and the ventral-most floorplate cells

of the CNS by 24hpf. In this way, *cdon* creates a dorsal sub-domain of *boc* expression (**Figure**

2D). Floorplate cells also express hedgehog ligand, and thus unlike *boc*, *cdon* is expressed in

hedgehog ligand producing and non-producing cells. This CNS expression at 24hpf is similar to

expression patterns reported in mouse embryos (Izzi et al., 2011; Kang et al., 2002; Mulieri et

al., 2002; MULIERI et al., 2000; Tenzen et al., 2006).

Prior studies have suggested that *boc* is expressed by slow-twitch muscle mesoderm at

24hpf (Bergeron et al., 2011), however detailed investigation of *cdon/boc* expression in NCCs

and slow-twitch muscle are not available. To further characterize expression, we used the

tg(*sox10*:mRFP) and tg(*prdm1a*:eGFP) transgenic reporter lines to isolate both tissues from the

trunks of 24hpf zebrafish embryos and sequenced the transcriptome of these tissues using bulk

RNA-seq (see methods). These data identify *cdon*/*boc* transcripts in both tissues supporting the

hypothesis that tNCCs and adaxial mesoderm express *cdon* and *boc* at low to moderate levels

(**Figure 2E**).

We then asked if this expression is observable by HCR. Using the tg(*sox10*:mRFP)

transgene to label NCCs, we segmented out *cdon* and *boc* HCR puncta that overlap the *sox10*

transgene confirming *cdon*/*boc* transcripts in tNCCs at 24hpf (**Figure 2F&G**), albeit at levels far

lower than the surrounding tissues.

To confirm expression in slow-twitch muscle/adaxial cell mesoderm, we used the

tg(*prdm1a*:eGFP) (Elworthy et al., 2008). At the 2 somite stage, adaxial cells are labeled with

the tg(*prdm1a*:eGFP) transgene, but we did not observe significant expression of *cdon*/*boc* in

these cells (**Figure 2H**). By the 10 somite stage, *cdon* and *boc* transcripts were observed

throughout the paraxial mesoderm. Using the tg(*prdm1a*:eGFP) transgene to segment out HCR

transcripts expressed by adaxial cells, we show *cdon* and *boc* HCR puncta in slow-twitch

muscle progenitor cells (**Figure 2I-J**). At 24hpf, differentiating slow-twitch muscle cells continue

to express both *cdon* and *boc* (**Figure 2K-L**) consistent with findings from our bulk RNA-seq

data (**Figure 2E**).

176   Taken together, these analyses indicate that *cdon* and *boc* are broadly expressed by

177   mesodermal tissues that include adaxial cell derived slow-twitch muscle cells, as well as tNCCs

178   at low levels, in addition to the developing neural tube.

180   *tNCC migration and slow-twitch muscle development are affected in cdon;boc mutants*

181   To determine whether *cdon* and *boc* mutations affect tNCC migration and slow-twitch

182   muscle development, we used the tg(*sox10*:mRFP) line to label migrating tNCCs and

183   immunostaining of a slow-twitch muscle specific myosin (DSHB F59) to label adaxial mesoderm

184   derivatives in 24hpf embryos.

185   In wildtype embryos, tNCCs collectively migrate in streams down the middle of each

body segment in concert with the lateral migration of slow-twitch muscle cells (**Figure 3A,B**).

We found that tNCC migration was normal in both single *cdon* and *boc* mutant embryos (**Figure**

3D-E). These data from homozygous mutant embryos are in contrast to those reported by

morpholino knockdown of *cdon* (Powell et al., 2015). However, in double *cdon;boc* mutants,

tNCCs failed to form distinguishable streams, and tNCC migration arrested at the ventral neural

tube/body midline (**Figure 3F**), a phenotype similar to that observed in *cdon* morphant embryos

and suggestive of genetic compensation between paralogs (Powell et al., 2015).

193   Similarly, we found that slow-twitch muscle differentiation/migration was normal in *cdon*

and *boc* (Bergeron et al., 2011) single mutants. Only in double mutants did we observe defects

in slow-twitch muscle differentiation (**Figure 3C-F**). Loss of slow-twitch muscle cell

differentiation was not complete, however, and posterior body segments tended to have fewer

slow-twitch muscle fibers than anterior segments. In double mutant embryos, slow-twitch

muscle fibers appeared dysmorphic with loosely packed myosin bundles relative to wildtype

embryos (e.g. compare myosin bundle width and packing in **Figure 3**).

Thus defects in tNCC migration and slow-twitch muscle cell differentiation are observed

in *cdon;boc* double mutants.

cdon and boc affect hedgehog signal transduction in slow-twitch muscle cells

Since *cdon* and *boc* have previously been implicated in hedgehog signaling (Allen et al.,

2011; Zhang et al., 2006), we hypothesized that defects in both tNCC migration and slow-twitch

muscle cell differentiation were due to loss of hedgehog signaling that is only apparent in the

double mutants. To test this, we measured hedgehog signal reception by using the transgenic,

tg(8xGliBS:mCherry-NLS-Odc1) reporter line (hereafter 8xGli:mCherry) that drives nuclear

localized mCherry under the control of 8 tandem Gli binding sites (Mich et al., 2014). The

mCherry fluorophore contains an Odc1 destabilization domain the facilitates proteasomal

degradation to promote rapid fluorophore turnover in expressing cells.

At 24hpf, we observed mCherry positive nuclei in ventral neural tube cells and in slow-

twitch muscle fiber nuclei (**Figure 3G**)(Kearns et al., 2021; Mich et al., 2014). We never

observed mCherry positive nuclei in tNCCs. In *cdon* and *boc* mutants, the number and signal

intensity of nuclei that are 8xGli:mCherry positive was lower as compared to wildtype siblings

(**Figure 3G-J**). This decrease in signal was true for both cells in the neural tube and slow-twitch

muscle. Importantly, 8xGli:mCherry intensity varied across genotypes, with *boc* mutants

displaying lower 8xGli:mCherry signal than *cdon* mutants, and double mutants exhibiting lower

signal than all other genotypes. These data suggest that *cdon* and *boc* mutations have

decreased hedgehog activity in slow-twitch muscle cells, which is consistent with a hypothesis

that perturbations to hedgehog signaling is responsible for the slow-twitch muscle differentiation

phenotype in *cdon* and *boc* mutant embryos (Barresi et al., 2000; Devoto et al., 1996).

Interestingly, the tg(8xGli:mCherry) transgene reported substantively lower hedgehog

signal reception in single *cdon* and *boc* mutant embryos that exhibit normal slow-twitch muscle

differentiation. These data suggest that slow-twitch muscle development is robust to moderate

perturbations to hedgehog signal. If this hypothesis is correct, we reasoned that *cdon* and *boc*

mutants should sensitize embryos to further pharmacological manipulation of smoothened

signaling with cyclopamine. To test this, we in-crossed double heterozygous *cdon;boc* parents

and treated embryos with a subthreshold dose of 20uM cyclopamine from 50% epiboly to 24hpf.

We then quantified the number of slow-twitch muscle fibers (F59) at 24hpf as a readout.

While single mutants treated with ethanol as a control had a similar number of slow-

twitch muscle fibers as their wildtype ethanol treated siblings, single mutants treated with 20μM

cyclopamine had significantly fewer slow-twitch muscle fibers than their similarly cyclopamine

treated wildtype siblings (**Figure 4**). This pattern was even more striking for the double mutant.

While double mutants treated with ethanol have an average of approximately 5-6 slow-twitch

muscle fibers per segment, double mutants lost all slow-twitch muscle fibers when treated with

20μM cyclopamine. We noted qualitative changes to slow-twitch muscle morphology in

cyclopamine treated *cdon* and *boc* embryos. Mutants treated with 20μM cyclopamine often

exhibited gaps or missing slow-twitch muscle fibers in body segments, and muscle fibers were

more likely to have loosely packed myosin.

We interpret these results to indicate that *cdon* and *boc* are modifying hedgehog signal

reception by slow-twitch muscles, and that loss of *cdon* or *boc* sensitize embryos to further

perturbations of hedgehog signal.

Trunk neural crest cells lose directionality but not migratory ability in cdon;boc mutants

The arrested tNCC migratory phenotype in the double mutants is striking and could result from

either a loss of NCC migratory ability or a loss of directionality. To investigate this further we

used the tg(*sox10*:mRFP) transgene to label migrating tNCCs and live imaged NCC migration

through the trunk of the zebrafish embryo beginning at 22hpf (**Figure 5**). Single tNCCs were

then manually tracked. We found that across all genotypes, tNCCs were qualitatively migratory,

and general measures of mobility (e.g. velocity etc.) in mutants were similar to that observed in

wildtype controls (**Figure 5D-G**). In contrast, we observed a significant loss of directional NCC

movement in double *cdon;boc* mutant embryos. In double mutants, tNCCs migrate to the body

midline and then move along the anterior-posterior axis (**Figure 5A-C**), which is in contrast to

the directed ventral movement typical of wildtype embryos. This is seen quantitatively as lower

migratory directionality and an increased proportion of movements in lateral directions (**Figure

5B&H**). Single *cdon* or *boc* mutants did not display migratory phenotypes that were different

from that of wildtype controls. Taken together, we interpret these results to indicate that loss of

cdon and *boc* affects tNCC directionality and path-finding ability, but that *cdon* and *boc* are not

required for overall tNCC mobility.

cdon/boc affect tNCC migration non-cell autonomously

The observed defects in tNCC migration are similar to phenotypes observed in other

hedgehog mutants that have previously been attributed to defects in slow-twitch muscle

differentiation (Banerjee et al., 2011; Banerjee et al., 2013; Honjo and Eisen, 2005). We

therefore hypothesized that tNCC migratory defects in *cdon;boc* mutants are a secondary

consequence of failed slow-twitch muscle differentiation leading to defects in the secretion or

deposition of tNCC guidance cues along the migratory path. In order to determine cell

autonomous vs. non-cell autonomous function we attempted to rescue tNCC migration in double

cdon;boc mutant embryos using two complementary methods.

First, we created genetic mosaics by transplanting double mutant NCCs into wildtype

hosts (**Figure 6**) and examined tNCC migration. The rationale for these experiments is that if

the observed tNCC migration defects are a non-cell autonomous consequence of slow-twitch

muscle differentiation, then we would expect transplanted mutant tNCCs to migrate normally in

wildtype hosts with normal muscle differentiation. This is exactly what we observed.

When wildtype cells from the region forming tNCCs are transplanted into wildtype hosts

we observed transplanted tNCCs to migrate in well-organized streams alongside host tNCCs

(**Figure 6A**). Similarly, in 3 of 3 cases where double mutant tNCCs were transplanted into

wildtype hosts we observed mutant tNCCs migrating alongside host tNCCs in organized

streams (**Figure 6C&D**). Cranial NCCs also migrated normally in all transplant conditions

(**Figure 6B,E,F**). These data suggest that double *cdon/boc* mutant tNCCs migrate normally in

wildtype hosts with normal mesoderm development suggesting a non-autonomous role for *cdon*

and *boc* in tNCC migration.

The second approach used tol2 transgenesis to rescue the *cdon*/*boc* mutations in a

tissue-specific manner. To do this, we used a -2.066kb fragment of the *unc45b* enhancer that

drives expression in adaxial cell mesoderm at early segmentation stages (Berger and Currie,

2013). Importantly, this early adaxial cell expression turns on prior to slow-twitch muscle cell

differentiation (**Figure 6G**). We also took advantage of a constitutively active smoothened

(*smoa1*-eGFP) protein (Ju et al., 2014; Ju et al., 2009), reasoning that if *cdon*/*boc* are affecting

hedgehog signaling then driving *smoa1*-eGFP in the slow-twitch muscle progenitor population

should rescue both slow-twitch muscle morphology and tNCC migration (**Figure 6G**).

Out of 8 double mutant embryos injected with the *smoa1a*-egfp construct, 4 exhibited at

least one tNCC stream that was qualitatively normal. This is in contrast to 4 double mutant

embryos injected with a control eGFP construct where only 1 exhibited a tNCC stream that

could be considered as a wildtype phenotype.

Importantly, these are F0 injected embryos, and thus tol2 transgenesis is mosaic, and

the rescue of phenotype is also expected to be mosaic in the embryo. For instance, in one

embryo injected with the *smoa1*-egfp construct, the left side of the embryos exhibited a typical

mutant phenotype, while the right side of the embryo exhibited tNCC streams that appeared

wildtype-like (**Figure 6J&K**). Further, the slow-twitch muscle morphology on the right side of this

embryo was similarly wildtype-like, while the non-rescued left side appeared dysmorphic with

loose packed myosin bundles.

To account for mosaic rescue, we analyzed 3 body segments over the yolk extension for

each embryo by (1) qualitatively categorizing tNCC streams into rescue, partial rescue, or no-

rescue categories based on tNCC stream formation, and (2) counting the number of slow-twitch

muscle fibers (F59) for that segment. These data show that body segments with rescued tNCC

streams tended to have more slow-twitch muscle fibers than either non-rescued or partial

rescued segments (**Figure 6I**). Further, we noted that slow-twitch muscle fiber morphology

appeared normal in these 'rescued' segments (note tight packed slow-twitch muscle fibers in

Figure 6 panels **K&L** compared to loosely packed fibers in panels **H&J**).

We did notice, however, that some control injected embryos had similar numbers of

slow-twitch muscle fibers to those observed in the *smoa1*-eGFP injected embryos (**Figure 6I**,

note range for control construct). Further, these body segments with greater numbers of slow-

twitch muscle cells in the controls were more likely to exhibit 'partial rescues', or in one case a

wildtype-like tNCC stream. We thus hypothesized that there is likely some degree of incomplete

penetrance in the *cdon*;*boc* mutants whereby occasional segments in some embryos exhibit

wildtype phenotypes. In this scenario, we interpret our mosaic expression of *smoa1*-eGFP to be

pushing *cdon;boc* mutants towards a wildtype phenotype that is interacting with background

variability (genetic or stochastic) in the penetrance of the *cdon*/*boc* alleles. Nonetheless, these

data support the idea of a non-cell autonomous role for *cdon/boc* in regulating directed tNCC

migration.

Deposition of a Col1a1a extracellular matrix is affected in cdon;boc mutants

Prior research suggests that slow-twitch muscle cells express or modify extracellular matrix

proteins including Col15a1b (Guillon et al., 2016), tenascin (Schweitzer et al., 2005), *col18a1*

(Banerjee et al., 2013; Schneider and Granato, 2006), and collagen modifying enzyme *plod3*/*lh3*

(Schneider and Granato, 2006). Unlike cranial NCCs, trunk NCCs are known to migrate on

collagen substrates (Lallier et al., 1992) suggesting that changes to a collagen extracellular

matrix (ECM) may be responsible for mediating the effects of *cdon*/*boc* on tNCC migration

(Banerjee et al., 2013).

To investigate this further we mined our RNAseq dataset of FAC sorted slow-twitch

muscle cells to find all collagen genes expressed by this tissue. These data identify many low to

moderate collagen genes expressed by slow-twitch muscle cells. In particular, *col1a1a* is highly

expressed by slow-twitch muscle, with more than twice the number of estimated transcripts than

the next two most highly expressed collagen genes (**Figure 7A**).

We looked at Col1a1a protein deposition using an antibody raised against the zebrafish

Col1a1a protein (GeneTex). In wildtype embryos at 24hpf, Col1a1a deposition was observed in

the presumptive skin ectoderm, vertical myosepta, and medial border of the mesoderm adjacent

to the neural tube and developing gut (**Figure 7B**). At 16.5 hours, we did not observe Col1a1a

fibers, suggesting that this ECM component is deposited after this time period (Figure 7B).

Intriguingly, this period between 16.5hpf and 24hpf coincides with slow-twitch muscle migration

(Elworthy et al., 2008).

Importantly, we observed Col1a1a fibers deposited as ECM along the medial path that

tNCCs migrate on (**Figure 7C**). In wildtype embryos, Col1a1a is richly deposited along the

vertical myosepta, and fibers extend into the medial region where tNCC streams are observed.

However, in double *cdon*;*boc* mutants, Col1a1a deposition in the tNCC stream pathway was

largely absent. Col1a1a was present in other areas of the embryos such as the vertical

myosepta and skin suggesting that loss of Col1a1a in the trunk is specific to the tNCC stream

path region (**Figure 7C**).

We interpret these data to suggest that tNCC migratory defects may be a consequence

of loss of the Col1a1a ECM along the migratory path (**Figure 7**). Supporting this hypothesis is

the observation that Col1a1a is still expressed dorsally in the double mutants, but stops at the

position where tNCCs often arrest. Thus the arresting migratory phenotype and anterior-

posterior movements of tNCCs along this axis are hypothesized as a consequence of the loss of

Col1a1a expression in the ventral-medial compartments of each body segment.

Discussion

cdon and boc affect tNCC migration non-cell autonomously

Here we show that _cdon_ and _boc_ affect tNCC migration and differentiation of slow-twitch muscle

cells. Despite our prior knockdown studies suggesting that _cdon_ alone functions in tNCC

migration (Powell et al., 2015), we find that only in double _cdon;boc mutants_ do we observe

noticeable tNCC migratory defects that are associated with loss of slow-twitch muscle. Prior

research suggests that slow-twitch muscle differentiation requires hedgehog signaling (Elworthy

et al., 2008), and that tNCC migration requires proper slow-twitch muscle differentiation (Honjo

and Eisen, 2005). Our data are consistent with these result and suggests that the tNCC

migratory defect in _cdon;boc_ mutants is a consequence of defects in mesodermal differentiation

(Banerjee et al., 2011; Banerjee et al., 2013; Honjo and Eisen, 2005). This hypothesis is further

supported by the rescue experiments, that emphasize the importance of non-autonomous

adaxial cell mesoderm differentiation for proper neural crest migration in zebrafish.

While mesodermal tissues are appreciated as critical for neural crest migration (Krull,

2010; Kulesa and Gammill, 2010; Theveneau and Mayor, 2012), the mechanisms underlying

how the mesoderm affects NCC migration vary among both taxa and different NCC

subpopulations. In many tetrapod taxa, mesoderm restricts cranial and trunk NCC migration

along conserved paths through neuropilin, semaphorin, and ephrin ligand-receptor interactions

(Davy and Soriano, 2007; Gammill et al., 2007; Gammill et al., 2006; Krull, 2010; Krull et al.,

1997; Santiago and Erickson, 2002). Thus, the localization of receptors and ligands within

tissues creates permissive/non-permissive zones that restrict NCC migratory paths. In addition,

mesodermal tissues can act as a source for chemoattractant molecules such as chemokine

signaling (Kasemeier-Kulesa et al., 2010; Olesnicky Killian et al., 2009). In zebrafish,

semaphorin/neuropilin interactions mediate cranial NCC migration (Yu and Moens, 2005),

though the role of this pathway in zebrafish tNCC migration is less established. To identify

complementary ligand-receptor pairs in FAC sorted tNCCs and slow-twitch muscle cells, we

searched our RNA-seq data. While some possible interacting molecules could be found, there

were no clear candidate signaling pathways, leading us to hypothesize that there is not a direct

ligand-receptor interaction between tNCCs and slow-twitch muscle mesoderm (not shown,

accessible in NCBI Geo:GSE213728).

Instead, we build on prior research in both tNCC migration (Banerjee et al., 2011;

Banerjee et al., 2013) and motor axon guidance (Guillon et al., 2016; Schneider and Granato,

2006; Zeller and Granato, 1999; Zhang et al., 2004) to suggest that slow-twitch muscle cells

affect tNCC stream formation through ECM deposition. Adaxial derived slow-twitch muscle cells

express Col15a1b (Guillon et al., 2016), *col18a1* (Schneider and Granato, 2006), as well as

chondroitin sulfate proteoglycans associated with collagen-rich ECMs (Zhang et al., 2004).

Morpholino knockdown of *col18a1* leads to tNCC migratory defects that are similar to those we

observe in *cdon;boc mutants* (Banerjee et al., 2013). Our data add to this literature, showing

that slow-twitch muscle express other collagen genes in addition to those previously studied,

including *col1a1a* at exceptionally high levels. Deposition of *col1a1a* along the medial somitic

boundary is lost in *cdon;boc mutants* consistent with a model where slow-twitch muscle cells are

responsible for depositing a collagen rich ECM along the tNCC migratory path.

How the ECM regulates NCC migration is an active area of research (Leonard and

Taneyhill, 2020; Vega-Lopez et al., 2017). One hypothesis is that ECM components create a

permissive zone that bias NCCs to move on specific migratory routes (Banerjee et al., 2013;

Leonard and Taneyhill, 2020). Our hypothesis of a collagen ECM creating a permissive

migratory route is supported by chick neural tube explant studies, which show that trunk, but not

cranial NCCs utilize collagen ECMs as preferred migratory substrates (Lallier et al., 1992). Thus

slow-twitch muscle cells may be depositing a complex collagen-rich ECM critical for tNCC

stream formation. However, other mechanisms may also be at play. For instance, cranial NCCs

utilize durotaxis to migrate along a stiffness gradient of pre-chordal mesoderm in *Xenopus*

(Shellard and Mayor, 2021). Muscle segment organization is disrupted in different slow-twitch

muscle mutants (Henry and Amacher, 2004; Snow and Henry, 2009) and thus mesoderm

stiffness may be affected in the *cdon;boc mutants* that exhibit defects in slow-twitch muscle

differentiation. Whether tNCCs utilize durotaxis has not been studied to our knowledge.

A possible cell autonomous hypothesis for cdon/boc function

Though we argue that *cdon*/*boc* affect tNCC migration non-cell autonomously through slow-

twitch muscle differentiation, *cdon* and *boc* may also play cell autonomous roles in NCC

migration. Indeed, NCC autonomous and non-autonomous function are not mutually exclusive

hypotheses. We point to data showing *cdon/boc* expression in both the neural plate border,

migrating tNCCs and mesoderm to suggest multiple roles for these genes. Recent work in

zebrafish showed that notch signaling among tNCC pre-migratory progenitors establishes a

unique 'leader cell' identity that is required for stream formation (Alhashem et al., 2022;

Richardson et al., 2016) (McLennan et al., 2015; Morrison et al., 2017). Intriguingly, notch

mutant embryos that lack leader cell identity have normal slow-twitch muscle differentiation and

tNCC migratory phenotypes that are remarkably similar to what we observe in *cdon;boc mutants*

(Alhashem et al., 2022). Whether *cdon*/*boc* also play a role in establishing a 'leader cell' identity,

or function in 'leader cell' guidance is unknown. It is worth noting that our transplant rescue

experiments cannot differentiate between a role for mesoderm or 'leader cell' identity, as leader

cells will be present in the wildtype hosts.

cdon and boc as modifiers of hedgehog signaling

Our data add to a growing literature showing that *cdon* and *boc* act synergistically during

zebrafish development to modify hedgehog signal reception and regulate the development of

multiple traits (Echevarria-Andino and Allen, 2020; Song et al., 2015; Zhang et al., 2011).

Interestingly, we found that *cdon* single mutants do not have noticeable phenotypes in

zebrafish, and we can recover adult homozygous mutant zebrafish (not shown). This is contrast

to mouse, where loss of CDON results in mild holoprosencephaly (Allen et al., 2011; Allen et al.,

2007; Echevarria-Andino and Allen, 2020; Zhang et al., 2011). While it is possible that our *cdon*

mutations do not result in full loss of function, we find this possibility unlikely. Both *cdon*

mutations lead to a predicted early stop prior to the transmembrane domain, we cannot fully rule

out the possibility that a late start rescues *cdon* protein resulting in a hypomorphic but functional

receptor or creates a dominant negative. Future research could investigate these possibilities.

An alternative possibility is that *cdon* and *boc* interact with other hedgehog pathway

members resulting in species specific phenotypic responses. Thus in zebrafish other genes may

provide robustness against *cdon* mutations and act as a buffer. The cyclopamine

pharmacological experiments support this hypothesis. Intriguingly, *cdon* expression is often

nested within *boc* expression suggesting that *cdon* and *boc* act together to modify hedgehog

morphogen gradients. Kearns et al. 2021 found that *patched-2* (*ptch2*) expression in the dorsal

neural tube is expanded in *boc*[co25] mutant embryos, suggesting that loss of *boc* alone leads to

more dorsal hedgehog reception (*ptch2* is a readout for hedgehog activity). These data were

interpreted to suggest that *boc* sequesters hedgehog ligand driving hedgehog signal reception

to ventral tissues. Expanding on this hypothesis, we note that *cdon* transcripts are positioned

even more distally in the neural tube from hedgehog ligand source than are *boc* transcripts.

Thus together *boc* and *cdon* gradients may act as strong sequestering agents to limit hedgehog

signal activation and inhibit ligand activity in dorsal neural tube/NCCs.

In summary, we show that *cdon* and *boc* act synergistically in zebrafish to regulate slow-twitch

muscle cell differentiation and affect tNCC migration non-cell autonomously through regulation

of ECM deposition. Our data add to a growing literature in zebrafish showing the critical role of

the slow-twitch muscle for regulating migration of cells. Intriguingly, homologous adaxial/slow-

twitch muscle tissues in other taxa are less understood (Grimaldi et al., 2004) (Hammond et al.,

2009), and to our knowledge there is no known tissue in amniotes that is homologous to adaxial

cells. Despite this, *cdon* and *boc* are known to be critical for myogenesis in amniotes (Jeong et

al., 2017; Lu and Krauss, 2010), and genes expressed by the slow-twitch muscle in zebrafish

also regulate tNCC migration in mouse (Banerjee et al., 2011). Further, we and others note that

cdon and *boc* are expressed in a broad range of tissues including the mesoderm and NCCs

among others. *cdon*, but not *boc*, transcripts are also co-expressed in hedgehog ligand

producing cells, and some studies have suggested that *cdon* plays a role in presenting

hedgehog ligand to responsive cells (Hall et al., 2021). Thus it is likely that *cdon* and *boc* are

playing different roles in different tissues, and that cellular context, expression levels, and

distance from ligand sources, may be critical for determining the biological function of these

receptors.

Materials and Methods

Fish stocks, husbandry, and generation of mutants

Adult fish and embryos were maintained at the University of Colorado Anschutz Medical

school according to standard zebrafish rearing protocols and guidelines for the rearing and

maintenance of vertebrate animals. Two *boc* mutations were investigated. The *boc*[co25] mutant

was previously recovered from an ENU screen (Kearns et al., 2021) and results in an non-

synonymous isoleucine to serine substitution in the third FNIII domain (**Figure 1C**). The

boc[uml/ty54z] mutant was acquired from the Zebrafish International Resource Center and results in

a predicted early stop (Bergeron et al., 2011). Both mutations are previously characterized as

loss of function.

We used CRISPR/Cas9 to generate novel mutations in *cdon*. Two sets of guide RNAs

(sgRNA) were designed to target either the 2nd and 3rd exons (ggcagUUgaggagcagagUa,

UggUaggagccUgUgagagc, UgcUUcagaaUcgcUgagag), or to target the 11th exon

(UcgccaUUgagaUUgUgggg, cUccacgcgaaaUgcagUga) of cdon. DNA oligos were annealed to a

common backbone scaffold, which was used to generate sgRNAs for injections (Williams et al.,

2018). For each sgRNA set, our final CRISPR injection solution consisted of sgRNAs diluted to

20ng/uL, 1ug Cas9 Protein (Invitrogen TrueCut V2), and 5ug Rhodamine Dextran in a total

volume of 10uL. Approximated 2-4nl were injected into the cell of a 1cell stage embryo, and F0

injected embryos were reared to adulthood.

We recovered a single allele for each sgRNA set. These include the cdon^{co1019} allele,

which is a 229bp deletion that spans the second to third exon of *cdon*, removes the intronic

sequence between these exons, and is predicted to result in an early stop (**Figure 1C**). The

cdon^{co1023} allele is a 5bp deletion resulting in a predicted early stop in the 11th exon that codes

for the second FNIII domain. F0 injected embryos were outcrossed for a minimum of two

generations to minimize effects of off target cutting.

Zebrafish *boc* and *cdon* mutants were crossed into existing transgenic reporter lines

including the tg(-7.2kb*sox10*:mRFP)^{vu234} that labels NCCs with cytosolic mRFP (Kucenas et al.,

2008), and the gli reporter tg(8xgli-Xla.Cryaa:NLS-d1mCherry) line that has 8 gli-binding sites

driving a destabilized nuclear localized mCherry (Mich et al., 2014). The tgPAC(*prdm1a*:eGFP)

line was used to label adaxial mesoderm for reference (Elworthy et al., 2008).

cdon/boc molecular conservation and gene tree

Amino acid coding sequences for select bilaterian taxa were downloaded from NCBI GenBank.

Orthologous sequences were searched directly or identified through BLAST searches and

manual curation. The final set of sequences was chosen to include taxonomic representatives

for major bilaterian taxa for which high confidence sequences are available. Effort was made to

identify two paralogs for each taxon. Amino acid sequences were aligned using MUSCLE

(v3.8.31) with default parameters (Edgar, 2004). Amino acid conservation (Figure 1A) was

calculated by pairwise comparisons of *cdon*/*boc* homologues from 11 taxa (human, mouse,

chicken, *Anolis*, frog, bichir, Gar, sheepshead minnow, medaka, catfish, zebrafish) and plotted

in R by smoothing conservation values across 20 amino acid windows. Substitution models

were selected using ProtTest3 by AIC (Darriba et al., 2011). A maximum likelihood gene tree

was built using phyml (v3.3) and a JTT+I+G+F model with 1000 bootstraps (Guindon and

Gascuel, 2003). Nodes with less than 60% bootstrap support were collapsed into polytomies.

We recovered similar results from filtered alignments that removed poorly aligned regions and

regions with gaps.

Immunohistochemistry, in situ hybridization, and stains

Adaxial mesoderm derivatives were stained by fluorescent immuno-histochemistry against a

slow-twitch muscle specific myosin (anti-MYH1A, F59, DSHB) at a concentration of 1:20 as

previously described (Elworthy et al., 2008; Nguyen-Chi et al., 2012). Hybrid Chain Reaction

(HCR) *in situ* hybridizations were performed as previously described (Choi et al., 2018; Lencer

et al., 2021). Unlike traditional enzymatic methods for *in situ* hybridization, HCR allows single

molecule measurement of gene expression at cellular resolution, which enables identifying cells

and cell types that express transcripts at relatively low levels (Kearns et al., 2021). Custom

HCR probes for zebrafish *cdon* and *boc* were acquired from Molecular Instruments and used at

a concentration of 2picoMol/500uL Hybridization Buffer.

RNA-seq of tNCCs and adaxial cell mesoderm

We used bulk RNA-seq to characterize the transcriptome of tNCCs and adaxial cell mesoderm

at 24 hpf. To do this we took advantage of embryos that were doubly transgenic for the

tg(*sox10*:mRFP) transgene that labels NCCs and tg(*prdm1a*:eGFP) transgene that labels

adaxial mesoderm. Trunks from 80 embryos were dissected and cells were dissociated by

incubating in Accumax (Stem cell technologies) at 37°C for 30 minutes with intermittent

agitation. Cells were cleaned of debris by passing through a 40µM mesh filter, and washed in

Dulbecco's phosphate buffered saline plus DNaseI (final 7500U/mL). NCCs and adaxial

mesoderm cells were isolated by FACS using a MoFlo XDP100 at the CU Anschutz Flow

Cytometry Core.

Total RNA was extracted from FAC sorted samples of tNCCs and adaxial mesoderm

using the RNAqueous-Micro Isolation Kit (ThermoFisher). Library preparation and Illumina

sequencing (PE 150bp) was performed by the CU Anschutz Genomics core on a

NovaSeq6000. Reads were aligned to the zebrafish genome (GRCz11, ensemble annotations)

using STAR aligner (Dobin et al., 2013) and gene counts were calculated using featurecounts

(Liao et al., 2014) with default parameters. DEseq2 (Love et al., 2014) was used to estimate

normalized expression levels using the normTransform function.

Imaging and tracking of migrating cells

For live-imaging and tracking, neural crest cells were labelled using the tg(*sox10*:mRFP)

line. Embryos were mounted in 0.4% low melt agarose and neural crest cells were imaged over

the yolk extension every 5 minutes from 22hpf to 30hpf on an Andor spinning disk confocal

microscope at 28°C. Neural crest cells were manually tracked along the xy axis from maximum

projection images using the ImageJ mTrackJ plugin (Meijering et al., 2012). Drift was accounted

for using the Correct_3D_Drift.py plugin as implemented in ImageJ (Parslow et al., 2014). As we

only observed a tNCC migratory defect in the medial migrating streams, we limited our tracking

of cells to tNCCs migrating along this route. Quantitative analysis of track mobility was

performed in R. A small number of short tracks with fewer than 30 points (2.5hrs) were

excluded. As we also noted that NCCs often stopped migrating upon reaching a ventral position

we further limited tracks to a maximum of 6 hours in order to capture cell migration. Limiting all

tracks to the same time (e.g. number of points) produces similar results. Rose plots depict cell

movements with a velocity greater than $0.3\mu M/min$, which was empirically determined as a

cutoff for capturing cell movements while excluding noise from drift and changes in cell shape.

Using different cutoffs produced similar results. Linear models were built in R by including track

length as a fixed effect, and embryo identity and date of imaging as random effects, with the

packages lme4 (Bates et al., 2015), lmerTest (Kuznetsova et al., 2017), and emmeans. Images

of tracks were produced in napari (Sofroniew et al., 2019).

Transplant Rescue Experiments

Transplant rescues were performed by transplanting cells from gastrula stage embryos

produced by in crossing double heterozygous *cdon*[co1019]*;boc*[co25] parents into gastrula stage

wildtype hosts. All donor embryos were transgenic for the tg(*sox10*:mRFP) transgene, while

hosts were transgenic for the tg(*sox10*:eGFP) transgene (Carney et al., 2006), allowing us to

differentiate transplanted NCCs (red) from host NCCs (green). Embryos were fixed at 24hpf,

genotyped, and imaged for tNCC migration.

Tol2 Rescue Experiments

Tol2 rescue experiments were performed following standard gateway cloning and tol2

transgenic methods (Kwan et al., 2007). Briefly,the primers 5'-

GCGTAAAACTGTGGCGCGTAAAAC-3' and 5'-CATTGAAGTCAATTCAGCTTCGTC-3' were

used to clone the Unc45b promoter from wildtype genomic DNA following (Berger and Currie,

2013). This produced a 2.066kb fragment of the 5' upstream region to the *unc45b* gene. This

fragment was placed into a tol2kit p5Entry vector using TA cloning (Invitrogen). A *smoa1*-eGFP

construct was obtained as generous gift from Michael Taylor and subcloned into a tol2 middle

entry vector. Expression constructs using the -2.066Unc45b enhancer to drive either *smoa1*-

eGFP, or eGFP as a control, in adaxial cell progenitors.

For rescues, double heterozygous *cdon*[co1019]*;boc*[co25] parents were in crossed and

embryos were injected at the 1 cell stage with tol2 constructs and tol2 mRNA at a total injection

volume of ~2nL. Embryos were reared to 24hpf and then stained for F59 by

immunofluorescence (see above) to assess tNCC stream formation and slow-twitch muscle cell

morphology.. Wildtype and double mutant embryos were imaged over the yolk extension. Sick

embryos were not used for analyses.

Pharmacological manipulation

For pharmacological manipulation, embryos were manually dechorionated and treated with

20uM cyclopamine in ethanol (5uL/mL) from 50% epiboly to 24 hpf. Embryos that appeared

developmentally delayed were excluded.

Imaging, Statistical Analysis, and biological replication

All imaging was performed on either a Leica Sp8 LSM or Andor spinning disk confocal

microscope. Image analysis and presentation was performed in ImageJ or napari. All statistical

analyses were performed in R. Experiments were repeated at least 3 times and biological

variation was accounted for in models and through inclusion of biological replicates.

Acknowledgements

We thank members of the Artinger and Prekeris laboratories for comments on this manuscript

and project. Special thanks to Caleb Doll, Bruce Appel, Christina Kearns for providing tools and

advice. We thank Michael Taylor for his generous gift of the smoa1-eGFP construct. This work

was in part supported by GM122768 (to RP) and a University of Colorado RNA Biosciences

Initiative award to EL and KA. EL was supported by National Institutes of Health Ruth L.

Kirschstein National Research Service Awards (T32CA17468 and F32HD103406). EL, RP, and

KA conceived of the study and wrote the manuscript. EL performed research and analyses.

Figure 1. *cdon* and *boc* are transmembrane receptors and share conserved domain

structure. (A) *cdon*/*boc* share a domain structure of 4-5 immunoglobin (IG) domains, 3

fibronectin (FNIII) domains, a transmembrane (TM) region, and a cytosolic C-terminal tail. A fifth

cdon IG domain (*) is found in some tetrapod taxa. Percent conservation (20 AA sliding window)

is shown for an alignment of *cdon* and *boc* amino acid sequences from 11 tetrapod and ray-

finned fish taxa. Note that the extracellular domains exhibit high sequence conservation, while

the cytosolic region is less conserved. **(B)** Maximum likelihood tree (phyml) for curated *cdon*/*boc*

ortholog amino acid sequences from multiple bilaterian taxa. Note independent duplications for

Ihog/Boi in drosophilids and Cdon/Boc in vertebrates. Node support was calculated based on

1000 bootstraps, and nodes with less than 60% bootstrap support were collapsed. Asterisks

mark taxa used to generate conservation in panel A. **(C)** Schematic of *cdon* and *boc* mutant

alleles characterized in the current study.

Figure 2. *cdon* **and** *boc* **are expressed by NCCs and adaxial mesoderm cells**. **(A-C)** HCR

shows *cdon* (magenta) and boc (green) expression in the developing neural tube of 2 somite

(A), 24 hpf **(B)**, and 48 hpf **(C)** embryos. Panels **B&C** are 20μM sections at the level of the yolk

extension (YE) in the trunk. Note expression at the border of the neural plate **(A)**, and in the

rdorsal neural tube at 24 and 48 hpf. Weak expression is observed throughout the mesoderm.

(D) Schematic summarizing expression of *cdon*/*boc* in the neural tube. Gradient of hedgehog

ligand is indicated at left in grey, while expression of *cdon*/*boc* are indicated by magenta and

green respectively. **(E)** Bulk RNA-seq from FAC-sorted tNCCs (blue) and slow-twitch muscle

(red) suggest both tissues express *cdon*/*boc*. Boxplots show *cdon*/*boc* expression levels in 2

replicate samples. Right panel shows histogram of gene expression levels for all samples for all

genes. Location of genes that mark NCCs or slow-twitch muscle are indicated. **(F&G)** HCR

expression of *cdon*/*boc* in tNCCs. **(F)** whole mount lateral view of 24 hpf embryo showing

expression of *cdon*/*boc* with NCCs (grey) labeled by the tg(*sox10*:mRFP) transgene. HCR

puncta overlapping the *sox10* transgene (*) are segmented out and shown in yellow **(F'&F'')** **(G)**

Cross section shows tNCCs expressing *cdon*/*boc* transcripts, with HCR puncta overlapping the

sox10 transgene **(G' and G'')**. **(H-L)** Slow-twitch muscle cells express *cdon* and *boc*. The

tg(*prdm1a*:eGFP) transgene labels adaxial cells at 2 somites **(H)**, and slow-twitch muscle cells

at later stages **(I-L)**. (**I**) 3D rendered cross section of whole mount imaged embryo shows HCR

puncta overlapping the tg(*prdm1a*:eGFP) transgene **(I' and I'')**. **(J)** 3D rendered lateral views

show HCR puncta overlapping the *prdm1a* transgene. **(K)** 24hpf slow-twitch muscle cells (grey)

continue to express *cdon*/*boc* transcripts. Panels **K'** and **K''** show overlapping puncta. **(L)** Cross

section of the trunk of a 24hpf embryo shows *cdon*/*boc* puncta in slow-twitch muscle.

Figure 3. *cdon* and *boc* double mutants have NCC migration and slow-twitch muscle

differentiation phenotypes and a reduction in hedgehog signal reception. (A) Schematic

cross sections show NCC (red) and adaxial mesoderm (green) migration from approximately 10

somites to 20 somites. Adaxial mesoderm cells are specified adjacent to the notochord (nc) and

migrate laterally to differentiate into slow-twitch muscle fibers. Concurrent with adaxial cell

migration, NCCs migrate ventrally from dorsal neural tube (NT). **(B)** 3D renderings of a 10uM

cross section from the trunk of a 15 somite (left) and a whole mount 24hpf embryo (right)

showing association between migrating adaxial mesoderm (green) and NCCs (red). **(C-F)**

Maximum projection images show NCC streams (tg(*sox10*:mRFP)) and immunostaining for

slow-twitch muscle (F59) for 24hpf wildtype embryos. Note loss of both streams and slow-twitch

muscle in the double mutant. All images taken over the yolk extension in the trunk. **(G-J)**

Maximum projection images of the tg(8xGli:mCherry) transgene in 24hpf embryos. Note

mCherry positive nuclei in the slow-twitch muscle mesoderm (yellow arrows) and ventral neural

tube (yellow arrowheads). mCherry signal progressively decreases in *cdon;boc mutants*

reflecting lower hedgehog signal reception.

Figure 4. *cdon* and *boc* sensitize zebrafish embryos to a reduction in hedgehog signal by

cyclopamine treatment. **(A)** Representative images show slow-twitch muscle fiber morphology

in 24hpf embryos treated with either 20uM cyclopamine or an equivalent volume of ethanol as a

control. Note *cdon* and *boc* mutants exhibit more severe phenotypes than their wildtype siblings

when treated with cyclopamine as evidenced by gaps in segments (arrows), and complete loss

of slow-twitch muscles in the double. **(B)** Quantification of average number of slow-twitch

muscle fibers per segment shows that *cdon* and *boc* mutations are sensitizing embryos to

cyclopamine treatment. Note that the number of slow-twitch muscle fibers in single *cdon* and

boc mutants are similar to wildtype siblings when treated with ethanol, but significantly lower

when treated with 20uM cyclopamine. Significance values are from Tukey pairwise post-hoc

tests: $<0.05 = *$, $<0.001 = ***$.

Figure 5. NCC directionality, but not migratory ability, is disrupted in the *cdon;boc*

mutants. (A) Maximum projection images show migrating tNCCs in representative wildtype,

cdon, *boc*, and double mutant embryos. Migratory tracks from individual cells are shown by

colored lines. Note that NCCs form horizontal tracks at the body midline in the double mutant.

(B) Matched roseplots (left panels) showing directional movement and line plots (right panel)

showing tracks of NCCs in each genotype. For both panels, dorsal is up and anterior is to the

left. Samples sizes reflect #tracks/#Embryos. Note that ventral directionality of NCC movement

is lower in the double mutant as best seen by the rose plot. **(C)** NCC migration tracks (colored

lines) are overlayed onto outlines of the body (grey dashed lines) for representative embryos.

Note the vertical aligned tracks in the wildtype, *cdon*, and *boc* mutants that reflects ventral

migration. Tracks form horizontal lines in the double mutant reflecting NCC migratory

movements along the anterior-posterior axis. Body outlines do not reflect growth. Somite

segments in the double mutant are wider and dysmorphic. **(D-H)** Boxplots show quantification of

NCC migration for maximum velocity **(D)**, mean velocity **(E)**, track length **(F)**, track displacement

(G), and track straightness **(H)**. Note that the double *cdon;boc mutants* are the only condition

that is significantly different from wildtype embryos in track straightness. Small grey points are

individual track values, large colored points are means for individuals, boxplots reflect variability

across tracks. **(I)** Schematic of track straightness which is calculated as track displacement

dived by track length.

Figure 6. *cdon/boc* affect tNCC migration non-cell autonomously. **(A-F)** Mutant tNCCs

transplanted into wildtype hosts migrate normally. For all transplants, donor embryos were

derived by in crossing double heterozygous *cdon;boc* parents where 1/16 will be double

mutants. All donor embryos were transgenic for the tg(*sox10*:mRFP) transgene, and wildtype

embryos carry the tg(*sox10*:eGFP) transgene. Wildtype NCCs transplanted into a wildtype

background migrate normally in both the trunk **(A)** and cranial **(B)** regions. Similarly, double

mutant *cdon;boc* tNCCs migrate normally in wildtype hosts when transplanted into the trunk

(C&D, 3/3 embryos) and cranial regions. For panels **(C,D)**, trunk streams are shown as both 3D

projections and 3D volume renderings **(C' & D')**. **(G-L)** Mosaic expression of *smoa1-eGFP* in

adaxial cell progenitors leads to a partial rescue of tNCC streams. **(G)** tol2 constructs used for

rescues. Dorsal view of a 10 somite embryo shows mosaic expression in adaxial cell

progenitors. **(H)** Representative image of slow-twitch muscle morphology (F59) and tNCC

streams in a double cdon/boc embryo injected with a control eGFP construct. **(I)** Number of

slow-twitch muscle fibers per segment in eGFP/control and *smoa1a-eGFP* injected embryos.

For *smoa1a-eGFP* injected embryos, points are separated by whether the tNCC stream in that

segment was qualitatively rescued. IControl injected, N=4 embryos. *smoa1a-eGFP* injected,

N=8. **(J-L)** Representative images of *smoa1a-eGFP* injected embryos. Panel **(J)** shows left side

of an embryo where tNCC migration was not rescued, while panel **(K)** shows the right side of

this same embryo where tNCC migration was rescued. Note the slow-twitch muscle fiber

morphology is only rescued on the right side of the embryo. Panel **(L)** shows *smoa1a-eGFP*

expression as mosaic. Arrows points to segments with qualitatively rescued tNCC streams.

Asterisks label segments with non-rescued streams, "p" labels a representative "partial rescue".

e = eye, ba1 = branchial arch 1, ba2 = branchial arch 2, pa3-5 = posterior arches.

Figure 7. Col1a1a extracellular matrix (ECM) deposition is disrupted in cdon;boc

mutants. (A) RNA-seq data from FAC sorted tg(*prdm1a*:eGFP) slow-twitch muscle cells at

24hpf identify numerous expressed collagen genes including *col1a1a*. **(B)** 3D projections of a

15somite and 24hpf embryo show Col1a1a protein deposition. **(C)** 3D projections (top rows) and

volume renderings (bottom row) show medial Col1a1a deposition in wildtype and double mutant

embryos. Note loss of Col1a1a ECM along the medial zone of each segment in the double

mutant associated with loss of slow-twitch muscle differentiation. Retention of Col1a1a around

dorsal neural tube (yellow arrows) coincides with location tNCCs often arrest migration. **(D)**

Schematic model whereby slow-twitch muscle cells deposit an ECM containing Colo1a1a

necessary for tNCC stream formation and migration.

References

Alhashem, Z., Feldner-Busztin, D., Revell, C., Alvarez-Garcillan Portillo, M., Camargo-Sosa, K., Richardson, J., Rocha, M., Gauert, A., Corbeaux, T., Milanetto, M., Argenton, F., Tiso, N., Kelsh, R.N., Prince, V.E., Bentley, K., Linker, C., 2022. Notch controls the cell cycle to define leader versus follower identities during collective cell migration. Elife 11.

Allen, B.L., Song, J.Y., Izzi, L., Althaus, I.W., Kang, J.S., Charron, F., Krauss, R.S., McMahon, A.P., 2011. Overlapping roles and collective requirement for the coreceptors GAS1, CDO, and BOC in SHH pathway function. Dev Cell 20, 775-787.

Allen, B.L., Tenzen, T., McMahon, A.P., 2007. The Hedgehog-binding proteins Gas1 and Cdo cooperate to positively regulate Shh signaling during mouse development. Genes Dev 21, 1244-1257.

Banerjee, S., Gordon, L., Donn, T.M., Berti, C., Moens, C.B., Burden, S.J., Granato, M., 2011. A novel role for MuSK and non-canonical Wnt signaling during segmental neural crest cell migration. Development 138, 3287-3296.

Banerjee, S., Isaacman-Beck, J., Schneider, V.A., Granato, M., 2013. A novel role for Lh3 dependent ECM modifications during neural crest cell migration in zebrafish. PLoS One 8, e54609.

Barresi, M.J.F., Stickney, H.L., Devoto, S.H., 2000. The zebrafish slow-muscle-omitted gene product is required fro Hedgehog signal transduction and the development of slow muscle identity. Development 127, 2189-2199.

Barriga, E.H., Maxwell, P.H., Reyes, A.E., Mayor, R., 2013. The hypoxia factor Hif-1alpha controls neural crest chemotaxis and epithelial to mesenchymal transition. J Cell Biol 201, 759-776.

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Usinglme4. Journal of Statistical Software 67.

Berger, J., Currie, P.D., 2013. 503unc, a small and muscle-specific zebrafish promoter. Genesis 51, 443-447.

Bergeron, S.A., Tyurina, O.V., Miller, E., Bagas, A., Karlstrom, R.O., 2011. Brother of cdo (umleitung) is cell-autonomously required for Hedgehog-mediated ventral CNS patterning in the zebrafish. Development 138, 75-85.

Bilioni, A., Sanchez-Hernandez, D., Callejo, A., Gradilla, A.C., Ibanez, C., Mollica, E., Carmen Rodriguez-Navas, M., Simon, E., Guerrero, I., 2013. Balancing Hedgehog, a retention and release equilibrium given by Dally, Ihog, Boi and shifted/DmWif. Dev Biol 376, 198-212.

Camp, D., Currie, K., Labbé, A., Meyel, D.J.v., Charron, F.d.r., 2010. Ihog and Boi are essential for Hedgehog signaling in Drosophila. Neural Development 5.

Cardozo, M.J., Sanchez-Arrones, L., Sandonis, A., Sanchez-Camacho, C., Gestri, G., Wilson, S.W., Guerrero, I., Bovolenta, P., 2014. Cdon acts as a Hedgehog decoy receptor during proximal-distal patterning of the optic vesicle. Nat Commun 5, 4272.

Carmona-Fontaine, C., Matthews, H.K., Kuriyama, S., Moreno, M., Dunn, G.A., Parsons, M., Stern, C.D., Mayor, R., 2008. Contact inhibition of locomotion in vivo controls neural crest directional migration. Nature 456, 957-961.

Carney, T.J., Dutton, K.A., Greenhill, E., Delfino-Machin, M., Dufourcq, P., Blader, P., Kelsh, R.N., 2006. A direct role for Sox10 in specification of neural crest-derived sensory neurons. Development 133, 4619-4630.

Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha,
A., Pierce, N.A., 2018. Third-generation in situ hybridization chain reaction: multiplexed,
quantitative, sensitive, versatile, robust. Development 145.
Clay, M.R., Halloran, M.C., 2011. Regulation of cell adhesions and motility during initiation of
neural crest migration. Curr Opin Neurobiol 21, 17-22.
Clay, M.R., Halloran, M.C., 2013. Rho activation is apically restricted by Arhgap1 in neural crest
cells and drives epithelial-to-mesenchymal transition. Development 140, 3198-3209.
Clay, M.R., Halloran, M.C., 2014. Cadherin 6 promotes neural crest cell detachment via F-actin
regulation and influences active Rho distribution during epithelial-to-mesenchymal transition.
Development 141, 2506-2515.
Connor, R.M., Allen, C.L., Devine, C.A., Claxton, C., Key, B., 2005. BOC, brother of CDO, is a
dorsoventral axon-guidance molecule in the embryonic vertebrate brain. J Comp Neurol 485,
32-42.
Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2011. ProtTest 3: fast selection of best-fit
models of protein evolution. Bioinformatics 27, 1164-1165.
Davy, A., Soriano, P., 2007. Ephrin-B2 forward signaling regulates somite patterning and neural
crest cell development. Dev Biol 304, 182-193.
Dehal, P., Boore, J.L., 2005. Two rounds of whole genome duplication in the ancestral
vertebrate. PLoS Biol 3, e314.
Devoto, S.H., Melancon, E., Eisen, J.S., Westerfield, M., 1996. Identification of separate slow
and fast muscle precursor cells in vivo, prior to somite formation. Development 122, 3371-
3380.
Diener, J., Sommer, L., 2021. Reemergence of neural crest stem cell-like states in melanoma
during disease progression and treatment. Stem Cells Transl Med 10, 522-533.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15-21.
Echevarria-Andino, M.L., Allen, B.L., 2020. The hedgehog co-receptor BOC differentially
regulates SHH signaling during craniofacial development. Development 147.
Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32, 1792-1797.
Elworthy, S., Hargrave, M., Knight, R., Mebus, K., Ingham, P.W., 2008. Expression of multiple
slow myosin heavy chain genes reveals a diversity of zebrafish slow twitch muscle fibres with
differing requirements for Hedgehog and Prdm1 activity. Development 135, 2115-2126.
Ferent, J., Giguere, F., Jolicoeur, C., Morin, S., Michaud, J.F., Makihara, S., Yam, P.T., Cayouette,
M., Charron, F., 2019. Boc Acts via Numb as a Shh-Dependent Endocytic Platform for Ptch1
Internalization and Shh-Mediated Axon Guidance. Neuron 102, 1157-1171 e1155.
Gammill, L.S., Gonzalez, C., Bronner-Fraser, M., 2007. Neuropilin 2/semaphorin 3F signaling is
essential for cranial neural crest migration and trigeminal ganglion condensation. Journal of
Neurobiology 67, 47-56.
Gammill, L.S., Gonzalez, C., Gu, C., Bronner-Fraser, M., 2006. Guidance of trunk neural crest
migration requires neuropilin 2/semaphorin 3F signaling. Development 133, 99-106.
Grimaldi, A., Tettamanti, G., Martin, B.L., Gaffield, W., Pownall, M.E., Hughes, S.M., 2004.
Hedgehog regulation of superficial slow muscle fibres in Xenopus and the evolution of tetrapod
trunk myogenesis. Development 131, 3249-3262.

Guillon, E., Bretaud, S., Ruggiero, F., 2016. Slow Muscle Precursors Lay Down a Collagen XV
Matrix Fingerprint to Guide Motor Axon Navigation. J Neurosci 36, 2663-2676.
Guindon, S., Gascuel, O., 2003. A simple, fast, and accurate algorithm to estimate large
phylogenies by maximum likelihood. Syst Biol 52, 696-704.
Hall, E.T., Dillard, M.E., Stewart, D.P., Zhang, Y., Wagner, B., Levine, R.M., Pruett-Miller, S.M.,
Sykes, A., Temirov, J., Cheney, R.E., Mori, M., Robinson, C.G., Ogden, S.K., 2021. Cytoneme
delivery of Sonic Hedgehog from ligand-producing cells requires Myosin 10 and a Dispatched-
BOC/CDON co-receptor complex. Elife 10.
Hammond, K.L., Baxendale, S., McCauley, D.W., Ingham, P.W., Whitfield, T.T., 2009. Expression
of patched, prdm1 and engrailed in the lamprey somite reveals conserved responses to
Hedgehog signaling. Evol Dev 11, 27-40.
Henry, C.A., Amacher, S.L., 2004. Zebrafish slow muscle cell migration induces a wave of fast
muscle morphogenesis. Dev Cell 7, 917-923.
Hong, M., Christ, A., Christa, A., Willnow, T., Krauss, R.S., 2020. Cdon Mutation and Fetal
Alcohol Converge on Nodal Signaling in a Gene-Environment Interaction Model of
Holoprosencephaly. BioRxiv.
Honjo, Y., Eisen, J.S., 2005. Slow muscle regulates the pattern of trunk neural crest migration in
zebrafish. Development 132, 4461-4470.
Izzi, L., Levesque, M., Morin, S., Laniel, D., Wilkes, B.C., Mille, F., Krauss, R.S., McMahon, A.P.,
Allen, B.L., Charron, F., 2011. Boc and Gas1 each form distinct Shh receptor complexes with
Ptch1 and are required for Shh-mediated cell proliferation. Dev Cell 20, 788-801.
Jeong, M.H., Kim, H.J., Pyun, J.H., Choi, K.S., Lee, D.I., Solhjoo, S., O'Rourke, B., Tomaselli, G.F.,
Jeong, D.S., Cho, H., Kang, J.S., 2017. Cdon deficiency causes cardiac remodeling through
hyperactivation of WNT/beta-catenin signaling. Proc Natl Acad Sci U S A 114, E1345-E1354.
Ju, B., Chen, W., Spitsbergen, J.M., Lu, J., Vogel, P., Peters, J.L., Wang, Y.D., Orr, B.A., Wu, J.,
Henson, H.E., Jia, S., Parupalli, C., Taylor, M.R., 2014. Activation of Sonic hedgehog signaling in
neural progenitor cells promotes glioma development in the zebrafish optic pathway.
Oncogenesis 3, e96.
Ju, B., Spitsbergen, J., Eden, C.J., Taylor, M.R., Chen, W., 2009. Co-activation of hedgehog and
AKT pathways promote tumorigenesis in zebrafish. Mol Cancer 8, 40.
Kang, J.-S., Mulieri, P.J., Hu, Y., Taliana, L., Krauss, R.S., 2002. BOC, an Ig superfamily member,
associates with CDO to positively regulate myogenic differentiation. EMBO J 21, 114-124.
Kang, J.S., Bae, G.U., Yi, M.J., Yang, Y.J., Oh, J.E., Takaesu, G., Zhou, Y.T., Low, B.C., Krauss, R.S.,
2008. A Cdo-Bnip-2-Cdc42 signaling pathway regulates p38alpha/beta MAPK activity and
myogenic differentiation. J Cell Biol 182, 497-507.
Kasemeier-Kulesa, J.C., McLennan, R., Romine, M.H., Kulesa, P.M., Lefcort, F., 2010. CXCR4
controls ventral migration of sympathetic precursor cells. J Neurosci 30, 13078-13088.
Kaufman, C.K., Mosimann, C., Fan, Z.P., Yang, S., Thomas, A.J., Ablain, J., Tan, J.L., Fogley, R.D.,
van Rooijen, E., Hagedorn, E.J., Ciarlo, C., White, R.M., Matos, D.A., Puller, A.C., Santoriello, C.,
Liao, E.C., Young, R.A., Zon, L.I., 2016. A zebrafish melanoma model reveals emergence of
neural crest identity during melanoma initiation. Science 351, aad2197.
Kearns, C.A., Walker, M., Ravanelli, A.M., Scott, K., Arzbecker, M.R., Appel, B., 2021. Zebrafish
spinal cord oligodendrocyte formation requires boc function. BioRxiv.

Krull, C.E., 2010. Neural crest cells and motor axons in avians: Common and distinct migratory
molecules. Cell Adh Migr 4, 631-634.
Krull, C.E., Lansford, R., Gale, N.W., Collazo, A., Marcelle, C., Yancopoulos, G.D., Fraser, S.E.,
Bronner-Fraser, M., 1997. Interactions of Eph-related receptors and ligands confer rostrocaudal
pattern to trunk neural crest migration. Current Biology 7, 571-580.
Kucenas, S., Snell, H., Appel, B., 2008. nkx2.2a promotes specification and differentiation of a
myelinating subset of oligodendrocyte lineage cells in zebrafish. Neuron glia biology 4, 71-81.
Kulesa, P.M., Gammill, L.S., 2010. Neural crest migration: patterns, phases and signals. Dev Biol
344, 566-568.
Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in Linear
Mixed Effects Models. Journal of Statistical Software 82, 1-26.
Kwan, K.M., Fujimoto, E., Grabher, C., Mangum, B.D., Hardy, M.E., Campbell, D.S., Parant, J.M.,
Yost, H.J., Kanki, J.P., Chien, C.B., 2007. The Tol2kit: a multisite gateway-based construction kit
for Tol2 transposon transgenesis constructs. Dev Dyn 236, 3088-3099.
Lallier, T., LeBlanc, G., Artinger, K.B., Bronner-Fraser, M., 1992. Cranial and trunk neural crest
cells use different mechanisms for attachment to extracellular matrices. Development 116, 531-
541.
Lencer, E., Prekeris, R., Artinger, K.B., 2021. Single-cell RNA analysis identifies pre-migratory
neural crest cells expressing markers of differentiated derivatives. Elife 10.
Leonard, C.E., Taneyhill, L.A., 2020. The road best traveled: Neural crest migration upon the
extracellular matrix. Semin Cell Dev Biol 100, 177-185.
Li, H., Coghlan, A., Ruan, J., Coin, L.J., Heriche, J.K., Osmotherly, L., Li, R., Liu, T., Zhang, Z.,
Bolund, L., Wong, G.K., Zheng, W., Dehal, P., Wang, J., Durbin, R., 2006. TreeFam: a curated
database of phylogenetic trees of animal gene families. Nucleic Acids Res 34, D572-580.
Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30, 923-930.
Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 15, 550.
Lu, M., Krauss, R.S., 2010. N-cadherin ligation, but not Sonic hedgehog binding, initiates Cdo-
dependent p38alpha/beta MAPK signaling in skeletal myoblasts. Proc Natl Acad Sci U S A 107,
4212-4217.
Mayor, R., Etienne-Manneville, S., 2016. The front and rear of collective cell migration. Nat Rev
Mol Cell Biol 17, 97-109.
McLennan, R., Schumacher, L.J., Morrison, J.A., Teddy, J.M., Ridenour, D.A., Box, A.C., Semerad,
C.L., Li, H., McDowell, W., Kay, D., Maini, P.K., Baker, R.E., Kulesa, P.M., 2015. Neural crest
migration is driven by a few trailblazer cells with a unique molecular signature narrowly
confined to the invasive front. Development 142, 2014-2025.
Meijering, E., D'zyubachyk, O., Smal, I., 2012. Methods for Cell and Particle Tracking. Methods
in Enzymology 504, 183-200.
Mich, J.K., Payumo, A.Y., Rack, P.G., Chen, J.K., 2014. In vivo imaging of Hedgehog pathway
activation with a nuclear fluorescent reporter. PLoS One 9, e103661.
Morrison, J.A., McLennan, R., Wolfe, L.A., Gogol, M.M., Meier, S., McKinney, M.C., Teddy, J.M.,
Holmes, L., Semerad, C.L., Box, A.C., Li, H., Hall, K.E., Perera, A.G., Kulesa, P.M., 2017. Single-cell

transcriptome analysis of avian neural crest migration reveals signatures of invasion and
molecular transitions. Elife 6.
Mulieri, P.J., Kang, J.S., Sassoon, D.A., Krauss, R.S., 2002. Expression of the boc gene during
murine embryogenesis. Dev Dyn 223, 379-388.
MULIERI, P.J., OKADA, A., SASSOON, D.A., MCCONNELL, S.K., KRAUSS, A.R.S., 2000.
Developmental Expression Pattern of the cdo Gene. Developmental Dynamics 219, 40-49.
Nguyen-Chi, M.E., Bryson-Richardson, R., Sonntag, C., Hall, T.E., Gibson, A., Sztal, T., Chua, W.,
Schilling, T.F., Currie, P.D., 2012. Morphogenesis and cell fate determination within the adaxial
cell equivalence group of the zebrafish myotome. PLoS Genet 8, e1003014.
Okada, A., Charron, F., Morin, S., Shin, D.S., Wong, K., Fabre, P.J., Tessier-Lavigne, M.,
McConnell, S.K., 2006. Boc is a receptor for sonic hedgehog in the guidance of commissural
axons. Nature 444, 369-373.
Okumura, F., Oki, N., Fujiki, Y., Ikuta, R., Osaki, K., Hamada, S., Nakatsukasa, K., Hisamoto, N.,
Hara, T., Kamura, T., 2021. ZSWIM8 is a myogenic protein that partly prevents C2C12
differentiation. Sci Rep 11, 20880.
Olesnicky Killian, E.C., Birkholz, D.A., Artinger, K.B., 2009. A role for chemokine signaling in
neural crest cell migration and craniofacial development. Dev Biol 333, 161-172.
Parslow, A., Cardona, A., Bryson-Richardson, R., 2014. Sample drift correctino following 3D
confocal time-lapse imaging. J. Vis. Exp. 12.
Perris, R., Perissinott, D., 2000. Role of the extracellular matrix during neural crest cell
migration. Mechanisms of Development 95, 3-21.
Powder, K.E., Cousin, H., McLinden, G.P., Craig Albertson, R., 2014. A nonsynonymous mutation
in the transcriptional regulator lbh is associated with cichlid craniofacial adaptation and neural
crest cell development. Mol Biol Evol 31, 3113-3124.
Powell, D.R., Williams, J.S., Hernandez-Lagunas, L., Salcedo, E., O'Brien, J.H., Artinger, K.B.,
2015. Cdon promotes neural crest migration by regulating N-cadherin localization. Dev Biol 407,
289-299.
Richardson, J., Gauert, A., Briones Montecinos, L., Fanlo, L., Alhashem, Z.M., Assar, R., Marti, E.,
Kabla, A., Hartel, S., Linker, C., 2016. Leader Cells Define Directionality of Trunk, but Not Cranial,
Neural Crest Cell Migration. Cell Rep 15, 2076-2088.
Roy, S., Wolff, C., Ingham, P.W., 2001. The u-boot mutation identifies a Hedgehog-regulated
myogenic switch for fiber-type diversification in the zebrafish embryo. Genes Dev 15, 1563-
1576.
Ruan, J., Li, H., Chen, Z., Coghlan, A., Coin, L.J., Guo, Y., Heriche, J.K., Hu, Y., Kristiansen, K., Li, R.,
Liu, T., Moses, A., Qin, J., Vang, S., Vilella, A.J., Ureta-Vidal, A., Bolund, L., Wang, J., Durbin, R.,
2008. TreeFam: 2008 Update. Nucleic Acids Res 36, D735-740.
Sanchez-Arrones, L., Cardozo, M., Nieto-Lopez, F., Bovolenta, P., 2012. Cdon and Boc: Two
transmembrane proteins implicated in cell-cell communication. Int J Biochem Cell Biol 44, 698-
702.
Santiago, A., Erickson, C.A., 2002. Ephrin-B ligands play a dual role in the control of neural crest
cell migration. Development 129, 3621-3632.
Scarpa, E., Szabo, A., Bibonne, A., Theveneau, E., Parsons, M., Mayor, R., 2015. Cadherin Switch
during EMT in Neural Crest Cells Leads to Contact Inhibition of Locomotion via Repolarization of
Forces. Dev Cell 34, 421-434.

Schneider, V.A., Granato, M., 2006. The myotomal diwanka (lh3) glycosyltransferase and type
XVIII collagen are critical for motor growth cone migration. Neuron 50, 683-695.
Schweitzer, J., Becker, T., Lefebvre, J., Granato, M., Schachner, M., Becker, C.G., 2005. Tenascin-
C is involved in motor axon outgrowth in the trunk of developing zebrafish. Dev Dyn 234, 550-
566.
Shellard, A., Mayor, R., 2021. Collective durotaxis along a self-generated stiffness gradient in
vivo. Nature 600, 690-694.
Shellard, A., Szabo, A., Trepat, X., Mayor, R., 2018. Supracellular contraction at the rear of
neural crest cell groups drives collective chemotaxis. Science 362, 339-343.
Snow, C.J., Henry, C.A., 2009. Dynamic formation of microenvironments at the myotendinous
junction correlates with muscle fiber morphogenesis in zebrafish. Gene Expr Patterns 9, 37-42.
Sofroniew, N., Lambert, T., Evans, K., Nunez-Iglesias, J., Bokota, G., Peña-Castellanos, G.,
Winston, P., Yamauchi, K., Bussonnier, M., Pop, D.D., Liu, Z., ACS, Pam, alisterburt, Buckley, G.,
Sweet, A., Gaifas, L., Rodríguez-Guerra, J., Migas, L., Hilsenstein, V., Bragantini, J., Lee, G.R.,
Hector, Freeman, J., Boone, P., Lowe, A.R., Gohlke, C., Royer, L., PIERRÉ, A., Har-Gil, H., 2019.
napari: a multi-dimensional image viewer for python.
Song, J.Y., Holtz, A.M., Pinskey, J.M., Allen, B.L., 2015. Distinct structural requirements for
CDON and BOC in the promotion of Hedgehog signaling. Dev Biol 402, 239-252.
Szabo, A., Melchionda, M., Nastasi, G., Woods, M.L., Campo, S., Perris, R., Mayor, R., 2016. In
vivo confinement promotes collective migration of neural crest cells. J Cell Biol 213, 543-555.
Tenzen, T., Allen, B.L., Cole, F., Kang, J.S., Krauss, R.S., McMahon, A.P., 2006. The cell surface
membrane proteins Cdo and Boc are components and targets of the Hedgehog signaling
pathway and feedback network in mice. Dev Cell 10, 647-656.
Theveneau, E., Mayor, R., 2012. Neural crest delamination and migration: from epithelium-to-
mesenchyme transition to collective cell migration. Dev Biol 366, 34-54.
Tobin, J.L., Di Franco, M., Eichers, E., May-Simera, H., Garcia, M., Yan, J., Quinlan, R., Justice,
M.J., Hennekam, R.C., Briscoe, J., Tada, M., Mayor, R., Burns, A.J., Lupski, J.R., Hammond, P.,
Beales, P.L., 2008. Inhibition of neural crest migration underlies craniofacial dysmorphology and
Hirschsprung's disease in Bardet-Biedl syndrome. Proc Natl Acad Sci U S A 105, 6714-6719.
Vega-Lopez, G.A., Cerrizuela, S., Aybar, M.J., 2017. Trunk neural crest cells: formation, migration
and beyond. Int J Dev Biol 61, 5-15.
Vega-Lopez, G.A., Cerrizuela, S., Tribulo, C., Aybar, M.J., 2018. Neurocristopathies: New insights
150 years after the neural crest discovery. Dev Biol 444 Suppl 1, S110-S143.
Williams, J.S., Hsu, J.Y., Rossi, C.C., Artinger, K.B., 2018. Requirement of zebrafish pcdh10a and
pcdh10b in melanocyte precursor migration. Dev Biol 444 Suppl 1, S274-S286.
Wolff, C., Roy, S., Ingham, P.W., 2003. Multiple Muscle Cell Identities Induced by Distinct Levels
and Timing of Hedgehog Activity in the Zebrafish Embryo. Current Biology 13, 1169-1181.
Yao, S., Lum, L., Beachy, P., 2006. The ihog cell-surface proteins bind Hedgehog and mediate
pathway activation. Cell 125, 343-357.
Yin, J., Lee, R., Ono, Y., Ingham, P.W., Saunders, T.E., 2018. Spatiotemporal Coordination of FGF
and Shh Signaling Underlies the Specification of Myoblasts in the Zebrafish Embryo. Dev Cell 46,
735-750 e734.
Yu, H.H., Moens, C.B., 2005. Semaphorin signaling guides cranial neural crest cell migration in
zebrafish. Dev Biol 280, 373-385.

999 Zeller, J.r., Granato, M., 1999. The zebrafish diwanka gene controls an early step of motor
1000 growth cone migration. Development 126, 3461-3472.
1001 Zhang, J., Lefebvre, J.L., Zhao, S., Granato, M., 2004. Zebrafish unplugged reveals a role for
1002 muscle-specific kinase homologs in axonal pathway choice. Nat Neurosci 7, 1303-1309.
1003 Zhang, W., Hong, M., Bae, G.U., Kang, J.S., Krauss, R.S., 2011. Boc modifies the
1004 holoprosencephaly spectrum of Cdo mutant mice. Dis Model Mech 4, 368-380.
1005 Zhang, W., Kang, J.S., Cole, F., Yi, M.J., Krauss, R.S., 2006. Cdo functions at multiple points in the
1006 Sonic Hedgehog pathway, and Cdo-deficient mice accurately model human holoprosencephaly.
1007 Dev Cell 10, 657-665.
1008

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

WT → WT

Cdon^co1019;Boc^co25 → WT

A 25uM

B e ba1 ba2 pa3-5 50uM

C c' 25uM

C'

D D' 25uM

D'

E e ba1 ba2 pa3-5 50uM

F ba1 ba2 pa3-5 50uM

G tg(Unc45b::eGFP) tol2 -2066 Unc45b Enhancer EGFP tol2

tg(Unc45b::smoa1-eGFP) tol2 -2066 Unc45b Enhancer smoa1 (activated smo) EGFP tol2

14 hpf/12 somites Adaxial Cells DAPI tg(Unc45b:eGFP) 100um

H tg(unc45b::eGFP) Cdon^co1019;Boc^co25 F59 tg(sox10:mRFP) 20 µm

I NCC Stream Present NCC Stream Partially Present No NCC Stream

Number of F59 Positive Muscles Per Segment

12.5 10.0 7.5 5.0 2.5

tg(Unc45b::eGFP) No Rescue Partial Rescue tg(Unc45b::smoa1-EGFP)

tg(unc45b::*smoa1*-eGFP)

J **No Rescue** (Left Side Yolk Extension) Cdon^co1019;Boc^co25 p * * Same Individual F59 tg(sox10:mRFP) 25uM

K **Rescue** (Right Side Yolk Extension) Cdon^co1019;Boc^co25 Same Individual F59 tg(sox10:mRFP) 25uM

L **Rescue** Cdon^co1019;Boc^co25 * F59 tg(unc45b:smoa1-EGFP) tg(sox10:mRFP) 25uM

Figure 7