1

cdon and _boc_ affect trunk neural crest cell migration through a non-cell autonomous

reduction of hedgehog signaling in zebrafish slow-twitch muscle

Ezra Lencer PhD[1,2]

Rytis Prekeris PhD[2,3]

Kristin Artinger PhD[1,3]

1. Department of Craniofacial Biology, University of Colorado, United States

2. Department of Cell and Developmental Biology, University of Colorado, United States

3. Co-corresponding authors

Abstract

The immunoglobin superfamily members *cdon* and *boc* are transmembrane proteins implicated in regulating hedgehog signaling during vertebrate development. Recent work showing roles for these genes in axon guidance and neural crest cell migration further suggest that *cdon*/*boc* may play additional functions in regulating directed cell movements during development. Here we use novel and existing mutants to investigate a role for *cdon* and *boc* in zebrafish neural crest cell migration. We find that single *cdon* or *boc* mutant embryos exhibit normal neural crest phenotypes, but that neural crest migration is strikingly disrupted in double *cdon*/*boc* mutant embryos. We further show that this neural crest migration phenotype is associated with defects to the differentiation of slow-twitch muscle cells, and that this slow-twitch muscle phenotype is a consequence of reduced hedgehog signaling in mutant fish. While neural crest migratory ability is not affected in double mutant embryos, neural crest directionality is severely affected. These data suggest that neural crest migration defects are likely to be secondary to defects in slow-twitch muscle differentiation. Combined, our data add to a growing literature showing that *cdon* and *boc* act synergistically to promote hedgehog signaling during vertebrate development, and provide a foundation for using zebrafish to further study the function of these hedgehog receptor paralogs.

Introduction

During vertebrate development, neural crest cells (NCCs) migrate ventrally from the dorsal neural tube along conserved pathways in the embryo to contribute to numerous traits

such as the craniofacial skeleton, peripheral nervous system, and pigment (Theveneau and

Mayor, 2012). Defects in NCC migration result in multiple human hereditary disorders (Tobin et

al., 2008; Vega-Lopez et al., 2018), and modifications to NCC migration is associated with the

evolution of morphological diversity among some vertebrate taxa (Powder et al., 2014).

Further, diseases, such as metastatic melanoma, have been shown to progress through a

reactivation of a NCC-like migratory genetic program (Diener and Sommer, 2021; Kaufman et

al., 2016). Thus understanding the genetic regulation of NCC migration is fundamental to

understanding the etiology to multiple human diseases as well as a major development event

that occurs in every vertebrate embryo.

Current thinking stresses interactions among multiple mechanisms that when combined

coordinate NCC migration along highly conserved routes in the embryo (Mayor and Etienne-

Manneville, 2016). These include NCC intrinsic factors, such as genes regulating cell adhesion

and actin cytoskeleton (Clay and Halloran, 2011, 2013, 2014; Williams et al., 2018), as well as

cell extrinsic factors that affect the extracellular matrix (Perris and Perissinott, 2000; Szabo et

al., 2016), tissue stiffness (Shellard and Mayor, 2021), and chemical gradients (Barriga et al.,

2013; Olesnicky Killian et al., 2009; Shellard et al., 2018), which attract and funnel NCCs along

conserved paths in the embryo. Mediating cell intrinsic and extrinsic regulation of NCC

migration are cell-cell signaling interactions (Carmona-Fontaine et al., 2008; Mayor and

Etienne-Manneville, 2016; Scarpa et al., 2015; Szabo et al., 2016; Theveneau and Mayor, 2012).

These include the chemical gradients that attract and inhibit migrating NCCs, as well as the cell-

cell adhesion molecules that adhere NCCs to each other and mediate many cell-cell interactions

(Carmona-Fontaine et al., 2008; Mayor and Etienne-Manneville, 2016; Scarpa et al., 2015;

Szabo et al., 2016; Theveneau and Mayor, 2012; Vega-Lopez et al., 2017).

Recent work has implicated a role for the immunoglobin superfamily member protein

cell adhesion associated, oncogene regulated (*cdon*) as a cell surface protein regulating trunk

NCC (tNCC) migration in zebrafish (Powell et al., 2015). Morpholino knockdown of *cdon* results

in failed (tNCC) migration, with tNCCs mislocalizing N-cadherin and arresting migration at the

body midline. *cdon* and its paralog *BOC, brother of cell adhesion associated, oncogene*

regulated (*boc*), are vertebrate members of a larger gene family with identifiable orthologs in

other metazoan taxa that includes the drosophila paralogs *ihog* and *boi*. Recent work on both

vertebrate *cdon*/*boc* and *Drosophila ihog*/*boi* has established a role for these transmembrane

proteins in activating hedgehog signaling. *cdon* and *boc* act as co-receptors binding to hedgehog

ligand and patched receptor in order to promote smoothened signaling within cells. Other work

suggests that *cdon*/*boc* may commonly interact with additional signaling pathways including

Wnt signaling and Nodal signaling (Hong et al., 2020; Jeong et al., 2017). Further, *cdon* has been

shown to form a complex with N-cadherin/Cdc42/Bnip-2 in a hedgehog independent manner to

regulate actin cytoskeletal dynamics during myogenic differentiation in cell culture (Kang et al.,

2008; Lu and Krauss, 2010). *boc* has been shown to play a role in axon guidance, where it

operates as a receptor for hedgehog ligand to regulate proper neuronal growth (Connor et al.,

2005; Okada et al., 2006). Both *cdon*/*boc* and *ihog*/*boi* are often localized to the ends of long

membrane extensions (e.g. cytonemes in *Drosophila*), suggesting that these receptors may help

cells sense the wider external environment and communicate over long distances (Bilioni et al.,

2013; Ferent et al., 2019; Hall et al., 2021). It is thus possible that these transmembrane

proteins are reiteratively used as cell surface proteins to integrate multiple signals during

processes such as cell migration and axon guidance. While a role for *cdon* in tNCC migration has

been suggested (Powell et al., 2015), the function of *cdon*/*boc* in regulating NCC movements is

not fully understood.

In zebrafish, hedgehog signaling is currently thought to affect tNCC migration in a non-

cell autonomous manner by regulating the differentiation of adaxial cell mesoderm (Honjo and

Eisen, 2005). Adaxial cells form at the border of the notochord during the end of gastrulation

and differentiate into slow-twitch muscle cells that migrate laterally through the somatic

mesoderm (Elworthy et al., 2008; Wolff et al., 2003). A non-migratory population of adaxial cell

derivatives, called muscle pioneers, remain at the body midline. In hedgehog mutant embryos,

adaxial cells fail to differentiate into slow-twitch muscle/muscle pioneer derivatives (Elworthy

et al., 2008; Roy et al., 2001; Yin et al., 2018). Through a mechanism not fully understood, but

likely to depend on extracellular matrix deposition/modification (Banerjee et al., 2011;

Banerjee et al., 2013; Guillon et al., 2016; Honjo and Eisen, 2005), this slow-twitch muscle

differentiation is critical for normal NCC migration. Whether *cdon*/*boc* affect slow-twitch

muscle differentiation, however, is unknown (Bergeron et al., 2011), and it is further possible

that *cdon*/*boc* affect NCC development through NCC autonomous in addition to non-cell

autonomous mechanisms.

Here we use novel CRISPR/Cas9 generated mutations and existing mutants to

investigate the role of *cdon* and *boc* in tNCC migration and slow-twitch muscle differentiation.

We show that *cdon* and *boc* are expressed by tNCCs at low levels and by the adaxial mesoderm

at moderate levels. Loss of either *cdon* or *boc* individually has no observable effect on tNCC

migration nor slow-twitch muscle differentiation. However, embryos that are double mutant

for both *cdon* and *boc* exhibit both tNCC migratory defects as well as loss of slow-twitch muscle

mesoderm. We show that tNCC mobility is not impaired by loss of *cdon*/*boc* suggesting that the

tNCC migration defect is a non-cell autonomous consequence of slow-twitch muscle

differentiation. We further provide data suggesting that *cdon*/*boc* are modifying hedgehog

signal reception during mesoderm differentiation. Combined, our data contribute to a growing

literature showing that *cdon*/*boc* play a critical role in modulating hedgehog signaling across

multiple tissues during embryogenesis.

Results

***cdon/boc* paralogs exhibit shared sequence conservation in extracellular functional domains**

The transmembrane robo-related proteins, *cdon* and *boc*, share a similar genomic

domain architecture that includes 4 immunoglobin (IG) domains, 3 fibronectin type 3 repeat

(FNIII) domains, and a cytosolic C-terminal tail (Figure 1A). A fifth IG domain is variable present

in *cdon* orthologs of tetrapod taxa (grey in Figure 1A). The FNIII domains have been implicated

in binding to hedgehog ligand and patched receptor (Izzi et al., 2011; Song et al., 2015), and

some evidence suggests that the c-terminal region is needed for protein localization in the cell

(Okumura et al., 2021).

As functional conservation is likely to be reflected by sequence conservation across

evolutionary time, we investigated the molecular evolution of *cdon*/*boc* orthologs across

vertebrate taxa. An alignment of *cdon* and *boc* amino acid sequences from 11 select jawed

vertebrate taxa shows that extracellular domains exhibit high conservation across taxa and

paralogs, suggesting that these extracellular domains are critical for gene function (Figure 1A).

In contrast, the C-terminal cytosolic tail exhibits relatively low sequence conservation (Kang et

al., 2002) suggesting that cytosolic region is evolutionarily labile to different selection

pressures.

Identifiable orthologs to *cdon* and *boc* are present in non-vertebrate bilaterian taxa.

These include *Drosophila* paralogs *ihog* and *boi*, whose function in hedgehog signaling has been

well characterized (Bilioni et al., 2013; Camp et al., 2010; Yao et al., 2006). However, the

orthology relationships between *cdon*/*boc* and non-vertebrate orthologs such as *ihog* and *boi*

are less well known. To explore the molecular evolution of this gene family we built a maximum

likelihood tree (phyml) using 44 amino acid sequences from taxa representative for major

bilaterian clades (Figure 1B). These data identify *cdon*/*boc* as shared vertebrate specific

paralogs that emerge from a duplication event at either the base of the vertebrate phylogeny

or along the gnathostome lineage following splitting from cyclostomes (hagfish and lamprey).

Importantly, these data are consistent with a hypothesis that *cdon*/*boc* originated during one of

the two vertebrate specific genome duplications (Dehal and Boore, 2005). Our analyses further

suggest that an independent gene duplication event led to the origins of *boi* and *ihog* paralogs

in drosophilids (Kang et al., 2002). Thus the co-orthologs *cdon*/*boc* and *boi*/*ihog* represent two

independent duplication events (Figure 1B)(Allen et al., 2011; Izzi et al., 2011; Sanchez-Arrones

et al., 2012), a finding that is concordant with gene trees made publicly available by the

TreeFam consortium (Li et al., 2006; Ruan et al., 2008).

Generation of cdon and boc null alleles

To further study the function of *cdon* and *boc* during zebrafish development we turned

to existing and novel mutations that are predicted to disrupt gene function (Figure 1C). Two

previously published ENU generated *boc* alleles were used in this study to characterize the

function of *boc* during zebrafish tNCC development. The *boc*/uml allele (Bergeron et al., 2011)

results in a predicted early stop (hereafter boc^{uml}). The boc^{co25} allele (Kearns et al., 2021) results

in an non-synonymous isoleucine to serine substitution in the third FNIII domain (Figure 1C). As

boc^{uml} and boc^{co25} alleles do not complement each other (Kearns et al., 2021), we focus our

analyses on boc^{co25} and note that we see similar results with boc^{uml}.

We used CRISPR/Cas9 to generate 2 novel mutations in *cdon*. Both alleles result in

predicted early stops (Figure 1C). The $cdon^{co1019}$ allele is a 229bp deletion that results in a

predicted early stop leading to a truncated protein prior to conserved IG domains. The

$cdon^{co1023}$ allele is a 5bp deletion in the FNIII(2) domain that similarly results in a predicted early

stop and a truncated protein prior to the transmembrane domain. We focus on the $cdon^{co1019}$

allele and note that we see similar results with the $cdon^{co1023}$ allele.

cdon and boc transcripts are expressed in tNCCs and slow-twitch muscle cells

We next used multiplexed hybrid chain reaction *in situ* hybridization (HCR) in order to

ask where *cdon* and *boc* transcripts are expressed, and whether transcripts are detectable in

tNCCs and slow-twitch muscle progenitors. Unlike traditional enzymatic methods for *in situ*

hybridization, HCR allows single molecule measurement of gene expression at cellular

resolution, which enables identifying cells and cell types that express transcripts at relatively

low levels (Kearns et al., 2021).

We found *cdon* and *boc* transcripts to be abundant in the dorsal central nervous system

(CNS) of 24hpf-48hpf zebrafish embryos (Figure 2A-D), consistent with similar data from prior

studies (Bergeron et al., 2011; Cardozo et al., 2014; Kearns et al., 2021). This pattern is

established early in development. At 2 somites, *cdon* is expressed in the presumptive

notochord, while both *cdon* and *boc* are both expressed along the border of the neural plate in

cells that we suspect contribute to the dorsal neural tube at 24hpf (Figure 2A). Importantly,

while *boc* transcripts in the neural tube form a broad dorsal-ventral gradient (Kearns et al.,

2021), *cdon* transcripts are restricted to the dorsal-most roofplate cells and the ventral most

floorplate cells of the neural tube by 24hpf. In this way, *cdon* creates a dorsal sub-domain of

boc expression in the neural tube (Figure 2D). Floorplate cells also express hedgehog ligand,

and thus unlike *boc*, *cdon* is expressed in both hedgehog ligand producing and non-producing

cells within the CNS. This neural tube expression at 24hpf is broadly similar to expression

patterns reported in mouse embryos (Izzi et al., 2011; Kang et al., 2002; Mulieri et al., 2002;

MULIERI et al., 2000; Tenzen et al., 2006) suggesting that *cdon*/*boc* function in the CNS is likely

to be conserved.

We next asked whether migrating tNCCs express *cdon* and *boc*. Using the

tg(*sox10*:mRFP) transgene, we FAC sorted NCCs from the dissected trunks of 24hpf zebrafish

embryos and performed bulk RNA sequencing. These data suggest that tNCCs do express *cdon*

and *boc* at low levels (Figure 2E). We then asked if this expression is observable by HCR. Using

the tg(*sox10*:mRFP) transgene to again label NCCs, we segmented out *cdon* and *boc* HCR puncta

that overlap the *sox10* transgene. These data identify *cdon*/*boc* transcripts expressed by NCCs

at 24hpf (Figure 2F&G). While these data indicate that *cdon* and *boc* are expressed by tNCCs,

we note that tNCCs are expressing *cdon*/*boc* at levels far lower than the surrounding tissues.

Of note is that mesoderm appears to express *cdon* and *boc* at relatively high levels

(Figure 2), and prior studies had suggested that *boc* is expressed by slow-twitch muscle

mesoderm at 24hpf (Bergeron et al., 2011). We thus, next sought to identify whether *cdon*/*boc*

transcripts are in fact expressed by the developing slow-twitch muscle/adaxial cell mesoderm.

To do this we used the tgBAC(*prdm1a*:eGFP) transgene to label adaxial cells and slow-twitch

muscle mesoderm (Elworthy et al., 2008). At the 2 somite stage, adaxial cells are labeled with

the *prdm1a*:eGFP transgene, but we did not observe significant expression of *cdon*/*boc* in these

adaxial cells (Figure 2H). Instead, *cdon* and *boc* transcripts were largely restricted to the

developing neural plate and notochord (see above). By the 10 somite stage, *cdon* and *boc*

transcripts were observed throughout the paraxial mesoderm. Using the tg(*prdm1a*:eGFP)

transgene to segment out HCR transcripts expressed by adaxial cells, we were able to confirm

that *cdon* and *boc* are expressed by slow-twitch muscle progenitor cells (Figure 2I-J). At 24hpf,

differentiating slow-twitch muscle cells continue to express both *cdon* and *boc* (Figure 2K-L).

These data are consistent with bulk RNA-seq data from FAC-sorted trunk tgBAC(prdm1a:egFP)

positive cells at 24hpf (Figure 2E).

Taken together, our analyses indicate that *cdon* and *boc* are broadly expressed by

mesodermal tissues that include adaxial cell derived slow-twitch muscle cells, as well as tNCCs

at low levels, in addition to the developing neural tube.

Trunk neural crest migration and adaxial mesoderm development are affected in cdon/boc

double mutants

We next asked whether *cdon* and *boc* mutations affect tNCC migration and slow-twitch

muscle development by using the tg(*sox10*:mRFP) line to label migrating tNCCs and

immunostaining of a slow-twitch muscle specific myosin (DSHB F59) to label adaxial mesoderm

derivatives in 24 hpf embryos.

In wildtype embryos, tNCCs collectively migrate in streams down the middle of each

body segment (Figure 3). We found that this tNCC migratory phenotype was largely normal in

both single *cdon* and *boc* mutant embryos (Figure 3D-E). These data from homozygous mutant

embryos are in contrast to those reported by morpholino knockdown of *cdon* where tNCC

migratory streams fail to form (Powell et al., 2015). However, in double cdon^{co1019};boc^{co25}

mutants, tNCCs did fail to form distinguishable streams, and tNCC migration arrested at the

ventral neural tube/body midline (Figure 3F), a phenotypes that is similar to that observed in

cdon morphant embryos (Powell et al., 2015).

Similar to tNCC migration, we found that slow-twitch muscle differentiation was largely

normal in *cdon* and *boc* (Bergeron et al., 2011) single mutants. Only in double mutants did we

observe defects in slow-muscle cell differentiation (Figure 3C-F). Loss of slow-twitch muscle cell

differentiation was not complete, however, and posterior body segments tended to exhibit

greater loss of slow-twitch muscle differentiation than anterior segments. In double mutant

embryos, cells that did express slow-twitch muscle specific myosin (F59) appeared dysmorphic

with loosely packed myosin bundles relative to wildtype embryos (e.g. compare myosin bundle

width and packing in Figure 3).

Thus defects in tNCC migration and slow-twitch muscle cell differentiation are only

observed in *cdon/boc* double mutants.

cdon and boc affect hedgehog signal transduction in slow-twitch muscle cells

Since *cdon* and *boc* have previously been implicated in hedgehog signaling (Allen et al.,

2011; Zhang et al., 2006), we hypothesized that defects in both tNCC migration and slow-twitch

muscle cell differentiation were due to loss of hedgehog signaling that is only apparent in the

double mutants. To test this, we measured hedgehog signal reception by using the transgenic,

tg(8xGliBS:mCherry-NLS-Odc1) reporter line (hereafter 8xGli:mCherry) that drives nuclear

localized mCherry under the control of 8 tandem Gli binding sites (Mich et al., 2014). The

mCherry fluorophore contains an Odc1 destabilization domain the facilitates proteasomal

degradation to promote rapid fluorophore turnover within expressing cells.

At 24hpf, we observed mCherry positive nuclei in ventral neural tube cells and in slow-

twitch muscle fiber nuclei (Figure 3G)(Kearns et al., 2021; Mich et al., 2014). We never observed

mCherry positive nuclei in tNCCs (data not shown). In *cdon^{co1019}* and *boc^{co25}* mutants, the

number and signal intensity of nuclei that are 8xGli:mCherry positive was lower as compared to

wildtype siblings (Figure 3G-J). This decrease in signal was true for both cells in the neural tube

and slow-twitch muscle cells. Importantly, 8xGli:mCherry signal intensity varied across

genotypes, with *boc^{co25}* mutants displaying lower 8xGli:mCherry signal than *cdon^{co1019}* mutants,

and double mutants exhibiting low to no signal than all other genotypes. These data suggest

that *cdon^{co1019}* and *boc^{co25}* mutations are decreasing hedgehog signal reception in slow-twitch

muscle cells, which is consistent with a hypothesis that perturbations to hedgehog signaling is

responsible for the slow-twitch muscle differentiation phenotype in *cdon* and *boc* mutant

embryos (Barresi et al., 2000; Devoto et al., 1996).

Interestingly, the 8xGli:mCherry transgene reported substantively lower hedgehog

signal reception in single *cdon*^{co1019} and *boc*^{co25} mutant embryos that exhibit normal slow-twitch

muscle differentiation. These data suggest that slow-twitch muscle development is robust to

moderate perturbations to hedgehog signal. If this hypothesis is correct, we reasoned that *cdon*

and *boc* mutants should sensitize embryos to further pharmacological manipulation of

smoothened signaling with cyclopamine. To test this, we in-crossed double heterozygous

cdon$^{co1019/+}$;boc$^{co25/+}$ parents and treated embryos with 20uM cyclopamine from 50% epiboly to

24hpf. We then quantified the number of slow-twitch muscle fibers at 24hpf as a readout for

hedgehog signaling.

While single *boc* and *cdon* mutants treated with ethanol as a controls had a similar

number of slow-twitch muscle fibers as their wildtype ethanol treated siblings, *cdon* and *boc*

mutants treated with 20uM cyclopamine had significantly fewer slow-twitch muscle fibers than

their similarly cyclopamine treated wildtype siblings (Figure 4). This pattern was even more

striking for the double mutant. While double mutants treated with ethanol have an average of

approximately 5-6 slow-twitch muscle fibers per segment, double mutants lost all slow-twitch

muscle fibers when treated with 20uM cyclopamine. Furthermore, we noted qualitative

changes to slow-twitch muscle morphology in cyclopamine treated *cdon* and *boc* embryos.

Mutants treated with 20uM cyclopamine often exhibited gaps or missing slow-twitch muscle

fibers in body segments, and slow-twitch muscle fibers often appeared to have loosely packed

myosin.

We interpret these results to indicate that *cdon* and *boc* are modifying hedgehog signal

reception by slow-twitch muscles, and that loss of *cdon* or *boc* sensitize embryos to further

perturbations of hedgehog signal.

<u>*Trunk neural crest cells lose directionality but not migratory ability in cdon/boc mutants*</u>

The arrested NCC migratory phenotype in the double mutants is striking and could result from

either a loss of NCC migratory ability or a loss of directionality. To investigate this further we

used the tg(*sox10*:mRFP) transgene to label migrating tNCCs and live imaged NCC migration

through the trunk of the zebrafish embryo beginning at 22hpf (Figure 5). Migrating tNCCs were

manually tracked in imageJ using the mTrackJ plugin (Meijering et al., 2012). As we only

observed a tNCC migratory defect in the medial migrating streams, we limited our tracking of

cells to NCCs migrating along the medial route. We found that tNCC mobility was not disrupted

in *cdon*, *boc*, or double mutants as compared to wildtype controls. Across all genotypes, tNCCs

were qualitatively migratory, and general measures of mobility (e.g. Max velocity etc.) in

mutants were similar to that observed in wildtype controls (Figure 5D-G). In contrast, we

observed a significant loss of directional NCC movement in double cdon[co1019];boc[co25] mutant

embryos. In double mutants, tNCCs migrate to the body midline and then often move along the

anterior-posterior axis (Figure 5A-C), which is in contrast to the directed ventral movement

typical of wildtype embryos. This is seen quantitatively as lower migratory directionality and an

increased proportion of movements in lateral directions (Figure 5B&H). Single *cdon* or *boc*

mutants did not display migratory phenotypes that were different from that of wildtype

controls. Taken together, we interpret these results to indicate that loss of *cdon* and *boc* affects

tNCC directionality and path-finding ability, but that *cdon* and *boc* are not required for tNCC

mobility.

cdon and boc mutants exhibit hedgehog-like craniofacial phenotypes

While our data suggest that *cdon* and *boc* affect tNCC migration, we did not observe any

defects in cranial NCC migration (data not shown). However we did observe severe defects to

cranial cartilage morphology in *boc* and double mutants that are suggestive of other hedgehog

mutants. Homozygous *cdon*^{co1019} and *cdon*^{co1023} mutants, however, do not display noticeable

phenotypes (Figure 6). Further, we recovered homozygous *cdon* mutant adults that are fully

viable. While these *cdon* alleles are apparently not deleterious to organismal viability, we note

that when in combination with *boc* mutations these *cdon* alleles produce severe phenotypes

(see below) supporting the prediction that these alleles disrupt normal *cdon* function based on

the introduction of early stop codons.

In contrast to mutations in *cdon*, homozygous mutant *boc* embryos display striking

craniofacial and curved body phenotypes that are characteristic of reduced hedgehog signaling

(Bergeron et al., 2011). In the developing head, homozygous *boc*^{co25} and *boc*^{uml} mutants display

neurocranial defects with fused trabecular cartilages (Figure 6A). Defects in the development of

the neurocranial cartilages are classic midline phenotypes associated with mutations in

hedgehog pathway genes (Bergeron et al., 2011; Eberhart et al., 2006; Wada et al., 2005).

Interestingly, while mutations in smoothened or hedgehog ligand lead to complete loss of the

anterior ethmoid plate and fusion of the posterior trabecular cartilages (Eberhart et al., 2006;

Swartz et al., 2012; Wada et al., 2005), loss of *boc* only results in a relatively mild fusion of the

trabecular cartilages. Further, *boc* mutants have shortened or absent symplectic cartilages and

a striking ventral oriented ceratohyal cartilage that attaches at an obtuse angle to the head

(Figure 6). In the smoothened mutant (Swartz et al., 2012) and Dispatched 1 mutant (Schwend

and Ahlgren, 2009) a rudimentary ceratohyal projects ventrally (Swartz et al., 2012), suggesting

this mis-orientation of the ceratohyal is a canonical hedgehog phenotype. However, unlike in

these other mutants where the ceratohyal is greatly hypomorphic and jaw joints are often

fused, in *boc* mutants the ceratohyal is correctly formed and joints appear normal. To ask if this

ceratohyal orientation defect was secondary to defects in soft tissue we stained craniofacial

muscles with phalloidin and used the tg(*sox10*:mRFP) transgene to label NCC derived cartilages

(Figure 6B). Craniofacial muscles appear normal, suggesting that the orientation of the

ceratohyal in the *boc* mutant is due to cartilage element patterning and growth and not a

secondary defect resulting from disrupted muscle differentiation and or attachment.

Double mutants homozygous for both *cdon*^{co1019} and *boc*^{co25} exhibit more extreme

phenotypes consistent with a hypothesis that functional disruption of both *boc* and *cdon* leads

to a more dramatic loss of hedgehog signal reception. In double mutants, the symplectic is

absent, all craniofacial elements are hypomorphic, and neurocranial defects are more severe.

Double mutants variably present mild to severe forms of cardiac edema and typically have

more curved bodies than that observed in the *boc* mutants alone. We note that despite having

more severe phenotypes, the ethmoid plate was present in all *cdon/boc* double mutant

embryos observed, a curious observation given that the ethmoid plate is often lost in other

hedgehog mutants. Studied *cdon* and *boc* mutations do not complement each other and

cdon^{co1019}/*cdon*^{co1023};*boc*^{uml}/*boc*^{co25} embryos present phenotypes similar to homozygous

cdon^{co1019};*boc*^{co25} mutant embryos (Figure 6E). Taken together these data indicate that

cdon/boc mutants display phenotypes similar to other characterized hedgehog mutants

supporting the hypothesis that *cdon* and *boc* are modifying hedgehog signaling in the

developing embryo head. That double *cdon/boc* mutants have more severe phenotypes and

supports a hypothesis that *cdon/boc* paralogs have synergistic function in promoting hedgehog

signal reception.

Discussion

Here we show that *cdon* and *boc* affect trunk NCC migration as well as differentiation of

slow-twitch muscle cells. Despite some prior data suggesting that *cdon* may act cell

autonomously to affect tNCC migration, we find that neither *cdon* nor *boc* mutations have a

noticeable effect on NCC mobility. Rather, only in double *cdon/boc* mutants do we observe a

noticeable NCC migratory defect that is associated with loss of slow-twitch muscle mesoderm.

Prior research has shown that slow-twitch muscle differentiation requires hedgehog signaling

(Elworthy et al., 2008), and our data show that this slow-twitch muscle phenotype is due to a

reduction in hedgehog signaling within *cdon/boc* double mutants. Taken together we interpret

our data to suggest that a tNCC migratory defect in *cdon/boc* double mutants is a consequence

of defects in mesodermal differentiation (Banerjee et al., 2011; Banerjee et al., 2013; Honjo and

Eisen, 2005).

Our data add to a growing literature showing that *cdon* and *boc* act synergistically

during zebrafish development to modify hedgehog signal reception and regulate the

development of multiple traits (Echevarria-Andino and Allen, 2020; Song et al., 2015; Zhang et

al., 2011). Interestingly we found that *cdon* mutations alone do not produce any noticeable

phenotypes in zebrafish, and we are able to recover adult homozygous mutant zebrafish. This is

contrast to mouse, where loss of CDON results in mild holoprocephaly (Allen et al., 2011; Allen

et al., 2007; Echevarria-Andino and Allen, 2020; Zhang et al., 2011). While it is possible that our

cdon mutations do not result in full loss of function, we find this possibility unlikely. Both *cdon*

mutations lead predicted early stops prior to the transmembrane domain, which is similar to

the *boc^{uml}* mutation that does lead to a severe loss of function phenotype. Further, our *cdon*

mutants do impact *cdon* function as double *cdon*/*boc* mutants exhibit extreme phenotypes

suggestive of loss of *cdon* function. As both mutations lead to an early stop prior to the

transmembrane domain, production of any truncated *cdon* protein in these mutants should

lead to an extracellular pool of *cdon* that would be more likely to result in a dominant negative

phenotype than in a mild hypomorphic allele. However, we cannot fully rule out the possibility

that a late start rescues *cdon* protein resulting in a hypomorphic but functional receptor.

Unfortunately antibodies do not reliably recognize zebrafish *cdon* protein, and thus future

research could investigate whether other alleles targeting the FNIII(3) domain might result in

more severe phenotypes.

An alternative possibility, however, is that *cdon* and *boc*, as modifiers of hedgehog

signal, interact with other hedgehog pathway members resulting in species (or fish/mouse line

specific) phenotypic responses to *cdon* mutations. Thus in zebrafish other hedgehog pathway

members may provide robustness against *cdon* mutations allowing generally normal

phenotypic development. This hypothesis is supported by our findings that *cdon* and *boc*

mutations both sensitize embryos to cyclopamine treatment. Thus presumably *cdon* mutations

also sensitize embryos to genetic perturbations such as loss of other hedgehog pathway

members (or modifications to their gene expression). Intriguingly, we observed that *cdon* gene

expression is often nested within *boc* expression suggesting that *cdon* and *boc* act together to

modify hedgehog morphogen gradients. Kearns et al. 2021 found that *patched-2* (*ptch2*)

expression in the dorsal neural tube is expanded in *boc*[co25] mutant embryos, suggesting that

loss of *boc* alone leads to more dorsal hedgehog reception (e.g. *ptch2* is a readout for hedgehog

signaling). These data were interpreted to suggest that *boc* sequesters hedgehog ligand driving

hedgehog signal reception to ventral tissues. Expanding on this hypothesis, we note that *cdon*

transcripts are positioned even more distally in the neural tube from hedgehog ligand source

than are *boc* transcripts. Thus together *boc* and *cdon* gradients may act as strong sequestering

agents to limit hedgehog signal activation to proper tissues and inhibit ligand activity in dorsal

neural tube cells.

Intriguingly, we and others note that *cdon* and *boc* are lowly expressed in a broad range

of tissues including the mesoderm, NCCs among others. Further *cdon*, but not *boc*, transcripts

are also co-expressed in hedgehog ligand producing cells, and some studies have suggested

that *cdon* plays a role in presenting hedgehog ligand to responsive cells (Hall et al., 2021). Thus

it is likely that *cdon* and *boc* are playing different roles in different tissues, and that cellular

context may be critical for determining the biological function of these receptors.

Here we present an early investigation into the phenotypes of *cdon* and *boc* mutations

in zebrafish as a foundation to further study the interact roles of these two paralogs. Future

work dissecting how these genes operate in different cellular contexts will address the diverse

roles of these genes in hedgehog pathway function.

Materials and Methods

410   *Fish stocks, husbandry, and generation of mutants*

411   Adult fish and embryos were maintained at the University of Colorado Anschutz Medical

412   school according to standard zebrafish rearing protocols and guidelines for the rearing and

413   maintenance of vertebrate animals. The boc[co25] mutant was previously recovered from an ENU

414   screen (Kearns et al., 2021). The boc[uml] mutant was acquired from the Zebrafish International

415   Resource Center (boc[ty54z]) and was described by (Bergeron et al., 2011).

416   We used CRISPR/Cas9 to generate novel mutations in *cdon*. Two sets of guide RNAs

(sgRNA) were designed to target either the 2nd and 3rd exons of cdon

(ggcagUUgaggagcagagUa, UggUaggagccUgUgagagc, UgcUUcagaaUcgcUgagag) or to target the

11[th] exon (UcgccaUUgagaUUgUgggg, cUccacgcgaaaUgcagUga). DNA oligos were annealed to a

common backbone scaffold that contains a T7 RNA polymerase site, which was used to

generate sgRNAs used for injections (Williams et al., 2018). For each sgRNA set, our final CRISPR

injection solution consisted of sgRNAs diluted to 20ng/uL, 1ug Cas9 Protein (Invitrogen TrueCut

V2), and 5ug Rhodamine Dextran in a total volume of 10uL. Approximated 2-4nl were injected

into the cell of a 1cell stage embryo, and F0 injected embryos were reared to adulthood. We

recovered a single allele for each sgRNA set. These include the cdon[co1019] allele, which is a

229bp deletion that spans the second to third exon of *cdon*, removes the intronic sequence

between these exons, and is predicted to result in an early stop. The cdon[co1023] allele is a 5bp

deletion resulting in a predicted early stop in the 11[th] exon that codes for the second FNIII

domain. F0 injected embryos were outcrossed for a minimum of two generations to minimize

effects of off target cutting.

Zebrafish *boc* and *cdon* mutants were crossed in to existing transgenic reporter lines

including the tg(*sox10*(7.2kb):mRFP)[vu234] that labels NCCs with cytosolic mRFP (Kucenas et al.,

2008), and the gli reporter tg(8xgli-Xla.Cryaa:NLS-d1mCherry) line that has 8 gli-binding sites

driving a destabilized nuclear localized mCherry (Mich et al., 2014). The tgPAC(*prdm1a*:EGFP)

line was used to label adaxial mesoderm for reference (Elworthy et al., 2008).

cdon/boc molecular conservation and gene tree

Amino acid coding sequences for select bilaterian taxa were downloaded from NCBI GenBank.

Orthologous sequences were searched directly or identified through BLAST searches and

manual curation. The final set of sequences was chosen to include taxonomic representatives

for major bilaterian taxa for which high confidence sequences are available. Effort was made to

identify two paralogs for each taxon. Amino acid sequences were aligned using MUSCLE

(v3.8.31) with default parameters (Edgar, 2004). Amino acid conservation (Figure 1A) was

calculated by pairwise comparisons of *cdon*/*boc* homologues from 11 taxa (human, mouse,

chicken, *Anolis*, frog, bichir, Gar, sheepshead minnow, medaka, catfish, zebrafish) and plotted in

R by smoothing conservation values across 20 amino acid windows. Substitution models were

selected using ProtTest3 by AIC (Darriba et al., 2011). A maximum likelihood gene tree was built

using phyml (v3.3) and a JTT+I+G+F model on the entire unfiltered alignment with 1000

bootstraps (Guindon and Gascuel, 2003). Nodes with less than 60% bootstrap support were

collapsed into polytomies. We recovered similar results from filtered alignments that removed

poorly aligned regions and regions with gaps.

Immunohistochemistry, in situ hybridization, and stains

Adaxial mesoderm derivatives were stained by fluorescent immuno-histochemistry against a

slow-twitch muscle specific myosin (anti-MYH1A, F59, DSHB) at a concentration of 1:20 as

previously described (Elworthy et al., 2008; Nguyen-Chi et al., 2012). Hybrid Chain Reaction

(HCR) _in situ_ hybridizations were performed as previously described (Choi et al., 2018; Lencer et

al., 2021). Custom HCR probes for zebrafish _cdon_ and _boc_ were acquired from Molecular

Instruments and used at a concentration of 2pMol/500uL Hybridization Buffer. Phalloidin-488

and phalloidin-647 stains (Thermo Fisher) were used at a concentration of 1:1000 to label

muscle tissue.

Live imaging and tracking of migrating cells

For live-imaging and tracking, neural crest cells were labelled using the tg(_sox10_:mRFP) line.

Embryos were mounted in 0.4% low melt agarose and neural crest cells were imaged over the

yolk extension every 5 minutes from ~22hpf to ~30hp on an Andor spinning disk confocal

microscope at 28C. Neural crest cells were manually tracked along the xy axis from maximum

projection images using the ImageJ mTrackJ plugin (Meijering et al., 2012). Drift was accounted

for using the Correct_3D_Drift.py plugin as implemented in ImageJ (Parslow et al., 2014).

Quantitative analysis of track mobility was performed in R. A small number of short tracks with

fewer than 30 points (2.5hrs) were excluded. As we also noted that NCCs often stopped

migrating upon reaching a ventral position we further limited tracks to a maximum of 6 hours in

order to capture cell migration. Limiting all tracks to the same time (e.g. number of points)

produces similar results. Roseplots depict cell movements with a velocity greater than

0.3uM/min, which empirically determined as a cutoff for capturing cell movements while also

excluding noise from drift and changes in cell shape. Using different cutoffs produced similar

results. Linear models were built in R by including track length as a fixed effect, and embryo

identity and date of imaging as random effects, with the packages lme4 (Bates et al., 2015),

lmerTest (Kuznetsova et al., 2017), and emmeans. Images of tracks were produced in napari

(Sofroniew et al., 2019).

Pharmacological manipulation

For pharmacological manipulation, embryos were manually dechorionated and treated with

20uM cyclopamine in ethanol (5uL/mL) from 50% epiboly to 24 hpf. Embryos that appeared

developmentally delayed were excluded.

Imaging, Statitiscal Analysis, and biological replication

All imaging was performed on either a Leica Sp8 Confocal, Andor spinning disk confocal, or

Olympus BX compound microscope. Image analysis and presentation was performed in ImageJ

or napari. All statistical analyses were performed in R. Experiments were repeated at least 3

times and biological variation was accounted for either in models and through inclusion of

biological replicates.

Acknowledgements

We thank members of the Artinger and Prekeris laboratories for comments on this manuscript and project. This work was in part supported by GM122768 (to RP) and a University of Colorado RNA Biosciences Initiateive award to EL and KA. EL was supported by National Institutes of Health Ruth L. Kirschstein National Research Service Awards (T32CA17468 and F32HD103406). EL, RP, and KA conceived of the study and wrote the manuscript. EL performed research and analyses.

Figure Legends

Figure 1. *cdon* and *boc* are transmembrane receptors and share conserved domain structure. **(A)** *cdon*/*boc* proteins share a domain structure that includes 4-5 immunoglobin (IG) domains, 3 fibronectin (FNIII) domains, a transmembrane (TM) region, and a cytosolic C-terminal tail. A fifth IG domain (*) in *cdon* is found in some tetrapod taxa and indicated in grey. Percent conservation (20 amino acid sliding window) is shown for an alignment of *cdon* and *boc* amino acid sequences from 11 tetrapod and ray-finned fish taxa. Note that the extracellular domains exhibit high sequence conservation, while the cytosolic region is less conserved. **(B)** Maximum likelihood tree (phyml) for curated *cdon* and *boc* amino acid sequences from multiple bilaterian taxa. Note independent duplications for Ihog/Boi in drosophilids and *cdon*/*boc* in vertebrates. Node support was calculated based on 1000 bootstraps, and nodes with less than 60% bootstrap support were collapsed. **(C)** Schematic of *cdon* and *boc* mutant alleles that are characterized in the current study. Three alleles result in a predicted early stop, while boc[co25]

contains a single nucleotide substitution that results in a non-synonymous isoleucine to serine

substitution.

Figure 2. *cdon* **and** *boc* **are expressed by NCCs and adaxial mesoderm cells**. (A-C) HCR shows

cdon (magenta) and boc (green) expression in the developing neural tube of 2 somite **(A)**, 24

hpf **(B)**, and 48 hpf **(C)** embryos. Note *cdon/boc* expression at the border of the neural plate **(A)**,

and in the roofplate/dorsal neural tube cells at 24 and 48 hpf. Additionally, *cdon* is expressed in

the ventral floorplate cells of the neural tube and notochord **(B&C)**. Weak *cdon/boc* expression

is observed throughout the mesoderm. Panels B&C are 20uM sections at the level of the yolk

extension (YE) in the trunk. **(D)** Schematic summarizing expression of *cdon/boc* in the neural

tube from panels B&C. Gradient of hedgehog ligand produced by notochord and floorplate cells

is indicated at left in grey, while expression of *cdon/boc* are indicated by magenta and green

respectively. (E) Bulk RNA-seq from FAC-sorted trunk NCCs (blue) and slow-twitch muscle cells

(red) suggests that both tissues express *cdon/boc*. Left & middle panels are boxplots showing

cdon/boc expression levels in 2 replicate samples of both cell types. Right panel shows

histogram of gene expression levels for all samples for all genes in the dataset aligned with

boxplots in left two panels. The location of genes that are markers for NCCs (*sox10*) or slow-

twitch muscle (*prdm1a*, *smyhc1*) are indicated by arrows. **(F&G)** HCR shows expression of

cdon/boc in trunk NCCs. **(F)** whole mount lateral view of the trunk of a 24 hpf embryo showing

expression of *cdon* (magenta), *boc* (green), and NCCs (grey) labeled by the tg(*sox10*:mRFP)

transgene. HCR puncta that overlap the *sox10* transgene were segmented out to visualize NCC

expression of *cdon* (F') or *boc* (F''). Segmented *cdon/boc* expression (*) are shown in yellow. (G)

Image from a 20uM cross section at the level of the yolk extension in the trunk shows NCCs

expressing *cdon/boc* transcripts. *cdon/boc* HCR puncta overlapping the tg(*sox10*:mRFP)

transgene (*) are shown in panels G' and G''. **(H-L)** Slow-twitch muscle cells express *cdon* and

boc. The tg(*prdm1a*:eGFP) transgene labeles adaxial cells at 2 somites **(H)**, and derivative slow-

twitch muscle cells at later stages **(I-L)**. *cdon/boc* transcripts do not overlap the adaxial cells at 2

somites **(H)**, but do overlap the slow-twitch muscle cells at 10 somites **(I-J)** and 24 hpf **(K-L)**.

Panel I shows 3D rendered of cross section from whole mount imaged embryos. HCR puncta

overlapping the tg(*prdm1a*:eGFP) transgene are shown in yellow in panels I' and I'' to highlight

slow-twitch muscle expression of *cdon/boc*. Panel J shows 3D rendered lateral view with HCR

puncta overlapping the *prdm1a* transgene (yellow). **(K)** At 24hpf slow-twitch muscle cells (grey)

continue to express *cdon/boc* transcripts. Panels K' and K'' show overlapping puncta. **(L)** 20uM

cross section of the trunk of a 24hpf embryo shows *cdon/boc* puncta overlapping the

tg(*prdm1a*:eGFP) transgene.

Figure 3. *cdon* and *boc* double mutants have NCC migration and slow-twitch muscle

differentiation phenotypes and a reduction in hedgehog signal reception. (A) Schematic cross

sections show NCC (red) and adaxial mesoderm (green) migration from approximately 10

somites to 20 somites. Adaxial mesoderm cells are specified adjacent to the notochord (nc) and

migrate laterally to differentiate into slow-twitch muscle fibers. Concurrent with adaxial cell

migration, NCCs migrate ventrally from dorsal neural tube (NT). **(B)** Left Panel shows 3D

rendering of a 10uM cross section from the trunk of a 15 somite embryo shows medially

positioned adaxial mesoderm (green) and ventrally migrating NCCs (red). NCCs are labeled with

the tg(*sox10*:mRFP) transgene, and adaxial mesoderm are labeled with the tg(*prdm1a*:eGFP)

transgene. Right panels show a lateral view of a 3D rendering of the trunk of a 24hpf embryo

shows slow-twitch muscle mesoderm (green) and migrating NCCs (red). Note the streams of

migrating NCCs and laterally position slow-twitch muscle. **(C-F)** Maximum projection images

show NCC streams (tg(*sox10*:mRFP)) and immunostaining for slow-twitch muscle specific

myosin (F59) for 24hpf wildtype embryos. Note loss of both streams and slow-twitch muscle

fibers in the double mutant. All images taken over the yolk extension in the trunk. **(G-J)**

Maximum projection images the tg(8xGli:mCherry) transgene in 24hpf embryos. Images taken

over the yolk extension. Note mCherry positive nuclei in the slow-twitch muscle mesoderm

(yellow arrows) and ventral neural tube (yellow arrowheads). 8xGli:mCherry signal

progressively decreases in *cdon*/*boc* mutants reflecting lower hedgehog signal reception.

Figure 4. *cdon* and *boc* sensitize zebrafish embryos to a reduction in hedgehog signal by

cyclopamine treatment. **(A)** Representative images show slow-twitch muscle fiber morphology

in 24hpf embryos treated with either 20uM cyclopamine or an equivalent volume of ethanol as

a control. Note *cdon* and *boc* mutants exhibit more severe phenotypes than their wildtype

siblings when treated with cyclopamine as evidenced by gaps in segments (arrows). Also note

complete loss of slow-twitch muscles in the double mutant treated with cyclopamine. **(B)**

Quantification of average number of slow-twitch muscle fibers per segment shows that *cdon*

and *boc* mutations are sensitizing embryos to cylclopamine treatment. Note that the number of

slow-twitch muscle fibers in single *cdon* and *boc* mutants are similar to wildtype siblings when

treated with ethanol, but significantly lower when treated with 20uM cyclopamine. Significance

values are from Tukey pairwise post-hoc tests: <0.05 = *, <0.001 = ***.

Figure 5. NCC directionality, but not migratory ability, is disrupted in the *cdon/boc* double

mutants. (A) Maximum projection images show migrating NCCs (tg(*sox10*:mRFP)) in the trunks

of representative wildtype, *cdon*, *boc*, and double mutant embryos. Migratory tracks from

individual cells are shown by colored lines. Note that NCCs form horizontal tracks at the body

midline in the double mutant. **(B)** Matched Roseplots (left panels) showing directional

movement and line plots (right panel) showing tracks for NCCs in each genotype. For both

panels, dorsal is up and anterior is to the left. Samples sizes reflect #tracks/#Embryos. Note that

ventral directionality of NCC movement is lower in the double mutant as best seen by the rose

plot. **(C)** NCC migration tracks (colored lines) are overlayed onto outlines of the body (grey

dashed lines) for representative embryos. Note the vertical aligned tracks in the wildtype, *cdon*,

and *boc* mutants that reflects ventral migration. Tracks form horizontal lines in the double

mutant reflecting NCC migratory movements along the anterior-posterior axis. Body outlines

were traced from the starting timepoint and do not reflect growth. Somite segments in the

double mutant are wider and dysmorphic. **(D-H)** Boxplots show quantification of NCC migration

for maximum velocity **(D)**, mean velocity **(E)**, track length **(F)**, track displacement **(G)**, and track

straightness **(H)**. Note that the double *cdon/boc* mutants are the only condidtion that is

significantly different from wildtype embryos in track straightness. Small grey points are

individual track values, large colored points are means for individuals, boxplots reflect

variability across tracks. **(I)** Schematic of track straightness which is calculated as track

displacement dived by track length.

Figure 6. *cdon* and *boc* are important for proper craniofacial development. (A) Representative

images of 7dpf larvae show craniofacial cartilages in *boc*[co25], *cdon*[co1019], and double mutants as

shown by alcian blue (cartilage, blue) and alizarin red (bone, red) stain. Note the medial

trabecular elements of the neurocranium are disrupted in *boc* and double mutants (arrows).

Also note a ventral projecting ceratohyal cartilage in the *boc* and double mutant (arrow head).

All phenotypes are more severe in the double mutant which also has hypomorphic craniofacial

cartilages and lacks the symplectic arm of the hyosymplectic cartilage. Scale bars are 20uM. ch

= ceratohyal, ep = ethmoid plate, hs = hyosymplectic, mc = Meckel's cartilage, op = opercle, pq

= palatoquadrate, pt = pterygoid process, sy = symplectic, tr = trabecula. **(B)** Representative

maximum projection images of 7dpf larvae stained with phalloidin (green) to label muscles.

Cartilages are labeled by the tg(*sox10*:mRFP) transgene that marks NCCs (magenta). Arrow

head points to ventral oriented ceratohyal in boc and double mutants. Note that craniofacial

muscles are normal across all genotypes suggesting that mis-orientation of the ceratohyal is

unlikely to be a secondary consequence of disruption to soft-tissue morphology. **(C)**

Quantification of ceratohyal angle shows that *boc*[co25] and double mutants have significantly

more ventral projecting ceratohyals. **(D)** Quantification of symplectic length showing that the

length of the symplectic is shorter in *cdon*/*boc* mutants. Significance in panels B&C is Tukey

post-hoc test relative to wildtype values. Sample sizes are wildtype = 6, cdon = 8, boc = 11,

Double = 6. **(E)** A cdon^{co1019}/cdon^{co1023};boc^{co25}/boc^{uml} embryo showing that different *cdon* and

boc alleles do not complement each other.

References

Allen, B.L., Song, J.Y., Izzi, L., Althaus, I.W., Kang, J.S., Charron, F., Krauss, R.S., McMahon, A.P.,
2011. Overlapping roles and collective requirement for the coreceptors GAS1, CDO, and BOC in
SHH pathway function. Dev Cell 20, 775-787.

Allen, B.L., Tenzen, T., McMahon, A.P., 2007. The Hedgehog-binding proteins Gas1 and Cdo
cooperate to positively regulate Shh signaling during mouse development. Genes Dev 21, 1244-
1257.

Banerjee, S., Gordon, L., Donn, T.M., Berti, C., Moens, C.B., Burden, S.J., Granato, M., 2011. A
novel role for MuSK and non-canonical Wnt signaling during segmental neural crest cell
migration. Development 138, 3287-3296.

Banerjee, S., Isaacman-Beck, J., Schneider, V.A., Granato, M., 2013. A novel role for Lh3
dependent ECM modifications during neural crest cell migration in zebrafish. PLoS One 8,
e54609.

Barresi, M.J.F., Stickney, H.L., Devoto, S.H., 2000. The zebrafish slow-muscle-omitted gene
product is required fro Hedgehog signal transduction and the development of slow muscle
identity. Development 127, 2189-2199.

Barriga, E.H., Maxwell, P.H., Reyes, A.E., Mayor, R., 2013. The hypoxia factor Hif-1alpha controls
neural crest chemotaxis and epithelial to mesenchymal transition. J Cell Biol 201, 759-776.

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models
Usinglme4. Journal of Statistical Software 67.

Bergeron, S.A., Tyurina, O.V., Miller, E., Bagas, A., Karlstrom, R.O., 2011. Brother of cdo
(umleitung) is cell-autonomously required for Hedgehog-mediated ventral CNS patterning in
the zebrafish. Development 138, 75-85.

Bilioni, A., Sanchez-Hernandez, D., Callejo, A., Gradilla, A.C., Ibanez, C., Mollica, E., Carmen
Rodriguez-Navas, M., Simon, E., Guerrero, I., 2013. Balancing Hedgehog, a retention and release
equilibrium given by Dally, Ihog, Boi and shifted/DmWif. Dev Biol 376, 198-212.

Camp, D., Currie, K., Labbé, A., Meyel, D.J.v., Charron, F.d.r., 2010. Ihog and Boi are essential for
Hedgehog signaling in Drosophila. Neural Development 5.

Cardozo, M.J., Sanchez-Arrones, L., Sandonis, A., Sanchez-Camacho, C., Gestri, G., Wilson, S.W.,
Guerrero, I., Bovolenta, P., 2014. Cdon acts as a Hedgehog decoy receptor during proximal-
distal patterning of the optic vesicle. Nat Commun 5, 4272.

Carmona-Fontaine, C., Matthews, H.K., Kuriyama, S., Moreno, M., Dunn, G.A., Parsons, M.,
Stern, C.D., Mayor, R., 2008. Contact inhibition of locomotion in vivo controls neural crest
directional migration. Nature 456, 957-961.

Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha,
A., Pierce, N.A., 2018. Third-generation in situ hybridization chain reaction: multiplexed,
quantitative, sensitive, versatile, robust. Development 145.

Clay, M.R., Halloran, M.C., 2011. Regulation of cell adhesions and motility during initiation of
neural crest migration. Curr Opin Neurobiol 21, 17-22.

Clay, M.R., Halloran, M.C., 2013. Rho activation is apically restricted by Arhgap1 in neural crest
cells and drives epithelial-to-mesenchymal transition. Development 140, 3198-3209.

Clay, M.R., Halloran, M.C., 2014. Cadherin 6 promotes neural crest cell detachment via F-actin
regulation and influences active Rho distribution during epithelial-to-mesenchymal transition.
Development 141, 2506-2515.

Connor, R.M., Allen, C.L., Devine, C.A., Claxton, C., Key, B., 2005. BOC, brother of CDO, is a
dorsoventral axon-guidance molecule in the embryonic vertebrate brain. J Comp Neurol 485,
32-42.

Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2011. ProtTest 3: fast selection of best-fit
models of protein evolution. Bioinformatics 27, 1164-1165.

Dehal, P., Boore, J.L., 2005. Two rounds of whole genome duplication in the ancestral
vertebrate. PLoS Biol 3, e314.

Devoto, S.H., Melancon, E., Eisen, J.S., Westerfield, M., 1996. Identification of separate slow
and fast muscle precursor cells in vivo, prior to somite formation. Development 122, 3371-
3380.

Diener, J., Sommer, L., 2021. Reemergence of neural crest stem cell-like states in melanoma
during disease progression and treatment. Stem Cells Transl Med 10, 522-533.

Eberhart, J.K., Swartz, M.E., Crump, J.G., Kimmel, C.B., 2006. Early Hedgehog signaling from
neural to oral epithelium organizes anterior craniofacial development. Development 133, 1069-
1077.

Echevarria-Andino, M.L., Allen, B.L., 2020. The hedgehog co-receptor BOC differentially
regulates SHH signaling during craniofacial development. Development 147.

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32, 1792-1797.

Elworthy, S., Hargrave, M., Knight, R., Mebus, K., Ingham, P.W., 2008. Expression of multiple
slow myosin heavy chain genes reveals a diversity of zebrafish slow twitch muscle fibres with
differing requirements for Hedgehog and Prdm1 activity. Development 135, 2115-2126.

Ferent, J., Giguere, F., Jolicoeur, C., Morin, S., Michaud, J.F., Makihara, S., Yam, P.T., Cayouette,
M., Charron, F., 2019. Boc Acts via Numb as a Shh-Dependent Endocytic Platform for Ptch1
Internalization and Shh-Mediated Axon Guidance. Neuron 102, 1157-1171 e1155.

Guillon, E., Bretaud, S., Ruggiero, F., 2016. Slow Muscle Precursors Lay Down a Collagen XV
Matrix Fingerprint to Guide Motor Axon Navigation. J Neurosci 36, 2663-2676.

Guindon, S., Gascuel, O., 2003. A simple, fast, and accurate algorithm to estimate large
phylogenies by maximum likelihood. Syst Biol 52, 696-704.

Hall, E.T., Dillard, M.E., Stewart, D.P., Zhang, Y., Wagner, B., Levine, R.M., Pruett-Miller, S.M.,
Sykes, A., Temirov, J., Cheney, R.E., Mori, M., Robinson, C.G., Ogden, S.K., 2021. Cytoneme
delivery of Sonic Hedgehog from ligand-producing cells requires Myosin 10 and a Dispatched-
BOC/CDON co-receptor complex. Elife 10.

Hong, M., Christ, A., Christa, A., Willnow, T., Krauss, R.S., 2020. Cdon Mutation and Fetal
Alcohol Converge on Nodal Signaling in a Gene-Environment Interaction Model of
Holoprosencephaly. BioRxiv.

Honjo, Y., Eisen, J.S., 2005. Slow muscle regulates the pattern of trunk neural crest migration in
zebrafish. Development 132, 4461-4470.

Izzi, L., Levesque, M., Morin, S., Laniel, D., Wilkes, B.C., Mille, F., Krauss, R.S., McMahon, A.P.,
Allen, B.L., Charron, F., 2011. Boc and Gas1 each form distinct Shh receptor complexes with
Ptch1 and are required for Shh-mediated cell proliferation. Dev Cell 20, 788-801.

Jeong, M.H., Kim, H.J., Pyun, J.H., Choi, K.S., Lee, D.I., Solhjoo, S., O'Rourke, B., Tomaselli, G.F.,
Jeong, D.S., Cho, H., Kang, J.S., 2017. Cdon deficiency causes cardiac remodeling through
hyperactivation of WNT/beta-catenin signaling. Proc Natl Acad Sci U S A 114, E1345-E1354.

Kang, J.-S., Mulieri, P.J., Hu, Y., Taliana, L., Krauss, R.S., 2002. BOC, an Ig superfamily member,
associates with CDO to positively regulate myogenic differentiation. EMBO J 21, 114-124.

Kang, J.S., Bae, G.U., Yi, M.J., Yang, Y.J., Oh, J.E., Takaesu, G., Zhou, Y.T., Low, B.C., Krauss, R.S.,
2008. A Cdo-Bnip-2-Cdc42 signaling pathway regulates p38alpha/beta MAPK activity and
myogenic differentiation. J Cell Biol 182, 497-507.

Kaufman, C.K., Mosimann, C., Fan, Z.P., Yang, S., Thomas, A.J., Ablain, J., Tan, J.L., Fogley, R.D.,
van Rooijen, E., Hagedorn, E.J., Ciarlo, C., White, R.M., Matos, D.A., Puller, A.C., Santoriello, C.,
Liao, E.C., Young, R.A., Zon, L.I., 2016. A zebrafish melanoma model reveals emergence of
neural crest identity during melanoma initiation. Science 351, aad2197.

Kearns, C.A., Walker, M., Ravanelli, A.M., Scott, K., Arzbecker, M.R., Appel, B., 2021. Zebrafish
spinal cord oligodendrocyte formation requires boc function. BioRxiv.

Kucenas, S., Snell, H., Appel, B., 2008. nkx2.2a promotes specification and differentiation of a
myelinating subset of oligodendrocyte lineage cells in zebrafish. Neuron glia biology 4, 71-81.

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in Linear
Mixed Effects Models. Journal of Statistical Software 82, 1-26.

Lencer, E., Prekeris, R., Artinger, K.B., 2021. Single-cell RNA analysis identifies pre-migratory
neural crest cells expressing markers of differentiated derivatives. Elife 10.

Li, H., Coghlan, A., Ruan, J., Coin, L.J., Heriche, J.K., Osmotherly, L., Li, R., Liu, T., Zhang, Z.,
Bolund, L., Wong, G.K., Zheng, W., Dehal, P., Wang, J., Durbin, R., 2006. TreeFam: a curated
database of phylogenetic trees of animal gene families. Nucleic Acids Res 34, D572-580.

Lu, M., Krauss, R.S., 2010. N-cadherin ligation, but not Sonic hedgehog binding, initiates Cdo-
dependent p38alpha/beta MAPK signaling in skeletal myoblasts. Proc Natl Acad Sci U S A 107,
4212-4217.

Mayor, R., Etienne-Manneville, S., 2016. The front and rear of collective cell migration. Nat Rev
Mol Cell Biol 17, 97-109.

Meijering, E., D'zyubachyk, O., Smal, I., 2012. Methods for Cell and Particle Tracking. Methods
in Enzymology 504, 183-200.

Mich, J.K., Payumo, A.Y., Rack, P.G., Chen, J.K., 2014. In vivo imaging of Hedgehog pathway
activation with a nuclear fluorescent reporter. PLoS One 9, e103661.

Mulieri, P.J., Kang, J.S., Sassoon, D.A., Krauss, R.S., 2002. Expression of the boc gene during
murine embryogenesis. Dev Dyn 223, 379-388.

MULIERI, P.J., OKADA, A., SASSOON, D.A., MCCONNELL, S.K., KRAUSS, A.R.S., 2000.
Developmental Expression Pattern of the cdo Gene. Developmental Dynamics 219, 40-49.

Nguyen-Chi, M.E., Bryson-Richardson, R., Sonntag, C., Hall, T.E., Gibson, A., Sztal, T., Chua, W.,
Schilling, T.F., Currie, P.D., 2012. Morphogenesis and cell fate determination within the adaxial
cell equivalence group of the zebrafish myotome. PLoS Genet 8, e1003014.

Okada, A., Charron, F., Morin, S., Shin, D.S., Wong, K., Fabre, P.J., Tessier-Lavigne, M.,
McConnell, S.K., 2006. Boc is a receptor for sonic hedgehog in the guidance of commissural
axons. Nature 444, 369-373.

Okumura, F., Oki, N., Fujiki, Y., Ikuta, R., Osaki, K., Hamada, S., Nakatsukasa, K., Hisamoto, N.,
Hara, T., Kamura, T., 2021. ZSWIM8 is a myogenic protein that partly prevents C2C12
differentiation. Sci Rep 11, 20880.

Olesnicky Killian, E.C., Birkholz, D.A., Artinger, K.B., 2009. A role for chemokine signaling in
neural crest cell migration and craniofacial development. Dev Biol 333, 161-172.

Parslow, A., Cardona, A., Bryson-Richardson, R., 2014. Sample drift correctino following 3D
confocal time-lapse imaging. J. Vis. Exp. 12.

Perris, R., Perissinott, D., 2000. Role of the extracellular matrix during neural crest cell
migration. Mechanisms of Development 95, 3-21.

Powder, K.E., Cousin, H., McLinden, G.P., Craig Albertson, R., 2014. A nonsynonymous mutation
in the transcriptional regulator lbh is associated with cichlid craniofacial adaptation and neural
crest cell development. Mol Biol Evol 31, 3113-3124.

Powell, D.R., Williams, J.S., Hernandez-Lagunas, L., Salcedo, E., O'Brien, J.H., Artinger, K.B.,
2015. Cdon promotes neural crest migration by regulating N-cadherin localization. Dev Biol 407,
289-299.

Roy, S., Wolff, C., Ingham, P.W., 2001. The u-boot mutation identifies a Hedgehog-regulated
myogenic switch for fiber-type diversification in the zebrafish embryo. Genes Dev 15, 1563-
1576.

Ruan, J., Li, H., Chen, Z., Coghlan, A., Coin, L.J., Guo, Y., Heriche, J.K., Hu, Y., Kristiansen, K., Li, R.,
Liu, T., Moses, A., Qin, J., Vang, S., Vilella, A.J., Ureta-Vidal, A., Bolund, L., Wang, J., Durbin, R.,
2008. TreeFam: 2008 Update. Nucleic Acids Res 36, D735-740.

Sanchez-Arrones, L., Cardozo, M., Nieto-Lopez, F., Bovolenta, P., 2012. Cdon and Boc: Two
transmembrane proteins implicated in cell-cell communication. Int J Biochem Cell Biol 44, 698-
702.

Scarpa, E., Szabo, A., Bibonne, A., Theveneau, E., Parsons, M., Mayor, R., 2015. Cadherin Switch
during EMT in Neural Crest Cells Leads to Contact Inhibition of Locomotion via Repolarization of
Forces. Dev Cell 34, 421-434.

Schwend, T., Ahlgren, S.C., 2009. Zebrafish con/disp1 reveals multiple spatiotemporal
requirements for Hedgehog-signaling in craniofacial development. BMC Dev Biol 9, 59.

Shellard, A., Mayor, R., 2021. Collective durotaxis along a self-generated stiffness gradient in
vivo. Nature 600, 690-694.

Shellard, A., Szabo, A., Trepat, X., Mayor, R., 2018. Supracellular contraction at the rear of
neural crest cell groups drives collective chemotaxis. Science 362, 339-343.

Sofroniew, N., Lambert, T., Evans, K., Nunez-Iglesias, J., Bokota, G., Peña-Castellanos, G.,
Winston, P., Yamauchi, K., Bussonnier, M., Pop, D.D., Liu, Z., ACS, Pam, alisterburt, Buckley, G.,
Sweet, A., Gaifas, L., Rodríguez-Guerra, J., Migas, L., Hilsenstein, V., Bragantini, J., Lee, G.R.,
Hector, Freeman, J., Boone, P., Lowe, A.R., Gohlke, C., Royer, L., PIERRÉ, A., Har-Gil, H., 2019.
napari: a multi-dimensional image viewer for python.

Song, J.Y., Holtz, A.M., Pinskey, J.M., Allen, B.L., 2015. Distinct structural requirements for
CDON and BOC in the promotion of Hedgehog signaling. Dev Biol 402, 239-252.

Swartz, M.E., Nguyen, V., McCarthy, N.Q., Eberhart, J.K., 2012. Hh signaling regulates patterning
and morphogenesis of the pharyngeal arch-derived skeleton. Dev Biol 369, 65-75.

Szabo, A., Melchionda, M., Nastasi, G., Woods, M.L., Campo, S., Perris, R., Mayor, R., 2016. In
vivo confinement promotes collective migration of neural crest cells. J Cell Biol 213, 543-555.

Tenzen, T., Allen, B.L., Cole, F., Kang, J.S., Krauss, R.S., McMahon, A.P., 2006. The cell surface
membrane proteins Cdo and Boc are components and targets of the Hedgehog signaling
pathway and feedback network in mice. Dev Cell 10, 647-656.

Theveneau, E., Mayor, R., 2012. Neural crest delamination and migration: from epithelium-to-
mesenchyme transition to collective cell migration. Dev Biol 366, 34-54.

Tobin, J.L., Di Franco, M., Eichers, E., May-Simera, H., Garcia, M., Yan, J., Quinlan, R., Justice,
M.J., Hennekam, R.C., Briscoe, J., Tada, M., Mayor, R., Burns, A.J., Lupski, J.R., Hammond, P.,
Beales, P.L., 2008. Inhibition of neural crest migration underlies craniofacial dysmorphology and
Hirschsprung's disease in Bardet-Biedl syndrome. Proc Natl Acad Sci U S A 105, 6714-6719.

Vega-Lopez, G.A., Cerrizuela, S., Aybar, M.J., 2017. Trunk neural crest cells: formation, migration
and beyond. Int J Dev Biol 61, 5-15.

Vega-Lopez, G.A., Cerrizuela, S., Tribulo, C., Aybar, M.J., 2018. Neurocristopathies: New insights
150 years after the neural crest discovery. Dev Biol 444 Suppl 1, S110-S143.

Wada, N., Javidan, Y., Nelson, S., Carney, T.J., Kelsh, R.N., Schilling, T.F., 2005. Hedgehog
signaling is required for cranial neural crest morphogenesis and chondrogenesis at the midline
in the zebrafish skull. Development 132, 3977-3988.

Williams, J.S., Hsu, J.Y., Rossi, C.C., Artinger, K.B., 2018. Requirement of zebrafish pcdh10a and
pcdh10b in melanocyte precursor migration. Dev Biol 444 Suppl 1, S274-S286.

Wolff, C., Roy, S., Ingham, P.W., 2003. Multiple Muscle Cell Identities Induced by Distinct Levels
and Timing of Hedgehog Activity in the Zebrafish Embryo. Current Biology 13, 1169-1181.

Yao, S., Lum, L., Beachy, P., 2006. The ihog cell-surface proteins bind Hedgehog and mediate
pathway activation. Cell 125, 343-357.

Yin, J., Lee, R., Ono, Y., Ingham, P.W., Saunders, T.E., 2018. Spatiotemporal Coordination of FGF
and Shh Signaling Underlies the Specification of Myoblasts in the Zebrafish Embryo. Dev Cell 46,
735-750 e734.

Zhang, W., Hong, M., Bae, G.U., Kang, J.S., Krauss, R.S., 2011. Boc modifies the
holoprosencephaly spectrum of Cdo mutant mice. Dis Model Mech 4, 368-380.

Zhang, W., Kang, J.S., Cole, F., Yi, M.J., Krauss, R.S., 2006. Cdo functions at multiple points in the
Sonic Hedgehog pathway, and Cdo-deficient mice accurately model human holoprosencephaly.
Dev Cell 10, 657-665.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6