Epigenomic mapping identifies a super-enhancer repertoire that regulates cell identity in bladder cancers through distinct transcription factor networks

Authors

Hélène Neyret-Kahn*[1,2], Jacqueline Fontugne*[1,2,3], Xiang Yu Meng[1,2,5], Clarice S. Groeneveld[1,2,6], Luc Cabel[1,2], Tao Ye[7], Elodie Guyon[1,4], Clémentine Krucker[1,2], Florent Dufour[1,2], Elodie Chapeaublanc[1,2], Audrey Rapinat[8], Daniel Jeffery[9], Yann Neuzillet[10], Thierry Lebret[10], David Gentien[8], Irwin Davidson[7], Yves Allory[1,2,3], Isabelle Bernard-Pierrot [#,1,2], François Radvanyi [#,1,2].

* These authors contributed equally to this work

# These authors co-supervised this work

Correspondance to helene.neyret-kahn@curie.fr and francois.radvanyi@curie.fr

Affiliations

[1] Molecular Oncology, PSL Research University, CNRS, UMR 144, Institut Curie, Equipe Labellisée Ligue Nationale Contre le Cancer, Paris, France

[2] Sorbonne Universités, UPMC Université Paris 06, CNRS, UMR144, 75005, Paris, France.

[3] Department of Pathology, Institut Curie, Saint-Cloud, France.

[4] Department of Pathology, Institut Curie, Paris, France.

[5] Department of Urology, Zhongnan Hospital of Wuhan University

[6] Université de Paris, Centre de Recherche des Cordeliers, Paris, France

[7] Department of Functional Genomics and Cancer, Institut de Genétique et de Biologie Moleculaire et Cellulaire, CNRS/INSERM/UDS, 67404 Illkirch Cedex, France.

[8] Department of Translational Research, Genomics Platform, Institut Curie, PSL Research University, Paris, France

[9] Urology Medico-Scientific Program, Department of Translational Research, Institut Curie, PSL Research University, Paris, France

[10] Urology Department, Hôpital Foch, Suresnes, France

Abstract

Background:

Muscle-invasive bladder cancer is a common aggressive disease with unmet clinical needs.

Recent work established a set of consensus bladder cancer transcriptomic subtypes that

distinguishes the cell identity of bladder cancers for improved diagnosis and treatment.

However, how these distinct subtypes are regulated remains unclear. Given the link between

super-enhancers and the regulation of cell identity, we hypothesized that epigenetic activation

of distinct super-enhancers could drive the transcriptional programs of the various bladder

cancer subtypes.

Results:

Through integrated RNA sequencing and epigenomic profiling of histone marks (H3K27ac,

H3K27me3, H3K9me3) in a diverse panel of 15 primary bladder tumours, seven bladder

cancer cell lines, and two primary cultures from normal human urothelia, we established the

first integrated epigenetic map of bladder cancer and demonstrate the link between bladder

cancer subtype and epigenetic control. Through H3K27ac analysis, we identify the repertoire

of activated super-enhancers in bladder cancer that distinguish molecular subtypes. Building

on these findings, we reveal the super-enhancer-regulated networks of candidate master

transcription factors for Luminal and Basal bladder cancer subgroups. We find that FOXA1, a

key pioneer factor in Luminal bladder cancers identified in our Luminal transcription factor

network, binds subgroup-specific bladder super-enhancers and correlates with their activation.

Furthermore, CRISPR-Cas9 inactivating mutation of *FOXA1* triggers a shift from Luminal to

Basal cell identity. This shift is accompanied by an overexpression of ZBED2, one of the

newly identified transcriptional regulators in the Basal-specific transcription factor network.

Finally, we show that both FOXA1 and ZBED2 play concordant roles in preventing

inflammatory response in bladder cancer cells through STAT2 inhibition and promote cancer

cell survival.

Conclusions:

Overall, our study provides new data for understanding epigenetic regulation of muscle-

invasive bladder cancer and identifies a coregulated network of super-enhancers and

associated transcription factors as new potential targets for the treatment of this aggressive

disease.

Keywords Three to ten keywords representing the main content of the article.

Epigenetic, Super-enhancer, Network, ZBED2, FOXA1, Cellular identity, Inflammation,
Bladder Cancer

Abbreviation

SE, Super-enhancer

TF, Transcription Factor

NMIBC, non-muscle-invasive bladder carcinoma

MIBC, muscle-invasive bladder carcinoma

BLCA, Bladder Cancer Carcinoma

Introduction

Bladder cancer is the tenth most common cancer worldwide, accounting for nearly 200 thousand cancer-related deaths globally in 2018 [1]. Urothelial carcinoma is classified as non-muscle-invasive bladder cancer (NMIBC, carcinoma *in situ*, pTa or pT1 stages) or the aggressive muscle-invasive bladder cancer (MIBC, stages pT2 to pT4), depending on the level of invasion into the bladder wall [2]. Molecular classifications of bladder carcinomas have been established using gene expression profiling studies [3–8]. A recent consensus classification of MIBC presents six subtypes, from which tumours can be coarsely divided into Luminal or Basal subgroups [3]. Luminal tumours, accounting for about 50% of MIBCs, present high expression of urothelial differentiation markers (GATA3, FOXA1, KRT20, uroplakins) and are enriched in activating mutations of *FGFR3*. Ba/Sq tumours are particularly aggressive and account for ~35% of MIBCs [3]. They are characterized by the overexpression of markers of the basal layer of the urothelium (including KRT5, KRT6) and the under-expression of markers of luminal differentiation. In NMIBC, the recent UROMOL studies group NMIBC into 4 classes including Class 1 associated with luminal differentiation and good prognosis, and a Class 2a comprising high risk tumours [7,8]. One hypothesis to explain the establishment of the different subtypes, and their potential plasticity, is that each subtype harbours a regulatory network in which various upstream genomic and epigenomic alterations lead to the activation of a core set of master transcription factors (TFs) that then determine a stable transcriptomic downstream program. While transcriptional regulators of urothelial differentiation, such as FOXA1, GATA3 and PPARG, have been established as key regulators of the Luminal phenotypes, the essential transcription regulators driving the Basal subgroup have not been elucidated [9–13].

Recent studies have demonstrated that altered enhancer activity drives changes in cell identity and oncogenic transformation, notably through large clusters of highly active enhancers called super-enhancers (SEs) [14–16]. Indeed, targeting SE-driven oncogenesis has become a novel therapeutic approach with the advent of BRD4 inhibitors, which inhibit SE activation [17]. By regulating the expression of a small number of master TFs, SEs can orchestrate cell- or cancer-specific transcriptional programs. The gold standard for identifying SEs is histone mark profiling [16]. The ENCODE roadmap, that profiled histone marks in normal and cancer cell lines, has become a valuable source of information to uncover chromatin organisation,

alteration, and subsequent regulation of master regulators [18]. However, ENCODE did not

include bladder models. Two recent studies began to fill this gap, profiling particular histone

marks in bladder cancer samples and cell lines [19,20]. However, to comprehensively

characterize bladder cancer epigenetic alterations, super-enhancer rewiring linked to

molecular subtypes and their associated master TFs, there is a need to integrate multiple

chromatin marks with transcriptomic data in normal bladder cells, cell lines and tumour

samples.

Here, we integrated transcriptomics and epigenomic profiling of multiple histone marks in

human bladder tumours, bladder cancer cell lines, and primary cultures from normal urothelia

to produce a comprehensive bladder cancer epigenetic map. With this map, we demonstrate

the link between molecular subtype and the underlying epigenetic landscape. Through

H3K27ac analysis, we establish a repertoire of SEs that are specific to distinct subgroups,

highlighting SE-associated genes with subgroup-specific clinical relevance. From there, we

identify the core SE-regulated networks of master TFs that distinguish luminal and basal

subgroups, including known and new candidate master TFs. Finally, through functional

knock-down and knock-out experiments, we reveal that one of these master TFs (FOXA1) is a

key factor in subtype determination antagonized by ZBED2, and that both FOXA1 and

ZBED2 present the ability to dampen inflammatory response. Overall, this work provides new

data characterizing epigenetic regulation in bladder cancer. We reveal important genes that

can be essentials for maintenance of bladder cancer cell identity and present potential new

targets to treat aggressive bladder cancers.

Results

Integrated bladder cancer chromatin landscape

To elucidate the contribution of chromatin landscape in bladder cancer biology, we generated ChIP-seq for active (H3K27ac) and repressive histone marks (H3K27me3, H3K9me3) in 24 bladder samples (Fig.1). In order to distinguish features of the non-cancerous stromal cells and of normal urothelial cells, we used not only human primary tumours (n=15) from the CIT (*Carte d'Identité des Tumeurs*) cohort [21,22], but also cellular models (7 bladder cancer cell lines) and patient-derived Normal Human Urothelium in proliferation (NHU, n=2). Of note, tumors were macrodissected to enrich for bladder cancer content (Fig. 1, Fig. S1A). Of the 15 primary tumours, we included 13 MIBCs and 2 NMIBCs to assess the stage-dependence of our results (Fig. 1A, Fig. S1A, and Table S1). With the aim of identifying subtype-specific epigenetic alterations/characteristics, we coupled our ChIP-seq with RNA-seq from the same extraction and classified them according to the current consensus subtypes [3]. Of the 13 MIBCs, 2 classified as stroma-rich, 4 classified as basal/squamous (Ba/Sq) and 7 as luminal, including 1 luminal papillary (LumP), 3 luminal unstable (LumU) and 2 luminal non-specified (LumNS). Of the 7 cell lines, 3 were classified Ba/Sq, 3 LumP and one could not be classified (Table S1)[23]. Using the recent UROMOL classifier, the two NMIBC samples classified as class 3 [7,8]. Further analysis using subtype deconvolution (WISP, [24]), and previously described regulatory signatures [3,7], revealed that one of the tumours originally classified as Ba/Sq (T391) was composed of a mixed population of LumP and Ba/Sq cells (Fig. S1B, S1C). Knowing it's important intra-tumoral heterogeneity, tumour T391 was excluded from differential analyses between subgroups. Peak calling using MACS showed that ChIP-seq for H3K27ac gave the most homogeneous and highest number of peaks across the 24 samples (Fig. S1D).

We integrated our multi-factorial ChIP-seq profiles using ChromHMM [25], reporting the first integrated epigenetic map in bladder cancer in both primary tumour samples and cell lines (Fig. 2A). Six chromatin states (E1–E6) were assigned according to histone mark enrichments, as previously described (ENCODE, Roadmap project [18]), where H3K27ac-enriched regions correspond to active promoters and enhancers (E2), H3K27me3 and H3K9me3-enriched states associate with repression (E4) or heterochromatin (E6), and regions enriched in both active and repressive marks define bivalent enhancers or promoters (E3). Regions without any marks or only weak H3K9me3 enrichment were designated as quiescent/no marks (E1) or quiescent/weakly repressed (E5), respectively (Fig. 2A, Fig. S2A,

B). Analysis of associated RNA-seq data confirmed that gene expression correlates with the

expected chromatin states (Fig. 2B). Briefly, genes with transcription start sites (TSSs) in E2

states (active enhancers / promoters) have the highest expression levels, followed by those in

E3 states (bivalent enhancers / promoters). Minimal expression was noted for genes with

TSSs in the remaining states.

Chromatin states classify bladder cancers by transcriptomic subgroups

Next, we sought to classify our samples based on chromatin states for comparison with

molecular subtypes. To do this we first performed an unsupervised analysis to select the most

distinguishing features from the chromatin profiles (see methods, Fig. S2C) and plotted all

samples by multiple correspondence analysis on the most varying regions (MCA, Fig. S2D).

Similar to Principal Component Analysis (PCA), but adapted for categorical data, this method

of dimensionality reduction separates samples in 2D space by proximity according to the

primary (Dim 1) and secondary (Dim 2) dimensions. Thus, greater differences in chromatin

profiles are represented by greater distances in the 2D plot. Dim 1 distinguished primary

tumour samples from cell lines, which could be indicative of chromatin changes associated

with cell culture or stroma content. Interestingly, Dim 2 distinguished Non-basal from Basal

subgroups. To confirm this distinction of molecular subtypes, we re-assessed the data with an

alternative dimensionality reduction method (MDS), coupled to a batch effect-like correction

(MNN), which eliminated most of the cell line vs primary tumour differences while

maintaining and strengthening the distinction between Non-Basal and Basal subgroups along

Dim 2 (Fig. 2C). Therefore, we identified two clusters derived from differences in chromatin

state that are associated with molecular subtypes; a "basal cluster" containing all Ba/Sq

samples (except the mixed T391), and a "non-basal cluster" including all luminal, stroma-rich

and NMIBC samples (Fig. 2C). Interestingly, NHU cells were located at the border between

the two groups (Fig. 2C). To explore the biological pathways associated with the chromatin

profiles that could distinguish Luminal from Basal bladder cancers, we ranked genes based on

the MCA outputs for Dim 2 and performed Gene Set Enrichment Analysis (GSEA [26,27],

Fig. 2D). As expected, for the basal cluster, we found that active chromatin (E2) was strongly

enriched at genes involved in decreased Luminal differentiation, while repressive chromatin

(E4) was strongly depleted for these genes (Fig. 2D). Interestingly, genes involved in

increased tumour aggressiveness, stemness, extracellular matrix, epithelial-mesenchymal

transition and invasion were also enriched in active chromatin and depleted for repressive

chromatin in the basal cluster (Fig. 2D). Taking an alternative approach, we derived Basal and

Luminal gene signatures from an independent publicly available scRNA-seq dataset
(GSM4307111) and compared these genes with the chromatin states associated with the basal
and non-basal clusters identified in Fig. 2C (Fig. 2E, see methods). Luminal signature genes
were enriched in active state in the non-basal chromatin cluster while Basal signature genes
were enriched in active state in the basal chromatin cluster, suggesting epigenetic regulation
of signature genes involved in urothelial differentiation. We further illustrate this relationship
with two well-described markers of bladder cancer subtypes: *FOXA1* and *KRT6A* (Fig. 2F).
FOXA1 expression is higher in Luminal than Basal tumours [10,13,28,29]. In agreement, our
results show that *FOXA1* was marked with active (E2) chromatin in Luminals (including
LumP, LumNS and LumU), NMIBC samples, and even NHU cells, but harboured repressive
chromatin in Ba/Sq tumour samples (Fig. 2F). On the other hand, *KRT6A*, commonly
expressed in Basal tumours [3]*,* had active chromatin marks in seven out of eight Ba/Sq
samples, as well as NHU, but not in any Luminal sample. Taken together, these results
demonstrate the importance of histone marks in the regulation of gene expression driving cell
identity in bladder cancer.
218

Identification of the bladder super-enhancer repertoire and subtype specificities
To determine if chromatin profiles identify SEs that could control bladder cancer subtype, we
determined and annotated typical enhancer and SE regions in our samples using the ROSE
algorithm (Table S2), which calls enhancers and particularly super enhancers according to
H3K27ac signal [30]. Despite signal correction using input, the number of called SEs was
notably lower in samples with gene amplification, owing to very high H3K27ac signal in
amplified regions, thus creating a bias in the ranked enhancer plot (Fig. S3A). To correct for
the copy-number bias, we set a threshold and defined the top 1000 enhancer regions in each
sample as SEs for all downstream analyses, approximately representing the mean number of
SEs per sample (mean = 956 SEs, Table S2).
To gain insight into subtype-specific enhancer alterations and assess sample similarity based
on these SE profiles, we determined the global repertoire of SEs in bladder by extracting a
consensus set of 2887 SEs present in at least 2 of our 24 samples (Table S3). Using PCA of
H3K27ac signal corresponding only to the consensus SE regions, we again found that samples
were grouped according to molecular subgroup, separating Basals from Luminals, NMIBC
segregating with differentiated tumors (Fig. 3A). This reveals that the variability in SE
profiles reflects the differences in Basal and Luminal transcriptional programs. We also
performed PCA using H3K27ac, H3K27me3 and H3K9me3 independently for peaks located

inside the SE consensus regions for the tumour samples alone (Fig. S3B). Interestingly, all

three profiles separated the Ba/Sq tumors from the other samples, indicating that all three

histone marks are likely linked to SE regulation of bladder cancer molecular subtypes.

We sought to further explore the functional pathway enrichment of SE-regulated genes. We

first assigned SEs to their closest and most transcriptionally correlated genes (see methods).

For example, several SEs mapped close to *KLF5*, whose expression was correlated with SE

activity mostly in Luminal samples, as previously reported [31] (Fig. 3B). In contrast, *KLF7*

was regulated by a large SE mostly active in Basal samples (Fig. 3B). We then performed

pairwise differential analyses between subtypes (Fig. 3C, S3D). The comparison between

Basal and Luminal samples identified 369 subgroup-specific SEs (Fig. 3C, Table S3). By

comparing the differential SEs to RNA-seq differential expression analysis, we confirmed that

luminal-gained SEs showed significantly higher expression levels in Luminal samples relative

to Basal samples and vice-versa for basal-gained SEs (Fig. 3D). We then validated the

subgroup-specific SE-associated genes identified in our samples in a larger dataset, leveraging

the gene expression profiles of the TCGA-BLCA MIBC cohort across molecular subtypes

(n=406) [32,33]. Hierarchical clustering of TCGA samples using the genes associated to the

most differentially regulated SEs in our consensus repertoire recovered the molecular

classification (Fig. S3C, Table S3). Strikingly, differential analysis between Basal and

Luminal SEs revealed that Luminal-specific SEs were attributed to known transcriptional

drivers of the luminal phenotype, namely *GATA3*, *PPARg*, *FOXA1* [13,21]. Luminal-gained

SEs were associated with "Signalling by TGF-beta family members", notably due to SEs

annotated close to negative regulators of TGF-beta signalling such as the E3 Ubiquitin ligase

SMURF1 or *SMAD6 (*Fig. 3C, E, Table S4). In contrast, SE regions significantly bound at

higher levels in the Basal tumours were associated with genes known to contribute to Basal

cancer biology such as *EGFR,* but also less characterized genes with regards to bladder cancer

biology, such as genes related to inflammation and FOXO signalling (*IL7R*, *FBX032*),

signalling by Interleukin or signalling by MET, the activation of which is often correlated

with BLCA progression [34] (Fig. 3E). We also identified genes encoding membrane

receptors (*IL7R*, *OSMR, EGFR*) and transcriptional regulators (*BNC2*, *HMGA2*, *KLF7*,

NR3C1) as enriched in Basal tumours (Fig. 3C). Taking advantage of the NHU samples in our

cohort, we extracted differential SEs in three comparisons (Ba/Sq *vs* NHU *vs* Luminals, Fig.

S3D, Fig. 3F). This analysis validated the identification of genes that could be specific to

cancer biology, such as *IL7R, OSMR, JUN, NR3C1* in Ba/Sq subtype, *NPAS2*, *FOXQ1*,

GRHL3 in Luminal samples and even *FGFR3* in both Basals and Luminals vs NHU (Fig. 3F).

Overall, we established a first SE repertoire for bladder cancer, highlighting subgroup-

specific, cancer-specific SE activation coupled with gene expression.

275   Super-enhancers regulate a network of candidate master transcription factors for

276   bladder cancer subgroups

SEs often regulate the expression of master TFs, forming autoregulatory loops and correlated

networks [35,36]. Having established the SE landscape in bladder cancer, we next sought to

determine which master regulators control the subtype-specific transcriptional programs. To

this end, we overlaid the genomic coordinates of subgroup-specific peaks inside SEs with

publicly available ChIP-seq datasets [37,38]. Our analysis revealed that Luminal-specific SEs

were significantly enriched in several TF binding sites (Fig. 4A, Table S5), including known

regulators of Luminal subtypes FOXA1, GATA3, and ESR1 [3,13,32]. Basal-specific SEs

were enriched in binding sites of a different set of regulators, including components of the

AP-1 complex (FOSL1, FOSL2, JUND, JUNB), as well as SMAD2/3, NFkB, and STAT3.

Further DNA motif enrichment analysis comparing Basal differential peaks inside subgroup-

specific SEs over Luminal ones, again identified AP-1 as a potential regulator of Basal SEs,

as well as FOXA1, FOXA1:AR, GATA, and GRHL1/2/3 for Luminal SEs (Fig. 4B, Table S5,

Homer [39]).

However, motif binding and ChIP-seq data are not available for all known TFs. To overcome

this issue, we designed a method to identify subgroup-associated TFs and their co-regulated

networks based on our differential SEs and the large transcriptomic cohort from the TCGA

[32] (Fig. 4C). We selected TFs that were regulated by differential SEs (Basal vs Luminal),

according to annotations from Lambert *et al*. [40], and that were differentially expressed in

TCGA Ba/Sq vs Luminal subgroups. Then, we used ARACNe [41] and VIPER [42] analysis

to identify and evaluate the regulons (group of genes regulated in response to one

transcription regulator) of the 75 resulting TFs. Hierarchical clustering of the resulting TF

regulons clustered scores into three groups, which were respectively associated with Luminal,

Ba/Sq or Stroma-Rich subtypes (Fig. 4D). Since master TFs form interconnected networks

with highly correlated levels of expression, we selected only TFs whose expression was

correlated with that of at least one of the other TF in TCGA-BLCA data (Pearson correlation

coefficient ≥0.5, n=55), and built the top correlated network based on subtype-specific SE-

associated TFs (Fig. 4F) (see methods). This strategy identified a large module of Luminal

TFs, including known Luminal-associated TFs (e.g., FOXA1/GATA3/PPARG), as well as

TFs with yet unexplored roles in Luminal bladder cancer biology (e.g., HES1, FOXQ1,
ZBTB7C, MECOM, GRHL2/3 and TBX3). Unlike for Luminals, few TFs have been
characterized as key regulators of Basal tumours. Our analysis revealed a network of TFs
whose activity could be essential to Basal bladder cancer biology, including HMGA2, KLF7,
NR3C1 and ZBED2. Notably, ZBED2 has recently been associated with basal identity in
keratinocytes [43] and regulation of inflammation in pancreatic cancer [44].

Combining analyses of tumour and cell lines SEs should avoid the identification of master
TFs expressed by the stroma. In fact, we found that TFs associated with the Luminal network
showed strong expression correlation in TCGA-BLCA and in CCLE bladder cell cohorts [45]
(Fig. S4A, B) while expression correlations of Stroma-Rich or Basal-associated TFs (e.g.,
ZEB1, SPI1) in the TCGA were lower for urothelial cell lines in the CCLE [46] (Fig.
S4A,B). This indicates that expression of those TFs could be dependent on growing
conditions and/or interactions with the tumour microenvironment.

To validate our Luminal- and Basal-specific TF networks, we analysed public single-cell
RNA-seq data of a tumour presenting both a Luminal and a Basal cell population
(GSM4307111, Fig. S4C). The Luminal-associated TFs FOXA1, GATA3 and PPARG were
mostly expressed in the Luminal cell cluster, whereas ZBED2, HMGA2, and KLF7, newly
identified as part of the Basal TF network, were mostly expressed in the Basal cell clusters
(Fig. 4G), validating our subgroup-specific networks. Together, these analyses identified a
targeted subset of interconnected candidate master TFs that could represent key regulators of
bladder cancer subgroup identity.

FOXA1 binds subgroup-specific bladder super-enhancers and correlates with their
activation

We identified FOXA1 as one of our candidate master TFs for the Luminal bladder cancer
subgroup. FOXA1 is a known pioneer factor with a demonstrated impact on Luminal bladder
cancer biology [29,32,47], though the mechanism for how FOXA1 regulates cell identity is
unknown. To better assess the role of FOXA1 in the regulation of bladder cancer SEs, we
mapped FOXA1, CTCF (Insulator/enhancers) and H3K4me3 (Promoter) binding by ChIP-seq
in two bladder cancer cell lines: SD48 (LumP) and 5637 (Ba/Sq). FOXA1 binding was mostly
found outside promoters (Fig. S5A), with 61,083 FOXA1 peaks detected in SD48 cells and
39,445 in 5637, an expected variation as FOXA1 was more abundant in Luminal cells.
Despite such differences, we identified three classes of FOXA1 peaks: SD48-specific peaks,

peaks overlapping in the two cell lines, and 5637-specific peaks (Fig. 5A, Fig. S5B), which
suggests that FOXA1 has specific targets in each cell line and subtype. Interestingly, when
analysing TF binding sites from publicly available ChIP-seq data, SD48-specific FOXA1
peaks were highly enriched not only for FOXA1 binding sites, but also GATA3 binding sites
(Fig. 5B), which could indicate a functional partnership between FOXA1 and GATA3 for
regulation of the Luminal program, as suggested by Warrick et al. and described in breast
cancer [13,48]. Surprisingly, 5637-specific FOXA1 peaks were mostly enriched at AP-1
binding sites and not FOXA1 sites (Fig. 5B, Fig. S5B). Both enrichments were confirmed by
Homer motif analysis of SD48-specific peaks versus 5637-specific peaks or vice and versa
(Fig. S5C and D, [39]). Ontology comparison of genes associated with the three classes of
FOXA1 peaks showed that 5637-specific peaks were enriched in terms associated with Ba/Sq
super-enhancers (e.g. Signalling by Tyrosine Kinase, Signalling by MET, Signalling by
Interleukin), indicating that FOXA1 might be involved in the regulation of both Luminal and
Basal bladder cancer subtypes (Fig. 5C). Indeed, FOXA1 binding in the two cell lines
overlapped with most (87%) of the total repertoire of bladder SEs (Fig. 5D) and correlated
strongly with H3K27ac levels at these loci (Fig. 5E, F), in line with a role for its regulation of
these SEs. Notably, FOXA1 bound at SEs associated with genes involved in regulating
urothelial differentiation and strongly correlated with increased H3K27ac at these loci. This
could clearly be observed for the Luminal-specific SE associated with *GATA3* in the Luminal
SD48 cells and in both cell lines for the non-specific *PPARG* SE. But we also found FOXA1
binding associated with high H3K27ac at certain Basal-specific SEs in the Basal 5637 cells,
such as *TGFB2* (Fig. 5G). This implies that FOXA1, even if expressed at a low level as in
Ba/Sq cells, could play an important role in BLCA biology, through enhancer/SE regulation.
In summary, FOXA1 may regulate bladder cell identity through binding of subgroup-specific
bladder SEs with partners such as GATA3 in Luminal cells and AP-1 in Basal cells.
363

FOXA1 regulates inflammation and cellular identity
To better understand FOXA1 function, we performed short-term (<72h) knock-down in both
Luminal and Basal models. Knock-down of FOXA1 by siRNA decreased clonogenicity and
proliferation of both Luminal and Basal cells (Fig. S6A, B) and it reduced cell viability in
both RT112 (LumP subtype) and SCaBER (Ba/Sq) cells, with a stronger impact in RT112
(Fig. 6A). Furthermore, FOXA1 knock-down in RT112 and SCaBER cell lines dramatically
altered gene expression (Fig. 6B, Table S6). The downregulated genes were related to cell

cycle and checkpoint pathways, consistent with the reductions in viability and proliferation

upon FOXA1 knock-down. Surprisingly, the upregulated genes in both cell lines were

strongly associated with inflammatory signalling and interferon response (Fig. 6C, Fig. S6D).

Notably, FOXA1 knock-down induced upregulation of master interferon response TFs,

STAT1 and STAT2, and key genes involved in the regulation of inflammation in human

cancer, including the immune checkpoint modulator *CD274* (PD-L1) (Fig. 6D), which we

also identified as a downregulated SE in both Luminal and Basal *vs* NHU cells (see earlier

Fig. 3F). While our FOXA1 ChIP-seq in Luminal and Basal cell lines showed FOXA1

binding at many interferon responsive genes, we did not observe strong FOXA1 enrichment

on *STAT1, STAT2* or *CD274* promoters or enhancers (Fig. S6E). This suggests that the

upregulation of these genes upon FOXA1 knock-down is independent of FOXA1 binding of

their regulatory elements, in agreement with recent work showing that FOXA1 directly binds

and inhibits the STAT2 protein to dampen inflammation in a chromatin-independent manner

[49]. Interestingly, if FOXA1 knock down triggered interferon response in both Luminal and

Basal models, its depletion affected the Luminal network of co-regulated TFs only in RT112

cells and not in SCaBER (Fig. 6E). PCA projection of TCGA-BLCA transcriptomes together

with that of our knock down cells on our scRNA-seq-derived Basal/Luminal signature space

confirmed that FOXA1 acute depletion induced a small but consistent shift from Luminal

towards Basal subtype only in RT112 cells (Fig. S6F, see methods). Therefore, in agreement

with a previous study [13], short-term knock-down of FOXA1 showed a consistent but mild

impact on cell identity, not sufficient to majorly alter the subtype of the luminal cells.

Altering the epigenetic landscape could indeed take a longer time. To determine if FOXA1,

through its binding to the SE repertoire, regulates the bladder cancer epigenetic landscape and

subsequently cellular identity, we produced *FOXA1* CRISPR mutant clones allowing long-

term FOXA1 inactivation in two Luminal cell lines (SD48 and RT112, Fig. S6G). Despite

fundamental differences between RT112 and SD48 cellular models and heterogeneity

between clones due to clonal selection, transcriptomic analysis of 3' RNA-seq data by PCA

distinguished CRISPR *FOXA1* mutant clones from wildtype (WT) (Fig. S6H). Importantly,

PCA projection of TCGA-BLCA transcriptomes together with that of our WT and mutant

clones on the Basal/Luminal signature space showed that mutation of *FOXA1* induced a

strong shift from the Luminal cluster to the Basal cluster (Fig. 6F). GSEA analysis confirmed

that *FOXA1* mutants were enriched for our Basal signature and depleted for our Luminal

signature (Fig. S6I)[26,27]. GSVA analysis further revealed that *FOXA1* mutant clones were

less differentiated than WT controls (Fig. 6G, [10]) and tended to express higher levels of TFs

associated with the Basal TF network (Fig. 6H, Fig. 4F). Differential gene expression analysis

revealed 1040 and 1102 Differentially Expressed Genes (DEGs) in RT112 and SD48,

respectively, when comparing *FOXA1* mutant clones to WT (Fig. S6K). *FOXA1* mutant DEGs

were associated with EMT, KRAS signalling and the inflammatory response pathway (Fig.

6I), all linked to Basal phenotypes. Intriguingly, differential analysis of *FOXA1* mutants vs

WT revealed increased expression of NR3C1 and ZBED2 in the mutants, two of the candidate

master TFs identified in our Basal TF network (Fig. 6J, Table S7). In summary, our results

demonstrate that loss of *FOXA1* promotes a clear shift from Luminal to Basal cell identity.

ZBED2, a novel Basal-associated TF involved in inflammation dampening

To further explore the interconnected network of candidate master TFs, we chose to examine

ZBED2 as one of the TFs in the Basal network since it was upregulated by *FOXA1* mutation,

and because of recent work in keratinocytes that identified a role for ZBED2 in the basal

phenotype [43]. ZBED2 expression in the TCGA-BLCA cohort is upregulated in the Ba/Sq

subtype (Fig. 7A) and correlates with poor survival prognosis (Fig. S7A). ZBED2 expression

is negatively correlated with FOXA1 expression in the TCGA cohort (Fig. 7G), but more

interestingly scRNA-seq in CCLE bladder cell lines shows that FOXA1 and ZBED2

expression are often mutually exclusive (Fig. 7C). As little is known about the ZBED2 TF, we

used ARACNE/VIPER algorithms to identify the ZBED2 regulon based on TCGA-BLCA

expression data. Interestingly, FOXA1 was predicted as a ZBED2 target, with the most

negative weight, whereas two genes associated with Basal-specific SEs (*IL7R* and *CAV1*)

were in the top 10 positive ZBED2 regulon weights (Fig. S7B). Using ZBED2 ChIP-seq data

from pancreatic cancer cell lines [44] (the only ZBED2 ChIP-seq reported so far), we found a

high confidence ZBED2 peak in the *FOXA1* promoter (Fig. 7D, left). Analysis of the RNA-

seq data from the same study revealed that ZBED2 overexpression triggered downregulation

of FOXA1 (Fig. S7C, *p*=0.004). In our data, we found that the *ZBED2* SE was highly

enriched in FOXA1 binding in SD48 luminal cells, whereas FOXA1 binding was

significantly decreased in 5637 Ba/Sq cells, and negatively correlated with ZBED2 expression

(Fig. 7D, right). Overall, these findings suggest that FOXA1 and ZBED2 could negatively

regulate each other to promote or maintain Luminal or Basal identity, respectively.

On the other hand, ZBED2 has been shown to inhibit STAT2 and dampen inflammation by

direct competition with IRF1 for Interferon Responsive Element binding in the pancreas [44].

We therefore sought to determine if ZBED2 is involved in the downregulation of interferon

signalling in bladder cancer, potentially through interfering with FOXA1-activated pathways.

Intriguingly, *ZBED2* expression in the TCGA-BLCA cohort positively correlated with
interferon gamma associated gene expression (Fig. S7D), which could be an indication that
ZBED2 increases in response to inflammation at the cell population level, or vice versa. We
then examined the correlation between ZBED2 expression and different cellular pathways at
the single-cell level using publicly available scRNA-seq data [50]. Our analysis revealed that
ZBED2 expression anti-correlates with interferon response and positively correlates with cell
cycle progression and *E2F* targets within the same cell (Fig. 7E), suggesting that the positive
correlation with interferon response in the bulk RNA-seq data reflects increased levels of
ZBED2 expression and interferon response genes in different subpopulations of cells. To test
this association further, we knocked down *ZBED2* by siRNA and performed bulk 3' RNA-seq
in two BLCA cell lines. Strikingly, downregulation of *ZBED2* increased expression of
interferon response genes and decreased expression of cell cycle progression and *E2F* target
genes (Fig. 7F). Furthermore, siZBED2 in both RT112 and SCaBER cells increased gene
expression of *STAT2* and *CD274* (Fig. 7G) and tended towards decreased cell viability (Fig.
S7E). Therefore, ZBED2 directly dampens interferon response in bladder cancer, in
agreement with its reported role in the pancreas [44]. Notably, siRNA of *FOXA1* induced
strong *STAT2* and *CD274* expression, while double knock-down of both *FOXA1* and *ZBED2*
partially dampened this response compared to siFOXA1 alone (Fig. 7G), suggesting that the
inflammatory response resulting from FOXA1 knock-down is partially dependant on ZBED2
target genes. In conclusion, both FOXA1 and ZBED2 inhibit inflammatory response and
promote bladder cancer cell survival.
460
461

Discussion:

Epigenetic mechanisms are essential for the establishment and maintenance of cellular identity notably through SE regulation of master transcriptional regulators [16]. Bladder cancer has been extensively studied at the transcriptomic level, but until very recently, very little was known about the epigenetic landscape. Here, we report the largest epigenetic profiling of both bladder cancer primary tumours and bladder cancer cell lines representative of the main molecular subtypes, as well as NHU cultures, using three histone marks ChIP-seq and paired RNA-seq. Using integrative analysis of H3K27ac, H3K27me3, H3K9me3 marks, we established a comprehensive chromatin state map of bladder cancer and showed that Basal and Luminal subgroups can be distinguished by their chromatin profiles. This map can be used to identify new genes or regulatory regions for diagnostic, prognostic or pharmacological targeting. We determined the bladder SE repertoire, and through differential analysis identified subgroup-specific activation of SEs regulating bladder cancer-associated pathways. Luminal-activated SEs were located in proximity to known key regulators of the Luminal phenotype, namely *FOXA1, GATA3,* and *PPARG* [3,32,51], validating our approach, as well as new Luminal-associated genes. In Basal samples, we identified higher enhancer activity associated with genes of interest in cancer biology, including cell surface receptors (*IL7R*, *OSMR*, *EGFR*, *MET*) and transcriptional regulators (*BNC2*, *HMGA2*, *KLF7*, *NR3C1*).

Two recent studies have also determined an enhancer landscape of bladder cancer [19,20]. Van der Vos *et al*. provided a histone methylation profiling of 19 bladder tumours, including H3K27me3, H3K4me3, and H3K4me1 ChIP-seq, another marker of active enhancers [20]. In agreement with our findings using H3K27ac, the authors reported that H3K4me1 profiles were associated with molecular subtypes and that Luminal enhancers were enriched with PPARG motifs. However, they were not able to identify any Basal-specific TF binding site enrichments. Iyyanki *et al*. conducted H3K27ac and ATAC-seq profiling in four bladder cells lines and five bladder tumours. This study also identified enrichment of PPARG binding sites in the open enhancer regions of Luminal cells and identified NPAS2 as a Luminal regulator of bladder cancer [19]. Our study corroborates and expands on the breakthrough work of these two important studies by adding RNA-seq and ChIP-seq of a new mark (H3K9me3) in a wider source of samples and by identifying subgroup-associated master regulators and co-regulated networks.

Master regulators have been linked to cellular identity, viability and resistance mechanisms,
but they can be difficult to identify, particularly for genes with low expression, unknown
binding motifs or multi-partner complexes. Algorithms such as CRCmapper [36] have been
designed to identify master TFs that regulate core regulatory circuits from SE data. However,
CRCmapper is limited to a specific set of TFs with known binding motifs and thus may
restrict the discovery of candidate master regulators. Here, to robustly reveal known and
unknown master regulators of major bladder cancer subgroups, we used two complementary
approaches. First, we identified TFs with enriched binding sites or DNA motifs in subgroup-
specific SEs. Second, we combined SE activity in our cohort with regulon analysis of TCGA
data to identify master regulator networks for Luminal and Basal subgroups. The first
approach, based on public ChIP-seq data, validated the role of TFs involved in urothelial
differentiation in Luminal SE activity, namely FOXA1 and GATA3, but also revealed that the
AP-1 complex regulates Basal SEs. AP-1 has been shown to drive reprogramming of breast
cancer cells from Luminal to Basal during resistance acquisition through high-order
assemblies of transcription factors [48]. Thus, a role for AP-1 in driving Basal enhancers and
cell identity in bladder cancer presents AP-1 inhibitors as potential therapeutic options for this
aggressive disease. Interestingly, our mapping of FOXA1 binding sites in two different cell
models indicated that the pioneer factor binds most bladder-associated SEs, even if its DNA
binding motif is mostly only found in the Luminal-specific SEs. The mapping of FOXA1
binding also confirmed that FOXA1 binding sites in Basal enhancers are associated with AP-1
localisation, suggesting that AP-1 could play an important role in the regulation of Basal
regulatory regions through FOXA1 recruitment—or trapping—at discrete chromatin loci. By
combining the ChIP-seq approach with regulon analysis, we were able to identify a Basal TF
network and highlighted new Luminal-associated TFs, in addition to known Luminal master
regulators (FOXA1, GATA3, PPARG) and the recently identified NPAS2 [19]. Here we
identified ZBED2, KLF7, HMGA2, NR3C1 as major regulators of the Basal subgroup, with
their expression restricted to the basal component of tumours, according to scRNA-seq data.
To our knowledge, the role of these TFs has not been elucidated in bladder cancer biology.
522
With regards to ZBED2, a scRNA-seq study of keratinocytes revealed that it promotes basal
cell identity of keratinocytes [43]. Another study demonstrated that in pancreatic cancer,
ZBED2 represses differentiation and dampens STAT2-mediated inflammatory response
through IRES binding competition with IRF1 [44]. Previous work showed that the three
master Luminal TFs (FOXA1, GATA3, and PPARG) had to be perturbed simultaneously to

induce a cell identity switch from luminal to basal [13]. However, we found that while short

term knock-down of FOXA1 had a mild effect on cell identity, the long-term inactivation of

FOXA1 alone through CRISPR mutation was sufficient to induce a shift from Luminal to

Basal subgroup, highlighting the role of FOXA1 in the regulation of cell fate. Moreover, we

demonstrated that this major shift is accompanied by activation of our newly identified Basal

network TF, ZBED2. Despite its known role as an activator of transcription, FOXA1 has also

been associated to direct repression of transcription [51]. ZBED2 is described as a

transcriptional repressor [44]. Therefore, FOXA1 and ZBED2 could repress each other,

defining a new cell identity regulatory loop. Through functional knock-down and knock-out

experiments, we verified that FOXA1 and ZBED2 have antagonistic but interconnected

functions in cell identity. However, ZBED2 is expressed at a very low level and additional

experiments, including overexpression models, are needed to validate its repressive function

on FOXA1, or vice versa, and its potential role in Luminal to Basal plasticity.

Finally, our work also uncovers a role for both FOXA1 and ZBED2 in the regulation of

inflammation in bladder cancer. While they play antagonistic roles in the regulation of cell

identity, we found that they share a common function in inhibiting inflammation. Short term

loss of either FOXA1 or ZBED2 triggers an inflammatory response, identified through

STAT2 overexpression, in agreement with the study of ZBED2 function in the pancreas [44].

The low FOXA1 binding enrichment at STAT2 in our FOXA1 ChIP-Seq experiments suggest

that FOXA1 could have a repressive function of inflammation presumably independent of its

chromatin binding. These conclusions are in accordance with the recent work of He et *al.* [57]

characterizing a chromatin independent function of FOXA1, which, by direct binding of

STAT2 protein, inhibited STAT2 mediated inflammation. This could explain the limited

infiltrate of luminal tumours, expressing high level of FOXA1.

Therefore, given the dual role of FOXA1 and ZBED2 in the regulation of cell identity and

inflammation, it will be important to study their link with tumour plasticity and in response to

immunotherapy. Although direct inhibitors do not yet exist, targeting FOXA1/ZBED2 or the

upstream or downstream signaling pathways, may improve sensitivity to immune based

therapies. Similarly, it will be worth studying the effect of interferon treatment on FOXA1

and ZBED2 inhibited inflammation as it could be used to overcome the inflammation

inhibition induced by these two master regulators.

If FOXA1 and ZBED2 revealed promising features, our study identifies numerous other super

enhancers, associated genes and master regulators that could be explored for pharmacological

targeting. Several selected master regulators will be studied using large scale screening in
both Luminal and Basal models.

General targeting of SE with BRD4 inhibitors has shown efficiency notably in cancers with
specific SE single mutation alteration or activation of MYC SE in leukemia or lymphoma
[17,52,53]. However, those treatments show mild efficiency in solid tumours. In addition,
enhancer rewiring has been associated to resistance to treatment. Targeting SEs could be used
alone or in combination to impair tumour viability, improve treatment or prevent resistance
acquisition.

Therefore, identification of SEs associated with bladder cancer and subgroups pave the way
for research into targeting either activated master regulators, upstream/downstream activated
pathway or even with the advent of RNA and CRispR technology, directly targeting SE.

Conclusions

Based on chromatin profiling and RNA-seq of human primary bladder tumours, bladder
cancer cell lines and normal human urothelium cultures, we provide an integrated epigenomic
and transcriptomic map of bladder cancer. With this map, we identify a consensus bladder SE
repertoire, revealing subsets of cancer-specific and subtype-specific SEs. Using
complementary approaches, we identify known and new master transcription factors of the
Luminal and Basal subgroups, with important clinical relevance. Through functional studies,
we reveal that FOXA1 maintains the Luminal cell identity, which is antagonized by the Basal
TF ZBED2. We also reveal that both FOXA1 and ZBED2 dampen inflammation signalling in
bladder cancer, through different mechanisms targeting STAT2, and promote cancer cell
survival. Overall, our findings demonstrate that the SEs and TF networks identified herein
represent prime targets for further pre-clinical investigation for bladder cancer treatment.

585
586
587
588

METHODS

Patient tumour tissue processing

We selected human tumours with an available OCT-compound frozen block from our CIT (Carte d'Identité des Tumeurs) cohort [21,22]. Each block was frozen-sectioned and stained with hematoxylin and eosin. Pathology review was performed to confirm the tumour stage and to select tumour areas, in order to enhance neoplastic content (estimated at 30 to 95%, median tumour cell content = 65%). For tumours with sufficient material, tumour-enriched areas were macrodissected from the frozen block and manually finely ground in a mortar. Frozen ground tumour tissue was kept at -80°C until further processing.

The characteristics of the tumours are shown in Table 1.

Tumour chromatin cross-linking and extraction

In order to obtain efficiently disrupted tissue, the frozen ground material (15mg) was further homogenized using a tube pestle or the TissueLyser II system (Qiagen). Disrupted tissue was then processed using the reagents from the iDeal ChIP-seq Kit for Histones (Diagenode), according to the manufacturer's instructions. Briefly, the tissue was homogenized and washed in 1ml PBS-protease inhibitor cocktail. DNA-protein cross-linking was ensured with an 8-minute incubation in 1% formaldehyde then quenched with 0.125 M glycine for 5 minutes. Cells were then washed and lysed. Centrifuged cell lysates were resuspended in shearing buffer and sonicated using the Pico Bioruptor device (Diagenode) for 15 minutes (30s ON/30s OFF). Following a centrifugation at 16000g for 10 minutes, an aliquot was reserved to control the sonication and the remaining supernatant was stored at -80°C. Sonication efficiency was controlled for each sample on the aliquot of sheared chromatin by overnight reverse cross-linking, DNA was purified using the phenol-chloroform method and 2% agarose gel electrophoresis was used to determine DNA fragment size.

Chromatin immunoprecipitation and DNA sequencing

Tumour samples with optimal chromatin fragment size (200-500 bp) were immunoprecipitated using the iDeal ChIP-seq Kit for Histones (Diagenode). Magnetic immunoprecipitation of sheared DNA-chromatin complexes (500ng) was performed overnight using a rabbit polyclonal histone H3K27acetyl ChIP Grade antibody (ab4729, Abcam), H3K27me3 (Active Motif, ref. 39155), and H3K9me3 (Active Motif, ref. 39161). Magnetic immunoprecipitation beads were washed the following day. The captured chromatin

as well as non-immunoprecipitated input chromatin underwent elution and reverse cross-

linking steps. DNA purification was performed using iPure magnetic beads.

Immunoprecipitation (IP) efficiency was verified by qPCR according to the manufacturer's

protocol. Library preparation from IP DNA and input DNA was performed using the

Diagenode MicroPlex Library Preparation kit v2. The resulting amplified libraries were

assessed using the Bioanalyzer system 2100 (Agilent) and sequenced using the HiSeq 4000

platform (Illumina) as single-read 50 base reads, following Illumina's instructions. Reads

were aligned to the reference genome (Hg19) using Bowtie 1.0.0.

Cell culture

The human bladder cancer derived cell lines RT112, 5637, KK47, and SCaBER were

obtained from DSMZ (Heidelberg, Germany). MGH-U3, KK47 and SD48 cell lines were

provided by Yves Fradet (CRC, Quebec), Jennifer Southgate laboratory (previously of Cancer

Research Unit, St James's University Hospital, Leeds, UK), and Henri Mondor Hospital

(Créteil, France), respectively. The L1207 cell line was derived from tumour T1207 (25).

RT112 and 5637 cells were cultured in RPMI medium, L1207 were cultured in DMEM-F12

and all the other cell lines were cultured in DMEM medium. All cell media were

supplemented with 10% fetal bovine serum (FBS). Normal human urothelium (NHU) cells

were obtained from normal ureter urothelium from healthy kidney donors from Foch hospital

and were cultured as previously described (26). All cells were cultured at 37°C in an

atmosphere of 5% CO_2 and were routinely tested for mycoplasma contamination.

Cell line ChIP-seq

Cell lines cultures were crosslinked directly in the growing medium with formaldehyde 1%

for 10 minutes at room temperature. The reaction was stopped by adding Glycine with a final

concentration of 0.125M for 10 minutes at room temperature. Fixed cells were rinsed 3 times

with PBS containing protease inhibitors, pelleted, and resuspended in lysis buffer (10mM

EDTA, pH8, 50mM Tris-HCl pH8, SDS 1%). After centrifugation, the ChIP was performed

using ChIP-IT High Sensitivity kit (Active Motif, Carlsbad, CA, USA), following the

manufacturer's instructions. Chromatin was sonicated in a bioruptor Pico device (Diagenode)

for 10 min (30s ON/30s OFF). Sheared chromatin was immunoprecipitated using an H3K27ac

antibody (Abcam ab4729). Sheared chromatin was used as input-DNA control.

654 ChIP-seq libraries were prepared using NEXTflex ChIP-Seq Kit (#5143-02, Bioo Scientific)

655 following the manufacturer's protocol (V12.10) with some modifications. Briefly, 10 ng of

656 ChIP enriched DNA were end-repaired using T4 DNA polymerase, Klenow DNA polymerase

657 and T4 PNK, then size selected and cleaned-up using Agencourt AMPure XP beads

658 (#A63881, Beckman). A single 'A' nucleotide was added to the 3' ends of the blunt DNA

659 fragments with a Klenow fragment (3' to 5'exo minus). The ends of the DNA fragments were

660 ligated to double stranded barcoded DNA adapters (NEXTflex ChIP-Seq Barcodes - 6,

661 #514120, Bioo Scientific) using T4 DNA Ligase. The ligated products were enriched by PCR

662 and cleaned-up using Agencourt AMPure XP beads. Prior to sequencing, DNA libraries were

663 checked for quality and quantified using a 2100 Bioanalyzer (Agilent). The libraries were

664 sequenced on the Illumina Hi-Seq 2500 as single-end 50 base reads following Illumina's

665 instructions. Sequence reads were mapped to reference genome hg19 using Bowtie 1.0.0.

667 **ChIP-seq data analysis**

668 Peak detection was performed using MACS2 (model-based analysis for ChIP-seq

669 v2.1.0.20140616) software under settings where input samples were used as a negative

670 control. We used a default cutoff and -B option for broad peaks.

671 To identify enhancer regions in each tumour we used ROSE (Ranked Ordering of Super-

672 Enhancers) algorithm [17,54], with the following parameters: 12.5kb stitching distance,

673 exclusion of promoter regions 2500 bp around TSS. For each sample, stitched enhancer

674 regions are normalized, ranked and plotted. The regions above the inflexion point are

675 considered super-enhancers by the algorithm. However, the number of called super-enhancers

676 was notably lower in cases with a known amplified gene. The very high H3K27ac signal in

677 the amplified region likely created a bias in the plot of ranked enhancers. To correct for this

678 bias, we set a threshold to the top 1000 ranked enhancers to select candidate super-enhancer

679 regions.

680 Heatmaps and PCA of ChIP-seq signal were performed using Diffbind R package (version

681 2.16.0) or Easeq [55]. For super-enhancers analysis, the top 1000 SE regions of either

682 tumours or cell lines were merged for a consensus using Diffbind. Then, H3K27ac signal was

683 calculated in the consensus peak for each sample. Differential analysis between molecular

684 subtypes was performed with Diffbind and DESeq2 default parameters using both IP and

685 input bam files, and a file containing the consensus super enhancer regions evaluated for

686 differential analysis as input. Regions with an pval <0.05 were considered differentially

687 bound.

Genomic annotation and pathway enrichment analyses were performed using ChIPseeker,
clusterProfiler and GREAT (28).

Chromatin Binding enrichment

Factor binding analyses were performed using public data available in Cistrome DB Toolkit
[37,38]. DNA binding motif analysis was performed using HOMER known motif function
[39].

Genomic annotation of the SE regions and cis-regulatory genes

SE activity and gene expression was jointly analysed to determine the cis-regulatory between
the SE and genes on proximity. In brief, the overlap / proximal genes corresponding to each
SE were annotated using GREAT/ROSE tools, as the candidate proximal genes regulated by
the SE [54,56]. The spearman correlation coefficients between SE activity (H3K27ac read
counts, log2RPKM normalized) and the expression of the candidate genes (RLE normalized)
were calculated in the tumours. The gene whose expression showing the highest correlation
with the activity of the corresponding SE was determined as the gene most likely regulated by
the SE. The SE-gene relationships within the top 1% were also given, not limited to the
proximal genes. The number of germline single nucleotide polymorphisms (SNPs) within a
given SE as well as their association with bladder cancer (median –log10 p-value) was
provided based on the UK biobank GWAS summary statistics (Neale lab Round 2, ukb-d-
C67, extracted from the MRC IEU OpenGWAS database) [57]. The germline SNPs falling
within the SEs and with published GWAS-level association with BCa or with –log10P.value
> 5 in GWAS summary statistics (PhenoScanner v2 database) were provided as GWAS SNPs
within the SEs [58]. For the genes most likely regulated by a given SE, we provided their
median CERES dependency score of all and urothelial cancer cells from the Cancer
Dependency Map database [45], as well as the p-value for difference between the two. We
checked if any bias compared to the background in mutation type (missense, non-sense,
synonymous, etc.) for the protein coding genes by Chi-square test. We checked if they were
within the list of established cancer genes, including the COSMIC Cancer Gene Census and
Network of Cancer Genes 6 [59,60].

Chromatin States Analysis

Chromatin state and correlation with expression

ChromHMM was used to identify chromatin states. The genome was analyzed at 1000 bp

intervals and the tool was used to learn models from the 3 histone marks ChIP-seq reads files

and corresponding Input controls. A model of 6 states was selected and applied on all

samples. The 6 states identified were then given functional annotation based on histone marks

enrichment and ENCODE published chromatin states.

We checked the genome-wide association between gene expression and chromatin states of

the TSS, in both tumour and cell line samples. In each tumour / cell line, we classified the

genes according to the chromatin states of the TSS. For genes with multiple TSS, the

chromatin states showing frequency dominance was considered. We then calculated the

median expression of the genes by their TSS categories in each sample, and assessed the

distribution of the median expression by chromatin state across all tumour and cell line

samples.

ChromHMM output files were concatenated using the unionbed function from BEDTools, by

which a consensus sample-by-states matrix was created, where in each cell the chromatin

state corresponding to the column's chromosome region in the row's sample, excluding

regions from sexual chromosomes, with all samples included (n = 24, including 15 tumours

and 9 cell lines).

We next performed unsupervised analysis of the integrated chromatin states in tumour and

cell line samples.

Selection of most informative features

We first looked for the most informative features in the consensus sample-by-states matrix

where in each cell the chromatin state corresponding to the column's chromosome region in

the row's sample, excluding regions from sexual chromosomes, with all samples included (n

= 24, including 15 tumours and 9 cell lines). We excluded genome regions of 'no mark' state

to enrich our feature selection with active regions and filtered features with top 1% Shannon's

entropy. Then to further select informative feature, we signed-rank transformed the data: For

each state, given the constitution of the histone marks and association with gene expression,

we performed a numeric transformation of the categorical states by assigning numeric values

to categorical states, as 3 to E2 (Active Enhancer / Promoter), -3 to E4 (Repressed

Chromatin), 2 to E3 (Bivalent Enhancer / Promoter), -2 to E6 (Heterochromatin / ZNF/

Repeats), 1 to E1 (Quiescent / No mark), and -1 to E5 (Quiescent / Weak repression). This

allow to further increased the selection power as the top 1% features by variance ranking. We

then used these top 1% variable features for subsequent analysis.

Dimension reduction and functional ontology analysis

We performed dimension reduction and visualization taking directly the categorical format of
the above described selected features using multiple correspondence analysis (MCA). To
explore the biological significance of the regions that contributed to the dimension that
distinguishes the non-basal and basal clusters, the chromosome segments' loading estimates
to the Dim 2 were extracted from the MCA outputs and regions with a p-value < 0.05 for the
loading estimate were included (n = 12,198). Genes mapped to Dim 2 contributing regions
were pre-ranked by loading estimates for gene-set enrichment analysis (GSEA) by which we
identified multiple biological gene sets / ontologies associated with Dim 2. The gene sets
collections were retrieved from the Broad Institute Molecular Signature Database, spanning
the H (hall mark gene sets), C2 (curated gene sets, eg. pathways), C3 (regulatory target gene
sets), C5 (ontology gene sets, eg. Gene Ontology), C6 (oncogenic signature gene sets), and C8
(cell type signature gene sets) categories, using the msigdbr R package [27,61].
As complementary exploration, we in the meantime performed dimensional reduction to the
numeric transformed data of the selected features, using MDS. Similar to what was observed
in MCA, the Dim 2 represents the dimension that distinguishes the basal versus non-basal
samples, and Dim 1 separates cell lines from tumours, suggesting potential batch effect and/or
in vitro culture-specific effect. We then adjusted for these latent effects to obtain a refined
clustering (basically on Dim 2), using the MNN algorithm implemented in the fastMNN
function of the batchelor R Bioconductor package [62].
We then calculated for the chromosomal segments (ie. features of the consensus sample-by-
states matrix), the difference in the numeric chromatin state scores between basal and non-
basal groups, named chromatin state score difference basal vs non-basal. A negative score
difference indicates stronger activation in the non-basal group, and a positive one indicates
stronger activation in the basal group. For subsequent function analysis, we performed
expression quantitative trait locus (eQTL) mapping to refine the segments to the ones
significantly linked with associated gene expression, and limited the analysis to the significant
eQTL pairs (p-value < 0.05, n = 4,377). We then analysed the distribution of the chromatin
state score difference of the segments corresponding to the luminal and basal cell type
signature genes.
784

Cell treatments, cell viability assay
For siRNA treatments, cells were reverse transfected using Lipofectamine RNAi max
(Invitrogen) using 10 ng of siRNA (siRNA table in additional information).

For CRispR mutant cell lines production, RT112 and SD48 cells were plated at 80%
confluence and the day after transfected with vectors expressing Cas9 an gRNA
(VectorBuilder, see additional information) using Fugene HD transfection reagent. 48h post
transfection, cells were selected using Puromycin (2µg/µL) during 4 days. After 2 weeks,
clonal selection was performed using clonal dilution. FOXA1 mutation was assessed by
Western Blot (anti-FOXA1 Abcam ab23738), PCR and genomic DNA sequencing.
Cell Viability was assessed in 96 well plates using CellTiter-Glo® Luminescent Cell Viability
Assay (Promega).
796

797 **RNA extraction**
798 Cell lines RNA (treated or not) were extracted using Qiagen RNeasy kit coupled with DNAse
799 treatment. Tumours RNA were extracted using Triple extraction protocol.
800

801 **RTqPCR**
802 Reverse transcription was performed with 0.5 to 1µg of total RNA, with the High-Capacity
803 cDNA reverse transcription kit (Applied Biosys- tems). Genes were amplified with the Roche
804 Taqman master mix (Roche) or SYBR Green PCR kit purchased from Applied Life
805 Technologies. See additional information for primers sequences and probes references.
806

RNAseq
Tumours RNA extraction, sequencing
RNA sequencing – Pool Nugen Ovation Solo – Run D345
RNA sequencing libraries: Kit Nugen
The pool of libraries was quantified using a qPCR method (KAPA library quantification kit,
Roche). The sequencing was carried out using paired-end mode (PE100) on a Illumina
Novaseq 6000 instrument, using a custom primer (provided into the Nugen kit) to initiate the
Read 1 sequencing. The target number of reads was about 50 million paired-reads per sample.
Cell lines RNAseq
RNA sequencing – 16mRNA - run B160-B163
RNA sequencing libraries were prepared from 1µg of total RNA using the Illumina TruSeq
Stranded mRNA Library preparation kit (Illumina) which allows to perform a strand specific
RNA sequencing. A first step of polyA selection using magnetic beads is done to focus
sequencing on polyadenylated transcripts. After fragmentation, cDNA synthesis was

performed and resulting fragments were used for dA-tailing and then ligated to the TruSeq

indexed adapters. PCR amplification was finally achieved to create the final cDNA library (12

cycles). The resulting barcoded libraries were then equimolarly pooled and quantified using a

qPCR method (KAPA library quantification kit, Roche). The sequencing was carried out

using paired-end mode (PE100) on a Illumina HiSeq2000 instrument. The sequencing

configuration was set to reach an average of 100 million paired-reads per sample.

3'RNA-seq (Lexogen 3'Seq)

RNA sequencing libraries were prepared from 200ng of total RNA using the QuantSeq FWD

3'mRNA Seq LEXOGEN Standard (CliniSciences). Libraries were prepared according to the

manufacturer's recommendations. The first step enables the synthesis of double strand cDNA,

by revers transcription, using oligo dT priming. A qPCR optimization step was performed in

order to estimate the most appropriate number of PCR cycles for library amplification. The

resulting amplified and barcoded libraries were then equimolarly pooled and quantified using

a qPCR method (KAPA library quantification kit, Roche). The sequencing was carried out

using single-read mode (SR100) on a Illumina Novaseq 6000 instrument. The sequencing

configuration was set to reach an average of 10 million reads per sample.

RNA-seq analysis

Our RNA-seq as well as RNA-seq from pubic data repository were integrated using Deseq2

default parameters and VST normalisation. 3'RNA-seq were analysed with Deseq2 and RPM

normalisation.

Assignment of MIBC and NMIBC subtypes

Gene expression data of the most tumour cases was previously generated and published

[21,22]. We assigned consensus classes using the previously generated gene expression data

using ConsensusMIBC (v1.1.0) R package [3] (Table S1). Given potential intra-tumour

molecular heterogeneity, we also to verify the subtype in our ChIP-seq sampled tumour area

using RNAseq from the same powder using the same ConsensusMIBC (v1.1.0) R package.

NMIBC samples (n=2) were classified using classifyNMIBC R Package [7].

General bioinformatics, statistical analyses and public data

Plots and statistical analyses were performed in R software version 3.6.1, using ggpubr

package. Wilcoxon and Kruskal-Wallis tests were used to test the association between

continuous and categorical variables, for 2 categories or >2 categories, respectively. P-values <0.05 were considered statistically significant. Pairwise correlation of gene expression was calculated using Pearson coefficient and plotted using complexHeatmap R package. All gene expression heatmaps show mean-centered log2-transformed normalized counts of each represented gene.

TCGA-BLCA MIBC RNA-seq data were downloaded from TCGA data portal using TCGAbiolinks package (R), raw counts were normalized to account for different library size and the variance was stabilized with VST function in the DESeq2 R-package [63]. TCGA-BLCA samples (n=404) were classified using the consensus system using consensusMIBC R package.

CCLE urinary tract cell line gene expression were downloaded from the DepMap portal (*https://depmap.org/portal/download/*). For consensus classification of CCLE bladder cancer cell lines, we adapted this classification considering only genes expressed by both tumours and cell lines. Ten cell lines were classified as Ba/Sq, all other were grouped as non-Ba/Sq.

MGHU3 RNA-seq bulk data were download from GEO (accession number, GSE171129).

Survival analysis

For Kaplan Meier survival analyses testing the association of gene expression and overall survival, we used http://tumorsurvival.org/index.html tool and divided the samples based on mean +/- sd. Log-rank P values were calculated to test the association between overall survival and low vs high expression groups.

Public scRNA-seq and Basal/Luminal signature

We downloaded the log2 TPM normalized gene expression of single cells from a Ba/Sq subtype MIBC tumour from the GEO database (accession number, GSM4307111). Initial quality control excluded genes expressed in less than 3 cells and cells with less than 200 genes. The top 2000 variable genes were used as features for subsequent PCA and the first 9 principal components were used for cell clustering and visualization by uniform manifold approximation and projection (UMAP) embedding. The marker genes of the luminal and basal tumour cells were calculated with Wilcoxon test based approach. The single cell RNA-seq data analyses were performed using the Seurat v4 package with default parameters unless otherwise specified.

Given the single-cell derived luminal and basal tumour cell signature was based on single-cell sequencing of primary *in vivo* tumour sample, and the *FOXA1* knock-out perturbation signature is likely limited to the genes regulated by *FOXA1* in an *in vitro* setting, it is important to adopt the cell subtype signatures to refine to the marker genes regulated by

FOXA1, as a *FOXA1*-depdent luminal-basal plasticity signature which could be then used for

further analyses involving *in vitro* transcriptomes. We first compared the perturbation and

single-cell signatures by GSEA (perturbation DEG effect for ranking, and luminal / basal

signatures as gene sets of interest) and found that in RT112 cell line, there was both

significant enrichment of luminal signature in genes down-regulated in *FOXA1 KO* clones

and significant enrichment of basal signature in genes up-regulated in *FOXA1 KO* clones. We

then took the leading edge genes as the adopted *FOXA1*-depdent plasticity signature. As

validation, this adopted signature showed similar enrichment in RT112 *FOXA1* KD assays

and SD48 *FOXA1* KO assays, while the original cell type signature failed.

Regulons

The regulatory network was reverse engineered by ARACNe-AP [41] from human urothelial

cancer tissue datasets profiled by RNA-seq from TCGA. ARACNe was run with 100

bootstrap iterations using all probe-clusters mapping to a set of 1,740 transcription factors.

Parameters used were standard parameters, with Mutual Information p-value threshold of 10^{-8}.

The VIPER (Virtual Inference of Protein-activity by Enriched Regulon analysis)[42](R

package viper 1.24), using the regulatory network obtained from ARACNE on urothelial

cancer, and we computed the enrichment of each regulon on the gene expression signature

using different implementations of the analytic Rank-based Enrichment Analysis algorithm.

SE Correlation Network

To build SE driven correlation network, we first selected genes regulated by SE defined as TF

in Lambert *et al.* [40]. Next using TCGA regulon VIPER score, we calculated the mean

regulon score by subtype (Luminal, Ba/Sq or Stroma-Rich and kept TFs with mean regulon >

2 or < -2 (n=75). We further restricted the list to TFs with a minimum expression correlation

of 0.5 in TCGA to build correlation network using igraph.

Declarations

Ethics approval and consent to participate

All patients provided written informed consent. All research in this study conformed to the principles of the Helsinki Declaration.

Consent for publication

All patients consent to participate

Availability of data and materials

The dataset(s) supporting the conclusions of this article are available in the GEO repository under accession numbers: [will be submitted upon acceptance]

Competing interests

The authors declare that they have no competing interests

Funding

The work was supported by grants from Ligue Nationale Contre le Cancer: (H.N-K, J.F., C.G., X-Y.M., L.C., F.D., C.K., Y.A., F.R., I. B-P.) as an associated team (Equipe labellisée), the "Carte d'Identité des Tumeurs" program initiated, developed and funded by Ligue Nationale Contre le Cancer, and a post-doctoral fellowship supporting H.N-K. J.F. was supported by the Fondation ARC pour la recherche sur le cancer, L.C. by FRM (Fondation Recherche Médicale) and X-Y.M. was supported by a fellowship from ITMO Cancer AVIESAN, within the framework of Cancer Plan. The work was also supported by a "PL-Bio" project funded by INCa (2016-146), the French Ministry of Education and Research, the CNRS, and the Institut Curie.

Authors' contributions

H. N-K, J. F, performed the ChIP-seq and RNA-seq experiments.

C.K established the NHU primary cells cultures.

Y. N. and T. L. provided clinical insight for analysis of the CIT cohort.

JF under Y.A supervision performed histo-pathological analysis and E.G the macrodissection of the tumors.

A.R under D.G supervision optimized RNA sequencing of primary tumour samples.

H.N-K performed the functional experiments, siRNA, CRispR, cellular assays.

H. N-K, X-Y.M., C.G., J.F, T.Y., L.C., F.D, E.C. carried out the bioinformatics analysis. In

particular, H.N-K analysed ChIP-seq data, performed ChromHMM integration, SE

identification with J.F., differential analysis, ontology/GSEA analysis, 3'RNA-seq analyses,

and established coregulatory network with the help of J.F., F.D and D.J. C.G. designed R

scripts and participated to several analyses (GSVA, SE differential analysis, classifications).

L.C performed the regulon analysis. X-Y. M. analysed ChromHMM output, publicly

available scRNA-seq, derived Basal/Luminal signature, 3'RNA-seq, performed GSEA

analysis, optimized SE annotation. T.Y under I.D. supervision processed the ChIPseq data.

E.C processed RNA-seq data of cell lines and centralized the data for bioinformatics analyses.

H. N-K, J.F., X-Y.M, C.G, D.J, I. B-P, F.R. wrote the manuscript.

The study was conceived by H. N-K, J.F and supervised by Y.A, I.B-P and F.R.

All authors gave critical insights and approved the final version for publication.

Acknowledgements

We thank the NGS platform of Institut Curie, directed by Sylvain Baulande.

References:

1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin [Internet]. Wiley; 2018 [cited 2021 Jan 25];68:394–424. Available from: https://pubmed.ncbi.nlm.nih.gov/30207593/

2. Humphrey PA, Moch H, Cubilla AL, Ulbright TM, Reuter VE. The 2016 WHO Classification of Tumours of the Urinary System and Male Genital Organs—Part B: Prostate and Bladder Tumours. Eur Urol [Internet]. Elsevier B.V.; 2016 [cited 2021 Jan 25];70:106–19. Available from: https://pubmed.ncbi.nlm.nih.gov/26996659/

3. Kamoun A, de Reyniès A, Allory Y, Sjödahl G, Robertson AG, Seiler R, et al. A Consensus Molecular Classification of Muscle-invasive Bladder Cancer[Formula presented]. Eur Urol [Internet]. Elsevier B.V.; 2020 [cited 2020 Nov 2];77:420–33. Available from: https://pubmed.ncbi.nlm.nih.gov/31563503/

4. Marzouka NAD, Eriksson P, Rovira C, Liedberg F, Sjödahl G, Höglund M. A validation and extended description of the Lund taxonomy for urothelial carcinoma using the TCGA cohort. Sci Rep [Internet]. Nature Publishing Group; 2018 [cited 2021 Jan 25];8. Available from: https://pubmed.ncbi.nlm.nih.gov/29487377/

5. Choi W, Porten S, Kim S, Willis D, Plimack ER, Hoffman-Censits J, et al. Identification of Distinct Basal and Luminal Subtypes of Muscle-Invasive Bladder Cancer with Different Sensitivities to Frontline Chemotherapy. Cancer Cell. 2014;25:152–65.

6. Rebouissou S, Hérault A, Letouzé E, Neuzillet Y, Laplanche A, Ofualuka K, et al. CDKN2A homozygous deletion is associated with muscle invasion in FGFR3-mutated urothelial bladder carcinoma. J Pathol [Internet]. 2012;227:315–24. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22422578

7. Viborg Lindskrog S, Prip F, Lamy P, Taber A, Groeneveld CS, Birkenkamp-Demtröder K, et al. An integrated multi-omics analysis identifies prognostic molecular subtypes of non-muscle-invasive bladder cancer. Available from: https://doi.org/10.1038/s41467-021-22465-w

8. Hedegaard J, Lamy P, Nordentoft I, Algaba F, Høyer S, Ulhøi BP, et al. Comprehensive Transcriptional Analysis of Early-Stage Urothelial Carcinoma. Cancer Cell. 2016;30:27–42.

9. Fishwick C, Higgins J, Percival-Alwyn L, Hustler A, Pearson J, Bastkowski S, et al. Heterarchy of transcription factors driving basal and luminal cell phenotypes in human urothelium. Cell Death Differ [Internet]. Nature Publishing Group; 2017 [cited 2021 Jan 25];24:809–18. Available from: https://pubmed.ncbi.nlm.nih.gov/28282036/

10. Eriksson P, Aine M, Veerla S, Liedberg F, Sjödahl G, Höglund M. Molecular subtypes of

urothelial carcinoma are defined by specific gene regulatory systems. BMC Med Genomics

[Internet]. BioMed Central Ltd.; 2015 [cited 2021 Jan 25];8. Available from:

https://pubmed.ncbi.nlm.nih.gov/26008846/

11. Yamashita H, Kawasawa YI, Shuman L, Zheng Z, Tran T, Walter V, et al. Repression of

transcription factor AP-2 alpha by PPARγ reveals a novel transcriptional circuit in basal-

squamous bladder cancer. Oncogenesis [Internet]. Springer Nature; 2019 [cited 2021 Jan

25];8. Available from: https://pubmed.ncbi.nlm.nih.gov/31772149/

12. Karni-Schmidt O, Castillo-Martin M, HuaiShen T, Gladoun N, Domingo-Domenech J,

Sanchez-Carbayo M, et al. Distinct expression profiles of p63 variants during urothelial

development and bladder cancer progression. Am J Pathol [Internet]. Elsevier Inc.; 2011

[cited   2021   Jan   25];178:1350–60.   Available   from:

https://pubmed.ncbi.nlm.nih.gov/21356385/

13. Warrick JI, Walter V, Yamashita H, Chung E, Shuman L, Amponsa VO, et al. FOXA1,

GATA3 and PPARIγ Cooperate to drive luminal subtype in bladder cancer: A molecular

analysis of established human cell lines. Sci Rep [Internet]. Nature Publishing Group; 2016

[cited 2020 Nov 2];6:1–15. Available from: http://www.nature.com/articles/srep38531

14. Chapuy B, McKeown MR, Lin CY, Monti S, Roemer MGM, Qi J, et al. Discovery and

Characterization of Super-Enhancer-Associated Dependencies in Diffuse Large B Cell

Lymphoma. Cancer Cell [Internet]. Elsevier Inc.; 2013;24:777–90. Available from:

http://dx.doi.org/10.1016/j.ccr.2013.11.003

15. Ott CJ, Federation AJ, Schwartz LS, Kasar S, Klitgaard JL, Lenci R, et al. Enhancer

Architecture and Essential Core Regulatory Circuitry of Chronic Lymphocytic Leukemia.

Cancer Cell [Internet]. Cell Press; 2018 [cited 2021 Jan 25];34:982-995.e7. Available from:

https://pubmed.ncbi.nlm.nih.gov/30503705/

16. Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-André V, Sigova AA, et al. Super-enhancers

in the control of cell identity and disease. Cell [Internet]. 2013 [cited 2014 Jul 10];155:934.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/24119843

17. Lovén J, Hoke HA, Lin CY, Lau A, Orlando DA, Vakoc CR, et al. Selective inhibition of

tumor oncogenes by disruption of super-enhancers. Cell [Internet]. 2013 [cited 2018 Jul

6];153:320–34.   Available   from:

http://linkinghub.elsevier.com/retrieve/pii/S0092867413003930

18. Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, et al. An integrated

encyclopedia of DNA elements in the human genome. Nature [Internet]. Nature Publishing

Group; 2012 [cited 2021 Jan 28];489:57–74. Available from: http://encodeproject.org/ENCODE/

19. Iyyanki T, Zhang B, Jin Q, Yang H, Liu T, Wang X, et al. Subtype-specific epigenomic landscape and 3D genome structure in bladder cancer. Genome Biology; 2020;1–20.

20. van der Vos KE, Vis DJ, Nevedomskaya E, Kim Y, Choi W, McConkey D, et al. Epigenetic profiling demarcates molecular subtypes of muscle-invasive bladder cancer. Sci Rep [Internet]. Nature Publishing Group UK; 2020 [cited 2020 Nov 24];10:1–10. Available from: https://doi.org/10.1038/s41598-020-67850-5

21. Biton A, Bernard-Pierrot I, Lou Y, Krucker C, Chapeaublanc E, Rubio-Pérez C, et al. Independent Component Analysis Uncovers the Landscape of the Bladder Tumor Transcriptome and Reveals Insights into Luminal and Basal Subtypes. Cell Rep [Internet]. Elsevier; 2014 [cited 2020 Nov 2];9:1235–45. Available from: https://pubmed.ncbi.nlm.nih.gov/25456126/

22. Rebouissou S, Bernard-Pierrot I, de Reyniès A, Lepage M-LL, Krucker C, Chapeaublanc E, et al. EGFR as a potential therapeutic target for a subset of muscle-invasive bladder cancers presenting a basal-like phenotype. Sci Transl Med [Internet]. American Association for the Advancement of Science; 2014 [cited 2020 Nov 2];6:244ra91. Available from: https://pubmed.ncbi.nlm.nih.gov/25009231/

23. Shi MJ, Meng XY, Fontugne J, Chen CL, Radvanyi F, Bernard-Pierrot I. Identification of new driver and passenger mutations within APOBEC-induced hotspot mutations in bladder cancer. Genome Med [Internet]. BioMed Central Ltd; 2020 [cited 2021 Dec 16];12:85. Available from: https://genomemedicine.biomedcentral.com/articles/10.1186/s13073-020-00781-y

24. Blum Y, Meiller C, Quetel L, Elarouci N, Ayadi M, Tashtanbaeva D, et al. Dissecting heterogeneity in malignant pleural mesothelioma through histo-molecular gradients for clinical applications. Nat Commun [Internet]. Springer US; 2019;10:1–12. Available from: http://dx.doi.org/10.1038/s41467-019-09307-6

25. Ernst J, Kellis M. ChromHMM: Automating chromatin-state discovery and characterization [Internet]. Nat. Methods. Nature Publishing Group; 2012 [cited 2021 Jan 28]. p. 215–6. Available from: http://compbio.mit.edu/ChromHMM/.

26. Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, et al. PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat Genet [Internet]. Nature Publishing Group; 2003 [cited 2021 Dec 17];34:267–73. Available from: http://www.nature.com/naturegenetics

27. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene
set enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles [Internet]. 2005. Available from:
www.pnas.orgcgidoi10.1073pnas.0506580102

28. Parolia A, Cieslik M, Chu S-CC, Xiao L, Ouchi T, Zhang Y, et al. Distinct structural
classes of activating FOXA1 alterations in advanced prostate cancer. Nature [Internet]. 2019
[cited 2019 Jul 5];571:413–8. Available from: http://www.nature.com/articles/s41586-019-
1347-4

29. Kim E, Choi S, Kang B, Kong JH, Kim Y, Yoon WH, et al. Creation of bladder
assembloids mimicking tissue regeneration and cancer. Nature [Internet]. Springer US;
2020;588:664–9. Available from: http://dx.doi.org/10.1038/s41586-020-3034-x

30. Panwala T, Rajdev S, Mulla S, Pérez Rodríguez NM, Galloway RL, Blau DM, et al. No
主観的健康感を中心とした在宅高齢者における 健康関連指標に関する共分散構造分
析 Title. PLoS Negl Trop Dis [Internet]. 2017;9:1–14. Available from:
https://doi.org/10.1016/j.tmaid.2020.101607%0Ahttps://doi.org/10.1016/j.ijsu.2020.02.034%
0Ahttps://onlinelibrary.wiley.com/doi/abs/10.1111/cjag.12228%0Ahttps://doi.org/10.1016/j.s
sci.2020.104773%0Ahttps://doi.org/10.1016/j.jinf.2020.04.011%0Ahttps://doi.o

31. Zhang X, Choi PS, Francis JM, Gao GF, Campbell JD, Ramachandran A, et al. Somatic
Superenhancer Duplications and Hotspot Mutations Lead to Oncogenic Activation of the
KLF5 Transcription Factor. Cancer Discov [Internet]. 2018 [cited 2018 Jul 6];8:108–25.
Available from: http://cancerdiscovery.aacrjournals.org/lookup/doi/10.1158/2159-8290.CD-
17-0532

32. Robertson AG, Kim J, Al-Ahmadie H, Bellmunt J, Guo G, Cherniack AD, et al.
Comprehensive Molecular Characterization of Muscle-Invasive Bladder Cancer. Cell. Cell
Press; 2017;171:540-556.e25.

33. Weinstein JN, Akbani R, Broom BM, Wang W, Verhaak RGWW, McConkey D, et al.
Comprehensive molecular characterization of urothelial bladder carcinoma. Nature [Internet].
Nature Publishing Group; 2014 [cited 2014 Jan 29];507:315–22. Available from:
https://www.nature.com/articles/nature12965

34. Sim WJ, Iyengar PV, Lama D, Lui SKL, Ng HC, Haviv-Shapira L, et al. c-Met activation
leads to the establishment of a TGFβ-receptor regulatory network in bladder cancer
progression. Nat Commun [Internet]. Nature Publishing Group; 2019 [cited 2021 Jan 25];10.
Available from: https://pubmed.ncbi.nlm.nih.gov/31554791/

35. Chen L, Huang M, Plummer J, Pan J, Jiang YY, Yang Q, et al. Master transcription
factors form interconnected circuitry and orchestrate transcriptional networks in oesophageal
adenocarcinoma. Gut [Internet]. 2020;69:630–40. Available from:
https://gut.bmj.com/lookup/doi/10.1136/gutjnl-2019-318325

36. Saint-André V, Federation AJ, Lin CY, Abraham BJ, Reddy J, Lee TI, et al. Models of
human core transcriptional regulatory circuitries. Genome Res. 2016;26:385–96.

37. Zheng R, Wan C, Mei S, Qin Q, Wu Q, Sun H, et al. Cistrome Data Browser: Expanded
datasets and new tools for gene regulatory analysis. Nucleic Acids Res [Internet]. Oxford
University Press; 2019 [cited 2021 Sep 13];47:D729–35. Available from: https://ncbi.

38. Mei S, Qin Q, Wu Q, Sun H, Zheng R, Zang C, et al. Cistrome Data Browser: A data
portal for ChIP-Seq and chromatin accessibility data in human and mouse. Nucleic Acids Res
[Internet]. Oxford University Press; 2017 [cited 2021 Sep 13];45:D658–62. Available from:
https://academic.oup.com/nar/article/45/D1/D658/2333932

39. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple Combinations of
Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for
Macrophage and B Cell Identities. Mol Cell [Internet]. Mol Cell; 2010 [cited 2021 Sep
13];38:576–89. Available from: https://pubmed.ncbi.nlm.nih.gov/20513432/

40. Lambert SA, Jolma A, Campitelli LF, Das PK, Yin Y, Albu M, et al. The Human
Transcription Factors. Cell [Internet]. Elsevier Inc.; 2018;172:650–65. Available from:
https://doi.org/10.1016/j.cell.2018.01.029

41. Lachmann A, Giorgi FM, Lopez G, Califano A. ARACNe-AP: gene network reverse
engineering through adaptive partitioning inference of mutual information. Bioinformatics
[Internet]. Oxford University Press; 2016 [cited 2021 Sep 16];32:2233–5. Available from:
https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btw216

42. Alvarez MJ, Shen Y, Giorgi FM, Lachmann A, Ding BB, Hilda Ye B, et al. Functional
characterization of somatic mutations in cancer using network-based inference of protein
activity. Nat Genet [Internet]. Nature Publishing Group; 2016 [cited 2021 Sep 16];48:838–47.
Available from: https://www.nature.com/articles/ng.3593

43. Finnegan A, Cho RJ, Luu A, Harirchian P, Lee J, Cheng JB, et al. Single-cell
transcriptomics reveals spatial and temporal turnover of keratinocyte differentiation
regulators. Front Genet. 2019;10:1–14.

44. Somerville TDD, Xu Y, Wu XS, Maia-Silva D, Hur SK, De Almeida LMN, et al. ZBED2
is an antagonist of interferon regulatory factor 1 and modifies cell identity in pancreatic
cancer. [cited 2021 Jan 26]; Available from: https://www.pnas.org/lookup/suppl/

45. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The
Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity.
Nature. 2012;483:603–7.

46. Ghandi M, Huang FW, Jané-Valbuena J, Kryukov G V., Lo CC, McDonald ER, et al.
Next-generation characterization of the Cancer Cell Line Encyclopedia. Nature. Nature
Research; 2019;569:503–8.

47. Choi W, Czerniak B, Ochoa A, Su X, Siefker-Radtke A, Dinney C, et al. Intrinsic basal
and luminal subtypes of muscle-invasive bladder cancer. Nat Rev Urol [Internet]. Nature
Publishing Group; 2014 [cited 2014 Dec 5];11:400–10. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/24960601

48. Bi M, Zhang Z, Jiang YZ, Xue P, Wang H, Lai Z, et al. Enhancer reprogramming driven
by high-order assemblies of transcription factors promotes phenotypic plasticity and breast
cancer endocrine resistance. Nat Cell Biol [Internet]. Springer US; 2020;22:701–15. Available
from: http://dx.doi.org/10.1038/s41556-020-0514-z

49. He Y, Wang LL, Wei T, Xiao Y-TT, Sheng H, Su H, et al. FOXA1 overexpression
suppresses interferon signaling and immune response in cancer. J Clin Invest [Internet].
American Society for Clinical Investigation; 2021 [cited 2021 Jun 10];131. Available from:
https://doi.org/10.1172/JCI147025

50. Lee HW, Chung W, Lee H-O, Jeong DE, Jo A, Lim JE, et al. Single-cell RNA sequencing
reveals the tumor microenvironment and facilitates strategic choices to circumvent treatment
failure in a chemorefractory bladder cancer patient. Available from:
https://doi.org/10.1186/s13073-020-00741-6

51. Song B, Park SH, Zhao JC, Fong KW, Li S, Lee Y, et al. Targeting FOXA1-mediated
repression of TGF-β signaling suppresses castration-resistant prostate cancer progression. J
Clin Invest. American Society for Clinical Investigation; 2019;129:569–82.

52. Mansour MR, Abraham BJ, Anders L, Gutierrez A, Durbin AD, Lawton L, et al. An
oncogenic super-enhancer formed through somatic mutation of a noncoding intergenic
element. Science (80-) [Internet]. 2014;346:1373–7. Available from:
http://www.sciencemag.org/content/346/6215/1373.abstract

53. Chapuy B, McKeown MRR, Lin CYY, Monti S, Roemer MGMGMM, Qi J, et al.
Discovery and Characterization of Super-Enhancer-Associated Dependencies in Diffuse
Large B Cell Lymphoma. Cancer Cell [Internet]. Elsevier Inc.; 2013 [cited 2014 Jan
13];24:777–90. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24332044

54. Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, et al. Master

transcription factors and mediator establish super-enhancers at key cell identity genes. Cell

[Internet]. Cell; 2013 [cited 2021 Jan 25];153:307–19. Available from:

https://pubmed.ncbi.nlm.nih.gov/23582322/

55. Lerdrup M, Johansen JV, Agrawal-Singh S, Hansen K. An interactive environment for

agile analysis and visualization of ChIP-sequencing data. Nat Struct Mol Biol [Internet].

Nature Publishing Group; 2016 [cited 2021 Sep 3];23:349–57. Available from:

https://www.nature.com/articles/nsmb.3180

56. McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, et al. GREAT

improves functional interpretation of cis-regulatory regions. Nat Biotechnol [Internet]. Nat

Biotechnol; 2010 [cited 2021 Jan 25];28:495–501. Available from:

https://pubmed.ncbi.nlm.nih.gov/20436461/

57. Elsworth B, Lyon M, Alexander T, Liu Y, Matthews P, Hallett J, et al. The MRC IEU

OpenGWAS data infrastructure. Available from: https://doi.org/10.1101/2020.08.10.244293

58. Kamat MA, Blackshaw JA, Young R, Surendran P, Burgess S, Danesh J, et al. Databases

and ontologies PhenoScanner V2: an expanded tool for searching human genotype-phenotype

associations. Available from: www.phenoscanner.medschl.cam.ac.uk.

59. Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, et al. COSMIC: The

Catalogue Of Somatic Mutations In Cancer. Nucleic Acids Res [Internet]. Oxford University

Press; 2019 [cited 2021 Dec 10];47:D941–7. Available from: https://cancer.sanger.ac.uk

60. Sondka Z, Bamford S, Cole CG, Ward SA, Dunham I, Forbes SA. The COSMIC Cancer

Gene Census: describing genetic dysfunction across all human cancers [Internet]. Nat. Rev.

Cancer. Nature Publishing Group; 2018 [cited 2021 Dec 10]. p. 696–705. Available from:

https://www.nature.com/articles/s41568-018-0060-1

61. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The

Molecular Signatures Database Hallmark Gene Set Collection. Cell Syst [Internet]. Cell Press;

2015 [cited 2021 Dec 17];1:417–25. Available from:

http://dx.doi.org/10.1016/j.cels.2015.12.004

62. Haghverdi L, Lun ATL, Morgan MD, Marioni JC. Batch effects in single-cell RNA-

sequencing data are corrected by matching mutual nearest neighbors. Nat Biotechnol

[Internet]. Nature Publishing Group; 2018 [cited 2021 Dec 17];36:421–7. Available from:

https://pubmed.ncbi.nlm.nih.gov/29608177/

63. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol [Internet]. BioMed Central Ltd.; 2014 [cited 2021

Sep 16];15:550. Available from:

http://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8

1211

1212

1213

Figures legends

Figure 1: Methodology/Workflow

Figure 2: Chromatin states classify bladder cancers by subgroups
A. ChromHMM principle example and emission order dividing genome in 6 states based on combination of H3K27ac, H3K27me3 and H3K9me3 marks.
B. Expression level association with each chromatin state and each sample.
C. Two chromatin state clusters revealed by unsupervised analysis of top 1% varying regions using MDS for dimension reduction plus MNN for batch effect correction.
D. GSEA functional enrichment analysis of the genes mapped to the MCA Dim2 contributing features. A negative NES indicates significant enrichment in lower Dim2 coordinates (basal direction), and the reverse is in higher Dim2 coordinates (Luminal direction).
E. Luminal versus basal tumour cell signature genes identified with single cell RNA-seq analysis showing concordant enrichment in chromatin state clusters.
F. Genome Browser view of chromatin states at *FOXA1* and *KRT6* loci with corresponding RNAseq (VST normalized scaled expression).

Figure 3: Identification of the bladder super-enhancer repertoire and subtype specificities
A. PCA of H3K27ac signal inside ROSE consensus SE (n=2887) for all samples.
B. Representative examples of H3K27ac signal in Ba/Sq, Stroma-Rich, Luminal and NMIBC tumors with corresponding RNA-seq gene expression. Orange boxes represent SE localisation.
C. Fold Change plots for differentially bound SEs between Ba/Sq and Luminals samples. Significance by p-value <0.05
D. Plot comparing expression LogFC between Ba/Sq and Luminal samples for genes assigned to subgroup-enriched SE.
E. Reactome pathway enrichment analysis of genes associated with Ba/Sq vs Luminal SEs.
F. Venn diagram comparing 3 differential analyses of SE. NHU-enriched SEs are enriched in NHU vs Luminal or NHU vs Basal (pink circle). Basal-enriched SEs are enriched in Basal vs NHU or Basal vs Luminal (red circle). Luminal-enriched SEs are enriched in Luminal vs NHU or Luminal vs Basal (green circle).

Figure 4: Super-enhancers regulate a network of candidate master transcription factors for bladder cancer subgroups
A. Cistrome analysis of LumP and Ba/Sq specific SE.
B. Homer motif enrichment analysis in H3K27ac differential peaks inside differential SE in Luminal vs Basal and Basal vs Luminal.
C. Methodology to identify key coregulated SE-associated TFs
D. Heatmap of the top 75 TFs with high regulon score. Clustering identified 3 major clusters.
E. Heatmap of the top 75 TFs expression in TCGA-BLCA.
F. Correlation network of the top 55 TFs with an expression correlation coefficient of min 0.5 in TCGA-BLCA cohort

G. Single cell RNA-seq analysis of one Bladder Cancer tumors with both Basal and Luminal
Population (GSM4307111) Right panel, associated expression for key TF in each com-
partment.

Figure 5: FOXA1 binds subgroup-specific bladder super-enhancers and correlates with
their activation
A. Venn diagram comparing FOXA1 ChIP-seq peaks in SD48 and 5637 cell line. Heatmap
for 3 categories of peaks and associated mean profiles.
B. Cistrome analysis of motif enrichment analysis in SD48-specific and 5637-specific
FOXA1 peaks
C. Reactome pathway analysis of genes associated to the 3 categories of FOXA1 peaks.
D. Pie chart showing proportion of SE with an overlapping FOXA1 peak (merge of FOXA1
peaks in SD48 and 5637)
E. Correlation between H3K27ac peaks versus FOXA1 peaks inside SE.
F. Heatmap of FOXA1 and H3K27ac reads on FOXA1 peaks overlapping Super-enhancers
ranked by FOXA1 reads ration in SD48 vs 5637
G. Genome browser view of GATA3, PPARg and TGFB2 associated SE. SE are highlighted
with orange boxes.
1273

Figure 6: FOXA1 regulates inflammation and cellular identity
A. Cell viability in RT112 and SCaBER under siRNA treatment against FOXA1
B. Venn diagram comparing differentially expressed genes in RT112 and SCaBER FOXA1
KD.
C. GSEA plot of Msig Hallmark GSEA Analysis of genes differentially regulated in RT112
and SCaBER cell lines upon FOXA1 siRNA (2 independent siRNA, 2 replicates)
D. Heatmap of genes in Hallmark interferon gamma response genes that are differentially
regulated in FOXA1 KD *vs* Ct (min Fold Change = 1,5).
E. Heatmap of Top Luminal TFs expression in RT112 and SCaBER cell lines upon FOXA1
KD.
F. PCA projection of TCGA tumors and CRispR mutant clones on the Basal/Luminal signa-
tures.
G. GSVA analysis of FOXA1 CRispR mutant clones on Urothelial differentiation signature
from Eriksson et al.
H. GSVA analysis of FOXA1 CRispR mutant clones on Basal TFs identified in Fig. 4F
I. Overrepresentation analysis of DEG in FOXA1 mutant *vs* Controls.
J. Volcano plot of Deseq2 RNA-seq analysis comparing pooled CRispR mutant FOXA1
clones in SD48 and RT112 *versus* controls.
1292

1293 **Figure 7: ZBED2, a novel Basal-associated TF also involved inflammation dampening**
1294
A. TCGA expression of ZBED2 by Subtypes
B. TCGA expression Heatmap of ZBED2 and FOXA1 and TCGA correlation between
ZBED2 and FOXA1

C. Expression of FOXA1 and ZBED2 in single-cell transcriptomics from bladder cancer cell
lines in the Cancer Cell Line Encyclopedia (CCLE), highlighting the nearly mutually ex-
clusive expression of these genes.
D. Genome browser view of ZBED2 and FOXA1 loci in SD48 and 5637 cell lines
E. GSEA analysis (Hallmarlk) of ZBED2 correated genes in basal cells population of
GSM4307111 scRNAseq tumor.
F. GSEA analysis (Hallmarlk) of gene expression upon siZBED2 KD in RT112 (siZBED2-1
and siZBED2-2)
G. 3'seq STAT2 and CD274 (PD-L1) expression in RT112 and SCaBER after siZBED2 and
siFOXA1.

1308

Figure 1

ChIPseq + RNAseq
H3K27ac
H3K9me3
H3K27me3

Data integration: **BLCA : Epigenetic Map**

Feature extraction: **Super Enhancers repertoire**

→ SE regulated potential target
→ Corregulated network
→ Master Regulators

Functional validation

Figure 2

Figure 3

Figure 4: Specific regulators of each subgroup specific SE + CRC

Figure 5

Figure 6

Figure 7