Machine learning for exploring microbial inter-kingdom associations in

Cystic Fibrosis and Bronchiectasis

L. Cuthbertson*[1], J. Ish-Horowicz*[4], I.C. Felton[2], P. James[1,3], E. Turek[1], M.J. Cox[1],

M.R. Loebinger[1,2], N.J. Simmonds[1,2], S. Filippi [4], M.F. Moffatt**[1], W.O.C.

Cookson**[1,2]

[1]National Heart and Lung Institute Imperial College London, Genomic Medicine,

London, United Kingdom, [2]Royal Brompton & Harefield NHS Trust, London, United

Kingdom, [3]Oxford Nanopore Technologies Ltd, Oxford, United Kingdom,

[4]Department of Mathematics, Imperial College London, London, United Kingdom

* Authors contributed equally to this manuscript

** Joint senior authors

Corresponding author: L. Cuthbertson, National Heart and Lung Institute,

Imperial College London, London SW3 6LY, UK.

E-mail: l.cuthbertson@imperial.ac.uk

Abstract

Background

Cystic fibrosis (CF) and non-CF bronchiectasis (BX) are lung diseases

characterised by severe chronic infections. Fungal and bacterial components of

infection are both recognized. Recent molecular investigation of sputum from

patients with CF and BX has revealed a complex mycobiome. However, little is

known about how fungal and bacterial organisms interact or whether the

interactions impact on disease outcomes.

Methods

Quantitative PCR and next generation sequencing of ITS2 and 16S rRNA gene was

carried out on 107 patients with CF and BX and defined clinical fungal infection

status. Fungal and bacterial communities were explored using supervised and

unsupervised machine learning to understand associations between fungal and

bacterial communities and their relationship to disease.

Results

Fungal and bacterial communities both had significantly higher biomass and

lower diversity in CF compared to BX patients. Random forest modelling

demonstrated that the fungal and bacterial communities were distinct between CF

and BX patients. Within the CF group, bacterial communities contained no

predictive signal for fungal disease status. Neither bacterial nor fungal community

composition were predictive of the presence of CF pulmonary exacerbation

(CFPE). Intra-kingdom correlations were far stronger than those between the two

kingdoms. Dirichlet mixture components analysis identified two distinct clusters

of bacteria related to the relative abundance of *Pseudomonas*. Fungal community

composition contained no predictive signal for bacterial clusters.

Conclusions

Clear changes in diversity were observed between patients with different clinical

disease status. Although our results demonstrate that bacterial community

composition differs in the presence of fungal disease, no direct relationship

between bacterial and fungal OTUs was found.

Keywords: (10 words)

Mycobiome, Microbiome, machine learning, random forest, cystic fibrosis,

bronchiectasis

Introduction

Chronic respiratory infections are the leading causes of morbidity and mortality

for the chronic suppurative lung diseases (CSLD); cystic fibrosis (CF) and non-CF

bronchiectasis (BX) [1, 2]. Bacterial infections are a major pathophysiological

factor in disease progression in these patients [3, 4]. The impact of fungal

infections is, however, increasingly being recognised. Fungal infections have been

associated with higher disease burdens, increased exacerbation rates and

accelerated clinical decline [5, 6].

The development of robust microbial sequencing protocols has revealed complex

fungal communities within the lungs of patients with CF and BX, exhibiting a range

of clinical disease manifestations [7, 8]. The recognition of these fungal

communities has led to the investigation of bacterial and fungal interactions and

their roles in disease progression.

Insights into these complex associations and interactions are helped through an

evolving statistical methodology. A recent publication by Soret *et al.*, used adapted

Phy-Lasso models to explore the inter-kingdom relationships associated with CF pulmonary exacerbations (CFPE)[8]. The study suggested that fungal genera previously linked to decreasing lung function and poor clinical outcomes were also associated with exacerbations [9, 10].

Machine learning techniques, such as those used by Soret *et al.*, have proven to be effective for host phenotype prediction from microbiome profiles [8]. Studies by Zhou *et al.* and Topçuoğlu *et al.* have reported that random forest modelling achieved the best predictive performance for clinical phenotypes on a range of human gut 16S rRNA gene sequencing datasets, beating penalised linear models and support vector machines [11, 12].

Random forests (RFs) are popular in bioinformatics as they are non-parametric, non-linear and are able to achieve strong predictive performance on many types of data [13]. Furthermore, they often offer improved predictive performance relative to linear models due to their ability to model interactions between variables. In addition, random forests offer a degree of interpretability via variable importance measures such as Mean Decrease Gini and Mean Decrease Accuracy. This has resulted in random forests being used for association testing in genome-wide association studies [14] and biomarker identification in gene microarray studies [15].

In the field of microbiome studies random forests have been used to identify important taxa using both 16S rRNA gene (bacterial) [16, 17] and ITS2 (fungal) sequence data [18, 19]. Recognising the microbial predictors of disease status in chronic respiratory disease may prevent disease progression and allow the development of informed treatment regimens improving patient outcomes.

Therefore, in this present study we explored the bacterial and fungal communities

in CSLD. Using random forest modelling we aimed to explore the power of

bacterial and fungal taxa to predict the presence of CF (n=83) or BX (n=24). Within

subjects with CF, we also sought predictors of clinical diagnoses of fungal disease

(fungal bronchitis, FB) and the presence of CF pulmonary exacerbations (CFPE).

Because of the small number of patients with BX, we confined subgroup analyses

of fungal disease or exacerbations to CF patients only (Table 1).

Table 1. Patient demographics. Data for all samples included in this manuscript,

only samples with paired fungal and bacterial sequencing that passed all quality

control steps (see supplementary information) were taken forward for analysis.

Differences in continuous variables were calculated using a one-way t-test, while

a Chi-square test was used to explore categorical variables.

	BX (n = 24)	**CF** (n = 83)	P
Age/yrs (mean (SD))	59.79 (11.06)	32.00 (11.90)	<0.001
Male (%)	10 (41.7)	47 (56.6)	0.288
BMI (mean (SD))	26.51 (8.73)	22.51 (6.50)	0.029
FEV$_1$ % predicted (mean (SD))	58.24 (25.23)	48.81 (17.79)	0.047
Disease Group (%)			<0.001
ABPA	8 (33.3)	14 (16.9)	
CNPA	5 (20.8)	0 (0.0)	
FB	0 (0.0)	20 (24.1)	
NAFD	8 (33.3)	39 (47.0)	
NTM	3 (12.5)	10 (12.0)	
Exacerbation* (%)	1 (4.2)	36 (43.4)	0.001

CF: Cystic fibrosis, BX: Non-CF bronchiectasis, BMI: Body mass index, FEV$_1$: Forced

expiratory volume in 1 second, ABPA: Allergic bronchopulmonary aspergillosis,

CNPA: Chronic necrotising pulmonary aspergillosis, FB: Fungal bronchitis, NAFD:

No active fungal disease, NTM: Non-tuberculosis mycobacteria.

* Patients were defined as exacerbating by treating clinician as a change in clinical

status from baseline that results in a new pulmonary antibiotic treatment.

Methods

Study Design

A prospective, cross-sectional study of spontaneously expectorated sputum from

adults with CF and BX was carried out at the Royal Brompton NHS trust between

April 2013 and July 2014 [7]. Ethical approval was obtained from the Royal

Brompton and Harefield Hospital Biomedical Research Unit ethics committee

(Advanced Lung Disease Biobank study number: 10/H0504/9).

Patient details, including treatments were collected and subjects were partitioned

into defined clinical subgroups according to the diagnostic criteria defined in

Cuthbertson *et al*. 2020 [7](Table 1).

Sputum samples were collected and processed as previously described with half

the sample sent for routine clinical microbiological culture and the other half

stored at -80 °C for DNA analysis [7]. DNA extraction was performed using the

DNA fast spin kit for soil (MPBio, California, USA) according to the manufacturer's

instructions. Extraction controls were blinded and processed along with patient

samples [7].

Quantitative PCR

Bacterial biomass was quantified by SYBR green quantitative PCR (qPCR) [20].

Fungal biomass was estimated using a modified Taqman based qPCR assay as

previously described [7]. All qPCR reactions were performed in triplicate.

DNA sequencing

16S rRNA gene and ITS2 sequencing were performed on the Illumina MiSeq

platform using dual barcode fusion primers. Bacterial sequencing was performed

on the V4 region of the 16S rRNA gene as previously described [20]. ITS2

sequencing was performed using the primers, ITS2F (5'-CAR CAA YGG ATC TCT

TGG-3') and ITS2R (5'-GAT ATG CTT AAG TTC AGC GGG T-3') with ligated adaptors

[20]. Extraction controls, PCR negative and mock communities were included on

all sequencing runs.

Sequence processing of 16S rRNA gene and ITS2 data was carried out using QIIME

1.9 as described previously in Cuthbertson *et al.* 2017 [20] and Cuthbertson *et al.*

2020 [7] respectively.

All sequences were submitted to the European nucleotide database. Bacterial data

can be accessed under project number PRJEB33064 and the fungal data is

accessible under project number PRJEB33434.

Statistical analysis

Statistical analysis was carried out in R version 3.5.1. Data was analysed in

phyloseq version 1.24.2 [21]. Decontamination of the data was carried out using

decontam version 1.1.2 [22]. Full details are available in Supplementary Materials.

Differences between categorical variables were calculated using Wilcoxon rank

sum test and Kruskal-Wallis. Pearson correlation was used for tests between

continuous variables. Differences in microbial community composition were

tested with PERMANOVA using the Adonis function from vegan version 2.5-6 [23].

Random forests models were fitted using the caret (version 6.0) and ranger

(version 0.12) packages [24, 25]. Differential abundance was performed using the

ALDEx2 (version 1.26) [26], while the Dirichlet Multinomial package (version

1.36) was used for sample-wise clustering [27].

All *P*-values were adjusted for multiple corrections using false discovery rate

(FDR).

Predictive modelling using random forests

Random forests were used to investigate associations between microbial

community composition and the three clinical labels given and described in Table

2. The covariates for these models were the relative abundance of any bacterial

genera or fungal species with prevalence over 20% across all 107 samples. Taxa

accounting for fewer than 0.0001% of reads were also removed, with the

remaining 20 fungal species and 45 bacterial taxa accounting for 99.6% and 96.8%

of bacterial and fungal total reads respectively. Whittaker plots (Figure S1) show

that fungal species and bacterial genus provide similar resolutions with which to

identify microbes. Raw counts were normalised by dividing by the total number

of reads per sample before being transformed using log(x+1) for the random

forest analysis.

Table 2. Phenotypes investigated in this study using random forest. Phenotypes

were chosen to maximise sample sizes and power of the analyses.

The Group phenotype was limited to the FB (n = 20) and NAFD (n = 39) groups

within the CF population. BX samples were removed from the group analysis due

to the low numbers.

Phenotype	Description	Classes	n
Disease	Whether a patient has CF or BX	CF or BX	83 CF, 24 BX (107 total)

Group	Whether a CF patient has been diagnosed with FB or NAFD	FB or NAFD	20 FB, 39 NAFD (59 total)
CFPE	Whether a CF patient was experiencing pulmonary exacerbation when the sample was collected (clinician defined).	Yes or No	36 Yes, 47 No (83 total)

CF: cystic fibrosis, BX: non-cystic fibrosis bronchiectasis, FB: fungal bronchitis,

NAFD: no active fungal disease.

The predictive performance of each RF model was estimated using 5-fold nested

cross-validation with 5 folds in the inner loop. The inner loop performed model

selection using a random hyperparameter search of ten combinations of mtry and

splitrule. Folds were sampled in a stratified manner meaning that the class

proportions in each fold reflected the proportions of the entire dataset. The

predictive performance of the models were evaluated using area under receiver-

operating characteristic curves (AUC) and 95% confidence intervals on the mean

AUC were calculated using the method of LeDell *et al*. [28].

Random forest variable importance and differential abundance analysis

Phenotype-taxa associations were quantified using the debiased Mean Decrease

Gini (MDG) variable importance measure proposed by Nembrini *et al*. [29].

Uncorrected MDG scores are known to be artificially inflated in many settings and

this increases the likelihood of false positive associations [30]. Each random forest

was grown using 5,000 trees to achieve stability in the variable importance scores

[31, 32]. The statistical significance of each score was assessed using *P*-values

calculated using the permutation method of Altmann *et al.* with 5,000

permutations [33]. Random forest variable importance scores are unsigned (they

do not give an effect direction) although an approximate effect direction, using

partial dependence plots which visualise the marginal effect of a variable on the

predictions of a model, can be obtained [34].

ALDEx2 effect sizes and Welch's t-tests were used to complement the random

forest scores for each taxa having differential abundance. Unlike random forest

ALDEx2 explicitly models the compositional nature of sequencing data [35].

Combining multiple computational tools with different underlying assumptions

has been shown to lead to more accurate conclusions in various biological

contexts by reducing the chance of findings being due to statistical or technical

artefacts [36, 37].

Correlation analysis and sample-wise clustering

The correlation structure amongst the taxa included in the RF analysis was also

analysed. Samples were Centre Log Ratio (CLR)-transformed prior to calculating

correlations to account for the compositional nature of 16S rRNA gene and ITS2

data. The CLR transformation was applied separately to the 16S rRNA gene and

ITS2 samples.

Results

Description of data

After decontamination and removal of samples with less than 2,000 reads (n = 2),

107 samples were retained for downstream analyses (for demographics see Table

1). For analyses based on relative abundances, bacterial reads were rarefied to

2,357 while fungal reads were rarefied to 2,542.

Ecological analysis

Differences between diseases

Both bacterial and fungal diversity were significantly higher in patients with BX

than CF (Bacteria; richness, W = 1,507, effect size = 0.369, P < 0.001, Shannon, W

= 1,545, effect size = 0.396, P < 0.001, Simpson, W = 1,549, effect size = 0.399, P <

0.001. Fungi; richness, W = 1,603, effect size = 0.438, P < 0.001, Shannon, W =

1,522, effect size = 0.379, P < 0.001, Simpson, W = 1,465, effect size = 0.338, P <

0.001). No significant difference in dominance however was observed in either

bacterial or fungal communities between diseases (Bacteria; W = 786, effect size

= 0.151, P = 0.118. Fungi; W = 781, effect size = 0.155, P = 0.108). Although

bacterial biomass was significantly lower in the CF group compared to BX (W =

1,100, effect size = 0.241, P = 0.018), no significant difference in fungal biomass

was observed (W = 706, effect size = 0.112, P = 0.126). Differences in both bacterial

and fungal community composition, calculated by PERMANOVA, showed

significant differences between CF and BX (bacterial, R^2 = 0.066, P <0.001; fungal

R^2 = 0.028, P = 0.004).

Differences between clinical fungal disease groups

Participants within the study cohort were classified into four clinically defined

fungal disease groups (see Table 2): (i) fungal bronchitis (FB), (ii) Allergic

bronchopulmonary aspergillosis (ABPA), (iii) chronic necrotising pulmonary

aspergillosis (CNPA, BX only and (iv) non-tuberculosis mycobacteria (NTM). NTM

patients were included as a subgroup due to the strong link between NTM

infection and fungal disease [7, 38, 39]. A control group of patients presenting with

no active fungal disease (NAFD) were also included. Due to the small number of

patients with BX in the study (n = 24) the analysis was confined to the CF patient

samples.

We observed significant differences in fungal alpha diversity between the 39 CF

patients with no active fungal disease (NAFD) and the 20 with fungal bronchitis

(FB) (biomass: W = 522, effect size = 0.275, P = 0.03, richness; W = 211, effect size

= 0.374, P < 0.004; Shannon: W = 204, effect size = 0.387, p = 0.002; Simpson: W =

213, effect size = 0.368, P = 0.004).

Fungal beta diversity using Bray-Curtis similarity was also significantly different

between groups (PERMANOVA R^2 = 0.077, P = 0.004). We found no significant

differences in bacterial biomass or diversity between fungal disease groups.

CFPE

Thirty-six of the 83 CF patients in the study were defined as experiencing CFPE at

the time of sampling. There was no significant difference in bacterial or fungal

biomass or alpha diversity measures (P > 0.1) between CFPE subjects and the rest.

Random forest and differential abundance analysis

We investigated differences in fungal and bacterial community composition

between groups of patients using random forest modelling. Models with mean

AUC >0.7 that were trained on a single kingdom were then used for association

testing. The strength of association was measured using de-biased random forest

variable importance scores while statistical significance was assessed using the

method of Altmann *et al*. (see Methods). Effect directions for the RF variable

importance scores were assigned using partial dependence plots (Figure S2). We

note that RF variable importance scores cannot be compared across models and

so are best used for ranking taxa from the same model.

Disease status

Fungal and bacterial community composition were both good predictors of

Disease status (CF or BX, bacterial mean AUC=0.78, fungal mean AUC=0.74, Figure

1a, left). Including both kingdoms as predictors in the model increased the mean

AUC to 0.79. The confidence intervals on the mean AUC however overlap for all

three models, so we cannot conclude that any one set of covariates were

significantly better predictors of Disease status.

The Gram-negative bacteria genus *Neisseria* was the highest ranked bacterial taxa

by the RF model predicting Disease status from bacterial genera (Figure 1b, left)

and was statistically significantly associated with Disease status at 1% FDR.

Partial dependence plots show that a higher relative abundance of *Neisseria*

implies a higher probability of BX. Other highly ranked bacterial genera included

Megasphaera, Lactobacillus and *Treponema* (direction BX) as well as *Pseudomonas*

(direction CF).

Pseudomonas was highly ranked by the RF model and was also the most

differentially abundant bacterial taxa according to ALDEx2, with a higher

abundance being significantly associated with CF at 1% FDR (effect size=2.5,

Figure 1c, left panel) Fusobacterium, Gemella and *Leptrotrichia* were the bacterial

taxa for which a higher abundance was significantly associated with BX by ALDEx2

at 10% FDR.

Three *Penicillium* species, - *Penicillium thomii*, *Penicillium commune* and

Penicillium psychrosexualis - were highly ranked by the RF models and

significantly associated with disease status by the RF model at 5% FDR (direction

BX), as was *Malassezia restricta. Candida parapsilosis* was the only one of the five

top-ranked fungal taxa where an increased relative abundance increased the

probability of CF according to the RF model. In addition, it was significantly

differentially abundant according to ALDEx2 at 10% FDR (Figure 1c, centre panel)

The pathogenic *Exophiala dermatitidis* was also identified by ALDEx2 but was not

highly ranked by the corresponding RF model.

Figure 1. (a) Bacterial genera (BG), fungal species (FS) and a combination of both are all equally good predictors of CF/BX disease status (left panel), while fungal species are the best predictors of fungal bronchitis (centre panel). Neither fungal species nor bacterial genera are good predictors of CFPE (right panel). Individual fold AUCs are available in Figure S3. (b) Top five taxa by RF (unsigned) importance scores. Effect directions are taken from partial dependence plots (Figure S3) with NA indicating that the partial dependence is non-monotonic. Scores marked with **/*** are statistically significant at 5%/1% FDR. (c) ALDEx2 volcano plot showing effect size magnitudes and FDR-corrected *P*-values from Welch's t-test. NS: not statistically significant.

Clinically defined fungal disease

The random forest models were unable to predict clinically defined fungal disease from bacterial community composition, as an AUC of 0.5 was within the 95% confidence interval on the mean AUC (Figure 1a, centre panel). As expected, fungal community composition was a good predictor of fungal disease status (mean AUC=0.72). In this case, inclusion of both bacterial genera and fungal species in the RF model reduced the power of the model to predict the fungal disease group. RF models highlight three previously identified fungi as being associated with fungal disease status: *Aspergillus fumigatus* ($P < 0.05$); *Exophiala dermatitidis* ($P < 0.05$) and *Scedosporium boydii* ($P < 0.05$) [7]. *Candida albicans* was the highest ranked fungal taxa and was significantly associated with the absence of fungal disease ($P < 0.05$). *Aspergillus fumigatus, Exophiala dermatitidis* and *Candida albicans* also had the largest ALDEx2 effect sizes, however the associations were not significant at 10% FDR (Figure 1c, left panel).

Exacerbations

Neither the bacterial nor fungal taxa contained strong predictive signal for CFPE (Figure 1a, right panel). RF models predicting CFPE from bacterial genera had a mean AUC<0.5, indicating that the random forest model is exhibiting worse-than-random performance on unseen data. This may be a result of stratification bias in a small sample, low-signal dataset (the presence of different proportions of the two classes in different cross-validation folds) [40]. The poor predictive performance of these models meant that no differential abundance analysis was performed for CFPE.

Unsupervised learning/clustering analysis and clustering

Bacterial and fungal communities are structured, with strong positive and negative correlations between the abundance of different organisms. Correlation analysis found large blocks of co-correlated taxa from the same kingdom with few strong correlations between taxa in different kingdoms (Figure 2a and 2b).

Amongst bacterial genera, the largest block of dependent taxa included *Steptococcus*, *Veillonella*, *Rothia*, *Selemonas*, *Prevotella*, *Fusobacterium* and *Atopobium*.

Two clear groups of fungal taxa were observed. One contained four Penicilluim species (*psychrosexualis*, *thomii*, *commune* and *brevicompactum*) and the second contained the pathogenic species *Aspergillus flavus*, *Aspergillus subversicolor*, *Filobasidiella depauperata*, *Talaromyces marneffei* and *Trichosporon dematis*.

There were fewer significant inter-kingdom correlations than intra-kingdom correlations for all FDR thresholds. This further indicates that the main dependencies in this dataset exist within kingdoms rather than between them (Figure 2c).

Figure 2: (a) Pearson correlation between the taxa included in the RF analysis (those with prevalence over 20%). The bacterial and fungal abundances were separately centred log-ratio transformed prior to calculating correlations. (b): The same correlations visualised as a dependence network. Correlations with magnitude less than 0.5 are not shown and all displayed correlations are positive(c): There are a higher percentage of significant correlations within the bacterial and fungal communities (red and green lines) than between them (blue line) for any FDR threshold alpha.

Community structure within the bacterial and fungal kingdoms across samples

was further analysed with Dirichlet mixture components [41]. The latter groups

samples into distinct clusters with similar composition. In this context two

samples are similar if they are well-described by the same Dirichlet multinomial

distribution. This unsupervised approach can provide insight into the community-

level structure that may exist across all the samples and may or may not

correspond to pre-defined clinical labels. This complements the supervised RF

analysis in the previous section, which identified taxa that differentiated between

specified groups.

All 107 samples were clustered using Dirichlet Multinomial Mixture models. This

was done separately using the fungal species and bacterial genera, giving two sets

of cluster labels. Using information-theoretic goodness of fit measures (Figure S4)

2 distinct bacterial clusters (Figure 3a) and a single fungal cluster were identified.

Random forest variable importance scores and ALDEx2 effect sizes show that the

two bacterial clusters are characterised by the presence or absence of

Pseudomonas domination (Figure 3c). The two bacterial clusters did not

correspond to any of the clinical labels in the dataset (Table 3 and Figure 3d). The

dependence between the fungal and bacterial communities was then further

investigated using RF modelling with the bacterial cluster as the class label. Fungal

community composition contained low levels of predictive signal for bacterial

cluster (mean AUC=0.56, Figure 3b), indicating that this important feature of

bacterial community composition (*Pseudomonas* domination) is largely

independent of the fungal community.

Figure 3: (a) Clustering of samples based on bacterial genera abundances identifies two clusters that are separable in Bray-Curtis Principle co-ordinates analysis (PCoA) space. The clusters do not correspond to clinical labels (see Table 3). Clustering of samples based on fungal species abundance did not identify any distinct clusters. (b) Fungal species are poor predictors of bacterial cluster, suggesting limited interactions between the kingdoms in this dataset. Error bars denote 95% confidence interval on mean validation area under the curve (AUC). (c) Random Forest (RF) modelling and ALDEx2 differential expression analysis show that *Pseudomonas* abundance differentiates the two clusters, with Cluster 1 containing low-*Pseudomonas* samples and Cluster 2 containing high-*Pseudomonas* samples. Negative RF variable importance scores are set to zero. (d) The bacterial composition of the samples when ordered by cluster clearly shows that they correspond to presence or absence of domination by *Pseudomonas* species.

Table 3. Adjusted Rand Index (ARI) between bacterial cluster and clinical labels show that there is no similarity between the bacterial cluster labels and other clinical labels used in this study**.**

	Disease	Group	CFPE
ARI(bacterial cluster, clinical label)	-0.01	-0.02	0.02

Discussion

Identifying factors predicting the prevalence and severity of chronic respiratory infections may be crucial for improving clinical outcomes in CSLDs [1, 2]. To date most research has focused on bacterial pathogens., Clinical evidence however suggests that fungal infection plays a key role in chronic disease progression [5, 6]. As such, understanding the inter-kingdom association present within the lungs is an essential step towards effective antimicrobial treatments.

Despite many similarities in the symptoms and treatments of the underlying diseases (CF and BX), there are clearly fundamental differences in their microbial communities. These are likely to be driven by the physiological differences underlying the individual diseases [42].

To understand the microbial drivers that differ between CF and BX, we used random forest modelling to explore the power of bacterial and fungal communities to distinguish between CF and BX patients. In addition, we used equivalent analyses to investigate differences between CF patients with and without clinically defined fungal disease, and indicators of those experiencing CFPE at the time of sampling.

Using RF modelling, bacterial and fungal communities were both independently

found to be good predictors of underlying disease status. Including both fungal

and bacterial OTUs in the model did not however significantly improve the

predictive power, suggesting independence between the fungal and bacterial

communities. We then used the models to identify taxa associated with either CF

or BX. We found that the variable importance scores for fungal covariates were

more concentrated in a small number of highly discriminative taxa (*Penicillium*

thomii, *Penicillium commune*, *Penicillium psychrosexualis*, *Malassezia restricta* and

Candida parapsilosis). In contrast, RF models identified *Neisseria* as being highly

discriminative (direction BX) followed by 18 taxa with similar importance scores

(Figures S5(a) and S6). This point is further demonstrated by the fact that RF

models predicting Disease status identified 4 fungal species with P < 0.05, while

corresponding models identified 19 bacterial genera with P < 0.05 (Figure S5(b)).

RF models revealed *Neisseria* to have the highest variable importance scores and

to be associated with BX. Members of the *Neisseria* genus are commonly isolated

in the upper respiratory tract with some species being known pathogens [43]. Our

results may suggest that a pathogenic role for *Neisseria* spp. could be considered

for BX. This warrants further investigation.

Both RF modelling and ALDEx2 identified the *Pseudomonas* genus as significantly

associated with CF patients. This is consistent with the known role of

Pseudomonas aeruginosa as one of the most common pathogens associated with

CF lung disease. In non-CF bronchiectasis *Pseudomonas* infection may be

associated with more severe disease [44], but it was not a prominent factor in our

BX patients.

Fungal species associated with BX were primarily part of the *Penicillium* genus.

Symptomatic infections with *Penicillium* spp. are rare [45] and *Penicillum* spp. are

widely present in the air, making it a logical part of the normal respiratory flora.

Conversely *Exophaila dermatitidis* and *Candida parapsilosis*, which were identified

as significantly associated with CF using ALDEx2, have been shown to be

significant pathogens in CF. RF models identified *E. dermatitidis* alongside

Aspergillus fumigatus and *Scedosporium boydii*, as one of the major pathogens in

FB, while *Candida parapsilosis* was shown to be associated with CF patients with

NAFD. Despite the association with the NAFD group, *C. parapsilosis* has been

identified previously to be an opportunistic human pathogen [46].

We also explored differences in the microbiota between patients with and without

a clinical fungal infection. The relatively low numbers of patients with BX led us to

restrict analysis of fungal disease diagnoses to differences between FB and NAFD

in the CF patients. We did not find any predictive signal in the bacterial community

for the presence of FB. This provides further evidence for a lack of inter-kingdom

interactions driving fungal disease. The fungal pathogens *Aspergillus fumigatus,*

Exophiala dermatitidis and *Scedosporium boydii* were however associated with FB

by RF models, with *Aspergillus fumigatus* and *Exophiala dermatitidis* showing the

strongest effects by both the RF model and ALDEx2. Although the RF associations

were significant ($P < 0.05$) those from ALDEx2 were not ($P > 0.1$).

The most important taxon associated with the NAFD group in the CF cohort was

Candida albicans. *C. albicans* is a common commensal organism, however, in

immunocompromised patients these organisms have been shown to be

opportunistic pathogens. In CF adults *Candida spp.* colonization has been shown

to be associated with use of inhaled steroids, diabetes mellitus and antibiotic

treatment. Despite these observations the virulence potential of *C. albicans* in CF

is still being explored[47]. In the present study *Candida spp.* were present with a

lower relative abundance in the FB group suggesting that dominance of

filamentous fungi may out compete *Candida spp.* in these patients. More work is

therefore needed to understand the role *C. albicans* plays in CF disease

progression.

Pulmonary exacerbations are major clinical events in patients with CF resulting in

lung function decline and clinical disease progression [48]. The presence of

bacteria and viruses is commonly associated with these poor outcomes during

CFPE [8], but defining their exact role is challenging. Recent evidence has

suggested fungal infections are also associated with increased CFPE although to

date few studies have explored this area [49, 50]. A recent publication by Soret *et*

al. investigated CFPE using an adapted penalised linear model and cross-sectional

data and suggested that two fungal genera, *Aspergillus* and *Malassezia* were

associated with CFPE [8]. In our study however random forest models were not

able to distinguish between patients with and without CFPE. This may be because

the cross-sectional data in our study is unable to show intra-patient variation

within the microbial community. Longitudinal data is likely to be more

informative for predicting pulmonary exacerbations and should be considered for

future studies.

We used an unsupervised clustering analysis to detect interactions between

different genera, identifying two bacterial clusters which did not correspond

simply to the CF/BX groups. The bacterial clusters were however clearly driven

by dominance of *Pseudomonas* within individual samples (Figure 2c). This is in

agreement with the RF analysis, which found that a higher relative abundance of

Pseudomonas was associated with increased probability of CF. This effect was

weak with partial dependence plots (Figure S2) showing a small increase in

probability of CF when the relative abundance of *Pseudomonas* was >70%.

Our RF models were unable to predict bacterial clusters from fungal species,

providing additional evidence of independence between the bacterial and fungal

communities. Inter-kingdom correlations were generally weaker than those

within either kingdom (measured by proportion of significant correlations at

different significance thresholds). This further indicates that intra-kingdom

interactions may play a minor role in these subjects.

By utilising distinct statistical methods (RF variable importance scores and

ALDEx2 effect sizes) we were able to obtain a consensus result that is more robust

than results from either individual tool, as applying two statistical approaches

with distinct assumptions reduces the risk that findings are due to technical or

statistical artefacts [51]. ALDEx2 explicitly models the compositional nature of

sequencing data, while RFs do not make assumptions about the characteristics of

the covariates [35]. There was statistically significant agreement between the RF

variable importance scores and ALDEx2 effect sizes for all the models used for

differential abundance analysis (Figure S7).

Our analysis does though have several limitations that should be considered when

interpreting the results presented. Most importantly, associations identified by

machine learning models such as random forest do not imply causal links.

Furthermore, the importance scores from random forests should be interpreted

with care. Using debiased Gini importance scores and assessing statistical

significance via permutation reduces the danger of spurious associations but does

not provide a framework that allows quantitative statements to be made (as is

possible with effect sizes in linear models). Machine learning is a powerful tool for

exploring microbial interactions and drivers of disease, however, understanding

the limitations of the models is vital for interpretation. We have consequently

utilised multiple statistical tools to reduce the dangers of over-interpreting the

results of one approach.

A further limitation of this study is the use of 16S rRNA gene sequencing and ITS2

sequencing for the exploration of these communities. This technology allows us to

understand the microbial community present within the lung but give us no

information on their activity.

A recent study carried out by Hughes *et al.* established that using culture-based

methodologies, the known CF pathogens, *Pseudomonas aeruginosa* and *Aspergillus*

fumigatus are rarely cultured from the same sample [52]. Despite this, our culture

independent techniques clearly show the common presence of both *Pseudomonas*

aeruginosa and *Aspergillus fumigatus* reads in the samples. It is possible that

microbial interactions within the lungs may not be detected by DNA based

methods. Future work may require functional analyses to explore relative

microbial gene expression within the lung.

Conclusions

Our study suggests that the fungal microbiota play a key role in chronic

respiratory disease independently of bacterial infection. Further work is required

to understand the full impact of fungal infection in chronic respiratory disease.

Improvements in clinical diagnosis of fungal infections, whether by sequence

analysis, transcriptomics or advanced cultures, may underpin the improvement of

patient outcomes.

<u>Declarations</u>

Availability of data and material

Sequencing data is freely assessable through the European nucleotide database,

bacterial data can be accessed under project number PRJEB33064, while the

fungal data can be accessed under project number PRJEB33434.

Data analysis scripts are freely available on figshare, under the project "Machine

learning for exploring microbial inter-kingdom associations in Cystic Fibrosis

and Bronchiectasis" DOIs: https://doi.org/10.6084/m9.figshare.17897708 and

https://doi.org/10.6084/m9.figshare.17942543.

Funding

This project was supported by the Asmarley Trust, the Wellcome Trust and the

NIHR Respiratory Disease Biomedical Research Unit at the Royal Brompton and

Harefield NHS Foundation Trust, Imperial College London. IF was supported by an

NIHR PhD studentship. The funders had no role in study design, data collection

and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

All authors listed provided substantial contributions to this manuscript. Study

concept and design; MFM, WOCC, SF, LC, NJS, MRL, MJC. Acquisition of data; IF, PJ,

LC, MJC. Statistical analysis and interpretation of the data; LC, JIH. Drafting of the

manuscript; LC, JIH, WOCC, SF. Critical revision of the manuscript for was carried

out by all authors.

<u>References</u>

1.   O'Sullivan BP, Freedman SD: **Cystic fibrosis.** *Lancet* 2009, **373:**1891-
1904.
2.   King PT: **The pathophysiology of bronchiectasis.** *Int J Chron Obstruct*
Pulmon Dis 2009, **4:**411-419.
3.   Zemanick ET, Hoffman LR: **Cystic Fibrosis: Microbiology and Host**
Response. *Pediatr Clin North Am* 2016, **63:**617-636.
4.   Maselli DJ, Amalakuhan B, Keyt H, Diaz AA: **Suspecting non-cystic**
fibrosis bronchiectasis: What the busy primary care clinician needs
to know. *Int J Clin Pract* 2017, **71**.
5.   Chotirmall SH, O'Donoghue E, Bennett K, Gunaratnam C, O'Neill SJ,
McElvaney NG: **Sputum Candida albicans presages FEV(1) decline and**
hospital-treated exacerbations in cystic fibrosis. *Chest* 2010,
138:1186-1195.
6.   Amin R, Dupuis A, Aaron SD, Ratjen F: **The effect of chronic infection**
with Aspergillus fumigatus on lung function and hospitalization in
patients with cystic fibrosis. *Chest* 2010, **137:**171-176.
7.   Cuthbertson L, Felton I, James P, Cox MJ, Bilton D, Schelenz S, Loebinger
MR, Cookson WOC, Simmonds NJ, Moffatt MF: **The fungal airway**
microbiome in cystic fibrosis and non-cystic fibrosis bronchiectasis.
Journal of Cystic Fibrosis 2021, **20:**295-302.
8.   Soret P, Vandenborght LE, Francis F, Coron N, Enaud R, Avalos M,
Schaeverbeke T, Berger P, Fayon M, Thiebaut R, et al: **Respiratory**
mycobiome and suggestion of inter-kingdom network during acute
pulmonary exacerbation in cystic fibrosis. *Sci Rep* 2020, **10:**3589.
9.   Reece E, Segurado R, Jackson A, McClean S, Renwick J, Greally P: **Co-**
colonisation with Aspergillus fumigatus and Pseudomonas
aeruginosa is associated with poorer health in cystic fibrosis
patients: an Irish registry analysis. *BMC Pulm Med* 2017, **17:**70.
10.   Parize P, Boussaud V, Poinsignon V, Sitterle E, Botterel F, Lefeuvre S,
Guillemain R, Dannaoui E, Billaud EM: **Clinical outcome of cystic**
fibrosis patients colonized by Scedosporium species following lung
transplantation: A single-center 15-year experience. *Transpl Infect Dis*
2017, **19**.
11.   Zhou YH, Gallins P: **A Review and Tutorial of Machine Learning**
Methods for Microbiome Host Trait Prediction. *Frontiers in Genetics*
2019, **10**.

12. Topcuoglu BD, Lesniak NA, Ruffin MIV, Wiens J, Schlossa PD: **A**
Framework for Effective Application of Machine Learning to
Microbiome-Based Classification Problems. *Mbio* 2020, **11**.
13. Qi Y: *Random forest for bioinformatics. In Ensemble machine learning.*
Springer, Boston, MA.; 2012.
14. Szymczak S, Holzinger E, Dasgupta A, Malley JD, Molloy AM, Mills JL,
Brody LC, Stambolian D, Bailey-Wilson JE: **r2VIM: A new variable**
selection method for random forests in genome-wide association
studies. *Biodata Mining* 2016, **9**.
15. Ram M, Najafi A, Shakeri MT: **Classification and Biomarker Genes**
Selection for Cancer Gene Expression Data Using Random Forest.
Iran J Pathol 2017, **12:**339-347.
16. Roguet A, Eren AM, Newton RJ, McLellan SL: **Fecal source identification**
using random forest. *Microbiome* 2018, **6**.
17. Thompson J, Johansen R, Dunbar J, Munsky B: **Machine learning to**
predict microbial community functions: An analysis of dissolved
organic carbon from litter decomposition. *Plos One* 2019, **14**.
18. Nagpal R, Neth BJ, Wang SH, Mishra SP, Craft S, Yadav H: **Gut mycobiome**
and its interaction with diet, gut bacteria and alzheimer's disease
markers in subjects with mild cognitive impairment: A pilot study.
Ebiomedicine 2020, **59**.
19. Das A, O'Herlihy E, Shanahan F, O'Toole PW, Jeffery IB: **The fecal**
mycobiome in patients with Irritable Bowel Syndrome. *Scientific*
Reports 2021, **11**.
20. Cuthbertson L, Craven V, Bingle L, Cookson W, Everard ML, Moffatt MF:
The impact of persistent bacterial bronchitis on the pulmonary
microbiome of children. *PLoS One* 2017, **12:**e0190075.
21. McMurdie PJ, Holmes S: **phyloseq: an R package for reproducible**
interactive analysis and graphics of microbiome census data. *PLoS*
One 2013, **8:**e61217.
22. Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ: **Simple**
statistical identification and removal of contaminant sequences in
marker-gene and metagenomics data. *Microbiome* 2018, **6:**226.
23. Jari Oksanen FGB, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan
McGlinn, Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos,
M. Henry H. Stevens, Eduard Szoecs, Helene Wagner: **vegan: Community**
Ecology Package. R package version 2.5-6. 2019.
24. Kuhn M: **Building Predictive Models in R Using the caret Package.**
Journal of Statistical Software 2008, **28:**1-26.
25. Wright MN, Ziegler A: **ranger: A Fast Implementation of Random**
Forests for High Dimensional Data in C plus plus and R. *Journal of*
Statistical Software 2017, **77:**1-17.
26. Fernandes AD, Macklaim JM, Linn TG, Reid G, Gloor GB: **ANOVA-Like**
Differential Expression (ALDEx) Analysis for Mixed Population RNA-
Seq. *Plos One* 2013, **8**.
27. Morgan M: **DirichletMultinomial: Dirichlet-Multinomial Mixture**
Model Machine Learning for Microbiome Data. *R package version 1340*
2021.

28.   LeDell E, Petersen M, van der Laan M: **Computationally efficient**
confidence intervals for cross-validated area under the ROC curve
estimates. *Electronic Journal of Statistics* 2015, **9:**1583-1607.
29.   Nembrini S, Konig IR, Wright MN: **The revival of the Gini importance ?**
Bioinformatics 2018, **34:**3711-3718.
30.   Strobl C, Boulesteix AL, Zeileis A, Hothorn T: **Bias in random forest**
variable importance measures: Illustrations, sources and a solution.
Bmc Bioinformatics 2007, **8**.
31.   Wang H, Yang F, Luo Z: **An experimental study of the intrinsic stability**
of random forest variable importance measures. *BMC Bioinformatics*
2016, **17:**60.
32.   Behnamian A, Millard K, Banks SN, White L, Richardson M, Pasher J: **A**
Systematic Approach for Variable Selection With Random Forests:
Achieving Stable Variable Importance Values. *Ieee Geoscience and*
Remote Sensing Letters 2017, **14:**1988-1992.
33.   Altmann A, Tolosi L, Sander O, Lengauer T: **Permutation importance: a**
corrected feature importance measure. *Bioinformatics* 2010, **26:**1340-
1347.
34.   Friedman JH: **Greedy function approximation: A gradient boosting**
machine. *The Annals of Statistics* 2001, **29:**1189:1232.
35.   Quinn TP, Erb I, Gloor G, Notredame C, Richardson MF, Crowley TM: **A**
field guide for the compositional analysis of any-omics data.
Gigascience 2019, **8**.
36.   Marbach D, Costello JC, Kuffner R, Vega NM, Prill RJ, Camacho DM, Allison
KR, Consortium D, Kellis M, Collins JJ, Stolovitzky G: **Wisdom of crowds**
for robust gene network inference. *Nat Methods* 2012, **9:**796-804.
37.   Aghaeepour N, Finak G, Flow CAPC, Consortium D, Hoos H, Mosmann TR,
Brinkman R, Gottardo R, Scheuermann RH: **Critical assessment of**
automated flow cytometry data analysis techniques. *Nat Methods*
2013, **10:**228-238.
38.   Kunst H, Wickremasinghe M, Wells A, Wilson R: **Nontuberculous**
mycobacterial disease and Aspergillus-related lung disease in
bronchiectasis. *Eur Respir J* 2006, **28:**352-357.
39.   Joao I, Bujdakova H, Jordao L: **Opportunist Coinfections by**
Nontuberculous Mycobacteria and Fungi in Immunocompromised
Patients. *Antibiotics (Basel)* 2020, **9**.
40.   Parker BJ, Guenter S, Bedo J: **Stratification bias in low signal**
microarray studies. *Bmc Bioinformatics* 2007, **8**.
41.   Holmes I, Harris K, Quince C: **Dirichlet Multinomial Mixtures:**
Generative Models for Microbial Metagenomics. *Plos One* 2012, **7**.
42.   !!! INVALID CITATION !!! [39].
43.   Weyand NJ: **Neisseria models of infection and persistence in the**
upper respiratory tract. *Pathog Dis* 2017, **75**.
44.   Davies G, Wells AU, Doffman S, Watanabe S, Wilson R: **The effect of**
Pseudomonas aeruginosa on pulmonary function in patients with
bronchiectasis. *Eur Respir J* 2006, **28:**974-979.
45.   Lyratzopoulos G, Ellis M, Nerringer R, Denning DW: **Invasive infection**
due to penicillium species other than P. marneffei. *J Infect* 2002,
45:184-195.

46. Pendleton KM, Huffnagle GB, Dickson RP: **The significance of Candida in**
the human respiratory tract: our evolving understanding. *Pathog Dis*
2017, **75**.
47. Chotirmall SH, Greene CM, McElvaney NG: **Candida species in cystic**
fibrosis: A road less travelled. *Med Mycol* 2010, **48 Suppl 1:**S114-124.
48. de Boer K, Vandemheen KL, Tullis E, Doucette S, Fergusson D, Freitag A,
Paterson N, Jackson M, Lougheed MD, Kumar V, Aaron SD: **Exacerbation**
frequency and clinical outcomes in adult patients with cystic
fibrosis. *Thorax* 2011, **66:**680-685.
49. Nguyen LDN, Deschaght P, Merlin S, Loywick A, Audebert C, Van Daele S,
Viscogliosi E, Vaneechoutte M, Delhaes L: **Effects of Propidium**
Monoazide (PMA) Treatment on Mycobiome and Bacteriome
Analysis of Cystic Fibrosis Airways during Exacerbation. *Plos One*
2016, **11**.
50. Willger SD, Grim SL, Dolben EL, Shipunova A, Hampton TH, Morrison HG,
Filkins LM, Toole GAO, Moulton LA, Ashare A, et al: **Characterization and**
quantification of the fungal microbiome in serial samples from
individuals with cystic fibrosis. *Microbiome* 2014, **2**.
51. Costa-Silva J, Domingues D, Lopes FM: **RNA-Seq differential expression**
analysis: An extended review and a software tool. *Plos One* 2017, **12**.
52. Hughes DA, Rosenthal M, Cuthbertson L, Ramadan N, Felton I, Simmonds
NJ, Loebinger MR, Price H, Armstrong-James D, Elborn JS, et al: **P130**
Pseudomonas aeruginosa impairs growth of aspergillus from CF
airway samples. *Thorax* 2021, **76:**A159.
756