PFTK1 kinase regulates axogenesis during development via RhoA activation.

Yasmilde Rodríguez González[1], Fatemeh Kamkar[1], Paymaan Jafar-nejad[1#,] Suzi Wang[1], Dianbo Qu[1], Leticia
Sanchez Alvarez[1], Dina Hawari[1], Margaret Sonnenfeld[1†], Ruth S. Slack[1], Paul Albert[3], David S. Park[2*] and Alvin
Joselin[2*]

[1] Department of Cellular and Molecular Medicine, University of Ottawa, Ottawa, ON, Canada K1H 8M5,

[2] Hotchkiss Brain Institute, Department of Clinical Neurosciences University of Calgary, Calgary, AB, T2N 4N1

[3] Ottawa Hospital Research Institute and Brain and Mind Research Institute, University of Ottawa, Ottawa, ON,
Canada K1H 8M5

[#] Presently at Department of Core Antisense Research, Ionis Pharmaceuticals, Carlsbad, California 92010

[†] Presently at Faculty of Science, Thompson Rivers University, Kamloops BC V2C 0C8

[*] To whom correspondence should be addressed

Correspondence: Alvin.Joselin1@ucalgary.ca

Running Title: PFTK1 regulates axogenesis through Rho.

Key words: PFTAIRE; Eip63E; Rho; Cyclin dependent kinases; axogenesis; neuronal development.

Abstract

PFTK1/Eip63E is a member of the Cyclin-dependent kinases (CDKs) family and plays an important role in normal
cell cycle progression. Eip63E expresses primarily in postnatal and adult nervous system in Drosophila
melanogaster but its role in CNS development remains unknown. We sought to understand the function of Eip63E
in the CNS by studying the fly ventral nerve cord during development. Our results demonstrate that Eip63E
regulates axogenesis in neurons and its deficiency leads to neuronal defects. Functional interaction studies
performed using the same system identify an interaction between Eip63E and the small GTPase Rho1.
Furthermore, deficiency of Eip63E homolog in mice, PFTK1, in a newly generated PFTK1 knockout mice results
in increased axonal outgrowth confirming that the developmental defects observed in the fly model are due to
defects in axogenesis. Importantly, RhoA phosphorylation and activity is affected by PFTK1 in primary neuronal
cultures. We here report that GDP bound inactive RhoA is a substrate of PFTK1 and PFTK1 phosphorylation
maintains RhoA activity. In conclusion, our work establishes an unreported role of PFTK1 in axon development
mediated by phosphorylation and activation of GDP-bound RhoA. The results presented add to our understanding
of the role of Cdks in the maintenance of RhoA mediated axon growth and its impact on CNS development and
axonal regeneration.

1. **Introduction**

2. Cyclin Dependent Kinase (CDK) are involved in the regulation of several cellular processes including cell cycle

3. regulation, transcription, translation and neuronal differentiation and function (Malumbres and Barbacid, 2005).

4. Several of these kinases and their interaction partners are typically highly expressed and active in neuronal cells.

5. This includes CDK5, CDK16-18 (PCTAIRE 1-3/ PCTK1-3) and CDK14-15 (PFTAIRE/ PFTK1-2). The best

6. characterized of these CDKs, CDK5, is predominantly active in neurons owing to the tissue-specific expression of

7. its activators, p35/25 and p39. CDK5 deficiency in mice leads to perinatal lethality as a result of defective

8. neuronal migration, differentiation and survival. Both PFTK1 and PCTK1 have been shown to be activated by

9. cyclin Y (Jiang et al., 2009; Mikolcevic P, 2012). PFTK1, first identified in *Drosophila melanogaster* (Dmel) (Sauer

10. et al., 1996), and later in mice (Lazzaro and Julien, 1997) and humans (Yang and Chen, 2001), is highly

11. expressed in the Central Nervous System (CNS) (Besset et al., 1998; Lazzaro et al., 1997), predominantly in the

12. brain and likely plays an important role in development since PFTAIRE deficient flies die mostly at early larval

13. stages (Stowers et al., 2000). Notwithstanding, the function of PFTK1 in CNS development and the downstream

14. substrates PFTK1 regulates has remained largely undiscovered.

15. In mammals, PFTK exists as 2 separate genes: *PFTAIRE1/PFTK1/CDK14* (Yang and Chen, 2001) and

16. *PFTAIRE2/PFTK2/Als2cr7/CDK15* (PubMed Gene ID: 65061). PFTK1 is highly conserved among different

17. species from yeast to mammals (Liu and Kipreos, 2000). The fly and worm genome code for the gene known as

18. Eip63-Ecdysone-induced protein 63E which shares 70% homology to mouse PFTK1 (Liu and Kipreos, 2000;

19. Stowers *et al.*, 2000). The human *PFTK1* and mouse *Pftk1* are located at (7q21-q22) and (5 A1; 2.61 cM)

20. respectively (Lazzaro and Julien, 1997) and show 96% homology. *PFTAIRE2/ PFTK2/ Als2cr7/ CDK15* is

21. recognized in GenBank database, and has been reported for its role in conferring resistance to tumor necrosis

22. factor-related apoptosis-inducing ligand (TRAIL) in cancer cells (Park et al., 2014). In the adult mouse, *Pftk1*

23. mRNA is located both in neuroglia and neurons (Besset *et al.*, 1998; Lazzaro *et al.*, 1997) in different regions of

24. the brain such as cortex, hippocampus, thalamus, cerebellum, and in the spine. PFTK1 is localized in neuronal

25. cell bodies but not in neurite extensions. During mouse development, *Pftk1* mRNA is expressed at very low levels

26. at embryonic days 12.5, 15.5, and even 18.5 but soon rises after birth, at P 10.5, and reaches a maximum,

27. comparable to levels in adult mice. This is different from the expression pattern of CDK5 that has a continuous

28. increase from embryonic day 12, followed by a significant increase at birth (Lazzaro *et al.*, 1997). All this

29. evidence, together with the knowledge that PFTK1 and CDK5 share a high amino acid sequence similarity (~ 50-

30. 52%), led us to examine the role of PFTK1 during CNS development.

31. We first examined the role of PFTK1 homolog Eip63E in Dmel ventral nerve cord (VNC) in the regulation of axon

32. and neuronal structure. We show that Eip63E regulates axogenesis in neurons and Eip63E deficiency or

33. downregulation leads to defects in the axon and neuronal structure of the VNC. Furthermore, our functional

34. interaction studies using this model led us to identify Rho1 as important potential mediator of Eip63E function. To

35. further confirm this role of PFTK1 in CNS, we describe the characterization of a *Pftk1* knockout (KO) mice.

36. Although, no gross abnormalities were observed in morphology, fertility, life span or anatomical brain structures of

the PFTK1 KO mice within the parameters tested, perturbations in axonal growth were observed in cultured cortical neurons. Corroborating our observations in the flies, overexpression of a DN- PFTK1 in WT cultured cortical neurons and primary cortical neurons from *Pftk1* KO mice, exhibit faster-growing axons. In line with our observations in the flies, investigations show compelling evidence that PFTK1 phosphorylates GDP-RhoA, and this leads to RhoA activation. These data together point to an important and unexplored role of PFTK1 in axogenesis through its interaction with RhoA implicating PFTK1s role in CNS development.

Results

Eip63E KO embryos show defasciculation and misguidance of axons in VNC

To examine the role of PFTK1 in CNS development, we used the VNC of Dmel embryos, since it provides an

excellent model to study developmental processes (Garbe and Bashaw, 2004; Sanchez-Soriano et al., 2007).

Additionally, flies do not express other neuronal CDKs closely related to PFTK1 that could add a level of

complexity, due to possible compensation. We used two different Eip63E mutant Dmel lines, carrying the

Df(3L)E1 and Eip63E[81] alleles respectively (Stowers *et al.*, 2000). The nature of both these alleles is different in

that the former is a result of a large deletion that spans the transcription initiation region to the entire kinase

domain of Eip63E. The latter is a smaller 48 bp in-frame deletion that removes 16 codons within the conserved

kinase domain (213-228). To distinguish the mutant homozygous embryos from the rest of the genotypes in the

progeny, both mutant lines were balanced with the TM3krGFP,Sb chromosome which expresses GFP under the

control of Krüppell promoter from embryonic stage 9-10 to adulthood (Casso et al., 2000). BP102 or fasciclin-II

(1D4) were both used as different markers to assess the VNC structures of stage 14-15 embryos from the

crosses. Both are expressed in the axon bundles that define the commissural and longitudinal connectives of the

fly VNC (Fambrough et al., 1996; Hummel et al., 1999). Both markers confirmed axonal defects in the Eip63E KO

embryos (Fig 1). KO embryos (GFP negative) show a diffuse appearance of the axonal pathways (Fig. 1A and

1B) suggesting defasciculation of axon bundles. Additionally, axons appear not to follow the appropriate

pathways, leading to malformation of the ladder-like normal axonal structure of the VNC, with disruption of both

commissural and longitudinal pathways (Fig. 1A and 1B).

Since the larger deletion in Df(3L)E1 line is known to affect two other genes upstream of the Eip63E gene, namely

SDE3/armitage (Cook et al., 2004) and 1(3)63D (Stowers *et al.*, 2000), we tested any possible genetic variations

within these lines. We generated trans-heterozygous embryos containing one copy of each of the mutant alleles

and stained for BP102. As shown in Fig. 1A (bottom panel) and its corresponding scoring chart (Fig. 1A, bottom

right), axonal defect comparable to those of the single deficiency were observed. Together with the observation

that no defects were ever observed in GFP-positive embryos, this suggests a strong correlation between Eip63E

deficiency and axonal defects.

Axon outgrowth occurs prematurely in Eip63E KO embryos.

Axogenesis in Dmel embryos starts at ~7:30h into development (early stage 12) after specification of the midline

cells that will guide the pioneer axons contra-laterally to cross the midline (Hummel *et al.*, 1999; Jacobs and

Goodman, 1989). To elucidate how early in CNS development axon defects start to appear, we examined CNS

features using anti-horseradish peroxidase antibody which recognizes a carbohydrate epitope (called HRP for

simplicity) present on several cell-surface molecules expressed by neuronal cell bodies and axons (Snow et al.,

1987) including fasciclin II (Desai et al., 1994). 10% of stage 8-9 embryos mixed population from the Eip63E KO

line (n=48) showed premature HRP staining distinct from WT embryos (Fig. 2C, top 2 panels). Lateral views of the

embryos are presented since this abnormality became more evident when looking laterally. Stage 9-10 WT

embryos show faint, background staining along the ventral region of the embryo where VNC axons would later

appear. In contrast, Eip63E KO embryos show HRP-positive structures at stages 9-10 distinct from WT embryos

(Fig. 2C, middle 2 panels). By late stage 11, 36% of KO embryos (GFP-, n=52) show premature, and in some

cases misguided, axon outgrowth affecting several segments of the VNC (Fig. 2C, bottom 2 panels). These

results show that axon-related defects appear early, from the time axon structures tend to emerge and that the

incidence of axon defects in KO embryos increases during development.

Early neurogenesis events are defective in Eip63E mutants.

An early event in Dmel CNS development is the formation of the ventral midline. This has an essential impact in

axon outgrowth/guidance and the proper ladder-like scaffold of axons of the VNC (Hummel *et al.*, 1999; Klambt et

al., 1991). To assess midline formation in Eip63E mutant lines, we performed immunostaining of single-minded

(sim), a protein which is exclusively expressed in midline precursors and their progeny and is a master regulator

of the differentiation of midline neurons and glial cells (Nambu et al., 1990). Eip63E deficiency does not have any

effect on midline formation, since no defects were found in embryos from the Eip63E[81] line (Stages 5-10)

compared to WT embryos (Fig. 2A). Furthermore, HRP staining at stage 11 (Fig. 2C, bottom panel) suggests that

the ventral midline appears normal in Eip63E KO embryos even when the deficiency might lead to axon defects

and abnormal HRP staining at later stages.

Another hallmark of the Dmel embryo development is the segmentation of the embryo which is triggered early

(within the first 3h) leading to the formation of stripe-like segments along the anterior-posterior axis. Segment

polarity genes are part of the expression cascade that regulate this process and are expressed in 14 stripes at the

onset of gastrulation (~3h, stage 6-7) (Weigmann et al., 2003). These genes, which include engrailed (en) and

invected (inv), are major players in neurogenesis (Bhat, 1999; Bhat and Schedl, 1997). We examined the

expression pattern of en/inv proteins to evaluate if segmentation and/or neuroblasts delamination/segregation

was affected in Eip63E KO embryos. Immunostaining with the 4D9 antibody, that recognizes both en/inv proteins,

reveal that 18% of stage 8-9 embryos from the Eip63E81/TM3krGFP,Sb line show abnormal patterning (Fig. 2B).

Since genotyping is not practical at these stages, it is not possible to link this phenotype to Eip63E deficiency. It

should be noted that comparable defects were observed in similarly timed embryos from the

Df(3L)E1/TM3krGFP,Sb line (data not shown). In comparison, only 3% of WT embryos of similar stages show any

defects. Although abnormalities were also observed in older KO embryos (stages 10), their incidence decreased

(8% Eip63E[81]). This is likely due to the adverse effects of the defect on the survival of defective embryos. Taken

together, these results suggest a potential role for Eip63E in delamination of neuroblasts and/or their segregation.

Eip63E deficiency leads to other morphological defects in the CNS.

To further evaluate the effects of Eip63E deficiency on neuronal cell bodies and glia, we stained for the product of

elav (embryonic lethal abnormal visual system) gene which codes for an RNA-binding protein that is expressed

exclusively in the neurons (Robinow and White, 1991) and is required during the development and maintenance

of the CNS (Jimenez and Campos-Ortega, 1987). ELAV expression pattern in Eip63E KO embryos reveal defects

as early as stage 12 (Fig. 3A). ELAV staining is diffuse in the kO compared to WT embryos of similar stages.

Additionally, the morphology of neuromeres and hemisegments along the VNC of mutant embryos was disrupted

and abnormal, making it difficult to distinguish midline neurons from the rest in several VNC segments of stage

12-13 KO embryos (Fig. 3A).

Lateral views of elav and HRP co-staining (Fig. 3E) reveal that Eip63E deficiency resulted in a disorganized

neuronal cortex, since elav staining was detected dorsally underneath the axon scaffold. Similar abnormalities

were observed using a different neuronal marker, Neuroglian (long isoform, BP-104), a cell-adhesion molecule

expressed in a sub-population of neuronal cell bodies and axons (Hortsch et al., 1990). Malformation of

neuromeres and hemisegments linked to axon pathfinding defects are observed in Eip63E KO embryos (Fig. 3B).

The lack of symmetry between the hemisegments and the midline pattern of different neuromeres suggest a

disorganization of neuronal cell bodies consistent with the observations for elav immunostaining. Disarray of

neurons and axons was also observed when a VNC flat preparation of timed stage 11 embryos was co-stained for

Futsch and anti-HRP (Fig. 3F). Futsch is a microtubule-associated protein, necessary for axon and dendrite

development in Drosophila (Hummel et al., 2000). The axonal bundles in Eip63E KO embryos appear

defasciculated when compared with controls (HRP staining) and defective hemisegments coincide with misplaced

neurons (Futsch staining). Additionally, as can be seen with Futsch/HRP merged images, neuronal cell bodies are

abnormally located dorsally to the axon scaffold in KO embryos, comparable to elav staining (Fig. 3E).

As in vertebrates, glial cells play a key role in neuronal development in Dmel, particularly in axon guidance

(Chotard and Salecker, 2004; Freeman, 2006; Parker and Auld, 2004). Since the axon defects linked to Eip63E

deficiency appears early in CNS development, we explored the effect of the deficiency on glial cells. The reverse-

polarity (Repo) gene codes for a homeodomain protein that is expressed in all embryonic glial populations, except

midline glial cells, from stage 9 glioblasts to mature glial cells (Xiong et al., 1994). Co-staining of glial cells using

Repo and of axons using HRP antibody (Fig 3D), suggest a disorganization of glial cell bodies and the

appearance of structures that are distinct in KO embryos from control embryos. The combined penetrance of all

these relevant markers along the VNC structures suggests that Df(3L)E1 allele associated defects appear later in

development with more sever outcomes than those associated with Eip63E[81] allele (Fig S2). These results

suggest that axon defects in Eip63E KO embryos are accompanied by defects in neuronal cell body localization

and that these defects appear early in CNS development.

Specific downregulation of Eip63E in neurons leads to axon defects

To confirm that the CNS defects observed in the Eip63E KO mutant lines were indeed due to the loss of the role

of Eip63E in CNS development and to rule out any background effects of the germ-line deletion, we performed

tissue specific and timed downregulation of Eip63E using the Gal4-UAS system (Duffy, 2002). Utilizing the UAS-

RNAi fly lines for Eip63E developed by the Vienna Drosophila RNAi Centre, Eip63E was knocked down (KD)

using drivers for all cells (Act5-Gal4), neurons (elav-Gal4), glia (repo-Gal4), mesoderm/visceral muscles (mef2-

Gal4), dorsal epidermis/ peripheral nervous system (PNS) (GawB109.69-Gal4) and digestive system/ longitudinal

muscles (GawBtey5053A-Gal4) (Fig. 4). Since the Krüppel-GFP balancers used in previous experiments for

genotyping purposes also have an internal Gal4-UAS system, we used Cyo generic balancer instead when

needed. Ubiquitous down regulation of Eip63E using the Act5-Gal4 driver resulted in axon defects and lethality

similar to those observed with the germline deficient lines. Half of the embryonic population was expected to be control (no driver). Axon defects were observed in 59.7% of the embryonic population. This would suggest a 100% penetrance of defects for ubiquitous downregulation. Additionally, when embryos were left to complete the life cycle, all the flies that eclosed were positive for the Cyo marker, suggesting that like the germ-line mutants, ubiquitous downregulation of Eip63E is developmentally lethal and also corroborates the effectiveness of the RNAi construct used. VNC defects were observed in 49.4% of the embryonic population using the neuron-specific elav-Gal4 driver (Fig. 4C). These results are consistent with the notion that neuronal Eip63E function is required for axon development in Dmel. Intriguingly, when development was not interrupted for embryo collection, we observed an equivalent number of Cyo-positive and Cyo-negative flies eclosing (n=72 vs n=76). This suggests that although important in CNS development, Eip63E neuronal function alone does not play a role in developmental survival.

No noticeable defects were observed when Eip63E was specifically knocked down in glial, or any of the other tissues (Fig. 4 D to G). The total number of embryos per group and the percentage of embryos with observed axon defects are presented in Fig 4H. These results suggest that the functions of Eip63E in neurons, and possibly neuroblasts, are key to axon development in Drosophila, especially for the regulation of axon outgrowth and guidance.

Drosophila Eip63E functionally interacts with Rho1

Previous reports have shown that CDK5 and PCTAIREs regulate axon development by functionally interacting with the Rho family of GTPases (Matsuda et al., 2014; Nikolic M, 1998; Rashid T, 2001). However, whether such an interaction exists between PFTK1/Eip63E and Rho GTPases is not known. We used the Dmel line expressing dominant-negative Rho1^{72F} allele, balanced with the CyokrGFP chromosome. Characterization of this line revealed 100% lethality. Eip63E^{81} heterozygosity does not lead to axon defects (Fig. 5B; Fig. 1A, bottom left table and Fig. 1B bottom right table). Rho1^{72F} balanced line revealed severe VNC axon defects in GFP+ embryos, suggesting that heterozygosity was enough to cause the defects (Fig. 5C). 22.72% of double heterozygous (GFP negative) showed VNC defects that were of intermediate severity (Fig. 5D). Note that double heterozygosity also corrected Rho1^{72} lethality, since flies lacking the balancer phenotypic marker were present in the adult population when development was not interrupted for experimentation (multiple crosses, 155 live double-heterozygous adult flies out of 400 total n, data not shown).

Expression of constitutively active Rho1.V14 allele, driven in neurons also resulted in 100% lethality for heterozygous neuronal expression of the allele (n=131). 71.43% of embryos showed severe VNC defects with acutely decreased axonal density and some degree of misguidance (Fig. 5G). Specific neuronal downregulation of Eip63E resulted in VNC defects (Fig. 5F) like those observed previously (Fig. 4C, 4H). We then created a line expressing the balanced UASRho1.V14 allele in the second chromosome and the UASRNAiL63 in homozygosity on the third chromosome. Neuronal double downregulation by crossing this line with an elavGal4 driver fly revealed embryonic VNC axon defects that resembled a combination of both the controls (Fig. 5H). We detected less axons like those observed in Rho1.V14 control with acute defasciculation and misguidance, just like the

RNAiL63 control. Since all the individual control phenotypes were present, we conclude that there is a lack of

functional interaction between Eip63E deficiency and constitutively active Rho1. We also examined any functional

interaction between Eip63E and the other major members of the Rho family of GTPases, Rac1 and Cdc42 (Fig.

S3). Results show that that simultaneous Eip63E neuronal downregulation and activation of Rac1 and Cdc42

(constitutive active forms), but not downregulation (dominant negative form), lead to rescue of the phenotype.

This data suggests that in flies, the Rho family GTPases might be players in the same pathway as Eip63E

regulating axonal development. These results also point to a role for Eip63E upstream of Rho GTPases, by

regulating their activity or that of their regulators.

PFTK1 downregulation and deficiency leads to faster-growing axons in murine cortical neurons

Previous research has indicated that CDK5 inactivation leads to inhibition of axonal growth (Nikolic M, 1998;

Nikolic M, 1996) and that CDK related protein kinase PCTAIRE1-3 negatively regulates neurite outgrowth (Cole

A R, 2009). We investigated whether manipulation of PFTK1 activity had an impact on axon outgrowth in

mammalian cells. Primary cortical cultures from CD1 mouse embryos were infected with adenovirus expressing

GFP tagged-D228N PFTK1, GFP tagged-wild type PFTK1 or GFP control alone (Fig. 6). Both at 12hr after plating

and at 24 hours, a significant increase in axonal length was observed in cells expressing D228N but not the GFP

control or wt PFTK1 (Fig. 6a). Analysis of the mean neurite length at these time points suggest a noticeable

increase in neurite length relative to the control GFP and wt PFTK1. This suggests an increase in neurite length in

vitro upon downregulation of PFTK1 function, contrary to that observed with the loss of CDK5 (Nikolic M, 1996).

We generated a PFTK1 deficient mouse line to further examine the role of CDK14 in CNS development and

maintenance. Details of the generation of the PFTK1 KO mouse line is presented in Fig. S1 and described in the

methods section. Unlike PFTK1 deficiency in Drosophila, which leads to embryonic lethality, PFTK1 knockout

(KO) mice are viable, develop normally and do not show any obvious morphological defects compared to the wild

type littermates. Microscopic analysis of coronal sections of the adult mice brain on a mixed C57Bl6 background

(3rd generation) revealed no major abnormalities upon cresyl violet staining (Fig. S5A). Furthermore, no noticeable

deviation from the expected Mendelian ratios (Fig. S5B) were observed in embryonic or postnatal populations,

suggesting that PFTK1 loss alone, in mice, might not have the same effects observed with L63E deficiency in the

fly regarding lethality.

Next, we investigated whether the axonal outgrowth effects of D228N overexpression could be replicated by

PFTK1 deficiency, we studied populations of axons in primary cortical cultures from PFTK1 KO and WT embryos.

Cultures were prepared from E13.5 - 14.5 mouse embryos, fixed at 16 and 24 hours after plating and stained for

the microtubule marker Tau-1 and Hoescht. The axon length distribution at 16h in vitro (n=13-14

embryos/genotype, 236-469 cells/embryo) revealed a statistically significant increase in axons greater than 40 μm

in PFTK1 KO neurons and a decrease in shorter axons when compared to WT controls (Fig. 6C, 6D).

Nevertheless, the significant difference between KO and WT cultures at 16h is no longer detected after 24h in

vitro (Fig 6E). Together these results point to a role of PFTK1 in regulating axon outgrowth during early

axogenesis.

PFTK1 regulated axon outgrowth is dependent on RhoA

Based on the functional interaction of Eip63E and Rho GTPases, next we investigated whether manipulation of RhoA activity in PFTK1 KO neurons could rescue the effects on axonal outgrowth. PFTK1 KO or WT neurons were treated with RhoA activator CN03, which blocks intrinsic, and GAP stimulated GTPase activity leading to constitutive activation of RhoA by robustly increasing the level of GTP bound RhoA (Flatau et al., 1997). In the same experiment presented in Fig. 6C, treatment of the RhoA activator CN03 rescued the increase in axon lengths observed in PFTK1 KO neurons (Fig. 6C and E). These results present a strong case for the role of RhoA activity downstream of PFTK1 and the specific involvement of PFTK1 mediated RhoA regulation in axonal outgrowth.

We then assessed whether PFTK1 levels influence RhoA expression. Examination of the endogenous levels of RhoA protein revealed significantly higher levels of RhoA in PFTK1 KO neurons compared to WT (Fig. 7B and C; Fig. S6). Of note, no significant difference in Rac1 or Cdc42 levels were observed (Fig. S6), suggesting that, Rac1 and Cdc42 were unaffected by PFTK1 deficiency. We then assessed whether PFTK1 regulates RhoA gene expression at a transcriptional level. Reverse transcription polymerase chain reaction (RT-PCR) analysis show no significant differences in RhoA mRNA level (Fig. 7A and C).

To further investigate the relationship between the activity of RhoA and PFTK1, we assessed RhoA GTPase activity in PFTK1 KO mice. The activation level of RhoA was determined by Rho GTPase activity assay utilizing GTPase activation kit on cell lysates. Briefly, cell extracts were prepared from cortical cultures of PFTK1 wild type and KO littermates at E13.5-14.5 and then incubated with Rho-binding domain (RBD) of Rhotekin RBD beads provided with the kit, which pulls down the endogenous active form of RhoA from the cell lysate. Active RhoA levels were examined by western blotting. The detected signal is expected to correlate with the activity level of RhoA. Interestingly, despite RhoA protein upregulation in PFTK1 KO mice, we did not detect any RhoA activity in the KO extracts (Fig. 7D). Based on these results, we conclude that in PFTK1 deficient cells a) RhoA protein levels are upregulated despite no noticeable change in RhoA mRNA levels and b) RhoA activity remains unaffected despite upregulation in RhoA protein level.

RhoA phosphorylation and activity is affected by PFTK1

Next, we explored possible mechanisms by which PFTK1 might be regulating RhoA activity. We performed an *in vitro* kinase assay in the presence or absence of immunoprecipitated PFTK1 to examine the ability of PFTK1 to phosphorylate RhoA. As control, IgG immunoprecipitated samples were used. Change in RhoA phosphorylation status was analyzed by Western blot analysis for the resulting reactions using a phosphoserine antibody (Fig. 7E). Interestingly, a noticeable increase in serine phosphorylation of RhoA by PFTK1 was observed when GDP was externally added during incubation. These results suggest that GDP-RhoA is a direct in vitro substrate of PFTK1 (Fig. 7E).

Next, we investigated whether PFTK1 phosphorylation of GDP-bound RhoA had an effect on RhoA activity using an absorbance-based G-LISA assay. We used the products of the in vitro Kinase Assay described in the previous section and followed it up with a G-LISA reaction to test for RhoA activity. A significant increase in RhoA activity

1 was detected as a function of PFTK1 compared to IgG control. Additionally, a significant increase in RhoA activity

2 was observed when GDP bound RhoA was incubated with PFTK1. These results suggest that GDP-RhoA is not

3 only a substrate of PFTK1 but also that this phosphorylation results in increased RhoA activity (Fig. 7F). Taken

4 together, these results point to an as yet unidentified role of PFTK1 in regulating RhoA activity by phosphorylating

5 and thereby activating GDP bound RhoA.

Discussion

The proper development of the CNS requires tight coordination of several important processes including axon
development. This is a complex process involving a broad range of factors that are present in the neuronal
microenvironment. However, the precise role of PFTK1 in neurite morphogenesis and implications in CNS
development has not been fully characterized. Here we present evidence that PFTK1 (Eip63E in Drosophila), is
involved in this process and mediates its action on axon development by regulating its downstream effector via its
kinase activity.

Screening of Eip63E deficient embryos from two different and independent mutant alleles consistently show
defects in VNC structures. These axon defects occur along neuronal and, to a lesser extent, along glial cell body
disorganization. The penetrance values of neuron-related defects are comparable to those of axonal defects
already from stage 10 onward (Fig. S2). This does not seem to be case for glial defects. No consistent glial
perturbations were noticed between the two Eip63E mutant alleles. Taken together, these observations suggest
that the axon defects are related to neuronal, rather than glial-related Eip63E functions.

More than one of the CNS markers used to characterize the VNC phenotype of Eip63E KO embryos, point to a
potential defect in biological processes that regulate or mediate cell identity specification and/or neuronal
differentiation processes (Fig. 2B and Fig. 3). These defects could be described in general as an abnormal cell
cortex laminar organization along the VNC of mutant embryos, a process that is highly dependent on the timing
and pattern of progenitors' birth and divisions (Isshiki et al., 2001). It is possible that defects in Eip63E mutants
are related to temporal control of new neurons and/or their differentiation. Indeed, it has been shown before that
the absence of cell cycle progression or cytokinesis in neuroblasts leads to abnormal expression pattern of
hunchback (Hb) (Grosskortenhaus et al., 2005), one of the progenitor temporal transcription factors that regulate
first-born cell fates in Drosophila (Isshiki *et al.*, 2001). Previous reports show that Eip63E has a role in cell cycle
regulation, since its downregulation in embryonic cell cultures by RNAi leads to low mitotic index and therefore
larger G2 cells (Bettencourt-Dias et al., 2004). There is also evidence that Eip63E interacts with cyclinY to
phosphorylate the low-density lipoprotein receptor-related 6 (LRP6; 'arrow' in Drosophila) and mediate Wnt
signaling (Davidson et al., 2009); a pathway broadly recognized as essential for regulating progenitor proliferation,
cell lineage decisions and neuronal differentiation in insects and vertebrates (Grigoryan et al., 2008; Zechner et
al., 2003).

In addition to this early potential function for Eip63E, our data strongly support a later role for this kinase in
neurons to regulate axon development. It is evident that early deregulation of differentiation timing would set a
precedent for axon defects, since for example, pioneer neurons and axons are needed for correct pathfinding and
fasciculation of more mature axon pathways (Sanchez-Soriano and Prokop, 2005). Nevertheless, our results that
Eip63E elav-driven downregulation on a WT background leads to axon defects (Fig. 4C) suggests a more direct
involvement of Eip63E in mechanisms relevant to axon guidance.

A search for downstream regulators of Eip63E activity resulted in the identification of RhoA as a potential target.
RhoA belongs to the Rho subfamily of GTPases within the Ras superfamily, which also include Cdc42 and Rac1.

The family which now includes 20 total members (Hodge and Ridley, 2016) is involved in diverse cellular

functions such as the regulation of cytoskeletal rearrangements, cellular motility, cell polarity, axon guidance,

vesicle trafficking and the cell cycle (Heasman and Ridley, 2008). Most of the Rho family of proteins function as

switches that undergo conformational change between the inactive GTP-bound and the active GTP-bound state.

Guanine nucleotide exchange factors (GEFs) activate these Rho GTPases by regulating the dissociation of GDP

and the subsequent uptake of GTP from cytosol whereas GTPase-activating proteins (GAPs) facilitate their

inactivation by increasing the rate of hydrolysis of GTP. Inactive Rho GTPases in the cytosol are often bound to

guanine nucleotide dissociation inhibitors (GDIs) which prevent the GTPases from interacting with their Rho-

GEFs at the plasma membrane (Garcia-Mata et al., 2011), providing an additional layer of regulation. The role of

RhoA in neuronal morphogenesis has been long implicated and has become clearer over the past several years.

Although controversial, with initial overexpression studies suggesting inhibition of RhoA activity promoting neurite

outgrowth (Jin and Strittmatter, 1997; Kozma et al., 1997; Kranenburg et al., 1997), several others (Kuhn et al.,

1999; Threadgill et al., 1997; Zipkin et al., 1997) demonstrate that RhoA activity positively affects neurite initiation

and extension resulting in longer axons. Our results side with the former, pointing to a role for PFTK1 in regulating

activation of RhoA leading to more controlled axogenesis. Both expression of a dominant negative form of RhoA

and PFTK1 deficiency, resulting in reduced RhoA activity, leads to uncontrolled initiation of axon outgrowth and

hence longer neurites.

When Rho1 activity was overexpressed in specific neuronal populations with Eip63E deficiency, VNC defects in

axonal defasciculation and misguidance appeared to be additive as both features of Eip63E deficiency (Fig. 5F)

and Rho1 constitutive active overexpression (Fig. 5G) could be observed (Fig. 5H). In contrast, simultaneous

expression of the constitutive active forms of Rac1 (Fig. S5D) and Cdc42 (Fig. S5J) and Eip63E downregulation

in neurons resulted in an intermediate phenotype, implying that Eip63E may be regulating upstream factors that

activate Rac1 and Cdc42 and may need further investigations. Intriguingly, in mice neurons, no change in Rac1 or

Cdc42 expression levels were observed (Fig. S6). One possible explanation for this observed difference between

the flies and mouse could be compensation by PFTK2 and PCTAIREs in mice. In flies, in the absence of these

homologs, Eip63E emerges to have a dominant role in CNS development and regulation of the activity of Rho

family of GTPases. Nevertheless, based on our results with Rho1 activity in flies, we conclude that PFTK1

functions upstream of Rho1, and PFTK1 deficiency results in axonal defects due to perturbations in Rho1 activity.

CDK5 and PCTAIRE, two close family members of PFTK1 have been implicated in axon outgrowth (Nikolic M,

1998; Nikolic M, 1996). In our experiments, a noticeable change in axon lengths at earlier time-points (Fig. 6)

might indicate that PFTK1 is more important in regulating the initiation of axon outgrowth rather than being

involved in axon outgrowth. Interestingly, the percent of neurons with longer axons are higher at 16 hours in

comparison to 24 hours (Fig 6D). This indicates that over time, the rate of axonal outgrowth evens out and the

effect is lost, suggesting the participation of other downstream factors that regulate axon elongation. Note that this

change in axon lengths is not due to generalized effect on the health of the cell as the expression of the dominant

negative PFTK1 had no effect on the survival of neurons (Fig. S4).

The lack of any obvious gross CNS phenotype in PFTK1 deficient mice compared to those observed in flies, although intriguing, could be due, in part, to the presence of other PFTK1 related kinases. Flies merely possess the PFTK1 homologue Eip63E but not PFTK2 and PCTAIRE kinases. In comparison, mammals contain both PFTK1 and 2 that show 68 % homology, as well as PCTAIRE 1-3, which are 61 % homologous to PFTK1 and are all highly expressed in the mouse brain (Cole A R, 2009; Hirose T, 2000; Mikolcevic P, 2012; Okuda T, 1992). The presence of these highly similar genes in the mammalian system could compensate for the lack of PFTK1 and result in a lower level of abnormalities in the KO mice. However, the full extent of the effect of PFTK1 deficiency on brain morphology would require a closer examination of specific brain structures. Although the lack of gross developmental defects is not inconsistent with effects on axon length, future analysis of the PFTK1 mice would require a more careful examination.

We observe a noticeable increase in RhoA protein level in PFTK1 KO cortical neurons (Fig. 7B and C) despite no detectable increase in mRNA levels suggesting a posttranslational regulation of RhoA levels. It could be speculated that PFTK1 also plays a role in regulating cellular RhoA levels by regulating RhoA phosphorylation and ubiquitination. In this regard, previous reports have identified specific E3 ligases, Cullin-3, that targets GDP-bound RhoA for degradation (Chen et al., 2009) and further studies may provide clues to the role of PFTK1 in this pathway. Intriguingly, this increase in RhoA level did not translate to more active RhoA but rather a noticeable decrease in detectable active RhoA in PFTK1 KO neurons (Fig. 7D) suggesting that PFTK1 deficient neurons had reduced GTP-bound active RhoA. Additionally, PFTK1 deficiency also resulted in perturbation in axon length of cultured cortical neurons. These results are in line with previous studies that show that reduced RhoA activity leads to longer axons (Sebok A, 1999). Combined with the observations that pharmacological activation of RhoA resulted in the loss of the observed difference between PFTK1 WT and KO neurons (Fig. 6E), our data strongly supports a model where PFTK1 regulates RhoA activation which in turn mediates the initiation of axogenesis.

Observations via an in vitro kinase assay lend further support to this model (Fig. 7E). Serine phosphorylation of RhoA by PFTK1 in the presence of GDP indicates a preference for GDP-bound RhoA by PFTK1. Downstream targets or substrates of PFTK1, other than cyclin Y, that may play prominent role in its function in neurodevelopment are not currently known. However, our data clearly establishes RhoA as a downstream effector of PFTK1 and identifies regulation of GDP-bound RhoA phosphorylation and activation by PFTK1 as a critical factor in PFTK1 mediated axogenesis. Accumulating evidence suggests the importance of regulation of Rho GTPases phosphorylation in its function (Chang et al., 2011; Tong et al., 2016). In this regard, it will be critical in future studies to address potential phosphorylation sites on RhoA, regulated directly or indirectly by PFTK1. Based on our evidence, it is also difficult to rule out the regulation of other upstream factors by PFTK1 that may mediate RhoA activity. This adds a further mode of regulation of RhoA function by PFTK1. Taken together, the evidence presented here proposes a working model for PFTK1 in mediating neuronal polarization and axon growth via RhoA regulation.

Figure Legends

Figure 1. Eip63E deficiency leads to defasciculation and misguidance of VNC axons.
Ventral views of whole mount stage 14-15 embryos from both Eip63E mutant lines, co-stained for GFP and BPI
02 (A) or Fasciclin-II (1D4 mAb) (B). Penetrance of axon abnormalities was calculated as the % of embryos of a
given scoring phenotype that show any defect from the total population of such phenotype (GFP+ or GFP-
embryos). Arrowhead points to normal looking longitudinal connectives and arrows to disrupted structures in KO
embryos. L63[81]: Eip63E[81].

Figure 2. Eip63E deficiency leads to early embryonic developmental defects.
Ventral views (unless otherwise specified) of whole mount embryos are shown head to the left. Stage 5-8
embryos are from Eip63E[81]/TM3krGFP,Sb line are of unknown genotype. Panels for stages 10-11 show Eip63E[81]
homozygous mutant embryos. (A) Staining for single-minded at early embryonic stages (st 5-10) show no gross
difference in midline formation but cellular morphological differences are noticable. (B) Staining for engrailed/
invected (4D9 mAb). 18% (n=48) of total population of stage 8-9 embryos show acute segmentation defects. At
stage 10-11, cell bodies appear disorganized (arrowheads) to the extent of loss of hemi-segments division at the
midline (wavy arrow) compared to control (straight arrow). (C) Staining for the HRP antigen. In top 4 panels, a
lateral view of embryos is shown. 10% (n=48) of st 8-9 embryos show premature HRP staining (arrowheads). At
late stage 11, precocious axon outgrowth (arrows) seen in 36% of KO embryos (n=52).

Figure 3. Eip63E deficiency leads to early segmentation defects.
(A, B, C and D) Ventral views of whole mount embryos stage 12-15 shown head to the left. CTR show GFP+
embryos (wild-type and mutant heterozygous population) and Eip63E[81] represent homozygous mutant embryos
(GFP-). Immunostaining for Elav (A), Neuroglian (B), Neurotactin (C, top 2 panels) and BP102 (C, bottom 2
panels) showing axon defects (arrows). (D) Co-staining for glial populations (Repo) and axons (HRP) showing
disorganized cell bodies (arrowheads). (E) Merged elav and HRP staining images (lateral view) of stage 12
embryos shown in panel A. Elav-positive cells appear to be dorsally deeper than normal (arrowheads), under the
axon bundles. (F) VNC flat preparations of timed embryos (10-11h, stage 14) co-stained for Futsch and HRP are
shown dorsal up, head to the left. Arrowheads point to some of the structures where neuronal organization is
different from control embryos. Arrows point to examples of axon growth abnormalities

Figure 4. Only ubiquitous and neuronal downregulation of Eip63E leads to embryonic axon defects.
Ventral views of stage 14 whole mount embryos shown head to the left. A UAS-Gal4 approach was used to drive
Eip63E downregulation in specific tissues during embryonic development. Embryos negative for GFP (present in
balancers for both the Gal4 and the UAS lines) was considered as experimental embryos vs the GFP positive of
the same progeny. (A) A representative GFP+ embryo is shown. (B) Ubiquitous downregulation led to a replica of
the germline deficiency lines, regarding CNS defective phenotype and developmental lethality (see text for
details). (C) Neuronal-specific downregulation led to a robust and highly penetrant axonal phenotype. (D, E, F, G)
Downregulation of Eip63E in other tested tissue (indicated in red) did not cause any detectable axonal defects.

The number of embryos per genotype and the percentage of embryos per group showing axonal defects is

summarized in (H).

Figure 5. Eip63E functionally interacts with Rho1 to regulate axogenesis.

Ventral views of stage 14 whole mount embryos stained for BP102 antigen are shown head to the left. Panels

show representative images for each specified genotype, (A-D) Representative images of control embryos (A)

single heterozygous Eip63E[81] (B) or Rho1[72]F (C) from a cross between Eip63E[81] KO line and Dominant-Negative

Rho1 (Rho1[72]F). (B, C) Images of presumed genotype (see text for details). (D) Axon defects of intermediate

severity in 22.7% of double-heterozygous mutants (GFP-). (E-H) UAS-elavGal4 approach to drive Eip63E

downregulation and constitutive active Rho1 overexpression in neurons in two independent Dmel lines. Lines

were then crossed to generate the indicated genotypes. (F, G) Heterozygous neuronal downregulation of Eip63E

or Rho1 showing axon defects with specified penetrance. 86.96% of Double-heterozygous embryos (H) had axon

defects showing features of both heterozygous controls. Penetrance values from stages 13-17, n as follows: B)

>100, C) 86, D) 22, F) 89, G) 126, H) 69. n represents the total of screened embryos from several different

pairings/collection pools.

Figure 6. PFTK1 downregulation leads to faster-growing axons in murine cortical neurons.

Primary cortical neurons derived from CD1 embryos (E 14-15) were infected with 100 MOI of adenoviruses

expressing GFP alone, GFP and WT- PFTK1 or GFP and D228N- PFTK1. After 12 or 24 hours in culture, cells

were fixed and stained with Hoechst 33258 (nuclei visualization). Axons of GFP+ neurons were measured using

Image J. (A) Values are presented as averages ±SEM from 122-143 cells/treatment/timepoint. Statistics were

done using two-way ANOVA followed by Tuckey's post-hoc. (B). Same data as in (A) but presented as axon

length relative to the GFP control. (C) Representative images of primary cortical neurons from E13-14 embryos

from HET x HET crosses from WT and KO cultures fixed at the specified times and stained for Tau and Hoechst.

(D) Axon length distribution per genotype from the indicated number of embryos (236-469 cells/embryo) at 16

hours in vitro. Each column represents the percentage of cells in the specified range of length for the indicated

genotype. Non-treated (NT) cultures were also fixed after 16 and 24h in vitro. Graph represents the average axon

length/genotype/treatment. t-test was used for statistical analysis whenever p value is specified. (E) 16h cultures

were treated for the last 8h with 0.5ug/ml of RhoA activator (Cytoskeleton Inc. CN03). Graph represents the

average axon length treatment.

Figure 7. RhoA phosphorylation and activity is affected by PFTK1.

Neuronal cultures from E13-14 embryos from HET x HET crosses were examined after 18-20h *in vitro*. Total RNA

(A) or protein (B) was extracted followed by RT-PCR or western blot. Representative images from RT-PCR (A)

and western blot (B) assays are shown. (C) Bar graph represents the average band intensity from 3 KO and 6 WT

for mRNA and 7 WT and KO each for protein. t-test was used for statistical analysis. D) Assessment of RhoA

activity using an IP-based assay. Neuronal extracts were incubated with Rhotekin PBD beads to pull down active

RhoA. WT extract preincubated with $GTP_{\gamma}S$ (positive) or GDP (negative) were used as controls. IP was followed

by western blotting for RhoA. Two representative assays of four independent assays are shown. E) *In vitro* Kinase
assay. IPed protein (either with IgG control or PFTK1) from WT cortical neuronal culture extracts was used as
source of kinase and incubated with human recombinant RhoA (His-tagged) with or without GDP. Results were
analyzed for phospho-serine signal. F) RhoA activity assay following *in vitro* kinase assay. Kinase reactions were
analyzed by G-LISA to test for RhoA activity. Bar graph represents mean values ±SEM. Two-way ANOVA
analysis of grouped data and Sidak test for multiple comparisons are indicated on bar graph. RHO1: His-RhoA
human recombinant protein.

Supplemental figure legends

Figure S1. Strategy for generation of PFTK1 KO murine line.

The mouse was designed and developed by the Texas A&M Institute for Genomic Medicine (TIGM) (A) Strategy.

Exon 6 (from nucleotide 18-Exon6 to nucleotide l-Exon7) of the Cdk.14 gene at chromosomes (4,803,391-

5,380,19 7) was replaced for an IRES/bGeo/Poly A cassette by homologous recombination. (B) Southern Blot

screening of clones. (C). PCR strategy for genotyping. Bottom panel shows representative gel image with

samples from the three possible genotypes.

Figure S2. Eip63E deficiency leads to concerted defects in axons and neurons of the Drosophila VNC.

The combined penetrance of markers relevant to the same VNC structures are presented as one. Axon-

associated penetrance reflects those calculated using HRP, BP-102 and Fasciclin-II antibodies. Neuron-related

values include Neuroglian, Elav and Futsch-revealed penetrance. Penetrance was calculated as the % of

defective KO embryos from the total KO embryos screened. Both deficient lines show the same nature of defects.

Although it seems that defects associated to the Df(3L)E1 allele are more acute and appear later in development,

Eip63E81 shows a robust axonal phenotype. ND: Not determined.

Figure S3. Eip63E functionally interacts with Rac1 and Cdc42 in D. melanogaster to regulate axogenesis.

Ventral views of stage 14-15 whole mount embryos stained for BP102 antigen are shown head to the left. Panels

show representative images for each specified genotype. An UAS-elavGal4 approach was used to drive Eip63E

downregulation and either constitutively active or dominant negative forms of Rac1 or Cdc42 overexpression in

neurons of independent D mel. lines. Fly genotype following crosses are indicated. (A-F) Neuronal functional

interaction between Eip63E and Rac1. Only the combination of Eip63E deficiency and constitutively active Rac1

(UAS Rac1 V12; D), but not DN-Rac1 (UAS Rac1 N17; F), leads to an intermediate phenotype. (G-L) Functional

interaction between Eip63E and Cdc42. Ventral views of the indicated genotypes of crosses between Eip63E and

Cdc42 constitutive active form (UAS Cdc42 V12) or the dominant negative Cdc42 (UAS Cdc42 N17) are

presented.

Figure S4. Manipulation of PFTK1 levels has no impact on neuronal survival.

Primary cortical neurons derived from CD1 embryos (E 14-15) were infected with 100 MOI of adenoviruses

expressing GFP alone, GFP and WT- PFTK1 or GFP and D228N- PFTK1. After 12 or 24 hours in culture, cells

were fixed and stained with Hoechst 33258 (nuclei visualization). nuclear integrity of GFP+ neurons was used as

a criterion for vitality. Survival values represent the % of healthy nuclei of the total GFP+ cells. Table is showing

average +SEM from three independent experiments. Graph shows the same data but presented as value relative

to the GFP control.

Figure S5. PFTK1 deficiency does not appear to cause any gross brain defects or lethality.

(A) Brains from three-month-old mice (3rd generation backcrossed) were collected, perfused and fixed for

sectioning. 14µm sections were collected from cryopreserved samples, starting from the cortex at bregma: ~

2.710 mm up through the 4th ventricle at bregma: ~-4.20 mm. Sections were then stained with cresyl violet to
examine the gross anatomy of the brain. No abnormality was detected for KO mice (n=3/genotype).
Representative sections from bregma 1.98, 1.70, 0.74, 1.00, 1.64, and 2.75 mm are presented. (B). Embryos or
mice (=> 6th generation backcrossed) from HET x HET crosses were genotyped and incidence of the three
possible outcomes is presented as % of the total n. No gross deviation from the expected 25:50:25 Mendelian
ratio was detected.

Figure S5. Differential expression of Rho GTPases in brains of Wt vs KO E13-14 murine embryos.
Only RhoA is dysregulated in cortical neurons. Basal levels of RhoA, Rac1 and Cdc42 protein in WT and Pftk1-
defficient cortical neurons. Primary cortical neurons from E13-14 embryos from HET x HET crosses were
dissected and plated at 0.65-0.75x106 cells/ml. After 18-20h in vitro, proteins were extracted followed by western
blot. Top: Representative images from western bot are shown. Bottom, levels were assessed by densitometry
using Image J and Rho GTPases signals were standardized vs loading control (Actin). Each column represents
the average from 3 embryos/genotype +SEM. Significant difference between WT and KO neurons was only
found for RhoA levels. Each column represents the average from n embryos/genotype +SEM. Significant
difference between WT and KO neurons was only found for Cdc42 levels. t-test was used for statistical analysis
whenever a p value is specified

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Lead contact

All requests for further information, reagents and resources should be directed to the lead contact, Alvin Joselin (Alvin.Joselin1@ucalgary.ca).

Availability of Data and Materials

Source data are provided with this paper. All other data supporting the findings of this study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by grants from the Canadian Institutes of Health Research (grant numbers FRN # 15123 and 148402), Heart and Stroke Foundation of Ontario (HSFO), Canadian Stroke Network, and Centre for Stroke Recovery, Parkinson Society Canada, Parkinson's Disease Foundation, Parkinson's Research Consortium, and Michael J. Fox Foundation (DSP.). YRG. was supported by

grants from Ontario Graduate Scholarship in Science and Technology and Ontario Graduate Scholarship. DSP is an HSFO Career Investigator.

Author contributions

Materials and Methods

Drosophila strains:

Flies were kept under standard conditions. The following stocks were obtained from the Bloomington stock centre:

W1118, Eip6381/TM6tb, Df(3L)E1/TM6, green balancer flies (L2Pin1/CyokrGFP, D*gl3/TM3krGFP,

In(2LR)nocSco,b1/CyoActGFP and Sb/TM3ActGFP,Ser) and Gal4 driver lines (elav::Gal4, Repo::Gal4/TM3,Sb,

elav::Gal4;UAS::Dcr2.D and UAS::Dcr2.D;Act5::Gal4/Cyo). Three independent RNAi Drosophila lines were

obtained from the Vienna Drosophila Rnai Center (UAS::RNAiL63/Cyo, UAS::RNAiL63 and

UAS::RNAiL63/TM3,Sb). For the purpose of mutant embryos characterization, the TM6 balancer in both Eip63E

deficient lines (Stowers *et al.*, 2000) was replaced with the TM3.Kruppell-GFP balancer chromosome (Casso *et*

al., 2000). To perform Eip63E specific downregulation, the Gal4:UAS system was used (Duffy, 2002). Balancers

in all UAS and Gal4 lines involved in the crosses strategies were changed for the Actin5c-GFP balancer

chromosomes (Cyo or TM3 as needed) (Reichhart and Ferrandon, 1998). .

Drosophila Immunohistochemistry:

Drosophila embryos from Eip63E mutant lines were collected for 2 hours on grape juice-agar medium (Ashburner,

1989), and allowed to carry on development to embryonic stages 6-15 (Campos-Ortega and Hartenstein, 1985),

in batches including only 2-3 stages each, by further incubation at 25°C. This developmental window includes all

events of embryonic CNS development, from anlage formation to differentiation and axogenesis (Weigmann *et*

al., 2003). Whole mount embryos were prepared following standard procedures (Harlow and Lane, 2006).

Embryos were dechorionated with 50% bleach, followed by fixation for 20 minutes in heptane saturated with 4%

Paraformaldehyde in 136 mM NaCl, 2.68 mM KCl, 10.14 mM Na2HPO4, 1.76 mM KH2PO4, pH 7.4, (PBS).

Vitelline membrane was removed by vigorously shaking the fixed embryos in methanol, followed by further

methanol washes. Embryos were re-hydrated using PBS-0.1% Triton X-100 (PBT) and then blocked for 20

minutes using 20% Normal Goat Serum (NGS) (GE Healthcare, Amersham RPN410) in PBT. Fluorescence

immuno co-staining was performed against GFP (to amplify the presence of the balancer embryonic marker),

HRP and a CNS marker. Primary antibodies were as follows: mouse anti-GFP (JL-8, Living colors, 1/100), Cy3-

anti-HRP (1/25), mouse BP102 (1/10), mouse anti-En/Inv (4D9, 1/10), mouse anti-single-minded (1/5), mouse

anti-Futsch (22C10, 1/10), mouse anti-Neuroglian (BP-104, 1/10), mouse anti-elav (9F8A9, 1/14) and mouse anti-

Repo (8D12, 1/10). All monoclonal antibodies against CNS markers were obtained from the Developmental

Studies Hybridoma Bank developed under the auspices of the NICHD and maintained by The University of Iowa,

Department of Biology, Iowa City, IA 52242. Such antibodies were developed by C. Goodman, S. Crews, S.

Benzer and G.M. Rubin. Since the anti-HRP antibody was Cy3-conjugated, only Alexa 488-Goat anti-mouse

(1/100) was used as secondary antibody, after a second blocking step. Stained embryos were then directly

mounted in Vectashield (Vector laboratories, H-1000) and visualized under a fluorescent microscope.

Embryo dissections were performed essentially as described (Benveniste et al., 1998). 7-13 hours timed embryos

were transferred to a piece of double-sided tape mounted on a polylysine-covered microscope slide, and then

rolled using forceps to mechanically remove the chorions. Embryos were then covered with PBS and examined

under fluorescence light for GFP expression. Absence of GFP expression would indicate an Eip63E KO. After

genotyping and fine determination of developmental stage based on morphological features, a dorsal incision was

made in the vitelline membrane of each with a sharpened tungsten needle (Small Parts Inc TW-006-60). Each

embryo was then removed from the membrane and placed ventral side down directly on the polylysine-covered

surface of the slide, while still submerged in the PBS. Using the tungsten needle, an incision was made

longitudinally along the dorsal surface of the embryo and the body walls were pressed down onto the glass. The

gut and remaining yolk sac were then removed, resulting in a flat embryonic preparation. Dissected embryos were

then fixed for 30 minutes at room temperature in 4% paraformaldehyde in PBT. One-hour blocking was performed

using 10% NGS in PBT, followed by antibody staining. Fluorescence co- immunostaining was performed against

HRP and a CNS marker. Same antibodies mentioned above were used but at different concentrations: 1/50 for

primary antibodies and 1/300 for secondary. Preparations were mounted using Fluoromount™ mounting media

(Sigma F4680), sealed with clear nail polish and visualized under a fluorescent microscope.

Transgenic Mice Systems

All experimental animal studies were conducted in accordance with the guidelines of the University of Ottawa

Animal Care Committee and conformed to the guidelines set forth by the Canadian Council on Animal Care and

Canadian Institutes of Health Research.

Generation and analysis of PFTK1 Transgenic (knockout) Mice

PFTK1 heterozygote mice were generated by the Gene Targeting and Transgenic Facility of Texas A&M Institute

for Genomic Medicine (TIGM), on mixed (129/SvEvBrd x C57BL/6) background. To generate the mutant allele,

PFTK1 Gene (MGI: 894318) and the transcript sequence of PFTK1 or (Cdk14-002 ENSMUST00000030763)

located on chromosome5 (4,803,391-5,380,197) (reverse strand) was targeted. The intron-exon structure of the

PFTK1 gene obtained by genomic mapping is illustrated in Fig. S1A. PFTK1 heterozygotes were generated by

replacing the majority of exon 6 of the Mus Musculus PFTK1 transcript (ENSMUST00000030763) starting from

nucleotide 18 with the neomycin (Neo) resistance cassette via homologous recombination. The resulting insertion

in the PFTK1 gene causes a frame shift altering the open reading frame and leading to a premature stop codon

which produces a truncated protein without the entire kinase domain. The correct integration of the targeting

sequence within the Pifraire1 gene was confirmed by Polymerase Chain Reaction (PCR) and Southern blot

hybridization analysis in the 129/SvEvBrd embryonic cells that were transformed (Fig. S1B). Following Bgl II or

Acc65 I restriction digestion of the genomic DNA, Southern blot was performed using probes targeting 5' or 3'

regions outside of the targeting vector, respectively. The predicted band sizes of 15.9 kb after Bgl II digestion with

the 5' probe and 13.5 kb after Acc65 I digestion with the 3' probe confirms the homologous recombination of the

external cassette. The transformed PFTK1 allele was easily detectable in the transformed mice through the agouti

coat color of the chimeras. The produced chimeric mice were viable and fertile; after being crossed with wild type

mice they also produce the PFTK1 heterozygote deficient mice that were viable and fertile. PFTK1 F1 generation,

homozygote (null) mice, which were obtained by breeding heterozygous parents were also viable and fertile. To

establish PFTK1 knockout line on a pure C57BL6 background, we backcrossed mixed (129/SvEvBrd x C57BL/6)

heterozygote mice from the PFTK1 deficient line with C57BL6 for 10 generations, which continue to be viable.

PCR performed on tail genomic DNA samples obtained from PFTK1 wildtype, heterozygotes and knockout

PFTK1 mice and the corresponding primer pairs (fwd. 5'-TGACCCTTGTCTTTGAATACGTG-3' and rev. 5'-

GGAAGCCTAAATTGTAAGGATCAGG-3' generating a 418 bp wild-type fragment and fwd. 5'–

GCAGCGCATCGCCTTCTATC-3' and rev. 5'-GGAAGCCTAAATTGTAAGGATCAGG-3' generating a 341 bp

fragment) are presented in Fig. S1C.

PCR mix was run with GoTaq Green Master Mix 2x (Promega, M712B). The PCR amplification process was as

follows: initial amplification designed as touchdown PCR including: 6 cycles of denaturation 98 °C for 17 sec,

annealing 69 °C (touchdown, -1°C per cycle) for 15 sec, amplification 72°C for 15 sec, and further amplification of

the target genes was as follows: 35 cycles of 98 °C for 17 sec, 60 °C for 15 sec, and 72 °C for 15 sec. PCR

products were resolved on a 1.5 % agarose gel- Ultra Pure Agarose (Invitrogen, 15510-027) and visualized by

ethidium bromide (Ethyl hexadecyl dimethyl bromide $C_{20}H_{44}BrN$) (Sigma Ultra, C5335-1000).

Real Time PCR (Quantitative RT- PCR)

All steps were performed in RNAase-free environment. To ensure RNA-free surface, RNase away (MBP

Molecular Bioproducts, 7003) was applied. Total RNA was extracted from mouse embryo whole brains/cortices at

embryonic day E 13.5-14.5 using Trizol Reagent (Ambion, 15596026) based on Invitrogen protocols. 25 ng of

extracted RNA was used per reaction. To determine disruption of PFTK1 transcription the SuperScript III Platinum

SYBR Green One-Step RT-PCR kit (Invitrogen, 204174) was used. Negative controls, no RNA and no RT were

run in parallel with both the GAPDH housekeeping gene and PFTK1. Each sample was run in quadruplets and

n=3. Results were normalized against GAPDH. Primers used for amplification of the target genes were as

follows: (GAPDH-Fwd.: 5'-GGTGAAGGTCGGTGTGAACG-3', GAPDH-Rev: 5'-CTCGCTCCTGGAAGATGGTG-

3') to generate a 233 bp product and (PFTK1-Fwd: 5'-CAGCGATCTGCCTCCACGGC-3') designed within exon13.

(PFTK1-Rev: 5'-GGCCCTCATGCTCTCTCCAGC-3') designed within exon 14, to generate a 341 bp product. The

primers would also be able to produce a short heptamer within mutated region (exon 8); thus, it could not interfere

with mRNA levels for PFTK1-KO. The program for RT-PCR amplification was as follows: 48 °C for 30 min, 95 °C

for 10 min, and 40 cycles of 95 °C for 15 sec, 60 °C for 30 sec and 72°C for 30 sec. The RT-PCR products were

resolved on a 1.5 % agarose-ethidium bromide gel as well.

Cell Culture

Human Embryonic Kidney cells (HEK-293T) were cultured in Dulbecco's Modified Eagle Medium (DMEM) (High

Glucose 4500 mg/L with 4 mM L-Glutamine without Sodium Pyruvate 0.1 µm sterile filtered) (ThermoScientific,

SH30022-01) supplemented with 10 % Fetal Bovine Serum (FBS) (Hyclone, SH30397.03) and 1 % Antibiotic/

Antimicotic Solution100x (10,000 units/ml of penicillin, 10,000 µg/ml of streptomycin, and 25 µg/ml of

Amphotericin B , 0.2 µm filtered) (ThermoScientific, SV30079.01) at 37°C in a 5% CO_2 atmosphere.

Primary Cortical Neuron Cultures

Cortical neuron cultures were prepared either from wild-type CD1 mouse embryos purchased from Charles River
or from heterozygous crosses between PFTK1 mice, at gestational day /embryonic day E 13.5-14.5. Embryos
were considered E 0.5 d, at the time vaginal plug was detected. Embryos were removed from the uterine horns
and transferred to Hanks' Balanced Salt Solution (HBSS) (modified without Calcium and Magnesium) (Hyclone
Gibco, SH3003102). Brains were excised by cutting the skull open at the midline, from base of the head towards
the front. Two lateral incisions were made under the cerebral cortices and skull was peeled off, meninges were
removed and individual cortices were transferred to HBSS containing 8 µl Trypsin (25g/L in 0.9 % NaCl) (Sigma,
T4549), incubated for 20 min at 37 ☐C with agitation. 10.7 µl of Trypsin inhibitor in neurobasal medium was
added to stop trypsinization, and 13.34 µl DNase1 (10 mg / ml) (Boehringer Mannheim Roche Diagnostics,
10137100) to degrade extracellular DNA medium were added to the solution. Cells were pelleted at 1000 x, for 5
min at 4 °C and 10.7 µl of trypsin inhibitor and 13.34 µl of DNase were added. Cells were triturated; viable cells
were counted using sterile filtered trypan blue solution (0.4 %) (Sigma, T8154). (Number of cells per ml was
calculated by: cell count in the central square x dilution factor x 104). Cells were diluted to desired concentration
for culture, in Complete Neurobasal Medium containing: Neurobasal medium (Gibco LifeTechnologies, 21103-
049), B27 supplement (Invitrogen, 17504-044), N2 supplement (Invitrogen, 17502-048), L-Glutamine 200 mM
(Gibco, 25030-081) and Penicillin-Streptomycin. Primary cortical neuron cells were plated into 24 or 6 well plates.
Plates were coated with Poly-D-Lysine-Hydrobromide (1 mg/ml, sterile) (Sigma, P0899) prior to use. Poly-D-
Lysine incubation was performed for at least 30 min and then plates were washed. For immunostaining, sterilized
coverslips were placed into 24 well culture dishes and then coated with poly-D-lysine. Cultures were incubated at
37 °C in a 5% CO_2 atmosphere and fixed or lysed at varied times.

Expression Vectors
Plasmids for transfection: GFP subcloned into pCig2 vector (generously gifted by Carola Sherman), FLAG-
Tagged: WT- PFTK1 and D228N- PFTK1 cDNA (mutation of conserved aspartic acid residue to asparagine, by a
point mutation) (1284 kb) subcloned into pCMV vector (generously gifted by Dr. Paul Albert and Maribeth
Lazzaro) (Lazzaro MA, 1997); GST-Tagged: WT-Rac1, G12V-Rac1, S17N-Rac1 subcloned in pEBG vector; GST-
tagged: WT-Cdc42, G12V-Cdc42, S17N-Cdc42 subcloned in pEBG vector and MYC-tagged: WT-RhoA, CA-
RhoA Q63L, DN-RhoA L19N in pRK5 vector. Rho GTPases (generously gifted by Dr. Gary Bokoch and Dr.
Margaret Chou, University of Pennsylvania). Constructs were sequenced at Ottawa Genomic Center, using
following primers: (1) For Rac1 and Cdc42 constructs in the pEBG backbone: (pGEX 5' Sequencing Primer: 5'-
[GGGCTGGCAAGCCACGTTTGGTG]-3'   and   pGEX   3'   Sequencing   Primer   5'-
[CCGGGAGCTGCATGTGTCAGAGG]-3') (2) To sequence PFTK1, CMV virus backbone: (pCMV 5' Sequencing
Primer: 5'-CGCAAATGGGCGGTAGGCGTG-3'   PFTK1   GC   rich   region   at   3'   end: 5'-
GGGGGAGTTGAAGCTGGCAG-3'). Sequencing was performed 100 bp upstream of start codon and 100 bp
downstream of stop codon. Plasmids were transformed with Top10 competent cells. Transformed cells were
cultured on Ampicillin resistant agar plates (LB Agar Granulated, Molecular Genetics BP-1423-500). A single
colony was picked and grown in Luria-Bertani (LB) Broth (Miller Amresco, J-106-500G). DNA was purified using
the Pure Yield Plasmid Midi Prep System (Promega, A2495).

Transient Transfection with Lipofectamine

HEK 293T Cells were cultured to 70% confluency, then co-transfected with Flag tagged (WT- PFTK1 or D228N-

PFTK1) and GST tagged [WT-Rac1, G12V-Rac1 (constitutively active), S17N-Rac1 (dominant negative inactive)]

or GST tagged [WT-Cdc42, G12V-Cdc42 (constitutively active), S17N-Cdc-42 (dominant negative inactive)] or

GST tagged [WT-RhoA, Q63L- RhoA (constitutively active), L19N-RhoA (dominant negative inactive)], as well as,

the reporter vector (GST::PEGB, MYC::PRK5, Flag::CMV using lipofectamine 2000 reagent (1mg/ml) (Invitrogen,

11668-019). Opti-MEM (Minimal Essential Medium), 1x Reduced Serum (Gibco, 31985-070) was used to dilute

the plasmid and lipofectamine according to the manufacturer's instructions. Diluted plasmid and lipofectamine

were mixed and incubated for 20 minutes at room temperature. Cell culture medium was changed to media

without FBS at the time of transfection. The plasmid-lipofectamine complex was added to culture media in

droplets. 24 hours post transfection, cells were triturated off culture plates by scraping, cells were collected into

ice cold PBS and spun down at 1.4 x g for 1 min for pull down assay.

Viral Infection

Gene delivery via Adenoviral (AV) infection was performed at the time of plating. Adenovirus expressing WT-

PFTK1 (Ad ET-CMV-3x Flag-WT- PFTK1, Pfu: 1.00 x 1012 CsCl Pure, Plaque pure), D228- PFTK1 (Ad ET-CMV-

3x Flag-D228N- PFTK1, Pfu: 9.20 x 1011 CsCl Pure, Plaque pure), and GFP (Ad ET-CMV-EGFP, Pfu: 4.89 x

1011 CsCl Pure, 3x Plaque pure), was mixed with neuronal cortical culture cells at 50 -100 MOI (multiplicity of

infection). Cells were fixed or lysed, at different time points, post-culture, for further analysis, depending on the

nature of the experiment.

Neuronal Death Assay

Cortical culture cells from mouse embryos at E13.5-14.5 were seeded at (1 x 105) in 96 well plates and

infected/co-infected with AV viruses (GFP alone or GFP + WT- PFTK1, D228N- PFTK1) at MOI=100 at the time

of plating, for survival assay. Cells were washed with 1X PBS and then fixed with 4% PFA 12, 16 and 24 hours

post-infection and then stained with Hoechst 33258 (1:10000) (Sigma-Aldrich B2883) in 1x PBS and incubated for

the minimum of 30 min for nuclei staining. Intact nuclei were counted against condensed and fragmented nuclei to

determine live neurons versus dead ones. To determine survival the number of live GFP positive neurons were

assessed in each well, each treatment was performed in triplets, and the average for each treatment was

calculated. Survival was assessed for WT- PFTK1 or D228N- PFTK1 as the percent of live GFP positive neurons

in comparison to GFP. Survival percentage represents the ratio of GFP expressing neurons with morphologically

intact nuclei at (12, 16, and 24 h) in WT- PFTK1 or KO- PFTK1 to the total number of GFP expressing neurons.

The values were normalized to GFP alone as control. Data are presented as (Mean± SEM) of at least three

independent experiments.

Total Protein Extraction

To compare protein expression levels, whole brain lysates were prepared from mouse brains at E13.5-14-5 or

from mouse pups at P21. Basically, mouse brains were dissected and washed with ice-cold PBS. Brains were

homogenized in lysis buffer: [50mM Tris-HCl (pH=7.5), 150mM NaCl,1 mM MgCl2, 1mM EDTA, 1% triton X-100

(Boehringer Mannheim, 789704), 1mM L-Dithiothreitol (DTT , C4H10O2S2) (Sigma Aldrich, D0632-5G), protease

inhibitor (Halt Protease inhibitor Cocktail EDTA-free 100x (Roche, 87785)] for 20 min, at 4 °C, with agitation.

Lysates were clarified by centrifugation at 19500 x g, 4 °C, for 20 min. Supernatant was collected (approximately

2mg of protein was collected per brain), mixed with 0.1% w/v bromophenol blue/ DTT sample buffer and heated at

100°C, for 10 min.

Immunoprecipitation (IP)

All steps were performed on ice. Cell lysate was prepared as described, previously. Lysate was prepared using

lysis buffer 50 mM Tris-HCl (pH=7.5), 150 mM NaCl, 1 mM MgCl2, 1mM EDTA, 1% Triton X-100 (Boehringer

Mannheim, 789704), 1mM DTT (L-Dithiothreitol, C4H10O2S2) (Sigma Aldrich, D0632-5G), protease inhibitor, for

20 min, at 4°C with agitation. Lysates were clarified by centrifugation at 19500 x g, 4°C, for 20 min. Supernatant

was collected and incubated with 20 µl of washed bead per reaction. After immunoprecipitation, beads were

pelleted at 1000 x g , washed five times with wash buffer (Tris-HCl (pH≈7.3-7.5), NaCl, EDTA, Triton 10 % ,

DTT), then mixed with 0.1 % w/v bromophenol blue / DTT sample buffer, and heated at 100°C, for 10 min to

release the protein complex. Proteins were resolved on 12 % SDS-polyacrylamide gels.

For immunoprecipitation of ectopically expressed GST tagged proteins, GST beads - (Glutathione Sepharose 4b

beads, particle size 45 µm - 165 µm, GE Healthcare 17-0756-05) were incubated with the lysate overnight, at 4

°C, on a rotor. For immunoprecipitation of endogenous protein, Trueblot Anti Rabbit IgIP beads - (eBioScience,

00-8800-25) or Anti mouse IgIP beads - (eBioScience, 00-8811-25) were used, depending on the origin of the bait

antibody. Lysate was collected and precleared by incubating 20 µl of the washed bead that was to be used for IP,

for 30 min, at 4 °C, on a rotor. Supernatant was cleared by centrifugation at 1000 x g, 1 min, at 4°C. 4 - 8 µg of

antibody per reaction, normal mouse IgG (SantaCruz, sc-2025) or normal rabbit IgG (SantaCruz, sc-2027) and

Pftaire-1 (H-140) (1:500) rabbit polyclonal IgG (SantaCruz, sc-50475), RhoA (1:500) mouse monoclonal IgG

(SantaCruz, sc-418), Rac1 (C-11) (1:500) rabbit polyclonal IgG (SantaCruz, sc-95), Cdc42 (1:500) rabbit

polyclonal IgG (SantaCruz, sc-87) was incubated with the cleared supernatant, overnight, at 4°C, on a rotor.

PFTK1 was used to IP GTPase and vice versa. 20 µl of the washed beads were incubated with the antibody-

lysate mix for 3 hours, at 4 °C, on a rotor to precipitate the antibody.

SDS-PAGE and Western Blot

Protein extracts were resolved on a 10 or 12 % SDS - polyacrylamide gel, along with Molecular Marker, Blueye

Prestained Protein Ladder (GeneDirex, P14007-0500). Transfers were done on 0.45 µm PolyVinylidene

DiFluoride (PVDF) Transfer membrane (Immobilon-P, IPVH00010). Membrane was blocked in 10 % non-fat dry

milk in 1 x PBST [Phosphate Buffered Saline (pH=8.0) and 0.1 % Tween] at room temperature, for 1 hour.

Proteins were detected using antibodies. Primary antibodies were diluted in [3 % Bovine Serum Albumin (BSA)

fraction V (Fisher, BP1600-100) in PBST and 0.1 % Sodium Azide NaN3 (Sigma Ultra, s-8032)] as follows:

PFTK1 (H-140) (1:500) rabbit polyclonal IgG, (SantaCruz, sc-50475), Pftaire2 (CDK15) (1:500) rabbit polyclonal

(Life Span Biosciences, LS-C119219/34326), RhoA (1:500) mouse monoclonal IgG (SantaCruz, sc-418), Rac1

(C-11) (1:500) rabbit polyclonal IgG (SantaCruz, sc-95), Cdc42 (1:500) rabbit polyclonal IgG (SantaCruz, sc-87),

Cdk5 (C-8) (1:500) rabbit polyclonal IgG (SantaCruz, sc-173), Pctaire1 (CDK16) (1:500) rabbit polyclonal, (Life

Span Biosciences, LS-C112292/34325), Anti-Flag (1:500) rabbit monoclonal IgG (Sigma, F7425), Anti-GST

(1:500) mouse monoclonal IgG (SantaCruz, sc-138) and incubated with the membranes overnight, at 4 °C, with

agitation. Membranes were washed and then incubated with secondary antibodies diluted in 5 % non-fat milk, for

1 hour, at room temperature [goat anti-mouse 170-6516 or goat anti-rabbit 170-6515 IgG (H+L) HRP conjugate

(Bio-Rad)] were all used at 1:2000 in 3 % BSA in 0.1 % Tween 20 (PBS/T). Immunoreactivity was detected using

Immobilin Western Chemiluminescence HRP substrate (Millipore, WBKLS0500) or Pierce ECL Western Blotting

Substrate (Thermo Scientific, 32106) and visualization HyBlot CL Autoradiography film, Denville Scientific. Protein

levels were normalized against β-actin signals, and results were reported in reference to control values (untreated

control for each individual experiment).

PFTK1 Kinase Assay

Kinase assay was performed using immunoprecipitated WT- PFTK1 protein or IgG as a control. Briefly, Neuronal

cortical cells prepared from WT- PFTK1 mouse embryos E13.5-14.5 were cultured in Neurobasal media (Gibco,

21103-049) containing B27 (Invitrogen, 17504-044) and N2 (Invitrogen, 17502-048), in 12 well plates, at the ratio

of 1 embryo per well, for 18-20 hour, at 37°C, 5 % CO_2. After washing with ice-cold 1x PBS, lysates were

collected directly from plates by scraping into ice-cold immunoprecipitation (IP) buffer (Tris-HCL 25mM (pH 7.4) ,

Na4P2O7 10 mM, Na3VO4 25 mM, NaF 50 mM, EDTA 1 mM , Digitonin 2 mg/ml) and sonicated twice, for 3

cycles, with a 20 minute incubation in between the 2 set of cycles, and then supernatant was cleared by

centrifugation at 14000x g, 4°C, for 10 min. 1 mg of protein(adjusted to the volume of 300 µl was incubated with

4 µg antibody either Normal rabbit IgG (200 µg/0.5 ml) or PFTK1 (200 µg/ml) antibodies, overnight, at 4°C, on the

shaker. IP-ed, WT- PFTK1 and IgG were mixed with 40 µl of pre-equilibrated (performed washes with IP wash

buffer) Trueblot anti-rabbit IgIP beads per reaction and incubated for 2 hours, at 4°C, on the shaker. Beads were

then washed and pelleted at 2000rpm, for 1 min, at 4°C, with IP wash buffer and kinase assay buffer [HEPES 50

mM (pH 7.5) and 1mM DTT], respectively. Next, kinase reaction buffer (HEPES 50 mM (pH 7.5), 1 mM DTT, 10

mM MgCl, and 20 µM ATP)) was added to a final volume of 25 µl. 10 µg of RhoA substrate His-Recombinant

human RhoA (Prospec, pro-057), was added to reactions with or without (5 µl, 100 x) GDP. Reactions were

incubated overnight, at 30°C. Samples were then centrifuged at 2000 rpm, for 1 min, at 4°C, then supernatant

was collected and subjected to either western blot for rabbit polyclonal IgG phosphoserine (Invitrogen, 61-8100)

signal; or G-LISA assay to assess RhoA activity as described below.

Assessment of Rho GTPase Activity

Activation of RhoA, as a function of PFTK1 kinase activity was determined using commercially available "RhoA

Activation Assay Kit (CellBioLabs, STA-403-A)" according to the manufacturer's instructions. Primary Cortical

Culture Cells were prepared from Het x Het crosses at embryonic day 13.5-14.5, at the density of 1 embryo per

well, in 1.5 ml, cultured in complete Neurobasal media supplemented with B27 and N2, in 12 well plates, and

incubated at 37 °C and 5 % CO_2. 18-20 hours post-plating cells were washed with ice-cold PBS and GTPase

assay was performed according to the manufacturer's kit. To decrease rapid hydrolysis of active GTP bound to

inactive GDP bound protein all following procedures were performed on ice. Briefly, cells were incubated on ice,

for 20 minutes, with 1x Assay / lysis buffer provided with the kit (Part No. 240102) (containing 125 mM HEPES,

pH 7.5, 750 mM NaCl, 5% NP-40, 50 mM MgCl2, 5 mM EDTA, 10% Glycerol) and then collected by scraping.

Due to very low expression of RhoA same-genotype extracts were pooled to achieve large amounts of Protein.

Total protein was quantified by Bradford protein assay to achieve at least 800ug of protein for each sample/assay.

Cell lysates were clarified by centrifugation at 14,000 g x, at 4 °C, for 10 min. (at least 800 µg protein per

reaction). 100x GTPγS (Part No. 240103) and GDP (Part No. 240104) were added to the cells for positive and

negative control, respectively. Control samples were incubated for 30 minutes, at 30 □C, with agitation, and the

reaction was terminated by adding 1M MgCl2 to the samples. Volume of sample per reaction was adjusted to be

identical in all reactions and 40 µl of bead added per reaction. Active proteins were immunoprecipitated with

respective rhotekin RBD agarose beads for RhoA. Beads were pelleted by centrifugation at 14000 x g, for 10

seconds, and washed three times with 1x Assay buffer supplied in the kit. Afterwards, beads were resuspended in

SDS-PAGE sample buffer and boiled for 5 minutes for detection by western blotting. Separated proteins were

visualized using RhoA, Mouse Monoclonal antibody (Part No. 240302) provided with the kit.

To quantify Rho GTPase activity after the PFTK1 Kinase assay, in a more accurate/sensitive manner, G-LISA

RhoA activation assay biochem kit (Absorbance based) (Cytoskeleton, BK124) was used according to the

manufacturer's protocol. Samples from the kinase reactions, (IgG-IP: RhoA, IgG-IP: RhoA+GDP, PFTK1-IP:

RhoA and PFTK1-IP: RhoA+GDP) were diluted and pre-equalized with ice cold lysis buffer /binding buffer (Part

#GL37). To pre-equalize samples, protein concentration was determined at 600nm, using Precision RedTM

Advanced Protein Assay Reagent (Part # GL50) as suggested by the manufacturer. Same buffer mix (no protein)

and RhoA (provided by manufacturer), were respectively, used as blank and positive control for the assay. Next,

samples were transferred to 96 well Rho-GTP binding plates (Part #GL25), equipped with pre-washed GLISA

assay strips, incubated at 4°C, on an orbital microplate shaker at 400 rpm, for 30min. Samples were then washed

with wash buffer (Part # GL38) and incubated with Antigen Presenting Buffer (Part#GL45) for 2 mins, washed

again with wash buffer, then incubated at room temperature for 45 min, on orbital microplate shaker at 400rpm,

with anti-RhoA primary antibody (Part # GL01) (1:250) in Antigen dilution buffer (Part #GL40) and washed

subsequently. Samples were next incubated with (1:62.5) secondary horseradish peroxidase (HRP) conjugate

antibody (Part #GL02) at room temperature, for 45 min, on an orbital microplate shaker at 400 rpm and washed

afterwards. Then samples were incubated with HRP detection reagents A (Part # GL43) and B (Part # GL44) (1:1

ratio) for 15 min, at 37 °C, and reaction was stopped by adding HRP Stop Buffer (Part #GL80), and absorbance

measured at 490 nm using a microplate spectrophotometer. To measure RhoA activity, values were determined

as follows, $\left(\frac{\text{GLISA value(ng)}}{\text{ELISA value(ng)}} \times 100 = \text{normalized \% of active RhoA}\right)$. The detected signal was normalized against total

RhoA using G-LISA and E-LISA (BK#150) kits side by side. To determine if any significant difference existed in

the activity of RhoA among samples, two-way ANOVA was performed in the grouped data followed by

Bonferroni's post-hoc test. Results were presented as average of at least three independent experiments.

Brain Sectioning

Brains were collected from the Wt- PFTK1 and KO- PFTK1 mice at 8 weeks of age. Animals were humanely

anesthetized with euthasol and transcardially perfused with 0.9 % saline, followed by fixative solution 4 %

Paraformaldehyde (PFA) (EMD, PX0055-3) in 1x PBS at pH=7.4. Brains were then removed from the skull and

fixed further with 4 % PFA, at 4 °C, for 24 hrs. Fixed brain tissue was cryopreserved in 20 % sucrose in 0.1 M

phosphate buffer and stored at 4°C, over a 3-day course, and the solution was changed 3 times per day. Brains

were then snap-frozen with Nitrogen balanced with $CO2$ and then embedded with O.C.T. (Tissue - Tek /O.C.T

compound, 4583) for sectioning. 14 µm coronal sections were collected from the brains; starting at the frontal

cortex at Bregma: ~ 2.710 mm up to the 4th ventricle at Bregma: ~ - 4.20 mm. For collecting, 3 sections were

skipped in between each selection. Sections were directly mounted onto SuperFrost Plus microscopic slides 25 x

75 x 1.0 mm (Fisher brand, 12-550-15) and subjected to2 % cresyl violet staining as described below.

Cresyl Violet Staining

Sections were thawed on slide warmer for 15-20 min, at 35 °C, and then placed in distilled water for 5 min.

Sections were then stained with 2 % filtered cresyl violet (pH=3.5) cresyl violet acetate ($C18H15N3O3$, Sigma

C5042-10G), for 10-30 minutes depending on the stain concentration. Sections were then dehydrated in graded

ethanol solutions from 50 % to 100 %, followed by defatting with xylene. Sections were mounted with coverslips

using toluene.

Immunostaining

Cells plated on glass coverslips were fixed in 4 % PFA in 1 x PBS, for 15 min. Fixed cells were permeabilized and

blocked simultaneously in 0.1 % Triton X-100, and 5 % BSA in 1x PBS, respectively, for 20 min, at room

temperature. Coverslips were incubated with primary antibodies diluted in 5 % BSA in 1x PBS: Tau1 clone 46

(1:500) mouse monoclonal IgG (Sigma-Aldrich, T-9450), GFP (H-140) (1:500) mouse monoclonal IgG (Abcam,

ab1218), PFTK1 (H-140) (1:500) rabbit polyclonal IgG (SantaCruz, sc-50475), MAP2 (H-300) (1:500) rabbit

polyclonal IgG (SantaCruz, sc-20172) for 1hr, at room temperature. Coverslips were washed; incubated with

FITC-secondary antibodies: Alexafluor 488 (1:1000) Goat-Anti-mouse (Molecular Probes, A11001), Streptavidin

Alexafluor 488 (1:1000) Goat-Anti-mouse (Molecular Probes, S11223), Streptavidin Alexafluor 546 (1:1000) Goat-

Anti-mouse (Molecular Probes, S11225), Alexafluor 594 (1:1000) Goat-Anti-mouse (Molecular Probes, A11005),

Alexafluor 594 (1:1000) Goat-Anti-rabbit (Molecular Probes, A11037), Alexafluor 594 (1:1000) Donkey-Anti-Goat

(Molecular Probes, A11058) were diluted in 5% BSA in 1x PBS, and then incubated with coverslips for 30 min, at

room temperature. To detect intact nuclei, cells were stained with Hoechst 33258 (1:10000) (Sigma-Aldrich

B2883) in 1x PBS, incubated for the minimum of 30 min, at room temperature. Slides were immersed in distilled

water to remove excess salt and then mounted on slides applying approximately 25 µl of Vectashield Mounting

Medium (Vector laboratories, H-1200).

Microscopy and Imaging

Images were taken using upright Zeiss Axioskop 2 Mot or invert Zeiss AxioObserver-Z1 fluorescent microscope
with Northern Eclipse Software or Zeiss and Axiovision Rel. 4.8 software, captured in XY mosaic format, in
rectangle mode. Images were taken at 20x, 32x, and 40 x magnification lenses. To determine the total diameter of
the field of view for each objective, Ocular Field of View (F.O.V.) number (32) was divided by objective
magnification and following measurements were obtained: for 20x (1500 µm), for 32x (719 µm), and for 40x (575
µm). Images were processed for length measurements using Axiovision software Rel. 4.8, spline mode to trace
axon length or Stereo Investigator microscopy imaging system (version 6; MicroBrightField, Williston, VT)
software using contour mapping mode and tracing the length of axons. The distance from the cell body of the
GFP expressing neuron or Tau1 to the distal region of the longest neurite was measured as axon/lengthiest
neurite.

Quantification of Signal Density on Images
Immunoblots were further analyzed by Image J software to compare the density/intensity of the bands of an agar
gel or a western blot membrane. Following imaging of the blots, representative bands were selected, and their
intensity was quantified. Then quantified measurements were transferred to excel and results were normalized
against relative controls (β-actin for western blot and S12 for RT-PCR).

16
17

References

Ashburner, M. (1989). Drosophila: A Laboratoty Manual (Cold Spring Harbor Press).

Benveniste, R.J., Thor, S., Thomas, J.B., and Taghert, P.H. (1998). Cell type-specific regulation of the Drosophila FMRF-NH2 neuropeptide gene by Apterous, a LIM homeodomain transcription factor. Development *125*, 4757-4765.

Besset, V., Rhee, K., and Wolgemuth, D.J. (1998). The identification and characterization of expression of Pftaire-1, a novel Cdk family member, suggest its function in the mouse testis and nervous system. Mol Reprod Dev *50*, 18-29. 10.1002/(SICI)1098-2795(199805)50:1<18::AID-MRD3>3.0.CO;2-#.

Bettencourt-Dias, M., Giet, R., Sinka, R., Mazumdar, A., Lock, W.G., Balloux, F., Zafiropoulos, P.J., Yamaguchi, S., Winter, S., Carthew, R.W., et al. (2004). Genome-wide survey of protein kinases required for cell cycle progression. Nature *432*, 980-987.

Bhat, K.M. (1999). Segment polarity genes in neuroblast formation and identity specification during Drosophila neurogenesis. Bioessays *21*, 472-485.

Bhat, K.M., and Schedl, P. (1997). Requirement for engrailed and invected genes reveals novel regulatory interactions between engrailed/invected, patched, gooseberry and wingless during Drosophila neurogenesis. Development *124*, 1675-1688.

Campos-Ortega, J., and Hartenstein, V. (1985). The Embryonic Development of Drosophila melanogaster. (Springer Verlag).

Casso, D., Ramirez-Weber, F., and Kornberg, T.B. (2000). GFP-tagged balancer chromosomes for Drosophila melanogaster. Mech Dev *91*, 451-454.

Chang, F., Lemmon, C., Lietha, D., Eck, M., and Romer, L. (2011). Tyrosine Phosphorylation of Rac1: A Role in Regulation of Cell Spreading. PLoS ONE *6*, e28587. 10.1371/journal.pone.0028587.

Chen, Y., Yang, Z., Meng, M., Zhao, Y., Dong, N., Yan, H., Liu, L., Ding, M., Peng, H.B., and Shao, F. (2009). Cullin Mediates Degradation of RhoA through Evolutionarily Conserved BTB Adaptors to Control Actin Cytoskeleton Structure and Cell Movement. Molecular Cell *35*, 841-855. 10.1016/j.molcel.2009.09.004.

Chotard, C., and Salecker, I. (2004). Neurons and glia: team players in axon guidance. Trends Neurosci *27*, 655-661.

Cole A R (2009). PCTK Proteins:The Forgotten Brain Kinases? Neurosignals *17*, 288-297.

Cook, H.A., Koppetsch, B.S., Wu, J., and Theurkauf, W.E. (2004). The Drosophila SDE3 homolog armitage is required for oskar mRNA silencing and embryonic axis specification. Cell *116*, 817-829.

Davidson, G., Shen, J., Huang, Y.L., Su, Y., Karaulanov, E., Bartscherer, K., Hassler, C., Stannek, P., Boutros, M., and Niehrs, C. (2009). Cell cycle control of wnt receptor activation. Dev Cell *17*, 788-799.

Desai, C.J., Popova, E., and Zinn, K. (1994). A Drosophila receptor tyrosine phosphatase expressed in the embryonic CNS and larval optic lobes is a member of the set of proteins bearing the "HRP" carbohydrate epitope. J Neurosci *14*, 7272-7283.

Duffy, J.B. (2002). GAL4 system in Drosophila: a fly geneticist's Swiss army knife. Genesis *34*, 1-15.

Fambrough, D., Pan, D., Rubin, G.M., and Goodman, C.S. (1996). The cell surface metalloprotease/disintegrin Kuzbanian is required for axonal extension in Drosophila. patent application 23, Nov 12.

Flatau, G., Lemichez, E., Gauthier, M., Chardin, P., Paris, S., Fiorentini, C., and Boquet, P. (1997). Toxin-induced activation of the G protein p21 Rho by deamidation of glutamine. Nature *387*, 729-733. 10.1038/42743.

Freeman, M.R. (2006). Sculpting the nervous system: glial control of neuronal development. Curr Opin Neurobiol *16*, 119-125.

Garbe, D.S., and Bashaw, G.J. (2004). Axon guidance at the midline: from mutants to mechanisms. Crit Rev Biochem Mol Biol *39*, 319-341.

Garcia-Mata, R., Boulter, E., and Burridge, K. (2011). The 'invisible hand': regulation of RHO GTPases by RHOGDIs. Nature Reviews Molecular Cell Biology *12*, 493-504. 10.1038/nrm3153.

Grigoryan, T., Wend, P., Klaus, A., and Birchmeier, W. (2008). Deciphering the function of canonical Wnt signals in development and disease: conditional loss- and gain-of-function mutations of beta-catenin in mice. Genes Dev *22*, 2308-2341.

Grosskortenhaus, R., Pearson, B.J., Marusich, A., and Doe, C.Q. (2005). Regulation of temporal identity transitions in Drosophila neuroblasts. Dev Cell *8*, 193-202.

Harlow, E., and Lane, D. (2006). Preparing Early Whole-Mount Drosophila Embryos for Immunostaining (Cold Spring Harbor Protocols. doi:10.1101/pdb.prot4524).

Heasman, S.J., and Ridley, A.J. (2008). Mammalian Rho GTPases: new insights into their functions from in vivo studies. Nature Reviews Molecular Cell Biology *9*, 690-701. 10.1038/nrm2476.

Hirose T, K.M., Tamaru T, Okumura N, Nagai K, Okada M (2000). Identification of tudor repeat associator with PCTAIRE 2 (Trap). A novel protein that interacts with the N-terminal domain of PCTAIRE 2 in rat brain. Eur J Biochem. *267*, 2113-2121.

Hodge, R.G., and Ridley, A.J. (2016). Regulating Rho GTPases and their regulators. Nature Reviews Molecular Cell Biology *17*, 496-510. 10.1038/nrm.2016.67.

Hortsch, M., Bieber, A.J., Patel, N.H., and Goodman, C.S. (1990). Differential splicing generates a nervous system-specific form of Drosophila neuroglian. Neuron *4*, 697-709.

Hummel, T., Krukkert, K., Roos, J., Davis, G., and Klambt, C. (2000). Drosophila Futsch/22C10 is a MAP1B-like protein required for dendritic and axonal development. Neuron *26*, 357-370.

Hummel, T., Schimmelpfeng, K., and Klambt, C. (1999). Commissure formation in the embryonic CNS of Drosophila. Development *126*, 771-779.

Isshiki, T., Pearson, B., Holbrook, S., and Doe, C.Q. (2001). Drosophila neuroblasts sequentially express transcription factors which specify the temporal identity of their neuronal progeny. Cell *106*, 511-521.

Jacobs, J.R., and Goodman, C.S. (1989). Embryonic development of axon pathways in the Drosophila CNS. II. Behavior of pioneer growth cones. J Neurosci *9*, 2412-2422.

Jiang, M., Gao, Y., Yang, T., Zhu, X., and Chen, J. (2009). Cyclin Y, a novel membrane-associated cyclin, interacts with PFTK1. FEBS Letters *583*, 2171-2178. 10.1016/j.febslet.2009.06.010.

Jimenez, F., and Campos-Ortega, J.A. (1987). Genes in subdivision 1B of the Drosophila melanogaster X-chromosome and their influence on neural development. J Neurogenet *4*, 179-200.

Jin, Z., and Strittmatter, S.M. (1997). Rac1 Mediates Collapsin-1-Induced Growth Cone Collapse. The Journal of Neuroscience *17*, 6256-6263. 10.1523/jneurosci.17-16-06256.1997.

Klambt, C., Jacobs, J.R., and Goodman, C.S. (1991). The midline of the Drosophila central nervous system: a model for the genetic analysis of cell fate, cell migration, and growth cone guidance. Cell *64*, 801-815.

Kozma, R., Sarner, S., Ahmed, S., and Lim, L. (1997). Rho family GTPases and neuronal growth cone remodelling: relationship between increased complexity induced by Cdc42Hs, Rac1, and acetylcholine and collapse induced by RhoA and lysophosphatidic acid. Mol Cell Biol *17*, 1201-1211. 10.1128/mcb.17.3.1201.

Kranenburg, O., Poland, M., Gebbink, M., Oomen, L., and Moolenaar, W.H. (1997). Dissociation of LPA-induced cytoskeletal contraction from stress fiber formation by differential localization of RhoA. J Cell Sci *110 (Pt 19)*, 2417-2427.

Kuhn, T.B., Brown, M.D., Wilcox, C.L., Raper, J.A., and Bamburg, J.R. (1999). Myelin and collapsin-1 induce motor neuron growth cone collapse through different pathways: inhibition of collapse by opposing mutants of rac1. J Neurosci *19*, 1965-1975.

Lazzaro, M.A., Albert, P.R., and Julien, J.P. (1997). A novel cdc2-related protein kinase expressed in the nervous system. J Neurochem *69*, 348-364.

Lazzaro MA, A.P., Julien JP (1997). A novel cdc2-related protein kinase expressed in the nervous system. . J Neurochem. *69*, 348-364.

Lazzaro, M.A., and Julien, J.P. (1997). Chromosomal mapping of the PFTAIRE gene, Pftk1, a cdc2-related kinase expressed predominantly in the mouse nervous system. Genomics *42*, 536-537.

Liu, J., and Kipreos, E.T. (2000). Evolution of cyclin-dependent kinases (CDKs) and CDK-activating kinases (CAKs): differential conservation of CAKs in yeast and metazoa. Mol Biol Evol *17*, 1061-1074.

Malumbres, M., and Barbacid, M. (2005). Mammalian cyclin-dependent kinases. Trends Biochem Sci *30*, 630-641. 10.1016/j.tibs.2005.09.005.

Matsuda, S., Kominato, K., Koide-Yoshida, S., Miyamoto, K., Isshiki, K., Tsuji, A., and Yuasa, K. (2014). PCTAIRE Kinase 3/Cyclin-dependent Kinase 18 Is Activated through Association with Cyclin A and/or Phosphorylation by Protein Kinase A. Journal of Biological Chemistry *289*, 18387-18400. 10.1074/jbc.m113.542936.

Mikolcevic P, S.R., Rauch V, Hess MW, Pfaller K, Barisic M, Pelliniemi LJ, Boesl M, Geley S (2012). Cyclin-dependent kinase 16 /PCTAIRE kinase 1 is activated by cyclin Y and is essential for spermatogenesis. Mol Cell Biol. *32*, 868-879.

Nambu, J.R., Franks, R.G., Hu, S., and Crews, S.T. (1990). The single-minded gene of Drosophila is required for the expression of genes important for the development of CNS midline cells. Cell *63*, 63-75.

Nikolic M, C.M., Lu W, Mayer BJ, Tsai LH (1998). The p35/Cdk5 kinase is a neuron-specific Rac effector that inhibits Pak1 activity. . Nature *395*, 194-198.

Nikolic M, D.H., Kwon YT, Ramos YF, Tsai LH (1996). The cdk5/p35 kinase is essential for neurite outgrowth during neuronal differentiation. Genes Dev. *10*, 816-825.

Okuda T, C.J., Downing JR (1992). PCTAIRE-1 and PCTAIRE-3, two members of a novel cdc2/CDC28-related protein kinase gene family. Oncogene. *7*.

Park, M.H., Kim, S.Y., Kim, Y.J., and Chung, Y.-H. (2014). ALS2CR7 (CDK15) attenuates TRAIL induced apoptosis by inducing phosphorylation of survivin Thr34. *450*, 129-134. 10.1016/j.bbrc.2014.05.070.

Parker, R.J., and Auld, V.J. (2004). Signaling in glial development: differentiation migration and axon guidance. Biochem Cell Biol *82*, 694-707.

Rashid T, B.M., Nikolic M (2001). Phosphorylation of Pak1 by the p35/Cdk5 kinase affects neuronal morphology. J Biol Chem *276*, 49043-49052.

Reichhart, J.M., and Ferrandon, D. (1998). Green balancers. Drosophila Information Service *81*, 201-202.

Robinow, S., and White, K. (1991). Characterization and spatial distribution of the ELAV protein during Drosophila melanogaster development. J Neurobiol *22*, 443-461.

Sanchez-Soriano, N., and Prokop, A. (2005). The influence of pioneer neurons on a growing motor nerve in Drosophila requires the neural cell adhesion molecule homolog FasciclinII. J Neurosci *25*, 78-87.

Sanchez-Soriano, N., Tear, G., Whitington, P., and Prokop, A. (2007). Drosophila as a genetic and cellular model for studies on axonal growth. Neural Develop *2*, 9.

Sauer, K., Weigmann, K., Sigrist, S., and Lehner, C.F. (1996). Novel members of the cdc2-related kinase family in Drosophila: cdk4/6, cdk5, PFTAIRE, and PITSLRE kinase. Mol Biol Cell *7*, 1759-1769.

Sebok A, N.N., Debreceni B, Guo Z, Santos MF, Szeberenyi J, Tigyi G (1999). Different roles for RhoA during neurite initiation, elongation, and regeneration in PC12 cells. J Neurochem *73*, 949-960.

Snow, P.M., Patel, N.H., Harrelson, A.L., and Goodman, C.S. (1987). Neural-specific carbohydrate moiety shared by many surface glycoproteins in Drosophila and grasshopper embryos. J Neurosci *7*, 4137-4144.

Stowers, R.S., Garza, D., Rascle, A., and Hogness, D.S. (2000). The L63 gene is necessary for the ecdysone-induced 63E late puff and encodes CDK proteins required for Drosophila development. Dev Biol *221*, 23-40.

Threadgill, R., Bobb, K., and Ghosh, A. (1997). Regulation of dendritic growth and remodeling by Rho, Rac, and Cdc42. Neuron *19*, 625-634. 10.1016/s0896-6273(00)80376-1.

Tong, J., Li, L., Ballermann, B., and Wang, Z. (2016). Phosphorylation and Activation of RhoA by ERK in Response to Epidermal Growth Factor Stimulation. PLOS ONE *11*, e0147103. 10.1371/journal.pone.0147103.

Weigmann, K., Klapper, R., Strasser, T., Rickert, C., Technau, G., Jäckle, H., Janning, W., and Klämbt, C. (2003). FlyMove – a new way to look at development of Drosophila. Trends Genet *19*, 310.

Xiong, W.C., Okano, H., Patel, N.H., Blendy, J.A., and Montell, C. (1994). repo encodes a glial-specific homeo domain protein required in the Drosophila nervous system. Genes Dev *8*, 981-994.

Yang, T., and Chen, J.Y. (2001). Identification and cellular localization of human PFTAIRE1. Gene *267*, 165-172. 10.1016/s0378-1119(01)00391-2.

Zechner, D., Fujita, Y., Hulsken, J., Muller, T., Walther, I., Taketo, M.M., Crenshaw, E.B., 3rd, Birchmeier, W., and Birchmeier, C. (2003). beta-Catenin signals regulate cell growth and the balance between progenitor cell expansion and differentiation in the nervous system. Dev Biol *258*, 406-418.

Zipkin, I.D., Kindt, R.M., and Kenyon, C.J. (1997). Role of a new Rho family member in cell migration and axon guidance in C. elegans. Cell *90*, 883-894. 10.1016/s0092-8674(00)80353-0.

A

	Control (GFP+)	KO (GFP-)
;;Df(3L)E1		
;;Eip63E⁸¹		
;;Df(3L)E1 Eip63E⁸¹		

Scoring Phenotype	Possible Genotypes	BP102-associated phenotype (embryos)		Penetrance
		Normal	Abnormal	
GFP+	;;Df(L63)E1/TM3krGFP; ;;TM3krGFP/TM3krGFP;	75.3% (208)	0.3% (1)	0.50%
GFP-	;;Df(L63)E1/Df(L63)E1;	0% (0)	24.4% (67)	100%

Scoring Phenotype	Possible Genotypes	BP102-associated phenotype (embryos)		Penetrance
		Normal	Abnormal	
GFP+	;;L63^{81}/TM3krGFP; ;;TM3krGFP/TM3krGFP;	76.0% (45)	0.0% (0)	0%
GFP-	;;L63^{81}/L63^{81};	3% (2)	21.0% (17)	89.5%

Scoring Phenotype	Possible Genotypes	BP102-associated phenotype (embryos)		Penetrance
		Normal	Abnormal	
GFP+	;;Df(L63)E1/TM3krGFP; ;;L63^{81}/TM3krGFP; ;;TM3krGFP/TM3krGFP;	74.0% (43)	0.0% (0)	0%
GFP-	;;Df(L63)E1/L63^{81};	14.0% (8)	12.0% (7)	47.0%

B

	Control (GFP+)	;;Df(L63)E1 (GFP-)	;;Eip63E^{81} (GFP-)
Fasciclin II			

Scoring Phenotype	Possible Genotypes	1D4-associated phenotype (embryos)		Penetrance
		Normal	Abnormal	
GFP+	;;Df(L63)E1/TM3krGFP; ;;TM3krGFP/TM3krGFP;	77.5% (107)	0.0% (0)	0%
GFP-	;;Df(L63)E1/Df(L63)E1;	0% (0)	22.5% (31)	100%

Scoring Phenotype	Possible Genotypes	1D4-associated phenotype (embryos)		Penetrance
		Normal	Abnormal	
GFP+	;;L63^{81}/TM3krGFP; ;;TM3krGFP/TM3krGFP;	78.5% (101)	0.0% (0)	0%
GFP-	;;L63^{81}/L63^{81};	1.5% (2)	20.0% (31)	93.9%

Figure 1. Eip63E deficiency leads to defasciculation and misguidance of VNC axons.
Ventral views of whole mount stage 14-15 embryos from both Eip63E mutant lines, co-stained for GFP and BPI 02 (A) or Fasciclin-II (1D4 mAb) (B). Penetrance of axon abnormalities was calculated as the % of embryos of a given scoring phenotype that show any defect from the total population of such phenotype (GFP+ or GFP- embryos). Arrowhead points to normal looking longitudinal connectives and arrows to disrupted structures in KO embryos. L63^{81}: Eip63E^{81}.

Figure 2. Eip63E deficiency leads to early embryonic developmental defects.
Ventral views (unless otherwise specified) of whole mount embryos are shown head to the left. Stage 5-8 embryos are from Eip63E[81]/TM3krGFP,Sb line are of unknown genotype. Panels for stages 10-11 show Eip63E[81] homozygous mutant embryos. (A) Staining for single-minded at early embryonic stages (st 5-10) show no gross difference in midline formation but cellular morphological differences are noticable. (B) Staining for engrailed/ invected (4D9 mAb). 18% (n=48) of total population of stage 8-9 embryos show acute segmentation defects. At stage 10-11, cell bodies appear disorganized (arrowheads) to the extent of loss of hemi-segments division at the midline (wavy arrow) compared to control (straight arrow). (C) Staining for the HRP antigen. In top 4 panels, a lateral view of embryos is shown. 10% (n=48) of st 8-9 embryos show premature HRP staining (arrowheads). At late stage 11, precocious axon outgrowth (arrows) seen in 36% of KO embryos (n=52).

Figure 3. Eip63E deficiency leads to early segmentation defects.
(A, B, C and D) Ventral views of whole mount embryos stage 12-15 shown head to the left. CTR show GFP+ embryos (wild-type and mutant heterozygous population) and Eip63E81 represent homozygous mutant embryos (GFP-). Immunostaining for Elav (A), Neuroglian (B), Neurotactin (C, top 2 panels) and BP102 (C, bottom 2 panels) showing axon defects (arrows). (D) Co-staining for glial populations (Repo) and axons (HRP) showing disorganized cell bodies (arrowheads). (E) Merged elav and HRP staining images (lateral view) of stage 12 embryos shown in panel A. Elav-positive cells appear to be dorsally deeper than normal (arrowheads), under the axon bundles. (F) VNC flat preparations of timed embryos (10-11h, stage 14) co-stained for Futsch and HRP are shown dorsal up, head to the left. Arrowheads point to some of the structures where neuronal organization is different from control embryos. Arrows point to examples of axon growth abnormalities

A	Control Stage 14	**E**	;;RepoGal4/UASRNAiL63	glial other than midline
B	;Act5Gal4/+;UASRNAiL63/+	ubiquitous	**F**	;GawB109.69Gal4/+;UASRNAiL63/+ dorsal epidermis, some PNS
C	;elavGal4/+;;UASRNAiL63/+ neurons	**G**	;;GawBtey5053AGal4/UASRNAiL63 dig system, longitudinal muscles	
D	;;mef2Gal4/UASRNAiL63 mesoderm, visceral muscles			

H

Genotype	N	%
;Act5Gal4/+;UASRNAiL63/+	72	59.7
;elavGal4/+;;UASRNAiL63/+	89	49.4
;;mef2Gal4/UASRNAiL63	78	17.9
;;RepoGal4/UASRNAiL63	18	0
;GawB109.69Gal4/+;UASRNAiL63/+	51	13.7
;;GawBtey5053AGal4/UASRNAiL63	36	8.3

Figure 4. Only ubiquitous and neuronal downregulation of Eip63E leads to embryonic axon defects.
Ventral views of stage 14 whole mount embryos shown head to the left. A UAS-Gal4 approach was used to drive Eip63E downregulation in specific tissues during embryonic development. Embryos negative for GFP (present in balancers for both the Gal4 and the UAS lines) was considered as experimental embryos vs the GFP positive of the same progeny. (A) A representative GFP+ embryo is shown. (B) Ubiquitous downregulation led to a replica of the germline deficiency lines, regarding CNS defective phenotype and developmental lethality (see text for details). (C) Neuronal-specific downregulation led to a robust and highly penetrant axonal phenotype. (D, E, F, G) Downregulation of Eip63E in other tested tissue (indicated in red) did not cause any detectable axonal defects. The number of embryos per genotype and the percentage of embryos per group showing axonal defects is summarized in (H).

Figure 5. Eip63E functionally interacts with Rho1 to regulate axogenesis. Ventral views of stage 14 whole mount embryos stained for BP102 antigen are shown head to the left. Panels show representative images for each specified genotype, (A-D) Representative images of control embryos (A) single hetero-zygous Eip63E[81] (B) or Rho1[72F] (C) from a cross between Eip63E[81] KO line and Dominant-Negative Rho1 (Rho1[72F]). (B, C) Images of presumed genotype (see text for details). (D) Axon defects of intermediate severity in 22.7% of double-heterozygous mutants (GFP-). (E-H) UAS-elavGal4 approach to drive Eip63E downregulation and constitutive active Rho1 overexpression in neurons in two independent Dmel lines. Lines were then crossed to generate the indicated genotypes. (F, G) Heterozygous neuronal downregulation of Eip63E or Rho1 showing axon defects with specified penetrance. 86.96% of Double-heterozygous embryos (H) had axon defects showing features of both heterozygous controls. Penetrance values from stages 13-17, n as follows: B) >100, C) 86, D) 22, F) 89, G) 126, H) 69. n represents the total of screened embryos from several different pairings/collection pools.

Figure 6. PFTK1 (CDK14) downregulation leads to faster-growing axons in murine cortical neurons.
Primary cortical neurons derived from CD1 embryos (E 14-15) were infected with 100 MOI of adenoviruses expressing GFP alone, GFP and WT-PFTK1 or GFP and D228N-PFTK1. After 12 or 24 hours in culture, cells were fixed and stained with Hoechst 33258 (nuclei visualization). Axons of GFP+ neurons were measured using Image J. (A) Values are presented as averages ±SEM from 122-143 cells/treatment/timepoint. Statistics were done using two-way ANOVA followed by Tuckey's post-hoc. (B). Same data as in (A) but presented as axon length relative to the GFP control. (C) Representative images of primary cortical neurons from E13-14 embryos from HET x HET crosses from WT and KO cultures fixed at the specified times and stained for Tau and Hoechst. (D) Axon length distribution per genotype from the indicated number of embryos (236-469 cells/embryo) at 16 hours in vitro. Each column represents the percentage of cells in the specified range of length for the indicated genotype. Non-treated (NT) cultures were also fixed after 16 and 24h in vitro. Graph represents the average axon length/genotype/treatment. t-test was used for statistical analysis whenever p value is specified. (E) 16h cultures were treated for the last 8h with 0.5ug/ml of RhoA activator (Cytoskeleton Inc. CN03). Graph represents the average axon length treatment.

Figure 7. RhoA phosphorylation and activity is affected by PFTK1.
Primary neuronal cultures from E13-14 embryos from HET x HET crosses were examined after 18-20h in vitro. Total RNA (A) or protein (B) was extracted followed by RT-PCR or western blot. Representative images from RT-PCR (A) and western blot (B) assays are shown. (C) Band intensity was assessed by densitometry using ImageJ and RhoA signal was standardized relative to loading control (S12 or Actin). Bar graph represents the average ±SEM from 3 KO and 6 WT for mRNA and 7 WT and KO each for protein. t-test was used for statistical analysis. D) Assessment of RhoA activity using an IP-based assay. Neuronal extracts were incubated with Rhotekin PBD beads to pull down active RhoA. WT extract preincubated with GTPγS (positive) or GDP (negative) were used as controls. IP was followed by western blotting for RhoA. Two representative assays of four independent assays are shown. E) In vitro Kinase assay. IPed protein (either with IgG control or PFTK1) from WT cortical neurons extracts was used as source of kinase and incubated with human recombinant RhoA as substrate (His-tagged) with or without GDP. Results were analyzed for phospho-serine signal. F) RhoA activity assay following in-vitro kinase assay. Kinase reactions were analyzed by G-LISA to test for RhoA activity. Bar graph represents mean values ±SEM. Two-way ANOVA analysis of grouped data and Sidak test for multiple comparisons are indicated on bar graph. RHO1: His-RhoA human recombinant protein, Cytoskeleton.