EnteroBase: Hierarchical clustering of 100,000s of bacterial genomes

into species/sub-species and populations

Mark Achtman*, Zhemin Zhou*,[a], Jane Charlesworth and Laura Baxter

University of Warwick, Coventry CV4 7AL, UK

[a]Current address: Pasteurien College, Soochow University, No.199 Ren'ai Road,

Suzhou, Jiangsu 215123, China.

*Corresponding authors: MA, m.achtman@warwick.ac.uk; ZZ, zheminzhou@qq.com

MA, ORCID 0000-0001-6815-0070; ZZ, ORCID 0000-0001-9783-0366; JC, ORCID

0000-0001-9759-9838; LB, ORCID 0000-0003-3287-7547

Word count: 7559

Word Count incl. 103 Citations 10846

Plus:

4 Tables

5 Figures

Abstract

The definition of bacterial species is traditionally a taxonomic issue while defining bacterial populations is done with population genetics. These assignments are species specific, and depend on the practitioner. Legacy multilocus sequence typing is commonly used to identify sequence types (STs) and clusters (ST Complexes). However, these approaches are not adequate for the millions of genomic sequences from bacterial pathogens that have been generated since 2012. EnteroBase (http://enterobase.warwick.ac.uk) automatically clusters core genome MLST alleles into hierarchical clusters (HierCC) after assembling annotated draft genomes from short read sequences. HierCC clusters span core sequence diversity from the species level down to individual transmission chains. Here we evaluate HierCC's ability to correctly assign 100,000s of genomes to the species/subspecies and population levels for *Salmonella, Clostridoides, Yersinia, Vibrio* and *Streptococcus*. HierCC assignments were more consistent with maximum-likelihood super-trees of core SNPs or presence/absence of accessory genes than classical taxonomic assignments or 95% ANI. However, neither HierCC nor ANI were uniformly consistent with classical taxonomy of *Streptococcus.* HierCC was also consistent with legacy eBGs/ST Complexes in *Salmonella* or *Escherichia* and revealed differences in vertical inheritance of O serogroups. Thus, EnteroBase HierCC supports the automated identification of and assignment to species/subspecies and populations for multiple genera.

1. Introduction

"Microbiologists need large databases to identify and communicate about clusters of related bacteria, … Such databases should contain the reconstructed genomes of bacterial isolates,… together with metadata describing their sources and phenotypic properties… and we are currently developing EnteroBase, a genome-based successor to the *E. coli* and *S. enterica* MLST databases." Excerpted from Achtman and Zhou, 2014 [1].

The Linnean system of genus and species designations was applied to bacterial nomenclature in the late 19th century, soon after their ability to cause infectious diseases had been recognised. These taxonomic labels initially reflected disease specificity and phenotypic similarities. Over the following decades, phenotypic criteria for distinguishing bacterial taxa were extended to include serologically distinct groupings and even differential sensitivity to bacteriophages. A primary goal for taxonomic designations for bacterial pathogens was that they be useful for epidemiology and clinical diagnoses. As a result, *Salmonella enterica* serovars, which differ by dominant epitopes on lipopolysaccharide and flagella, were each assigned a distinct species designation, often referring to the disease syndrome and host e.g. *Salmonella typhimurium* [2]. Similarly, even though *Yersinia pestis* is a clone of *Yersinia pseudotuberculosis* [3,4], it was designated as a distinct species because *Y. pestis* causes plague whereas *Y. pseudotuberculosis* causes gastroenteritis. The use of species designations for *Salmonella* serovars is now disparaged, although it is still in common use [5], but *Y. pestis* has retained its species designation.

Percentage DNA-DNA hybridization levels reflect genomic relationships and became the "Gold Standard" metric for new taxonomic designations after 1987 [6],

but distinctive phenotypic differences have remained a requirement for the definition

of a novel species until the present. As a result, the international committee which

controls taxonomic designations continues to reject the validity of taxonomic

designations based solely on DNA similarities [7,8].

a) **ANI and MLST**

An alternative genomic approach for taxonomic designations was proposed in

2005 by Konstantinidis [9], namely the definition of species on the basis of average

nucleotide identity (ANI). Pairs of genomes with 95% ANI usually belong to the same

species whereas pairs with lower ANI values belong to different species. Fast

computational methods based on k-mer searches, such as FastANI [10], can perform

ANI-like calculations on large numbers of genomes, and these calculations are now

routinely used by bioinformaticians for taxonomic assignments. An approach based

on ANI is very enticing because it could provide defined criteria for the definition of a

species across all Bacteria, and allow species assignment based exclusively on

genome sequences. However, the use of ANI for species assignments has been

criticized because it does not completely correlate with DNA-DNA hybridization

[11,12]. Furthermore, multiple taxa that are designated as single species each contain

multiple 95%ANI groups [13,14]. For example, *Streptococcus mitis* is designated a

single species because those strains colonise the human oropharyngeal tract and have

similar phenotypes. However, *S. mitis* encompasses myriad, genetically distinct

strains [15-17], and encompasses at least 44 distinct 95% ANI clusters [17]. Other

problematic genera include *Pseudomonas* [12,18] and *Aeromonas* [19]. Furthermore,

large databases including 100,000s of genomes per genus would struggle to

implement methods such as pairwise clustering by ANI because each single new entry

would require testing against all genomes. There is also no consensus on ANI criteria

for recognizing lower taxonomic entities, such as sub-species and populations.

Indeed, there is not even a consensus on the definitive properties of what constitutes a

bacterial species [20].

Bottom-up population genetic approaches such as multilocus sequence typing

(MLST) can provide an alternative to top-down taxonomy, and deal efficiently with

large numbers of bacterial strains. Legacy MLST based on the sequence differences

of several housekeeping genes was introduced in 1998 [21], and has now been applied

to more than 100 bacterial species [22]. MLST defines sequence types (STs),

consisting of unique integer designations for each unique sequence (allele) of each of

the MLST loci. Some STs mark individual clones with special pathogenic properties

and which seem to have arisen fairly recently, such as *Escherichia coli* ST131 which

is a globally prominent cause of urinary tract infection (UTI) and invasive disease

[23]. Similarly, *Salmonella enterica* subsp. *enterica* serovar Typhimurium ST313 is a

common cause of extra-intestinal, invasive salmonellosis in Africa [24]. Clusters of

related STs are also well known. Such clusters can be recognized by eBurst analyses

[25], and are referred to as ST Complexes in *E. coli* [26] and eBGs (eBurst groups) in

S. enterica [27]. ST Complexes and eBGs seem to reflect natural populations, but

their broad properties are still not well understood, in part because the resolution of

legacy MLST is fairly low.

The principles of legacy MLST were extended to rMLST, which uses sequences

of 53 universal bacterial genes encoding ribosomal proteins [28] and provides a

universal MLST scheme for all Bacteria [28]. For both *Escherichia* and *Salmonella,*

rMLST offers a slight improvement in resolution over legacy MLST, and the

identification of *Salmonella* eBGs is reasonably consistent across both approaches

[29]. rMLST has been used to identify sets of representative genomes from large

collections for calculating phylogenetic trees [28] and pan-genomes [17].

MLST has also been extended to cgMLST, which encompasses all the genes in

a soft core genome [29-31], and cgMLST nomenclatures have been implemented for

multiple bacterial genera [28]. The large number of loci encompassed by cgMLST

schemes results in enormous numbers of cgMLST STs (cgSTs), but these can be

clustered into groups of bacterial genomes at multiple levels of genomic diversity

(hierarchical clustering) [32]. cgMLST provides considerably higher resolution than

legacy MLST or rMLST, and initial analyses indicated that it is ideal for investigating

transmission chains within single source outbreaks or for identifying population

structures up to the genus level [28]. Here we focus on the automated assignments of

genome assemblies from six genera of important bacterial pathogens (Table 1) to

taxonomic and population structures by hierarchical clustering of cgSTs within

EnteroBase with the HierCC pipeline [32]. For five of those genera, cgMLST is a full

solution for taxonomic designations of species, and incorrect taxonomic designations

have now been unilaterally changed within EnteroBase.

b) EnteroBase and HierCC

EnteroBase (http://enterobase.warwick.ac.uk) automatically downloads short read

sequences and their accompanying metadata from the European Nucleotide Archive

(ENA). Metadata are then converted to consistent categories by a dedicated

EnteroBase parser with the EToKi pipeline [28]. Draft genomes are assembled from

the short reads, and annotated according to a genus-wide pan-genome scheme created

with PEPPAN [17] (Table 2). In September 2021, EnteroBase contained over 600,000

draft genomes from the six genera in Table 1, and likely provides a nearly

comprehensive overview of their global diversity. Many of these samples reflect a

focus on food-borne disease in the U.S. and United Kingdom, but this bias is

increasingly being reduced by the global sources of many genomes (Supplemental

Text, Sample Bias).

One of the primary goals for EnteroBase was a hierarchical overview of the

population structure of the genera in Table 1. We therefore developed HierCC [32]

based on cgMLST assignments to support the rapid recognition and deeper

investigation of differing levels of population structure, and EnteroBase provides

cluster assignments and designations for 10-13 levels of allelic differences for all six

genera [28] (Table 1). Some single source outbreaks of food-borne disease are caused

by uniform clades [33], and HC5 – HC10 HierCC have been used to identify tight

cgMLST clusters of *S. enterica* and *E. coli/Shigella* from single source outbreaks

across Europe [34-37]. Pathogen species can also include more distantly related

bacterial populations that are endemic in a single location or that have spread to

multiple countries [38-40]. Higher HierCC levels can be used to track such

populations, both locally and globally [28,41-43]. EnteroBase has even been used to

classify all *Shigella* [44], which correspond to discrete lineages of *E. coli* [26,45], and

to define novel sub-species of *S. enterica* [46].

The highest level of HierCC in EnteroBase can assign genomes to species/sub-species

for *Salmonella, Clostridioides, Yersinia* and *Vibrio* (Table 1). Here we compare those

assignments with assignments by classical taxonomic approaches and by 95% ANI.

The results illustrate problems with both classical approaches and 95% ANI, and we

conclude that HierCC is preferable for automated assignment of genomes to

species/subspecies for those genera. However, neither ANI nor HierCC is universally

satisfactory for species/sub-species assignments within *Streptococcus*. This

manuscript also provides an initial overview of the abilities of HierCC to assign

genomes to populations and Lineages [41,47] within *Salmonella* and

Escherichia/*Shigella*, and compare those assignments with the distributions of O

antigens within lipopolysaccharide. Detailed analysis of *Escherichia* will be presented

elsewhere.

Results

a. Species and sub-species

Classical taxonomic approaches for assigning individual strains and genomes to

species depend on human expertise, and are not suitable for automated pipelines in

real-time databases such as EnteroBase. This problem is acute because many short

read sequences in ENA have incorrect taxonomic assignments, or none at all, and

phenotypic distinctions are inappropriate for databases of 100,000s of genomes. We

therefore annotated the branch topologies within super-trees calculated by a divide

and conquer maximum likelihood approach (henceforth ML super-trees) from core

genome SNPs of representative genomes of *Salmonella, Clostridioides, Yersinia*,

Vibrio and *Streptococcus* with the groupings according to a 95% ANI cutoff

calculated with FastANI [10]. We also identified HierCC levels that corresponded

closely to peak values in normalized mutual information and silhouette plots, and

which identified well-defined species/sub-species in each genus [32]. For example,

for *Salmonella*, this optimal HC level was HC2850, which corresponds to a maximum

internal branch length of 2850 different alleles among the 3,002 loci used for

cgMLST within a minimal spanning tree of pair-wise allelic distances. The optimal

HC levels for the other five genera ranged from HC363 – HC2500 (Table 1).

Salmonella. ML super-trees based on 1,410,331 core SNPs from 10,002

representative genomes showed a strong concordance between traditional taxonomic

assignments and clustering according to either HC2850 (Fig. 1A) or 95% ANI (Fig.

1B). High concordance was also obtained with ML super-trees based on the presence

or absence of accessory genes (https://enterobase.warwick.ac.uk/ms_tree/53258).

Both HC2850 and ANI distinguished *S. enterica* from *S. bongori*, and confirmed the

recent conclusion by Pearce *et al*. [48] based on 569 genomes that former subspecies

IIIa corresponds to a separate species, *S. arizonae.* HC2850 also identified a new

Salmonella species, cluster HC2850_215890, and that conclusion was confirmed by

95% ANI. HC2850_215890 consists of five strains that have been isolated from

humans in the U.K. since 2018, and a gene for gene comparison indicated that

roughly half of the core genes were more similar to *S. enterica* and the other half to *S.*

bongori.

95% ANI did not detect any subspecies structure within *S. enterica* whereas HC2850

differentiated a total of 12 clusters that corresponded to distinct phylogenetic branches

in the ML super-tree (Fig. 1). Most of these have been designated as subspecies by

traditional taxonomy [46,48], but one of them, a singleton genome in

HC2850_222931, likely represents a new subspecies. Despite the correct

identification of so many sub-species, HC2850 did not support Pearce's definition of

subspecies XI, and did not distinguish those genomes from subsp. II. Similarly,

HC2850 did not differentiate Pearce's subsp. VIII from subsp. IV. The prior

definition of these subspecies was largely dependent on branch topologies within

phylogenetic trees based on rMLST loci, and was also contradicted by population

statistics [48]. HierCC did differentiate HC2850_7171, which consists of a short

phylogenetic branch within subsp. I, and which would not have been assigned to a

subspecies according to its ML topology. Genomic comparisons of multiple single

genes from genomes within HC2850_7171 revealed that some gene sequences were

more similar to subsp. I than to other subspecies while others were most similar to

subsp. II. These observations suggest that HC2850_7171 may be a hybrid between

those two subspecies, but do not address the question whether HC2850_7171 is a

novel subspecies or not. With this sole possible exception, HierCC seems to be

suitable for the automated detection of new *Salmonella* species and subspecies, and

for routinely assigning novel genomes to the appropriate taxa.

Clostridioides. Clustering according to 95% ANI differentiated *C. difficile* from *C.*

mangenotii, and also identified five other clusters that seem to be novel species on the

basis of the phylogeny of an ML SNP super-tree (Fig. 2). HC2500 identified the same

clusters, and also separated out two additional distinct clusters which resemble novel

sub-species (arrows: HC2500_15334, HC2500_15408). A subset of the genomes in

these clusters (Fig. 2) were assigned to cryptic clades C-I, CII and C-III by a recent

publication [49], which also concluded that they represent novel species. Thus,

HierCC is suitable for the automated detection of new *Clostridioides* species and

subspecies, and for routinely assigning novel genomes to the appropriate taxa.

Similar comparisons were also performed with *Escherichia.* These will be presented

in detail elsewhere.

Yersinia **and** ***Vibrio.*** Detailed results with *Yersinia* and *Vibrio* can be found in

Supplemental Text. The general take-home lessons from those analyses resemble

those with *Salmonella* and *Clostridioides*: 95% ANI clustering is largely congruent

with existing taxonomic designations, and HierCC identifies both species and sub-

species with better resolution than 95% ANI. However, multiple, unanticipated

taxonomic problems were evident in both genera. Firstly, ENA uses the strain

designation as a species designation when a species is not specified during the

submission process. The resulting public metadata is bloated with inaccurate species

names. Secondly, multiple distinct phylogenetic branches were found without any

species identifier. Thirdly, in multiple cases the species identifier did not correspond

with ANI and/or HierCC, and comparisons with both the SNP and presence/absence

ML topologies confirmed that those species designations in ENA were incorrect. We

addressed these problems within EnteroBase by performing radical manual curation to

the metadata to ensure that the taxonomic species designations correspond to the

phylogenies and population structures.

95% ANI supports the taxonomic convention that *Y. enterocolitica* represents a single

species, and all genomes designated as *Y. enterocolitica* were in a single 95% ANI

cluster. However, in accord with the conclusions by Reuter *et al.* [50], HierCC

clustering assigned the non-pathogenic biotype 1A genomes to HC1490_73 and

HC1490_764, the highly pathogenic biotype 1B to HC1490_2, and pathogenic

biotypes 2-5 to HC1490_10. We therefore renamed these four groups by adding the

HierCC cluster to the species name, e.g. *Y. enterocolitica* HC1490_2.

In contrast to *Y. enterocolitica,* the *Yersinia pseudotuberculosis* Complex corresponds

to a single phylogenetic cluster according to both HC1490 and 95% ANI

(Supplemental Fig. S1). Taxonomists have split these bacteria into *Y.*

pseudotuberculosis, *Y. pestis, Y. similis* and *Y. wautersii* [51,52]. DNA-DNA

hybridization [53] and gene sequences of several housekeeping genes [4] previously

demonstrated that *Y. pestis* is a clade of *Y. pseudotuberculosis,* and the new

observations demonstrate that a distinct species status is also inappropriate for *Y.*

similis and *Y. wautersii*. We have therefore downgraded all three taxa within

EnteroBase to the category of subspecies of *Y. pseudotuberculosis,* and extended the

designation of *Y. pseudotuberculosis* to *Y. pseudotuberculosis sensu stricto*. These

assignments are not reflected by distinct HC1490 clusters, and HC1490 clustering can

only be used to automatically assign new genomes to the *Y. pseudotuberculosis*

Complex.

We also defined eight other novel species/sub-species designations within *Yersinia* by

unique designations based on HierCC clusters: e.g. *Yersinia* HC1490_419. These

clusters had previously been incorrectly designated with the names of other species

which formed distinct ANI and HC1490 clusters, or had not been named at all

(SupplementalText, Fig. S1).

General concordance was also found for *Vibrio* between phylogenetic trees based on

either core SNPs or presence/absence of accessory genes, and also with the clustering

according to 95% ANI and HC1090 (SupplementalText). *Vibrio* encompassed 152

HC1090 clusters (Supplemental Fig. S2, SupplementalText), which corresponds to

much greater taxonomic diversity than for the other genera dealt with above. The

concordance between HC1090 and 95% ANI clusters was absolute for most clusters,

including a large numbers of genomes from *V. cholerae, V. parahaemolyticus*, *V.*

vulnificans, and *V. anguillarum* (Supplemental Table S1), and we did not find any

definitive evidence indicating the existence of sub-species. However, eleven HC1090

clusters each encompassed between 2 and 5 ANI clusters (Supplemental Table S2)

and three ANI clusters each encompassed 2-3 HC1090 clusters (Supplemental Table

S3). In order to support the automated assignment of genomes to named species, we

implemented taxonomic assignments according to HC1090, and renamed the species

of all genomes that were contradictory to this principle. The resulting dataset contains

109 HC1090 clusters with a unique, classical species designation as well as 43 other

species level clusters designated as *Vibrio* HC1090_xxxx. Seventeen species names

were eliminated because they were contradictory to the phylogenetic topologies or

were incoherently applied (Supplemental Text, Supplemental Table 4). Our radical

approach now permits future automated assignment of novel genomes to species

designations, and the recognition of novel species, and has provided a clean and

consistent basic taxonomy that can be progressively expanded.

Prior work has assigned many *Vibrio* species into so-called higher order clades of

species on the basis of MLSA (multilocus sequence analysis) [54]. These clades are

also apparent in the ML super-tree of core SNPs, and Fig. S2 indicates the three

largest (Cholerae, Harveyii, Splendidus).

Streptococcus. In all five genera summarized above, clustering according to HierCC

was largely concordant with the ML super-trees based on core SNPs or

presence/absence of accessory genes. 95% ANI and taxonomic designations were also

largely concordant with the phylogenetic trees. Similar concordance with the trees

based on core SNPs (Fig. 3) and presence/absence of accessory genes (Supplemental

Fig. S3) was also found for most of the named species within *Streptococcus*. 102

HC363 clusters were concordant with 95% ANI clusters, and each of those HierCC

clusters was specific for a single species after eliminating *S. milleri, S. periodonticum*

and *S. ursoris* (Supplemental Table S5). These results once again demonstrate strong

agreement between the two methods with classical taxonomy. However, genomes

from a large number of species within *Streptococcus* were not clustered satisfactorily

by either method.

Prior results had demonstrated that *Streptococcus mitis* and *oralis* each represents

multiple related sequence clusters rather than two single species [15,16,55,56], and

similar results were obtained with 95% ANI clustering [17]. The current analyses

confirmed these observations. 23 of the 102 concordant HC363 and 95% ANI clusters

from the species *S. cristatus, S. infantis, S. mitis, S. oralis, S. sanguinis,* and *S. suis*

corresponded to only a sub-set of the named species. Furthermore, HC363 identified

single clusters for 18 species or groups of species which were subdivided extensively

by 95% ANI for a total of 155 discrepancies (Supplemental Table S6). These

discrepant clusters were within the species *S. suis, S. criceti, S. sanguinis*, *S. suis, S.*

oralis, S. hyointestinalis, S. parasanguinis, S. australis, S. infantis, and *S. cristatus.*

95% ANI was not able to differentiate *S. pneumoniae* from *S. pseudopneumoniae*, *S.*

pyogenes from *S. dysgalactiae, S. salivarius* from *S. vestibularis*, or between *S.*

equinus, *S. infantarius,* and *S. lutentiensis.* Contrariwise, three other 95% ANI

clusters each encompassed two HC363 clusters (Supplemental Table S7). HC363

could not distinguish *S. pneumoniae* or *S. pseudopneumoniae* from fifty-seven 95%

ANI clusters within *S. mitis.* In addition to these problems, eight additional type

strains should not have been given a species name because they clearly belonged to

one of these chaotic clusters. We conclude that numerous species that have been

defined on taxonomic grounds cannot be correctly identified by either 95% ANI or

HierCC, and that multiple discrepancies exist between the two approaches within

Streptococcus.

b. Populations, eBGs and Lineages

For each genus in which it has been implemented, HierCC supports the identification

of populations at multiple hierarchical levels. The highest HC level was aimed at

species/sub-species and the lowest levels at deducing transmission chains. However,

an important measure of the success of HierCC would be its reliable detection of the

natural populations previously recognized by legacy MLST. These were initially

designated as clonal complexes [25], but are now referred to as ST Complexes in

Escherichia [26] and eBGs (eBurst Groups) in *Salmonella* [27]. These populations

should correspond to intermediate HierCC clustering levels, lower than for sub-

species and higher than for transmission chains. Here we focus on examples of their

from *Salmonella*, *E.coli/Shigella,* and *S. pneumoniae* where they have been examined

in greatest detail.

Salmonella. Phylogenetic trees and population genetic analyses of core genome SNP

diversity identified a limited number of deep subdivisions within *S. enterica* subsp.

enterica, which were variably designated as lineages [57], Clades [58-61] or Branches

[62]. Their phylogenetic branching topology may be unreliable because of extensive

inter-branch exchange of nucleotide diversity by homologous recombination [47,62].

We also failed to find any HierCC clustering level in *Salmonella* that was consistently

concordant with these branch structures [32]. Instead, populations were most clearly

associated with HC2000 and HC900 clusters, which represent later sub-divisions

within the phylogenetic trees. At those levels, recombination is much less prominent,

or even largely lacking [47].

Traditional nomenclature of *Salmonella* subgroupings below the sub-species level has

been predominantly based on assigning serovar designations by serotyping. Serovar

designations consist of common names that have been assigned to a total of more than

2500 unique antigenic formulas based on epitopes within lipopolysaccharide (O

antigen) and two alternately expressed flagellar subunits (H1, H2). These antigenic

formulas are written in the form O epitopes:H1 epitopes:H2 epitopes in the revised

Kauffmann-White scheme [63,64]. Detailed metabolic maps have been elucidated for

the synthesis of the LPS from 47 O serogroups which result in the expression of the O

epitopes [65,66]. These O groups have distinct numeric designations, e.g. O:4 for the

O group of serovar Typhimurium. Serogroups can be determined for natural isolates

using specific antisera for serological tests or *in silico* from genomic sequences with

the programs SeqSero [67] or SISTR [68]. In practice. each of these methods has an

error rate of a few percent [27,69].

Several hundred legacy eBGs within subsp. *enterica* were relatively uniform for

serovar [27], with multiple exceptions, and many serovars were associated with

multiple, apparently unrelated eBGs, likely due to the extensive exchange of genes

encoding LPS and flagellar epitopes between eBGs early in their evolutionary history.

We proposed that legacy MLST was an even better metric for identifying the natural

populations and could replace serotyping completely [27]. Our new data indicate that

HC900 clusters are even more reliable than legacy eBGs for recognizing natural

populations within *Salmonella*.

Several phylogenetic comparisons indicated that HC900 clusters based on cgMLST

are concordant with eBGs in *Salmonella* [32,41]. Here we provide independent

analyses confirming these observations. There is a very strong quantitative

concordance between eBG and HC900 clusters according to the Adjusted Mutual

Information Score [AMI] (Table 4), a metric that is suitable for samples with a

heterogeneous size distribution [70]. Comparisons of eBG or HC900 with HC2000

clustering yielded slightly lower AMI scores (Table 4). We also tested whether

HC900 clusters are uniform for serovar. First, we corrected false serological data in

the metadata Serovar field in EnteroBase by manual curation of 790 HC900 clusters

that contained at least 5 entries in late 2019. After manual curation, 97% (767/790) of

HC900 clusters with at least five entries were uniform (≥95%) for serovar

(Supplemental Table S8), as were most HC2000 clusters (342/409 clusters; 83.4%).

Changes in the O group between HC900 clusters within an HC2000 cluster was even

rarer: in 95.3% of 404 HC2000 clusters within *S. enterica* subsp. *enterica*, the O

group was uniform across all internal HC900 clusters and the same was the case for

80.5% of 77 HC2000 clusters in other *Salmonella* species and sub-species (Fig. 4;

Supplemental Table S9). These observations confirm and extend the prior conclusions

about concordance between natural populations and serogroup within *Salmonella.*

Escherichia coli. Patterns of multilocus isoenzyme electrophoresis were used in the

1980s to provide an overview of the genetic diversity of *E. coli* (see overview by

Chaudhuri and Henderson [71]) . Those analyses yielded a representative collection of

72 isolates [72], the EcoR collection, whose deep phylogenetic branches were

designated haplogroups A, B1, B2, C, D and E [73]. Several haplogroups have since

been added [74,75]. The presence or absence of several accessory genes can be used

for the assignment of genomes to haplogroups with the Clermont scheme [76,77], and

the haplogroup can also be predicted *in silico* from genomic assemblies with

ClermontTyping [78] or EZClermont [79], both of which are implemented within

EnteroBase. However, the Clermont scheme ignores *Shigella*, which consists of *E.*

coli clades despite its differing genus designation [26,44,45], and makes multiple

discrepant assignments according to phylogenetic trees [28]. The Clermont scheme

also does not properly handle the entire diversity of species and environmental

clades/subspecies within the genus *Escherichia* [28,80-82].

EnteroBase HierCC automatically assigns genomes within the *Escherichia/Shigella*

database to the cgMLST equivalents of Lineages (HC2000 clusters) and ST

Complexes (HC1100 clusters) (Table 1). It also perpetuates the ST Complexes that

were initially defined for legacy MLST by Wirth *et al.* [26]. However, the numbers

and composition of legacy ST Complexes have not been updated since 2009 because

additional sequencing data defined intermediate, recombinant genotypes that were

equidistant and similar to more than one ST Complex, and threatened to merge

existing ST Complexes. In contrast, HC1100 clusters do not merge: Intermediate

genotypes are rare because cgMLST involves 2512 loci while legacy MLST is based

on only seven. Secondly, new genotypes which are similar to and equidistant to

multiple clusters would not trigger merging because HierCC arbitrarily assigns them

to the oldest of the existing alternatives. Finally, unlike legacy MLST which has not

been actively pursued in *Escherichia* since 2006, HierCC clustering is an ongoing,

automated process for all new genotypes.

Frequent recombination in *E. coli* results in poor bootstrap support for the deep

branches in phylogenetic trees of concatenated genes [26], and we were unable to

identify a unique HierCC level which was largely concordant with the Clermont

scheme. However, similar to *Salmonella,* HC1100 clusters identify clear population

groups, which are highly concordant with legacy ST Complexes (Table 4). Unlike

Salmonella, HC2000 clusters did not in general mark recognizable additional

population structure beyond that which was provided by ST Complexes, and HC2000

clusters are only moderately concordant with either legacy ST Complexes or their

cgMLST equivalent, HC1100 clusters (Table 4). Exceptionally, legacy ST Complexes

which originally grouped multiple *Shigella* species [26] equate to HC2000 clusters

rather than HC1100 clusters according to phylogenetic trees. HC2000_305 replaces

ST152 Cplx, and encompasses all *S. sonnei*. HC2000_192 replaces ST245 Cplx, and

contains many *S. flexneri*. However, *S. flexneri* O:group F6 is in an HC1100 cluster

together with *S. boydii* B2 and B4 within HC2000_1465 (ST243 Cplx).

HC2000_1465 also include a second HC1100 cluster with *S. boydii* O groups B1 and

B18, and a third with *S. dysenteriae* D3, D9 and D13 (Fig. 5). HC2000_4118 replaces

the combination of ST250 Complex and ST149 Complex, which have merged, and

contains both *S. dysenteriae* as well as *S. boydii.*

Unlike *Salmonella*, O serogroups are remarkably variable within HC1100 clusters of

E. coli (Fig. 5). HC1100_63 (ST11 Complex) combines classical EHEC strains of

serovar O157:H7 and atypical EPEC isolates of serovar O55:H7, which are thought to

be ancestral [83]. Multiple other HC1100 clusters that contain EHEC strains also

encompass multiple O groups, including HC1100_2 (ST29 Cplx; O26, O103, O111)

[84] and HC1100_3 (ST20 Cplx; O45, O103, O111, O128). Similarly, multiple

HC1100 clusters of *E. coli* that cause extra-intestinal diseases also contain a variety of

O groups, including HC1100_7 (ST131 Cplx; O16, O25) [23,85,86] and HC1100_44

(K1-encapsulated bacteria of the ST95 Cplx; O1, O2, O18, O25, O45) [26,87,88].

Indeed, the primary impression from Fig. 5 is that O group variation within an

HC1100 cluster is nearly universal throughout *E. coli*, confirming prior conclusions

about the high frequency of homologous recombination in this species [26,74].

Other genera. Correlations have been described in *Clostridioides difficile* [42]

between intra-specific HierCC clusters and population groupings at multiple levels,

including ST Complexes and ribotypes. It remains to be proven whether HierCC can

identify populations within *Yersinia* because bacterial populations in *Y.*

pseudotuberculosis are largely obscured by recombination [89]. Similarly, we are not

aware of extensive efforts to identify bacterial populations within *Vibrio* which could

be compared with HierCC. However, recent work has described the assignment of

large numbers of genomes of *Streptococcus pneumoniae* to legacy Clonal Complexes

as well as GSPC (Global Pneumococcal Sequence Clusters) [90], initially with

PopPunk [91] and more recently with Mandrake [92]. Most of the Illumina short reads

456 for those genomes had been uploaded to ENA, and were automatically imported into

457 EnteroBase, which then assembled genomic sequences and assigned them to HierCC

458 clusters. We therefore compared the HierCC and GSPC assignments for 18,147

459 genomes, and also compared them to Clonal Complexes (CCs) from legacy MLST

460 (13,396 genomes). HC160 clusters yielded the greatest AMI and ARI scores with

461 HC160 clusters while CC was most concordant with HC100 clusters (Table 4). Visual

462 examination of of nodes color-coded by the different assignments within a Neighbor-

463 Joining tree based on cgMLST distances supported these conclusions (Supplemental

464 Fig. S4).

Discussion

We initially anticipated that EnteroBase might contain as many as 10,000 genomes of

Salmonella and *Escherichia*. Instead, EnteroBase currently contains >600,000

assembled genomes of *Salmonella, Escherichia/Shigella, Streptococcus*, *Vibrio*,

Clostridioides and *Yersinia*, and has become one of the primary global sources of

their genomic data. An analyses of their genomic properties is facilitated by software

tools (Table 2) that support indexing of genetic diversity, and searching as well as

investigating stable population structures. Here we focused on the automated

identification and designation of species/subspecies and populations on the basis of

HierCC of core genome MLST genotypes. The data demonstrate that HierCC can

completely replace classical taxonomic methods for genomes of *Salmonella,*

Escherichia/Shigella, Vibrio, *Clostridioides* and *Yersinia*, with only few minor

exceptions. HierCC is also a complete replacement for legacy MLST regarding the

assignments to populations within *Salmonella, Escherichia/Shigella* and

Streptococcus pneumoniae.

Taxonomic assignments. Disjointed tree merging (DTM) within a divide-and-

conquer approach allows large phylogenetic trees to be calculated in a reasonable time

[93]. We developed cgMLSA, a novel DTM approach that enabled analyses of the

core and accessory genome diversity of up to 10,000 representative genomes from

each of six bacterial genera. ML super-trees were calculated for each genus based on

core genome SNPs and from a presence/absence matrix for accessory genes. The

paired trees were generally topologically concordant. Therefore, species/subspecies

are well defined and reproducible in these genera. We compared current taxonomic

species designations with their topologies in the paired trees, and against clusters

based on ANI and HierCC patterns. HierCC was at least as compatible with current

taxonomy as 95% ANI for *Salmonella, Escherichia/Shigella, Vibrio, Clostridioides*

and *Yersinia,* and was much better for detecting sub-species. Classical taxonomical

designations had multiple, glaring problems in each genus and we introduced an

automated HierCC-based species/subspecies taxonomy in EnteroBase for

problematical branches. We also defined novel species/sub-species and gave them

HierCC designations. Why did we find so many obvious discrepancies? This may

reflect the additional novel possibilities introduced by EnteroBase and HierCC, and

the previously unprecedented analysis of so many representative genomes.

Our taxonomic changes in *Yersinia* are likely to be highly controversial. According to

both ML trees and HierCC, the current designation "*Y. enterocolitica*" encompasses

four, previously unnamed species/sub-species. These taxa are not distinguished by

95% ANI and had not previously been assigned distinct names. We maintain

continuity with the traditional designation of *Y. enterocolitica*, and simply add

HierCC affiliations to the species name, e.g. *Y. enterocolitica* HC1490_2.

Alternatively, they could be considered to represent sub-species and their name

modified slightly, e.g. *Y. enterocolitica* subspecies HC1490_2, similar to our

downgrading of the named species *Y. pestis, Y. similis* [52] and *Y. wautersii* [51] on

phylogenetic grounds to #to subspecies of the *Y. pseudotuberculosis* Complex (Fig.

S1)*. Finally, we defined eight new species/sub-species within *Yersinia* that were not

previously differentiated, and named them after their HC cluster, e.g. *Yersinia*

HC1490_4399, without submitting traditional evidence for simple phenotypic

differences. We also followed comparable strategies for the other genera analyzed

here.

In *Salmonella,* HierCC and 95% ANI identified a new species, *Salmonella*

HC2850_215890, the first since decades (Fig. 1). HierCC also identified a new

subspecies, *S. enterica* subsp. HC2850_222931. Five distinct new species were

identified within *C. difficile* by ANI as well as by HierCC, and HierCC identified two

additional new sub-species. *Escherichia* taxonomy has already been dealt with

previously [28], and an updated, more comprehensive analysis including HierCC

assignments will be published elsewhere.

The ML trees presented here (Fig. S2) supported prior definitions of high order clades

containing multiple species in *Vibrio* on the basis of MLSA [54]. We also observed a

general concordance between ANI and HierCC for 109 named *Vibrio* species,

However, 43 HierCC clusters of species rank had not previously been properly

classified with species designations and 17 other species names were superfluous.

These changes are fully described in Supplemental text, allowing closer scrutiny by

the *Vibrio* taxonomic community for consistency with other properties.

HierCC was so effective in clarifying the taxonomies of these five species that we had

hoped that it could also facilitate the taxonomic classification of *Streptococcus*. 102

HC363 clusters were indeed concordant with 95% ANI and taxonomic designations,

with only minor exceptions. However, extensive discrepancies existed between 95%

ANI and HierCC for numerous other clusters, and both approaches showed multiple

discrepancies with classical taxonomic designations. The existence of multiple 95%

ANI clusters within *S. mitis* and *S. oralis* and other species of *Streptococcus* has

previously been commented on by Kilian and his colleagues [15,16,55,56]. Our

observations extend these taxonomic problems to multiple additional species in which

HC363 does not distinguish between pairs of named species (Fig. 3). Neither 95%

ANI nor cgMLST HierCC seems to provide a suitable general strategy for elucidating

the taxonomy of all of *Streptococcus*, and we failed to find a general solution to this

problem.

HierCC *versus* classical taxonomy. Classical microbiological species taxonomy

involves identifying a type strain whose phenotype can be distinguished from all other

type strains, identifying additional isolates with similar phenotypes, demonstrating

distinct clustering from other known species in phylogenetic trees based on DNA or

amino acid sequences, and publishing a report in one of a very limited number of

acceptable journals. Such species definitions are then considered tentative until an

international committee has approved them. Other forms of identifying species that

include sole reliance on DNA sequence differences are not acceptable [7,8]. The

metagenomics community and scientists working with uncultivated organisms from

the environment have largely liberated themselves from such intellectually obsolete

regulations, and tend to use operational taxonomic units (OTUs) as taxonomic

entities. However, the taxonomic species structure of many microbes from

environmental sources remains fuzzy, or does not clearly correspond to classical

taxonomy (e.g. *Prochlorococcus* [94] or *Synechococcus* [95]). Furthermore, we are

not aware of a prior method that has been adequate to the tasks of assigning 1000s of

genomes per day to existing taxa and also identifying new taxa automatically as they

appear. HierCC performs this task with bravura for the genera in Table 1, with the

notable exception of *Streptococcus*.

We have taken the liberty of dropping all attempts to reconcile HierCC

species/subspecies clusters with classical prescriptions for how to define a species.

Instead we have adopted the practice of using HierCC cluster designations within

EnteroBase for the nomenclatures for species/subspecies groupings which had not yet

been identified by others. We split *Y. enterocolitica* into four related taxa and changed

the nomenclatures of species within the *Y. pseudotuberculosis* Complex to that of

subspecies. We also eliminated from EnteroBase multiple species designations of type

strains in *Vibrio* and *Streptococcus* where they did not match the ML tree topologies

and HierCC clusters. These actions provided a uniform base for the future additions of

additional species and ensured that future genomes can be correctly assigned to

uniform clusters of related strains. Scientists wishing to use these schemes to identify

the species of their bacterial isolates can upload their sequenced genomes to

EnteroBase. The HierCC assignments will be available within hours. We also

welcome additional curators of these databases who are willing to test and improve

the current taxonomic assignments we have implemented. But we reject the concept

of designating our assignments as tentative until they are confirmed in several years

by an international committee.

HierCC and Populations. The first bacterial taxonomic designations were assigned

over 100 years ago. Bacterial population genetics is much younger, and has many

fewer practitioners. Initial population genetic analyses in the early 1980s subdivided

multiple bacterial species into intra-specific lineages based on multilocus enzyme

electrophoresis [72]. This methodology was replaced in the late 1990s by legacy

MLST based on several housekeeping genes [21], which is currently being replaced

by cgMLST based on all common genes in the soft core genome [22,30]. EnteroBase

calculates genotypes for both types of MLST, as well as for rMLST [28]. Legacy STs

differ from cgSTs due to different levels of resolution. However, the boundaries of ST

Complexes/eBGs are highly concordant independent of the genes used for MLST

(Table 4). eBGs in *S. enterica* correspond to HC900 clusters [27,46], ST Complexes

in *E. coli* correspond to HC1100 clusters and Clonal Complexes (CCs) in *S.*

pneumoniae to HC100 clusters (Table 4). Similarly, serovars are largely uniform

within HC900 clusters/eBGs in *S. enterica* [27] as are serotypes across CCs in *S.*

pneumoniae [90].

We interpret these consistencies between legacy, rMLST and cgMLST as reflecting

the existence of natural populations. We previously claimed that legacy ST

Complexes in *E. coli* were unstable due to frequent recombination [26], and were

therefore not surprised at their tendency to merge as additional isolates were

sequenced. However, a major problem with legacy MLST is that it is based on only

seven genes. The high resolution of cgMLST and the decision to assign new

genotypes that are equidistant from multiple HierCC clusters to the older cluster have

negated these problems, and *E. coli* HC1100 clusters correspond to bacterial

groupings that have been independently identified by multiple phenotypic patterns. In

the early 1980s, MA began his academic research on bacterial pathogens with *E. coli*

that expressed the K1 polysaccharide capsule [87]. He observed unusually uniform

patterns of electrophoretic mobility of major outer membrane proteins across multiple

isolates, and interpreted those bacteria as representing "clones". These "clones"

correspond to HC1100 clusters, including HC1100_44 (ST95 Complex) which is still

a major cause of extra-invasive disease in humans and animals around the globe [88].

The 1980s analyses showed that these K1 bacteria were variably O1, O2, or O18, and

that the serotype variants differed both in their invasiveness and in the hosts which

they infected [96]. As indicated in Fig. 5, HC1100_44 includes O groups O25 and

O45 in addition to O1, O2 and O18. Similarly, HC1100_7 (ST131) represents another

major cause of extraintestinal disease in humans [23,85,86]. EHEC bacteria are

notorious for their association with hemolytic uremic syndrome (HUS). Many of them

belong to distinctive HC1100 clusters, including O157:H7/O55:H7 (HC1100_63,

ST11 Cplx [83]) and O26, O103 and O111 EHEC bacteria (HC1100_2, ST29 Cplx

[84]) (Fig. 5).

In contrast to such correlations with epidemiological groupings, we were unable to

identify any HierCC levels that were consistently concordant with other intra-specific

subdivisions based on phylogenetic SNP trees, including haplogroups [73] or

Clermont typing [76,77] in *E. coli* [28] and Clades [58-61], lineages [57] or Branches

[62] in *S. enterica.* Those intra-specific phylogenetic sub-divisions may not

necessarily be informative across the entire species.

Lineages. Instead of phylogenetic sub-divisions, we introduce Lineages, a subset of

population structures that are of higher order than ST Complexes/eBGs and lower

than sub-species. In *E. coli*, HC2000 Lineages were particularly appropriate for

defining the clustering level of seven groups of *Shigella* genomes (Fig. 5). In *S.*

enterica subspecies *enterica*, several HC2000 Lineages combined multiple serovars.

These included the Para C Lineage (HC2000_1272; serovars Paratyphi C, Typhisuis,

Choleraesuis and Lomita [47]) which emerged about 3500 years ago (Supplemental

Text) [47,62]. The observation that the Para C Lineage is uniformly serogroup O:7 is

not particularly surprising because all its serovars have identical antigenic formulas,

and are only distinguished by biotype. The Enteritidis Lineage (HC2000_12) consists

of serovars Enteritidis (1,9,12:g,m:-), Gallinarum and its variant Pullorum (1,9,12:-:-),

and Dublin (1,9,12:g,p:-:-) [41] (Fig. 4), and it is also not surprising that it is uniform

for O:9. However, we were not prepared for the Typhimurium Lineage (HC2000_2),

which includes serovars Typhimurium, Heidelberg, Reading, Saintpaul, Haifa,

Stanleyville and others; [41]) All these serovars are O:4, and this Lineage is therefore

also uniform for O group. The Mbandaka Lineage (HC2000_4) is uniformly

serogroup O:7 because serovar Mbandaka is 6,7,14: z10:e,n,z15 and serovar Lubbock

is 6,7:g,m,s:e,n,z15. However, three Lineages did contain more than one O serogroup.

HC2000_299 is a mixture of O:4 (HC900_299; serovar Agama [28]) and O:7

(HC900_127074; serovar Nigeria). HC900 clusters within HC2000_54 are O:4

(HC900_54: Bredeney, Schwarzengrund; HC24937: Kimuenza) or O:9,46

(HC900_57: Give). Similarly, HC900 groups within HC2000_106 are O:3,10; O:9;

O:9,46; O:4: or O:8. These exceptional HC2000 Lineages might be worth exploring in

greater detail to reconstruct the recombination that has resulted in such differences in

LPS epitopes.

HC160 clusters are the equivalent of Lineages in *S. pneumonia*, and HC100 the

equivalent of Clonal Complexes. HC160 clusters were strongly concordant with

PopPunk GSPC clustering. Otherwise, little is yet known about the general properties

of Lineages, except that they may vary across multiple species or genera. For

example, whereas *S. enterica* HC2000 Lineages were uniform for O group, O

serogroups were heterogeneous within HC1100 clusters/ST Complexes within *E. coli*

(Fig. 5).

Future prospects. EnteroBase was conceived to satisfy a need that MA and ZZ

perceived in 2014 [1]. We contend that it is now fit for purpose for microbial

epidemiology and population genetics of multiple bacterial genera for scientists

ranging from beginners through to experts in those areas. Its original creators have

now all left this project, but EnteroBase is being maintained as a service for the global

community by the University of Warwick. Maintenance of its technical functions and

databases are thereby assured for the near future. Further functional developments

will, however, depend on increased participation and ownership by its users. We

perceive a general trend to focus on insular solutions that can satisfy the demands of

individual bioinformaticians and regional diagnostic laboratories. Such approaches

can yield relatively rapid progress in solving short-term needs. However, a global

overview of genomic diversity needs central databases to ensure definitive

terminology. Pooling efforts on a central endeavor at the scale represented by

EnteroBase would ensure that it continues to function over decades, is representative

over all continents, and serves the global community even better. We therefore

welcome additional curators and scientific experts as well as bioinformatics

collaborations to improve EnteroBase even more.

Methods (see Supplemental Text)

Funding

MA was supported by grant 202792/Z/16/Z from the Wellcome Trust (https://wellcome.org/).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Code.

cgMLSA is available for downloading at https://github.com/zheminzhou/cgMLSA.

All trees in figures and genomic data are freely available within EnteroBase for interactive examination. Other code is described in Supplemental Text

Acknowledgements

We are very grateful to François-Xavier Weill for permission to include Table S8 for public access in Supplemental Material. We are very grateful to François-Xavier Weill and Tandy Warnow for their comments on an earlier draft of this manuscript.

687   References

689   1.   Achtman, M. and Zhou, Z. 2014 Distinct genealogies for plasmids and
690   chromosome. *PLoS Genet* **10**, e1004874. (doi:10.1371/journal.pgen.1004874
691   [doi];PGENETICS-D-14-02780 [pii])

692   2.   Kauffmann, F. 1961 *Die Bakteriologie der Salmonella-Species*. Copenhagen,
693   Denmark: Munksgaard.

694   3.   Morelli, G. *et al.*. 2010 *Yersinia pestis* genome sequencing identifies patterns of
695   global phylogenetic diversity. *Nature Genet* **42**, 1140-1143.

696   4.   Achtman, M., Zurth, K., Morelli, G., Torrea, G., Guiyoule, A., and Carniel, E.
697   1999 *Yersinia pestis*, the cause of plague, is a recently emerged clone of
698   *Yersinia pseudotuberculosis*. *Proc Natl Acad Sci USA* **96**, 14043-14048.

699   5.   Pulford, C. V. *et al.*. 2021 Stepwise evolution of *Salmonella* Typhimurium
700   ST313 causing bloodstream infection in Africa. *Nat Microbiol* **6**, 327-338.
701   (doi:10.1038/s41564-020-00836-1 [doi];10.1038/s41564-020-00836-1 [pii])

702   6.   Wayne, L. G. *et al.*. 1987 Report of the ad hoc committee on reconciliation of
703   approaches to bacterial systematics. *In J Syst Bacteriol* **37**, 463-464.

704   7.   Murray, A. E. *et al.*. 2020 Roadmap for naming uncultivated Archaea and
705   Bacteria. *Nat Microbiol* **5**, 987-994. (doi:10.1038/s41564-020-0733-x
706   [doi];10.1038/s41564-020-0733-x [pii])

707   8.   Sutcliffe, I. C., Dijkshoorn, L., Whitman, W. B., and Executive Board. 2020
708   Minutes of the International Committee on Systematics of Prokaryotes online
709   discussion on the proposed use of gene sequences as type for naming of
710   prokaryotes, and outcome of vote. *Int J Syst Evol Microbiol* **70**, 4416-4417.
711   (doi:10.1099/ijsem.0.004303 [doi])

712   9.   Konstantinidis, K. T. and Tiedje, J. M. 2005 Genomic insights that advance the
713   species definition for prokaryotes. *Proc Natl Acad Sci USA* **102**, 2567-2572.

714   10.   Jain, C., Rodriguez, R., Phillippy, A. M., Konstantinidis, K. T., and Aluru, S.
715   2018 High throughput ANI analysis of 90K prokaryotic genomes reveals
716   clear species boundaries. *Nat Commun* **9**, 5114. (doi:10.1038/s41467-018-
717   07641-9 [doi];10.1038/s41467-018-07641-9 [pii])

718   11.   Auch, A. F., von, J. M., Klenk, H. P., and Goker, M. 2010 Digital DNA-DNA
719   hybridization for microbial species delineation by means of genome-to-
720   genome sequence comparison. *Stand Genomic Sci* **2**, 117-134.
721   (doi:10.4056/sigs.531120 [doi])

722   12.   Garrido-Sanz, D., Meier-Kolthoff, J. P., Goker, M., Martin, M., Rivilla, R., and
723   Redondo-Nieto, M. 2016 Genomic and Genetic Diversity within the
724   *Pseudomonas fluorescens* Complex. *PLoS One* **11**, e0150183.
725   (doi:10.1371/journal.pone.0150183 [doi];PONE-D-15-47408 [pii])

726   13.   Richter, M. and Rosselló-Mora, R. 2009 Shifting the genomic gold standard for
727   the prokaryotic species definition. *Proc Natl Acad Sci U S A* **106**, 19126-
728   19131. (doi:0906412106 [pii];10.1073/pnas.0906412106 [doi])

729   14.   Maderankova, D., Jugas, R., Sedlar, K., Vitek, M., and Skutkova, H. 2019 Rapid
730   bacterial species delineation based on parameters derived from genome
731   numerical representations. *Comput Struct Biotechnol J* **17**, 118-126.
732   (doi:10.1016/j.csbj.2018.12.006 [doi];S2001-0370(18)30122-3 [pii])

733   15.   Jensen, A., Scholz, C. F., and Kilian, M. 2016 Re-evaluation of the taxonomy of
734   the Mitis group of the genus *Streptococcus* based on whole genome
735   phylogenetic analyses, and proposed reclassification of *Streptococcus*
736   *dentisani* as *Streptococcus oralis* subsp. *dentisani* comb. nov., *Streptococcus*
737   *tigurinus* as *Streptococcus oralis* subsp. *tigurinus* comb. nov., and
738   *Streptococcus oligofermentans* as a later synonym of *Streptococcus cristatus*.
739   *Int J Syst Evol Microbiol* **66**, 4803-4820. (doi:10.1099/ijsem.0.001433 [doi])

16.   Kilian, M., Poulsen, K., Blomqvist, T., Havarstein, L. S., Bek-Thomsen, M.,
Tettelin, H., and Sorensen, U. B. 2008 Evolution of *Streptococcus*
pneumoniae and its close commensal relatives. *PLoS ONE* **3**, e2683.

17.   Zhou, Z., Charlesworth, J., and Achtman, M. 2020 Accurate reconstruction of
bacterial pan- and core genomes with PEPPAN. *Genome Res* **30**, 1667-1679.
(doi:DOI 10.1011/gr.260828.120)

18.   Gomila, M., Pena, A., Mulet, M., Lalucat, J., and Garcia-Valdes, E. 2015
Phylogenomics and systematics in *Pseudomonas*. *Front Microbiol* **6**, 214.
(doi:10.3389/fmicb.2015.00214 [doi])

19.   Beaz-Hidalgo, R., Hossain, M. J., Liles, M. R., and Figueras, M. J. 2015
Strategies to avoid wrongly labelled genomes using as example the detected
wrong taxonomic affiliation for *Aeromonas* genomes in the GenBank
database. *PLoS ONE* **10**, e0115813. (doi:10.1371/journal.pone.0115813
[doi];PONE-D-14-36626 [pii])

20.   Achtman, M. and Wagner, M. 2008 Microbial diversity and the genetic nature
of microbial species. *Nat Rev Microbiol* **6**, 431-440.

21.   Maiden, M. C. J. *et al.*. 1998 Multilocus sequence typing: A portable approach
to the identification of clones within populations of pathogenic
microorganisms. *Proc Natl Acad Sci USA* **95**, 3140-3145.

22.   Jolley, K. A., Bray, J. E., and Maiden, M. C. 2018 Open-access bacterial
population genomics: BIGSdb software, the PubMLST.org website and their
applications. *Wellcome Open Res* **3**, 124. (doi:doi:
10.12688/wellcomeopenres.14826.1)

23.   Stoesser, N. *et al.*. 2016 Evolutionary history of the global emergence of the
Escherichia coli epidemic clone ST131. *MBio* **7**, e02162. (doi:mBio.02162-
15 [pii];10.1128/mBio.02162-15 [doi])

24. Kingsley, R. A. *et al.*. 2009 Epidemic multiple drug resistant *Salmonella*
Typhimurium causing invasive disease in sub-Saharan Africa have a distinct
genotype. *Genome Res* **19**, 2279-2287.

25. Feil, E. J., Li, B. C., Aanensen, D. M., Hanage, W. P., and Spratt, B. G. 2004
eBURST: Inferring patterns of evolutionary descent among clusters of related
bacterial genotypes from Multilocus Sequence Typing data. *J Bacteriol* **186**,
1518-1530.

26. Wirth, T. *et al.*. 2006 Sex and virulence in *Escherichia coli*: an evolutionary
perspective. *Mol Microbiol* **60**, 1136-1151.

27. Achtman, M. *et al.*. 2012 Multilocus sequence typing as a replacement for
serotyping in *Salmonella enterica*. *PLoS Pathog* **8**, e1002776.

28. Zhou, Z., Alikhan, N. F., Mohamed, K., Fan, Y., and Achtman, M. 2020 The
EnteroBase user's guide, with case studies on *Salmonella* transmissions,
Yersinia pestis phylogeny, and *Escherichia* core genomic diversity. *Genome*
Res **30**, 138-152. (doi:gr.251678.119 [pii];10.1101/gr.251678.119 [doi])

29. Mellmann, A. *et al.*. 2011 Prospective genomic characterization of the German
enterohemorrhagic *Escherichia coli* O104:H4 outbreak by rapid next
generation sequencing technology. *PLoS ONE* **6**, e22751.

30. Maiden, M. C., van Rensburg, M. J., Bray, J. E., Earle, S. G., Ford, S. A., Jolley,
K. A., and McCarthy, N. D. 2013 MLST revisited: the gene-by-gene
approach to bacterial genomics. *Nat Rev Microbiol* **11**, 728-736.

31. Moura, A. *et al.*. 2016 Whole genome-based population biology and
epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol* **2**,
16185. (doi:nmicrobiol2016185 [pii];10.1038/nmicrobiol.2016.185 [doi])

32. Zhou, Z., Charlesworth, J., and Achtman, M. 2021 HierCC: A multi-level
clustering scheme for population assignments based on core genome MLST.
Bioinformatics **37**, 3645-3646. (doi:10.1093/bioinformatics/btab234 [doi])

33.   Bielaszewska, M., Mellmann, A., Zhang, W., Kock, R., Fruth, A., Bauwens, A.,
Peters, G., and Karch, H. 2011 Characterisation of the *Escherichia coli* strain
associated with an outbreak of haemolytic uraemic syndrome in Germany,
2011: a microbiological study. *Lancet Infect Dis* **11**, 671-676.

34.   Jones, G. *et al.*. 2019 Outbreak of *Salmonella enterica* serotype Poona in infants
linked to persistent *Salmonella* contamination in an infant formula
manufacturing facility, France, August 2018 to February 2019. *Euro Surveill*
24, 13. (doi:doi.org/10.2807/1560-7917.ES.2019.24.13.1900161)

35.   Jones, G. *et al.*. 2019 Outbreak of *Shiga* toxin-producing *Escherichia coli*
(STEC) O26 paediatric haemolytic uraemic syndrome (HUS) cases
associated with the consumption of soft raw cow's milk cheeses, France,
March to May 2019. *Euro Surveill* **24**, 1900305. (doi:10.2807/1560-
7917.ES.2019.24.22.1900305)

36.   Van den Bossche, A., Ceyssens, P. J., Denayer, S., Hammami, N., van den Beld,
M., Dallman, T. J., and Mattheus, W. 2021 Outbreak of Central American
born *Shigella sonnei* in two youth camps in Belgium in the summer of 2019.
Eur J Clin Microbiol Infect Dis **40**, 1573-1577. (doi:10.1007/s10096-021-
04164-y [doi];10.1007/s10096-021-04164-y [pii])

37.   European Centre for Disease Prevention and Control, E. F. S. A. 2020 Multi-
country outbreak of *Salmonella* Typhimurium and *S*. Anatum infections
linked to Brazil nuts - 21 October 2020. *EFSA Supporting Publications* **17**,
1944E. (doi:https://doi.org/10.2903/sp.efsa.2020.EN-1944)

38.   Hawkey, J. *et al.*. 2021 Global population structure and genotyping framework
for genomic surveillance of the major dysentery pathogen, *Shigella sonnei*.
Nat Commun **12**, 2684. (doi:10.1038/s41467-021-22700-4
[doi];10.1038/s41467-021-22700-4 [pii])

39.   Park, S. E. *et al.*. 2018 The phylogeography and incidence of multi-drug
820   resistant typhoid fever in sub-Saharan Africa. *Nat Commun* **9**, 5094.
821   (doi:10.1038/s41467-018-07370-z [doi];10.1038/s41467-018-07370-z [pii])

822   40.   Weill, F. X. *et al.*. 2017 Genomic history of the seventh pandemic of cholera in
823   Africa. *Science* **358**, 785-789. (doi:358/6364/785
824   [pii];10.1126/science.aad5901 [doi])

825   41.   Achtman, M. *et al.*. 2020 Genomic diversity of *Salmonella enterica* -The
826   UoWUCC 10K genomes project. *Wellcome Open Res* **5**, 223.
827   (doi:10.12688/wellcomeopenres.16291.2 [doi])

828   42.   Frentrup, M. *et al.*. 2020 A publicly accessible database for *Clostridioides*
829   *difficile* genome sequences supports tracing of transmission chains and
830   epidemics. *Microbial Genomics*, mgen.0.000410.
831   (doi:10.1099/mgen.0.000410 [doi])

832   43.   Achtman, M., van den Broeck, F., Cooper, K. K., Lemey, P., Parker, C. T.,
833   Zhou, Z., and the ATCC14028s Study Group. 2021 Genomic population
834   structure associated with repeated escape of *Salmonella enterica*
835   ATCC14028s from the laboratory into nature. *PLoS Genet* **17**, e1009820.
836   (doi:DOI: 10.1371/journal.pgen.1009820)

837   44.   Yassine, I. *et al.*. 2021 Population structure analysis and laboratory monitoring
838   of *Shigella* with a standardised core-genome multilocus sequence typing
839   scheme: a validation study. *BioRxiv*, 2021.

840   45.   Pupo, G. M., Lan, R., and Reeves, P. R. 2000 Multiple independent origins of
841   Shigella clones of *Escherichia coli* and convergent evolution of many of their
842   characteristics. *Proc Natl Acad Sci USA* **97**, 10567-10572.

843   46.   Alikhan, N.-F., Zhou, Z., Sergeant, M. J., and Achtman, M. 2018 A genomic
844   overview of the population structure of *Salmonella*. *PLoS Genet* **14**,

e1007261. (doi:10.1371/journal.pgen.1007261 [doi];PGENETICS-D-18-
00122 [pii])

47.   Zhou, Z. *et al.*. 2018 Pan-genome analysis of ancient and modern *Salmonella*
enterica demonstrates genomic stability of the invasive Para C Lineage for
millennia. *Curr Biol* **28**, 2420-2428.
(doi:https://doi.org/10.1016/j.cub.2018.05.058)

48.   Pearce, M. E., Langridge, G. C., Lauer, A. C., Grant, K., Maiden, M. C. J., and
Chattaway, M. A. 2021 An evaluation of the species and subspecies of the
genus *Salmonella* with whole genome sequence data: Proposal of type strains
and epithets for novel *S. enterica* subspecies VII, VIII, IX, X and XI.
Genomics **113**, 3152-3162. (doi:S0888-7543(21)00281-0
[pii];10.1016/j.ygeno.2021.07.003 [doi])

49.   Knight, D. R., Imwattana, K., Kullin, B., Guerrero-Araya, E., Paredes-Sabja, D.,
Didelot, X., Dingle, K. E., Eyre, D. W., Rodríguez, C., and Riley, T. V. 2021
Major genetic discontinuity and novel toxigenic species in *Clostridioides*
difficile taxonomy. *eLife* **10**, e64325.

50.   Reuter, S. *et al.*. 2014 Parallel independent evolution of pathogenicity within the
genus *Yersinia*. *Proc Natl Acad Sci U S A* **111**, 6768-6773. (doi:1317161111
[pii];10.1073/pnas.1317161111 [doi])

51.   Savin, C. *et al.*. 2014 The *Yersinia pseudotuberculosis* complex:
characterization and delineation of a new species, *Yersinia wautersii*. *Int J*
Med Microbiol **304**, 452-463. (doi:S1438-4221(14)00012-5
[pii];10.1016/j.ijmm.2014.02.002 [doi])

52.   Sprague, L. D., Scholz, H. C., Amann, S., Busse, H. J., and Neubauer, H. 2008
Yersinia similis sp. nov. *Int J Syst Evol Microbiol* **58**, 952-958.

53.   Bercovier, H., Mollaret, H. H., Alonso, J. M., Brault, J., Fanning, G. R.,
Steigerwalt, A. G., and Brenner, D. J. 1980 Intra- and interspecies relatedness

872   of *Yersinia pestis* by DNA hybridization and its relationship to *Yersinia*

873   *pseudotuberculosis. Curr Microbiol* **4**, 225-229.

874   54. Gomez-Gil, B., Thompson, C. C., Matsumura, Y., Sawabe, T., Iida, T., Christen,

875   R., Thompson, F., and Sawabe, T. 2014 The family *Vibrionaceae*. In *The*

876   *prokaryotes - Gammaproteobacteria* (eds. E. Rosenberg, E. F. DeLong, S.

877   Lory, E. Stackebrandt, and F. Thompson), pp. 660-747. Berlin Heidelberg.

878   55. Kilian, M., Riley, D. R., Jensen, A., Bruggemann, H., and Tettelin, H. 2014

879   Parallel evolution of *Streptococcus pneumoniae* and *Streptococcus mitis* to

880   pathogenic and mutualistic lifestyles. *MBio* **5**, e01490-14. (doi:mBio.01490-

881   14 [pii];10.1128/mBio.01490-14 [doi])

882   56. Bishop, C. J., Aanensen, D. M., Jordan, G. E., Kilian, M., Hanage, W. P., and

883   Spratt, B. G. 2009 Assigning strains to bacterial species via the internet. *BMC*

884   *Biol* **7**, 3.

885   57. Didelot, X. *et al.*. 2011 Recombination and population structure in *Salmonella*

886   *enterica*. *PLoS Pathog* **7**, e1002191.

887   58. den Bakker, H. C. *et al.*. 2011 Genome sequencing reveals diversification of

888   virulence factor content and possible host adaptation in distinct

889   subpopulations of *Salmonella enterica*. *BMC Genomics* **12**, 425.

890   59. Timme, R. E., Pettengill, J. B., Allard, M. W., Strain, E., Barrangou, R.,

891   Wehnes, C., Van Kessel, J. S., Karns, J. S., Musser, S. M., and Brown, E. W.

892   2013 Phylogenetic diversity of the enteric pathogen *Salmonella enterica*

893   subsp. *enterica* inferred from genome-wide reference-free SNP characters.

894   *Genome Biol Evol* **5**, 2109-2123. (doi:evt159 [pii];10.1093/gbe/evt159 [doi])

895   60. Parsons, S. K., Bull, C. M., and Gordon, D. M. 2011 Substructure within

896   *Salmonella enterica* subspecies *enterica* isolated from Australian wildlife.

897   *Appl Environ Microbiol* **77**, 3151-3153.

61. Worley, J., Meng, J., Allard, M. W., Brown, E. W., and Timme, R. E. 2018
Salmonella enterica phylogeny based on whole-genome sequencing reveals
two new clades and novel patterns of horizontally acquired genetic elements.
MBio **9**. (doi:mBio.02303-18 [pii];10.1128/mBio.02303-18 [doi])

62. Key, F. M. *et al.*. 2020 Emergence of human-specific *Salmonella enterica* is
linked to the Neolithization process. *Nature Ecology & Evolution* **4**, 324-333.
(doi:https://doi.org/10.1038/s41559-020-1106-9)

63. Guibourdenche, M., Roggentin, P., Mikoleit, M., Fields, P. I., Bockemuhl, J.,
Grimont, P. A., and Weill, F.-X. 2010 Supplement 2003-2007 (No. 47) to the
White-Kauffmann-Le Minor scheme. *Res Microbiol* **161**, 26-29.

64. Issenhuth-Jeanjean, S., Roggentin, P., Mikoleit, M., Guibourdenche, M., De, P.
E., Nair, S., Fields, P. I., and Weill, F.-X. 2014 Supplement 2008-2010 (no.
48) to the White-Kauffmann-Le Minor scheme. *Res Microbiol* **165**, 526-530.
(doi:S0923-2508(14)00106-5 [pii];10.1016/j.resmic.2014.07.004 [doi])

65. Grimont, P. A. and Weill, F.-X. 2007 *Antigenic formulae of the Salmonella*
serovars--9th edition edition. Paris, France: WHO Collaborating Centre for
Reference and Research on *Salmonella*.

66. Seif, Y., Monk, J. M., Machado, H., Kavvas, E., and Palsson, B. O. 2019
Systems biology and pangenome of *Salmonella* O-Antigens. *MBio* **10**.
(doi:mBio.01247-19 [pii];10.1128/mBio.01247-19 [doi])

67. Zhang, S., den Bakker, H. C., Li, S., Chen, J., Dinsmore, B. A., Lane, C., Lauer,
A. C., Fields, P. I., and Deng, X. 2019 SeqSero2: rapid and improved
Salmonella serotype determination using whole genome sequencing data.
Appl Environ Microbiol. (doi:AEM.01746-19 [pii];10.1128/AEM.01746-19
[doi])

68. Robertson, J., Yoshida, C., Kruczkiewicz, P., Nadon, C., Nichani, A., Taboada,
E. N., and Nash, J. H. E. 2018 Comprehensive assessment of the quality of

Salmonella whole genome sequence data available in public sequence

databases using the *Salmonella in silico* Typing Resource (SISTR). *Microb*

Genom, 1-11. (doi:10.1099/mgen.0.000151 [doi])

69.   Uelze, L., Borowiak, M., Deneke, C., Szabo, I., Fischer, J., Tausch, S. H., and

Malorny, B. 2020 Performance and accuracy of four open-source tools for *in*

silico serotyping of *Salmonella* spp. based on whole-genome short-read

sequencing data. *Appl Environ Microbiol* **86**. (doi:AEM.02265-19

[pii];10.1128/AEM.02265-19 [doi])

70.   Romano, S., Vinh, N. X., Bailey, J., and Verspoor, K. 2016 Adjusting for

chance clustering comparison measures. *Journal of Machine Learning*

Research **17**, 1-32.

71.   Chaudhuri, R. R. and Henderson, I. R. 2012 The evolution of the *Escherichia*

coli phylogeny. *Infect Genet Evol* **12**, 214-226.

72.   Ochman, H. and Selander, R. K. 1984 Standard reference strains of *Escherichia*

coli from natural populations. *J Bacteriol* **157**, 690-693.

73.   Selander, R. K., Caugant, D. A., and Whittam, T. S. 1987 Genetic structure and

variation in natural populations of *Escherichia coli*. In *Escherichia coli and*

Salmonella typhimurium cellular and molecular biology Vol.II (eds. F. C.

Neidhardt, J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H.

E. Umbarger), pp. 1625-1648. Washington, D.C.: American Society for

Microbiology.

74.   Touchon, M., Perrin, A., de Sousa, J. A. M., Vangchhia, B., Burn, S., O'Brien,

C. L., Denamur, E., Gordon, D., and Rocha, E. P. 2020 Phylogenetic

background and habitat drive the genetic diversification of Escherichia coli.

PLoS Genet **16**, e1008866. (doi:10.1371/journal.pgen.1008866

[doi];PGENETICS-D-20-00247 [pii])

75.   Denamur, E., Clermont, O., Bonacorsi, S., and Gordon, D. 2021 The population
genetics of pathogenic Escherichia coli. *Nat Rev Microbiol* **19**, 37-54.
(doi:10.1038/s41579-020-0416-x [doi];10.1038/s41579-020-0416-x [pii])

76.   Gordon, D. M., Clermont, O., Tolley, H., and Denamur, E. 2008 Assigning
Escherichia coli strains to phylogenetic groups: multi-locus sequence typing
versus the PCR triplex method. *Environ Microbiol* **10**, 2484-2496.

77.   Clermont, O., Christenson, J. K., Denamur, E., and Gordon, D. M. 2013 The
Clermont *Escherichia coli* phylo-typing method revisited: improvement of
specificity and detection of new phylo-groups. *Environ Microbiol Rep* **5**, 58-
65. (doi:10.1111/1758-2229.12019 [doi])

78.   Beghain, J., Bridier-Nahmias, A., Le, N. H., Denamur, E., and Clermont, O.
2018 ClermonTyping: an easy-to-use and accurate *in silico* method for
Escherichia genus strain phylotyping. *Microb Genom* **4**.
(doi:10.1099/mgen.0.000192 [doi])

79.   Waters, N. R., Abram, F., Brennan, F., Holmes, A., and Pritchard, L. 2020 Easy
phylotyping of *Escherichia coli* via the EzClermont web app and command-
line tool. In (Anon.), pp. 1-5.

80.   Walk, S. T. 2015 The "Cryptic" Escherichia. *EcoSal Plus* **6**.
(doi:10.1128/ecosalplus.ESP-0002-2015 [doi])

81.   Shen, Z., Koh, X. P., Yu, Y., Woo, C. F., Tong, Y., and Lau, S. C. K. 2019
Draft genome sequences of 16 strains of *Escherichia* cryptic clade II isolated
from intertidal sediment in Hong Kong. *Microbiol Resour Announc* **8**.
(doi:8/29/e00416-19 [pii];10.1128/MRA.00416-19 [doi])

82.   van der Putten, B. C. L., Matamoros, S., COMBAT consortium, and Schultsz,
C. 2021 *Escherichia ruysiae* sp. *nov.,* a novel Gram-stain-negative bacterium,
isolated from a faecal sample of an international traveller. *Int J Syst Evol*
Microbiol **71**, 004609. (doi:DOI 10.1099/ijsem.0.004609)

83.   Wick, L. M., Qi, W., Lacher, D. W., and Whittam, T. S. 2005 Evolution of
979   genomic content in the stepwise emergence of *Escherichia coli* O157:H7. *J*
980   *Bacteriol* **187**, 1783-1791.

981   84.   Eichhorn, I., Semmler, T., Mellmann, A., Pickard, D., Anjum, M. F., Fruth, A.,
982   Karch, H., and Wieler, L. H. 2018 Microevolution of epidemiological highly
983   relevant non-O157 enterohemorrhagic *Escherichia coli* of serogroups O26
984   and O111. *Int J Med Microbiol* **308**, 1085-1095. (doi:S1438-4221(18)30241-
985   8 [pii];10.1016/j.ijmm.2018.08.003 [doi])

986   85.   Liu, C. M. *et al.*. 2018 *Escherichia coli* ST131-H22 as a foodborne uropathogen.
987   *MBio* **9**. (doi:mBio.00470-18 [pii];10.1128/mBio.00470-18 [doi])

988   86.   Price, L. B. *et al.*. 2013 The epidemic of extended-spectrum-beta-lactamase-
989   producing *Escherichia coli* ST131 is driven by a single highly pathogenic
990   subclone, H30-Rx. *MBio* **4**, e00377-13. (doi:mBio.00377-13
991   [pii];10.1128/mBio.00377-13 [doi])

992   87.   Achtman, M., Mercer, A., Kusecek, B., Pohl, A., Heuzenroeder, M., Aaronson,
993   W., Sutton, A., and Silver, R. P. 1983 Six widespread bacterial clones among
994   *Escherichia coli* K1 isolates. *Infect Immun* **39**, 315-335.

995   88.   Gordon, D. M. *et al.*. 2017 Fine-scale structure analysis shows epidemic patterns
996   of Clonal Complex 95, a cosmopolitan *Escherichia coli* lineage responsible
997   for extraintestinal infection. *mSphere* **2**. (doi:10.1128/mSphere.00168-17
998   [doi];mSphere00168-17 [pii])

89.   Laukkanen-Ninios, R. *et al.*. 2011 Population structure of the *Yersinia*
pseudotuberculosis complex according to multilocus sequence typing.
Environ Microbiol **13**, 3114-3127.

90.   Gladstone, R. A. *et al.*. 2019 International genomic definition of pneumococcal
lineages, to contextualise disease, antibiotic resistance and vaccine impact.
EBioMedicine **43**, 338-346. (doi:doi: 10.1016/j.ebiom.2019.04.021)

91.   Lees, J. A., Harris, S. R., Tonkin-Hill, G., Gladstone, R. A., Lo, S. W., Weiser,
J. N., Corander, J., Bentley, S. D., and Croucher, N. J. 2019 Fast and flexible
bacterial genomic epidemiology with PopPUNK. *Genome Res* **29**, 304-316.
(doi:gr.241455.118 [pii];10.1101/gr.241455.118 [doi])

92.   Lees, J. A., Tonkin-Hill, G., Yang, Z., and Corander, J. 2021 Mandrake:
visualising microbial population structure by embedding millions of genomes
into a low-dimensional representation. *BioRxiv*, 2021. (doi:doi:
https://doi.org/10.1101/2021.10.28.466232)

93.   Zaharias, P. and Warnow, T. 2022 Recent progress on methods for estimating
and updating large phylogenies. In (Anon.).

94.   Tschoeke, D., Salazar, V. W., Vidal, L., Campeao, M., Swings, J., Thompson,
F., and Thompson, C. 2020 Unlocking the genomic taxonomy of the
Prochlorococcus collective. *Microb Ecol* **80**, 546-558. (doi:10.1007/s00248-
020-01526-5 [doi];10.1007/s00248-020-01526-5 [pii])

95.   Salazar, V. W., Tschoeke, D. A., Swings, J., Cosenza, C. A., Mattoso, M.,
Thompson, C. C., and Thompson, F. L. 2020 A new genomic taxonomy
system for the *Synechococcus* collective. *Environ Microbiol* **22**, 4557-4570.
(doi:10.1111/1462-2920.15173 [doi])

96.   Achtman, M. and Pluschke, G. 1986 Clonal analysis of descent and virulence
among selected *Escherichia coli*. *Annu Rev Microbiol* **40**, 185-210.

97.   Griffiths, D. *et al.*. 2010 Multilocus sequence typing of *Clostridium difficile*. *J*
Clin Microbiol **48**, 770-778.

98.   Zhou, Z., Alikhan, N.-F., Sergeant, M. J., Luhmann, N., Vaz, C., Francisco, A.
P., Carrico, J. A., and Achtman, M. 2018 GrapeTree: Visualization of core
genomic relationships among 100,000 bacterial pathogens. *Genome Res* **28**,
1395-1404. (doi:DOI: 10.1101/gr.232397.117)

99.   Achtman, M. and Zhou, Z. 2020 Metagenomics of the modern and historical

1032   human oral microbiome with phylogenetic studies on *Streptococcus mutans*

1033   and *Streptococcus sobrinus*. *Phil Trans R Soc B* **375**, 20190573.

1034   (doi:http://dx.doi.org/10.1098/rstb.2019.0573)

1035   100. Zhou, Z., Luhmann, N., Alikhan, N.-F., Quince, C., and Achtman, M. 2018

1036   Accurate reconstruction of microbial strains from metagenomic sequencing

1037   using representative reference genomes. In *RECOMB 2018*, pp. 225-240:

1038   Springer, Cham.

1039   101. Luhmann, N., Holley, G., and Achtman, M. 2021 BlastFrost: fast querying of

1040   100,000s of bacterial genomes in Bifrost graphs. *Genome Biol* **22**, 30.

1041   (doi:10.1186/s13059-020-02237-3 [doi];10.1186/s13059-020-02237-3 [pii])

1042   102. Pedregosa, F. *et al.*. 2011 Scikit-learn: Machine learning in python. *Journal of*

1043   *Machine Learning Research* **12**, 2825-2830.

1044   103. Bostock, M., Ogievetsky, V., and Heer, J. 2011 D^3: Data-Driven Documents.

1045   *IEEE Trans Vis Comput Graph* **17**, 2301-2309.

1046   (doi:10.1109/TVCG.2011.185 [doi])

1047

1048

Table 1. Legacy strains, newly assembled genomes and hierarchical cgST clusters in EnteroBase (09/2021).

Genus	Legacy MLST	cgMLST	# Genomic Loci		# HierCC clusters (HierCC level)		
	(# strains)	(# genomes)	wgMLST	cgMLST	ST Complexes	Lineages	Species/ Sub-species
Salmonella	4,930	312,196	21,065	3,002	3648 (900)	2185 (2000)	15 (2850)
Escherichia/ Shigella	9,525	175,256	25,002	2,512	1379 (1100)	343 (2000)	16 (2350)
Streptococcus		76,718	33,887	372	3681 (100)		132 (363)
Clostridioides		23,197	11,490	2,556	299 (950)		12 (2500)
Vibrio		12,267	152,249	1,128	2000 (800)		155 (1090)
Yersinia	4,286	4,023	19,591	1,553	451 (600)		34 (1490)

NOTE: # Genomic Loci: numbers of CDS that are automatically called for the wgMLST and cgMLST schemes. cut-off: maximum numbers of allelic differences between cgSTs that define HC clusters. EnteroBase automatically calculates Legacy MLST for all *Clostridioides* genomes according to the Griffiths scheme [97], but does not maintain a database of legacy data from ABI sequencing. Additional public databases are presented by EnteroBase for *Helicobacter* (>3500 genomes) and *Moraxella* (>2350 genomes), but these lack a cgMLST scheme. EnteroBase also has a database for >73,000 *Mycobacterium* genomes, but this is not yet publicly available, and still lacks a cgMLST scheme.

Table 2. EnteroBase-related software tools that support the analysis of large numbers of bacterial genomes

Name	Purpose	Citation	URL for Stand-alone version
GrapeTree	GUI for depicting and analysing minimal spanning and NJ trees of character data	[98]	https://github.com/achtman-lab/GrapeTree
PEPPAN	Pan-genome calculation including pseudo-genes from numerous representatives of an entire genus	[17]	https://github.com/zheminzhou/PEPPAN
SPARSE	Assignment of metagenomic reads to individual taxa	[99,100]	https://github.com/zheminzhou/SPARSE
EToKi	EnteroBase toolkit of useful functions and pipelines	[28]	https://github.com/zheminzhou/EToKi
BlastFrost	Efficient k-mer based search for and sequence extraction from genes in large datasets	[101]	https://github.com/nluhmann/BlastFrost
HierCC	Automated hierarchical clustering of cgSTs to existing or novel clusters	[32]	https://github.com/zheminzhou/pHierCC

NOTE: PEPPAN is a stand-alone programme that was used to generate the wgMLST schemes used in EnteroBase. SPARSE is a second stand-alone programme that has been used to extract taxon-specific reads from metagenomes of ancient DNA that were then used with EToKi to define pseudo-MAGs that were uploaded to EnteroBase for phylogenetic comparisons. HierCC was used as a stand-alone programme in development mode to define an initial set of clusters from a representative set of genomes. Subsequent assignments to existing clusters or to novel clusters were performed automatically within EnteroBase in production mode. BlastFrost is used by EnteroBase to determine the presence/absence of toxins and other pathovar characteristics of gastrointestinal pathogens within *E. coli*, and to assign gastrointestinal pathovar designations.

Table 3. Parameters of datasets used for calculating super-trees.

Genus	# Type strains	Total # Genomes	# Strict core genes	# Accessory genes	# Soft core genes (cgMLST)	# SNPs
Salmonella	7	10,002	1409	19,486	3002	1,410,331
Streptocococcus	84	5937	90	31,630	372	263,080
Clostridioides	1	6725	515	10,975	2556	725,240
Vibrio	119	5032	124	146,041	1128	776,439
Yersinia	23	1847	744	18,799	1553	656,143

Table 4. Quantitative concordance between eBG and HierCC clustering

Salmonella			
Test statistic	eBG-HC900	eBG-HC2000	HC900-HC2000
Adjusted Mutual Information Score (AMI)	0.985	0.948	0.942
Adjusted Rand Index (ARI)	0.982	0.856	0.866
Escherichia/Shigella			
Test statistic	ST Cplx-HC1100	ST Cplx-HC2000	HC1100-HC2000
Adjusted Mutual Information Score (AMI)	0.940	0.688	0.692
Adjusted Rand Index (ARI)	0.871	0.299	0.337
Streptococcus pneumoniae			
Test statistic	CC-HC100	GSPC-HC100	GSPC-HC160
Adjusted Mutual Information Score (AMI)	0.959	0.956	0.980
Adjusted Rand Index (ARI)	0.987	0.906	0.949

NOTE: *Salmonella*: Calculations were performed on 319,490 entries in EnteroBase which had been assigned to eBGs, HC900 and HC2000 clusters by December, 2021. New eBGs have not been created in recent years, and the dataset only represented 411 eBGs, 690 HC900 clusters and 312 HC2000 clusters.

Escherichia/Shigella: HC1100 and HC2000 assignments were from 143,520 genomes which had been assigned to ST Complexes as well as to HC1100 and HC2350 clusters in December, 2021. At that time 186,852 genomes had been assigned to HC1100 and HC2000 clusters, with AMI of 0.59 and ARI of 0.2.

*Streptococcus pneumonia*e consisted of GSPC assignments for 18147 genomes from Supplemental Table 2 and CC assignments for 13396 genomes from Supplemental Table 1 to Gladstone et al. [90] which were imported into User defined fields in EnteroBase as described [28].

ARI and AMI were calculated with the functions adjusted_rand_score() and adjusted_mutual_info_score() in the Python 3 library sklearn.metrics V1.0.1 [102].

Figure Legends

Figure 1. Species and sub-species assignments within *Salmonella* with HierCC and ANI. ML super-tree of 1,410,331 SNPs among 3002 core genes from 10,002 representative genomes of *Salmonella*. Former sub-species IIIa is designated *S. arizonae* in accordance with Pearce *et al.,* 2021 [48]. A. Partitions coloured by HC2850 clusters (legend). Arrows indicate a new species, *HC2850_215890* (five strains from the U.K., 2018-2020), and a new sub-species, HC2850_222931 (one strain from France, 2018). All other HC2850 clusters correspond to species (*S. bongori* and *S. arizonae*) or subspecies, except for HC2850_7171 (starred), which is subsp. *enterica* (I) according to the ML tree. B. Partitions differentiated by ANI 95% clusters (legend) correspond to species (*S. enterica, S. bongori, S. arizonae, S. HC2850_215890*), as indicated by arrows, and sub-species are not differentiated. An interactive version of this GrapeTree rendition can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53257. The corresponding presence/absence tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53258.

Figure 2. Species and sub-species assignments within *Clostridioides* according to ANI (A) and HierCC (B). ML super-tree of 725,240 SNPs among 2556 core genes from 6,724 representative genomes of *Clostridioides difficile* and one genome of *Clostridioides mangenotii.* A. ANI 95% clusters (legend) differentiate *C. mangenotii* (cluster 1010) and five other clusters (clusters 255, 1370, 373, 2147, 1011) from *C. difficile* (cluster 0). As indicated, four of these correspond to cryptic clades C-I [clusters 255, 373], C-II [cluster 1370] and C-III [cluster 2147]) according to Knight *et al.* [49]. Arrows indicate two additional clusters that were distinguished by HierCC

in part B. B. Partitions coloured by HierCC assigns the same genomes as ANI to the

corresponding HC2500 clusters, except that it defines HC2500_15334 and

HC2500_15408 to one genome each, which might represent sub-species. An

interactive version of this GrapeTree rendition can be found at

https://enterobase.warwick.ac.uk/ms_tree?tree_id=53253 and a presence/absence tree

can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53254.

Figure 3. Comparison of HC363 clusters with taxonomic designations in

Streptococcus. ML super-tree of 263,080 SNPs among 372 core genes from 5937

representative *Streptococcus* genomes. Species names are indicated next to the

phylogenetic clusters where genomes mapped for type strains, and according to

published metadata. Nodes are colored by HC363 clusters, and exceptional

assignments are indicated by asterisks next to *S. pneumoniae* and *S.*

pseudopneumoniae, which were both HC363_99; multiple phylogenetic and HierCC

clusters grouped within *S. suis*; *S. salivarius* and *S. vestibularis*, which were both

HC353_202; *S. lutetiensis* and *S. equinus*, which were both HC363_181; and *S.*

dysgalactiae and *S. pyogenes*, which were both HC363_139. An interactive version of

the GrapeTree rendition of the SNP tree can be found at

https://enterobase.warwick.ac.uk/ms_tree?tree_id=53261 and the presence/absence

tree at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53262.

Figure 4. Hierarchical population structure of O serogroups in *Salmonella*.

Hierarchical bubble plot of species/subspecies (HC2850; light grey circles), Lineages

(HC2000; dark grey circles) and eBurst groups (HC900; other O:group specific

colours) of 274,824 *Salmonella* genomes in 790 HC900 clusters for which a

consensus O serogroup could be deduced by metadata, or bioinformatic analyses with

SeqSero V2 [67] or SISTR 1.1.1 [68]. Additional information is indicated by yellow

text for selected HC2000 and HC900 circles which are specifically mentioned in the

Discussion. The diameters of HC900 circles are proportional to the numbers of

genomes. An interactive version of this figure can be found at

https://observablehq.com/@laurabaxter/salmonella-serovar-piechart from which the

representation, raw data and d3 Java code [103] for generating the plot can be

downloaded.

Figure 5. Hierarchical population structure of O serogroups in *Escherichia*

coli/Shigella. Hierarchical bubble plot for 167,312 genomes of *Escherichia coli* or

Shigella in HC2350_1 (large grey circle) that were available in EnteroBase in April,

2021. Seven HC2000 Lineages encompassing 15 HC1100/ST Complexes of *Shigella*

are arbitrarily indicated at the right. HC2350_1 also includes 15 other *E. coli* HC2000

Lineages that encompass 144 other HC1100 clusters that each contains at least 50 *E.*

coli genomes. The figure shows only the HC1100 clusters and not the corresponding

HC1100 clusters. Numbers of genomes assigned to individual O serogroups (legend)

are indicated by pie chart wedges within the HC1100 circles. Selected HC1100

clusters are drawn on larger scale with indications of phenotype and nomenclature

outside the main circle, connected to the original circles by lines. An interactive

version of this figure can be found at

https://observablehq.com/@laurabaxter/escherichia-serovar-piechart, from which the

representation, raw data and d3 Java code [103] for generating the plot can be

downloaded.

Supplemental Material

Supplemental Text:

Correspondence between species designations and ANI or HierCC

Yersinia

Vibrio

Historical reconstructions

Sample bias

Methods

Collections of representative genome sequences

ML super-trees

ASTRID guide tree

Frequencies of O groups in *E. coli* and *Salmonella*

Supplemental Table S1. Consistency and inconsistency between HC1090 and
ANI95% in *Vibrio*.

Supplemental Table S2. Instances within *Vibrio* of one HC1090 cluster with
multiple 95% ANI clusters.

Supplemental Table S3. Instances within *Vibrio* of one 95% ANI cluster vs multiple
HC1090 clusters.

Supplemental Table S5. Consistent clustering in *Streptococcus* for 102 clusters
according to HC363 and ANI95%.

Supplemental Table S6. One HC363 cluster vs multiple ANI95% clusters in
Streptococcus.

Supplemental Table S7. One 95% ANI cluster vs two HC363 clusters in

Streptococcus.

Supplemental Table S8. Lookup table for *Salmonella* from HC2000 through HC900

to Serovar.

Supplemental Table S9. Detailed analysis of O groups in 790 HC900 clusters of

Salmonella with at least 5 entries.

Supplemental Table S10 (Supplemental Text). Lookup table from Serovar name to

antigenic epitopes and O:groups (Institut Pasteur, Paris, with permission from F.X.

Weill).

Supplemental Figure S1. Species and sub-species assignments to 1847

representative genomes of *Yersinia* with 95% ANI (A,C) and HC1490 (B,D).

Maximum Likelihood ML presence/absence trees of 18,799 accessory genes (A,B) or

656,143 SNPs within 1553 soft core cgMLST genes (C,D). The most common color

codes are indicated in legends next to each plot. Within each tree, curated species

names in EnteroBase are indicated next to the appropriate clusters. Note that *Y.*

enterocolitica is split into four species by HC1490 cluster as confirmed by 95% ANI

clustering. Similarly, asterisks in A and C indicate sub-species *similis*, *wautersii* and

pestis of the *Y. pseudotuberculosis* Complex that were previously assigned to the

invalid former species *Y. similis, Y. wautersii* and *Y. pestis.* The novel *Yersinia*

species *HC1490_1007, HC1490_440, HC1490_435,* and *HC1490_449* are also

included in the figure. Further details can be found in Supplemental Text and

interactive versions of the GrapeTree renditions can be found at

https://enterobase.warwick.ac.uk/ms_tree?tree_id=53270   (A,B)   and

https://enterobase.warwick.ac.uk/ms_tree?tree_id=53269 (C,D).

Supplemental Figure S2. Curated species, clade, HC1090 and ANI95% assignments

to 5032 representative genomes of *Vibrio*. ML SNP super-tree of 776, 439 SNPs

among 1128 soft core genes, colour-coded by HC1090 cluster The most common

colour codes are indicated in the legend at the bottom. The scope of three higher order

groups of species, Cholerae, Harveyii and Splendidus, that were defined by MLSA by

Gomez-Gil *et al.,* 2014 are indicated by arcs at the periphery of the plot. Further

details can be found in Supplemental Text and interactive version of the GrapeTree

renditions can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53265

(SNP   tree)   and   https://enterobase.warwick.ac.uk/ms_tree?tree_id=53266

(presence/absence tree).

Supplemental Figure S3. Comparison of HC363 clusters with taxonomic

designations in *Streptococcus* in an ML super-tree of 31,630 accessory genes from

5937 representative *Streptococcus* genomes. Species names are indicated next to the

phylogenetic clusters where genomes mapped together with type strains, and

according to published metadata. Nodes are colored by HC363 clusters, and

exceptional assignments are indicated by asterisks next to *S. mitis*, which could not be

distinguished by HC363 clustering from *S. pneumoniae* or *S. pseudopneumoniae,*

multiple phylogenetic and HierCC clusters within *S. suis*; *S. salivarius* and *S.*

vestibularis, which were both HC353_202; *S. lutetiensis* and *S. equinus*, which were

both HC363_181; and *S. dysgalactiae* and *S. pyogenes*, which were both HC363_139.

An interactive version of this GrapeTree rendition can be found at

https://enterobase.warwick.ac.uk/ms_tree?tree_id=53262.

Supplemental Figure S4. GrapeTree NJ plot of cgMLST allelic distances between

18,147 genomes of *Streptococcus pneumoniae* which had been assigned to GSPC

clusters [90]. An interactive version of this figure may be examined at

https://enterobase.warwick.ac.uk/ms_tree?tree_id=67948 and color-coded by HC100

cluster level, legacy Clonal Complex (CC), GPSC designation of HC100 cluster

number or by other metadata stored for those genomes within EnteroBase. Key

Legend: HC160 cluster numbers and their color-codes. Legend: Measure of 40 allelic

distances.

Supplemental Figure S5. Cartoon of the workflow of cgMLSA

(https://github.com/zheminzhou/cgMLSA). (A) Construct gene trees for each core

gene with RaXML (B) Build an initial guide super-tree with ASTRID and calculate

local branch supports with ASTRAL. (C) Split the initial guide tree into pairs of

disjoint subsets at each highly supported branch, and append an artificial branch

(dotted black line) to both members of the subset to allow later reconstruction of this

branch. (D) Split all gene trees into the same sets of genomes for each pair of disjoint

subsets. (E) Calculate a constraint super-tree for each subset with ASTRAL. (F)

Construct the final super-tree by merging constraint trees via the connections marked

in C.

Figure 1. Species and sub-species assignments within *Salmonella* with HierCC and ANI. ML super-tree of 1,410,331 SNPs among 3002 core genes from 10,002 representative genomes of *Salmonella*. Former sub-species IIIa is designated *S. arizonae* in accordance with Pearce et al., 2021 [48]. A. Partitions coloured by HC2850 clusters (legend). Arrows indicate a new species, HC2850_215890 (five strains from the U.K., 2018-2020), and a new sub-species, HC2850_222931 (one strain from France, 2018). All other HC2850 clusters correspond to species (*S. bongori* and *S. arizonae*) or subspecies, except for HC2850_7171 (starred), which is subsp. *enterica* (I) according to the ML tree. B. Partitions differentiated by ANI 95% clusters (legend) correspond to species (*S. enterica, S. bongori, S. arizonae, S.* HC2850_215890), as indicated by arrows, and sub-species are not differentiated. An interactive version of this Grape-Tree rendition can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53257. The corresponding presence/absence tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53258.

Figure 2. Species and sub-species assignments within *Clostridioides* according to ANI (A) and HierCC (B). ML super-tree of 725,240 SNPs among 2556 core genes from 6,724 representative genomes of *Clostridioides difficile* and one genome of *Clostridioides mangenotii*. A. ANI 95% clusters (legend) differentiate *C. mangenotii* (cluster 1010) and five other clusters (clusters 255, 1370, 373, 2147, 1011) from *C. difficile* (cluster 0). As indicated, four of these correspond to cryptic clades C-I [clusters 255, 373], C-II [cluster 1370] and C-III [cluster 2147]) according to Knight et al. [49]. Arrows indicate two additional clusters that were distinguished by HierCC in part B. B. Partitions coloured by HierCC assigns the same genomes as ANI to the corresponding HC2500 clusters, except that it defines HC2500_15334 and HC2500_15408 to one genome each, which might represent sub-species. An interactive version of this GrapeTree rendition can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53253 and a presence/absence tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53254.

Figure 3. Comparison of HC363 clusters with taxonomic designations in *Streptococcus*. ML super-tree of 263,080 SNPs among 372 core genes from 5937 representative *Streptococcus* genomes. Species names are indicated next to the phylogenetic clusters where genomes mapped for type strains, and according to published metadata. Nodes are colored by HC363 clusters, and exceptional assignments are indicated by asterisks next to *S. pneumoniae* and *S. pseudopneumoniae,* which were both HC363_99; multiple phylogenetic and HierCC clusters grouped within *S. suis*; *S. salivarius* and *S. vestibularis*, which were both HC353_202; *S. lutetiensis and S. equinus*, which were both HC363_181; and *S. dysgalactiae and S. pyogenes*, which were both HC363_139. An interactive version of the GrapeTree rendition of the SNP tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53261 and the presence/absence tree at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53262.

O antigens

Figure 4. Hierarchical population structure of O serogroups in *Salmonella*. Hierarchical bubble plot of species/subspecies (HC2850; light grey circles), Lineages (HC2000; dark grey circles) and eBurst groups (HC900; other O:group specific colours) of 274,824 *Salmonella* genomes in 790 HC900 clusters for which a consensus O serogroup could be deduced by metadata, or bioinformatic analyses with SeqSero V2 [67] or SISTR 1.1.1 [68]. Additional information is indicated by yellow text for selected HC2000 and HC900 circles which are specifically mentioned in the Discussion. The diameters of HC900 circles are proportional to the numbers of genomes. An interactive version of this figure can be found at https://observablehq.com/@laurabaxter/salmonella-serovar-piechart from which the representation, raw data and d3 Java code [103] for generating the plot can be downloaded.

HC1100_2
(ST29 Cplx;
EHEC)

O103
O111
O26

HC1100_44
(ST95 Cplx;
NMEC, UPEC, APEC)

O45
O25
O1
O2
O18

B2 B4 D13
D3
D9
F6
B1
B18

HC2000_1463
S. dysenteriae

HC2000_1465 (ST243 Cplx)
S. flexneri, boydii
& dysenteriae

HC2000_305
(ST152 Cplx) *S. sonnei*

HC2000_192
(ST245 Cplx)

O55

HC1100_63
(ST11 Cplx;
EHEC)

O45
O128 O111
HC1100_3
(ST20 Cplx;
EHEC)

HC1100_7
(ST131 Cplx;
UPEC)

O antigens

■ B1	■ D9	■ O13	■ O25	■ O83
■ B13	■ F1-5	■ O145	■ O26	■ O86
■ B16	■ F6	■ O146	■ O4	■ O88
■ B18	■ SS	■ O15	■ O45	■ O89
■ B2	■ O1	■ O153	■ O5	■ O9
■ B4	■ O102	■ O157	■ O55	■ O91
■ B9	■ O103	■ O16	■ O6	■ Others
■ D1	■ O11	■ O17	■ O7	□ Missing
■ D13	■ O111	■ O18	■ O75	
■ D2	■ O121	■ O2	■ O78	
■ D3	■ O128	■ O21	■ O8	

Figure 5. Hierarchical population structure of O serogroups in *Escherichia coli/Shigella*. Hierarchical bubble plot for 167,312 genomes of *Escherichia coli* or *Shigella* in HC2350_1 (large grey circle) that were available in EnteroBase in April, 2021. Seven HC2000 Lineages encompassing 15 HC1100/ST Complexes of *Shigella* are arbitrarily indicated at the right. HC2350_1 also includes 15 other *E. coli* HC2000 Lineages that encompass 144 other HC1100 clusters that each contains at least 50 *E. coli* genomes. The figure shows only the HC1100 clusters and not the corresponding HC1100 clusters. Numbers of genomes assigned to individual O serogroups (legend) are indicated by pie chart wedges within the HC1100 circles. Selected HC1100 clusters are drawn on larger scale with indications of phenotype and nomenclature outside the main circle, connected to the original circles by lines. An interactive version of this figure can be found at https://observableh-q.com/@laurabaxter/escherichia-serovar-piechart, from which the representation, raw data and d3 Java code [103] for generating the plot can be downloaded.