EnteroBase: Hierarchical clustering of 100,000s of bacterial genomes

into species/sub-species and populations

Mark Achtman*, Zhemin Zhou*,a, Jane Charlesworth and Laura Baxter

University of Warwick, Coventry CV4 7AL, UK

aCurrent address: Pasteurien College, Soochow University, No.199 Ren'ai Road,

Suzhou, Jiangsu 215123, China.

*Corresponding authors: MA, m.achtman@warwick.ac.uk; ZZ, zheminzhou@qq.com

MA, ORCID 0000-0001-6815-0070; ZZ, ORCID 0000-0001-9783-0366; JC, ORCID

0000-0001-9759-9838; LB, ORCID 0000-0003-3287-7547

Word count: 7559

Word Count incl. 103 Citations 10846

Plus:

4 Tables

5 Figures

Abstract

The definition of bacterial species is traditionally a taxonomic issue while bacterial populations are identified by population genetics. These assignments are species specific, and depend on the practitioner. Legacy multilocus sequence typing is commonly used to identify sequence types (STs) and clusters (ST Complexes). However, these approaches are not adequate for the millions of genomic sequences from bacterial pathogens that have been generated since 2012. EnteroBase (http://enterobase.warwick.ac.uk) automatically clusters core genome MLST allelic profiles into hierarchical clusters (HierCC) after assembling annotated draft genomes from short read sequences. HierCC clusters span core sequence diversity from the species level down to individual transmission chains. Here we evaluate HierCC's ability to correctly assign 100,000s of genomes to the species/subspecies and population levels for *Salmonella, Escherichia, Clostridoides, Yersinia, Vibrio* and *Streptococcus*. HierCC assignments were more consistent with maximum-likelihood super-trees of core SNPs or presence/absence of accessory genes than classical taxonomic assignments or 95% ANI. However, neither HierCC nor ANI were uniformly consistent with classical taxonomy of *Streptococcus*. HierCC was also consistent with legacy eBGs/ST Complexes in *Salmonella* or *Escherichia* and with O serogroups in *Salmonella*. Thus, EnteroBase HierCC supports the automated identification of and assignment to species/subspecies and populations for multiple genera.

1. Introduction

"Microbiologists need large databases to identify and communicate about clusters of related bacteria, … Such databases should contain the reconstructed genomes of bacterial isolates,… together with metadata describing their sources and phenotypic properties… and we are currently developing EnteroBase, a genome-based successor to the *E. coli* and *S. enterica* MLST databases." Excerpted from Achtman and Zhou, 2014 [1].

The Linnean system of genus and species designations was applied to bacterial nomenclature in the late 19[th] century, soon after their ability to cause infectious diseases had been recognised. These taxonomic labels initially reflected disease specificity and phenotypic similarities. Over the following decades, phenotypic criteria for distinguishing bacterial taxa were extended to include serologically distinct groupings and even differential sensitivity to bacteriophages. A primary goal for taxonomic designations for bacterial pathogens was that they be useful for epidemiology and clinical diagnoses. As a result, serovars of *Salmonella enterica*, which differ by dominant epitopes on lipopolysaccharide and flagella, were each assigned a distinct species designation, often referring to the disease syndrome and host e.g. *Salmonella typhimurium* [2]. Similarly, even though *Yersinia pestis* is a clone of *Yersinia pseudotuberculosis* [3,4], it was designated as a distinct species because *Y. pestis* causes plague whereas *Y. pseudotuberculosis* causes gastroenteritis. The use of species designations for *Salmonella* serovars is now disparaged, although it is still in common use [5], but *Y. pestis* has retained its species designation.

Percentage DNA-DNA hybridization levels became the "Gold Standard" metric for new taxonomic designations after 1987 [6]. DNA-DNA hybridization reflects

63 genomic relationships but distinctive phenotypic differences have remained a

64 requirement for the definition of a novel species until the present. Indeed, the

65 international committee which controls taxonomic designations continues to reject the

66 validity of taxonomic designations based solely on DNA similarities [7,8].

a) ANI and MLST

An alternative genomic approach for taxonomic designations was proposed in

2005 by Konstantinidis [9], namely the definition of species on the basis of average

nucleotide identity (ANI). Pairs of genomes with 95% ANI usually belong to the same

species whereas pairs with lower ANI values belong to different species.

Computational methods based on k-mer searches, such as FastANI [10], can rapidly

perform ANI-like calculations on large numbers of genomes, and these calculations

are now routinely used by bioinformaticians for taxonomic assignments. An approach

based on ANI is enticing because it could provide defined criteria for the definition of

a species across all Bacteria, and allow species assignment based exclusively on

genome sequences. However, the use of ANI for species assignments has been

criticized because it does not completely correlate with DNA-DNA hybridization

[11,12]. Furthermore, multiple taxa that are designated as single species each contain

multiple 95%ANI groups [13,14]. For example, strains of *Streptococcus mitis*

colonise the human oropharyngeal tract, have similar phenotypes and are considered

to represent a single species. However, *S. mitis* encompasses myriad, genetically

distinct strains [15-17], and encompasses at least 44 distinct 95% ANI clusters [17].

Other problematic genera include *Pseudomonas* [12,18] and *Aeromonas* [19].

Furthermore, large databases including 100,000s of genomes per genus would

struggle to implement methods such as pairwise clustering by ANI because each new

entry would require testing against all genomes. There is also no consensus on ANI

criteria for recognizing lower taxonomic entities, such as sub-species and populations.

Indeed, there is not even a consensus on the definitive properties of what constitutes a

bacterial species [20].

Bottom-up population genetic approaches such as multilocus sequence typing

(MLST) can provide an alternative to top-down taxonomy, and deal efficiently with

large numbers of bacterial strains. Legacy MLST based on the sequence differences

of several housekeeping genes was introduced in 1998 [21], and has now been applied

to more than 100 bacterial species [22]. MLST defines sequence types (STs),

consisting of unique integer designations for each unique sequence (allele) of each of

the MLST loci. Some STs mark individual clones with special pathogenic properties

and which seem to have arisen fairly recently, such as *Escherichia coli* ST131 which

is a globally prominent cause of urinary tract infection (UTI) and invasive disease

[23]. Similarly, *Salmonella enterica* subsp. *enterica* serovar Typhimurium ST313 is a

common cause of extra-intestinal, invasive salmonellosis in Africa [24]. Higher order

clusters of related STs are also well known. Such clusters can be recognized by eBurst

analyses [25], and are referred to as ST Complexes in *E. coli* [26] and eBGs (eBurst

groups) in *S. enterica* [27]. ST Complexes and eBGs seem to reflect natural

populations, but their broad properties are still not well understood.

The principles of legacy MLST were extended to rMLST, which uses sequences

of 53 universal bacterial genes encoding ribosomal proteins [28] and provides a

universal MLST scheme for all Bacteria [28]. For both *Escherichia* and *Salmonella,*

rMLST offers a slight improvement in resolution over legacy MLST, and the

identification of *Salmonella* eBGs is reasonably consistent across both approaches

[29]. rMLST has been used to identify tractable sets of representative genomes from

large collections for calculating phylogenetic trees [28] and pan-genomes [17].

MLST has also been extended to cgMLST, which encompasses all the genes in

a soft core genome [29-31], and cgMLST nomenclatures have been implemented for

multiple bacterial genera [28]. The large number of loci encompassed by cgMLST

schemes results in enormous numbers of cgMLST STs (cgSTs), but these can be

clustered into groups of bacterial genomes at multiple levels of genomic diversity

(hierarchical clustering) [32]. cgMLST provides considerably higher resolution than

legacy MLST or rMLST, and initial analyses indicated that it is ideal for investigating

transmission chains within single source outbreaks or for identifying population

structures up to the genus level [28]. Here we focus on the automated assignments of

genome assemblies from six genera of important bacterial pathogens to taxonomic

and population structures by hierarchical clustering of cgSTs with the EnteroBase

HierCC pipeline (Table 1) [32]. For five of those genera, HierCC is a full solution for

taxonomic designations of species.

b) EnteroBase and HierCC

EnteroBase (http://enterobase.warwick.ac.uk) includes tools for automatic

downloading of short read sequences and their metadata from the public domain,

assembly into draft genomes and population genetic analyses of the core and

accessory genomes (Table 2) [28]. Draft genomes are annotated according to genus-

wide pan-genome schemes created with PEPPAN [17]. In September 2021,

EnteroBase contained over 600,000 draft genomes from the six genera in Table 1, and

likely provides a nearly comprehensive overview of their global diversity. Many of

these samples reflect a focus on food-borne disease in the U.S. and United Kingdom,

but this bias is increasingly being reduced by the global sources of many genomes

(Supplemental Text, Sample Bias).

One of the primary goals for EnteroBase was a hierarchical overview of the

population structure of the genera in Table 1. We therefore developed HierCC [32]

based on cgMLST assignments to support the rapid recognition and detailed

investigation of differing levels of population structure. EnteroBase reports cluster

assignments and designations at 10-13 levels of allelic differences for all six genera

[28] (Table 1). HC5 – HC10 clusters with maximal internal pair-wise distances within

minimal spanning trees of 5 or 10 alleles, respectively, have been used to identify

short-term, single source outbreaks of *S. enterica* and *E. coli/Shigella* that extended to

multiple European countries [33-37].. Pathogen species can also include higher level

clusters, which can correspond to somewhat more distantly related bacterial

populations that cause endemic or epidemic disease over longer time periods in one or

more countries [38-41]. EnteroBase HierCC has even been used to classify all

Shigella [42], which correspond to discrete Lineages of *E. coli* [26,43], and to define

novel sub-species of *S. enterica* [44].

Classical taxonomic approaches for assigning individual strains and genomes to

species depend on human expertise, and are not suitable for automated pipelines in

real-time databases such as EnteroBase. This problem is acute because many short

read sequences in ENA have incorrect taxonomic assignments, or none at all, and

phenotypic distinctions are inappropriate for databases containing 100,000s of

genomes. Zhou *et al.* compared taxonomic designations with peak normalized mutual

information and silhouette scores for HierCC clusters, and identified HierCC levels

that corresponded with well-defined species/sub-species in each genus [32]. For

example, for *Salmonella* with a total of 3002 loci in the cgMLST scheme, HC2850

(94.9%) was chosen as the optimal HC level for identifying species and sub-species.

The optimal HC levels for the other five genera with cgMLST schemes ranged from

HC363 – HC2500 (93.6-97.8% of all cgMLST alleles) (Table 1). Here we compare

the consistency of those HierCC assignments with assignments based on classical

taxonomic approaches and 95% ANI. The results illustrate problems with classical

approaches and with 95% ANI, and we conclude that HierCC is preferable for

automated assignment of genomes to species/subspecies for those genera. However,

neither ANI nor HierCC is universally satisfactory for species/sub-species

assignments within *Streptococcus*. This manuscript also provides an initial overview

of the abilities of HierCC to assign genomes to populations and Lineages [45,46]

within *Salmonella* and *Escherichia*/*Shigella*, and compares those assignments with the

distributions of O antigens within lipopolysaccharide.

Results

a. **Species and sub-species**

In order to test the efficacy of HierCC at identifying species and sub-species, we

extracted collections of representative genomes from all six genera in EnteroBase for

which HierCC clusters had been implemented (Table 3). We wished to calculate

maximum likelihood phylogenetic topologies of these genomes based on their core

genome SNPs or the presence/absence of accessory genes from the pan-genome. Such

ML trees are very slow to calculate with large datasets, but disjointed tree merging

(DTM) within a divide-and-conquer approach allows large phylogenetic trees to be

calculated in a reasonable time [47]. We therefore developed cgMLSA (see Methods

in Supplemental Text), a novel DTM approach based on ASTRID [48] and ASTRAL

[49] that enabled the calculations of ML super-trees from up to 10,000 representative

genomes within hours to days. and applied it to each of the datasets. The resulting ML

super-trees were annotated with taxonomic designations and cluster designations

according to a 95% ANI cutoff calculated with FastANI [10] (95% ANI clusters). We

then compared those annotations with cluster assignments based on the species-

specific HierCC levels in Table 1.

Salmonella. The topologies of ML super-trees based on 1,410,331 core SNPs from

10,002 representative genomes were in large part concordant with traditional

taxonomic assignments, and with clustering according to 95% ANI (Fig. 1A) or

HC2850 (Fig. 1B). The three sets of assignments were also largely concordant with

the topology of an ML super-tree based on the presence or absence of accessory genes

(https://enterobase.warwick.ac.uk/ms_tree/53258). HC2850 and 95%ANI

distinguished *S. enterica* from *S. bongori*, and from former subspecies IIIa, which was

recently designated *S. arizonae* by Pearce *et al*. [50]. HC2850 also identified another

new *Salmonella* species, cluster HC2850_215890, and that identification was

confirmed by 95% ANI. HC2850_215890 consists of five strains that have been

isolated from humans since 2018 in the U.K., and a gene for gene comparison

indicated that roughly half of their core genes were more similar to *S. enterica* and the

other half to *S. bongori*.

95% ANI did not detect any subspecies structure within *S. enterica* whereas HC2850

differentiated a total of 12 clusters that corresponded to distinct phylogenetic branches

in the ML super-tree (Fig. 1B). Most of these have been designated as subspecies by

traditional taxonomy [44,50], except for a singleton genome in HC2850_222931,

which likely represents a novel subspecies. Despite the correct identification of so

many sub-species, HC2850 did not support Pearce's definition of subspecies XI, and

did not distinguish those genomes from subsp. II. Similarly, HC2850 did not

differentiate Pearce's subsp. VIII from subsp. IV. The prior definition of these

subspecies was largely dependent on branch topologies within phylogenetic trees

based on rMLST loci, and was also contradicted by population statistics [50].

HierCC also differentiated HC2850_7171, which consists of a short phylogenetic

branch within subsp. I that would not have been assigned to a subspecies according to

its ML topology. Genomic comparisons of multiple single genes from genomes within

HC2850_7171 suggest that HC2850_7171 may be a hybrid between subsp. I and II

because some gene sequences were most similar to the former and others were most

similar to the latter. With this sole possible exception, HierCC seems to be suitable for

the automated detection of new *Salmonella* species and subspecies, and for routinely

assigning novel genomes to the appropriate taxa.

Escherichia. In addition to *Escherichia coli, Escherichia* includes the named species

albertii [51-53], *fergusonii* [54,55], *marmotae* [56,57] and *ruysiae* [58]. And despite

their apparently distinct genus and species names, *Shigella boydii, Shigella*

dysenteriae, Shigella flexneri and *Shigella sonnei,* all common causes of dysentery,

corresponds to phylogenetic lineages within *E. coli* [43] rather than to discrete

taxonomic units. Still other, unusual *Escherichia* strains from lake and ocean water

are associated with long phylogenetic branches, and were designated as 'cryptic

clades' I-VIII by population geneticists [59-65]. The branch leading to Clade I is

simply a long phylogenetic branch within *E. coli* [59]. However, Clade V

encompasses *E. marmotae* [56,57] and *E. ruysiae* consists of the union of clades III

and IV [58]. Initial analyses with the EcoRPlus collection of 9479 genomes [28]

(Table 3) yielded results that were compatible with these interpretations. Almost all *E.*

coli or *Shigella* genomes were within HierCC cluster HC2350_1, and genomes with

other taxonomic or Clade designations belonged to other HC2350 clusters. However,

soon after the definition of the EcoRPlus collection, additional genomes of

Escherichia which were related to Clade II were described from inter-tidal marine and

fresh water sediments near Hong Kong [61,66]. Furthermore, due to its numerical

predominance, *E. coli* overshadows other *Escherichia* species and sub-species within

EcoRPlus. We therefore created *Escherichia* reps in Jan 2021, a novel set of 967

representative genomes which included one genome from each of the 160 most

common HC1100 clusters in HC2350_1 and all 807 genomes from other HC2350

clusters which existed in EnteroBase at that time.

Fig. 2 shows that *E. fergusonii* and *E. albertii* form discrete 95% ANI clusters as do

E. marmotae and other Clade V genomes. As previously reported [58], *E. ruysiae*

consists of the distinct clusters of Clades III and IV. Three distinct 95% ANI clusters

were found within Clades II, VI, and VIII while the Hong Kong genomes represented

multiple, related phylogenetic branches, one of which has previously been designated

Clade VII. All *E. coli* and *Shigella* genomes are in a common 95% ANI group, as is

Clade I. Comparable clustering results were also found in an ML tree of

presence/absence of accessory genes, albeit with different topology of the deepest

branches resulting in *E. fergusonii* and Clade VIII being most closely related to *E. coli*

(https://enterobase.warwick.ac.uk/ms_tree?tree_id=71125). These observations

indicate that the taxonomy and population genetic designations for *Escherichia* are

incomplete, and also partially inconsistent.

HierCC clustering of the same data provided a consistent and uniform nomenclature.

Each discrete phylogenetic cluster received its own HC2850 designation, including

Clades III and IV and four phylogenetic clusters among the Hong Kong isolates

which were designated according to their HC cluster number: e.g. *Escherichia*

HC2350_89353. *E. coli, Shigella* and Clade I all clustered together in HC2350_1, and

the only discordance between topological clustering and HierCC clustering among the

967 genomes in *Escherichia* reps was a single genome which belonged to *E. coli* by

phylogenetic topology but was assigned to HC2350_11465 by HierCC. These results

were so convincing that HierCC groupings were used in late 2021 to curate and

update all species designations for *Escherichia* entries within EnteroBase.

Clostridioides. *C. difficile* clustered distinctly from *C. mangenotii* according to 95%

ANI, and ANI also distinguished five other clusters among genomes that were

designated *C. difficile*. These clusters seem to be novel species according to their

phylogenetic topology in the ML SNP super-tree (Fig. 3). HC2500 identified the same

clusters, and also separated out two additional distinct clusters which resemble novel

sub-species (arrows: HC2500_15334, HC2500_15408). A subset of the genomes in

these clusters (Fig. 3) were assigned to cryptic clades C-I, CII and C-III by a recent

publication [67], which also concluded that they represent novel species. Thus,

HierCC is also suitable for the automated detection of new *Clostridioides* species and

subspecies, and for routinely assigning novel genomes to the appropriate taxa.

Yersinia **and** *Vibrio.* Detailed results with *Yersinia* and *Vibrio* can be found in

Supplemental Text. The general take-home lessons from those analyses resemble

those summarized above: 95% ANI clustering is largely congruent with existing

taxonomic designations, and HierCC identifies both species and sub-species with

better resolution than 95% ANI. However, multiple, unanticipated taxonomic

problems were evident in both genera. Firstly, ENA uses the strain designation as a

species designation when a species is not specified during the submission process.

The resulting public metadata is bloated with inaccurate species names. Secondly,

multiple distinct phylogenetic branches were found without any species identifier.

Thirdly, in multiple cases the species identifier did not correspond with ANI and/or

HierCC, and comparisons with both the SNP and presence/absence ML topologies

confirmed that the species designations in ENA were incorrect. We addressed these

problems within EnteroBase by performing radical manual curation to the metadata to

ensure that the taxonomic species designations correspond to the phylogenies and

population structures. We also encountered multiple, problematical taxonomic

designations which are described in the next paragraphs.

Yersinia. 95% ANI supports the taxonomic convention that *Y. enterocolitica*

represents a single species, and all genomes designated as *Y. enterocolitica* were in a

single 95% ANI cluster. However, in accord with the conclusions by Reuter *et al.*

[68], HierCC clustering assigned the non-pathogenic biotype 1A genomes to

HC1490_73 and HC1490_764, the highly pathogenic biotype 1B to HC1490_2, and

pathogenic biotypes 2-5 to HC1490_10. We therefore renamed these four groups by

adding the HierCC cluster to the species name, e.g. *Y. enterocolitica* HC1490_2.

In contrast to *Y. enterocolitica,* the *Yersinia pseudotuberculosis* Complex corresponds

to a single phylogenetic cluster according to both HC1490 and 95% ANI

(Supplemental Fig. S1). Taxonomists have split these bacteria into *Y.*

pseudotuberculosis, *Y. pestis, Y. similis* and *Y. wautersii* [69,70]. DNA-DNA

hybridization [71] and gene sequences of several housekeeping genes [4] previously

demonstrated that *Y. pestis* is a clade of *Y. pseudotuberculosis,* and the new

observations demonstrate that a distinct species status is not consistent with the

genomic data for *Y. similis* and *Y. wautersii*. We have therefore downgraded all three

taxa within EnteroBase to the category of subspecies of *Y. pseudotuberculosis,* and

extended the designation of *Y. pseudotuberculosis* to *Y. pseudotuberculosis sensu*

stricto. These assignments are not reflected by distinct HC1490 clusters, and HC1490

clustering can only be used to automatically assign new genomes to the *Y.*

pseudotuberculosis Complex.

We also defined eight other novel species/sub-species designations within *Yersinia* by

unique designations based on HierCC clusters: e.g. *Yersinia* HC1490_419. These

clusters were previously unnamed, or had been incorrectly designated with the names

of other species which formed distinct ANI and HC1490 clusters (SupplementalText,

Fig. S1).

Vibrio. *Vibrio* encompassed 152 HC1090 clusters (Supplemental Fig. S2,

SupplementalText), which corresponds to much greater taxonomic diversity than for

the other genera dealt with above. The concordance between HC1090 and 95% ANI

clusters was absolute for most clusters, including a large numbers of genomes from *V.*

cholerae, V. parahaemolyticus, *V. vulnificans,* and *V. anguillarum* (Supplemental

Table S1), and did not support the existence of additional sub-species. However,

eleven HC1090 clusters each encompassed between two and five ANI clusters

(Supplemental Table S2) and three ANI clusters each encompassed 2-3 HC1090

clusters (Supplemental Table S3). In order to support the automated assignment of

genomes to named species, we implemented taxonomic assignments according to

HC1090, and renamed the species of all genomes that were contradictory to this

principle. The resulting dataset contains 109 HC1090 clusters with a unique, classical

species designation as well as 43 other species level clusters designated as *Vibrio*

HC1090_xxxx. Seventeen species names were eliminated because they were

contradictory to the phylogenetic topologies or were incoherently applied

(Supplemental Text, Supplemental Table 4). These taxonomic changes now permit

future automated assignment of novel genomes to species designations, and the

recognition of novel species, and have provided a clean and consistent basic

taxonomy that can be progressively expanded.

Prior work has assigned many *Vibrio* species into so-called higher order clades of

species on the basis of MLSA (multilocus sequence analysis) [72]. These clades are

also apparent in the ML super-tree of core SNPs, and Fig. S2 indicates the three

largest (Cholerae, Harveyii, Splendidus).

Streptococcus. For all five genera summarized above, clustering according to HierCC

was largely concordant with the ML super-trees based on core SNPs or

presence/absence of accessory genes. 95% ANI and taxonomic designations were also

largely concordant with the phylogenetic trees, although to a lesser degree. Similar

concordances with the trees based on core SNPs (Fig. 4) and presence/absence of

accessory genes (Supplemental Fig. S3) were also found for a majority of the named

species within *Streptococcus*. 102 HC363 clusters were concordant with 95% ANI

clusters, and each of those HierCC clusters was specific for a single species after

eliminating *S. milleri, S. periodonticum* and *S. ursoris* (Supplemental Table S5).

These results also demonstrated strong agreement between the two methods with

classical taxonomy. However, genomes from a large number of other species within

Streptococcus were not clustered satisfactorily by either method.

Earlier publications [15,16,73,74], had demonstrated that *Streptococcus mitis* and

oralis each represents multiple related sequence clusters rather than two single

species, and similar results were obtained with 95% ANI clustering [17]. These

observations were confirmed by the current data. Of the 102 concordant HC363 and

95% ANI clusters, 23 clusters from *S. cristatus, S. infantis, S. mitis, S. oralis, S.*

sanguinis or *S. suis* each corresponded to only a sub-set of the topological diversity

within their species. Furthermore, single HC363 clusters from 18 species or groups of

species were subdivided extensively by 95% ANI for a total of 155 discrepancies

between the two methods (Supplemental Table S6). These discrepant clusters were

within the species *S. suis, S. criceti, S. sanguinis*, *S. suis, S. oralis, S. hyointestinalis,*

S. parasanguinis, S. australis, S. infantis, and *S. cristatus*. 95% ANI was not able to

differentiate *S. pneumoniae* from *S. pseudopneumoniae*, *S. pyogenes* from *S.*

dysgalactiae, S. salivarius from *S. vestibularis*, or between *S. equinus, S. infantarius,*

and *S. lutentiensis*. Contrariwise, three other 95% ANI clusters each encompassed two

HC363 clusters (Supplemental Table S7). HC363 could not distinguish *S. pneumoniae*

or *S. pseudopneumoniae* from fifty-seven 95% ANI clusters within *S. mitis*. In

addition to these problems, the eight additional species should not have been given a

species name because their type strains belonged to one of these chaotic clusters. We

conclude that numerous species have been defined on taxonomic grounds that cannot

be correctly identified by either 95% ANI or HierCC, and that multiple discrepancies

exist between the two approaches within *Streptococcus*.

b. Populations, eBGs and Lineages

HierCC supports the identification of populations at multiple hierarchical levels

ranging from the species/sub-species down to individual transmission chains. We also

expected that intermediate HierCC cluster levels could reliably detect the natural

populations defined by legacy MLST. Here we focus on examples of such populations

from *Salmonella*, *E.coli/Shigella,* and *S. pneumonia,* taxa where they have been

examined in greatest detail.

***Salmonella*.** eBGs (eBurstGroups) defined by legacy MLST in *Salmonella* generally

correspond to HierCC HC900 clusters [32]. In some cases, genetically related groups

of eBGs correspond to HC2000 clusters [45]; we refer to those as Lineages. In

contrast, none of the HierCC levels corresponded consistently with other, prior intra-

species subdivisions of *S. enterica* subsp. *enterica* into lineages [75], Clades [76-79]

or Branches [80]. This failure may reflect a high frequency of extensive homologous

recombination at these deeper branches, which can obscure topological relationships

[46,80], whereas HC900 (and HC2000 clusters) diverged more recently, and have not

yet undergone extensive recombination [46].

Traditional subgrouping nomenclature of *Salmonella* below the sub-species level

predates DNA sequencing, and is predominantly based on serovar designations.

Serovar designations consist of common names that have been assigned to a total of

more than 2500 unique antigenic formulas based on epitopes within

lipopolysaccharide (O antigen) and two alternately expressed flagellar subunits (H1,

H2). These antigenic formulas are written in the form O epitopes:H1 epitopes:H2

epitopes in the revised Kauffmann-White scheme [81,82], and O epitopes are

summarized as O serogroups with distinct numeric designations, e.g. O:4 for the O

group of serovar Typhimurium. Detailed metabolic maps for LPS synthesis have been

elucidated for 47 O serogroups [83,84]. Serogroups from natural isolates can be

determined serologically by agglutination reactions with specific antisera or *in silico*

from genomic sequences with the programs SeqSero [85] or SISTR [86]. In practice.

each of these methods has an error rate of a few percent [27,87].

Several hundred legacy eBGs within subsp. *enterica* were relatively uniform for

serovar [27], with multiple exceptions, and many serovars were associated with

multiple, apparently unrelated eBGs, likely due to the extensive exchange of genes

encoding LPS and flagellar epitopes between eBGs early in their evolutionary history.

We proposed that legacy MLST was a better metric for identifying the natural

populations than serotyping, and could completely replace this traditional method

[27]. Our new data indicate that HC900 clusters are even more reliable than legacy

eBGs for recognizing natural populations within *Salmonella*.

Several phylogenetic comparisons indicated that HC900 clusters based on cgMLST

are concordant with eBGs in *Salmonella* [32,45]. Here we test this concordance

quantitatively over a total of 319,490 genomes (Table 4). The concordance between

eBG and HC900 clusters was 0.985 according to the Adjusted Mutual Information

Score [AMI] (Table 4), a metric that is suitable for samples with a heterogeneous size

distribution [88], and almost as high with the Adjusted Rand Index (ARI: 0.982),

which is less suitable for heterogeneous data. Comparisons of eBG or HC900 with

HC2000 clustering yielded slightly lower AMI scores (Table 4). We also tested

whether HC900 clusters are uniform for serovar. In late 2019, we corrected false

serological data in the metadata Serovar field in EnteroBase by manual curation of

790 HC900 clusters that contained at least 5 entries. 97% (770/790) of those HC900

clusters were uniform (≥95%) for serovar (Supplemental Table S8), as were most

HC2000 clusters (437/473 clusters; 92%). The data indicated that the predominant O

group was uniform over the multiple HC900 clusters within 95% (382/403) of

HC2000 clusters within *S. enterica* subsp. *enterica* and 79% (58/70) of HC2000

clusters in other *Salmonella* species and sub-species (Fig. 5; Supplemental Table S9).

HC2000 Lineages with uniform O group included the Para C Lineage

(HC2000_1272; serovars Paratyphi C, Typhisuis, Choleraesuis and Lomita [46])

which emerged about 3500 years ago (Supplemental Text) [46,80]. The Para C

Lineage is uniformly serogroup O:7, which is not particularly surprising because all

its serovars have identical antigenic formulas, and are only distinguished by biotype.

The Enteritidis Lineage (HC2000_12) consists of serovars Enteritidis (1,9,12:g,m:-),

Gallinarum and its variant Pullorum (1,9,12:-:-), and Dublin (1,9,12:g,p:-:-) [45] (Fig.

5), and is uniform for O:9. The Typhimurium Lineage (HC2000_2) includes serovars

Typhimurium, Heidelberg, Reading, Saintpaul, Haifa, Stanleyville and others [45].

This Lineage is also uniform for O group because all these serovars are O:4. The

Mbandaka Lineage (HC2000_4) includes serovars Mbandaka (6,7,14: z10:e,n,z15)

and Lubbock (6,7:g,m,s:e,n,z15), and is uniformly O:7. However, three Lineages were

exceptional, and did contain more than one O serogroup. HC2000_299 is a mixture of

O:4 (HC900_299; serovar Agama [28]) and O:7 (HC900_127074; serovar Nigeria).

HC900 clusters within HC2000_54 are O:4 (HC900_54: Bredeney, Schwarzengrund;

HC24937: Kimuenza) or O:9,46 (HC900_57: Give). Similarly, HC900 clusters within

HC2000_106 are O:3,10; O:9; O:9,46; O:4: or O:8. These observations confirm and

extend the prior conclusions about concordance between natural populations and

serogroup within *Salmonella.*

Escherichia coli. Patterns of multilocus isoenzyme electrophoresis were used in the

1980s to provide an overview of the genetic diversity of *E. coli* (see overview by

Chaudhuri and Henderson [89]). Those analyses yielded a representative collection of

72 isolates [90], the EcoR collection, whose deep phylogenetic branches were

designated haplogroups A, B1, B2, C, D and E [91]. Several haplogroups have since

been added [92,93]. The presence or absence of several accessory genes can be used

for the assignment of genomes to haplogroups with the Clermont scheme [94,95], and

the haplogroup can also be predicted *in silico* from genomic assemblies with

ClermontTyping [96] or EZClermont [97], both of which are implemented within

EnteroBase. However, the Clermont scheme ignores *Shigella*, which consists of *E.*

coli clades despite its differing genus designation [26,42,43], and makes multiple

discrepant assignments according to phylogenetic trees [28]. The Clermont scheme

also does not properly handle the entire diversity of species and environmental

clades/subspecies within the genus *Escherichia* [28,58,59,61].

EnteroBase HierCC automatically assigns genomes within the *Escherichia/Shigella*

database to the cgMLST equivalents of ST Complexes (HC1100 clusters) and

Lineages (HC2000 clusters) (Table 1). It also perpetuates the ST Complexes that were

initially defined for legacy MLST by Wirth *et al.* [26]. However, the numbers and

composition of legacy ST Complexes have not been updated since 2009 because

additional sequencing data defined intermediate, recombinant genotypes that were

equidistant to multiple ST Complexes, and threatened to merge existing ST

Complexes. In contrast, HC1100 clusters do not merge: Intermediate genotypes are

rare because cgMLST involves 2512 loci while legacy MLST was based on only

seven. Secondly, new genotypes which are similar to and equidistant to multiple

clusters do not trigger merging because HierCC arbitrarily assigns them to the oldest

of the existing alternatives [32]. Finally, unlike ST Complex designations in legacy

MLST, which have not been actively updated in *Escherichia* since 2006, HierCC

clustering is an ongoing, automated process for all new genotypes.

Frequent recombination in *E. coli* results in poor bootstrap support for the deep

branches in phylogenetic trees of concatenated genes [26], and we were unable to

identify a unique HierCC level which was largely concordant with haplogroups

according to the Clermont scheme. However, similar to *Salmonella,* HC1100 clusters

identify clear population groups, which are highly concordant with legacy ST

Complexes (AMI = 0.94) (Table 4). Unlike *Salmonella*, HC2000 clusters did not in

general mark recognizable additional population structure beyond that which was

provided by ST Complexes, and HC2000 clusters are only moderately concordant

with either legacy ST Complexes (AMI = 0.69) or their cgMLST equivalent, HC1100

clusters (AMI = 0.69) (Table 4). Multiple *Shigella* species are exceptions to this rule

and their legacy ST Complexes [26] equate to cgMLST HC2000 clusters rather than

HC1100 clusters according to phylogenetic trees. HC2000_305 replaces ST152 Cplx,

and encompasses all *S. sonnei* (Fig. 6). HC2000_192 replaces ST245 Cplx, and

contains many *S. flexneri*. However, *S. flexneri* O group F6 is in an HC1100 cluster

together with *S. boydii* B2 and B4 within HC2000_1465 (ST243 Cplx).

HC2000_1465 includes a second HC1100 cluster with *S. boydii* O groups B1 and

B18, and a third with *S. dysenteriae* D3, D9 and D13 (Fig. 6). HC2000_4118 replaces

the combination of ST250 Complex and ST149 Complex, which have merged, and

contains both *S. dysenteriae* as well as *S. boydii.*

Unlike *Salmonella*, O serogroups are remarkably variable within HC1100 clusters of

E. coli (Fig. 6). HC1100_63 (ST11 Complex) combines classical EHEC strains of

serovar O157:H7 as well as O55:H7 [98], and other, atypical ancestral EPEC isolates

of serovar O55:H7 [99]. Multiple other HC1100 clusters that contain EHEC strains

also encompass multiple O groups, including HC1100_2 (ST29 Cplx; O26, O103,

O111) [100] and HC1100_3 (ST20 Cplx; O45, O103, O111, O128). Similarly,

multiple HC1100 clusters of *E. coli* that cause extra-intestinal diseases also contain a

variety of O groups, including HC1100_7 (ST131 Cplx; O16, O25) [23,101,102] and

HC1100_44 (K1-encapsulated bacteria of the ST95 Cplx; O1, O2, O18, O25, O45)

[26,103,104]. Indeed, the primary impression from Fig. 6 is that O group variation

within an HC1100 cluster is nearly universal throughout *E. coli*, confirming prior

conclusions about the high frequency of homologous recombination in this species

[26,92].

Other genera. Correlations between intra-specific HierCC clusters and population

groupings at multiple levels, including ST Complexes and ribotypes have been

described in *Clostridioides difficile* [105]. It remains to be tested whether HierCC can

identify populations within *Yersinia* because bacterial populations in *Y.*

pseudotuberculosis are largely obscured by recombination [106]. Similarly, we are

not aware of extensive efforts to identify bacterial populations which could be

compared with HierCC within *Vibrio*. However, recent work has described the

assignment of large numbers of genomes of *Streptococcus pneumoniae* to legacy

Clonal Complexes as well as GSPC (Global Pneumococcal Sequence Clusters) [107],

initially with PopPunk [108] and more recently with Mandrake [109]. Most of those

genomes have been assembled within EnteroBase, and assigned to HierCC clusters.

We therefore compared the HierCC and GSPC assignments for 18,147 genomes, and

also compared them to Clonal Complexes (CCs) based on legacy MLST (13,396

genomes). HC160 clusters yielded the highest AMI and ARI scores with GSPC

clusters while CC was most concordant with HC100 clusters (Table 4). These

conclusions were also supported by visual comparisons of the colour-coding of nodes

within a Neighbor-Joining tree based on cgMLST distances according to the various

clustering criteria (Supplemental Fig. S4). Thus, HC100 clusters seem to be

concordant with Clonal Complexes within *S. pneumonia* and Mandrake/PopPunk

clustering equates roughly with HC160 clusters.

Discussion

In 2013, we anticipated that EnteroBase might eventually contain as many as 10,000

genomes of *Salmonella* and *Escherichia*. By the end of 2021 it hosted >600,000

assembled genomes of *Salmonella, Escherichia/Shigella, Streptococcus*, *Vibrio*,

Clostridioides and *Yersinia*, and has become one of the primary global sources of

their genomic data. An analyses of their genomic properties is facilitated by software

tools (Table 2) that support indexing of genetic diversity, searching for genomes with

specific metadata or genetic relations and investigating stable population structures at

multiple levels. Here we focused on the automated identification and designation of

species/subspecies and populations on the basis of HierCC of core genome MLST

genotypes. The data demonstrate that HierCC can completely replace classical

taxonomic methods for genomes of *Salmonella, Escherichia/Shigella, Vibrio*,

Clostridioides and *Yersinia*, with only few minor exceptions. HierCC is also a

complete replacement for legacy MLST regarding the assignments to populations

within *Salmonella*, *Escherichia/Shigella* and *Streptococcus pneumoniae.*

Taxonomic assignments. The paired ML super-trees based on core SNPs and

presence/absence of accessory genes from each genus were generally concordant in

their clustering patterns, indicating that those genera contain well defined and

reproducible species/subspecies independent of the genetic criteria. Those

phylogenetic clustering were then used to resolve the most approrpriate designations

for genomes which were assigned contradictory designations by their taxonomic

metadata, and clusters based on 95% ANI or HierCC. The frequency of agreement

with current taxonomy was comparable for HierCC and 95% ANI at the species level

within *Salmonella, Escherichia/Shigella, Vibrio*, *Clostridioides* and *Yersinia* but 95%

ANI was generally unable to resolve sub-species whereas HierCC did not distinguish

between sub-species and species clusters. Classical taxonomical designations

exhibited multiple, glaring problems in each genus and we replaced those

problematical designations in EnteroBase with labels based on the automated,

HierCC-based species/subspecies taxonomy. We also defined novel species/sub-

species, and labelled them with HierCC-based designations.

Why were there so many obvious discrepancies between the different approaches?

One obvious reason is that we have discarded the medical tradition of retaining

discrete species names for pathogens that cause particular diseases, and recommend

designating *Y. pestis* as a sub-species of *Y. pseudotuberculosis.* Other discrepancies

may reflect technical errors in manipulating data or uploading information to public

databases, or simple strain mix-ups [41,45]. Possibly the discrepancies were

particularly obvious because the scale of our analyses were performed at an

unprecedented scale over multiple genera. Many insights can be attributed to the

facile ability to investigate large databases within a graphic framework that is

provided by EnteroBase, and to our optimization of HierCC levels for each genus.

However, we failed in a similar attempt to optimise ANI levels for recognizing both

species and sub-species.

Our taxonomic changes in *Yersinia* are likely to be highly controversial. According to

both ML trees and HierCC, the current designation "*Y. enterocolitica*" encompasses

four, previously unnamed species/sub-species. These taxa are not distinguished by

95% ANI and had not previously been assigned distinct names. We maintain

continuity with the traditional designation of *Y. enterocolitica*, and simply add

HierCC affiliations to the species name, e.g. *Y. enterocolitica* HC1490_2.

Alternatively, these clusters of strains could be considered to represent sub-species

and their name modified slightly, e.g. *Y. enterocolitica* subspecies HC1490_2, similar

to our downgrading of the named species *Y. pestis, Y. similis* [70] and *Y. wautersii*

[69] on phylogenetic grounds to subspecies of *Y. pseudotuberculosis* (Fig. S1).

Finally, we defined eight new species/sub-species within *Yersinia* that were not

previously differentiated without submitting traditional evidence for simple

phenotypic differences, and named them after their HC cluster, e.g. *Yersinia*

HC1490_4399. We also followed comparable strategies for the other genera analyzed

here.

In *Salmonella,* HierCC and 95% ANI identified a new species, *Salmonella*

HC2850_215890, and a new subspecies, *S. enterica* subsp. HC2850_222931 (Fig. 1).

Five new species were identified within *C. difficile* by HierCC as well as ANI*,* and

HierCC identified two additional new sub-species. In *Escherichia* we provide HierCC

designations for multiple clusters of bacteria from environmental sources near Hong

Kong that were previously unnamed.

The ML trees presented here (Fig. S2) support prior definitions of high order clades

containing multiple species in *Vibrio* on the basis of MLSA [72]. We also observed a

general concordance between ANI and HierCC for 109 named *Vibrio* species*,*

However, 43 HierCC clusters of species rank had not previously been properly

classified with species designations, and 17 other species names were superfluous.

These changes are fully described in Supplemental text, allowing closer scrutiny by

the *Vibrio* taxonomic community for consistency with other properties.

HierCC was so effective in clarifying the taxonomies of these five species that we had

hoped that it could also facilitate the taxonomic classification of *Streptococcus.* 102

HC363 clusters were indeed concordant with 95% ANI and taxonomic designations,

with only minor exceptions. However, extensive discrepancies existed between 95%

ANI and HierCC for numerous other clusters, and both approaches showed multiple

discrepancies with classical taxonomic designations. The existence of multiple 95%

ANI clusters within *S. mitis* and *S. oralis* and other species of *Streptococcus* has

previously been commented on by Kilian and his colleagues [15,16,73,74]. Our

observations extend these taxonomic problems to multiple additional species in which

HC363 does not distinguish between pairs of named species (Fig. 4). Neither 95%

ANI nor cgMLST HierCC seems to provide a suitable general strategy for elucidating

the taxonomy of all of *Streptococcus*, and we failed to find a general solution to this

problem.

HierCC *versus* classical taxonomy. Classical microbiological species taxonomy

involves identifying a type strain whose phenotype can be distinguished from all other

type strains, identifying additional isolates with similar phenotypes, demonstrating

distinct clustering from other known species in phylogenetic trees based on DNA or

amino acid sequences, and publishing a report in one of a very limited number of

acceptable journals. Such species definitions are then considered tentative until an

international committee has approved them. Other forms of identifying species that

include sole reliance on DNA sequence differences are not acceptable [7,8]. The

metagenomics community and scientists working with uncultivated organisms from

the environment have largely liberated themselves from such regulations, and tend to

use operational taxonomic units (OTUs) as taxonomic entities. However, the

taxonomic species structure of many microbes from environmental sources remains

fuzzy, or does not clearly correspond to classical taxonomy (e.g. *Prochlorococcus*

[110] or *Synechococcus* [111]). Furthermore, we are not aware of any other method

that able to assign 1000s of genomes per day to existing taxa and also reliably identify

new taxa automatically as they appear. HierCC performs this task with bravura for the

genera in Table 1, with the notable exception of *Streptococcus*.

We have taken the liberty of dropping all attempts to reconcile HierCC

species/subspecies clusters with classical prescriptions for how to define a species.

Instead we have adopted the practice of using HierCC cluster designations within

EnteroBase for the nomenclatures for species/subspecies groupings which had not yet

been identified by others, and eliminated from EnteroBase multiple species

designations of type strains in *Vibrio* and *Streptococcus* which did not match the ML

tree topologies and HierCC clusters. These actions provide a uniform base for the

future additions of additional species and ensured that future genomes can be

correctly assigned to uniform clusters of related strains. Scientists wishing to use

these schemes to identify the species of their bacterial isolates can upload their

sequenced genomes to EnteroBase. The HierCC assignments will be available within

hours. We also welcome additional curators of these databases who are willing to test

and improve the current taxonomic assignments we have implemented. But we reject

the concept of designating our assignments as tentative until they are confirmed in

several years by an international committee.

HierCC and Populations. The first bacterial taxonomic designations were assigned

over 100 years ago. Bacterial population genetics is much younger, and has many

fewer practitioners. Initial population genetic analyses in the early 1980s subdivided

multiple bacterial species into intra-specific lineages based on multilocus enzyme

electrophoresis [90]. This methodology was replaced in the late 1990s by legacy

MLST based on several housekeeping genes [21], which is currently being replaced

by cgMLST based on all genes in the soft core genome [22,30]. EnteroBase calculates

genotypes for both types of MLST, as well as for rMLST [28]. Legacy STs differ

from cgSTs due to different levels of resolution. However, the boundaries of ST

Complexes/eBGs are highly concordant with HierCC clusters, with AMI indices of

0.985 for eBGs *versus* HC900 *in Salmonella* and 0.94 for ST Complexes *versus*

HC1100 clusters in *Escherichia/Shigella* (Table 4). We recommended previously that

serovars should be replaced by eBGs in *S. enterica* [27] and the data presented here

show that HC900 clusters are an even better replacement. Our data also indicate that

Lineages in *S. enterica* correspond to HC2000 clusters, and that these tend to be

highly uniform for O group. The data presented here also indicate that HC1100

groups are a good replacement for detecting populations within *Escherichia* as are

HC100 clusters for CCs in *S. pneumoniae* [107] (Table 4).

We interpret these consistencies between legacy MLST and cgMLST as reflecting the

existence of natural populations. We previously claimed that legacy ST Complexes in

E. coli were unstable due to frequent recombination [26], and were therefore not

surprised at their tendency to merge as additional isolates were sequenced. However,

legacy MLST is based on only seven genes. The high resolution of cgMLST and the

decision to assign new genotypes that are equidistant from multiple HierCC clusters

to the oldest cluster have largely negated these problems, and *E. coli* HC1100 clusters

correspond to bacterial groupings that have been independently identified by multiple

phenotypic patterns. In the early 1980s, MA began his academic research on bacterial

pathogens with *E. coli* that expressed the K1 polysaccharide capsule [103]. He

observed unusually uniform patterns of electrophoretic mobility of major outer

membrane proteins across multiple isolates, and interpreted those bacteria as

representing "clones". These "clones" correspond to HC1100 clusters, including

HC1100_44 (ST95 Complex) which continues to cause invasive disease in humans

and animals around the globe [104]. The 1980s analyses showed that these K1

bacteria were variably O1, O2, or O18, and that the serotype variants differed both in

their invasiveness and in the hosts which they infected [112]. As indicated in Fig. 6,

HC1100_44 includes O groups O25 and O45 in addition to O1, O2 and O18.

Similarly, HC1100_7 (ST131) represents another major cause of extra-intestinal

disease in humans [23,101,102]. EHEC bacteria are notorious for their association

with hemolytic uremic syndrome (HUS). Many of them belong to distinctive HC1100

clusters, including O157:H7/O55:H7 (HC1100_63, ST11 Cplx [99,113]) and O26,

O103 and O111 EHEC bacteria (HC1100_2, ST29 Cplx [100]) (Fig. 5).

In contrast to these correlations with epidemiological groupings, we were unable to

identify any other HierCC levels that were consistently concordant with other intra-

specific phylogenetic subdivisions based on SNP trees, including haplogroups [91] or

Clermont typing [94,95] in *E. coli* [28] and Clades [76-79], lineages [75] or Branches

[80] in *S. enterica.* Instead, we found that a subdivision we refer to as Lineages are

marked by some HC2000 clusters in *Salmonella* and *Escherichia/Shigella.*

Lineages. In *E. coli*, HC2000 Lineages were particularly appropriate for defining the

clustering level of seven groups of *Shigella* genomes (Fig. 6). Lineage designations

did not add additional insights into other populations because were almost entirely

encompassed by HC1100 clusters or occasionally even lower level clusters, such as

HC400. In *S. enterica* subspecies *enterica*, multiple HC2000 Lineages corresponded

to genetically related combinations of multiple HC900 clusters, in some cases with

distinctive serovar designations. However, all but three of the major Lineages were

predominantly uniform in O group. The three exceptional HC2000 Lineages

(HC2000_54, HC2000_106, HC2000_299) might be worth exploring in greater detail

to reconstruct the recombination that has resulted in their differing LPS epitopes.

HC160 clusters, the Lineage equivalent in *S. pneumonia,* were strongly concordant

with PopPunk GSPC clustering. Otherwise, little is yet known about the general

properties of Lineages in other genera, except that their properties are likely to vary

with species or genus. For example, *S. enterica* HC2000 Lineages were uniform for O

group whereas O serogroups were already heterogeneous within HC1100 clusters/ST

Complexes within *E. coli* (Fig. 6).

Future prospects. EnteroBase was conceived to satisfy a need that MA and ZZ

perceived in 2014 [1]. We contend that it is now fit for purpose for investigations of

multiple bacterial genera by scientists ranging from beginners through to experts in

the areas of microbial epidemiology and population genetics. Its original creators have

now all left this project, but EnteroBase is being maintained as a service for the global

community by the University of Warwick. Maintenance of its technical functions and

databases are thereby assured for the near future. Further functional developments

will, however, depend on increased participation and perception of ownership by its

users. We perceive a general trend to focus on insular solutions that can satisfy the

demands of individual bioinformaticians and regional diagnostic laboratories. Such

approaches can yield relatively rapid progress in solving short-term needs. However,

a global overview of genomic diversity needs central databases to ensure definitive

terminology. Pooling efforts on a central endeavor at the scale represented by

EnteroBase would ensure that it continues to function over decades, is representative

over all continents, and serves the global community even better. We therefore

welcome additional curators and scientific experts as well as bioinformatics

collaborations to help improve EnteroBase even more.

Methods (see Supplemental Text)

Funding

MA was supported by grant 202792/Z/16/Z from the Wellcome Trust (https://wellcome.org/).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Code.

cgMLSA is available for downloading at https://github.com/zheminzhou/cgMLSA.

All trees in figures and genomic data are freely available within EnteroBase for interactive examination. Other code is described in Supplemental Text.

Data

Datasets used for the analyses presented here have been deposited for permanent public access at http://wrap.warwick.ac.uk/162247/

Acknowledgements

We are very grateful to François-Xavier Weill for permission to include Table S8 for public access in Supplemental Material. We are very grateful to François-Xavier Weill and Tandy Warnow for their comments on an earlier draft of this manuscript.

References

1. Achtman, M. and Zhou, Z. 2014 Distinct genealogies for plasmids and

chromosome. *PLoS Genet* **10**, e1004874. (doi:10.1371/journal.pgen.1004874

[doi];PGENETICS-D-14-02780 [pii])

2. Kauffmann, F. 1961 *Die Bakteriologie der Salmonella-Species*. Copenhagen,

Denmark: Munksgaard.

3. Morelli, G. *et al.*. 2010 *Yersinia pestis* genome sequencing identifies patterns of

global phylogenetic diversity. *Nature Genet* **42**, 1140-1143.

4. Achtman, M., Zurth, K., Morelli, G., Torrea, G., Guiyoule, A., and Carniel, E.

1999 *Yersinia pestis*, the cause of plague, is a recently emerged clone of

Yersinia pseudotuberculosis. *Proc Natl Acad Sci USA* **96**, 14043-14048.

5. Pulford, C. V. *et al.*. 2021 Stepwise evolution of *Salmonella* Typhimurium

ST313 causing bloodstream infection in Africa. *Nat Microbiol* **6**, 327-338.

(doi:10.1038/s41564-020-00836-1 [doi];10.1038/s41564-020-00836-1 [pii])

6. Wayne, L. G. *et al.*. 1987 Report of the ad hoc committee on reconciliation of

approaches to bacterial systematics. *In J Syst Bacteriol* **37**, 463-464.

7. Murray, A. E. *et al.*. 2020 Roadmap for naming uncultivated Archaea and

Bacteria. *Nat Microbiol* **5**, 987-994. (doi:10.1038/s41564-020-0733-x

[doi];10.1038/s41564-020-0733-x [pii])

8. Sutcliffe, I. C., Dijkshoorn, L., Whitman, W. B., and Executive Board. 2020

Minutes of the International Committee on Systematics of Prokaryotes online

discussion on the proposed use of gene sequences as type for naming of

prokaryotes, and outcome of vote. *Int J Syst Evol Microbiol* **70**, 4416-4417.

(doi:10.1099/ijsem.0.004303 [doi])

766 9. Konstantinidis, K. T. and Tiedje, J. M. 2005 Genomic insights that advance the
767   species definition for prokaryotes. *Proc Natl Acad Sci USA* **102**, 2567-2572.

768 10. Jain, C., Rodriguez, R., Phillippy, A. M., Konstantinidis, K. T., and Aluru, S.
769   2018 High throughput ANI analysis of 90K prokaryotic genomes reveals
770   clear species boundaries. *Nat Commun* **9**, 5114. (doi:10.1038/s41467-018-
771   07641-9 [doi];10.1038/s41467-018-07641-9 [pii])

772 11. Auch, A. F., von, J. M., Klenk, H. P., and Goker, M. 2010 Digital DNA-DNA
773   hybridization for microbial species delineation by means of genome-to-
774   genome sequence comparison. *Stand Genomic Sci* **2**, 117-134.
775   (doi:10.4056/sigs.531120 [doi])

776 12. Garrido-Sanz, D., Meier-Kolthoff, J. P., Goker, M., Martin, M., Rivilla, R., and
777   Redondo-Nieto, M. 2016 Genomic and Genetic Diversity within the
778   *Pseudomonas fluorescens* Complex. *PLoS One* **11**, e0150183.
779   (doi:10.1371/journal.pone.0150183 [doi];PONE-D-15-47408 [pii])

780 13. Richter, M. and Roselló-Mora, R. 2009 Shifting the genomic gold standard for
781   the prokaryotic species definition. *Proc Natl Acad Sci U S A* **106**, 19126-
782   19131. (doi:0906412106 [pii];10.1073/pnas.0906412106 [doi])

783 14. Maderankova, D., Jugas, R., Sedlar, K., Vitek, M., and Skutkova, H. 2019 Rapid
784   bacterial species delineation based on parameters derived from genome
785   numerical representations. *Comput Struct Biotechnol J* **17**, 118-126.
786   (doi:10.1016/j.csbj.2018.12.006 [doi];S2001-0370(18)30122-3 [pii])

787 15. Jensen, A., Scholz, C. F., and Kilian, M. 2016 Re-evaluation of the taxonomy of
788   the Mitis group of the genus *Streptococcus* based on whole genome
789   phylogenetic analyses, and proposed reclassification of *Streptococcus*
790   *dentisani* as *Streptococcus oralis* subsp. *dentisani* comb. nov., *Streptococcus*
791   *tigurinus* as *Streptococcus oralis* subsp. *tigurinus* comb. nov., and
792   *Streptococcus oligofermentans* as a later synonym of *Streptococcus cristatus*.
793   *Int J Syst Evol Microbiol* **66**, 4803-4820. (doi:10.1099/ijsem.0.001433 [doi])

16. Kilian, M., Poulsen, K., Blomqvist, T., Havarstein, L. S., Bek-Thomsen, M.,
Tettelin, H., and Sorensen, U. B. 2008 Evolution of *Streptococcus*
pneumoniae and its close commensal relatives. *PLoS ONE* **3**, e2683.

17. Zhou, Z., Charlesworth, J., and Achtman, M. 2020 Accurate reconstruction of
bacterial pan- and core genomes with PEPPAN. *Genome Res* **30**, 1667-1679.
(doi:DOI 10.1011/gr.260828.120)

18. Gomila, M., Pena, A., Mulet, M., Lalucat, J., and Garcia-Valdes, E. 2015
Phylogenomics and systematics in *Pseudomonas*. *Front Microbiol* **6**, 214.
(doi:10.3389/fmicb.2015.00214 [doi])

19. Beaz-Hidalgo, R., Hossain, M. J., Liles, M. R., and Figueras, M. J. 2015
Strategies to avoid wrongly labelled genomes using as example the detected
wrong taxonomic affiliation for *Aeromonas* genomes in the GenBank
database. *PLoS ONE* **10**, e0115813. (doi:10.1371/journal.pone.0115813
[doi];PONE-D-14-36626 [pii])

20. Achtman, M. and Wagner, M. 2008 Microbial diversity and the genetic nature
of microbial species. *Nat Rev Microbiol* **6**, 431-440.

21. Maiden, M. C. J. *et al.*. 1998 Multilocus sequence typing: A portable approach
to the identification of clones within populations of pathogenic
microorganisms. *Proc Natl Acad Sci USA* **95**, 3140-3145.

22. Jolley, K. A., Bray, J. E., and Maiden, M. C. 2018 Open-access bacterial
population genomics: BIGSdb software, the PubMLST.org website and their
applications. *Wellcome Open Res* **3**, 124. (doi:doi:
10.12688/wellcomeopenres.14826.1)

23. Stoesser, N. *et al.*. 2016 Evolutionary history of the global emergence of the
Escherichia coli epidemic clone ST131. *MBio* **7**, e02162. (doi:mBio.02162-
15 [pii];10.1128/mBio.02162-15 [doi])

24.   Kingsley, R. A. *et al.*. 2009 Epidemic multiple drug resistant *Salmonella*
Typhimurium causing invasive disease in sub-Saharan Africa have a distinct
genotype. *Genome Res* **19**, 2279-2287.

25.   Feil, E. J., Li, B. C., Aanensen, D. M., Hanage, W. P., and Spratt, B. G. 2004
eBURST: Inferring patterns of evolutionary descent among clusters of related
bacterial genotypes from Multilocus Sequence Typing data. *J Bacteriol* **186**,
1518-1530.

26.   Wirth, T. *et al.*. 2006 Sex and virulence in *Escherichia coli*: an evolutionary
perspective. *Mol Microbiol* **60**, 1136-1151.

27.   Achtman, M. *et al.*. 2012 Multilocus sequence typing as a replacement for
serotyping in *Salmonella enterica*. *PLoS Pathog* **8**, e1002776.

28.   Zhou, Z., Alikhan, N. F., Mohamed, K., Fan, Y., and Achtman, M. 2020 The
EnteroBase user's guide, with case studies on *Salmonella* transmissions,
Yersinia pestis phylogeny, and *Escherichia* core genomic diversity. *Genome*
Res **30**, 138-152. (doi:gr.251678.119 [pii];10.1101/gr.251678.119 [doi])

29.   Mellmann, A. *et al.*. 2011 Prospective genomic characterization of the German
enterohemorrhagic *Escherichia coli* O104:H4 outbreak by rapid next
generation sequencing technology. *PLoS ONE* **6**, e22751.

30.   Maiden, M. C., van Rensburg, M. J., Bray, J. E., Earle, S. G., Ford, S. A., Jolley,
K. A., and McCarthy, N. D. 2013 MLST revisited: the gene-by-gene
approach to bacterial genomics. *Nat Rev Microbiol* **11**, 728-736.

31.   Moura, A. *et al.*. 2016 Whole genome-based population biology and
epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol* **2**,
16185. (doi:nmicrobiol2016185 [pii];10.1038/nmicrobiol.2016.185 [doi])

32.   Zhou, Z., Charlesworth, J., and Achtman, M. 2021 HierCC: A multi-level
clustering scheme for population assignments based on core genome MLST.
Bioinformatics **37**, 3645-3646. (doi:10.1093/bioinformatics/btab234 [doi])

33. Jones, G. *et al.*. 2019 Outbreak of *Salmonella enterica* serotype Poona in infants
linked to persistent *Salmonella* contamination in an infant formula
manufacturing facility, France, August 2018 to February 2019. *Euro Surveill*
24, 13. (doi:doi.org/10.2807/1560-7917.ES.2019.24.13.1900161)

34. Jones, G. *et al.*. 2019 Outbreak of *Shiga* toxin-producing *Escherichia coli*
(STEC) O26 paediatric haemolytic uraemic syndrome (HUS) cases
associated with the consumption of soft raw cow's milk cheeses, France,
March to May 2019. *Euro Surveill* **24**, 1900305. (doi:10.2807/1560-
7917.ES.2019.24.22.1900305)

35. Van den Bossche, A., Ceyssens, P. J., Denayer, S., Hammami, N., van den Beld,
M., Dallman, T. J., and Mattheus, W. 2021 Outbreak of Central American
born *Shigella sonnei* in two youth camps in Belgium in the summer of 2019.
Eur J Clin Microbiol Infect Dis **40**, 1573-1577. (doi:10.1007/s10096-021-
04164-y [doi];10.1007/s10096-021-04164-y [pii])

36. European Centre for Disease Prevention and Control, E. F. S. A. 2020 Multi-
country outbreak of *Salmonella* Typhimurium and *S.* Anatum infections
linked to Brazil nuts - 21 October 2020. *EFSA Supporting Publications* **17**,
1944E. (doi:https://doi.org/10.2903/sp.efsa.2020.EN-1944)

37. Bielaszewska, M., Mellmann, A., Zhang, W., Kock, R., Fruth, A., Bauwens, A.,
Peters, G., and Karch, H. 2011 Characterisation of the *Escherichia coli* strain
associated with an outbreak of haemolytic uraemic syndrome in Germany,
2011: a microbiological study. *Lancet Infect Dis* **11**, 671-676.

38. Hawkey, J. *et al.*. 2021 Global population structure and genotyping framework
for genomic surveillance of the major dysentery pathogen, *Shigella sonnei*.
Nat Commun **12**, 2684. (doi:10.1038/s41467-021-22700-4
[doi];10.1038/s41467-021-22700-4 [pii])

39.   Park, S. E. *et al.*. 2018 The phylogeography and incidence of multi-drug
resistant typhoid fever in sub-Saharan Africa. *Nat Commun* **9**, 5094.
(doi:10.1038/s41467-018-07370-z [doi];10.1038/s41467-018-07370-z [pii])

40.   Weill, F. X. *et al.*. 2017 Genomic history of the seventh pandemic of cholera in
Africa. *Science* **358**, 785-789. (doi:358/6364/785
[pii];10.1126/science.aad5901 [doi])

41.   Achtman, M., van den Broeck, F., Cooper, K. K., Lemey, P., Parker, C. T.,
Zhou, Z., and the ATCC14028s Study Group. 2021 Genomic population
structure associated with repeated escape of *Salmonella enterica*
ATCC14028s from the laboratory into nature. *PLoS Genet* **17**, e1009820.
(doi:DOI: 10.1371/journal.pgen.1009820)

42.   Yassine, I. *et al.*. 2022 Population structure analysis and laboratory monitoring
of *Shigella* by core-genome multilocus sequence typing. *Nat Commun* **13**,
551. (doi:10.1038/s41467-022-28121-1 [doi];10.1038/s41467-022-28121-1
[pii])

43.   Pupo, G. M., Lan, R., and Reeves, P. R. 2000 Multiple independent origins of
Shigella clones of *Escherichia coli* and convergent evolution of many of their
characteristics. *Proc Natl Acad Sci USA* **97**, 10567-10572.

44.   Alikhan, N.-F., Zhou, Z., Sergeant, M. J., and Achtman, M. 2018 A genomic
overview of the population structure of *Salmonella*. *PLoS Genet* **14**,
e1007261. (doi:10.1371/journal.pgen.1007261 [doi];PGENETICS-D-18-
00122 [pii])

45.   Achtman, M. *et al.*. 2020 Genomic diversity of *Salmonella enterica* -The
UoWUCC 10K genomes project. *Wellcome Open Res* **5**, 223.
(doi:10.12688/wellcomeopenres.16291.2 [doi])

46.   Zhou, Z. *et al.*. 2018 Pan-genome analysis of ancient and modern *Salmonella*
enterica demonstrates genomic stability of the invasive Para C Lineage for

millennia. *Curr Biol* **28**, 2420-2428.

(doi:https://doi.org/10.1016/j.cub.2018.05.058)

47.   Zaharias, P. and Warnow, T. 2022 Recent progress on methods for estimating

and updating large phylogenies. *Phil Trans R Soc B*. (doi:doi:

10.20944/preprints202110.0258.v1)

48.   Vachaspati, P. and Warnow, T. 2015 ASTRID: Accurate Species TRees from

Internode Distances. *BMC Genomics* **16 Suppl 10**, S3. (doi:1471-2164-16-

S10-S3 [pii];10.1186/1471-2164-16-S10-S3 [doi])

49.   Sayyari, E. and Mirarab, S. 2016 Fast coalescent-based computation of local

branch support from quartet frequencies. *Mol Biol Evol* **33**, 1654-1668.

(doi:msw079 [pii];10.1093/molbev/msw079 [doi])

50.   Pearce, M. E., Langridge, G. C., Lauer, A. C., Grant, K., Maiden, M. C. J., and

Chattaway, M. A. 2021 An evaluation of the species and subspecies of the

genus *Salmonella* with whole genome sequence data: Proposal of type strains

and epithets for novel *S. enterica* subspecies VII, VIII, IX, X and XI.

Genomics **113**, 3152-3162. (doi:S0888-7543(21)00281-0

[pii];10.1016/j.ygeno.2021.07.003 [doi])

51.   Huys, G., Cnockaert, M., Janda, J. M., and Swings, J. 2003 *Escherichia albertii*

sp. nov., a diarrhoeagenic species isolated from stool specimens of

Bangladeshi children. *Int J Syst Evol Microbiol* **53**, 807-810.

(doi:10.1099/ijs.0.02475-0 [doi])

52.   Hyma, K. E., Lacher, D. W., Nelson, A. M., Bumbaugh, A. C., Janda, J. M.,

Strockbine, N. A., Young, V. B., and Whittam, T. S. 2005 Evolutionary

genetics of a new pathogenic *Escherichia* species: *Escherichia albertii* and

related *Shigella boydii* strains. *J Bacteriol* **187**, 619-628.

53.   Ooka, T. *et al.*. 2015 Defining the genome features of *Escherichia albertii*, an
926   emerging enteropathogen closely related to *Escherichia coli*. *Genome Biol*
927   *Evol* **7**, 3170-3179. (doi:evv211 [pii];10.1093/gbe/evv211 [doi])

928   54.   Farmer, J. J. 3., Fanning, G. R., Davis, B. R., O'Hara, C. M., Riddle, C.,
929   Hickman-Brenner, F. W., Asbury, M. A., Lowery, V. A. 3., and Brenner, D.
930   J. 1985 *Escherichia fergusonii* and *Enterobacter taylorae*, two new species of
931   Enterobacteriaceae isolated from clinical specimens. *J Clin Microbiol* **21**, 77-
932   81. (doi:10.1128/jcm.21.1.77-81.1985 [doi])

933   55.   Gaastra, W., Kusters, J. G., van, D. E., and Lipman, L. J. 2014 *Escherichia*
934   *fergusonii*. *Vet Microbiol* **172**, 7-12. (doi:S0378-1135(14)00216-8
935   [pii];10.1016/j.vetmic.2014.04.016 [doi])

936   56.   Liu, S., Jin, D., Lan, R., Wang, Y., Meng, Q., Dai, H., Lu, S., Hu, S., and Xu, J.
937   2015 *Escherichia marmotae sp. nov.*, isolated from faeces of *Marmota*
938   *himalayana*. *Int J Syst Evol Microbiol* **65**, 2130-2134.
939   (doi:10.1099/ijs.0.000228 [doi];ijs.0.000228 [pii])

940   57.   Liu, S. *et al.*. 2019 Genomic and molecular characterisation of *Escherichia*
941   *marmotae* from wild rodents in Qinghai-Tibet plateau as a potential
942   pathogen. *Sci Rep* **9**, 10619. (doi:10.1038/s41598-019-46831-3
943   [doi];10.1038/s41598-019-46831-3 [pii])

944   58.   van der Putten, B. C. L., Matamoros, S., COMBAT consortium, and Schultsz,
945   C. 2021 *Escherichia ruysiae* sp. *nov.,* a novel Gram-stain-negative bacterium,
946   isolated from a faecal sample of an international traveller. *Int J Syst Evol*
947   *Microbiol* **71**, 004609. (doi:DOI 10.1099/ijsem.0.004609)

948   59.   Walk, S. T. 2015 The "Cryptic" Escherichia. *EcoSal Plus* **6**.
949   (doi:10.1128/ecosalplus.ESP-0002-2015 [doi])

950   60.   Vignaroli, C., Di, S. L., Magi, G., Luna, G. M., Di, C. A., Pasquaroli, S.,
951   Facinelli, B., and Biavasco, F. 2015 Adhesion of marine cryptic *Escherichia*

952    isolates to human intestinal epithelial cells. *ISME J* **9**, 508-515.

953    (doi:ismej2014164 [pii];10.1038/ismej.2014.164 [doi])

954 61. Shen, Z., Koh, X. P., Yu, Y., Woo, C. F., Tong, Y., and Lau, S. C. K. 2019

955    Draft genome sequences of 16 strains of *Escherichia* cryptic clade II isolated

956    from intertidal sediment in Hong Kong. *Microbiol Resour Announc* **8**.

957    (doi:8/29/e00416-19 [pii];10.1128/MRA.00416-19 [doi])

958 62. Zhou, Z., Alikhan, N.-F., Mohamed, K., Fan, Y., Agama Study Group, and

959    Achtman, M. 2020 The EnteroBase user's guide, with case studies on

960    *Salmonella* transmissions, *Yersinia pestis* phylogeny, and *Escherichia* core

961    genomic diversity. *Genome Res* **30**, 138-152. (doi:gr.251678.119

962    [pii];10.1101/gr.251678.119 [doi])

963 63. Luo, C., Walk, S. T., Gordon, D. M., Feldgarden, M., Tiedje, J. M., and

964    Konstantinidis, K. T. 2011 Genome sequencing of environmental *Escherichia*

965    *coli* expands understanding of the ecology and speciation of the model

966    bacterial species. *Proc Natl Acad Sci USA* **108**, 7200-7205.

967 64. Walk, S. T., Alm, E. W., Gordon, D. M., Ram, J. L., Toranzos, G. A., Tiedje, J.

968    M., and Whittam, T. S. 2009 Cryptic lineages of the genus *Escherichia*. *Appl*

969    *Environ Microbiol* **75**, 6534-6544.

970 65. Walk, S. T., Alm, E. W., Calhoun, L. M., Mladonicky, J. M., and Whittam, T. S.

971    2007 Genetic diversity and population structure of *Escherichia coli* isolated

972    from freshwater beaches. *Environ Microbiol* **9**, 2274-2288.

973 66. Shen, Z. Y., Koh, X. P., Yu, Y. P., and Lau, S. C. K. 2020 Genetic variation and

974    preliminary indications of divergent niche adaptation in cryptic clade II of

975    *Escherichia*. *Microorganisms* **8**. (doi:microorganisms8111713

976    [pii];10.3390/microorganisms8111713 [doi])

977 67. Knight, D. R., Imwattana, K., Kullin, B., Guerrero-Araya, E., Paredes-Sabja, D.,

978    Didelot, X., Dingle, K. E., Eyre, D. W., Rodríguez, C., and Riley, T. V. 2021

979   Major genetic discontinuity and novel toxigenic species in *Clostridioides*

980    *difficile* taxonomy. *eLife* **10**, e64325.

981 68. Reuter, S. *et al.*. 2014 Parallel independent evolution of pathogenicity within the

982    genus *Yersinia*. *Proc Natl Acad Sci U S A* **111**, 6768-6773. (doi:1317161111

983    [pii];10.1073/pnas.1317161111 [doi])

984 69. Savin, C. *et al.*. 2014 The *Yersinia pseudotuberculosis* complex:

985    characterization and delineation of a new species, *Yersinia wautersii*. *Int J*

986    *Med Microbiol* **304**, 452-463. (doi:S1438-4221(14)00012-5

987    [pii];10.1016/j.ijmm.2014.02.002 [doi])

988 70. Sprague, L. D., Scholz, H. C., Amann, S., Busse, H. J., and Neubauer, H. 2008

989    *Yersinia similis* sp. nov. *Int J Syst Evol Microbiol* **58**, 952-958.

990 71. Bercovier, H., Mollaret, H. H., Alonso, J. M., Brault, J., Fanning, G. R.,

991    Steigerwalt, A. G., and Brenner, D. J. 1980 Intra- and interspecies relatedness

992    of *Yersinia pestis* by DNA hybridization and its relationship to *Yersinia*

993    *pseudotuberculosis*. *Curr Microbiol* **4**, 225-229.

994 72. Gomez-Gil, B., Thompson, C. C., Matsumura, Y., Sawabe, T., Iida, T., Christen,

995    R., Thompson, F., and Sawabe, T. 2014 The family *Vibrionaceae*. In *The*

996    *prokaryotes - Gammaproteobacteria* (eds. E. Rosenberg, E. F. DeLong, S.

997    Lory, E. Stackebrandt, and F. Thompson), pp. 660-747. Berlin Heidelberg.

998 73. Kilian, M., Riley, D. R., Jensen, A., Bruggemann, H., and Tettelin, H. 2014

999    Parallel evolution of *Streptococcus pneumoniae* and *Streptococcus mitis* to

1000    pathogenic and mutualistic lifestyles. *MBio* **5**, e01490-14. (doi:mBio.01490-

1001    14 [pii];10.1128/mBio.01490-14 [doi])

1002 74. Bishop, C. J., Aanensen, D. M., Jordan, G. E., Kilian, M., Hanage, W. P., and

1003    Spratt, B. G. 2009 Assigning strains to bacterial species via the internet. *BMC*

1004    *Biol* **7**, 3.

1005   75.   Didelot, X. *et al.*. 2011 Recombination and population structure in *Salmonella*
1006   *enterica*. *PLoS Pathog* **7**, e1002191.

1007   76.   den Bakker, H. C. *et al.*. 2011 Genome sequencing reveals diversification of
1008   virulence factor content and possible host adaptation in distinct
1009   subpopulations of *Salmonella enterica*. *BMC Genomics* **12**, 425.

1010   77.   Timme, R. E., Pettengill, J. B., Allard, M. W., Strain, E., Barrangou, R.,
1011   Wehnes, C., Van Kessel, J. S., Karns, J. S., Musser, S. M., and Brown, E. W.
1012   2013 Phylogenetic diversity of the enteric pathogen *Salmonella enterica*
1013   subsp. *enterica* inferred from genome-wide reference-free SNP characters.
1014   *Genome Biol Evol* **5**, 2109-2123. (doi:evt159 [pii];10.1093/gbe/evt159 [doi])

1015   78.   Parsons, S. K., Bull, C. M., and Gordon, D. M. 2011 Substructure within
1016   *Salmonella enterica* subspecies *enterica* isolated from Australian wildlife.
1017   *Appl Environ Microbiol* **77**, 3151-3153.

1018   79.   Worley, J., Meng, J., Allard, M. W., Brown, E. W., and Timme, R. E. 2018
1019   *Salmonella enterica* phylogeny based on whole-genome sequencing reveals
1020   two new clades and novel patterns of horizontally acquired genetic elements.
1021   *MBio* **9**. (doi:mBio.02303-18 [pii];10.1128/mBio.02303-18 [doi])

1022   80.   Key, F. M. *et al.*. 2020 Emergence of human-specific *Salmonella enterica* is
1023   linked to the Neolithization process. *Nature Ecology & Evolution* **4**, 324-333.
1024   (doi:https://doi.org/10.1038/s41559-020-1106-9)

1025   81.   Guibourdenche, M., Roggentin, P., Mikoleit, M., Fields, P. I., Bockemuhl, J.,
1026   Grimont, P. A., and Weill, F.-X. 2010 Supplement 2003-2007 (No. 47) to the
1027   White-Kauffmann-Le Minor scheme. *Res Microbiol* **161**, 26-29.

1028   82.   Issenhuth-Jeanjean, S., Roggentin, P., Mikoleit, M., Guibourdenche, M., De, P.
1029   E., Nair, S., Fields, P. I., and Weill, F.-X. 2014 Supplement 2008-2010 (no.
1030   48) to the White-Kauffmann-Le Minor scheme. *Res Microbiol* **165**, 526-530.
1031   (doi:S0923-2508(14)00106-5 [pii];10.1016/j.resmic.2014.07.004 [doi])

1032   83.   Grimont, P. A. and Weill, F.-X. 2007 *Antigenic formulae of the Salmonella*

1033   *serovars--9th edition* edition. Paris, France: WHO Collaborating Centre for

1034   Reference and Research on *Salmonella*.

1035   84.   Seif, Y., Monk, J. M., Machado, H., Kavvas, E., and Palsson, B. O. 2019

1036   Systems biology and pangenome of *Salmonella* O-Antigens. *MBio* **10**.

1037   (doi:mBio.01247-19 [pii];10.1128/mBio.01247-19 [doi])

1038   85.   Zhang, S., den Bakker, H. C., Li, S., Chen, J., Dinsmore, B. A., Lane, C., Lauer,

1039   A. C., Fields, P. I., and Deng, X. 2019 SeqSero2: rapid and improved

1040   *Salmonella* serotype determination using whole genome sequencing data.

1041   *Appl Environ Microbiol.* (doi:AEM.01746-19 [pii];10.1128/AEM.01746-19

1042   [doi])

1043   86.   Robertson, J., Yoshida, C., Kruczkiewicz, P., Nadon, C., Nichani, A., Taboada,

1044   E. N., and Nash, J. H. E. 2018 Comprehensive assessment of the quality of

1045   *Salmonella* whole genome sequence data available in public sequence

1046   databases using the *Salmonella* in silico Typing Resource (SISTR). *Microb*

1047   *Genom*, 1-11. (doi:10.1099/mgen.0.000151 [doi])

1048   87.   Uelze, L., Borowiak, M., Deneke, C., Szabo, I., Fischer, J., Tausch, S. H., and

1049   Malorny, B. 2020 Performance and accuracy of four open-source tools for *in*

1050   *silico* serotyping of *Salmonella* spp. based on whole-genome short-read

1051   sequencing data. *Appl Environ Microbiol* **86**. (doi:AEM.02265-19

1052   [pii];10.1128/AEM.02265-19 [doi])

1053   88.   Romano, S., Vinh, N. X., Bailey, J., and Verspoor, K. 2016 Adjusting for

1054   chance clustering comparison measures. *Journal of Machine Learning*

1055   *Research* **17**, 1-32.

1056   89.   Chaudhuri, R. R. and Henderson, I. R. 2012 The evolution of the *Escherichia*

1057   *coli* phylogeny. *Infect Genet Evol* **12**, 214-226.

1058   90.   Ochman, H. and Selander, R. K. 1984 Standard reference strains of *Escherichia*
1059   *coli* from natural populations. *J Bacteriol* **157**, 690-693.

1060   91.   Selander, R. K., Caugant, D. A., and Whittam, T. S. 1987 Genetic structure and
1061   variation in natural populations of *Escherichia coli*. In *Escherichia coli and*
1062   *Salmonella typhimurium cellular and molecular biology Vol.II* (eds. F. C.
1063   Neidhardt, J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H.
1064   E. Umbarger), pp. 1625-1648. Washington, D.C.: American Society for
1065   Microbiology.

1066   92.   Touchon, M., Perrin, A., de Sousa, J. A. M., Vangchhia, B., Burn, S., O'Brien,
1067   C. L., Denamur, E., Gordon, D., and Rocha, E. P. 2020 Phylogenetic
1068   background and habitat drive the genetic diversification of *Escherichia coli*.
1069   *PLoS Genet* **16**, e1008866. (doi:10.1371/journal.pgen.1008866 [doi])

1070   93.   Denamur, E., Clermont, O., Bonacorsi, S., and Gordon, D. 2021 The population
1071   genetics of pathogenic Escherichia coli. *Nat Rev Microbiol* **19**, 37-54.
1072   (doi:10.1038/s41579-020-0416-x [doi];10.1038/s41579-020-0416-x [pii])

1073   94.   Gordon, D. M., Clermont, O., Tolley, H., and Denamur, E. 2008 Assigning
1074   *Escherichia coli* strains to phylogenetic groups: multi-locus sequence typing
1075   versus the PCR triplex method. *Environ Microbiol* **10**, 2484-2496.

1076   95.   Clermont, O., Christenson, J. K., Denamur, E., and Gordon, D. M. 2013 The
1077   Clermont *Escherichia coli* phylo-typing method revisited: improvement of
1078   specificity and detection of new phylo-groups. *Environ Microbiol Rep* **5**, 58-
1079   65. (doi:10.1111/1758-2229.12019 [doi])

1080   96.   Beghain, J., Bridier-Nahmias, A., Le, N. H., Denamur, E., and Clermont, O.
1081   2018 ClermonTyping: an easy-to-use and accurate *in silico* method for
1082   *Escherichia* genus strain phylotyping. *Microb Genom* **4**.
1083   (doi:10.1099/mgen.0.000192 [doi])

97.   Waters, N. R., Abram, F., Brennan, F., Holmes, A., and Pritchard, L. 2020 Easy
phylotyping of *Escherichia coli* via the EzClermont web app and command-
line tool. *Access Microbiology* **2**, 1-5. (doi:DOI 10.1099/acmi.0.000143)

98.   Sawyer, C. *et al.*. 2021 Epidemiological investigation of recurrent outbreaks of
haemolytic uraemic syndrome caused by Shiga toxin-producing *Escherichia*
coli serotype O55:H7 in England, 2014-2018. *Epidemiol Infect* **149**, e108.
(doi:10.1017/S0950268821000844 [doi];S0950268821000844 [pii])

99.   Wick, L. M., Qi, W., Lacher, D. W., and Whittam, T. S. 2005 Evolution of
genomic content in the stepwise emergence of *Escherichia coli* O157:H7. *J*
Bacteriol **187**, 1783-1791.

100. Eichhorn, I., Semmler, T., Mellmann, A., Pickard, D., Anjum, M. F., Fruth, A.,
Karch, H., and Wieler, L. H. 2018 Microevolution of epidemiological highly
relevant non-O157 enterohemorrhagic *Escherichia coli* of serogroups O26
and O111. *Int J Med Microbiol* **308**, 1085-1095. (doi:S1438-4221(18)30241-
8 [pii];10.1016/j.ijmm.2018.08.003 [doi])

101. Liu, C. M. *et al.*. 2018 *Escherichia coli* ST131-H22 as a foodborne uropathogen.
MBio **9**. (doi:mBio.00470-18 [pii];10.1128/mBio.00470-18 [doi])

102. Price, L. B. *et al.*. 2013 The epidemic of extended-spectrum-beta-lactamase-
producing *Escherichia coli* ST131 is driven by a single highly pathogenic
subclone, H30-Rx. *MBio* **4**, e00377-13. (doi:mBio.00377-13
[pii];10.1128/mBio.00377-13 [doi])

103. Achtman, M., Mercer, A., Kusecek, B., Pohl, A., Heuzenroeder, M., Aaronson,
W., Sutton, A., and Silver, R. P. 1983 Six widespread bacterial clones among
Escherichia coli K1 isolates. *Infect Immun* **39**, 315-335.

104. Gordon, D. M. *et al.*. 2017 Fine-scale structure analysis shows epidemic patterns
of Clonal Complex 95, a cosmopolitan *Escherichia coli* lineage responsible

for extraintestinal infection. *mSphere* **2**. (doi:10.1128/mSphere.00168-17
[doi];mSphere00168-17 [pii])

105. Frentrup, M. *et al.*. 2020 A publicly accessible database for *Clostridioides*
difficile genome sequences supports tracing of transmission chains and
epidemics. *Microbial Genomics*, mgen.0.000410.
(doi:10.1099/mgen.0.000410 [doi])

106. Laukkanen-Ninios, R. *et al.*. 2011 Population structure of the *Yersinia*
pseudotuberculosis complex according to multilocus sequence typing.
Environ Microbiol **13**, 3114-3127.

107. Gladstone, R. A. *et al.*. 2019 International genomic definition of pneumococcal
lineages, to contextualise disease, antibiotic resistance and vaccine impact.
EBioMedicine **43**, 338-346. (doi:doi: 10.1016/j.ebiom.2019.04.021)

108. Lees, J. A., Harris, S. R., Tonkin-Hill, G., Gladstone, R. A., Lo, S. W., Weiser,
J. N., Corander, J., Bentley, S. D., and Croucher, N. J. 2019 Fast and flexible
bacterial genomic epidemiology with PopPUNK. *Genome Res* **29**, 304-316.
(doi:gr.241455.118 [pii];10.1101/gr.241455.118 [doi])

109. Lees, J. A., Tonkin-Hill, G., Yang, Z., and Corander, J. 2021 Mandrake:
visualising microbial population structure by embedding millions of genomes
into a low-dimensional representation. *BioRxiv*, 2021. (doi:doi:
https://doi.org/10.1101/2021.10.28.466232)

110. Tschoeke, D., Salazar, V. W., Vidal, L., Campeao, M., Swings, J., Thompson,
F., and Thompson, C. 2020 Unlocking the genomic taxonomy of the
Prochlorococcus collective. *Microb Ecol* **80**, 546-558. (doi:10.1007/s00248-
020-01526-5 [doi];10.1007/s00248-020-01526-5 [pii])

111. Salazar, V. W., Tschoeke, D. A., Swings, J., Cosenza, C. A., Mattoso, M.,
Thompson, C. C., and Thompson, F. L. 2020 A new genomic taxonomy

system for the *Synechococcus* collective. *Environ Microbiol* **22**, 4557-4570.
(doi:10.1111/1462-2920.15173 [doi])

112. Achtman, M. and Pluschke, G. 1986 Clonal analysis of descent and virulence
among selected *Escherichia coli*. *Annu Rev Microbiol* **40**, 185-210.

113. Carroll, K. J., Jenkins, C., Harvey-Vince, L., Mohan, K., and Balasegaram, S.
2021 Shiga toxin-producing *Escherichia coli* diagnosed by Stx PCR:
assessing the public health risk of non-O157 strains. *Eur J Public Health* **31**,
576-582. (doi:6067613 [pii];10.1093/eurpub/ckaa232 [doi])

114. Griffiths, D. *et al.*. 2010 Multilocus sequence typing of *Clostridium difficile*. *J*
Clin Microbiol **48**, 770-778.

115. Zhou, Z., Alikhan, N.-F., Sergeant, M. J., Luhmann, N., Vaz, C., Francisco, A.
P., Carrico, J. A., and Achtman, M. 2018 GrapeTree: Visualization of core
genomic relationships among 100,000 bacterial pathogens. *Genome Res* **28**,
1395-1404. (doi:DOI: 10.1101/gr.232397.117)

116. Achtman, M. and Zhou, Z. 2020 Metagenomics of the modern and historical
human oral microbiome with phylogenetic studies on *Streptococcus mutans*
and *Streptococcus sobrinus*. *Phil Trans R Soc B* **375**, 20190573.
(doi:http://dx.doi.org/10.1098/rstb.2019.0573)

117. Zhou, Z., Luhmann, N., Alikhan, N.-F., Quince, C., and Achtman, M. 2018
Accurate reconstruction of microbial strains from metagenomic sequencing
using representative reference genomes. In *RECOMB 2018*, pp. 225-240:
Springer, Cham.

118. Luhmann, N., Holley, G., and Achtman, M. 2021 BlastFrost: fast querying of
100,000s of bacterial genomes in Bifrost graphs. *Genome Biol* **22**, 30.
(doi:10.1186/s13059-020-02237-3 [doi])

119. Pedregosa, F. *et al.*. 2011 Scikit-learn: Machine learning in python. *Journal of*
Machine Learning Research **12**, 2825-2830.

120. Bostock, M., Ogievetsky, V., and Heer, J. 2011 D^3: Data-Driven Documents.

IEEE Trans Vis Comput Graph **17**, 2301-2309.

(doi:10.1109/TVCG.2011.185 [doi])

1166

1167

Table 1. Legacy data, newly assembled genomes and hierarchical cgST clusters in EnteroBase (09/2021).

Genus	Legacy MLST	cgMLST	# Genomic Loci		# HierCC clusters (HierCC level)		
	(# strains)	(# genomes)	wgMLST	cgMLST	ST Complexes	Lineages	Species/ Sub-species
Salmonella	4,930	312,196	21,065	3,002	3648 (900)	2185 (2000)	15 (2850)
Escherichia/ Shigella	9,525	175,256	25,002	2,513	1379 (1100)	343 (2000)	16 (2350)
Streptococcus		76,718	33,887	372	3681 (100)		132 (363)
Clostridioides		23,197	11,490	2,556	299 (950)		12 (2500)
Vibrio		12,267	152,249	1,128	2000 (800)		155 (1090)
Yersinia	4,286	4,023	19,591	1,553	451 (600)		34 (1490)

NOTE: # Genomic Loci: numbers of CDS whose alleles are automatically called for the wgMLST and cgMLST schemes. cut-off: maximum numbers of allelic differences within minimal spanning trees that define HC clusters of cgSTs. EnteroBase automatically calculates existing Legacy MLST STs according to legacy schemes for *Escherichia/Shigella* (Wirth *et al.*, 2006 [26], *Salmonella* (Achtman *et al.,* 2012 [27], and *Clostridioides* (Griffiths *et al.,* 2010 [114]), but does assign new STs nor does it maintain a database of legacy data from ABI sequencing.

Additional public databases are presented by EnteroBase for *Helicobacter* (>3500 genomes) and *Moraxella* (>2350 genomes), but these lack a cgMLST scheme. EnteroBase also has a database for >80,000 *Mycobacterium* genomes, but this currently (March, 2022) lacks a cgMLST scheme and a curator, and is not yet publicly available.

Table 2. EnteroBase-related software tools that support the analysis of large numbers of bacterial genomes

Name	Purpose	Citation	URL for Stand-alone version
GrapeTree	GUI for depicting and analysing minimal spanning and NJ trees of character data	[115]	https://github.com/achtman-lab/GrapeTree
PEPPAN	Pan-genome calculation including pseudo-genes from numerous representatives of an entire genus	[17]	https://github.com/zheminzhou/PEPPAN
SPARSE	Assignment of metagenomic reads to individual taxa	[116,117]	https://github.com/zheminzhou/SPARSE
EToKi	EnteroBase toolkit of useful functions and pipelines	[28]	https://github.com/zheminzhou/EToKi
BlastFrost	Efficient k-mer based search for DNA sequences in large genomic datasets	[118]	https://github.com/nluhmann/BlastFrost
HierCC	Automated hierarchical clustering of cgSTs to existing or novel clusters	[32]	https://github.com/zheminzhou/pHierCC

NOTE: PEPPAN is a stand-alone program that was used to generate the wgMLST schemes used in EnteroBase. SPARSE is a second stand-alone program that has been used to extract taxon-specific reads from metagenomes of ancient DNA that were then used with EToKi to define pseudo-MAGs that were uploaded to EnteroBase for phylogenetic comparisons. HierCC was used as a stand-alone program in development mode to define an initial set of clusters from a representative set of genomes. Subsequent assignments to existing clusters or to novel clusters were performed automatically within EnteroBase using pHierCCin production mode. BlastFrost is used by EnteroBase to determine the presence/absence of toxins and other pathovar characteristics of gastrointestinal pathogens within *E. coli*, and to assign gastrointestinal pathovar designations.

Table 3. Parameters of datasets used for calculating ML super-trees.

Genus	# Type strains	Total # Genomes	# Strict core genes	# Accessory genes	# Soft core genes (cgMLST)	# SNPs
Salmonella	7	10,002	1409	19,486	3002	1,410,331
Escherichia (EcoRPlus)	8	9479	134	20,294	2513	1,172,200
Escherichia reps	2	967	192	17,983	2513	971,516
Streptococcus	84	5937	90	31,630	372	263,080
Clostridioides	1	6725	515	10,975	2556	725,240
Vibrio	119	5032	124	146,041	1128	776,439
Yersinia	23	1847	744	18,799	1553	656,143

Each dataset consists of genomes representing the entire diversity of a genus as described in Methods (Supplemental Text).

Table 4. Quantitative concordance between eBG and HierCC clustering

Salmonella			
Test statistic	eBG-HC900	eBG-HC2000	HC900-HC2000
Adjusted Mutual Information Score (AMI)	0.985	0.948	0.942
Adjusted Rand Index (ARI)	0.982	0.856	0.866
Escherichia/Shigella			
Test statistic	ST Cplx-HC1100	ST Cplx-HC2000	HC1100-HC2000
Adjusted Mutual Information Score (AMI)	0.940	0.688	0.692
Adjusted Rand Index (ARI)	0.871	0.299	0.337
Streptococcus pneumoniae			
Test statistic	CC-HC100	GSPC-HC100	GSPC-HC160
Adjusted Mutual Information Score (AMI)	0.959	0.956	0.980
Adjusted Rand Index (ARI)	0.987	0.906	0.949

NOTE: *Salmonella*: Calculations were performed on 319,490 entries in EnteroBase which had been assigned to eBGs, HC900 and HC2000 clusters by December, 2021. The dataset only represented 411 eBGs, 690 HC900 clusters and 312 HC2000 clusters, and consists of a subset of all genomes because new eBGs have not been created in recent years.

Escherichia/Shigella: HC1100 and HC2000 assignments were from 143,520 genomes which had been assigned to ST Complexes as well as to HC1100 and HC2350 clusters in December, 2021. At that time 186,852 genomes had been assigned to HC1100 and HC2000 clusters, with AMI of 0.59 and ARI of 0.2.

Streptococcus pneumoniae consisted of GSPC assignments for 18,147 genomes from Supplemental Table 2 and CC assignments for 13,396 genomes from Supplemental Table 1 to Gladstone et al. [107] which were imported into User defined fields in EnteroBase as described [28].

ARI and AMI were calculated with the functions adjusted_rand_score() and adjusted_mutual_info_score() in the Python 3 library sklearn.metrics V1.0.1 [119].

Figure Legends

Figure 1. A comparison of species and sub-species assignments within *Salmonella* with HierCC and ANI. The figure shows an ML super-tree of 1,410,331 SNPs among 3002 core genes from 10,002 representative genomes of *Salmonella* (Table 3). Former sub-species IIIa is designated *S. arizonae* in accordance with Pearce *et al.,* 2021 [50]. A. Partitions differentiated by ANI 95% clusters (legend) correspond to species *S. enterica, S. bongori*, *S. arizonae* and a new species, *S. HC2850_215890* (five strains from the U.K., 2018-2020), as indicated by arrows, and sub-species are not differentiated. B. Partitions coloured by HC2850 clusters (legend). Arrows indicate HC2850_215890, and a new sub-species, HC2850_222931 (one strain from France, 2018). All other HC2850 clusters correspond to species (*S. bongori* and *S. arizonae*) or subspecies, except for HC2850_7171 (starred), which is subsp. *enterica* (I) according to the ML tree. An interactive version of this GrapeTree rendition can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53257. The corresponding presence/absence tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53258.

Figure 2. Maximum-likelihood core SNP tree of 967 *Escherichia* genomes consisting of one genome from each of 161 HC1100 clusters containing *E. coli* or *Shigella* as well as all 806 other *Escherichia* genomes in Enterobase as of November 2020. The tree is colored by A) pairwise FastANI values clustered at the 95% level and B) HC2350 cluster designations. The key legends indicates taxonomic designations in the literature which best match the cluster groupings. In part B, HC2350 cluster designations were used to mark novel taxonomic groupings in HC2350 clusters 89353, 89356, 89359 and 137132. An interactive version of this GrapeTree rendition

can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=52101. The

corresponding presence/absence tree can be found at

https://enterobase.warwick.ac.uk/ms_tree?tree_id=71125.

Figure 3. Species and sub-species assignments within *Clostridioides* according to

95% ANI (A) and HierCC (B). ML super-tree of 725,240 SNPs among 2556 core

genes from 6,724 representative genomes of *Clostridioides difficile* and one genome

of *Clostridioides mangenotii* (Table 3)*. A. ANI 95% clusters differentiate *C.

mangenotii* (cluster 1010) and five other clusters (clusters 255, 1370, 373, 2147,

1011) from *C. difficile* (cluster 0). Four of the 95% ANI clusters correspond to cryptic

clades C-I (clusters 255, 373), C-II (cluster 1370) and C-III (cluster 2147) in the

designations by Knight *et al.* [67]. Arrows indicate two additional clusters that were

distinguished by HierCC in part B. B. Partitions colored by HierCC assign the same

genomes to HC2500 clusters as ANI, except that HierCC assigns HC2500_15334 and

HC2500_15408 designations to one genome each. An interactive version of this

GrapeTree rendition can be found at

https://enterobase.warwick.ac.uk/ms_tree?tree_id=53253 and a presence/absence tree

can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53254.

Figure 5. Hierarchical population structure of O serogroups in *Salmonella*.

Hierarchical bubble plot of 310,901 *Salmonella* genomes in 790 HC900 clusters for

which a consensus O serogroup could be deduced by metadata, or bioinformatic

analyses with SeqSero V2 [85] or SISTR 1.1.1 [86]. Taxonomic level (HC level;

colors): species/subspecies (HC2850; light grey circles), Lineages (HC2000; dark

grey circles) and eBurst groups (HC900; O:group specific colors). Additional

information is indicated by yellow text for selected HC2000 and HC900 circles which

are specifically mentioned in the text. The diameters of HC900 circles are

proportional to the numbers of genomes. An interactive version of this figure can be

found at https://observablehq.com/@laurabaxter/salmonella-serovar-piechart from

which the representation, raw data and d3 Java code [120] for generating the plot can

be downloaded.

Figure 6. Hierarchical population structure of O serogroups in *Escherichia*

coli/Shigella. Hierarchical bubble plot for 167,312 genomes of *Escherichia coli* or

Shigella in HC2350_1 (large light grey circle) that were available in EnteroBase in

April, 2021. Seven HC2000 Lineages encompassing 15 HC1100/ST Complexes of

Shigella are shown at the right. HC2350_1 also includes 15 other *E. coli* HC2000

Lineages that each contains at least 50 *E. coli* genomes and encompass 144 other

HC1100 clusters. The remainder of the figure shows those HC1100 clusters and not

the corresponding HC2000 clusters. Numbers of genomes assigned to individual O

serogroups (legend) are indicated by pie chart wedges within the HC1100 circles.

Selected HC1100 clusters are also depicted with indications of phenotype and

nomenclature at a larger scale outside the main circle, connected to the original circles

by lines. An interactive version of this figure can be found at

https://observablehq.com/@laurabaxter/escherichia-serovar-piechart, from which the

representation, raw data and d3 Java code [120] for generating the plot can be

downloaded.

Figure 1. Species and sub-species assignments within *Salmonella* with HierCC and ANI. ML super-tree of 1,410,331 SNPs among 3002 core genes from 10,002 representative genomes of *Salmonella*. Former sub-species IIIa is designated *S. arizonae* in accordance with Pearce et al., 2021 [48]. A. Partitions differentiated by ANI 95% clusters (legend) correspond to species (*S. enterica, S. bongori, S. arizonae, S.* HC2850_215890), as indicated by arrows, and sub-species are not differentiated. B. Partitions coloured by HC2850 clusters (legend). Arrows indicate a new species, HC2850_215890 (five strains from the U.K., 2018-2020), and a new sub-species, HC2850_222931 (one strain from France, 2018). All other HC2850 clusters correspond to species (*S. bongori* and *S. arizonae*) or subspecies, except for HC2850_7171 (starred), which is subsp. *enterica* (I) according to the ML tree. An interactive version of this GrapeTree rendition can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53257. The corresponding presence/absence tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53258.

Figure 2. Maximum-likelihood core SNP tree of 967 *Escherichia* genomes consisting of one genome from each of 161 HC1100 clusters containing *E. coli* or *Shigella* as well as all 806 other *Escherichia* genomes in Enterobase as of November 2020. The tree is colored by A) pairwise FastANI values clustered at the 95% level and B) HC2350 cluster designations. The key legends indicates taxonomic designations in the literature which best match the cluster groupings. In part B, HC2350 cluster designations were used to mark novel taxonomic groupings in HC2350 clusters 89353, 89356, 89359 and 137132. An interactive version of this GrapeTree rendition can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=52101. The corresponding presence/absence tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=71125.

Figure 2. Species and sub-species assignments within *Clostridioides* according to ANI (A) and HierCC (B). ML super-tree of 725,240 SNPs among 2556 core genes from 6,724 representative genomes of *Clostridioides difficile* and one genome of *Clostridioides mangenotii*. A. ANI 95% clusters (legend) differentiate *C. mangenotii* (cluster 1010) and five other clusters (clusters 255, 1370, 373, 2147, 1011) from *C. difficile* (cluster 0). As indicated, four of these correspond to cryptic clades C-I [clusters 255, 373], C-II [cluster 1370] and C-III [cluster 2147]) according to Knight et al. [49]. Arrows indicate two additional clusters that were distinguished by HierCC in part B. B. Partitions coloured by HierCC assigns the same genomes as ANI to the corresponding HC2500 clusters, except that it defines HC2500_15334 and HC2500_15408 to one genome each, which might represent sub-species. An interactive version of this GrapeTree rendition can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53253 and a presence/absence tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53254.

Figure 3. Comparison of HC363 clusters with taxonomic designations in *Streptococcus*. ML super-tree of 263,080 SNPs among 372 core genes from 5937 representative *Streptococcus* genomes. Species names are indicated next to the phylogenetic clusters where genomes mapped for type strains, and according to published metadata. Nodes are colored by HC363 clusters, and exceptional assignments are indicated by asterisks next to *S. pneumoniae* and *S. pseudopneumoniae,* which were both HC363_99; multiple phylogenetic and HierCC clusters grouped within *S. suis*; *S. salivarius* and *S. vestibularis*, which were both HC353_202; *S. lutetiensis and S. equinus*, which were both HC363_181; and *S. dysgalactiae and S. pyogenes*, which were both HC363_139. An interactive version of the GrapeTree rendition of the SNP tree can be found at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53261 and the presence/absence tree at https://enterobase.warwick.ac.uk/ms_tree?tree_id=53262.

S. enterica subsp. I HC2850_2

HC2000_299 Agama Lineage

HC900_127074 Nigeria

HC900_299 Agama

HC2000_106 Lineage

HC2000_2 Typhimurium Lineage

HC900_2 Typhimurium

HC2000_12 Enteritidis Lineage

HC900_12 Enteritidis

Other species/subspecies

HC2000_1272 Para C Lineage

HC900_4381 Paratyphi C

HC2000_54 Lineage

HC2000_4 Mbandaka Lineage

HC900_4 Mbandaka

O antigens

O:9	O:16	O:11	O:55
O:61	O:48	O:40	O:65
O:4	O:50	O:44	O:51
O:8	O:45	O:41	O:39
O:7	O:18	O:58	O:54
O:3,10	O:21	O:53	O:66
O:13	O:1,3,19	O:43	O:62
O:9,46	O:2	O:9,46,27	O:56
O:35	O:42	O:38	O:57
O:30	O:47	O:60	O:67
O:28	O:6,14	O:17	-

Figure 5. Hierarchical population structure of O serogroups in *Salmonella*. Hierarchical bubble plot of 310,901 *Salmonella* genomes in 790 HC900 clusters for which a consensus O serogroup could be deduced by metadata, or bioinformatic analyses with SeqSero V2 [85] or SISTR 1.1.1 [86]. Taxonomic level (HC level; colors): species/subspecies (HC2850; light grey circles), Lineages (HC2000; dark grey circles) and eBurst groups (HC900; O:group specific colors). Additional information is indicated by yellow text for selected HC2000 and HC900 circles which are specifically mentioned in the text. The diameters of HC900 circles are proportional to the numbers of genomes. An interactive version of this figure can be found at https://observablehq.com/@laurabaxter/salmonella-serovar-piechart from which the representation, raw data and d3 Java code [120] for generating the plot can be downloaded.

HC1100_2
(ST29 Cplx;
EHEC)

O103

O111

O26

HC1100_44
(ST95 Cplx;
NMEC, UPEC, APEC)

O45

O25

O2

O1

O18

B2 B4 D13

F6

D3

D9

B1

B18

HC2000_1463
S. dysenteriae

HC2000_1465 (ST243 Cplx)
S. flexneri, boydii
& dysenteriae

HC2000_305
(ST152 Cplx) *S. sonnei*

HC2000_192
(ST245 Cplx)

O55

HC1100_63
(ST11 Cplx;
EHEC)

O45

O128 O111

HC1100_3
(ST20 Cplx;
EHEC)

HC1100_7
(ST131 Cplx;
UPEC)

O antigens

■ B1	■ D9	■ O13	■ O25	■ O83
■ B13	■ F1-5	■ O145	■ O26	■ O86
■ B16	■ F6	■ O146	■ O4	■ O88
■ B18	■ SS	■ O15	■ O45	■ O89
■ B2	■ O1	■ O153	■ O5	■ O9
■ B4	■ O102	■ O157	■ O55	■ O91
■ B9	■ O103	■ O16	■ O6	■ Others
■ D1	■ O11	■ O17	■ O7	□ Missing
■ D2	■ O111	■ O18	■ O75	
■ D3	■ O121	■ O2	■ O78	
	■ O128	■ O21	■ O8	

Figure 5. Hierarchical population structure of O serogroups in *Escherichia coli/Shigella*. Hierarchical bubble plot for 167,312 genomes of *Escherichia coli* or *Shigella* in HC2350_1 (large grey circle) that were available in EnteroBase in April, 2021. Seven HC2000 Lineages encompassing 15 HC1100/ST Complexes of *Shigella* are arbitrarily indicated at the right. HC2350_1 also includes 15 other *E. coli* HC2000 Lineages that encompass 144 other HC1100 clusters that each contains at least 50 *E. coli* genomes. The figure shows only the HC1100 clusters and not the corresponding HC1100 clusters. Numbers of genomes assigned to individual O serogroups (legend) are indicated by pie chart wedges within the HC1100 circles. Selected HC1100 clusters are drawn on larger scale with indications of phenotype and nomenclature outside the main circle, connected to the original circles by lines. An interactive version of this figure can be found at https://observableh-q.com/@laurabaxter/escherichia-serovar-piechart, from which the representation, raw data and d3 Java code [103] for generating the plot can be downloaded.