Background selection from unlinked sites causes non-independent evolution of deleterious mutations

Joseph Matheson[1,2], Joanna Masel[1]

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721, USA

[2]Department of Ecology, Behavior, and Evolution, University of California San Diego, San Diego, CA, 92093, USA

Corresponding author: Joanna Masel

ABSTRACT

Background selection describes the reduction in neutral diversity caused by selection against deleterious alleles at other loci. It is typically assumed that the purging of deleterious alleles affects linked neutral variants, and indeed simulations typically only treat a genomic window. However, background selection at unlinked loci also depresses neutral diversity. In agreement with previous analytical approximations, in our simulations of a human-like genome with a realistically high genome-wide deleterious mutation rate, the effects of unlinked background selection exceed those of linked background selection. Background selection reduces neutral genetic diversity by a factor that is independent of census population size. Outside of genic regions, the strength of background selection increases with the mean selection coefficient, contradicting the linked theory but in agreement with the unlinked theory. Neutral diversity within genic regions is fairly independent of the strength of selection. Deleterious genetic load among haploid individuals is underdispersed, indicating non-independent evolution of deleterious mutations. Empirical evidence for underdispersion was previously interpreted as evidence for global epistasis, but we recover it from a non-epistatic model.

Keywords: linkage disequilibrium, population genetics, nearly neutral theory, forward time simulation, effective population size, expected heterozygosity

SIGNIFICANCE

As individuals bearing deleterious alleles are removed from a population, other alleles are removed with

them, some that are tightly linked near the deleterious allele on a chromosome and some that aren't linked

at all. When the deleterious mutation rate is realistically high, unlinked pairs of loci are a more important

influence on the removal of genetic variation. Simulations that assume independent evolution cannot

capture removal just by using a lower "effective population size", because the probabilities of having

deleterious alleles on different chromosomes are negatively correlated rather than independent.

31   INTRODUCTION

The neutral theory of molecular evolution postulates that i) most genetic diversity observed in natural

populations is neutral with respect to an organism's fitness (Kimura 1968; King and Jukes 1969), and ii)

dynamics are well-described by models of a single neutral locus in a population of a specified "effective"

population size (Charlesworth 2009; Ewens 2004; Kern and Hahn 2018; Masel 2011). A crucial extension

of this theory uses a reduction in the effective population size to incorporate the fact that deleterious

mutations depress genetic diversity as they are removed, in a process known as background selection

(Charlesworth, Morgan, and Charlesworth 1993). While the removal of a deleterious allele will depress

genetic diversity at neutral loci across the entire genome, neutral loci linked to the deleterious allele will

be particularly affected (Charlesworth 2012). Indeed, the effect of unlinked background selection is often

considered small enough to be ignored (Charlesworth 2012). Here we revisit this assumption given

realistically high genome-wide deleterious mutation rates, by using simulations to evaluate previously

derived equations for linked vs. unlinked background selection.

In practice, observed genetic diversity at putatively neutral sites is used to estimate the coalescent

effective population size N_e as the census size of an idealized population that produces the same neutral

genetic diversity given demography (Charlesworth 2009) or background selection (Charlesworth et al.

1993). In an idealized population obeying Wright-Fisher or Moran dynamics, genetic diversity depends

only on the product of the neutral mutation rate at that locus and the census size of the population

(Kimura 1969). Neutral theory considers mutations that either are strictly neutral or so deleterious that

they are purged quickly enough as to leave no impact. Nearly neutral theory retains the binary distinction

between rapidly purged versus neutral mutations, but allows the ratio of mutations of these two types to

vary among species, according to that species' value of an effective population size (Ohta 1973). Both

models assume many independent single loci.

The problem with this binary distinction and independent loci approach is that slightly deleterious

mutations are purged only slowly from populations. During this removal process, they depress genetic

variation in the genome (Charlesworth et al. 1993). This depression in genetic variation caused by

background selection is typically modeled as a decrease in the effective population size for a neutral locus

linked to deleterious variants (Comeron 2014; Hudson and Kaplan 1995; Lohmueller et al. 2011). In a

population with no recombination and sufficiently large effect deleterious mutations ($N_e s \gg 1$), the

coalescent effective population size would decrease from N_e to $f_0 N_e$, where f_0 is the equilibrium

frequency of individuals with no deleterious mutations, because any neutral variants linked to deleterious

variants would be doomed (Charlesworth et al. 1993).

Recombination can decouple neutral variants from deleterious variants, reducing the degree to which

background selections depresses neutral variation (Cutter and Payseur 2013). For a single neutral locus

linked to a single locus where deleterious mutations with fixed heterozygous effect size sh occur at rate u

per diploid individual per generation, and with recombination between the loci occurring at rate r,

heterozygosity at the neutral locus is reduced by a factor $F \approx 1 - \frac{ush}{2(sh+r)^2}$ (Hudson and Kaplan 1994).

This result can be straightforwardly extended to any number of deleterious sites linked to the focal neutral

site by assuming i) that mutation and recombination rates are uniform across a genomic window centered

on the neutral site, ii) that the window is small enough that recombination rate and map length are linearly

related, and iii) that there is linkage equilibrium among deleterious variants e.g. because multiple

significantly linked deleterious mutations are not present at the same time (Hudson and Kaplan 1995;

Nordborg, Charlesworth, and Charlesworth 1996). In this case, in a genomic window we have

74
$$\frac{N_e}{N} = e^{-U_w/2sh+R_w} \quad (1)$$

75   where U_w is the total diploid deleterious mutation rate across the entire window, and R_w is the map length

76   between the ends of the window (Hudson and Kaplan 1995). Assuming a sufficiently large window size

77   (such that $R_w \gg s$), this equation simplifies to

78
$$\frac{N_e}{N} = e^{-u/r} \quad (2)$$

where u/r is the ratio of deleterious mutation rate to recombination rate (Charlesworth 2012). With the

same assumptions, similar results can be obtained in the case where both deleterious and beneficial

mutations occur (Kim and Stephan 2000).

The results above do not apply across different chromosomes or given free recombination between the

ends of the window (Hudson and Kaplan 1995). However, deleterious variants do still depress neutral

diversity even at unlinked sites: at a bare minimum, a deleterious mutation will eliminate any unique

neutral genetic variants in a single individual who dies due to that deleterious mutation. The ratio of N_e to

N with free recombination has been derived as

87
$$\frac{N_e}{N} = e^{-8Ush} \quad (3)$$

where U is the total deleterious mutation rate at unlinked loci and sh is the fitness effect of a deleterious

allele when in the heterozygous state (Charlesworth 2012).

It has previously been presumed that the reduction in diversity due to unlinked deleterious loci will be

much smaller than that from linked loci (Charlesworth 2012). That is, for any given neutral locus, even

though U across all unlinked deleterious loci is much larger than U_w across the window of linked

deleterious loci, $1/(2sh + R_w)$ will be so much larger than $8sh$ that the latter comparison will overwhelm

the former comparison, and linked background selection will dominate. Indeed, models of background

selection often simply omit the effects of unlinked background selection (Elyashiv et al. 2016; Ewing and

Jensen 2016; McGee et al. 2022; Torres, Szpiech, and Hernandez 2018).

But omitting the effects of unlinked background selection may be a concern given the sheer quantity of

deleterious mutations entering populations, and the greater dependence of Equation 3 on this rate. For

example, the average number of new deleterious mutations per human is estimated to be at least 2.1

(Lesecque, Keightley, and Eyre-Walker 2012). This estimate is derived from a point mutation rate of

1.1×10^{-8}, mutations only being deleterious in the 55% of the 6×10^{9} diploid genome that is not dominated

by the remnants of transposable elements, which evolves due to this constraint at 94.3% of the rate. This

estimate is conservative because some mutations to transposable element regions are deleterious, because

more recent estimates of the human point mutation rate are slightly higher at $\sim1.341\times10^{-8}$ (Wang and

Obbard 2023), and because non-point mutations and beneficial mutations are neglected. Some therefore

argue that deleterious mutation rates are even higher, closer to 10 new deleterious mutations per person in

humans (Kondrashov 2017). High deleterious mutation rate estimates are not unique to humans (Haag-

Liautard et al. 2007; Popovic et al. 2023).

Here we numerically compare Equations 2 and 3 given human parameters, confirming that unlinked

selection has the larger effect. This wouldn't matter for some purposes, so long as unlinked selection were

well described by one-locus models of genetic drift with a lower effective population size. To test this, we

perform a multi-locus simulation using the fwdpy11 package (Thornton 2014, 2019), which efficiently

handles large numbers of non-neutral mutations in relatively large census size populations (Haller and

Messer 2017). These simulations agree reasonably well with past analytic results, confirming the

importance of unlinked background selection. More importantly, simulations show that deleterious

mutations are not well described by one-locus models, with underdispersion of genetic load among

haploid genotypes. This pattern of underdispersion has previously been observed empirically for humans

and fruit flies, and interpreted as evidence for epistasis (Sohail et al. 2017); we recover it in a non-

epistatic model.

Unlinked background selection reduces neutral diversity more than linked background selection does

(Figure 1A) when the genome-wide deleterious mutation rate U is high, specifically when $U > 1$, as is

estimated to be the case for humans (Lesecque et al. 2012). This main result is independent of N and s. A

five-fold change in census population size N has no significant effect on the $\frac{N_e}{N}$ ratio (Figure 1A). This

contradicts models that imply that the impact of background selection is likely to be greater in larger

populations (Corbett-Detig, Hartl, and Sackton 2015; Cutter and Payseur 2013), although deviations from

such models have been previously found (Gillespie 2001; Kaplan, Hudson, and Langle 1989; Santiago

and Caballero 2016).

Figure 1. At high deleterious mutation rates, unlinked background selection reduces neutral

diversity to a greater extent than linked background selection does. Where not shown, $sh = -0.01$

and $N = 5000$. Each point represents a single simulation — we chose to allocate computation to a denser

grid of parameter values rather than to replicates of the same parameter values. (A) Unlinked background

selection alone (dashed line) is a closer match than linked background selection alone (solid line) to the

joint model (dot-dashed line) and to simulations (points). Census population size has no effect on the $\frac{N_e}{N}$

ratio across a five-fold change in simulated census population size. (B) Background selection at high

mutation rates is stronger with larger selection coefficients, although our joint model exaggerates this

effect. The model with linked background selection alone predicts the opposite dependence on sh, and

strongly underestimates background selection. (C) The relationship between deleterious mutation rate and

N_e is similar whether the effect size of new deleterious mutations is constant versus drawn from a

distribution of effect sizes with the same mean value of -0.01.

Larger selection coefficients result in stronger background selection for high U but not low U (Figure 1B).

This is consistent with a dominant role for unlinked background selection (Eq. 3); linked background

selection predicts the opposite effect (Eq. 1). Our approximate analytical joint model (see Methods)

exaggerates the impact of selection strength; for our focal value of $sh = -0.01$, our joint model slightly

underestimates background selection (Figure 1A and C), distinctly underestimates it for smaller selection

coefficients (Figure 1B green), and slightly overestimates it for large selection coefficients (Figure 1B

blue). The underestimation for smaller selection coefficients has been noted before, and likely occurs

because the small coefficients are near a weakly deleterious range, violating the assumption that $N_e s \gg 1$

(Johri, Charlesworth, and Jensen 2020). A distribution of selective effect sizes behaves similarly to a

single sh value with the same mean (Figure 1C).

The simulations above assume that deleterious mutations occur uniformly at random across the genome.

A more realistic scenario would be for deleterious mutations to be clustered within a functional subset of

the genome. We modify our simulations to model genomes where only 10% of the genome is made up of

'genes' subject to deleterious mutations. Concentrating deleterious mutations into more tightly linked

'genes' results in a smaller reduction in neutral diversity than found in simulations where deleterious

mutations occur across the genome (Figure 2, red vs. blue circles). This does not depend on population

size (vertical comparison on Figure 2A, B, D).

Figure 2. Concentrating deleterious mutations into 'genes' slightly weakens overall background

selection, and for genic sites, removes the dependence on selection coefficients. As in Figure 1,

background selection becomes significant only at high mutation rates, and census population size does not

affect results (A vs B vs D). For large selection coefficients, background selection is similar at genic vs

intergenic sites (E, red triangles vs squares). For small selection coefficients, the gap is considerable (C,

red circles vs squares). The dependence on the strength of selection (circles in C vs D vs E, blue matches

Figure 1) does not apply to genic sites (red triangles in C vs D vs E). In all panels, the solid black line is

the theoretical expectation incorporating both linked and unlinked background selection, blue dots are

simulations where deleterious mutations occur uniformly at random on the genome, and red dots are

simulations where deleterious mutations are clustered into 'genes' (see Methods). Within the genes

condition, neutral sites within genes are indicated as red triangles, and outside them are indicated as red

squares.

Simulations quantify the degree to which neutral diversity is more depressed in genes than in intergenic

regions (Figure 2, red squares vs. red triangles in panels C, D, and E). At low mutation rates, there is little

depression in neutral diversity, with no appreciable difference between linked vs. unlinked sites. There is

also little difference with strong selection (Figure 2, panel E); this is expected as we approach the limit at

which each deleterious mutation immediately dooms the genome it appears on (Charlesworth et al. 1993).

While larger selection coefficients increase overall background selection (discussed above for Fig. 1 and

seen for circles in Figure 2 C-E), background selection in genic regions is relatively independent of

selection coefficient (red triangles in Figure 2 C-E) and corresponds well to the joint model for $sh =$

-0.01 (Figure 2D).

Strong background selection might be well captured by traditional one-locus models with lower N_e, if

deleterious sites evolve independently. However, at high deleterious mutation rates, linkage

disequilibrium occurs (Barton 1998; Barton and Otto 2005), potentially breaking this assumption. To test

for independence, we asked whether the variance in the number of deleterious alleles is lower than that

expected from the mean under a Poisson distribution (Figure 3). We find that independent evolution of

sites breaks down for $U > 1$. Underdispersion among deleterious alleles has been found empirically for

humans and fruit flies (Lee 2022; Sohail et al. 2017) (both with $U > 1$ (Haag-Liautard et al. 2007;

Lesecque et al. 2012)). The index of dispersion was 0.94 for missense variants in the human "crucial

genome" in which these variants are most likely to be deleterious (Sohail et al. 2017). Note that Sohail et

al. (2017) and Lee (2022) interpreted underdispersion as evidence for synergistic epistasis, but we find

underdispersion even in a non-epistatic model.

Figure 3. Deleterious mutation rates above 1 create non-independence among segregating sites. The

index of dispersion is calculated as the variance in the number of deleterious alleles per haplotype divided

by the mean number of deleterious alleles per haplotype. A haplotype is defined as all the chromosomes

from one parent, i.e. each individual contains two haplotypes. A Poisson distribution has an index of

dispersion of 1, shown as a horizontal line. Points are the mean index of dispersion from eleven replicate

simulations with $N = 10,000$, $sh = -0.01$, and bars show the standard error in the estimate of the mean.

199   DISCUSSION

High deleterious mutation rates are well established empirically in humans and a variety of other species.

Our simulations confirm previous analytic results, showing that when deleterious mutation rates are

realistically high, unlinked background selection reduces neutral diversity more than linked background

selection does. These findings do not depend on the census population size. Background selection is

stronger with larger selection coefficients when deleterious sites are distributed uniformly at random, but

when deleterious mutations are clustered into genes, this dependence disappears at neutral sites within

genic regions. Existing analytical models exaggerate the dependence on selection coefficients,

highlighting our incomplete understanding of unlinked background selection. The original view was that

high-s mutations at sites that are unlinked to the focal site would exclude an effectively random set of

individuals in each generation from the effective population size. This heuristic does not easily explain

our findings. However, the approximate fit of our simulation results to models suggest that we are not

operating in the interference selection regime characterized by Good et al. (2014).

We focused on the index of dispersion as our genome-wide metric of linkage disequilibrium. There have

been recent attempts to disentangle pair-wise measures of linkage disequilibrium from their dependence

on allele frequency (Garcia and Lohmueller 2021; Good 2022; Potapova and Kondrashov 2023), and the

complexities of unphased data (Ragsdale and Gravel 2020). It would be interesting for such work to

include unlinked controls in future.

Our multilocus simulations neglect some population features known to affect neutral diversity (e.g.

adaptive evolution (Maynard Smith and Haigh 1974) and temporal changes in population size (Torres et

al. 2020)), and simplify others (e.g. variation in dominance coefficients among deleterious variants

(Gilbert et al. 2020) and heterogeneity in recombination rates (Kulathinal et al. 2008)). The purpose of

our simulations is to isolate the effects of background selection with high mutation rates, rather than to

accurately reflect the genetics of specific populations. Incorporating additional complications into the

model might change the quantitative strength of background selection. However, we do not expect adding

new complications to change the broader conclusion that unlinked background selection cannot be safely

ignored.

It is already known that the effects of linked background selection are critical for explaining differences in

neutral genetic diversity among genomic regions, and that failure to do so is a problem for demographic

inference (Ewing and Jensen 2016; Johri et al. 2021; Pouyet et al. 2018). Our results bring up the

possibility that unlinked background selection could also be important. At low mutation rates ($U \ll 1$, see

Figure 3), the effects of unlinked background selection are trivial in magnitude. But at realistically high

mutation rates ($U > 1$), unlinked background selection will result in non-independent evolution among

sites. Non-independence among loci cannot be accounted for with a fudge factor in the effective

population size of a one-locus model. In a one-locus model, random changes to allele frequencies are

modelled as white noise, i.e. there is no autocorrelation over time. When sites are not independent, being

on a bad/good genetic background in one generation will predict being on a bad/good genetic background

in the next generation, producing colored noise (Masel 2011). We therefore cannot assume that a one-

locus model with lower N_e can accurately capture background selection at unlinked loci.

To save computational cost, previous simulations of background selection treated only a section of a

chromosome. The local deleterious mutation rate is set to a value corresponding to high genome-wide U,

but mutations outside the local window are neglected. We have shown that neglecting them is

problematic. This could matter both in the context of demographic inference, and when the goal is to use

neutral diversity to distinguish between the effects of linked background selection vs. selective sweeps

(Elyashiv et al. 2016; McVicker et al. 2009; Murphy et al. 2022). Neglecting unlinked background

selection is also a concern for papers investigating only negative selection; e.g. Torres *et al*. (2018) and

Beissinger *et al*. (2016) look for differences in background selection between human or between maize

populations, but their measure of background selection assumes a reference class of neutral sites that are

unaffected, to be compared to sites subject to strong background selection.

Another common practice in evolutionary simulations is to rescale parameters (e.g. N, s, U, r) in a

manner that keeps products of interest (e.g. Ns, Nr) constant (Campos and Charlesworth 2019; Comeron

and Kreitman 2002; Hill and Robertson 1966; Hoggart et al. 2007; Kaiser and Charlesworth 2009;

Uricchio and Hernandez 2014). We found that the strength of background selection is independent of N,

but is stronger for large s (at least outside of genic regions). Reducing N while increasing s will therefore

exaggerate the effect of background selection, i.e. the reduction in N_e will be greater than that expected as

proportional to the reduction in simulated N. The net result of rescaling as it is normally conducted would

thus be a reduction in sN_e in populations with high U.

Accounting for background selection is often conceived as a necessary step prior to inferring demography

and/or adaptation. But unlinked background selection in species with high mutation rates globally

suppresses neutral diversity to such a degree that it is worthy of scientific attention in its own right. E.g.

discussions of whether background selection is sufficient to resolve Lewontin's paradox (Buffalo 2021)

should consider high U_d and the magnitude of the unlinked background selection that it causes.

To the extent that currently neutral genetic diversity might become relevant in new environments,

background selection imposes a genome-wide penalty caused by the constant deluge of deleterious

mutations. Massive background selection is thus a sister question to that of overwhelming deleterious

mutation load, which has not yet been fully resolved (Agrawal and Whitlock 2012; Goyal et al. 2012;

Kondrashov 1995, 2017).

<div align="center">MATERIALS AND METHODS</div>

Joint model

To calculate the expected combined effects of background selection at linked and unlinked sites, we

simply multiply together the respective reductions in neutral diversity from Equations 2 and 3. To avoid

double counting, we subtract from unlinked selection those mutations that fall within a window of

presumed linked selection. Considering constant sh for convenience, this joint model is given by

$$\frac{N_e}{N} = e^{-8(U-U_w)sh}e^{-U/r} \quad (4)$$

where U is the total genome-wide deleterious mutation rate. We assume 20,000 windows (this number is

chosen to roughly match the number of genes in humans); using different window numbers has minimal

impact (Supplementary Figure 1A). Results are qualitatively the same if we use a joint model constructed

from equations 1 and 3, with slightly more dependence on window size (Supplementary Figure 1B).

Multi-locus simulations

All simulations were written in Python using fwdpy11 (Thornton 2014, 2019). We simulated populations

of *N* diploid individuals undergoing selection against deleterious mutations using a standard Wright-

Fisher infinite-sites model for 10*N* generations. Each individual's genome was made up of 23

chromosomes of equal length, with recombination occurring via exactly two cross-overs per chromosome,

matching data for humans (Pardo-Manuel De Villena and Sapienza 2001).

Deleterious mutations occur with genome-wide rate *U*. In the "no genes" condition, they are located

uniformly at random along the chromosomes, while in the "genes" condition they occur only in "genes".

We simulate 1,000 genes, accounting for 10 percent of the genome, interspersed at regular intervals

throughout the genome. Assumptions about "genes" were not chosen to be representative of any

particular species, but simply to capture the qualitative consequences of clustering the sites that are

subject to deleterious mutations and hence background selection.

A recent study of a large sample of modern European humans estimated a gamma distribution of fitness

effects of new non-synonymous mutations with mean $sN_e = -224.33$ and $2N_e = 23,646$, implying a

mean $sh \approx$ -0.01 (Kim, Huber, and Lohmueller 2017). In our main results, we simplify to use a constant

sh = -0.01 to avoid complications from deleterious mutations with sN_e so near to 1 that they are

effectively neutral. We also explore higher and lower values of *s*, and a gamma distribution with the same

mean and shape parameter $\alpha = 0.169$ (Kim et al. 2017). All mutations have $h = 0.5$ and fitness is

calculated multiplicatively with no epistasis.

While our forward-time simulations track only deleterious mutations, tree-sequence recording (Kelleher

et al. 2018) during the simulation allows neutral mutations to be projected backwards onto the

genealogical histories of different genomic regions, enabling us to later compute neutral genetic diversity

and hence effective population size. In all simulations, neutral mutations occur uniformly at random on

the entire genome at an arbitrary rate 10^{-4} per genomic 'unit', for a total rate of 0.23 per genome. This

low value provides sufficient resolution of N_e at low computational cost. We use msprime (Kelleher,

Etheridge, and McVean 2016) to calculate neutral diversity θ on the resulting tree sequence using an

infinite-alleles model, and then calculate the effective population size for a simulation using $\theta = \frac{4N_e\mu}{1+4N_e\mu}$

and solving for N_e. When calculating effective population size within genes vs. intergenic regions (red

squares vs. triangles in Figure 2C-E) we use an infinite-sites model to avoid complications with the

distribution of finite neutral sites in genes vs. intergenic regions. In these cases, we calculate the effective

population size using $\theta = 4N_e\mu$ and solving for N_e.

We simulated census population sizes N ranging from 2,000 to 10,000. This is compatible with the range

of inferred estimates for human effective population sizes (McEvoy et al. 2011; Takahata 1993; Tenesa et

al. 2007). We calculate the 'observed' value of $\frac{N_e}{N}$ from the neutral diversity, to compare with analytical

expectations.

312   DATA AVAILABILITY

313   Simulation code written in python, and graphs produced with R. Scripts available on github at

314   www.github.com/MaselLab/BackgroundSelection.

315   ACKNOWLEDGEMENTS

We thank Bruce Walsh for pointing us to Hudson & Kaplan's analytical results on linked background

selection that sparked this work; Denis Roze for a wealth of advice (including pointing us to the equation

for unlinked background selection), detailed comments on the manuscript, and catching many errors on

our part; two anonymous reviewers of an earlier version of this work who also pointed us to unlinked

background selection; Kevin Thornton for pointing out a possible connection with implausibly high

empirical estimates of the deleterious mutation rate, and for assistance with fwdpy simulations; and Ryan

Gutenkunst, Parul Johri, and David Enard for helpful discussions.

323   FUNDING

324 This work was supported by the John Templeton Foundation [62028].

325   CONFLICT OF INTEREST

326 The authors declare no conflict of interest.

328   BIBLIOGRAPHY

Agrawal, Aneil F., and Michael C. Whitlock. 2012. 'Mutation Load: The Fitness of Individuals in
Populations Where Deleterious Alleles Are Abundant'. *Annu. Rev. Ecol. Evol. Syst* 43:115–35. doi:
10.1146/annurev-ecolsys-110411-160257.

Barton, N. H. 1998. 'The Effect of Hitch-Hiking on Neutral Genealogies'. *Genetical Research* 72(2):123–
33. doi: 10.1017/S0016672398003462.

Barton, N. H., and Sarah P. Otto. 2005. 'Evolution of Recombination Due to Random Drift'. *Genetics*
169(4):2353–70. doi: 10.1534/genetics.104.032821.

Beissinger, Timothy M., Li Wang, Kate Crosby, Arun Durvasula, Matthew B. Hufford, and Jeffrey Ross-
Ibarra. 2016. 'Recent Demography Drives Changes in Linked Selection across the Maize Genome'.
Nature Plants 2(7):1–7. doi: 10.1038/NPLANTS.2016.84.

Buffalo, Vince. 2021. 'Quantifying the Relationship between Genetic Diversity and Population Size
Suggests Natural Selection Cannot Explain Lewontin's Paradox'. *ELife* 10:1–30. doi:
10.7554/eLife.67509.

Campos, José Luis, and Brian Charlesworth. 2019. 'The Effects on Neutral Variability of Recurrent
Selective Sweeps and Background Selection'. *Genetics* 212(1):287–303. doi:
10.1534/genetics.119.301951.

Charlesworth, B., M. T. Morgan, and D. Charlesworth. 1993. 'The Effect of Deleterious Mutations on
Neutral Molecular Variation'. *Genetics* 134(4):1289–1303.

Charlesworth, Brian. 2009. 'Effective Population Size and Patterns of Molecular Evolution and Variation'.
Nature Reviews Genetics 10(3):195–205. doi: 10.1038/nrg2526.

Charlesworth, Brian. 2012. 'The Effects of Deleterious Mutations on Evolution at Linked Sites'. *Genetics*
190(1):5–22. doi: 10.1534/genetics.111.134288.

Comeron, Josep M. 2014. 'Background Selection as Baseline for Nucleotide Variation across the
Drosophila Genome'. *PLoS Genetics* 10(6):e1004434. doi: 10.1371/journal.pgen.1004434.

Comeron, Josep M., and Martin Kreitman. 2002. 'Population, Evolutionary and Genomic Consequences
of Interference Selection'. *Genetics* 161(1):389–410. doi: 10.1093/genetics/161.1.389.

Corbett-Detig, Russell B., Daniel L. Hartl, and Timothy B. Sackton. 2015. 'Natural Selection Constrains
Neutral Diversity across A Wide Range of Species'. *PLoS Biology* 13(4):1–25. doi:
10.1371/journal.pbio.1002112.

Cutter, Asher D., and Bret A. Payseur. 2013. 'Genomic Signatures of Selection at Linked Sites: Unifying
the Disparity among Species'. *Nature Reviews Genetics* 14(4):262–74. doi:
10.1038/nrg3425.Genomic.

Elyashiv, Eyal, Shmuel Sattath, Tina T. Hu, Alon Strutsovsky, Graham McVicker, Peter Andolfatto,
Graham Coop, and Guy Sella. 2016. 'A Genomic Map of the Effects of Linked Selection in
Drosophila'. *PLoS Genetics* 12(8):e1006130. doi: 10.1371/journal.pgen.1006130.

Ewens, Warren. 2004. *Mathematical Population Genetics*. 2nd ed. New York: Springer US.

Ewing, Gregory B., and Jeffrey D. Jensen. 2016. 'The Consequences of Not Accounting for Background
Selection in Demographic Inference'. *Molecular Ecology* 25(1):135–41. doi: 10.1111/mec.13390.

Garcia, Jesse A., and Kirk E. Lohmueller. 2021. 'Negative Linkage Disequilibrium between Amino Acid
Changing Variants Reveals Interference among Deleterious Mutations in the Human Genome'.
PLoS Genetics 17(7):1–25. doi: 10.1371/journal.pgen.1009676.

Gilbert, Kimberly J., Fanny Pouyet, Laurent Excoffier, and Stephan Peischl. 2020. 'Transition from
Background Selection to Associative Overdominance Promotes Diversity in Regions of Low
Recombination'. *Current Biology* 30(1):101-107.e3. doi: 10.1016/j.cub.2019.11.063.

Gillespie, John H. 2001. 'Is the Population Size of a Species Relevant To Its Evolution?' *Evolution*
55(11):2161. doi: 10.1554/0014-3820(2001)055[2161:itpsoa]2.0.co;2.

Good, Benjamin H. 2022. 'Linkage Disequilibrium between Rare Mutations'. *Genetics* 220(4). doi:
10.1093/genetics/iyac004.

Good, Benjamin H., Aleksandra M. Walczak, Richard A. Neher, and Michael M. Desai. 2014. 'Genetic
Diversity in the Interference Selection Limit'. *PLoS Genetics* 10(3). doi:
10.1371/journal.pgen.1004222.

Goyal, Sidhartha, Daniel J. Balick, Elizabeth R. Jerison, Richard A. Neher, Boris I. Shraiman, and Michael
M. Desai. 2012. 'Dynamic Mutation–Selection Balance as an Evolutionary Attractor'. *Genetics*
191:1309–19. doi: 10.1534/genetics.112.141291.

Haag-Liautard, Cathy, Mark Dorris, Xulio Maside, Steven Macaskill, Daniel L. Halligan, Brian
Charlesworth, and Peter D. Keightley. 2007. 'Direct Estimation of per Nucleotide and Genomic
Deleterious Mutation Rates in Drosophila'. *Nature* 445(7123):82–85. doi: 10.1038/nature05388.

Haller, Benjamin C., and Philipp W. Messer. 2017. 'SLiM 2: Flexible, Interactive Forward Genetic
Simulations'. *Molecular Biology and Evolution* 34(1):230–40. doi: 10.1093/molbev/msw211.

Hill, W., and A. Robertson. 1966. 'The Effect of Linkage on Limits to Artificial Selection.' *Genetical*
Research 8(3):269–94. doi: 10.1017/S001667230800949X.

Hoggart, Clive J., Marc Chadeau-Hyam, Taane G. Clark, Riccardo Lampariello, John C. Whittaker, Maria
De Iorio, and David J. Balding. 2007. 'Sequence-Level Population Simulations over Large Genomic
Regions'. *Genetics* 177(3):1725–31. doi: 10.1534/genetics.106.069088.

Hudson, Richard R., and Norman L. Kaplan. 1994. 'Gene Trees with Background Selection'. Pp. 140–53 in
Non-neutral Evolution: Theories and Molecular Data, edited by B. Golding. Boston, MA: Springer
US.

Hudson, Richard R., and Norman L. Kaplan. 1995. 'Deleterious Background Selection with
Recombination'. *Genetics* 141(4):1605–17.

Johri, Parul, Brian Charlesworth, and Jeffrey D. Jensen. 2020. 'Toward an Evolutionarily Appropriate Null
Model⍰': *Genetics* 215(May):173–92.

Johri, Parul, Kellen Riall, Hannes Becher, Laurent Excoffier, Brian Charlesworth, and Jeffrey D. Jensen.
2021. 'The Impact of Purifying and Background Selection on the Inference of Population History:
Problems and Prospects'. *Molecular Biology and Evolution* 38(7):2986–3003. doi:
10.1093/molbev/msab050.

Kaiser, Vera B., and Brian Charlesworth. 2009. 'The Effects of Deleterious Mutations on Evolution in
Non-Recombining Genomes'. *Trends in Genetics* 25(1):9–12.

Kaplan, Norman L., Richard R. Hudson, and Charles H. Langle. 1989. 'The "Hitchhiking Effect" Revisited'.
Genetics 123(4):887–99.

Kelleher, Jerome, Alison M. Etheridge, and Gilean McVean. 2016. 'Efficient Coalescent Simulation and
Genealogical Analysis for Large Sample Sizes'. *PLoS Computational Biology* 12(5):e1004842. doi:
10.1371/journal.pcbi.1004842.

Kelleher, Jerome, Kevin R. Thornton, Jaime Ashander, and Peter L. Ralph. 2018. 'Efficient Pedigree
Recording for Fast Population Genetics Simulation'. *PLoS Computational Biology* 14(11):e1006581.

Kern, Andrew D., and Matthew W. Hahn. 2018. 'The Neutral Theory in Light of Natural Selection'.
Molecular Biology and Evolution 35(6):1366–71. doi: 10.1093/molbev/msy092.

Kim, Bernard, Christian Huber, and Kirk Lohmueller. 2017. 'Inference of the Distribution of Selection
Coefficients for New Nonsynonymous Mutations Using Large Samples'. *Genetics* 206:345–61. doi:
10.1534/genetics.116.197145/-/DC1.1.

Kim, Yuseob, and Wolfgang Stephan. 2000. 'Joint Effects of Genetic Hitchhiking and Background
Selection on Neural Variation'. *Genetics* 155(3):1415–27. doi: 10.1093/genetics/155.3.1415.

Kimura, Motoo. 1968. 'Evolutionary Rate at the Molecular Level'. *Nature* 217:624–26.

Kimura, Motoo. 1969. 'The Number of Heterozygous Nucleotide Sites Maintained in a Finite Population
Due to Steady Flux of Mutations.' *Genetics* 61(4):893–903.

King, Jack Lester, and Thomas H. Jukes. 1969. 'Non-Darwinian Evolution'. *Science* 164(3881):788–98.

Kondrashov, Alexey S. 1995. 'Contamination of the Genome by Very Slightly Deleterious Mutations: Why
Have We Not Died 100 Times Over?' *Journal of Theoretical Biology* 175(4):583–94. doi:
10.1006/jtbi.1995.0167.

Kondrashov, Alexey S. 2017. *Crumbling Genome: The Impact of Deleterious Mutations on Humans*.
Hoboken, NJ: John Wiley & Sons, Inc.

Kulathinal, Rob J., Sarah M. Bennett, Courtney L. Fitzpatrick, and Mohamed A. F. Noor. 2008. 'Fine-Scale
Mapping of Recombination Rate in Drosophila Refines Its Correlation to Diversity and Divergence'.
Proceedings of the National Academy of Sciences of the United States of America 105(29):10051–
56. doi: 10.1073/pnas.0801848105.

Lee, Yuh Chwen G. 2022. 'Synergistic Epistasis of the Deleterious Effects of Transposable Elements'.
Genetics 220(2). doi: 10.1093/genetics/iyab211.

Lesecque, Yann, Peter D. Keightley, and Adam Eyre-Walker. 2012. 'A Resolution of the Mutation Load
Paradox in Humans'. *Genetics* 191(4):1321–30. doi: 10.1534/genetics.112.140343.

Lohmueller, Kirk E., Anders Albrechtsen, Yingrui Li, Su Yeon Kim, Thorfinn Korneliussen, Geng Tian,
Emilia Huerta-sanchez, Alison F. Feder, and Niels Grarup. 2011. 'Natural Selection Affects Multiple
Aspects of Genetic Variation at Putatively Neutral Sites across the Human Genome'. *PLoS Genetics*
7(10):e1002326. doi: 10.1371/journal.pgen.1002326.

Masel, Joanna. 2011. 'Genetic Drift'. *Current Biology* 21(20):R837–38. doi: 10.1016/j.cub.2011.08.007.

Maynard Smith, John, and John Haigh. 1974. 'The Hitch-Hiking Effect of a Favourable Gene'. *Genetics
Research* 23(1):23–35. doi: 10.1017/S0016672308009579.

McEvoy, Brian P., Joseph E. Powell, Michael E. Goddard, and Peter M. Visscher. 2011. 'Human
Population Dispersal "Out of Africa" Estimated from Linkage Disequilibrium and Allele Frequencies
of SNPs'. *Genome Research* 21(6):821–29. doi: 10.1101/gr.119636.110.

McGee, Ryan Seamus, Olivia Kosterlitz, Artem Kaznatcheev, Benjamin Kerr, and Carl T. Bergstrom. 2022.
'The Cost of Information Acquisition by Natural Selection'. *Biorxiv* 2022.07.02. doi:
10.1101/2022.07.02.498577.

McVicker, Graham, David Gordon, Colleen Davis, and Phil Green. 2009. 'Widespread Genomic Signatures
of Natural Selection in Hominid Evolution'. *PLoS Genetics* 5(5):e1000471. doi:
10.1371/journal.pgen.1000471.

Murphy, David, Eyal Elyashiv, Guy Amster, and Guy Sella. 2022. 'Broad-Scale Variation in Human Genetic
Diversity Levels Is Predicted by Purifying Selection on Coding and Non-Coding Elements'. *ELife* 11.
doi: 10.7554/eLife.76065.

Nordborg, Magnus, Brian Charlesworth, and Deborah Charlesworth. 1996. 'The Effect of Recombination
on Background Selection'. *Genetics Research* 67(2):159–74.

Ohta, Tomoko. 1973. 'Slightly Deleterious Mutant Substitutions in Evolution'. *Nature* 246(5428):96–98.

Pardo-Manuel De Villena, Fernando, and Carmen Sapienza. 2001. 'Recombination Is Proportional to the
Number of Chromosome Arms in Mammals'. *Mammalian Genome* 12(4):318–22. doi:
10.1007/s003350020005.

Popovic, Iva, Lucie Bergeron, Yves-Marie Bozec, Ann-Marie Waldvogel, Samantha Howitt, Katarina
Damjanovic, Frances Patel, Maria G. Cabrera, Gert Worheide, Sven Uthicke, and Cynthia Riginos.
2023. 'High Germline Mutation Rates but Not Extreme Population Size Outbreaks Influence Genetic
Diversity in Crown-of-Thorns Sea Stars'. *BioRxiv* 2023.06.28. doi: 10.1101/2023.06.28.546961.

Potapova, Nadezhda, and Alexey S. Kondrashov. 2023. 'Positive Association between Alleles at
Selectively Neutral Loci'. *Biorxiv* 1–12. doi: 10.1101/2023.06.14.544962.

Pouyet, Fanny, Simon Aeschbacher, Alexandre Thiéry, and Laurent Excoffier. 2018. 'Background
Selection and Biased Gene Conversion Affect More than 95% of the Human Genome and Bias
Demographic Inferences'. *ELife* 7:1–21. doi: 10.7554/eLife.36317.

Ragsdale, Aaron P., and Simon Gravel. 2020. 'Unbiased Estimation of Linkage Disequilibrium from
Unphased Data'. *Molecular Biology and Evolution* 37(3):923–32. doi: 10.1093/molbev/msz265.

Santiago, Enrique, and Armando Caballero. 2016. 'Joint Prediction of the Effective Population Size and
the Rate of Fixation of Deleterious Mutations'. *Genetics* 204(3):1267–79. doi:
10.1534/genetics.116.188250.

Sohail, Mashaal, Olga A. Vakhrusheva, Jae Hoon Sul, Sara L. Pulit, Laurent C. Francioli, Leonard H. Van
Den Berg, Jan H. Veldink, Paul I. W. De Bakker, Georgii A. Bazykin, Alexey S. Kondrashov, and
Shamil R. Sunyaev. 2017. 'Negative Selection in Humans and Fruit Flies Involves Synergistic
Epistasis'. *Science* 356(6337):539–42. doi: 10.1126/science.aah5238.

Takahata, N. 1993. 'Allelic Genealogy and Human Evolution'. *Molecular Biology and Evolution* 10(1):2–
22. doi: 10.1093/oxfordjournals.molbev.a039995.

Tenesa, Albert, Pau Navarro, Ben J. Hayes, David L. Duffy, Geraldine M. Clarke, Mike E. Goddard, Peter
M. Visscher, Human Genetics Unit, Western General Hospital, Edinburgh Eh, United Kingdom, and
Evolutionary Biology. 2007. 'Recent Human Effective Population Size Estimated from Linkage
Disequilibrium'. *Cold Spring Harbor Laboratory Press Human* 2:520–26. doi: 10.1101/gr.6023607.8.

Thornton, Kevin R. 2014. 'A C++ Template Library for Efficient Forward-Time Population Genetic
Simulation of Large Populations'. *Genetics* 198(1):157–66. doi: 10.1534/genetics.114.165019.

Thornton, Kevin R. 2019. 'Polygenic Adaptation to an Environmental Shift: Temporal Dynamics of
Variation under Gaussian Stabilizing Selection and Additive Effects on a Single Trait'. *Genetics*
213(December):1513–30. doi: 10.1101/505750.

Torres, Raul, Markus G. Stetter, Ryan D. Hernandez, and Jeffrey Ross-Ibarra. 2020. 'The Temporal
Dynamics of Background Selection in Nonequilibrium Populations'. *Genetics* 214(4):1019–30.

Torres, Raul, Zachary A. Szpiech, and Ryan D. Hernandez. 2018. 'Human Demographic History Has
Amplified the Effects of Background Selection across the Genome'. *PLoS Genetics* 14(6):1–27. doi:
10.1371/journal.pgen.1007387.

Uricchio, Lawrence H., and Ryan D. Hernandez. 2014. 'Robust Forward Simulations of Recurrent
497   Hitchhiking'. *Genetics* 197(1):221–36. doi: 10.1534/genetics.113.156935.

498   Wang, Yiguan, and Darren J. Obbard. 2023. 'Experimental Estimates of Germline Mutation Rate in
499   Eukaryotes: A Phylogenetic Meta-Analysis'. *Evolution Letters* 7(4):216–26. doi:
500   10.1093/evlett/qrad027.

501