An M protein coiled coil unfurls and exposes its hydrophobic core to capture LL-37

Piotr Kolesiński[1], Kuei-Chen Wang[1], Yujiro Hirose[2], Victor Nizet[1], Partho Ghosh[1,*]

[1]Department of Chemistry & Biochemistry, University of California, San Diego, La Jolla, CA

92093,

[2]Division of Host-Microbe Systems and Therapeutics, Department of Pediatrics, University of

California, San Diego, La Jolla, CA 92093

*To Whom Correspondence Should be Addressed: pghosh@ucsd.edu

ABSTRACT

Surface-associated, coiled-coil M proteins of *Streptococcus pyogenes* (Strep A) disable human

immunity through interaction with select proteins. However, coiled coils lack features typical of

protein-protein interaction sites, and it is therefore challenging to understand how M proteins

achieve specific binding, for example, with the human antimicrobial peptide LL-37, leading to its

neutralization. The crystal structure of a complex of LL-37 with M87 protein, an antigenic M

protein variant from a strain that is an emerging threat, revealed a novel interaction mode. The

M87 coiled coil unfurled and asymmetrically exposed its hydrophobic core to capture LL-37. A

single LL-37 molecule bound M87 in the crystal, but in solution recruited additional LL-37

molecules, consistent with a 'protein trap' neutralization mechanism. The interaction mode

visualized crystallographically was verified to contribute significantly to LL-37 resistance in an

M87 Strep A strain, and was identified to be conserved in a number of other M protein types that

are prevalent in human populations. Our results provide specific detail for therapeutic inhibition

of LL-37 neutralization by M proteins.

INTRODUCTION

M proteins are the major surface-localized virulence factor of the widespread and potentially deadly bacterial pathogen *Streptococcus pyogenes* (Group A *Streptococcus* or Strep A) (1). One of the primary functions of M proteins is to enable Strep A to evade human innate and adaptive immune responses. This is brought about by interaction of M proteins with select human proteins. M proteins are antigenically sequence variable, with over 220 different types having been identified (2). Despite this variation, the primary sequences of M proteins generally have a propensity to form dimeric, α-helical coiled coils, as verified by direct experimental evidence (3-5). This propensity is easily distinguishable by the presence of heptad repeats (6), in which amino acids in the *a* and *d* positions of the heptad are usually small and hydrophobic, and form the hydrophobic core of the coiled coil. In contrast to the usually complex topography of globular proteins, demarcated by pockets and cavities that enable specific protein-protein interactions, the simple fibrillar structure of M protein coiled coils raises the question of how M proteins achieve specific binding with their human targets.

A particular challenge lies in understanding how M proteins specifically bind and thereby neutralize the human antimicrobial peptide LL-37. This is because LL-37 also has a simple topography: LL-37 consists of a short amphipathic α-helix. LL-37 is a member of the cathelicidin antimicrobial peptide family, and constitutes a major host immune defense against Strep A as well as other pathogens (7-9). The 37-amino acid peptide is proteolytically generated from the precursor protein hCAP-18, which is produced by neutrophils, macrophages, mast cells, and keratinocytes along with other epithelial cell types (7, 10, 11). Like other amphipathic α-helical antimicrobial peptides, LL-37 functions by inserting into and lysing bacterial plasma membranes (12-14). Notably, M1 protein was shown to neutralize the antimicrobial activity of LL-37 by sequestering it in a 'protein trap' on the Strep A surface and away from its target of action, the bacterial membrane (8, 15). Not only does M1 protein attached to bacterial surface have this neutralization capacity, but so too does M1 protein released in soluble form from the Strep A

surface, as occurs during infection (8, 16). M1 protein also binds the LL-37 precursor hCAP-18,

and consequently prevents the proteolytic generation of LL-37 (8).

To determine the mechanism of LL-37 binding and neutralization by M proteins, we

pursued co-crystallization. While M1 protein proved recalcitrant to co-crystallization, several new

M protein types that bind and neutralize LL-37 were identified. LL-37 was co-crystallized with

M87 protein, which is an M type from strain that is an emerging health threat (17). The structure

revealed a remarkable and novel mode of interaction for a coiled coil, in which a portion of the

M87 protein coiled coil unfurled and exposed its hydrophobic core for interaction with LL-37.

The LL-37-binding motif visualized in M87 protein was identified in a number of other M protein

types, many of which are prevalent in human populations. Experimental evidence showed that

some of these other M types bound and neutralized LL-37 similarly to M87 protein. Our results

provide specific detail for inhibiting the interaction of M proteins with LL-37.

RESULTS

Structure of the M87/LL-37 Complex

To determine the mechanism of binding of LL-37 by M proteins, we pursued co-

crystallization beginning with M1 protein but were unable to obtain co-crystals despite various

attempts. A number of other M protein constructs (M4, M5, M22, M28, M44, M58, M77, M87,

and M89), which were available in the laboratory for other purposes, were tried next. These

consisted of the N-terminal 100 amino acids (denoted below with superscripted "N100") of these

M proteins. M58^{N100} and M87^{N100} bound LL-37, as did the M1^{HB} fragment (Figure 1A), while the

others did not bind LL-37 above background level (Figure 1 — figure supplement 1). The level

of LL-37 binding by M58^{N100}, M87^{N100}, M1^{HB} was approximately similar. As a point of

comparison, the K_D for the M1/LL-37 complex is ~1 µM (9).

To our knowledge, M58 and M87 proteins are the first M proteins besides M1 protein

identified to bind LL-37. To determine whether this binding was functionally significant, M58^{N100}

or M87[N100] was exogenously added to an M1 Strep A strain in which *emm1* had been deleted

(*Δemm1*) (15). As previously shown (8), *Δemm1* was sensitive to the antimicrobial action of LL-

37, and exogenously added M1[HB] increased resistance against LL-37, although in this particular

experiment the effect was not statistically significant (Figure 1B). Likewise, M58[N100] or M87[N100]

exogenously added to *Δemm1* increased the LL-37 MIC, and in this case, these effects were

statistically significant and M87[N100] had the greatest effect (Figure 1B).

87   Various fragments of M58 and M87 were tried in co-crystallization trials with LL-37.

Success was had with a version of M87 protein (amino acids, aa 68-105) that had a portion of

the canonical coiled-coil protein GCN4 (aa 250-278) (8, 15, 18) fused in register to its N-

terminus. GCN4 fusion was pursued to stabilize coiled-coil formation in the M87 protein

fragment, a technique that has proven successful in the crystallization of a number of coiled-coil

proteins (19, 20). The structure of the GCN4-M87/LL-37 complex was determined through

molecular replacement to 2.1 Å resolution limit (Table I, Figure 2, Figure 2 — figure supplement

1A). The GCN4-M87 fusion protein was verified to bind and neutralize LL-37, as gauged by co-

precipitation and MIC assays, respectively (Figures 1B, C). GCN4-M87 bound LL-37 and led to

a statistically significant increase in the LL-37 MIC, to an even greater extent than observed for

M87[N100].

98   The structure revealed a single LL-37 molecule bound to a dimer of GCN4-M87 (Figure

2A). Two nearly identical 1:2 complexes occupied the asymmetric unit of the crystal (Figure 2 —

figure supplement 1B, RMSD 0.70 Å). Almost the entire length of LL-37, which was

predominantly in α-helical conformation, was visible in the crystal structure. GCN4-M87 was

likewise in α-helical conformation throughout, but strikingly, only the GCN4 portion formed a

coiled coil while the M87 α-helices were unfurled and asymmetrically disposed. This was best

appreciated by comparing the structures of the complexed and free form of GCN4-M87. The

structure of the latter was determined to 2.4 Å resolution limit (Table I, Figure 2 — figure

supplement 1C). Free GCN4-M87 formed a dimeric coiled coil throughout (Figure 2B), indicating

that the unfurling of the M87 α-helices was unique to the bound form (Figure 2C). The greatest

extent of unfurling occurred at Phe91, in which the distance between Cα atoms was 8.7 Å in the

free form but 15.6 Å in the bound form. Only the M87 portion contacted LL-37, and therefore we

will refer only to M87 rather than GCN4-M87 hereafter. The two M87 α-helices and the LL-37 α-

helix together formed a parallel, three-helix bundle. One of the M87 α-helices made more

contact with LL-37 (1016 Å^2 total buried surface area, average of the two complexes in the

asymmetric unit of the crystal) (Figure 3, dark blue) while the other made less but still

substantial contact (491 Å^2 average total buried surface area) (Figure 3, cyan); to differentiate

between the two M87 α-helices, the one making greater contact will be denoted M87α1 and the

other M87α2.

Interface dominated by hydrophobic interactions

Overall, hydrophobic interactions dominated the interaction site. Most notably, a number

of hydrophobic M87 amino acids that occupied the core *a* or *d* positions and contacted each

other in the free form, as is typical of coiled coils, were exposed in the bound form and instead

contacted the hydrophobic face of the amphipathic LL-37 α-helix. Nearly five helical turns of LL-

37 (Phe5-Val21) and M87 protein (Leu74-Trp92) engaged one another. Near the N-terminus of

LL-37, a consecutive pair of phenylalanines, Phe5 and Phe6, were surrounded by hydrophobic

a and *d* position amino acids of M87 — Leu74 (*a*, usual position in heptad), Ala77 (*d*), and

Tyr81 (*a*) from M87α along with Ala77 from M87α2 (Figure 3A). The two LL-37 Phe's and M87

Tyr81 formed a series of π-stacks. Two helical turns later in LL-37 was Ile13 which engaged in a

ring of isoleucines, with M87 contributing Ile84 (*d*) from each of its helices (Figure 3B). A further

helical turn later in LL-37 was Phe17 which packed against a pair of M87 leucines, Leu88 (*a*),

one from each of the M87 helices, and π-stacked against M87α2 Phe91 (*d*) (Figure 3C). Lastly,

one more helical turn further in LL-37 were Ile20, which was surrounded by M87α1 Phe91 (*d*)

and Trp92; Val21, which packed against M87α2 Phe91; and Arg23 whose aliphatic side chain

atoms packed against M87α1 Trp92 (Figure 3D). These hydrophobic contacts were

supplemented by a sparingly few polar ones, which occurred all within a small polar break in the

hydrophobic face of LL-37, near its N-terminus: LL-37 Ser9 and Lys12 formed a hydrogen bond

and salt bridge, respectively, with M87α1 Glu85 (Figure 3E), and in one of the two complexes,

LL-37 Lys12 also formed a salt bridge with M87α1 Glu89.

Importance of Hydrophobic Interactions

The role of the contacts observed structurally were evaluated through site-directed

mutagenesis of M87 protein. To ensure that our structural observations were not limited to a

fragment of M87 protein, these experiments were carried out with intact M87 protein. Intact His-

tagged M87 protein and LL-37 were incubated at 37 °C, and their interaction was determined

through a Ni^{2+}-NTA agarose bead co-precipitation assay. Dual alanine-substitutions of M87

Tyr81 (*a*) and Ile84 (*d*), which interacted with LL-37 Phe5, Phe6, and Ile13 (Figures 3A and B),

significantly decreased LL-37 interaction (Figures 4A and B). M87 Y81A/I84A was verified

through circular dichroism (CD) to have a similar structure to wild-type M87 protein at 37 °C

(Figure 4 — supplement 1A). Additionally, M87 Y81A/I84A had a temperature-induced unfolding

profile similar to that of wild-type M87 protein (Figure 4 — supplement 1B), also as monitored by

CD. These results indicated that alanine-substitutions of Tyr81 and Ile84 affected LL-37 binding

directly rather than indirectly through compromised structure, stability, or both. Alanine-

substitution of M87 Leu88 (*a*) and Phe91 (*d*), which interacted hydrophobically with LL-37

Phe17, Ile20, and Val21 (Figures 3C and D), likewise markedly decreased LL-37 interaction

(Figures 4A and B). The secondary structure and stability of M87 L88A/F91A resembled that of

wild-type M87 protein as well (Figure 4 — figure supplements 1A and B). A substantial amount

of the surface area of M87 Trp92 was buried by contact with LL-37, but surprisingly, substitution

of this amino acid with alanine increased LL-37 interaction (Figures 3C and D). M87 Trp92 was

adjacent to LL-37 Arg23 (Figure 3D), and thus we asked whether Arg-substitution of M87 Trp92

would interfere with LL-37 binding. Indeed, M87 W92R was almost entirely deficient in LL-37

interaction (Figures 4A and B), while showing no changes in secondary structure or stability

(Figure 4 — figure supplements 1A and B).

The only M87 amino acid seen to make polar contacts in both complexes in the

asymmetric unit of the crystal was Glu85 (Figure 3E). Ala-substitution of M87 Glu85 had little

effect on LL-37 binding (Figures 4A and B), suggesting it did not contribute to affinity. To ask

whether it instead contributed to specificity, we asked whether Arg-substitution of this amino

acid would decrease interaction with LL-37, as M87 Glu85 was adjacent to LL-37 Lys12.

Consistent with our structural observations and a role in specificity, M87 E85R had significantly

decreased interaction with LL-37 (Figures 4A and B). This decrease was a direct effect, as M87

E85R had greater α-helical content than wild-type M87 protein and similar stability (Figure 4 —

figure supplements 1A and B).

These results validated the structural observations regarding the mode of interaction

between M87 protein and LL-37, and indicated the importance of the M87 *a* and *d* heptad

position amino acids to the interaction. They further indicated that the polar contact conferred by

M87 Glu85 conferred specificity rather than binding affinity.

Stoichiometry of Interactions

For an LL-37 neutralization mechanism that involves an M protein trap (8), the 1:2

LL-37:M87 stoichiometry was puzzling. LL-37 forms variably sized oligomers in solution (21-23),

and thus we undertook solution phase studies of the complex through size-exclusion

chromatography (SEC) coupled to multiangle light scattering (MALS). The molecular weight of

intact M87 protein alone was 73.6 ± 1.2 kDa (calc. 72.4 kDa) (Figure 4C, black). A 10-fold

excess of LL-37 to the intact M87 dimer was added and the sample was applied to SEC-MALS

almost immediately after mixing. The complex had a mass of 80.4 ± 0.6 kDa, which

corresponded to an M87 protein dimer bound to one or two molecules of LL-37 (calc. 4.5 kDa)

(Figure 4C, blue). Notably, after one hour of incubation of LL-37 with M87 protein, the mass of

the complex was 97.1 ± 0.6 kDa (Figure 4C, red), which corresponded to an M87 protein dimer

bound to five or six molecules of LL-37. Incubation of up to four hours also resulted in the same

increased mass, indicating self-limiting growth of the complex (Figure 4 — figure supplement 2).

These results suggested that the single LL-37 molecule bound to an M87 dimer could recruit

four or five additional LL-37 molecules.

emm87 **Confers Resistance to LL-37**

We asked whether the mode of interaction visualized through crystallography was

applicable to M87 protein in its native conformation on the Strep A surface. An isogenic

Δemm87 strain was constructed, and complemented with a plasmid expressing either wild-type

emm87 or *emm87* containing the E85R substitution, the latter having greatly diminished LL-37

binding in solution (Figure 4A). The wild-type M87 strain was resistant to LL-37 (Figure 4D), and

indeed had a greater LL-37 MIC than the M1 strain (Figure 1C). Deletion of *emm87* led to

significantly increased sensitivity to LL-37 (Figure 4D). Notably, the *Δemm87* strain

complemented with wild-type M87 was restored to the LL-37 resistance of the wild-type parental

M87 strain, while complemented with M87 E85R remained sensitive to LL-37, similar to the level

seen for the uncomplemented *Δemm87* strain. These LL-37 susceptibility results provide

physiological validation for our structural observations.

Conservation of M87 motif in other M types

The LL-37 binding motif visualized in M87 protein was identified in the sequence variable

N-terminal regions of 14 other M protein types (Figure 5A). The motif consisted of two

consecutive ideal heptads, that is, with *a* and *d* positions occupied by canonical hydrophobic

amino acids (24-26), including a strictly conserved Tyr at the *a* position of the first heptad.

These hydrophobic amino acids in M87 protein (Tyr81, Ile84, Leu88, and Phe91) were shown

above to be crucial to interaction with LL-37 (Figures 4A and B). Along with these, the motif had

a hydrogen bond acceptor at the *e* position of the first heptad, which in M87 protein (Glu85)

provided specificity by contacting LL-37 Lys12 (Figure 3). In addition, a positively charged

amino acid was excluded from the *e* position of the second heptad, which in M87 protein was

Trp92 and proximal to LL-37 Arg23 (Figure 3). Preceding these two ideal heptads was a nearly

ideal heptad, with the *a* position occupied by canonical amino acids (i.e., Leu or Tyr) while the *d*

position was tolerant of less than ideal amino acids (i.e., Ala77 as in M87, and also Gln) (25,

26).

219   Among the M proteins identified to have this motif were M25, M58, and M68 proteins.

We showed above that M58^{N100} bound LL-37 (Figure 1A). Similar fragments consisting of the N-

terminal 100 amino acids were constructed for M25 and M68. M68^{N100} bound LL-37 as did

M25^{N100} (Figure 5B). To test whether the LL-37 binding mode observed for M87 protein was

conserved in these M proteins, Arg-substitutions of the equivalent of M87 E85 were constructed.

As the E85R substitution had been evaluated only in intact M87 protein, it was introduced into

the M87^{N100} fragment, which resulted in significantly decreased LL-37 binding as well (Figures

5B and D). Arg-substitution of the equivalent Glu in M25^{N100} (E87) and M68^{N100} (E79) also led to

significantly decreased LL-37 binding (Figures 5B and D). While Arg-substitution of M58^{N100}

(E97) decreased LL-37 binding, this effect was not statistically significant. Modeling suggested

that an Arg at this position in M58 was capable of salt bridging with M58 D94, thereby

attenuating its LL-37 repulsion. In the case of M25 and M58 proteins, we noticed that the amino

acid preceding the strictly conserved Tyr was Gly, which is a helix-breaking amino acid.

Therefore, we substituted the Gly in M25 and M58 proteins with Asp, which is the equivalent

amino acid in M87 (D80), and found that M25^{N100} G82D and M58^{N100} G92D had significantly

higher levels of LL-37 binding as compared to their wild-type counterparts (Figures 5C and D).

235   These results are consistent with M25, M58, and M68 binding LL-37 in the same mode

as observed for M87 protein, and support the hypothesis that the M87 protein LL-37 binding

motif is conserved in at least fourteen other M proteins, which belong to the E2 or E3 cluster of

M proteins (27).

DISCUSSION

We sought to understand how M proteins achieve specific binding of LL-37, which is

essential to neutralization of this human antimicrobial peptide by the M1 strain of Strep A and

likely by other strains as well (7, 8, 15). This question was challenging as the fibrillar structure of

M proteins excludes features commonly present in globular proteins that enable specific

binding, such as pockets or cavities. Adding to the challenge, the structure of LL-37 lacks

complexity, consisting simply of an amphipathic α-helix. We discovered a remarkable mode of

interaction of M87 protein with LL-37 through structure determination. The two α-helices of M87

protein in the free state formed a coiled coil, with amino acids in the core *a* and *d* positions

engaged in 'knobs-into-holes' packing. In the bound state, the two helices did not form a coiled

coil, which was instead unfurled with the helices asymmetrically disposed. Intriguingly, analysis

of the ideality of 'knobs-into-holes' packing of M87 in the free state showed that A77 and Y81

were not ideally packed (28). This suggested that the segment containing A77 and Y81 may

initiate unfurling to bind LL-37. Most significantly, in the bound state, hydrophobic amino acids at

the M87 *a* and *d* positions were exposed and formed a continuous patch that contacted the

hydrophobic face of the LL-37 α-helix. These hydrophobic contacts dominated the interaction

interface and were shown to be essential by Ala-substitution mutagenesis. Recently, a coiled-

coil dimer that also binds an α-helical partner was reported (29), but in that complex coiled-coil

character is maintained, unlike the loss of coiled-coil character observed for the M87/LL-37

complex. In addition, while coiled-coil asymmetry coupled to cognate partner binding has been

noted in some proteins (30, 31), this has taken the form of a helical stagger with coiled-coil

character maintained. To our knowledge, the unfurling and asymmetric exposure of consecutive

a and *d* position amino acids to form a continuous interaction site is a novel binding mechanism

for a coiled-coil protein.

The LL-37 binding site in M87 protein was formed by two ideal heptads preceded by a

nearly ideal heptad (Ala and Gln being tolerated at the *a* position). M proteins identified or

predicted to bind LL-37 in the same mode also had at least three such consecutive repeats, with

the first heptad being ideal in most cases (Figures 6A and Figure 6 — figure supplement 1A). In

contrast, M proteins identified not to bind LL-37 lacked consecutive ideal or near ideal heptads.

Instead ideal heptads occurred as isolated singletons in the midst of non-ideal heptads. Notably,

M1 protein has only an isolated ideal repeat (Figure 6A), indicating that there are additional

mechanisms for binding LL-37, different from those observed for M87 protein. In general, the

occurrence of two or more consecutive ideal heptads is rare in M protein variable regions

(Figure 6 — figure supplement 1), whose sequences frequently have coiled-coil destabilizing

amino acids at the *a* and *d* positions (e.g., Glu or Lys) (3, 32). In the case of the M1 B-repeats

region, non-ideal heptads have been shown to be functionally essential for creating protein

dynamics, which are required for binding fibrinogen in a "capture-and-collapse" mechanism (33).

Dynamics in the consecutive ideal repeats of M87 and related M proteins are likely to be much

lower in magnitude than in the non-ideal M1 B-repeats, but nevertheless, it appears that enough

breathing motion exists even in these ideal coiled coils for the infiltration of LL-37 and disruption

of coiled-coil structure. The loose 'knobs-into-holes' packing of Y81 in the free form of M87

along with the conservation of the Tyr in M proteins shown or implicated to bind LL-37 suggests

that this may be a general site for initiating unfurling.

The single molecule of LL-37 bound to M87 protein acted as a nucleator for the

recruitment of additional LL-37 molecules. One or two molecules of LL-37 molecule bound to

M87 protein in solution at an initial time point, but grew to five or six molecules over time, as

evidenced by SEC-MALS analysis. This is consistent with observations that LL-37 oligomerizes

in solution (21-23). Plausible mechanisms for the recruitment of additional LL-37 molecules are

suggested by crystal structures of LL-37 alone. In one crystal structure, the polar faces of two

LL-37 molecules contact one another in anti-parallel orientation (13), and in another, the

hydrophobic faces do likewise (12). Thus, it is possible that the single LL-37 molecule bound

through its hydrophobic face to M87 protein is able to recruit a second molecule of LL-37

through polar face-polar face interactions. This second LL-37 molecule would then be able to

recruit a third molecule of LL-37 through hydrophobic face-hydrophobic face interactions (Figure

6B). While this sort of growth is not self-limiting, we observed that the LL-37/M87 complex was

self-limiting at five or six molecules of LL-37 per M87 dimer, and thus the specific details of LL-

37/LL-37 interactions are likely to differ. The excess of LL-37 bound by M87 protein is consistent

with a 'protein trap' neutralization mechanism (8).

The physiological relevance of the mode of LL-37 binding by M87 protein was

established through deletion and complementation experiments. A deletion of *emm87* resulted

in significant sensitivity to LL-37, as seen in the decrease in the LL-37 MIC. Resistance against

LL-37 was restored to the *Δemm87* strain by a plasmid encoding wild-type *emm87* but not

*emm*87 (E85R). *In vitro* studies using purified proteins showed that M87 E85R does not bind LL-

37, and structural studies provided an explanation for this — M87 E85 is positioned next to LL-

37 Lys12.

M87 Strep A strains are prevalent in human populations and are an emerging cause of

human clinical disease (17, 34, 35). The genomes of almost all M87 strain isolates contain a

recombination event that increases the expression of Strep A toxin genes (NADase and

streptolysin O) (17). Importantly, the same recombination event is present in the M1 strain that

caused a global pandemic starting in the 1980's and an M89 strain that is responsible for an

ongoing epidemic (36). A few isolates of the M82 strain also carry this recombination event (17).

The M82 strain, along with a number of the other M types identified to have the M87 motif, are

prevalent in human populations, including M25, M68, M90, and M103 (17, 34, 35). These

observations provide motivation for pursuing therapeutic inhibition of LL-37 binding by M

proteins, and our results provide specific detail to achieve this end.

Materials and methods

Bacterial strains and culture conditions. *Streptococcus pyogenes* (Strep A) strains M87

20161436 (NCBI SRA accession: SAMN07154152) and its isogenic Δ*emm87* strain (37), and

M1T1 5448 and its isogenic Δ*emm1* strain (15) were used. *S. pyogenes* was grown as standing

cultures in Todd-Hewitt broth in ambient atmosphere at 37 ˚C. *Escherichia coli* was cultured in

Lysogeny Broth at 37 ˚C with agitation. For selection and maintenance of strains, antibiotics

were added to the medium at the following concentrations: erythromycin 500 µg/mL for *E. coli*

and 2 µg/mL for *S. pyogenes*; chloramphenicol, 2 µg/mL for *S. pyogenes*.

DNA manipulation. Coding sequences for M proteins were cloned into a modified pET28b

vector (Novagen) that contained sequences encoding an N-terminal His$_6$-tag followed by a

PreScission protease cleavage site (5). Amino acid substitutions and deletions were introduced

into pET28b vectors with the QuickChange II Site-Directed mutagenesis kit (Stratagene),

according to the manufacturer's directions, and into pM87 E85R (37), which was used for

expression in *S. pyogenes*, with the Phusion Site Directed Mutagenesis Kit (ThermoScientific,

Waltham, MA, USA). The coding sequence for GCN4 250-278 was subcloned from

Saccharomyces cerevisiae, and was fused to M87 68-105 through strand overlap extension

PCR. An M87 protein-expressing vector (pM87) was constructed by insertion of *emm87* into

pDCerm (37).

Peptides and Proteins. LL-37, which was chemically synthesized and lyophilized as a fluoride

salt (Genscript, 95% purity), was solubilized at 5 mg/ml in sterile deionized water for MIC assays

or 100 mM NaCl, 20 mM HEPES-NaOH, pH 7.5 (HS) for other experiments.

Expression and purification of M proteins constructs was carried out as previously

described (3, 5), except for the following minor modifications. After PreScission protease

digestion, GCN4-M87 was subjected to gel filtration chromatography using a Superdex 200 (GE

Healthcare) column equilibrated with HS buffer. For formation of the GCN4-M87/LL-37 complex,

GCN4-M87 (2 mg/ml) was mixed with a three-fold molar excess of LL-37 (3 mg/ml), both in HS

buffer, and the complex was purified by gel filtration chromatography using a Superdex 200

column that had been equilibrated with 100 mM NaCl, 20 mM MES-NaOH, pH 6.5. Intact wild-

type and mutant M87 proteins, following Ni^{2+}-NTA agarose bead purification, were subjected to

gel filtration chromatography on a Superdex 200 column that had been equilibrated with HS

buffer. For CD measurements, the His-tag on intact wild-type and mutant M87 proteins was

removed by PreScission protease digestion, and the cleaved product was further purified by

reverse Ni^{2+}-NTA chromatography. Protein concentrations of M proteins were determined by

measuring A_{280} with the sample in 6 M guanidine hydrochloride, 20 mM Tris, pH 8.5, and using

a calculated molar 280 nm extinction coefficient. The concentration of LL-37 and GCN4-M87/LL-

37 complex was measured using the Bradford assay (Bio-Rad) with BSA as a standard.

Co-precipitation assay. One and half nmol of His_6-tagged M protein constructs (2-10 µl) were

added to 50 µl of Ni^{2+}-NTA agarose beads that had been pre-equilibrated with HS buffer and

incubated with gentle agitation for 10 min at RT. Beads were centrifuged (3,000 x g, 30 s, RT)

and the supernatant was removed. Six nmol of LL-37 in 150 µl of HS buffer was added to the

beads and incubated with gentle agitation for 30 min at 37 °C. The beads were washed three

times each with 1 ml HS buffer containing 5 mM imidazole, pH 8.0. For the washes, the resin

was mixed with the wash solution by gentle agitation, incubated for 1 min at RT and then

centrifuged (3,000 x g, 30 s, RT). Bound proteins were eluted with 50 µl HS containing 400 mM

imidazole, pH 8.0. Protein samples were resolved by SDS-PAGE and stained with InstaBlue

(APExBIO). The intensity of gel bands was quantified with ImageJ (38) and confirmed to be

within the linear range of detection. As the intensity on each gel depended on the length of time

of staining and destaining, a normalization factor was used. For this, the ratio of the LL-37 band

intensity to wild-type M protein band intensity from a particular gel lane was determined from

three or more independent gels, and the mean of these ratios (mean of LL-37/M_{WT}) was used as

the normalization factor. Accordingly, for each data point (i.e., gel lane), the ratio of the LL-37

band intensity to M protein (wild-type or mutant) band intensity was quantified, and divided by

the normalization factor (mean of LL-37/M_{WT}).

Minimal Inhibitory Concentration. *S. pyogenes* that had been grown overnight were

inoculated into Todd-Hewitt broth at 1:100 dilution and grown at 37 °C to an OD_{600} of 0.4. The

culture was diluted to an OD_{600} of 0.1, and 5 µl (~10^5 CFU) was mixed into 100 µl of RPMI 1640

medium with glutamine, which contained 0, 2, 4, 8, 12, 18, or 32 µM LL-37. In some

experiments, the medium also contained 10 µM $M1^{HB}$, $M58^{N100}$, or $M87^{N100}$ protein. *S. pyogenes*

were grown in individual wells of a 96-well plate for 24 h at 37 °C. *S. pyogenes* viability was

assessed at this time point by the color of the RPMI medium, where yellow indicated bacterial

growth and red no bacterial growth. The MIC was defined as the LL-37 concentration at which

no growth was detectable at 24 h.

Molecular mass determination. Intact His-tagged M87 protein (2.5 mg/ml) alone or mixed with

LL-37 (1.6 mg/ml; 10-fold molar excess over M87 dimer) in HS (100 mM NaCl, 20 mM HEPES-

NaOH, pH 7.5) was centrifuged (10 min, 20,000 x *g*, 20 °C) to remove aggregates. Samples

(100 µl) were then either immediately applied to a Superdex 200 10/300 column that had been

pre-equilibrated in HS, or incubated 1-4 h at RT before application to the column. Samples

eluting from the column were monitored with a light scattering detector (DAWN HELEOS II,

Wyatt Technology, Santa Barbara, CA, USA) and a differential refractometer (Optilab T-rEX;

Wyatt Technology). Data processing and molecular mass calculation were performed with

ASTRA software (Wyatt Technology).

Crystallization and data collection. Crystallization trials were carried out at 293 K using the

hanging drop vapor diffusion method. The GCN4-M87/LL-37 complex was concentrated by

ultrafiltration using a 3,500 MWCO membrane (Millipore; 4,500 x g, 30 min, 15 °C) to 8 mg/ml.

GCN4-M87 alone was concentrated to 10 mg/ml by ultrafiltration through 3,500 MWCO

membrane (Millipore; 4,500 x g, 30 min, 4 °C) The complex was brought to RT before

introduction into crystallization drops to overcome its low solubility at 4 °C.

The GCN4-M87/LL-37 complex (0.8 µl) was mixed in a 1:1 ratio with 5% (v/v)

acetonitrile, 0.1 M MES-NaOH, pH 6.5. Microclusters of plates (ca. 20 x 20 x 5 µm^3) that had

grown after 2-4 days were crushed, diluted 125-fold with the precipitant solution, and

centrifuged (2,000 x g, 30 s, RT). The supernatant was collected and used as a seed stock.

GCN4-M87/LL-37 (0.9 µl) was mixed with 0.9 µl of 10% (v/v) acetonitrile, 0.1 M MES-NaOH, pH

6.5 and 0.2 µl of the seed stock. Clusters of diffraction-sized crystals (50–150 µm in each

dimension) which were obtained after 3-7 days were crushed once again and used for a

subsequent round of seeding, carried out in the same manner as the first round. These two

rounds of seeding yielded single crystals (200 x 200 x 10 µm^3) that were cryo-preserved by

three serial transfers to the precipitant solution supplemented with 10, 20, and 30% ethylene

glycol, respectively, and flash-cooled in liquid N$_2$.

GCN4-M87 (0.8 µl) was mixed in a 1:1 ratio with 0.6 M monobasic ammonium

phosphate, 0.1 M Tris-HCl, pH 8.0,. Single crystals (300 x 150 x 150 µm^3), which were obtained

after 4-7 days, were cryo-protected in the precipitant solution supplemented with 30% ethylene

glycol and flash cooled in liquid N$_2$ prior to data collection.

Diffraction data were collected at SSRL (beamline 9-2) at 0.979 Å, integrated with

Mosflm (GCN4-M87/LL-37) (39) or DIALS (GCN4-M87) (40) and scaled with Aimless (41)

(Table I). Because of the highly anisotropic nature of both datasets, the resolution cutoffs for

both were determined using anisotropic $CC_{1/2}$.

Structure determination and refinement. Phases for the GCN4-M87/LL-37 complex and free

GCN4-M87 were determined by molecular replacement using Phaser (42). The search model

was generated from the coiled-coil dimer structure of GCN4 fused to the coiled-coil dimer

structure of striated muscle α-tropomyosin (PDB 1KQL) using Sculptor (43) with default settings.

Extensive model modification and building were performed with Coot and guided by the

inspection of σA-weighted 2mFo-DFc and mFo-DFc omit maps (Figure 2 — figure supplement

1).

LL-37 in the GCN4-M87/LL-37 complex was manually modelled into well-defined

difference electron density that was visible after a few rounds of refinement of the search model.

The asymmetric unit of the GCN4-M87/LL-37 crystal contained two heteromeric assemblies,

each composed of two chains of GCN4-M87 and one of LL-37. Refinement was performed

using Refine from the Phenix suite (44) with default settings. At the final stages of refinement,

TLS parameters were applied. TLS groups were applied as follows: chain A; chain B; chain C

aa 41-55 and 56-93' chain D aa 38-55 56-104; chain E; and chain F. Side chains with no

corresponding electron density were truncated to Cβ. Interaction interfaces between M87 and

LL-37 were analyzed with PISA (45).

The GCN4-M87 dimer was modeled into continuous electron density. The asymmetric

unit of the GCN4-M87 crystal contained a single coiled-coil dimer. Refinement of GCN4-M87

was performed using Refine with default settings in addition to the application of two-fold NCS

restraints and TLS parameters. Each chain constituted a single TLS group. Three and six amino

acids at N- and C-termini, respectively, of chain A and a single N-terminal amino acid of chain B

lacked electron density and were not modeled.

Molecular figures were generated with PyMol (http://pymol.sourceforge.net).

Structures have been deposited in the PDB: 7SAY for GCN4-M87/LL-37 and 7SAF for

GCN4-M87.

CD spectroscopy. CD spectra were measured on an Aviv 215 Circular Dichroism

Spectrometer using a quartz cell with 1 mm path length. Protein samples were ~0.125-0.250

mg/ml in 5 mM sodium phosphate, pH 7.9. Wavelength spectra were recorded in a range of

190-260 nm at 37 °C at 0.5 nm intervals with a 0.5 s averaging time per data point. Melting

curves were determined at 222 nm between 20-75 °C with 1 °C increments and 30 s

equilibration time for each temperature point. Two independent experiments were carried out for

each sample, and the data were averaged and presented as a mean molar residue ellipticity.

Identification of M87 LL-37-binding motif in other M proteins. The sequence of the N-

terminal 250 amino acids of mature M87 was aligned against the sequence of the N-terminal

250 amino acids of the mature form of 179 M proteins using Clustal Omega (46). Alignments

were manually curated for the presence of a Tyr (or Phe, although none were found) occupying

the position equivalent to M87 Tyr81, the occurrence of hydrophobic amino acids at the *d, a,*

and *d* positions following the Tyr, and a hydrogen bond acceptor at the *e* position following the

Tyr.

Acknowledgements

We thank S. Rees for his help with crystallographic data collection. This work was supported by

NIH 1R21AI144901 (P.G.).

Competing Interest Statement

The authors declare that they have no competing interests.

References

1.　　Ghosh P. Variation, Indispensability, and Masking in the M protein. Trends Microbiol.
2018;26(2):132-44. PMC5794653. PubMed 28867148

2.　　McMillan DJ, Dreze PA, Vu T, Bessen DE, Guglielmini J, Steer AC, Carapetis JR, Van
Melderen L, Sriprakash KS, Smeesters PR. Updated model of group A *Streptococcus* M
proteins based on a comprehensive worldwide study. Clin Microbiol Infect.
2013;19(5):E222-9. PubMed 23464795

3.　　McNamara C, Zinkernagel AS, Macheboeuf P, Cunningham MW, Nizet V, Ghosh P.
Coiled-coil irregularities and instabilities in group A *Streptococcus* M1 are required for
virulence. Science. 2008;319(5868):1405-8. PMC2288698. PubMed 18323455

4.　　Macheboeuf P, Buffalo C, Fu CY, Zinkernagel AS, Cole JN, Johnson JE, Nizet V, Ghosh
P. Streptococcal M1 protein constructs a pathological host fibrinogen network. Nature.
2011;472(7341):64-8. PMC3268815. PubMed 21475196

5.　　Buffalo CZ, Bahn-Suh AJ, Hirakis SP, Biswas T, Amaro RE, Nizet V, Ghosh P.
Conserved patterns hidden within group A Streptococcus M protein hypervariability
recognize human C4b-binding protein. Nat Microbiol. 2016;1:16155. PMC5014329.
PubMed 27595425

6.　　Manjula BN, Fischetti VA. Tropomyosin-like seven residue periodicity in three
immunologically distinct streptococal M proteins and its implications for the
antiphagocytic property of the molecule. J Exp Med. 1980;151(3):695-708.
PMC2185799. PubMed 6987328

7.　　Nizet V, Ohtake T, Lauth X, Trowbridge J, Rudisill J, Dorschner RA, Pestonjamasp V,
Piraino J, Huttner K, Gallo RL. Innate antimicrobial peptide protects the skin from
invasive bacterial infection. Nature. 2001;414(6862):454-7. PubMed 11719807

8.　　LaRock CN, Dohrmann S, Todd J, Corriden R, Olson J, Johannssen T, Lepenies B,
Gallo RL, Ghosh P, Nizet V. Group A Streptococcal M1 Protein Sequesters Cathelicidin
to Evade Innate Immune Killing. Cell Host Microbe. 2015;18(4):471-7. PMC4636435.
PubMed 26468750

9.　　LaRock CN, Nizet V. Cationic antimicrobial peptide resistance mechanisms of
streptococcal pathogens. Biochim Biophys Acta. 2015;1848(11 Pt B):3047-54.
PMC4539303. PubMed 25701232

10.　Zaiou M, Nizet V, Gallo RL. Antimicrobial and protease inhibitory functions of the human
cathelicidin (hCAP18/LL-37) prosequence. J Invest Dermatol. 2003;120(5):810-6.
PubMed 12713586

11.　Wong JH, Ye XJ, Ng TB. Cathelicidins: peptides with antimicrobial, immunomodulatory,
anti-inflammatory, angiogenic, anticancer and procancer activities. Curr Protein Pept Sci.
2013;14(6):504-14. PubMed 23968350

12.   Sancho-Vaello E, Francois P, Bonetti EJ, Lilie H, Finger S, Gil-Ortiz F, Gil-Carton D,
506   Zeth K. Structural remodeling and oligomerization of human cathelicidin on membranes
507   suggest fibril-like structures as active species. Sci Rep. 2017;7(1):15371. PMC5684418.
508   PubMed 29133814

509   13.   Sancho-Vaello E, Gil-Carton D, Francois P, Bonetti EJ, Kreir M, Pothula KR,
510   Kleinekathofer U, Zeth K. The structure of the antimicrobial human cathelicidin LL-37
511   shows oligomerization and channel formation in the presence of membrane mimics. Sci
512   Rep. 2020;10(1):17356. PMC7562864. PubMed 33060695

513   14.   Schneider VA, Coorens M, Ordonez SR, Tjeerdsma-van Bokhoven JL, Posthuma G, van
514   Dijk A, Haagsman HP, Veldhuizen EJ. Imaging the antimicrobial mechanism(s) of
515   cathelicidin-2. Sci Rep. 2016;6:32948. PMC5021996. PubMed 27624595

516   15.   Lauth X, von Kockritz-Blickwede M, McNamara CW, Myskowski S, Zinkernagel AS,
517   Beall B, Ghosh P, Gallo RL, Nizet V. M1 protein allows Group A streptococcal survival in
518   phagocyte extracellular traps through cathelicidin inhibition. J Innate Immun.
519   2009;1(3):202-14. PubMed 20375578

520   16.   Herwald H, Cramer H, Morgelin M, Russell W, Sollenberg U, Norrby-Teglund A,
521   Flodgaard H, Lindbom L, Bjorck L. M protein, a classical bacterial virulence determinant,
522   forms complexes with fibrinogen that induce vascular leakage. Cell. 2004;116(3):367-79.
523   PubMed 15016372

524   17.   Turner CE, Holden MTG, Blane B, Horner C, Peacock SJ, Sriskandan S. The
525   Emergence of Successful Streptococcus pyogenes Lineages through Convergent
526   Pathways of Capsule Loss and Recombination Directing High Toxin Expression. mBio.
527   2019;10(6):e02521-19. PMC6904876. PubMed 31822586

528   18.   O'Shea EK, Klemm JD, Kim PS, Alber T. X-ray structure of the GCN4 leucine zipper, a
529   two-stranded, parallel coiled coil. Science. 1991;254(5031):539-44. PubMed 1948029

530   19.   De N, Navarro MV, Raghavan RV, Sondermann H. Determinants for the activation and
531   autoinhibition of the diguanylate cyclase response regulator WspR. J Mol Biol.
532   2009;393(3):619-33. PMC2760619. PubMed 19695263

533   20.   Li Y, Mui S, Brown JH, Strand J, Reshetnikova L, Tobacman LS, Cohen C. The crystal
534   structure of the C-terminal fragment of striated-muscle alpha-tropomyosin reveals a key
535   troponin T recognition site. Proc Natl Acad Sci U S A. 2002;99(11):7378-83.
536   PMC124239. PubMed 12032291

537   21.   Johansson J, Gudmundsson GH, Rottenberg ME, Berndt KD, Agerberth B.
538   Conformation-dependent antibacterial activity of the naturally occurring human peptide
539   LL-37. J Biol Chem. 1998;273(6):3718-24. PubMed 9452503

540   22.   Oren Z, Lerman JC, Gudmundsson GH, Agerberth B, Shai Y. Structure and organization
541   of the human antimicrobial peptide LL-37 in phospholipid membranes: relevance to the
542   molecular basis for its non-cell-selective activity. Biochem J. 1999;341 (Pt 3):501-13.
543   PMC1220385. PubMed 10417311

23. Xhindoli D, Pacor S, Benincasa M, Scocchi M, Gennaro R, Tossi A. The human
cathelicidin LL-37--A pore-forming antibacterial peptide and host-cell modulator. Biochim
Biophys Acta. 2016;1858(3):546-66. PubMed 26556394

24. Wagschal K, Tripet B, Lavigne P, Mant C, Hodges RS. The role of position a in
determining the stability and oligomerization state of alpha-helical coiled coils: 20 amino
acid stability coefficients in the hydrophobic core of proteins. Protein Sci.
1999;8(11):2312-29. PMC2144206. PubMed 10595534

25. Tripet B, Wagschal K, Lavigne P, Mant CT, Hodges RS. Effects of side-chain
characteristics on stability and oligomerization state of a de novo-designed model coiled-
coil: 20 amino acid substitutions in position "d". J Mol Biol. 2000;300(2):377-402.
PubMed 10873472

26. Kwok SC, Hodges RS. Stabilizing and destabilizing clusters in the hydrophobic core of
long two-stranded alpha-helical coiled-coils. J Biol Chem. 2004;279(20):21576-88.
PubMed 15020585

27. Sanderson-Smith M, De Oliveira DM, Guglielmini J, McMillan DJ, Vu T, Holien JK,
Henningham A, Steer AC, Bessen DE, Dale JB, Curtis N, Beall BW, Walker MJ, Parker
MW, Carapetis JR, Van Melderen L, Sriprakash KS, Smeesters PR, Group MPS. A
systematic and functional classification of *Streptococcus pyogenes* that serves as a new
tool for molecular typing and vaccine development. J Infect Dis. 2014;210(8):1325-38.
PubMed 24799598

28. Kumar P, Woolfson DN. Socket2: A Program for Locating, Visualising, and Analysing
Coiled-coil Interfaces in Protein Structures. Bioinformatics. 2021. PMC8652024. PubMed
34498035

29. Dimitrova-Paternoga L, Jagtap PKA, Cyrklaff A, Vaishali, Lapouge K, Sehr P, Perez K,
Heber S, Low C, Hennig J, Ephrussi A. Molecular basis of mRNA transport by a kinesin-
1-atypical tropomyosin complex. Genes Dev. 2021;35(13-14):976-91. PMC8247599.
PubMed 34140355

30. Sato Y, Fukai S, Ishitani R, Nureki O. Crystal structure of the Sec4p·Sec2p complex in
the nucleotide exchanging intermediate state. Proceedings of the National Academy of
Sciences. 2007;104(20):8305-10.

31. Noell CR, Loh JY, Debler EW, Loftus KM, Cui H, Russ BB, Zhang K, Goyal P, Solmaz
SR. Role of Coiled-Coil Registry Shifts in the Activation of Human Bicaudal D2 for
Dynein Recruitment upon Cargo Binding. J Phys Chem Lett. 2019;10(15):4362-7.
PMC7243283. PubMed 31306018

32. Nilson BH, Frick IM, Akesson P, Forsen S, Bjorck L, Akerstrom B, Wikstrom M. Structure
and stability of protein H and the M1 protein from Streptococcus pyogenes. Implications
for other surface proteins of gram-positive bacteria. Biochemistry. 1995;34(41):13688-
98. PubMed 7577960

33. Stewart CM, Buffalo CZ, Valderrama JA, Henningham A, Cole JN, Nizet V, Ghosh P.
Coiled-coil destabilizing residues in the group A Streptococcus M1 protein are required

for functional interaction. Proc Natl Acad Sci U S A. 2016;113(34):9515-20.
PMC5003295. PubMed 27512043

34.   Steer AC, Law I, Matatolu L, Beall BW, Carapetis JR. Global *emm* type distribution of
group A *streptococci*: systematic review and implications for vaccine development.
Lancet Infect Dis. 2009;9(10):611-6. PubMed 19778763

35.   Li Y, Rivers J, Mathis S, Li Z, Velusamy S, Nanduri SA, Van Beneden CA, Snippes-
Vagnone P, Lynfield R, McGee L, Chochua S, Metcalf BJ, Beall B. Genomic
Surveillance of Streptococcus pyogenes Strains Causing Invasive Disease, United
States, 2016-2017. Front Microbiol. 2020;11:1547. PMC7396493. PubMed 32849323

36.   Zhu L, Olsen RJ, Nasser W, Beres SB, Vuopio J, Kristinsson KG, Gottfredsson M, Porter
AR, DeLeo FR, Musser JM. A molecular trigger for intercontinental epidemics of group A
Streptococcus. J Clin Invest. 2015;125(9):3545-59. PMC4588293. PubMed 26258415

37.   Hirose Y, Kolesinski P, Hiraoka M, Uchiyama S, Zurich RH, Kumaraswamy M, Bjanes E,
Ghosh P, Kawabata S, Nizet V. Contributions of *Stretpococcus pyogenes* M87 protein to
innate immune resistance and virulence. Microb Pathog. 2022;In Review.

38.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nat Methods. 2012;9(7):671-5. PubMed 22930834

39.   Battye TGG, Kontogiannis L, Johnson O, Powell HR, Leslie AGW. iMOSFLM: a new
graphical interface for diffraction-image processing with MOSFLM. Acta
Crystallographica Section D. 2011;67(4):271-81.

40.   Winter G, Waterman DG, Parkhurst JM, Brewster AS, Gildea RJ, Gerstel M, Fuentes-
Montero L, Vollmar M, Michels-Clark T, Young ID, Sauter NK, Evans G. DIALS:
implementation and evaluation of a new integration package. Acta Crystallographica
Section D. 2018;74(2):85-97.

41.   Evans PR, Murshudov GN. How good are my data and what is the resolution? Acta
Crystallographica Section D. 2013;69(7):1204-14.

42.   McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. Phaser
crystallographic software. Journal of Applied Crystallography. 2007;40:658-74.

43.   Bunkoczi G, Read RJ. Improvement of molecular-replacement models with Sculptor.
Acta Crystallogr D Biol Crystallogr. 2011;67(Pt 4):303-12. PMC3069745. PubMed
21460448

44.   Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M,
Terwilliger TC, Urzhumtsev A, Zwart PH, Adams PD. Towards automated
crystallographic structure refinement with phenix.refine. Acta Crystallographica Section
D. 2012;68(4):352-67.

45.   Krissinel E, Henrick K. Inference of macromolecular assemblies from crystalline state. J
Mol Biol. 2007;372(3):774-97. PubMed 17681537

46.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
622   Remmert M, Söding J, Thompson JD, Higgins DG. Fast, scalable generation of high-
623   quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol.
624   2011;7:539. PMC3261699. PubMed 21988835

Table I. Crystallographic Data Collection and Model Refinement

	GCN4-M87/LL-37	GCN4-M87
Data Collection		
Wavelength (Å)	0.979	0.979
Resolution range (Å)	45.02 - 2.10 (2.18 - 2.10)[a]	43.48 - 2.45 (2.54 - 2.45)
Space group	$P2_1$	$P4_32_12$
Cell dimensions		
a, b, c (Å)	50.2, 57.1, 71.5	97.2, 97.2, 40.7
α, β, γ (°)	90.0, 102.7, 90.0	90.0, 90.0, 90.0
Total reflections	134,344 (9,365)	91,495 (8,197)
Unique reflections	22,092 (1,694)	7,580 (738)
Multiplicity	6.1 (5.5)	12.1 (11.1)
Completeness (%)	94.51 (89.58)	99.75 (100.00)
$I/\sigma(I)$	8.50 (2.10)	31.40 (0.78)
Wilson B-factor (Å^2)	26.42	55.23
R_{meas}[b]	0.128 (0.735)	0.404 (4.911)
$CC_{1/2}$[c]	0.996 (0.878)	0.987 (0.556)
Refinement		
Resolution range (Å)	45.02 - 2.10 (2.20 - 2.10)	43.48 - 2.45 (2.80 - 2.45)
No. of reflections (work/test set)	21,958/2,426	7,563/383
R_{work}/R_{free}[d]	0.22/0.27 (0.27/0.34)	0.26/0.27 (0.29/0.38)
No. of non-hydrogen atoms	2673	1138
macromolecules	2556	1126
ligands	16 (ethylene glycol)	2 (PO_4^{3-})
solvent	101	2
R.m.s. deviations		
bonds (Å)	0.003	0.007
angles (°)	0.46	0.95
Ramachandran plot		
favoured (%)	100.00	100.00
outliers (%)	0.00	0.00
Rotamer outliers (%)	0.00	0.00
Clashscore	3.37	3.98
Average B-factor (Å^2)	46.59	76.91
macromolecules	46.80	75.60
ligands	37.52	110.74
solvent	42.81	64.27
Number of TLS groups	8	2
PDB code	7SAY	7SAF

[a]Values in parentheses are for the highest resolution shell.

$^{b}R_{meas} = \sum_{hkl} \sqrt{\frac{n}{n-1}} \sum_{j=1}^{n} \left| I_{hkl\,j} - < I_{hkl} > \right| / \sum_{hkl} \sum_{j} I_{hkl\,j}$

$^{c}CC_{1/2}$ is the Pearson correlation coefficient between two random half datasets.

$^{d}R_{work} = \sum_{hkl} \left| F_{obs} - F_{calc} \right| / \sum_{hkl} F_{obs}$

Formulas for R_{work} and R_{free} are identical except 95% of the total number of reflections was used

to calculate R_{work} whereas the remaining 5% of reflection was used to calculate R_{free}.

FIGURE LEGENDS

Figure 1. LL-37 binding and detoxification.

A. LL-37 binding to His-tagged M1^{HB}, M58^{N100}, or M87^{N100} as determined by Ni^{2+}-NTA agarose

bead co-precipitation assay at 37 °C. The last lane contains no M protein. Bound fractions were

resolved by SDS-PAGE and visualized by Coomassie staining. The gel is representative of at

least three experiments. Figure 1 — figure supplement 1 shows LL-37 binding to other M types.

B. LL-37 MIC for wild-type Strep A *emm1* and isogenic *Δemm1* alone or supplemented with 10

µM M1^{HB}, M58^{N100}, M87^{N100}, or GCN4-M87. Data from four independent experiments are

presented with means and standard deviations. Statistical significance was calculated using

one-way ANOVA with Dunnett post-hoc test. P values are as follows: ns > 0.05, * ≤ 0.05, ** ≤

0.01, *** ≤ 0.001 and **** ≤ 0.0001."

C. LL-37 binding to His-tagged M87^{N100} or GCN-M87, determined as in panel A. The last lane

contains no M protein. The gel is representative of at least three experiments.

Figure 2. Structures of GCN4-M87/LL-37 complex and free GCN4-M87.

A. Cartoon representation of the GCN4-M87/LL-37 complex. LL-37 is in magenta. The GCN4

portion of GCN4-M87 is in gray, and the M87 portion in blue for the chain that makes more

contacts to LL-37 (M87α1), and cyan for the chain that makes fewer contacts to LL-37 (M87α2).

The N- and C-terminal amino acids for LL-37 and M87 are indicated. Figure 2 — figure

supplement 1 shows electron density and superposition of the two GCN4-M87/LL-37 complexes

in the asymmetric unit.

B. Cartoon representation of free GCN4-M87. The GCN4 portion is colored gray and the M87

portion green.

C. Superposition (based on GCN4, not depicted) of the M87 portions in bound (blue and cyan)

and free (green) states.

Figure 3. M87/LL-37 Interface.

LL-37 is in magenta, and the M87 α-helix that forms a greater number of contacts with LL-37 is in blue (M87α1) and the one that forms fewer contacts in cyan (M87α2). Shown are contacts formed with LL-37 (**A**) Phe5 and Phe6; (**B**) Ile 13; (**C**) Phe17; (**D**) Ile20, Val21, and Arg23; and (**E**) Ser9 and Lys12.

Figure 4. Evaluation of M87/LL-37 Interactions.

A. LL-37 binding by His-tagged, intact wild-type M87 or M87 Y81A/I84A, L88A/F91A, E85A, E85R, W92A, or W92R at 37 °C as determined by Ni^{2+}-NTA agarose bead co-precipitation. The last lane contains no M protein. Bound fractions were resolved by SDS-PAGE and visualized by Coomassie staining. The gel is representative of four experiments. Figure 4 — figure supplement 1 shows CD spectra and melting curves of wild-type and mutant M87 proteins.

B. Quantification of LL-37 binding to intact wild-type and mutant M87 proteins. The ratio between LL-37 and wild-type M87 protein band intensities was quantified in four independent experiments, and used to determine a mean LL-37/wt-M87 ratio. Points shown are ratios between LL-37 and (wild-type or mutant) M87 protein band intensities, normalized by the mean LL-37/wt-M87 ratio. For each sample, means and SD are shown. Statistical significance was calculated using one-way ANOVA with Dunnett post-hoc test. P values are as follows: ns > 0.05, * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001 and **** ≤ 0.0001.

C. Molecular weight determination of intact M87 alone (black), M87 and LL-37 added together and immediately analyzed (blue), or M87 and LL-37 incubated for 1 h before being analyzed (red) by size-exclusion chromatography (SEC) coupled to multiangle light scattering (MALS). Dotted lines indicate calculated molecular weights across the profile. Data are representative of three experiments. Figure 4 — figure supplement 2 shows SEC-MALS analysis of M87 protein and LL-37 incubated for 1 or 4 h.

D. LL-37 MIC for *emm87* and *Δemm87* alone or expressing wild-type M87 (pM87) or M87 E85R

pM87(E85R) from a plasmid. Data from eight independent experiments are presented as mean

± SD. The statistical test and P values are the same as described for panel B.

Figure 5. Conservation of the M87LL-37-binding motif.

A. M87 amino acids that make hydrophobic contact to LL-37 are boxed in red and those that

disrupt interaction with LL-37 through Arg-substitution are in blue boxes, as are amino acids that

are similarly positioned in other M proteins. The arrow indicates the position of M87 D80.

B. LL-37 interaction with His-tagged wild-type or mutant $M87^{N100}$, $M25^{N100}$, $M58^{N100}$, $M68^{N100}$

proteins determined by Ni^{2+}-NTA agarose bead co-precipitation at 37 °C. Bound fractions were

resolved by SDS-PAGE and visualized by Coomassie staining. Data are representative of six

experiments.

C. LL-37 interaction with His-tagged $M25^{N100}$, $M25^{N10}$ G82D, $M58^{N100}$, and $M58^{N100}$ G02D,

determined as in panel B. Data are representative of six experiments.

D. Quantification of the effect of mutagenesis on LL-37 binding. The ratio between LL-37 and

wild-type M protein (i.e., M25, M58, M68, or M87) band intensities was quantified in six

independent experiments, and used to determine a mean LL-37/wt-M ratio. Points shown are

ratios between LL-37 and mutant M protein band intensities, normalized by the corresponding

mean LL-37/wt-M ratio. For each sample, means and SD are shown. Statistical significance was

calculated using Student's t-test to compare mutant and the corresponding wild-type protein. P

values are as follows: ns > 0.05, * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001 and **** ≤ 0.0001.

Figure 6. LL-37 binding motif and higher-order complex formation.

A. Occurrence of ideal heptads (dark grey boxes; Ile, Leu, Met, Phe, Tyr, or Val at both *a* and *d*

positions) and near ideal heptads (light grey boxes; ideal amino acid at *a* position and Ala at *d*

position, with M25 being an exception with Gln at this position) within the N-terminal regions of

various M protein types. Within these boxes, yellow circles indicate Tyr at an *a* position; green

circles Asp, Asn, Glu, or Gln at an *e* position; and red circles Arg or Lys at an *e* position. Yellow

and red circles favor LL-37 binding, while the red circles disfavor LL-37 binding. Figure 6 —

figure supplement 1 shows the occurrence of ideal and near ideal heptads within the N-terminal

regions of additional M types.

B. Model of higher-order M87/LL-37 complex. M87α1 and M87α2 are in blue and cyan,

respectively, and LL-37 is in magenta. Left, LL-37 bound to M87 protein has its polar face free

to form polar contacts with a second LL-37 molecule (light gray, as seen in a structure of LL-37

alone, PDB 7PDC). Right, the light gray LL-37 has its hydrophobic face free to form hydrophobic

contacts with yet another LL-37 molecule (dark gray, as seen in another structure of LL-37

alone, PDB 5NNM). The light and dark gray LL-37 molecules do not contact M87 protein.

    FIGURE SUPPLEMENT LEGENDS

Figure 1 — figure supplement 1. LL-37 Binding.

LL-37 binding to His-tagged $M1^{HB}$, $M4^{N100}$, $M5^{N100}$, $M22^{N100}$, $M28^{N100}$, $M44^{N100}$, $M77^{N100}$, or

$M89^{N100}$ as determined by a Ni^{2+}-NTA agarose bead co-precipitation assay at 37 °C. The last

lane contains no M protein. Bound fractions were resolved by SDS-PAGE and visualized by

Coomassie staining. The gel is representative of three experiments.

Figure 2 — figure supplement 1. Structures of the GCN4-M87/LL-37 complex and free

GCN4-M87.

A. Electron density for GCN4-M87 from a simulated annealing σA-weighted 2mFo-DFc

composite omit map contoured at 1.4σ. Part of the interface between M87 (left) and LL-37

(right) is shown.

B. Superposition of the two GCN4-M87/LL-37 complexes present in the asymmetric unit of the

crystal. LL-37 is in different shades of magenta. The GCN4 portions are in shades of gray, and

the M87 portions are in blue and cyan.

C. Electron density for free GCN-M87 from a simulated annealing σA-weighted 2mFo-DFc

composite omit map contoured at 1.4σ.

Figure 4 — figure supplement 1. Structure and Stability of M87 mutant proteins.

A. CD spectra of intact wild-type and mutant M87 proteins at 37 °C. The data are an average of

two independent measurements.

B. Melting curves of intact wild-type and mutant M87 proteins, as monitored by the CD mean

molar residue ellipticity at 222 nm. The data are an average of two independent measurements.

Figure 4 — figure supplement 2. Self-limiting growth of the M87/LL-37 complex.

Molecular weights of intact M87 protein and LL-37 added together and incubated for 1 h (black)

or 4 h (blue) were determined by size-exclusion chromatography coupled to multiangle light

scattering. Dotted lines indicate calculated molecular weights across the profile.

Figure 6 — figure supplement 1. LL-37 binding motif.

Depiction as in Fig. 6A. The first grouping contains M proteins that have the LL-37 binding motif

identified in M87 protein (experimentally verified for M87, M25, M58, and M68, and predicted for

M79, M82, M90, M96, M103, M107, M144, M168, M180, M209, and M231). M1 protein, which

binds LL-37 but does not have the motif identified in M87 protein, is next, followed by M proteins

that have been shown experimentally to not bind LL-37. Lastly, eleven M proteins randomly

selected from the remaining pool of M proteins are shown (M12, M15, M18, M42, M63, M71,

M73, M101, M114, M176, and M178).

Figure 1

Figure 2

Figure 3

A Phe5 / Helical turn 1: hydrophobic / Tyr81 / Phe6 / Ala77 / Leu74 / Ala77

B Helical turn 3: hydrophobic / Ile13 / Ile84 / Ile84

C Helical turn 4: hydrophobic / Leu88 / Phe17 / Phe91 / Leu88

D Helical turn 5: hydrophobic / Trp92 / Arg23 / Phe91 / Ile20 / Val21 / Phe91

E Helical turn 2: polar / Lys12 / Glu85 / Ser9

68 / 68 / 2 / 93 / 36 / 104

Figure 4

Figure 5

Figure 6

Figure 1 — figure supplement 1

Figure 2 — figure supplement 1

Figure 4 — figure supplement 1

Figure 4 — figure supplement 2

Figure 6 — figure supplement 1