Fibromodulin Ablation Exacerbates the Severity of Acute DSS Colitis

Mariana Halasi, PhD[1]; Mor Grinstein, PhD[2]; Avner Adini, PhD[3]; Irit Adini, PhD[1,*]

1. Harvard Medical School, Department of Surgery, Center for Engineering in Medicine & Surgery, Massachusetts General Hospital, 51 Bloosm Street, Boston, MA 02114 USA
2. Harvard Medical School, Center for Regenerative Medicine, Massachusetts General Hospital, Boston, MA 02114. Email: mgrinstein@mgh.harvard.edu
3. Harvard Medical School, Department of Medicine, Boston Children's Hospital, Karp Research Building, 11th Floor 300 Longwood Ave, Boston, MA 02115, Email: avner.adini@childrens.harvard.edu

* Correspondence to: Irit Adini, Department of Surgery, Harvard Medical School, Center for Engineering in Medicine & Surgery, Massachusetts General Hospital, 51 Bloosm Street, Boston, MA 02114 USA. Phone: 617.371.4874; Fax: 617.573.9471; Email: iadini@mgh.harvard.edu

ABSTRACT

Epidemiological studies have associated pigment production with protection against certain human diseases. In contrast to African Americans, European descendants are more likely to suffer from angiogenesis-dependent and inflammatory diseases, such as wet age-related macular degeneration (ARMD) and ulcerative colitis (UC), respectively. In this study, albino mice that produced high levels of fibromodulin (FMOD) developed less severe acute colitis compared with mice lacking in FMOD as assessed by clinical symptoms and histopathological changes. In a mouse model of dextran sodium sulfate (DSS)-induced acute colitis, FMOD depletion affected the expression and localization of tight junction proteins, contributing to the destruction of the epithelial barrier. Furthermore, this study revealed a stronger inflammatory response after DSS treatment in the absence of FMOD, where FMOD depletion led to an increase in activated T cells, plasmacytoid dendritic cells (pDCs), and type I IFN production. These findings point to FMOD as a potential biomarker of disease severity in UC among light-skinned individuals of European descent.

Keywords: FMOD, colitis, tight junction, inflammation, dendritic cells, T cells

INTRODUCTION

Ulcerative colitis (UC) is an inflammatory bowel disease (IBD) that affects primarily the mucosal surface of the large intestine (colon and rectum). It is characterized by destruction of the intestinal barrier, disappearance of the intestinal crypts, and infiltration of inflammatory cells [1,2]. Epidemiological studies indicate the prevalence of UC continues to increase worldwide, particularly in developed countries. Individuals with North American and Northern European ancestry have the highest incidence and prevalence rates with 20/100,000 and 156 - 291/100,000 cases per year, respectively. Multiple factors contribute to the pathogenesis of UC, including environmental triggers, genetic predisposition, changes in intestinal microbiota, and immune system dysfunction [3,4].

UC progression is mainly affected by the dysregulation of tight and adherens junction proteins, which form the apical junction complex and maintain the epithelial barrier. Damage to these proteins results in a dysfunctional mucosal barrier and increased intestinal permeability [5]. The molecular composition of the tight junctions (TJs) includes transmembrane proteins (claudin, occludin), cytoplasmic proteins (zonula occludens-1, ZO-1), and cytoskeletal components [6]. The tight junctions serve to prevent the internalization of intestinal pathogens. Interestingly, dendritic cells (DCs), the most effective antigen-presenting cells, not only play a key role in maintaining immunological homeostasis by orchestrating innate and adaptive immune responses, but also express TJs. DCs are found in the intestinal mucosa at the interface with the external environment, where they act as sentinels for external pathogens [7-9].

Fibromodulin (FMOD), a small leucine-rich proteoglycan, plays a role in extracellular matrix (ECM) composition. FMOD was shown to activate the classical complement cascade by direct binding to C1q and by the deposition of C3b, C4b, and C9. FMOD can also silence the complement cascade by binding to factor H [10,11]. Our previous work showed that FMOD up-regulates TGF-β1 secretion [12-14]. TGF-β1 appears to enhance intestinal epithelial barrier function by inducing the production of TJ proteins, such as claudin-1, occludin, and ZO-1 [15]. Furthermore, TGF-β1 is well known for its role in maintaining DCs in an immature state [16]. During activation and migration of immature DCs to inflammatory sites, the DCs interact directly with ECM microenvironment proteins, such as FMOD, resulting in DC differentiation and maturation [17,18].

To advance our understanding of the pathogenesis of UC, we investigated the role of FMOD in a DSS-induced murine model of acute colitis, as well as in biopsy samples from healthy and diseased individuals with IBD.

RESULTS

FMOD expression is affected by UC in human subjects and mice

The expression of FMOD was assessed by immunostaining the colon biopsy samples from healthy control individuals and patients diagnosed with UC or Crohn's disease (CD). As shown in Fig.1A (left panel), in healthy individuals, FMOD was mainly detected in the brush border on the luminal cell surface of the colon. In contrast, a significant increase in FMOD expression was observed in the crypts of colon biopsy samples from UC patients (Fig.1A, right panel).

Since dextran sulfate sodium (DSS)-induced colitis is a well-established and commonly used animal model [19,20] that mirrors key aspects of human IBD, especially UC, FMOD level and localization was evaluated in animals with DSS-induced colitis. FMOD expression demonstrated the same distribution observed in the human samples (Fig.1B).

Figure 1. FMOD is upregulated and re-localized in colitis. **(A)** Colon biopsy samples from healthy individuals and patients diagnosed with UC were stained with H&E or subjected to

immunostaining for FMOD. Representative images were captured at x20 magnification. **(B)** FMOD+/+ mice were given 3% DSS in drinking water for 7 days. Representative samples were H&E stained or immunostained for FMOD. Images were obtained at x20 magnification.

Depletion of FMOD aggravated the DSS-induced clinical symptoms of colitis

To determine the role of FMOD in the development of acute colitis, age-matched transgenic FMOD knockout (FMOD-/-) male mice, along with FMOD+/+ animals, were exposed to 3% DSS for 7 days as described in Materials and Methods. The clinical signs of DSS-induced acute colitis, which include changes in body weight, bleeding, and fecal consistency, were recorded daily to monitor onset and progression of disease. Both FMOD+/+ and FMOD-/- DSS-treated experimental animals showed similar signs of acute colitis, including body weight loss (Fig. 2A) and shortening of the colon (Fig. 2B, C). However, DSS-treated FMOD-/- mice began losing weight 3 days after DSS was administered, while weight loss in FMOD+/+ animals began 5 days after DSS was administered. During the remainder of the DSS treatment, all of the treated animals showed substantial weight loss, amounting to 8.37% and 10% of their original weight in FMOD+/+ and FMOD-/- animals, respectively (Fig. 2A).

One of the hallmarks of chemically induced colitis is colon shortening. As with the weight loss findings, colon shortening was similar between the treated groups on the day of necroscopy (19.71% in FMOD+/+ and 20.29% in FMOD-/-) but was significantly shorter compared to the respective untreated controls (Fig. 2B, C). Furthermore, both DSS-treated groups were visually checked for signs of rectal bleeding. Like weight loss, bleeding onset occurred earlier in the DSS-treated FMOD-/- animals, which started bleeding on day 3, compared to day 4 in the FMOD+/+ mice (Fig. 2D). Nevertheless, rectal bleeding remained more prominent in the FMOD-/- animals over the period of DSS treatment. The disease activity index (DAI) score, (i.e., the sum of the average weight loss score, the average stool consistency score, and the average bleeding score divided by 3), as shown in Fig 2E, indicated the FMOD-/- animals showed signs of colitis earlier and developed more severe disease compared with FMOD+/+ mice.

Together, these results demonstrated that the severity and onset of clinical symptoms of colitis, including weight loss, rectal bleeding, and diarrhea, were exacerbated in the FMOD knockout animals.

Figure 2. Ablation of FMOD facilitates DSS-related clinical signs of colitis. (A-C)
FMOD+/+ (n=8) and transgenic FMOD-/- (n=10) mice were exposed to 3% DSS in drinking
water for 7 days, and disease progression was monitored every day. Body weight was recorded
daily **(A)**, gross pictures of the colons were taken **(B)**, and colon length was measured and
plotted **(C)** on the day of sacrifice. **(D)** Rectal bleeding was recorded daily and scored according
to the bleeding score criteria detailed in Materials and Methods after visually examining the anal
area of treated animals. **(E)** DAI scores consisted of the sum of the average weight loss score, the
average stool consistency score, and the average bleeding score divided by 3. Data are the
mean $\pm$ SEM from two independent experiments. Statistical analysis was performed by
unpaired, two-tailed *t* test with Welch's correction for (D) and one-way ANOVA with Tukey's
multiple comparisons test for (A and C): *$P<0.05$, **$P<0.001$, ***$P<0.0001$.

**FMOD-ablated mice exhibited severe pathological changes of the colon after DSS
treatment**

In addition to the phenotypical signs of colitis, histopathological changes in the colon
were examined to further evaluate the possible role of FMOD in DSS-induced acute colitis. For
histological H&E staining, the flushed and longitudinally cut colons were prepared using Swiss
roll technique to permit a fuller assessment of the colonic pathology compared to the limited
scope of transverse or longitudinal sections. Histological observations were carried out to assess
the structural integrity of the intestinal epithelium (Fig. 3A). Colon sections from animals in both
groups exhibited a fully intact epithelium with well-defined crypts, no edema, no leukocyte
infiltration in the mucosa or sub-mucosa, and no ulceration or erosions (Fig. 3A). In contrast, the
colon sections of DSS-treated animals showed different degrees of gut barrier compromise in the
FMOD+/+ and FMOD-/- groups. Typical histological changes, including the presence of
mononuclear inflammatory cells, disappearance of goblet cells, degradation of the epithelium,
ulceration, edema, and leukocyte infiltration in the mucosa and sub-mucosa, were noted at
different levels among the DSS-treated groups (Fig. 3A).

In addition, the colon samples were scored blindly for the following criteria: destruction
of architecture (0-4), ulceration (0-4), and edema (0-4) (Fig. 3B). The detailed quantitative
histological analysis revealed more advanced disease in the DSS-treated FMOD-/- mice with an

overall histological score average of 2.9 compared to the FMOD+/+ mice, which had an overall histological score average of 2 (Fig. 3B).

Taken together, these results suggested that although FMOD+/+ and FMOD -/- mice showed similar physical characteristics of acute colitis, they demonstrated marked differences in colonic histopathology after DSS treatment, in which the FMOD-/- animals showed higher susceptibility to DSS and developed more severe colitis. The histological analysis underscores FMOD's role in moderating the damage inflicted by DSS on the epithelium, since knockout of FMOD resulted in substantial further damage.

Figure 3. FMOD depletion facilitates DSS-induced pathological changes in the colon. (A) FMOD+/+ (n=8) and FMOD-/- (n=10) mice were given 3% DSS in drinking water for 7 days. Paraffin-embedded sections of the colon using Swiss roll technique were H&E stained. Images were obtained with x4 magnification. **(B)** Histological scores for the destruction of architecture (0-4), edema (0-4), ulceration (0-4), and overall score (0-3) of the full-length colon given in a blinded fashion are plotted. Data are represented as the mean $\pm$ SEM of 3 control and 8 treated FMOD+/+, and 4 control and 10 treated FMOD-/- animals. *P<0.05 by unpaired, two-tailed t test with Welch's correction.

FMOD ablation facilitated epithelial barrier erosion

By its nature, DSS has the capacity to compromise the structural integrity of the epithelial cell barrier, possibly due to its interference with TJ proteins [21]. To evaluate the effects of FMOD on the composition and expression of TJ proteins in DSS-induced colitis, colonic samples of untreated and DSS-treated groups were analyzed by immunofluorescence and qRT-PCR.

Our analysis focused on intracellular protein ZO-1 and transmembrane proteins occludin and claudin-1 on account of their well-established involvement in the impairment of intestinal barrier function in IBD disease models [22,23]. ZO-1 protein expression was reduced, while claudin-1 expression was substantially upregulated (Fig. 4A) and occludin expression did not show a significant change (data not shown) in the DSS-treated colon sections of FMOD-/- mice compared to treated FMOD+/+ mice. Furthermore, quantification of fluorescence intensity by ImagJ 8 showed a 40% decrease in ZO-1 protein expression and a 1.6-fold increase of claudin-1 expression in DSS-treated FMOD-/- mice compared to treated FMOD+/+ mice (Fig.4B). To account for changes in protein expression levels, we also validated these findings at the mRNA level. The results showed that the mRNA level of ZO-1 was significantly reduced, claudin-1 mRNA was substantially upregulated, while occludin mRNA expression showed no statistically significant change in the DSS-treated FMOD-/- mice (Fig. 4C).

Poritz *et. al.*[21] found that occludin levels decrease and claudin-1 levels increase in the colon of human subjects with UC, resulting in a higher claudin-1: occludin (C:O) ratio compared to healthy colon. Using the C:O ratio calculated from the respective mRNA expression levels, we

observed a 3-fold increase in disease severity in the absence of FMOD after DSS treatment (Fig. 4D).

Fig. 4

Figure 4. Knockout of FMOD amplifies the changes in expression of TJ proteins following DSS treatment. (A) Paraffin-embedded sections of DSS-treated colon Swiss rolls were stained with anti-mouse ZO-1 or claudin-1. Images were captured at x20 magnifications. **(B)** The graph shows the quantification of mean fluorescence intensity (MFI). **(C)** The transcriptional expression of TJ proteins ZO-1, claudin-1, and occludin was measured by qRT-PCR in the colon of untreated and DSS-treated FMOD+/+ and FMOD-/- mice. The graphs show quantification as fold of mRNA expression levels compared to untreated FMOD+/+ mice (n=3), mean ± SEM 3 (FMOD+/+) or 4 (FMOD-/-) control and 8 (FMOD+/+) or 10 (FMOD-/-) treated mice. **(D)** The claudin-1 : occludin ratio was calculated using the respective mRNA expression levels. Statistical analysis was performed by unpaired, two-tailed *t* test with Welch's correction: *$P<0.05$, **$P<0.01$

FMOD alleviated DSS-induced colonic inflammation

Leukocyte infiltration has been associated with the pathogenesis of human IBD. It has also been implicated in the acute phase of DSS-induced colitis [24-26]. The presence of leukocytes in the colonic tissue was assessed by immunostaining for CD45 on colon sections from DSS-treated mice. Higher expression of CD45 (Fig. 5A) was observed in DSS-treated FMOD-/- animals, suggesting inflammation becomes more robust in the absence of FMOD.

Colonic inflammation was further examined by assessing cytokine production in the colonic tissue and the immune response in mesenteric lymph nodes (mLNs), which are known to drain the colonic tissue[27]. First, the production of inflammatory mediators was measured in colon tissue samples by qRT-PCR. Markedly higher mRNA expression of immunomodulatory cytokine interferon α/β (IFN α/β) was observed in the DSS-treated FMOD-/- animals, underscoring the development of a potentially more robust inflammatory response (Fig. 5B).

To characterize the recruited and activated immune cell populations in systemic organs following DSS exposure, mLNs and spleens were harvested and analyzed by flow cytometry. A decrease in the splenic dendritic cell (DC) and T cell populations in both DSS-treated groups was detected (data not shown). Significantly more plasmacytoid dendritic cells (pDCs) with increased MHC II (major histocompatibility complex II) maturation markers on their surface were observed in the mLNs of treated FMOD-/- animals (Fig. 5C). Analysis of T cells revealed significantly more activated T cells in the DSS-treated FMOD-/- group, while increased

mobilization of FoxP3+ T reg cells was observed in the FMOD+/+ DSS-treated mice (Fig. 5D), corroborating the presence of a stronger inflammatory response in the background of FMOD KO after DSS administration.

These results suggest that in the absence of FMOD, acute inflammation, represented by activated pDCs and reduced numbers of T regulatory cells, is significantly increased.

Figure 5. FMOD moderates DSS-induced colonic inflammation. (A) Paraffin-embedded sections of the DSS-treated colon Swiss rolls were stained with anti-mouse CD45. Images were captured at x20 magnifications. (B) The transcriptional expression of IFNα/β was measured by qRT-PCR in the colon. The graph shows quantification as fold of mRNA expression levels compared to untreated FMOD+/+ mice (n=3), mean□±□SEM 3 (FMOD+/+) or 4 (FMOD-/-) control and 8 (FMOD+/+) or 10 (FMOD-/-) treated mice. *$P<0.05$ by unpaired, two-tailed *t* test with Welch's correction. (C) The mesenteric lymph nodes (mLN) of DSS-treated FMOD+/+

(n=5) and FMOD-/- (n=5) mice were analyzed by flow cytometry for the frequency (*left panel*) and the maturity (*right panel*) of pDCs. Data are shown as a dot plot (frequency) or a density plot (maturity), and numbers represent the percentage of CD11c^{int}PDCA-1^{+} pDCs or CD11c^{int}PDCA-1^{+}MHC II^{+} pDCs among the live CD45^{+} leukocytes, respectively. **(D)** Cells from the mLNs of treated FMOD+/+ and FMOD-/- mice were stained for the expression of CD25 and FoxP3 and analyzed by flow cytometry. Dot plots show CD25 and CD4 staining pre-gated on live CD45^{+}CD3^{+} cells. The histogram represents FoxP3 staining pre-gated on live CD45^{+}CD3^{+}CD4^{+}CD25^{+} cells.

DISCUSSION

Ulcerative colitis is an idiopathic IBD limited to the colon. It is a multifactorial in nature, characterized by dysregulation of the immune system, epithelial barrier defects, and changes in the intestinal microbiota. Based on our earlier findings [12], we hypothesize that FMOD, a prominent ECM protein, plays a role in inflammation and in the maintenance of the epithelial barrier. This study investigates the potential role of FMOD in the pathogenesis of UC utilizing the widely used DSS-induced acute colitis mouse model.

FMOD, a cytosolic protein with a known secretory sequence but no transmembrane domain, is characterized as a class II member of the small leucine-rich proteoglycan family [28,29]. It was first described in human articular cartilage [30]. Although it is ubiquitously expressed in all tissues among vertebrates, its expression is most abundant in the extracellular matrix (ECM) of the connective tissues, where it plays a key role in the organization of collagen fibrils [31]. In recent years, additional tissue-specific activities have been reported, such as its role in muscle development [32], cell reprogramming [33], angiogenesis, and regulation of TGFβ-1 secretion [12,34,35]. FMOD has also been associated with inflammatory diseases of the joint and anti-inflammatory responses in wound healing [36]. To date, the role of FMOD in inflammatory and autoimmune diseases has been modestly explored. Increased expression of FMOD has been observed in patients with fibrotic kidney disease [37], heart failure, and atherosclerosis [36].

Through our experiments, we observed upregulated FMOD expression in both IBD patients and animals with DSS-induced acute colitis (Fig. 1). This observation may not be too surprising, since colitis causes oxidative stress, which is well-known to stimulate the expression of FMOD [38]. Interestingly, however, deletion of FMOD in a transgenic acute colitis mouse

model did not alleviate disease burden. Rather, it intensified the onset and severity of the clinical colitis symptoms as measured by weight loss, rectal bleeding, and DAI score (Fig. 2). In addition, histopathological examination of full-length colon sections underscored the development of more severe colitis in the absence of FMOD (Fig. 3).

The epithelial barrier physically separates the host immune cells in the intestinal mucosa from the luminal microbiota. Damage to the integrity of the colonic epithelia increases mucosal permeability due to the dysregulation of TJ proteins [2,39]. Although, epithelial barrier erosion is a key feature of clinical IBD, it is still debated whether such dysfunction is a manifestation or a cause of UC [40,41]. Loss and redistribution of tight junction proteins are commonly reported in acute DSS colitis [21]. In agreement with these findings, marked differences in mRNA and protein expression of TJ proteins, ZO-1 and claudin-1, were observed between the DSS-treated groups (Fig. 4), showing more robust changes in the absence of FMOD.

Further investigation into the effect of FMOD depletion on colitis revealed the presence of a stronger inflammatory immune response in DSS-treated FMOD-/- mice (Fig. 5). Studies have shown that FMOD plays an important role in inflammatory responses through activating the classical and alternative complement pathways by directly binding to C1q and C3b [10]. The activated complement system contributes to the adaptive and cellular immune responses by interacting with toll-like receptors (TLRs), antigen-presenting cells, and immune cells such as polymorphonuclear leukocytes, DCs, B cells, as well as T lymphocytes [10,42]. Interestingly, the deposition of C1q can also inhibit molecular processes required for DC differentiation and IFN-α production by pDCs [43-45]. Our results align with these previously described findings. Specifically, the presence of FMOD in FMOD+/+ animals, probably via complement activation, induced a more protective immune response as evidenced by the higher number of Foxp3+ T reg cells. Likewise, the absence of FMOD in FMOD-/- animals was associated with increased production of type I IFN, higher numbers of pDCs, and activated T cells indicating a stronger inflammatory response (Fig. 5).

Here we provide insight into the connection between FMOD and the increased susceptibility to colitis by employing genetic and biochemical approaches. In the acute phase of DSS-induced colitis, the presence of FMOD alleviates disease symptoms by maintaining the integrity of the epithelial barrier, reducing pDC activation, and inducing regulatory T cell

responses. Future studies are warranted to examine the potential utility of tissue FMOD as a biomarker to gauge the severity of UC among light-skinned individuals of European descent.

REFERENCES

1.   Zhou, J., *et al.* BLT1 in dendritic cells promotes Th1/Th17 differentiation and its deficiency ameliorates TNBS-induced colitis. *Cell Mol Immunol* **15**, 1047-1056 (2018).
2.   Ordas, I., Eckmann, L., Talamini, M., Baumgart, D.C. & Sandborn, W.J. Ulcerative colitis. *Lancet* **380**, 1606-1619 (2012).
3.   Zou, J., Liu, C., Jiang, S., Qian, D. & Duan, J. Cross Talk between Gut Microbiota and Intestinal Mucosal Immunity in the Development of Ulcerative Colitis. *Infect Immun* **89**, e0001421 (2021).
4.   Torres, J., Mehandru, S., Colombel, J.F. & Peyrin-Biroulet, L. Crohn's disease. *Lancet* **389**, 1741-1755 (2017).
5.   Bischoff, S.C., *et al.* Intestinal permeability--a new target for disease prevention and therapy. *BMC Gastroenterol* **14**, 189 (2014).
6.   Paris, L., Tonutti, L., Vannini, C. & Bazzoni, G. Structural organization of the tight junctions. *Biochim Biophys Acta* **1778**, 646-659 (2008).
7.   Banchereau, J. & Steinman, R.M. Dendritic cells and the control of immunity. *Nature* **392**, 245-252 (1998).
8.   Rescigno, M., Granucci, F., Citterio, S., Foti, M. & Ricciardi-Castagnoli, P. Coordinated events during bacteria-induced DC maturation. *Immunol Today* **20**, 200-203 (1999).
9.   Huang, F.P., *et al.* A discrete subpopulation of dendritic cells transports apoptotic intestinal epithelial cells to T cell areas of mesenteric lymph nodes. *J Exp Med* **191**, 435-444 (2000).
10. Sjoberg, A., Onnerfjord, P., Morgelin, M., Heinegard, D. & Blom, A.M. The extracellular matrix and inflammation: fibromodulin activates the classical pathway of complement by directly binding C1q. *J Biol Chem* **280**, 32301-32308 (2005).
11. Merle, N.S., Noe, R., Halbwachs-Mecarelli, L., Fremeaux-Bacchi, V. & Roumenina, L.T. Complement System Part II: Role in Immunity. *Front Immunol* **6**, 257 (2015).
12. Adini, I., *et al.* Melanocyte-secreted fibromodulin promotes an angiogenic microenvironment. *J Clin Invest* **124**, 425-436 (2014).

13.   Adini, I.*, et al.* Melanocyte pigmentation inversely correlates with MCP-1 production and angiogenesis-inducing potential. *FASEB J* **29**, 662-670 (2015).

14.   Kontos, C.D. More than skin deep: connecting melanocyte pigmentation and angiogenic diseases. *J Clin Invest* **124**, 76-79 (2014).

15.   Howe, K.L., Reardon, C., Wang, A., Nazli, A. & McKay, D.M. Transforming growth factor-beta regulation of epithelial tight junction proteins enhances barrier function and blocks enterohemorrhagic Escherichia coli O157:H7-induced increased permeability. *Am J Pathol* **167**, 1587-1597 (2005).

16.   Kel, J.M., Girard-Madoux, M.J., Reizis, B. & Clausen, B.E. TGF-beta is required to maintain the pool of immature Langerhans cells in the epidermis. *J Immunol* **185**, 3248-3255 (2010).

17.   Schultz, H.S.*, et al.* Collagen induces maturation of human monocyte-derived dendritic cells by signaling through osteoclast-associated receptor. *J Immunol* **194**, 3169-3179 (2015).

18.   Brand, U.*, et al.* Influence of extracellular matrix proteins on the development of cultured human dendritic cells. *Eur J Immunol* **28**, 1673-1680 (1998).

19.   Hall, L.J.*, et al.* Induction and activation of adaptive immune populations during acute and chronic phases of a murine model of experimental colitis. *Dig Dis Sci* **56**, 79-89 (2011).

20.   Arimura, K.*, et al.* Crucial role of plasmacytoid dendritic cells in the development of acute colitis through the regulation of intestinal inflammation. *Mucosal Immunol* **10**, 957-970 (2017).

21.   Poritz, L.S.*, et al.* Loss of the tight junction protein ZO-1 in dextran sulfate sodium induced colitis. *J Surg Res* **140**, 12-19 (2007).

22.   Noth, R.*, et al.* Increased intestinal permeability and tight junction disruption by altered expression and localization of occludin in a murine graft versus host disease model. *BMC Gastroenterol* **11**, 109 (2011).

23.   Poritz, L.S., Harris, L.R., 3rd, Kelly, A.A. & Koltun, W.A. Increase in the tight junction protein claudin-1 in intestinal inflammation. *Dig Dis Sci* **56**, 2802-2809 (2011).

24.   te Velde, A.A.*, et al.* Comparative analysis of colonic gene expression of three experimental colitis models mimicking inflammatory bowel disease. *Inflamm Bowel Dis* **13**, 325-330 (2007).

25.   Yan, Y.*, et al.* Temporal and spatial analysis of clinical and molecular parameters in dextran sodium sulfate induced colitis. *PLoS One* **4**, e6073 (2009).

26.   Arseneau, K.O. & Cominelli, F. Targeting leukocyte trafficking for the treatment of inflammatory bowel disease. *Clin Pharmacol Ther* **97**, 22-28 (2015).

27. Houston, S.A*., et al.* The lymph nodes draining the small intestine and colon are anatomically separate and immunologically distinct. *Mucosal Immunol* **9**, 468-478 (2016).

28. Mikaelsson, E*., et al.* Fibromodulin, an extracellular matrix protein: characterization of its unique gene and protein expression in B-cell chronic lymphocytic leukemia and mantle cell lymphoma. *Blood* **105**, 4828-4835 (2005).

29. Iozzo, R.V. & Schaefer, L. Proteoglycan form and function: A comprehensive nomenclature of proteoglycans. *Matrix Biol* **42**, 11-55 (2015).

30. Heinegard, D*., et al.* Two novel matrix proteins isolated from articular cartilage show wide distributions among connective tissues. *J Biol Chem* **261**, 13866-13872 (1986).

31. Lee, Y.H. & Schiemann, W.P. Fibromodulin suppresses nuclear factor-kappaB activity by inducing the delayed degradation of IKBA via a JNK-dependent pathway coupled to fibroblast apoptosis. *J Biol Chem* **286**, 6414-6422 (2011).

32. Lee, E.J*., et al.* Fibromodulin: a master regulator of myostatin controlling progression of satellite cells through a myogenic program. *FASEB J* **30**, 2708-2719 (2016).

33. Zheng, Z*., et al.* Reprogramming of human fibroblasts into multipotent cells with a single ECM proteoglycan, fibromodulin. *Biomaterials* **33**, 5821-5831 (2012).

34. Zheng, Z*., et al.* Fibromodulin Enhances Angiogenesis during Cutaneous Wound Healing. *Plast Reconstr Surg Glob Open* **2**, e275 (2014).

35. Jan, A.T., Lee, E.J. & Choi, I. Fibromodulin: A regulatory molecule maintaining cellular architecture for normal cellular function. *Int J Biochem Cell Biol* **80**, 66-70 (2016).

36. Zeng-Brouwers, J., Pandey, S., Trebicka, J., Wygrecka, M. & Schaefer, L. Communications via the Small Leucine-rich Proteoglycans: Molecular Specificity in Inflammation and Autoimmune Diseases. *J Histochem Cytochem* **68**, 887-906 (2020).

37. Schaefer, L*., et al.* Small proteoglycans of normal adult human kidney: distinct expression patterns of decorin, biglycan, fibromodulin, and lumican. *Kidney Int* **58**, 1557-1568 (2000).

38. Mormone, E., Lu, Y., Ge, X., Fiel, M.I. & Nieto, N. Fibromodulin, an oxidative stress-sensitive proteoglycan, regulates the fibrogenic response to liver injury in mice. *Gastroenterology* **142**, 612-621 e615 (2012).

39. Perse, M. & Cerar, A. Dextran sodium sulphate colitis mouse model: traps and tricks. *J Biomed Biotechnol* **2012**, 718617 (2012).

40.   Martini, E., Krug, S.M., Siegmund, B., Neurath, M.F. & Becker, C. Mend Your Fences: The Epithelial Barrier and its Relationship With Mucosal Immunity in Inflammatory Bowel Disease. *Cell Mol Gastroenterol Hepatol* **4**, 33-46 (2017).

41.   Lee, J.Y.*, et al.* Molecular Pathophysiology of Epithelial Barrier Dysfunction in Inflammatory Bowel Diseases. *Proteomes* **6**(2018).

42.   Heeger, P.S. & Kemper, C. Novel roles of complement in T effector cell regulation. *Immunobiology* **217**, 216-224 (2012).

43.   Son, M., Santiago-Schwarz, F., Al-Abed, Y. & Diamond, B. C1q limits dendritic cell differentiation and activation by engaging LAIR-1. *Proc Natl Acad Sci U S A* **109**, E3160-3167 (2012).

44.   Lood, C.*, et al.* C1q inhibits immune complex-induced interferon-alpha production in plasmacytoid dendritic cells: a novel link between C1q deficiency and systemic lupus erythematosus pathogenesis. *Arthritis Rheum* **60**, 3081-3090 (2009).

45.   Bencze, D., Fekete, T. & Pazmandi, K. Type I Interferon Production of Plasmacytoid Dendritic Cells under Control. *Int J Mol Sci* **22**(2021).