Emergence of Heartbeat Frailty in Advanced Age: Perspectives from Life-Long EKG Recordings in Mice

Jack M Moen[1,5], Christopher H. Morrell[1,3], Ismayil Ahmet[1], Michael G. Matt[1,6], Moran Davoodi[4], Michael Petr[2,7], Shaquille Charles[1], Raphael deCabo[2], Yael Yaniv[4], and Edward G Lakatta[1]‡

[1]Laboratory of Cardiovascular Science, [2]Laboratory of Experimental Gerontology Intramural Research Program, National Institute on Aging, National Institutes of Health, Baltimore, Maryland, USA [3]Department of Mathematics and Statistics, Loyola University Maryland, Baltimore, Maryland. [4]Biomedical Engineering Faculty, Technion-IIT, Haifa, Israel.

Present addresses:

[5]Cellular and Molecular Physiology, Yale University, New Haven, Connecticut, USA.

[6]Pediatric Residency Program, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

[7]Center for Healthy Aging, University of Copenhagen, Copenhagen, Denmark

‡**Corresponding Author**:

Edward Lakatta, MD

Email: lakattae@grc.nia.nih.gov

Telephone: 1-410-558-8202

Total Word Count (8983)

Abstract 244

Intro 834   Methods 1321

Results 4270   Discussion 2314

Abstract (244 words)

SAN failure, aka sick-sinus syndrome, which features sinus bradycardia, SAN impulse pauses, and irregularity of RR interval rhythms are manifestations of SAN cell dysfunction that increases exponentially with advanced age, i.e., SAN frailty. Abnormalities in intrinsic RR interval variability may be the earliest signatures of SAN cell dysfunction leading to SAN frailty in late life. We measured RR interval variability within EKG time-series prior to and during double autonomic blockade in long-lived C57/BL6 mice at 3 month intervals from 6 months of age until the end of life.

Long-lived mice (those that achieved the median cohort lifespan of 24 months and beyond) displayed relatively minor changes in intrinsic RR interval variability prior to 21 months of age. Between 21 and 30 months of age, marked changes in intrinsic RR interval variability signatures in time, frequency, non-linear, and fragmentation domains result in a marked increase in the mean intrinsic RR interval. The effects of autonomic input partially compensated for the prolongation of the mean RR interval by impacting the age-associated deterioration in the RR interval variability signatures toward a youthful pattern. Cross-sectional analyses of other subsets of mice at ages at or beyond the median life span of our longitudinal cohort demonstrated increased non-cardiac, constitutional, whole body frailty, a decrease in energetic efficiency, and an increase in respiratory exchange ratio. In this context, we interpret the progressive increase in intrinsic RR interval variability beyond 21 months of age to be an indication of heartbeat frailty.

Introduction (834 words)

The neuro-visceral autonomic axis continually conducts impulses at ms to sec time scales to and from the heart and other viscera via parasympathetic and sympathetic nerves. Beat to beat afferent signals (both neuronal and mechanical) originating within the heart signal to other parts of the heart and back out to the spinal cord, brain stem, and higher CNS structures, which in turn, elicit autonomic reflexes via efferent input of the central nervous system to the heart [1]. This autonomic input modulates the heart's beating rate and rhythm by impacting the rate and rhythm of spontaneous action potential firing generated by mechanisms intrinsic to sinoatrial nodal (SAN) pacemaker cells. Although by convention, the number of heartbeats per minute is casually referred to as the *in vivo* heart rate (HR), analyses of successive R-R intervals within an EKG time-series detect substantial beat to beat interval variability, indicating that the sub-cellular and cell-wide mechanisms that underlie SAN impulse generation never achieve equilibrium [2, 3]

The combined influences of SAN pacemaker cell automaticity and its response to autonomic input determine the heart's beating interval variability and mean beating rate. When the heartbeats in the absence of autonomic input, intrinsic base physiologic functions of SAN cells determine beat-to-beat interval variability from which the mean beating rate is calculated. SAN failure, aka "sick-sinus syndrome", manifests as sinus bradycardia, SAN impulse pauses, and irregularity of rhythms of RR intervals within EKG time-series that result from SAN cell dysfunction and increases exponentially with advanced age often requiring electronic pacemaker implantation. Although the resting heart rate in humans changes very little beyond young adulthood, the mean **intrinsic** RR interval, measured during dual sympathetic and parasympathetic autonomic blockade, begins to decrease substantially between young adulthood and middle age [4], but has not been studied in humans of older ages. Because autonomic input to the SAN masks

age-associated intrinsic dysfunction, intrinsic RR interval variability is likely to inform on the

overall health status of the SAN in the absence of confounding effects from autonomic input, and

changes in RR interval variability may be the earliest markers of SAN dysfunction, that ultimately

progresses to SAN failure in advanced age. In short, **intrinsic** RR interval variability reflects the

status of functions intrinsic to SAN cells that regulate the rate and rhythm of impulse emergence

from the SAN.

The aging mouse manifests abnormalities in HR and heartbeat interval variability that

recapitulate many of those described in humans [5-9]. **Cross-sectional** studies that compare

average data in different mice that have survived to different ages indicate that the intrinsic

heartbeat interval variability and mean intrinsic HR are relatively constant between young

adulthood and middle age but become substantially altered later in life [7]. The increased

sympathetic input to the SAN of older mice helps to reduce intrinsic heartbeat interval variability

towards that of mice at younger ages. Other cross-sectional studies have associated changes in

SAN automaticity in older mice with whole-body frailty [8, 10]. However, these cross-sectional

studies lack **mouse-specific, longitudinal perspectives** on when and how changes in heartbeat

interval variability leads to marked SAN dysfunction in individual long-lived mice, at an age when

whole body frailty becomes manifest at older ages [11]. Longitudinal data on heart rhythm frailty

in mice (or in humans) are mainly lacking because the longitudinal study design requires repeated

measures of HRV that are implemented over a large part of the life course. The most important

measures of cardiac frailty can be extended to mice, enabling the identification of SAN longevity

markers in animals that live to or beyond the median life span.

We reasoned that changes in longitudinal measurements of intrinsic heartbeat interval

variability in a given mouse will inform on the signature of late-life functional deterioration within

and among SAN pacemaker cells. This functional deterioration leads to progressive changes in RR

interval variability that increase the mean intrinsic RR interval in that mouse. To this end, we

designed and implemented a longitudinal study that assessed heartbeat interval variability

repeatedly in mice at 3-month intervals, beginning at 6 months of age and continuing to the end of

life. We recorded EKG time-series at each age prior to (basal state) and during double autonomic

blockade (intrinsic state) throughout the entire lifespan months in a cohort of 56 C57/BL6 black

mice. This design permitted the elucidation of how the signature of mechanisms that underlie

intrinsic SAN cell pacemaker function deteriorate over the life course. This design enabled the

creation of the autonomic index, a novel set of measurements for the differences between mean

basal and mean intrinsic state parameters measured at each age in a given mouse. This elucidated

the rates at which autonomic signatures compensate for the deterioration in SAN intrinsic

signatures during advanced age as the intrinsic SAN functions progress toward frailty. The unique

nature of this experimental design enabled comparisons on the deterioration of intrinsic SAN

function and compensatory autonomic signatures of heartbeat interval variability in advanced age

to more traditional non-cardiac, whole body frailty indices, movement-based energetic efficiency,

and energy substrate utilization.

Methods (1321 words)

Electrocardiogram

C57/BL6 mice, male, 3 months old (n=58), were obtained from Charles River Laboratories Inc. (Wilmington, MA). All studies were performed in accordance with *the Guide for the Care and Use of Laboratory Animals* published by the National Institutes of Health (NIH Publication no. 85-23, revised 1996). Experimental protocols were approved by the Animal Care and Use Committee of the National Institutes on Aging (protocol #441-LCS-2016).

Under light anesthesia with isoflurane (2% in oxygen) through nosecone at 0.4 ml/min, ECG electrode needles were inserted under the skin, a standard lead II ECG were recorded using Power Lab System (AD Instruments Inc.) at a sampling rate of 1,000 Hz. A heat lamp was positioned at a constant distance (25 cm) from the mouse during recordings to prevent heat loss. After let mice adapt to these conditions for twenty minutes, ECG was recorded continuously for 50 minutes: 10 minutes prior to, and 40 minutes after intraperitoneal injection of a saline solution (400 uL/30g body weight) with atropine (0.5 mg/kg) and propranolol (1.0 mg/kg). A representative time-series of ECG RR interval recordings is illustrated in Supplemental Figure 1. ECG time-series were recorded in each mouse at three-month intervals, starting at six months of age, and continuing for the entire lifespan of each mouse.

The basal heart rate (HR) in each mouse was calculated as an average HR of 10 min recordings prior to drug and was reported as the mean basal RR interval (BRR). During the 40 min recordings after atropine and propranolol administration, HR was calculated as an average of each minute recordings, and the lowest HR was identified as the intrinsic HR (IHR). IHR is usually reached around 20 min after injection of atropine and propranolol. HR then returns to normal value around 40 min

134   after injection. IHR reported as the mean intrinsic RR interval (IRR).

EKG RR Interval Time-series Analyses

EKG time-series of RR intervals were analyzed using Physiozoo [12]. A mouse preset with

rodent T waves was used to obtain a block average. Initially one-minute averages were obtained

from the 10 minutes prior to injection and 40 minutes following injection. From the one-minute

averages a block average was made for the set of basal recordings, and recordings in the presence

of pharmacological autonomic blockade. Recordings of a single mouse were removed from the

entire dataset, being at outlier with a resting heart rate >700 BPM.

Heart rate variability analysis also employed Physiozoo. ECGs were first analyzed to

identify segments that fit strict selection criteria. This was mainly dependent on whether or not the

heart rate was stationary for a sufficiently long period of time. Stationarity was determined based

on the absence of linear trends, a high ectopic number, or a non-stable heart rate. Segments that

met these eligibility criteria were included in the subset of tracings used for heart rate variability

(for specific N numbers refer to Table S1). Generally, a 1.5-2-minute segment was selected,

containing between 512 to 1,024 intervals. Rather, larger subsets were obtained for frequency

domain analysis. The software was set up such that segments could be auto-analyzed based on

length and segmented into similar sizes. Thus, our data was reported on files clipped to 512

intervals. After this we applied an automatic ectopic removal correction [12] in which any values

outside of two standard deviations were removed.

Statistics

156   All statistical analyses were implemented in R 3.2.3 [13] using RStudio [14].

Mixed ANOVA Analyses

To test for age-associated differences in mean response, age was set as a factor under the assumption that recording intervals were approximately the same, and a mixed-effects ANOVA was conducted (Table S2).

Linear mixed effects (LME) statistical analyses

A linear mixed effect (LME) model (lmerTest) was used to evaluate the effect of time following the initial measurement at 6 months of age on basal and intrinsic RR interval parameters, and autonomic index HR and HRV parameters.

Estimating Mouse-Specific Rates of Change in RR Interval Variability

The average age trajectory is displayed using a loess smooth curve [15]. For many EKG time-series parameters, the trajectory over the age span is highly non-linear, exhibiting a sharp change around 21 months of age. This sharp change cannot be adequately modeled by a simple polynomial. Consequently, when needed, the trajectories are modeled in two parts: ages 6 – 21 months, and 21 to 30 months. To accommodate the curvature in the trajectories within each of these age-spans, a quadratic model in age is adopted, when needed. In this case, the full LME model becomes:

$y_{ij} = (\beta_0 + b_{i0}) + (\beta_1 + b_{i1})\, Age + (\beta_2 + b_{i2})\, Age^2 + \varepsilon_{ij}$

LME models contain both **fixed** and **random** terms. The β's are the regression parameters for the **fixed-effects** variables, while b_{i0}, b_{i1}, and b_{i2} are the corresponding random effects. ε_i is the usual

error term. The fixed-effects terms of LME models are the usual regression-type explanatory

variables and their parameter estimates provide estimates of **average** rates of change in the

response variable for changes in the associated variable. The **random effects** terms of LME

models, however, allow for variability in initial value and trajectories among the mice. To obtain

mouse-specific rates of change, age must now be considered as a numerical variable, and the

trajectory over the age span must be appropriately modeled. The intercept random effect (b_{i0})

allows for variability in individual mouse intercepts or starting points while the additional random

effects, b_{i1} and b_{i2}, allow for the linear and quadratic terms to vary among the animals.

In the full model, the rate of change in the response for individual mouse *i* is the derivative

of the model function with respect to age and is given by:

$Rate_i = (\beta_1 + b_{i1}) + 2 \times (\beta_2 + b_{i2}) \times Age$

where the β's and b_i's are replaced by the estimates obtained from the data. When terms are

eliminated from the model, the appropriate remaining terms from the final model are used to obtain

the mouse-specific rates of change. For many variables, the data could not accommodate the

random effect for Age^2. Consequently, in these cases, this random effect was removed from the

model.

Frailty Assessment

In order to compare the onset and severity of heart beat frailty derived from our longitudinal

EKG analyses, we conducted a cross-sectional study to assess traditional, non-cardiac specific,

organism wide frailty in mice of advanced age.

Frailty Index

We used a validated mouse frailty index consisting of non-cardiac factors [11]. Variability

among mice in heart functions that change at older age have been shown to correlate better with

this frailty index score than with chronologic age [16]. The frailty index was administered to a sub-

set of older aged awake mice. A mouse Frailty Assessment Form was administered in parallel [6].

The assessment was performed in parallel by two different investigators in order to deter bias. The

scores from each assessment were summed from the 31 parametric observations to calculate the

mouse's frailty score, and the scores determined by each investigator were averaged to generate

final frailty score.

210   Energetic Efficiency

Energetic efficiency, defined as kinetic work performed/oxygen consumption/unit time,

was assessed cross sectionally in another sub-set of older mice. Work was measured as the distance

traveled per unit of time and Energetic Efficiency. Oxygen consumption was measured through an

enclosed treadmill coined as "metabolic treadmill", where the mouse is subjected to run at its age

groups' average gait speed determined by over- ground walking analysis in earlier TSE MotoRater

assessments. In order to ensure familiarity, mice were acclimated on the metabolic treadmills

(Columbus Instruments International, Columbus, OH) at 5 m/min for 30 min the day prior to

testing. The next day mice ran at the previously defined age-matched natural walking gait speed

for 45 min [17]. The external motivators included electric shock from an electrified metal grid

located near the moving belt to entice mice to run. Mice that were unwilling or incapable to

continue running after being shocked 5 consecutive times within a few seconds met criterion for

exhaustion and the testing ends.

Results (4270 words)

Longitudinal Cohort Longevity

The Kaplan-Meier Survival Curve of our entire cohort of C57/BL6 is shown in Fig 1A. The median life span of the total C57/BL6 cohort (n = 58) of our study was approximately 24 months (Fig 1). Because the main focus of our study was to identify how HR and HRV become altered in advanced age, we analyzed the life-long (6-30 month) EKG time series of each mouse in the basal and intrinsic states. We took 30 months of age as the maximum total life span of our mouse cohort because only 3 of 58 (5.2% of the total cohort) survived to this age. We focused on long-lived (LL) mice (n = 29), which we defined as those that achieved the median cohort lifespan of 24 months. Thus, the number of LL mice that survived from 6 to 24 months of age is constant at all ages and that beyond 24 months of age, the number of surviving mice becomes reduced as aging progresses, i.e., n = 17, and 3 at 27 and 30 months respectively (Table S1).

The survival curve for the LL subset of our total mouse cohort is also shown in Fig 1A. Note its relative rectangularity compared to the survival curve of the entire cohort, and its steep rate of decline relative to that of the entire cohort The average life span of LL mice was 27.4 months (834 days or 119 weeks and 1 day) and relative to the entire cohort, LL mice at 24 months of age had already achieved about 75% of the mouse maximum life span.

RR Interval Signature buried within EKG Time-Series

We assessed the EKG time-series RR interval variability patterns in the time, frequency, non-linear, and fragmentation domains. Mean intrinsic RR interval variability data for all measured parameters at each age are listed in Table S1, and the corresponding mixed-effects ANOVA analyses of these data are presented in Table S2. The kinetics and degrees of

synchronization within SAN cells determine the intrinsic rhythms buried within an EKG time

series during double autonomic blockade.

Representative EKG RR interval time-series recorded during autonomic blockade of a LL

mouse at 6 and 30 months of age are shown in Figures 1B&D. RR interval variability (figures

C&E) confirms that the SAN does not function as a metronome [18], as the heart beats in real

time it has no a priori knowledge of its **mean** beating rate; instead, the time at which the next beat

is generated is based on its memory of intrinsic mechanisms affected by prior beats [3]. Thus, a

mean inter-beat within a time series does not exist in real-time but is calculated post hoc as the

sum of individual and variable RR intervals in an EKG time series over a fixed period of time.

For example, the mouse in Figure 1B at 6 months of age generated QRS complexes with 16 RR

intervals, whereas at 30 months, only 8 RR intervals were generated within the same duration of

time. The mean RR interval of an EKG time series does not capture this exquisite RR interval

variability because the mean RR, calculated post hoc from RR intervals, assumes that all intervals

are equal.

Non-constant interval variability among RR intervals within an EKG time series generates

short and long-range correlations, nonlinearity components and variable frequency distributions

of RR intervals within the time series [19]. The age-associated shifts in the RR interval variability

pattern in the intrinsic state for the representative mouse (Figure 1B compared to Figure 1C, right

panels) is striking, not only in the difference for the range of RR intervals but also differences in

the synchronization of RR interval clusters. These RR interval rhythms buried within an EKG

time series inform on the beat-to-beat variability in the kinetics of molecular mechanisms and the

extent to which they are synchronized within and among cells [3].

SAN Signatures within EKG Time Series Rhythms Measured Longitudinally Over the Life Course

The Poincaré diagram is a convenient, informative method to visualize frequency distributions of successive RR intervals within a time series that underlie the mean RR interval that relate to each other in the time domain. In a Poincaré diagram, a given RR interval (N) is plotted against the next RR interval (N+1). Fig 2A, inset, shows the Poincaré diagram of the LL mouse RR intervals presented in Fig 1. The points within the Poincaré plot can be fit to an ellipsoid, with the spread of the data measured as the SD1 or SD2;the Poincaré SD1 informs on short-range correlations among RR intervals within the EKG time-series, while SD2 informs on long-range correlations among intervals [20]. The intersection of SD1 and SD2 at the center of the ellipse is the mean RR interval, and the SD1:SD2 informs on the nonlinearity of the correlations of RR intervals within a given EKG time-series [20].

Fig 2A shows the distributions of intrinsic RR intervals in the time domain (during autonomic blockade) for the same LL mouse at 6 and 30 months of age depicted in Fig 1 and Fig 2A inset. Note that not only is the RR interval (ms) distribution broader at 30 months than at 6 months of age, but the number of RR intervals within the distributions that are synchronized within narrow bins of intervals is substantially less at 30 months than at 6 months.

Intrinsic SD1, SD2, SD1:SD2, and the mean intrinsic RR interval throughout the life span of the LL mouse in Figs 1&2A are shown in Figs 2B&C. Note how the SD1 declines while SD2 increase beyond 21 months, creating a decreasing SD1:SD2. The average intrinsic SD1, SD2, and SD1:SD2 mean intrinsic RR intervals of all the LL cohort over its entire life span are shown in Figs 2D&E. On average, SD1, SD2 and the mean RR interval of LL mice markedly increased beyond 21 months of age, with the SD2 increasing to a much greater extent than SD1 (Fig 2D),

reflecting nonlinearity (SD1:SD2) of short and long-range RR interval correlations as age

increases beyond 21 months (Fig 2E). The intrinsic mean RR in all LL mice increases markedly

beyond 21 months (Fig 2F).

Our experimental design permitted the calculation of mouse-specific rates of change

(ROC) within aggregate RR interval data. In addition to providing the rate at which signatures

change during a given mouse's life course (i.e., mouse-specific rates), the ROC also provides

insight into the rates at which inter-mouse-to-mouse variability of rates at which RR interval

signatures change throughout the life course. ROC of intrinsic RR individual LL mice are

illustrated in Fig 2G, the most significant increase occurring between 21 and 24 months of age.

SD of the mean RR interval (SDRR) can be calculated from the distribution of RR intervals.

The **basal** SDRR has generally been attributed to the magnitude of parasympathetic input to the

SAN [21], but the **intrinsic** SDRR measured during double autonomic blockade reflects beat-to-

beat variability in mechanisms **intrinsic** to SAN cells. Beyond 21 months, the average increase in

the intrinsic SDRR of the entire LL cohort (Fig 2H) is due to an increased age-associated beat-to-

beat variability of kinetic functions that determine action potential firing rate within pacemaker

cells of the SAN [3]. The ROC of intrinsic SDRR individual LL mice is illustrated in Fig 2I, the

most significant increase occurring between 21 and 24 months.

Like the SD1:SD2, the slope coefficients α_1 and α_2 derived from detrended fluctuation

analyses (DFA) together inform on nonlinearity of short-range DFA (slope coefficient α_1) and

long-range (DFA slope coefficient α_2) RR interval correlations within an EKG time-series [19].

Examples of DFA analyses at 6 and 30 months in the same long-lived mouse depicted in Figures

1-2 are shown in Figure 3A. The calculated slope coefficients, α_1 and α_2, throughout the entire

life course of this mouse are shown in Figure 3B, Tables S1 and S2. The average intrinsic α_1 and

α_2 of the entire LL cohort also increased throughout the life span, with the increase accelerating

sharply (nearly doubling) between 21 and 30 months (Fig 3C, Table S1, and S2). Mouse-specific

ROC for α_1 and α_2 for all LL mice are shown in Figure 3D.

SAN Aging Signature of RR Interval Rhythms in the Frequency Domain

Fast Fourier transforms provide power spectral densities (PSD) of RR intervals within an

EKG interval time series, identifying different frequencies of hidden rhythms that are missed in

EKG time-series analyses [22]. A greater total power informs on greater complexity, i.e., less

coherence among RR intervals within the RR interval time series. Between 6 and 21 months, the

mean intrinsic total power of the LL cohort did not vary significantly (Fig 4A, Table S2); but,

beyond 21 months of age, the mean intrinsic total power **increased** by about 2-fold, similar to

ROC for SDRR (Fig 2I). ROC of both total power and SDRR varied substantially among LL mice

between 21 and 30 months (Fig 4B).

The total power can be partitioned into regions that differ in their approximate

frequencies, including the high frequency (HF), low frequency (LF), and very low frequency

(VLF) power components (Fig 4C, Table S1). Examples of power spectra of RR intervals within

the EKG time-series at 6 months and 30 months of age in the same long-lived mouse depicted in

Figs 1-3, are shown in Fig 4C. Average values of intrinsic VLF power components normalized

to total power in all mice increased modestly between 6 and 21 months (Fig 4D, Table S2).

Beyond 21 months, the intrinsic %VLF power increased exponentially as intrinsic %HF power

plummeted (Fig 4D, Table S2). Beyond 27 months, the **intrinsic** %VLF power exceeded that of

intrinsic %HF power. The contribution of intrinsic HF power to intrinsic total power declined by

about five-fold between 6 and 30 months in LL mice (Fig. 4D, Table S2).

Intrinsic RR Interval Fragmentation Signatures

RR interval fragmentation analyses that attempt to quantify the erratic beating of the SAN by detecting small, aperiodic long-range fluctuations in RR intervals are an expansion of the aforementioned mentioned RR interval variability analyses [23]. RR interval fragmentation parameters assessed include: percentage of inflection points (PIP), or points at which the first RR difference changes sign; the percentage of short RR segments (PSS); the inverse of the average RR segment length (IALS); and the percentage of alternating RR segments (PAS). A larger value of each parameter denotes a more fragmented RR interval structure within the RR interval time series.

Changes in PIP, IALS, and PSS in our LL cohort resembled each other across the mouse lifespan (Fig S2). At ages less than 21 months, intrinsic IALS PIP and PSS displayed slight but statistically significant age-associated trends. Beyond 21 months, neither mean intrinsic PIP nor IALS varied significantly with age (Table S2); but, intrinsic PSS increased beyond 21 months (Fig S2, Table S2). The average LL cohort intrinsic PAS increased sharply beyond 21 months of age (Fig 5A). The individual ROC for the LL intrinsic PAS can be seen in Fig 5B. The Mean RR and PAS across the entire age span for the same mouse highlighted in Figs 1-4 are shown in Fig 5C.

Relationships of the rates at which RR intervals change during the life course to the rates at which the mean RR interval changes

Because a crucial aspect of our RR interval variability analyses was to discover which intrinsic RR interval variability rhythms measured over the entire life course of a given mouse

underlie the life course changes in the mean intrinsic RR interval in that mouse. In other terms,

we sought to discover which RR interval variability signatures of aging in a given mouse define

the rate at which the mean intrinsic RR interval increases with age in that mouse. We performed

variable cluster analyses using mouse-specific ROC of all measured RR interval variability

parameters between 21 and 30 months of age in each mouse. The dendrogram in Figure 6A

illustrates the clustering of the mouse-specific ROC of the mean intrinsic RR interval and the

intrinsic RR variability parameters between 21-30 months. ROC of intrinsic RR interval

variability parameters that clustered with the ROC of mean intrinsic RR intervals in a given

mouse (Fig 6A red cluster) were VLF, HF, α_2, PAS, and SD1:SD2. Importantly, this cluster also

includes bodyweight ROC, a component of the frailty assessment index (see below). Bivariate

correlations for mouse-specific ROC of intrinsic RR variability signatures that clustered together

throughout life (Fig 6A) are listed in Table 1 and illustrated in Fig S1. ROC of the intrinsic RR in

individual mice as they age are illustrated in Fig 2G, with the most significant increase occurring

between 21 and 24 months of age.

It is important to note that the ROC of RR interval variability signatures excluded from

the red cluster (Fig 6A blue cluster) did **not** correlate with the ROC of the mean intrinsic RR

interval between 21 and 30 months of age (Table S3). The age-associated increase in nonlinearity

of RR interval variability (SD1:SD2) within short (SD1) and long (SD2) range RR interval

correlations is a crucial factor of the marked increase in mean intrinsic RR interval prolongation

between 21 and 30 months. Unlike the ROC SD1:SD2 (Table 1, Fig S3) neither the SD1, nor the

SD2 ROC, per se, correlated with the age-associated increase in mean RR interval ROC (Table

S3). Similarly, ROC for the standard deviation of the mean intrinsic RR intervals (SDRR) did not

correlate with the ROC of the mean intrinsic RR during late life (Table S3). Unlike the SD1:SD2

(Table 1, S1), the SDRR does not provide information regarding the increase in the nonlinearity

of RR interval variability within long and short-range RR interval correlations that are **directly**

linked to the rate at which the mean intrinsic RR interval increases in advanced age (Table 1, Fig

S1).

The rate at which the mean intrinsic RR increases over time in a given mouse between 21

and 30 months was significantly correlated with concurrent increases in the rates at which α_2 and

%VLF power increased, as well as the rates at which %HF power and SD1:SD2 became reduced.

The rates at which these RR interval parameters were underlying the increase in mean intrinsic RR

interval during late-life were strongly correlated with each other: reductions in the rate of %HF

power and SD1:SD2 are positively correlated, and the rates at which %VLF power and α_2 increase

are positively correlated (Table 1, Fig S1). Thus, ROC of relative %HF and %VLF power during

the late-life course were inversely correlated with each other (Table 1).

Mouse-specific ROC of %HF and %VLF power were mirror images and therefore the rates

at which they change are inversely related to each other. The rate at which the mean intrinsic RR

increased over time in a given mouse between 21 and 30 months was significantly and positively

correlated with concurrent increases in the rates at which α_2 and %VLF power increased, and

inversely with the rates at which %HF power and SD1:SD2 became reduced. The rates at which

these RR interval parameters underlying the increase in mean intrinsic RR interval during late-life

were strongly correlated with each other: reductions in the rate of %HF power and SD1:SD2 are

positively correlated, as were the rates at which %VLF power and α_2 increased (Fig S1, Table 1).

Thus, ROC %HF and %VLF power were inversely correlated with each other during the late-life

course (Table 1).

The ROC of intrinsic PAS (Fig 5B) in a given mouse was the only RR interval

fragmentation index that correlated with the rate at which mean intrinsic RR interval increased

during late life (Table 1, Fig S1).

Thus, between 21 and 30 months of age, the rate at which the mean intrinsic RR interval

increased in a given mouse was associated with: (1) the rates at which HF and VLF components

within the EKG time-series decrease and increase respectively; (2) positively with PAS; (3)

increases in the rate at which nonlinearity of RR interval variability (SD1:SD2) between short-

range (SD1) and long-range (SD2) RR interval correlations decreased; (4) the rates at which RR

interval variability increased within long-range RR interval correlations (SD2, α_2). Because the

increase in α_2 ROC in a given mouse was significantly correlated with the increase in mean

intrinsic RR interval of that mouse (Tables 1 & S1), the ROC of α_2 and SD1:SD2 during late life

are inversely correlated with each other (Table 1, Fig S1).

Quantitative Comparisons of the Relative Magnitudes of Life-Long Changes in RR Interval

Variability Signatures and Mean RR Interval

To compare the relative magnitudes of changes in RR interval variability parameters

(signature of aging in Table 1 and Fig 3A) that underlie the marked prolongation of the intrinsic

RR in advanced age, we measured the values at 3-month intervals in each mouse, which were

normalized to the 6 month values (Fig 6B). Intrinsic mean RR increased on average by 50%

between 6 and 30 months, with much of the increase occurring beyond 21 months. Intrinsic

%VLF power increased by about 2.5 fold; in contrast, intrinsic %HF power became reduced by

about 20%, and the Poincaré SD1/SD2 was reduced by 50% between 6 and 30 months. PAS

increased by 50% and DFA long-range intrinsic α_2 increased by 25%. Note that beyond 21

months of age the increase in mean intrinsic RR interval is paralleled by sharp increases in α_2,

PAS, and %VLF power with the relative change in %VLF power greater than PAS, greater than

α_2; and by sharp reductions in SD1:SD2 and %HF power (beyond 24 months in Fig 6B).

In summation, our results show the impact of advanced age on the signature of RR interval

variability and mean RR interval. The ROC of SD1: SD2, intrinsic HF, intrinsic VLF, and α_2 in a

given mouse are related to each other during late life, and these ROC of non-linear and frequency

domain RR interval variability signatures underlie the mean intrinsic RR interval in advanced

age.The unique clustering of mouse-specific RR interval variability signatures (Fig 6A) and

correlations of these ROC with the rate at which the mean intrinsic RR interval in a given mouse

increases in advanced age (Table 1, Fig S3) specifically inform on how the aging signatures of

rhythms within EKG time series define the mean intrinsic RR interval as age advances.

Impact of Autonomic Input on Age-Associated Changes in Intrinsic SAN RR Interval

Signatures

Modulation of intrinsic heartbeat intervals imparted by autonomic input to the SAN reflects

the combined effects of autonomic neurotransmitter impulses delivered to SAN cells and the

responses of these pacemaker cells to that input. We next sought to investigate the effect of

autonomic input (the difference between the mean intrinsic value and basal value of RR interval

parameters in a given mouse) to glean additional insight into heartbeat dysfunction during late life.

It is important to note that

Table S2 and Fig 7 show the signatures of autonomic input on RR interval variability

signatures that underlie the rate at which the mean intrinsic of RR interval of LL mice increased

during aging. Note that between 6 and 21 months of age, the difference between intrinsic and basal

states of RR interval variability signatures did not differ (Tables 1 & S2). Between 21 and 30 months

of age, the difference between intrinsic and basal state RR interval variability signatures differed

markedly, during aging and among each other (Fig 7A-F, Table S2).

Before 21 months of age, SD1:SD2 was substantially less than the average SD1:SD2 of all

mice in the intrinsic state (Fig 7A). Beyond 21 months, the average intrinsic SD1:SD2 began to

decline while its basal state counterpart increased, with the average basal and average intrinsic

SD1:SD2 converging at 27 months of age. Thus, up until 27 months of age, the effect of autonomic

input was to reduce the intrinsic SD1:SD2. Note that average basal and intrinsic SD1 and SD2 in

the three mice surviving up to 30 months of age crossed beyond 27 months, with the intrinsic

SD1:SD2 reducing, while the basal SD1:SD2 increased, and that the crossing pattern of basal and

intrinsic SD1:SD2 between 27 and 30 months of age is a general feature of other RR variability

parameters that correlated with the mean RR intervals in very old mice e.g., %HF power (Fig 7B)

%VLF power (Fig 7C) and DFA slope coefficient α_2 (Fig 7D).

The effects that autonomic input imparts to %HF power (Fig 7B) is strikingly similar to that

of its effect on the nonlinearity of short to long-range RR interval correlations within the time-series

informed by SD1:SD2 (Fig 7A). Recall (Table 1, Fig S1) that the ROC of intrinsic %HF power and

of SD1:SD2 in individual mice were positively correlated with each other. The effects of autonomic

input on %VLF power (Fig 7C) and DFA α_2 (Fig 7D) are also strikingly similar across the entire

age span from 6 to 30 months. Recall also (Table 1, Fig S1) that: (1) the ROC for intrinsic %VLF

power and α_2 between 21 and 30 months were positively correlated to each other and to the rates at

which the mean RR interval in individual mice changed over this age range; and (2) that the ROC

for %VLF power and α_2 were inversely correlated with the ROC of %HF power and SD1:SD2

(Table 1, Fig S2). Importantly, note in Figure 7 that the shapes of the mean %VLF power and α_2

(Fig 7C-D) are also mirror images of those of %HF power and SD1:SD2 (Fig 7A-B); and also note

the effects that autonomic signatures imparts to intrinsic %HF power and SD1:SD2 are strikingly

similar.

PAS was the only RR interval fragmentation index throughout the entire life span that was

not statistically significantly impacted by autonomic input (Table S2, S3). Average basal and

intrinsic PAS in all LL mice beyond 21 months of age (Fig 7E) resembled a muted version of mean

intrinsic and basal RR interval (Fig 7F), but unlike average intrinsic RR interval, the response to

autonomic input was not significant. (Table S2). Although the average basal RR interval of all LL

mice did not significantly differ between 6 and 21 months of age, the mean intrinsic RR manifested

small but statistically significant changes (Table S2); and beyond 21 months of age, while average

mean intrinsic RR precipitously increased (Fig 7F), autonomic input had a marked effect of

reducing the mean RR interval (Table S2).

We may deduce that either autonomic input to the SAN increases, the responsiveness of

SAN pacemaker cells to this input increases with age, or both mechanisms are involved. Because

SAN responsiveness to autonomic neurotransmitters is known to be reduced in advanced age [24,

25], the reduction in mean intrinsic RR interval in late-life while autonomic input is intact reflects

a **net** sympathetic response (increase sympathetic or reduced parasympathetic) to autonomic input,

because of the directionality of the response (to reduce the RR interval or increase HR) (Fig 7F).

Note, however, that beyond 21 months of age, autonomic input fails to restore the mean youthful

basal RR interval that prevailed between 6 and 21 months (dashed line in Fig 7F), reflecting an

inability of the net sympathetic autonomic input to overcome the high degree of disorder within the

SAN tissue later in life. The impact of autonomic signatures on the intrinsic RR interval signatures

over the entire life course of the representative mouse shown in Figs 1-7 are shown in Fig 8. Figs

S2 & S3 illustrate the average basal and intrinsic RR intervals and RR variability parameters of

the entire LL cohort having ROC that were not significantly correlated with mean RR interval

ROC in individual mice.

Body Weight, Frailty Index and Energetic Efficiency

Loss of body weight is essential in non-cardiac frailty and may serve as an important

biomarker in longevity [26]. Bodyweight in the long-lived mouse cohort changed in a non-linear

manner, increasing significantly with age up to 18 months and significantly declining after 21

months until the end of life (Fig 9A), with the variability in mouse-specific ROC increasing beyond

18 months (Fig 9B). Considering body weight loss as a sign of frailty, it appears that the earliest

signs of organismal frailty begin to emerge as early as 18 months (Fig 9A-B). Interestingly, the

the ROC of the autonomic effect on intrinsic mean RR correlates (r = -0.3387) with the change in

body weight.

We assessed body weight and whole organism frailty index that excluded cardiac

parameters, in another cross-sectional sub-set of C57/BL6 mice at younger and older ages. Body

weight increased substantially as age increased during late life (Fig 9C). As in our LL longitudinal

cohort, bodyweight declined cross-sectionally in advanced age and was paralleled by an increase

in frailty score (Fig 9D).

It is well known that the decline in health status is accompanied by a reduction in the

efficiency of oxygen utilization [27, 28], which can be quantified as energetic efficiency measured

as oxygen consumed during walking/speed of kinetic work performed [17]. Energetic efficiency

measured cross-sectionally in another subset of mice also became reduced with advanced age (Fig

9E). An increase in the respiratory exchange ratio accompanied a reduction in energy efficiency

(Fig 9F).

Thus, deterioration of the SAN interval variability signatures that underlie the marked

increase in mean RR, i.e., heartbeat frailty observed during advanced age in our longitudinal cohort

(Fig 1-8) are paralleled by increases in non-cardiac constitutional indices of frailty measured in

other cross-sectional subsets (Fig 9). These additional perspectives on age-associated increases in

the frailty index and reduced efficiency of O_2 utilization establish that C57/BL6 mice at or beyond

strain median life span of 24 months manifested constitutional signs of non-cardiac frailty.

Discussion (WC 2314)

We conducted a unique longitudinal study of mouse RR interval variability and mean RR interval analyses in EKG time-series recorded at 3-month intervals from 6 months to the end of life (approximately 30 months). By measuring RR interval variability prior to and during double autonomic blockade in a sub-set of mice that achieved the median cohort life span (24 months) or survived beyond this age we were able to assess how **intrinsic** SAN RR interval variability changed over the entire life course, particularly in advanced age. We specifically assessed mouse-specific rates of change for intrinsic RR interval variaility parameters to ascertain how these signatures vary within the same mouse over time, determining the rate at which the mean intrinsic RR interval increases during late-life in that mouse (SAN frailty), paralleling organismal frailty that emerges during late life.

The Signatures of SAN Frailty in Long-Lived Mice of Advanced Age

Most measured RR interval variability parameters and mean RR interval remained stable up to about 18 months, but beyond 21 months of age changed dramatically. Between 21 and 30 months, most **intrinsic** RR variability parameters (measured during dual autonomic blockade) manifested progressive changes, reflecting dysfunction of mechanisms **intrinsic** to pacemaker cells residing within the SAN [3], causing the mean intrinsic RR interval to increase with advanced age.

Cluster analyses of mouse-specific rates of change (ROC) for the intrinsic RR interval variability correlations among longitudinal ROC of the clustered parameters enabled the identification of signature components of intrinsic RR interval variability that underlie the marked

increase in the mean intrinsic RR interval that emerged beyond 21 months of age in LL mice (Table

S2 Fig 6). These signatures of functional deterioration that occur within and among cells

comprising SAN tissue underlying the marked increase in the mean intrinsic RR interval beyond

21 months of age are broadcast to the body surface as "heart beat music" [29] and can be heard on

numerous EKG RR interval variability "channels," including those that broadcast signals in the

time, frequency, non-linear and fragmentation domains. Beyond 21 months of age cluster analyses

pointed to ROC of intrinsic VLF, α2, HF, SD1/SD2 ratio and PAS being the signatures that sub–

tended the rate at which the mean RR interval over this age range increased in individual mice.

The time at which successive RR intervals occur within a time series requires **memory**

within SAN pacemaker cells created during prior beats [3]. Because each heartbeat is an emergent

phenomenon [2], i.e., the SAN is not a metronome with a pre-written score. Increases of both

intrinsic SD1 and SD2 means that RR interval variability within both short-range and long-range

correlations of RR intervals generated within and among populations of SAN pacemaker cells

become increased in advanced age in LL mice as SAN frailty begins to emerge. The nonlinearity

of RR interval variability within short and long-range RR interval correlations also increase with

aging, as reflcted in shifts in intrinsic SD1:SD2 over the late life course. A youthful profile of SD1,

SD2, and SD1:SD2 likely requires high frequency signal processing, because the rate at which

intrinsic %HF power is lost between 21 and 30 months strongly correlates with the rate at which

intrinsic SD1:SD2 changes. Significant increases in SD1, SD2 and correlations between the rates

at which intrinsic SD1:SD2 and %HF power components and decline in a given mouse are

associated with the rate at which mean intrinsic RR interval increases in that mouse would appear

to reflect, in part at least, a loss of memory with respect to when to generate the next AP as SAN

frailty emerges. The corollary is that high frequency and nonlinearity of signals emerging from

within and among populations of SAN cells informs on the memory that is required to maintain

short (high frequency) and fairly regular (synchronized) RR intervals. The strong positive

correlations of the rates at which HF components of RR intervals within the time-series decreased

between 21 and 30 months in each mouse and the rates at which SD1:SD2 decreased in that mouse

suggests that a reduction in the kinetics of pacemaker cell clock functions and desynchronization

among these functions [3] may be linked to increased nonlinearity of RR interval variability within

both short (SD1) and long-term (SD2) RR interval correlations. Rates at which intrinsic %HF

power is lost and nonlinearity in (SD1:SD2) increase in a given mouse during advanced age were

correlated with the rate at which mean intrinsic RR interval increased in that mouse.

The tight coupling between the rate at which the reduction in intrinsic %HF power and the

rate at which intrinsic %VLF power increased, in the absence of a change in the rate of total power

is likely attributable to a (Fig 7) reduction in the kinetics of intracellular Ca^{2+} and membrane

potential transitions, leading to variability in mechanisms within and among SAN cells that relate

to SAN pacemaker cell memory loss during the emergence of SAN frailty [3, 30, 31]. The shift in

intrinsic HF power to VLF power of mice in advanced age is likely attributable to reductions in

the kinetics of coupled-clock mechanisms that drive SAN pacemaker cell function [3, 31]. A

reduction in these kinetics leads to desynchronization of molecular actions within SAN pacemaker

cells associated with a loss of memory of synchronization effects created by the prior action

potential [3].

The rates at which intrinsic DFA, slope coefficient α_2, another measure of long term RR

interval variability measure, and intrinsic %VLF power increased beyond 21 months of age in a

given mouse were also significantly correlated with the rates at which the mean RR interval

increases. Directionally opposite changes in the rate at which intrinsic α_2 (increased) and SD1:SD2

decreased during advanced age means that the rate at which α_2 increases reflects the rate at which HF signal processing is lost within and among populations of SAN cells. Thus, similar to the rate at which %VLF power increases beyond 21 months of age, the rate at which α_2 increases is also correlated with the rate at which the mean intrinsic RR interval increases. The strong positive correlations between the rates at which %VLF power and α_2 increased in a given mouse between 21 and 30 months of age indicate that the rate at which α_2 increases reflects an increase in RR variability of VLF components within long-range RR interval (α_2) correlations. In short, the most characteristic signature of SAN frailty is a loss in the ability of the SAN pacemaker cells to generate HF signals, resulting in reduced synchronization among the ensemble of cells. This translates into a markedly prolonged interval at which a given heartbeat follows the previous beat, resulting in a marked increase in the mean RR interval or reduction in mean intrinsic HR.

Increased SD1, SD2, SD1:SD2, increases in the DFA and slope coefficient α_2, and shifts from intrinsic %HF power to intrinsic %VLF power underlie the increase in **intrinsic** percent alternating segments (PAS) within the EKG time-series of LL mice of advanced age that characterize SAN frailty beyond 21 months of age. This increase in the percentage of alternating long-short segments within the EKG interval time-series in LL mice of advanced age (i.e., increase in PAS) is reminiscent of action potential and calcium alternans that characterize dysfunctions within and among SAN pacemaker cells that underlie cardiac arrhythmias [32].

Cellular and Molecular Basis of RR Intveral Variability Signatures of the Fraile SAN

Both experimental and theoretical perspectives have led to the idea that a coupled-oscillator system intrinsic to individual pacemaker cells drives normal, youthful automaticity [33]. Briefly,

a chemical oscillator ("Calcium-Clock") and an ensemble of current oscillators within the cell

surface membrane ("Membrane Clock") mutually entrain each other in a feed-forward manner to

generate the rhythmic electrochemical gradients underlying action potential cycles [33].

Constitutive activation of adenylyl cyclase type 8 drives calcium and phosphorylation-dependent

modulation of molecules that operate within the pacemaker cell clocks [34]. The effectiveness or

fidelity of clock-coupling, determined by synchronization of mechanisms that regulate Ca^{2+} and

membrane potential, controls the rhythm of action potential firing. The frail intrinsic RR interval

variability signatures in LL mice of advanced age stem, at least in part, from disorder

(desynchronization) among pacemaker cell clock molecules [3, 31], resulting in desynchronization

of rhythms within and among clusters of pacemaker cells residing within SAN tissue [2].

Both age-associated reductions in clock molecule expression and reduced synchronization

of molecular activation within individual pacemaker cells may be root causes of SAN failure in

advanced age [31, 35]. But normal SAN function goes well beyond the function of individual

pacemaker cells because rhythmic impulses that emerge from the SAN result from synchronization

of heterogeneous, sub-cellular Ca^{2+} signals, not only within but also among SAN tissue resident

pacemaker cells [2]. Thus, the initiation of each heartbeat within the SAN is an emergent

phenomenon, and in addition to age-associated deficits within individual SAN cells, the ability to

synchronize heterogeneous local signals among cells may be compromised in advanced age.

Further, age-associated by pacemaker cell-matrix remodeling, including fibrosis [36] compounds

the disordered state within the frail SAN..

The Signatures of Autonomic Input to the Frail SAN

Adrenergic receptor activation increases cell calcium levels and protein phosphorylation,

thereby increasing the entertainment of clock coupling [3]. When the effectiveness of clock-

coupling increases in response to sympathetic autonomic input, RR variability and mean RR

become reduced [34, 37]. Conversely, reduced effectiveness of clock coupling in response to

parasympathetic autonomic input leads to prolonging the mean RR interval and increasing RR

interval variability [3]. Thus, RR interval variability and the mean RR interval in the **basal** state

result from modulation of the mechanisms intrinsic to SAN pacemaker cell that determine the

intrinsic RR interval variability and mean RR interval.

Our results demonstrate that autonomic modulation of intrinsic RR interval variability

differed over the life course, not only in magnitude but also in direction, particularly beginning

around 21 months of age. As the rate of decline in intrinsic SD1:SD2 accelerated beyond 21

months of age, that of the basal SD1:SD2 increased with the basal and intrinsic values converging

at 27 months of age. Thus, between 21 and 27 months, autonomic modulation of SAN pacemaker

cells mechanisms markedly reduced the SD1:SD2 (Fig 7A), indicating that the impact of this

autonomic input signature was to reduce the nonlinearity of RR interval variability that had

developed within short to long-range intrinsic RR interval correlations in LL mice over this age

range.

The signature of autonomic input that restored intrinsic %HF power was strikingly similar

to the autonomic signature of SD1:SD2 between 21 and 27 months of age. Conversely, the

signature of autonomic input was to reduce the increased intrinsic %VLF power and DFA α_2,

leading to preservation of their basal levels throughout the life course up to 27 months of age (Fig

7C-D). PAS was the only RR interval fragmentation parameter [38, 39] that did not have a

statistically significant signature, i.e., was not significantly modified by autonomic input. The net

result of these autonomic signatures on the intrinsic RR interval signatures was to reduce the basal

RR interval to a more youthful level. Further, because the RR interval became reduced (i.e., HR

was increased) this autonomic signature was net sympathetic in nature. Thus, age-associated

changes in intrinsic RR interval signatures (heartbeat music broadcast to the body surface)

reflected in the EKG time series is the net result of multiple changes within and among SAN cells

and their responsiveness to autonomic input. Nevertheless, even after this autonomic input the

basal state mean RR interval within a given mouse was not fully restored to its value at younger

ages because the autonomic signatures could not fully restore the RR interval variability signatures

to their youthful levels.

The Relationship of SAN Frailty to Body-Wide Frailty

We showed in an additional cross-sectional C57/BL6 mouse cohort that over the age range

in which marked disorder of SAN function emerged (in our LL longitudinal cohort beyond about

21 months of age), a body-wide frailty index not inclusive of cardiac parameters also sharply

increased during late life (Fig 8D). An increased frailty score with advanced age has also been

demonstrated in prior cross-sectional studies [17]. Of note, a prominent element within this frailty

index is bodyweight loss, which not only occurred in our cross-sectional cohort, but also began to

occur in all mice in our LL **longitudinal** cohort at 18 months of age (Fig 9). The increase in the

frailty index and loss of body weight during late-life in our cross-sectional cohort was

accompanied by a loss of energy efficiency (Fig 8E) and an increase in the respiratory exchange

ratio (Fig 8F).

In summary, in the context of the presence of these other non-cardiac frailty markers in

mice of advanced age, we may interpret the marked changes in increased intrinsic SAN RR

variability that underlies the marked increase in the intrinsic mean RR interval that emerges in late

life to be a signature of progressive SAN frailty. Specific aspects of this SAN frailty signature

include: (1) increased non-linear intrinsic RR interval variability within short and long-range

correlations of RR intervals; (2) reductions in the %HF power within intrinsic RR interval rhythms;

(3) increase in %VLF intrinsic power within RR rhythms; (4) increased long-range detrended RR

interval fluctuations; (5) progressive increase in the percentage of intrinsic RR interval segments

exhibiting alternans. In toto, these RR interval variability signatures reflect an inability to generate

high frequency RR intervals due to desynchronization of molecular functions within SAN

pacemaker cells that require memory of prior RR intervals. The emergence and progression of this

altered pattern of intrinsic RR interval variability beyond about 21 months of age cause the mean

intrinsic RR interval to progressively increase over this age range. Autonomic modulation of this

intrinsic SAN RR interval variability signature masks, to a large extent, SAN frailty cell that

emerges during late life concurrently with the emergence of non-cardiac, body-wide, organismal

frailty [40].

 Figure Legends

Figure 1. A. Kaplan-Meier Survival curves for entire cohort of mice (n = 58) and for long-lived

subset of mice (n = 30). Intrinsic EKGs and Intervalograms at 6 months (B&C) and at 30 months (D&E).

Figure 2. A. Distribution of intrinsic RR intervals at 6 and 30 months for mouse ZA1730. Inset,

Poincare plot of intrinsic RR interval data at 6 and 30 months for mouse ZA1730. B. Intrinsic SD1 and SD2

for Mouse ZA1730. C. Intrinsic SD1/SD2 and Mean RR for Mouse ZA1730. D. Average loess smooth curves

of intrinsic SD1 and SD2. E. Average loess smooth curves of intrinsic SD1/SD2. F. Average loess smooth

curve for intrinsic mean RR. G. Mouse-specific rates of change for intrinsic mean RR in long-lived mice H.

Average loess smooth curve of intrinsic SDRR. I. Mouse-specific rates of change of SDRR in long-lived mice.

Figure 3. A. Detrended Fluctuation Analysis (DFA) illustrating intrinsic α_1 and α_2 at 6 and 30 months

for mouse ZA1730. B. Trajectories of Intrinsic α_1 and α_2 in mouse ZA1730. C. Average loess smooth curves

of intrinsic α_1 and α_2. D. Mouse-specific rates of change of intrinsic α_1 and α_2 in long-lived mice.

Figure 4. A. Average loess smooth curve of intrinsic Total Power (PSD). B. Mouse-specific rates of

change of intrinsic Total Power (PSD) in long-lived mice. C. Intrinsic frequency domain data at 6 and 30

months for mouse ZA1730. D. Average loess smooth curves of intrinsic VLF and HF. E. Mouse-specific rates

of change of intrinsic VLF and HF in long-lived mice.

Figure 5. A. Average loess smooth curve of intrinsic PAS. B. Mouse-specific rates of change of

intrinsic PAS in long-lived mice. C. Trajectories of Intrinsic PAS in mouse ZA1730.

Figure 6. A. Cluster analysis dendogram of intrinsic rates of change in the late time period (after

21 months) in long lived mice. B. Relative change of intrinsic Mean RR and RR interval variability from 6

month baseline.

Figure 7. Average loess smooth curves for basal and intrinsic RR interval variablilty signatures and

on the mean RR interval during the enitre life course of all long-lived mice for SD1:SD2 (A), %HF of total

power (B), %VLF of total power (C), α_2 (D), % alternating sgements (E), and mean RR (F). Arrows show the

effect of autonomic modulation.

Figure 8. Impact of autonomic signature on intrinsic RR interval variability signatures; SD1:SD2

(A), %HF power (B), %VLF power (C), α_2 (D), % PAS, and on the mean RR interval (F) during the entire

lifespan of the representative mouse featured in Fig 1 – 6.

Figure 9. A. Average loess smooth curve for body weight. B. Mouse-specific rates of change of

body weight. C. Body Weight. D. Frailty Index, E. Energetic efficiency, and F. Respiratory Exchange Ratio

vs. Age.

Acknowledgements

735 This study was partially supported by the NIA Intramural Research Program and partially by

736 ISF330/19. This research was partially supported by the Technion Hiroshi Fujiwara Cyber Security

737 Research Center and the Israel Cyber Directorate (M.D).

739 We wish to acknowledge the indefatigable editorial assistance of Loretta E. Lakatta, R.N., B.S.N.,

740 and Syevda Tagirova Ph.D. for assistance with fitting Poincaré plot ellipses.

References

1. Armour, J.A., *Potential clinical relevance of the 'little brain' on the mammalian heart.* Exp Physiol, 2008. **93**(2): p. 165-76.
2. Bychkov, R., et al., *Synchronized Cardiac Impulses Emerge From Heterogeneous Local Calcium Signals Within and Among Cells of Pacemaker Tissue.* JACC Clin Electrophysiol, 2020. **6**(8): p. 907-931.
3. Yang, D., et al., *Ca2+ and Membrane Potential Transitions During Action Potentials Are Self-Similar to Each Other and to Variability of AP Firing Intervals Across the Broad Physiologic Range of AP Intervals During Autonomic Receptor Stimulation.* Frontiers in Physiology, 2021. **12**(953).
4. Jose, A.D. and D. Collison, *The normal range and determinants of the intrinsic heart rate in man.* Cardiovasc Res, 1970. **4**(2): p. 160-7.
5. Freeling, J.L. and Y. Li, *Age-related attenuation of parasympathetic control of the heart in mice.* Int J Physiol Pathophysiol Pharmacol, 2015. **7**(3): p. 126-35.
6. Whitehead, J.C., et al., *A clinical frailty index in aging mice: comparisons with frailty index data in humans.* J Gerontol A Biol Sci Med Sci, 2014. **69**(6): p. 621-32.
7. Yaniv, Y., et al., *deterioration of autonomic neuronal receptor signaling and mechanisms intrinsic to heart pacemaker cells contribute to age-associated alterations in heart rate variability in vivo.* Aging Cell, 2016. **15**(4): p. 716-24.
8. Dorey, T.W., et al., *Impacts of frailty on heart rate variability in aging mice: Roles of the autonomic nervous system and sinoatrial node.* Heart Rhythm, 2021.
9. Peters, C.H., E.J. Sharpe, and C. Proenza, *Cardiac Pacemaker Activity and Aging.* Annu Rev Physiol, 2020. **82**: p. 21-43.
10. Moghtadaei, M., et al., *The impacts of age and frailty on heart rate and sinoatrial node function.* J Physiol, 2016. **594**(23): p. 7105-7126.
11. Howlett, S.E. and K. Rockwood, *Factors that influence reliability of the mouse clinical frailty index.* J Gerontol A Biol Sci Med Sci, 2015. **70**(6): p. 696.
12. Behar, J.A., et al., *PhysioZoo: A Novel Open Access Platform for Heart Rate Variability Analysis of Mammalian Electrocardiographic Data.* Front Physiol, 2018. **9**: p. 1390.
13. Team, R.C., *R: A language and environment for statistical computing.* 2020.
14. RStudio Team, *RStudio: Integrated Development for R*. 2020, RStudio.
15. Wickham, H., *ggplot2 : Elegant Graphics for Data Analysis*, in *Use R!,*. 2016, Springer International Publishing : Imprint: Springer,: Cham. p. XVI, 260 pages 232 illustrations, 140 illustrations in color.
16. Kane, A.E., et al., *Age, Sex and Overall Health, Measured As Frailty, Modify Myofilament Proteins in Hearts From Naturally Aging Mice.* Sci Rep, 2020. **10**(1): p. 10052.
17. Petr, M.A., et al., *A cross-sectional study of functional and metabolic changes during aging through the lifespan in male mice.* Elife, 2021. **10**.
18. Shaffer, F., R. McCraty, and C.L. Zerr, *A healthy heart is not a metronome: an integrative review of the heart's anatomy and heart rate variability.* Front Psychol, 2014. **5**: p. 1040.
19. Peng, C.K., et al., *Quantification of scaling exponents and crossover phenomena in nonstationary heartbeat time series.* Chaos, 1995. **5**(1): p. 82-7.
20. Piskorski, J. and P. Guzik, *Filtering Poincaré plots.* computational methods in science and technology, 2005. **11**: p. 39-48.
21. Hayano, J., et al., *Accuracy of assessment of cardiac vagal tone by heart rate variability in normal subjects.* Am J Cardiol, 1991. **67**(2): p. 199-204.
22. Bhargava, V. and A.L. Goldberger, *Effect of exercise in healthy men on QRS power spectrum.* Am J Physiol, 1982. **243**(6): p. H964-9.

23. Lensen, I.S., et al., *Heart rate fragmentation gives novel insights into non-autonomic
mechanisms governing beat-to-beat control of the heart's rhythm.* JRSM Cardiovasc Dis, 2020. **9**:
p. 2048004020948732.
24. Lakatta, E.G., *Deficient neuroendocrine regulation of the cardiovascular system with advancing
age in healthy humans.* Circulation, 1993. **87**(2): p. 631-6.
25. Lakatta, E.G., *Cardiovascular regulatory mechanisms in advanced age.* Physiol Rev, 1993. **73**(2):
p. 413-67.
26. Alley, D.E., et al., *Changes in weight at the end of life: characterizing weight loss by time to death
in a cohort study of older men.* Am J Epidemiol, 2010. **172**(5): p. 558-65.
27. Davies, L.C., et al., *Enhanced prognostic value from cardiopulmonary exercise testing in chronic
heart failure by non-linear analysis: oxygen uptake efficiency slope.* Eur Heart J, 2006. **27**(6): p.
684-90.
28. Laine, H., et al., *Myocardial oxygen consumption is unchanged but efficiency is reduced in
patients with essential hypertension and left ventricular hypertrophy.* Circulation, 1999. **100**(24):
p. 2425-30.
29. Lakatta, E.G., *Heartbeat music.* Heart Rhythm, 2021. **18**(5): p. 811-812.
30. Tagirova Sirenko, S., et al., *Self-Similar Synchronization of Calcium and Membrane Potential
Transitions During Action Potential Cycles Predict Heart Rate Across Species.* JACC Clin
Electrophysiol, 2021.
31. Liu, J., et al., *Age-associated abnormalities of intrinsic automaticity of sinoatrial nodal cells are
linked to deficient cAMP-PKA-Ca(2+) signaling.* Am J Physiol Heart Circ Physiol, 2014. **306**(10): p.
H1385-97.
32. Weiss, J.N., et al., *Early afterdepolarizations and cardiac arrhythmias.* Heart Rhythm, 2010.
7(12): p. 1891-9.
33. Lakatta, E.G., V.A. Maltsev, and T.M. Vinogradova, *A coupled SYSTEM of intracellular Ca2+ clocks
and surface membrane voltage clocks controls the timekeeping mechanism of the heart's
pacemaker.* Circ Res, 2010. **106**(4): p. 659-73.
34. Moen, J.M., et al., *Overexpression of a Neuronal Type Adenylyl Cyclase (Type 8) in Sinoatrial
Node Markedly Impacts Heart Rate and Rhythm.* Front Neurosci, 2019. **13**: p. 615.
35. Lakatta, E.G., *Beyond Bowditch: the convergence of cardiac chronotropy and inotropy.* Cell
Calcium, 2004. **35**(6): p. 629-42.
36. Monfredi, O. and E.G. Lakatta, *Complexities in cardiovascular rhythmicity: perspectives on
circadian normality, ageing and disease.* Cardiovasc Res, 2019. **115**(11): p. 1576-1595.
37. Yaniv, Y., et al., *Synchronization of sinoatrial node pacemaker cell clocks and its autonomic
modulation impart complexity to heart beating intervals.* Heart Rhythm, 2014. **11**(7): p. 1210-9.
38. Costa, M.D., R.B. Davis, and A.L. Goldberger, *Heart Rate Fragmentation: A Symbolic Dynamical
Approach.* Front Physiol, 2017. **8**: p. 827.
39. Stein, P.K., et al., *Development of more erratic heart rate patterns is associated with mortality
post-myocardial infarction.* J Electrocardiol, 2008. **41**(2): p. 110-5.
40. Fried, L.P., et al., *Untangling the concepts of disability, frailty, and comorbidity: implications for
improved targeting and care.* J Gerontol A Biol Sci Med Sci, 2004. **59**(3): p. 255-63.

Table 1. Correlations and p-values among Intrinsic Rates of Change in long-lived mice

	Body Weight	Mean RR	VLF	HF	SD1/SD2	α_2	PAS
Body Weight: r		0.0914	0.2669	-0.1983	-0.2627	0.3155	0.0779
Body Weight: p		0.6312	0.1539	0.2934	0.1608	0.0895	0.6823
Mean RR : r	-0.1567		-0.0091	-0.0091	0.0601	0.1198	0.0603
Mean RR : p	0.4084		0.9621	0.9618	0.7524	0.5284	0.7515
VLF : r	0.2353	0.4774		-0.8964	-0.903	0.7772	-0.3217
VLF : p	0.2106	0.0076		2.1E-11	8.7E-12	4.4E-07	0.0829
HF : r	0.2452	-0.3914	-0.7248		0.9507	-0.5804	0.3271
HF : p	0.1915	0.0324	5.9E-06		8.9E-16	0.0008	0.0776
SD1/SD2 : r	-0.1620	-0.4264	-0.7533	0.8818		-0.6469	0.2817
SD1/SD2 : p	0.3924	0.0188	1.6E-06	1.2E-10		0.0001	0.1315
α_2 : r	-0.4078	0.4231	0.8189	-0.4662	-0.6444		-0.2243
α_2 : p	0.0253	0.0198	3.2E-08	0.0094	0.0001		0.2334
PAS : r	-0.1567	0.4223	0.2074	-0.2803	-0.2927	0.1397	
PAS : p	0.4084	0.0201	0.2716	0.1335	0.1165	0.4615	

The upper triangle contains the correlations in the early time period (before 21 months) and the lower triangle contains the correlations in the late time period (after 21 months).

Supplement Table S1A. Descriptive statistics of HR and HRV parameters in Long Lived mice.

		Intrinsic State (during double autonomic blockade)								
	Age	6	9	12	15	18	21	24	27	30
Time Domain										
	n*	30	28	29	30	28	29	28	16	3
Mean RR (ms)	mean	135.86	133.81	142.93	131.73	139.33	138.09	151.32	196.1	154.75
	sd	7.5543	6.9557	6.3442	6.7944	10.108	9.7206	9.8015	14.478	69.227
	n	30	28	29	30	28	29	28	16	3
SD RR (ms)	mean	1.91	1.616	1.8154	1.5356	2.1781	1.802	2.2878	5.1416	29.159
	sd	0.5472	0.5025	0.604	0.6912	2.306	0.8568	1.3248	3.585	40.273
Frequency Domain										
	n	30	28	29	30	28	29	28	16	3
Total Power PSD (ms^2)	mean	1.7269	1.2817	1.6601	1.3501	10.286	2.2217	5.43	19.707	44.669
	sd	1.0202	0.7197	1.0052	1.6521	34.37	2.6958	12.486	23.287	37.742
	n	30	28	29	30	28	29	28	16	3
VLF (%)	mean	8.5982	10.944	14.587	17.682	15.438	15.317	18.881	29.891	40.329
	sd	4.6037	8.8516	13.031	12.701	10.15	10.48	10.178	15.225	30.462
	n	30	28	29	30	28	29	28	16	3
LF (%)	mean	8.9183	10.891	12.756	11.72	14.401	16.543	13.281	21.088	65.082
	sd	5.8251	6.2324	9.6027	7.3111	12.266	10.575	11.947	13.919	31.832
	n	30	28	29	30	28	29	28	16	3
HF (%)	mean	77.513	72.832	65.299	62.606	59.341	53.457	48.169	26.461	47.794
	sd	10.752	15.446	20.821	20.871	22.597	22.212	22.361	20.61	45.237
	n	30	28	29	30	28	29	28	16	3
LF_TO_HF Ratio	mean	0.1327	0.1759	0.2616	0.2495	0.2704	0.3572	0.5631	2.1338	27.815
	sd	0.1024	0.1239	0.2192	0.179	0.2061	0.2518	1.2613	4.2574	41.361
	n	30	28	29	30	28	29	28	16	3
β (N.U.)	mean	-1.383	-1.315	-1.233	-1.371	-1.325	-1.231	-1.289	-1.282	24.405
	sd	0.22	0.2014	0.3093	0.2907	0.3397	0.3892	0.3999	0.5426	44.214
Non-linear Domain										
	n	30	28	29	30	28	29	28	16	3
SD1 (ms)	mean	2.0242	1.7053	1.8506	1.4973	2.1036	1.6488	2.0917	3.7887	27.904
	sd	0.6206	0.5874	0.7486	0.8172	2.4967	0.8243	1.4499	3.3762	41.252
	n	30	28	29	30	28	29	28	16	3
SD2 (ms)	mean	1.769	1.5027	1.7363	1.5254	2.0875	1.861	2.3579	5.7731	30.013
	sd	0.4923	0.4351	0.5198	0.6275	1.9782	0.9276	1.2871	3.9549	39.644
	n	30	28	29	30	28	29	28	16	3
SD Ratio	mean	1.1393	1.114	1.0434	0.9656	0.9819	0.8964	0.8476	0.6729	0.7431
	sd	0.1411	0.2005	0.2751	0.2686	0.2441	0.2751	0.2726	0.3452	0.2346
	n	30	28	29	30	28	29	28	16	3
α$_1$ (N.U.)	mean	0.267	0.3123	0.3605	0.3713	0.3633	0.4176	0.3727	0.531	25.642
	sd	0.1011	0.1207	0.166	0.1412	0.1132	0.1439	0.1956	0.3097	43.143
	n	30	28	29	30	28	29	28	16	3
α$_2$ (N.U.)	mean	0.5837	0.6022	0.6663	0.6865	0.6718	0.6429	0.7434	0.8097	25.578
	sd	0.1465	0.1759	0.203	0.2273	0.1965	0.1916	0.1883	0.2572	43.198
	n	30	28	29	30	28	29	28	16	3
Sample Entropy (N.U.)	mean	1.3289	1.3638	1.3767	1.4454	1.447	1.4457	1.332	1.0989	25.79
	sd	0.2012	0.2971	0.2781	0.3429	0.3453	0.2827	0.3164	0.4287	43.016

Heart Rate Fragmentation

	n	30	28	29	30	28	29	28	16	3
PIP (%)	mean	56.331	59.513	61.148	59.923	60.791	59.672	59.179	59.168	60.231
	sd	4.8508	5.1295	4.6558	4.3934	4.9786	4.3829	5.2608	5.1327	17.504
	n	30	28	29	30	28	29	28	16	3
IALS (N.U.)	mean	0.5645	0.5963	0.6128	0.6004	0.6092	0.598	0.5931	0.5935	25.505
	sd	0.0486	0.0513	0.0466	0.044	0.0499	0.0439	0.0527	0.0514	43.261
	n	30	28	29	30	28	29	28	16	3
PSS (%)	mean	66.12	73.606	75.775	74.967	74.546	69.684	62.799	63.905	60.173
	sd	11.908	13.117	10.695	9.0841	10.007	11.084	15.702	7.9964	21.741
	n	30	28	29	30	28	29	28	16	3
PAS (%)	mean	10.326	12.559	13.706	11.966	14.555	14.479	20.067	27.342	40.851
	sd	7.1928	10.623	11.989	9.0871	11.381	9.5201	10.646	7.7683	31.79

Supplement Table S1B. Descriptive statistics of HR and HRV parameters in Long Lived mice.

	Age	Basal State								
		6	9	12	15	18	21	24	27	30
Time Domain										
	n*	30	28	29	30	28	29	28	17	3
Mean RR (ms)	mean	135.8	135.7	140.8	136.3	136.2	138.4	139.6	151.2	147.3
	sd	8.234	9.498	10.56	11.23	13.63	12.74	14.19	15.59	22.16
	n	30	28	29	30	28	29	28	17	3
SD RR (ms)	mean	2.876	3.146	2.895	2.816	2.83	3.159	3.697	4.504	4.826
	sd	1.128	2.352	1.186	1.523	1.777	2.009	2.86	3.176	5.607
Frequency Domain										
	n	30	28	29	30	28	29	28	17	3
Total Power PSD	mean	4.224	6.77	4.571	4.578	5.15	6.646	11.09	15.46	29.69
(ms^2)	sd	3.053	14.23	3.708	5.816	9.062	9.796	20.68	19.38	48.39
	n	30	28	29	30	28	29	28	17	3
VLF (%)	mean	28.06	33.73	33.41	35.48	36.04	33.74	36.29	26.1	25.86
	sd	11.38	13.6	14.43	14.2	17.82	16.9	17.74	15.79	1.46
	n	30	28	29	30	28	29	28	17	3
LF (%)	mean	8.607	7.895	10.63	7.737	8.464	13.53	11.7	26.54	51.58
	sd	7.211	4.228	9.871	5.082	7.682	12.78	10.15	15.2	18.1
	n	30	28	29	30	28	29	28	17	3
HF (%)	mean	47.77	41.38	41.24	35.73	37.9	34.9	34.32	42.16	18.32
	sd	16.12	17.7	22.16	20.53	21.96	19.66	23.73	27.57	8.872
	n	30	28	29	30	28	29	28	17	3
LF_TO_HF Ratio	mean	0.207	0.329	0.427	0.561	0.345	0.502	0.509	1.41	3.731
	sd	0.22	0.525	0.654	1.483	0.366	0.543	0.534	1.651	3.04
	n	30	28	29	30	28	29	28	17	3
β (N.U.)	mean	-1.64	-1.66	-1.5	-1.71	-1.67	-1.46	-1.35	-1.06	-0.97
	sd	0.197	0.234	0.353	0.168	0.251	0.504	0.529	0.463	0.722
Non-linear Domain										
	n	30	28	29	30	28	29	28	17	3
SD1 (ms)	mean	2.261	2.18	2.218	2.014	2.069	2.063	2.738	3.591	2.493
	sd	0.964	1.148	1.204	1.56	1.694	1.176	3.013	3.033	2.646
	n	30	28	29	30	28	29	28	17	3
SD2 (ms)	mean	3.317	3.781	3.324	3.309	3.32	3.827	4.238	4.892	6.338
	sd	1.364	3.225	1.372	1.709	2.024	2.664	2.959	3.723	7.468
	n	30	28	29	30	28	29	28	17	3
SD Ratio	mean	0.703	0.649	0.68	0.596	0.632	0.591	0.597	0.81	0.442
	sd	0.195	0.229	0.291	0.265	0.276	0.254	0.331	0.412	0.064
	n	30	28	29	30	28	29	28	17	3
α_1 (N.U.)	mean	0.346	0.394	0.443	0.427	0.4	0.501	0.496	0.599	0.975
	sd	0.147	0.224	0.235	0.245	0.205	0.206	0.247	0.3	0.285
	n	30	28	29	30	28	29	28	17	3
α_2 (N.U.)	mean	1.008	1.097	1.041	1.096	1.088	1.035	1.081	0.742	0.616
	sd	0.203	0.13	0.2	0.216	0.257	0.268	0.271	0.255	0.155
	n	30	28	29	30	28	29	28	17	3
Sample Entropy	mean	1.468	1.449	1.436	1.389	1.412	1.402	1.34	1.328	1.353
(N.U.)	sd	0.301	0.372	0.287	0.33	0.31	0.319	0.339	0.399	0.511

Heart Rate Fragmentation										
PIP (%)	n	30	28	29	30	28	29	28	17	3
	mean	56.14	58.85	60.29	61.08	60.49	60.08	56.8	56.9	45.24
	sd	4.899	4.479	3.975	4.816	5.397	4.215	5.473	7.813	3.703
IALS (N.U.)	n	30	28	29	30	28	29	28	17	3
	mean	0.563	0.59	0.604	0.612	0.606	0.602	0.569	0.57	0.454
	sd	0.049	0.045	0.04	0.048	0.054	0.042	0.055	0.078	0.037
PSS (%)	n	30	28	29	30	28	29	28	17	3
	mean	74.53	79.53	81.49	79.9	78.64	76.43	67.73	61.76	42.35
	sd	9.979	10.26	10.3	12.43	11.7	10.68	12.42	12.81	5.667
PAS (%)	n	30	28	29	30	28	29	28	17	3
	mean	11.16	10.86	10.29	14.83	14.18	12.95	15.9	23.88	10.42
	sd	9.803	9.033	8.032	10.37	13.53	7.534	11.49	12.48	5.318

Supplement Table S1C. Descriptive statistics of HR and HRV parameters in Long Lived mice.

	Basal - Intrinsic = Effect of Autonomic Input								
Age	6	9	12	15	18	21	24	27	30
Time Domain									
	30	28	29	30	28	29	28	16	3
Mean RR (ms)	-0.04	1.85	-2.18	4.57	-3.13	0.27	-11.70	-46.52	-7.42
	5.72	8.34	7.80	8.57	11.40	7.68	8.26	8.10	85.35
	30	28	29	30	28	29	28	16	3
SD RR (ms)	0.97	1.53	1.08	1.28	0.65	1.36	1.41	-0.72	-24.33
	1.04	2.18	0.94	1.09	1.64	1.58	2.22	4.39	34.89
Frequency Domain									
	30	28	29	30	28	29	28	16	3
Total Power PSD	2.50	5.49	2.91	3.23	-5.14	4.42	5.66	-4.91	-14.98
(ms^2)	2.84	14.07	3.27	4.58	33.22	8.54	10.83	30.19	35.61
	30	28	29	30	28	29	28	16	3
VLF (%)	19.46	22.78	18.82	17.80	20.60	18.43	17.41	-4.44	-14.47
	12.53	13.50	11.68	13.69	16.01	14.56	15.47	16.63	31.76
	30	28	29	30	28	29	28	16	3
LF (%)	-0.31	-3.00	-2.12	-3.98	-5.94	-3.01	-1.58	6.21	-13.50
	8.83	5.53	8.89	6.16	10.76	8.84	8.14	22.54	25.71
	30	28	29	30	28	29	28	16	3
HF (%)	-29.74	-31.46	-24.06	-26.88	-21.44	-18.56	-13.85	18.07	-29.47
	20.64	17.23	20.16	22.50	26.98	26.14	28.10	22.61	53.10
	30	28	29	30	28	29	28	16	3
LF_TO_HF Ratio	0.07	0.15	0.17	0.31	0.07	0.14	-0.05	-1.05	-24.08
	0.24	0.49	0.59	1.38	0.34	0.47	1.16	4.56	38.54
	30	28	29	30	28	29	28	16	3
β (N.U.)	-0.261	-0.341	-0.268	-0.336	-0.348	-0.228	-0.058	0.228	-25.378
	0.294	0.307	0.370	0.303	0.346	0.392	0.550	0.687	43.528
Non-linear Domain									
	30	28	29	30	28	29	28	16	3
SD1 (ms)	0.236	0.474	0.367	0.517	-0.034	0.414	0.646	-0.104	-25.411
	0.884	0.859	0.848	0.955	1.352	0.854	2.627	3.173	38.714
	30	28	29	30	28	29	28	16	3
SD2 (ms)	1.548	2.278	1.588	1.784	1.233	1.966	1.880	-1.073	-23.675
	1.284	3.085	1.182	1.365	1.792	2.181	2.172	5.404	32.533
	30	28	29	30	28	29	28	16	3
SD Ratio	-0.437	-0.465	-0.363	-0.370	-0.350	-0.306	-0.251	0.171	-0.301
	0.239	0.209	0.249	0.256	0.306	0.311	0.335	0.357	0.295
	30	28	29	30	28	29	28	16	3
α$_1$ (N.U.)	0.079	0.082	0.082	0.056	0.037	0.084	0.123	0.031	-24.667
	0.176	0.208	0.182	0.191	0.181	0.206	0.217	0.401	42.873
	30	28	29	30	28	29	28	16	3
α$_2$ (N.U.)	0.424	0.495	0.374	0.410	0.416	0.392	0.337	-0.062	-24.962
	0.226	0.203	0.169	0.282	0.290	0.238	0.262	0.248	43.302
	30	28	29	30	28	29	28	16	3
Sample Entropy	0.14	0.09	0.06	-0.06	-0.03	-0.04	-0.01	0.26	-24.44
(N.U.)	0.23	0.36	0.30	0.38	0.36	0.32	0.34	0.60	43.53

Heart Rate Fragmentation

	30	28	29	30	28	29	28	16	3
PIP (%)	-0.19	-0.66	-0.86	1.15	-0.30	0.41	-2.38	-1.42	-14.99
	4.35	4.66	4.98	4.59	4.85	4.71	5.57	9.61	20.88
	30	28	29	30	28	29	28	16	3
IALS (N.U.)	-0.002	-0.007	-0.009	0.012	-0.003	0.004	-0.024	-0.015	-25.051
	0.044	0.047	0.050	0.046	0.049	0.047	0.056	0.096	43.275
	30	28	29	30	28	29	28	16	3
PSS (%)	8.41	5.92	5.71	4.93	4.10	6.75	4.93	-0.72	-17.82
	13.55	11.16	9.88	7.60	10.75	10.72	13.61	16.80	26.93
	30	28	29	30	28	29	28	16	3
PAS (%)	0.84	-1.70	-3.41	2.87	-0.37	-1.53	-4.17	-2.83	-30.43
	9.14	11.92	11.05	10.11	10.63	10.13	10.07	14.11	33.47

Supplement Table S2. Statistical Analyses (*) the effects of age on Basal, Intrinsic, and Autonomic Input on Intrinsic RR interval mean and RR interval variability in long lived mice.

	Intrinsic State		Basal State		Effect of Autonomic Input on Intrinsic	
Age (months)	6 to 21	21 to 30	6 to 21	21 to 30	6 to 21	21 to 30
Body Weight (g)			< 2.2e-16 ↑	3.5e-15 ↓		
Time Domain						
Mean RR (ms)	1.51E-06 ⌢	< 2.2e-16 ↑		0.01172 ↑	0.0008 ⌢	< 2.2e-16 ↑
SD RR (ms)		3.22e-6 ↑				0.0748 ↑
Frequency Domain						
Total Power PSD (ms^2)		0.0005 ↑		0.0984 ↑		
VLF (%)	0.0008 ↑	0.00577 ↑				1.31e-5 ↑
LF (%)	0.0021 ↑	0.0034 ↑	0.0385 ↑	1.04e-6 ↑		0.0576 ↓↑
HF (%)	7.33e-7 ↓	4.35e-5 ↓	0.0657 ↓↑			4.35e-05 ↓
LF TO HF Ratio	9.00e-6 ↑	0.0047 ↑		3.97e-6 ↑		
β (N.U.)			0.0071 ↑↓	0.05417 ↑		0.0316 ↓
Non-linear Domain						
SD1 (ms)		0.0044 ↑				
SD2 (ms)		5.39e-7 ↑				0.0317 ↑
SDRatio	3.95e-5 ↓	0.0253 ↓		0.0772 ⌢		7.31e-5 ↓
α$_1$ (N.U.)	1.77e-5 ↑	0.0059 ↑	0.0355 ↑	0.0120 ↑		
α$_2$ (N.U.)		0.05697 ↑		7.29e-5 ↓		1.90e-7 ↑
Sample Entropy (N.U.)		0.00681 ↓				0.0793 ↓↑
Heart Rate Fragmentation						
PIP (%)	0.0002 ↑→		6.15e-5 ↑	0.0002 ↓		0.0444 ↑
IALS (N.U.)	0.0002 ↑→		6.22e-5 ↑	0.0002 ↓		0.0429↑
PSS (%)	0.0007 ↑↓	0.0318 ↓	0.0336 ↑↓	3.61e-6 ↓		0.0284 ↑
PAS (%)		0.00037 ↑		0.0053 ↑		

Notes:

* 1. p-values from repeated-measures mixed ANOVA models; 2. Effect of Autonomic Input on Intrinsic = Basal - Intrinsic in each mouse;
3. N.U. = no units. 4. ↑ variable is significantly increasing with age, ↓ variable is significantly decreasing with age, and ⌢ the variable is significantly oscillating.

Table S3. Correlations and p-values of Mean RR with additional variables excluded from those that clustered with the increased mean RR interval in Fig. 6.

	Mean RR	SDRR	Total Power	LF	LF/HF	β	SD1	SD2	α₁	Sample Entropy	PIP	IALS	PSS
Mean RR : r		0.2485	0.0963	0.0926	0.0893	-0.0534	0.2391	0.2430	-0.0258	0.1376	0.3537	0.3648	0.2809
Mean RR : p		0.1854	0.6129	0.6266	0.6390	0.7794	0.2031	0.1958	0.8922	0.4684	0.0552	0.0475	0.1327
SDRR : r	0.2587		0.5317	0.0957	-0.0175	0.3991	0.9566	0.9692	-0.1033	-0.4317	0.0001	0.1899	0.1279
SDRR : p	0.1675		0.0025	0.6149	0.9269	0.0289	0.0000	0.0000	0.5869	0.0172	0.9997	0.3148	0.5006
Total Power : r	0.1554	0.9314		0.4895	0.3407	0.2988	0.4144	0.4752	0.2520	-0.1372	0.0347	0.0505	0.0091
Total Power : p	0.4122	8.2E-14		0.0060	0.0654	0.1087	0.0228	0.0080	0.1792	0.4698	0.8556	0.7911	0.9619
LF : r	-0.1159	0.3861	0.5291		0.8305	0.5644	-0.0094	0.1119	0.7343	-0.1685	-0.0540	-0.0202	-0.1941
LF : p	0.5421	0.0351	0.0026		1.4E-08	0.0012	0.9608	0.5559	3.9E-06	0.3734	0.7767	0.9154	0.3041
LF/HF : r	0.0393	0.4427	0.6258	0.7216		0.3529	-0.2171	0.1055	0.9259	-0.1505	0.0260	0.0954	-0.1583
LF/HF : p	0.8367	0.0143	0.0002	6.8E-06		0.0558	0.2492	0.5789	2.3E-13	0.4272	0.8917	0.6162	0.4035
β : r	0.0106	0.5640	0.6156	0.6493	0.5574		0.3978	0.3573	0.3180	-0.4953	-0.0080	0.1423	0.0889
β : p	0.9558	0.0012	0.0003	0.0001	0.0014		0.0295	0.0526	0.0868	0.0054	0.9667	0.4533	0.6404
SD1 : r	0.1446	0.8441	0.6910	0.2142	0.0355	0.4208		0.8685	-0.2847	-0.3777	0.0165	0.1527	0.1578
SD1 : p	0.4458	4.6E-09	2.4E-05	0.2558	0.8521	0.0206		5.0E-10	0.1273	0.0396	0.9310	0.4204	0.4051
SD2 : r	0.2798	0.9502	0.9531	0.4783	0.6192	0.5750	0.6451		0.0171	-0.4415	-0.0331	0.2089	0.0913
SD2 : p	0.1343	8.9E-16	4.4E-16	0.0075	0.0003	0.0009	0.0001		0.9285	0.0146	0.8620	0.2680	0.6313
α₁ : r	0.0177	0.4036	0.5433	0.5485	0.8007	0.4574	-0.0352	0.5920		-0.1303	0.0024	0.0481	-0.0751
α₁ : p	0.9260	0.0270	0.0019	0.0017	1.1E-07	0.0110	0.8537	0.0006		0.4927	0.9899	0.8006	0.6931
Sample Entropy : r	0.1937	-0.6270	-0.5650	-0.3840	-0.3243	-0.6726	-0.5670	-0.5778	-0.3180		-0.0095	-0.1335	-0.2226
Sample Entropy : p	0.3050	0.0002	0.0011	0.0362	0.0804	4.7E-05	0.0011	0.0008	0.0868		0.9604	0.4818	0.2371
PIP : r	0.1205	-0.2876	-0.3709	-0.5315	-0.4529	-0.3071	-0.1704	-0.3343	-0.3228	0.3053		0.8606	0.7326
PIP : p	0.5258	0.1233	0.0436	0.0025	0.0120	0.0988	0.3679	0.0710	0.0819	0.1009		1.1E-09	4.2E-06
IALS : r	0.1227	-0.2867	-0.3701	-0.5316	-0.4529	-0.3070	-0.1697	-0.3334	-0.3229	0.3055	1.0000		0.7631
IALS : p	0.5184	0.1245	0.0441	0.0025	0.0120	0.0989	0.3701	0.0718	0.0818	0.1006	0.0000		9.4E-07
PSS : r	-0.1424	0.3475	0.3579	0.3711	0.4174	0.3070	0.3041	0.3144	0.2964	-0.2977	-0.6038	-0.6043	
PSS : p	0.4530	0.0599	0.0521	0.0435	0.0217	0.0989	0.1023	0.0907	0.1117	0.1101	0.0004	0.0004	

Upper triangle: contains the correlations in the early time period (before 21 months); lower triangle contains the correlations in the late time period (after 21 months).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A.

B.

Figure 6

Figure 7

Figure 8

Figure 9

Supplemental Figures

Figure S1

Figure S2

Figure S3