Hippocampal protein aggregation signatures fully distinguish pathogenic and wildtype *UBQLN2* in amyotrophic lateral sclerosis

Kyrah M. Thumbadoo[1,2], Birger V. Dieriks[2,3], Helen C. Murray[2,3], Molly E. V. Swanson[1,3], Ji Hun Yoo[2,4], Nasim F. Mehrabi[2,3], Clinton Turner[2,3,5], Michael Dragunow[2,4], Richard L. M. Faull[2,3], Maurice A. Curtis[2,3], Teepu Siddique[6], Christopher E. Shaw[2,7], Lyndal Henden[8], Kelly L. Williams[8], Garth A. Nicholson[8,9,10,11], Emma L. Scotter[1,2]

School of Biological Sciences, University of Auckland, Auckland, New Zealand

Centre for Brain Research, University of Auckland, Auckland, New Zealand

Department of Anatomy and Medical Imaging, University of Auckland, Auckland, New Zealand

Department of Pharmacology and Clinical Pharmacology, University of Auckland, Auckland, New Zealand

Department of Anatomical Pathology, LabPlus, Auckland City Hospital, Auckland, New Zealand

Departments of Neurology, Cell and Developmental Biology and Pathology, Northwestern University Feinberg School of Medicine, Chicago, USA

UK Dementia Research Institute Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, United Kingdom

Macquarie University Centre for Motor Neuron Disease Research, Macquarie Medical School, Faculty of Medicine, Health and Human Sciences, Macquarie University, Sydney, New South Wales, Australia

Northcott Neuroscience Laboratory, Australian and New Zealand Army Corps (ANZAC) Research Institute, Concord, New South Wales, Australia

Faculty of Medicine, University of Sydney, Sydney, New South Wales, Australia

Molecular Medicine Laboratory, Concord Repatriation General Hospital, Concord, New South Wales, Australia

Corresponding author:

Email, emma.scotter@auckland.ac.nz; Telephone,　+64 9 923 1350

Acknowledgements and funding

This publication is dedicated to the incredible patients and families who contribute to our research. We thank Marika Eszes at the Centre for Brain Research, University of Auckland, New Zealand; Nailah Siddique at the Northwestern University Feinberg School of Medicine, Chicago, USA; Sashika Selvaduncko and Claire Troakes at the London Neurodegenerative Diseases Brain Bank and Brains for Dementia; and the Neurological Foundation of New Zealand for their ongoing financial support of the Human Brain Bank. We also thank Fairlie Hinton and Dr. Catriona McLean at the Victorian Brain Bank, which is supported by The Florey Institute of Neuroscience and Mental Health, The Alfred and the Victorian Forensic Institute of Medicine and funded in part by Parkinson's Victoria, MND Victoria, FightMND and Yulgilbar Foundation. The imaging data reported in this paper were obtained at the Biomedical Imaging Research Unit (BIRU), operated by the Faculty of Medical and Health Sciences' Technical Services at the University of Auckland. KT was funded by a doctoral scholarship from Amelia Pais-Rodriguez and Marcus Gerbich. BVD and MD were funded by the Michael J Fox Foundation [Grant ID: 16420]. BVD was also funded by a Health Research Council Sir Charles Hercus Health Research Fellowship [21/034]. ELS was supported by Marsden FastStart and Rutherford Discovery Fellowship funding from the Royal Society of New Zealand [15-UOA-157, 15-UOA-003]. This work was also supported by grants from Motor Neuron Disease NZ, Freemasons Foundation of New Zealand, Matteo de Nora, Sir Thomas and Lady Duncan Trust and the Coker Family Trust (to MD), and PaR NZ Golfing. No funding body played any role in the design of the study, nor in the collection, analysis, or interpretation of data nor in writing the manuscript.

Authors' contributions

KMT, BVD, HCM, MEVS, JHY, NFM conducted or designed experiments; KMT, BVD, HCM, MEVS, ELS performed data analysis or designed analysis methods; CT conducted neuropathological diagnostics; MD provided study supervision and resourcing; RLMF, MAC, TS, CES coordinated the banking and use of human tissue for study; TS, CES, GAN identified the *UBQLN2* patients; KLW, LH performed relatedness analysis; KMT, BVD,

61   ELS wrote the manuscript; GAN, ELS conceived of the study. ELS designed the study. All

62   authors read, edited, and approved the final manuscript.

Abstract

Mutations in the *UBQLN2* gene cause X-linked dominant amyotrophic lateral sclerosis (ALS) and/or frontotemporal dementia (FTD) characterised by ubiquilin 2 aggregates in neurons of the motor cortex, hippocampus, and spinal cord. However, ubiquilin 2 neuropathology is also seen in sporadic and familial ALS or FTD cases not caused by *UBQLN2* mutations, particularly *C9orf72*-linked cases. This makes the mechanistic role of ubiquilin 2 mutations and the value of ubiquilin 2 pathology for predicting genotype unclear. Here we examine a cohort of 41 genotypically diverse ALS cases with or without FTD, including five cases with *UBQLN2* variants (resulting in p.S222G, p.P497H, p.P506S, and two cases with p.T487I). Using multiplexed (5-label) fluorescent immunohistochemistry, we mapped the co-localisation of ubiquilin 2 with phosphorylated TDP-43 (pTDP-43), dipeptide repeat aggregates, and p62, in the hippocampus of controls (n=5), or ALS with or without FTD in sporadic (n=20), unknown familial (n=3), *SOD1*-linked (n=1), *FUS*-linked (n=1), *C9orf72*-linked (n=5), and *UBQLN2*-linked (n=5) cases. We differentiate between i) ubiquilin 2 aggregation together with pTDP-43 or dipeptide repeat proteins, and ii) ubiquilin 2 self-aggregation promoted by *UBQLN2* gene mutations that cause ALS/FTD. Overall, we describe a hippocampal protein aggregation signature that fully distinguishes mutant from wildtype ubiquilin 2 in ALS with or without FTD, whereby mutant ubiquilin 2 is more prone than wildtype to aggregate independently of driving factors. This neuropathological signature can be used to assess the pathogenicity of *UBQLN2* gene variants and to understand the mechanisms of *UBQLN2*-linked disease.

Keywords: *UBQLN2*, ubiquilin 2, amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), neuropathology, hippocampus, human, TDP-43, polyGP, polyGA

Introduction

UBQLN2 gene mutations are a rare cause of amyotrophic lateral sclerosis (ALS) with or without dementia, and the only known causal mutations on the X-chromosome. *UBQLN2* [NM_013444] encodes the ubiquilin 2 protein, the best studied of five human ubiquilins, which contains a unique PXX domain comprising 12 proline-rich tandem repeats [1,2]. Like other ALS- and FTD-linked degradation proteins (sequestosome 1/p62 (hereafter p62), valosin-containing protein, optineurin, and TANK-binding kinase 1), a key role for ubiquilin 2 is to bind ubiquitinated, misfolded, and aggregated protein cargos, triaging them between the proteasome and autophagy intracellular degradation pathways [3–6]. New evidence suggests that ubiquilin 2 is intrinsically prone to self-assembly and undergoes reversible liquid-liquid phase separation, allowing it to bind ubiquitin-labelled proteins within phase-separated stress granules. It is postulated that upon ubiquitinated protein binding, ubiquilin 2 comes out of the liquid-droplet phase, bringing its cargo from the stress granule for delivery to the proteasome; ubiquilin 2 is therefore also implicated in stress granule disassembly [7–10].

These roles at the interface between RNA processing and protein degradation — two key pathways in ALS and FTD pathogenesis — may underpin why ubiquilin 2 mutations cause disease. ALS and/or FTD (ALS/FTD)-causing *UBQLN2* mutations that alter residues flanking and within the PXX domain impair ubiquilin 2 binding to the proteasome and promote self-oligomerisation and liquid-to-solid rather than liquid-liquid phase transition [9,11–14]. The mutation c.1490C>A, resulting in p.P497H, was the first disease-causing *UBQLN2* mutation identified, leading to the discovery in unrelated ALS/FTD families of mutations c.1489C>T (p.P497S), c.1516C>A (p.P506T), c.1525C>T (p.P509S) and c.1573C>T (p.P525S) all altering the PXX domain [1]. A c.1516C>T mutation, resulting in p.P506S also within the PXX domain, was later identified in a family with young onset ALS, ALS with FTD onset, or pure spastic paraplegia [15] while a c.1460C>T mutation, resulting in p.T487I just upstream of the PXX region, was found in two families with members in New Zealand and Australia [16,17]. *UBQLN2* mutations/variants have also been identified in ALS/FTD cases altering other residues within or flanking the PXX domain; in the ubiquitin-like domain; the stress-induced protein 1-like domains; or outside

of known domains [18–37]. However, it is currently uncertain whether all of these
UBQLN2 variants are pathogenic (disease-causing).

The neuropathology of ALS/FTD caused by pathogenic *UBQLN2* mutations is
characterised by aggregated ubiquilin 2 in the motor cortex, spinal cord, and hippocampus
[1,15,17,23,38]. We previously reported ubiquilin 2-positive aggregates in the
hippocampus of an individual with p.T487I *UBQLN2*-linked ALS+FTD, but not in
sporadic ALS [38]. However, ubiquilin 2 inclusions are not specific to *UBQLN2*-linked
ALS/FTD cases and have been identified previously in sporadic and familial ALS, and
ALS-dementia, regardless of whether ubiquilin 2 is wildtype or mutant [1,15,17,23,38,39].
Some of these ubiquilin 2 inclusions are immunopositive for other ALS/FTD-linked
proteins such as TDP-43 or phosphorylated TDP-43 (pTDP-43), FUS, p62, optineurin, or
ubiquitin [1,17,40–44]. In addition, deposition of ubiquilin 2 in the hippocampus is a
characteristic feature of ALS/FTD caused by *C9orf72* hexanucleotide repeat expansions,
in which ubiquilin 2 is seen in the presence of dipeptide repeat (DPR) proteins [39]. Thus,
while ubiquilin 2 is clearly involved in ALS/FTD pathogenesis regardless of aetiology, the
role of *UBQLN2* mutations and the predictive value of ubiquilin 2 labelling in relation to
genotype is still unclear.

There is a clear need to determine whether *UBQLN2* mutations cause ubiquilin 2
neuropathology that is distinct from the wildtype ubiquilin 2 neuropathology that is seen
in ALS/FTD with other genotypes. If they do, then defining a pathological signature for
UBQLN2-linked disease will provide mechanistic insights and aid assessment of the
pathogenicity of *UBQLN2* genetic variants which are not yet confirmed to be causative.
Because ubiquilin 2 aggregation across a range of ALS/FTD genotypes occurs in the
hippocampus, this region may provide insight into the requirements for mutant and
wildtype ubiquilin 2 aggregation. Here we map the hippocampal ubiquilin 2 protein
deposition signature with respect to pTDP-43, the two most abundant *C9orf72*-linked DPR
proteins, and p62, in ALS/FTD with and without *UBQLN2* mutation.

Materials and methods

Systematic review of *UBQLN2*-linked ALS/FTD neuropathology

Journal articles were identified using PubMed that were published between Jan 1993 and May 2023. Search terms were *UBQLN2*, ubiquilin 2, amyotrophic lateral sclerosis, and ALS, combined with neuropathology, tissue, or immunohistochemistry. Articles which did not contain ubiquilin 2 neuropathological information in post-mortem ALS/FTD human brain tissue, were not primary research articles, or were not published in English were excluded. Seven papers were identified from which neuropathological data for TDP-43, ubiquilin 2, ubiquitin, p62, *C9orf72*-linked DPR proteins, FUS, and SOD1 protein aggregates in the spinal cord and hippocampus were extracted and tabulated.

Patient demographics and hippocampal brain tissue

Formalin-fixed paraffin-embedded (FFPE) post-mortem hippocampal tissue from 6 neurologically normal and 30 ALS cases with or without FTD, processed as previously described [45] were obtained from the Neurological Foundation Human Brain Bank at the Centre for Brain Research, Auckland, New Zealand. These included twenty cases with sporadic ALS (one with co-morbid FTD), three with familial ALS of unknown genotype, one with *SOD1*-linked ALS (p.E101G), five with *C9orf72*-linked ALS, and one with ALS+FTD with the *UBQLN2* p.T487I mutation (pedigree ID FALS5 IV:18 in Fig. 1A of [17] and in this report; coded MN17 in Figure 1D of [38] and in this report), giving a total cohort of 41 cases. Note that MN17 was stored in fixative for 7 years before embedding. All non-*SOD1*-linked NZ cases had confirmed pTDP-43 proteinopathy in the motor cortex. FFPE hippocampal tissue was also obtained from: the Victoria Brain Bank from a relative of the *UBQLN2* p.T487I case (p.T487I, ALS+FTD; pedigree ID V:7 in Fig. 1A of [17] and in this report); the London Neurodegenerative Diseases Brain Bank and Brains for Dementia from an unrelated *UBQLN2*-linked case (p.P506S, ALS+FTD:, pedigree ID II.2 in Fig. 1B of [15]), and a case with a *UBQLN2* variant of unknown significance (p.S222G, progressive supranuclear palsy; Supplemental_variant_data of [46]), and a *FUS*-linked case (p.P525L, ALS); and from the Northwestern University Feinberg School of Medicine, USA from another unrelated *UBQLN2*-linked case (p.P497H, ALS; pedigree ID V:I of Family #186 in Fig. 1A of [1]). All clinical and neuropathological diagnoses were

183   conducted as described previously [1,15,17,38]. Patient demographic and clinical
184   information is summarised in Supplementary Table 1.

Fluorescent immunohistochemistry and image acquisition

Multiplex fluorescent immunohistochemistry

Multiplex (five-label) immunohistochemistry was performed as described previously [47–
49]. Briefly, tissue sections were cut with a microtome in the transverse plane at a thickness
of 7-10 μm and mounted onto Superfrost Plus slides (Thermo Fisher Scientific). Mounted
sections were dried at room temperature for a minimum of 1 week before
immunohistochemistry. Slides were heated to 60 °C for 1 h on a hot plate, then dewaxed
and rehydrated through a xylene-alcohol-water series: 100% xylene, 2x 30 min; 100%
ethanol, 2x 15 min; 95%, 85%, 75% ethanol, 5 min each; water 3x 5 min. Antigens were
retrieved through immersion in 10 mM sodium citrate buffer (0.05% Tween 20, pH 6.0) in
a pressure cooker (Retriever 2100, Electron Microscopy Sciences) at 120 °C for 20 min
and cooled to room temperature for 100 min. Sections were washed 3x 5 min in 1x
phosphate-buffered saline (PBS) and wax borders were drawn with an ImmEdge
Hydrophobic Barrier PAP pen. Sections were permeabilised in PBS-T (PBS with 0.2%
Triton™ X100) for 15 min at 4 °C, followed by 3x 5-min PBS washes. Lipofuscin
autofluorescence was quenched using TrueBlack® Lipofuscin quencher (Biotium) 1:20 in
70% ethanol for at least 30 seconds followed by three vigorous water washes. Sections
were blocked with 10% normal donkey serum (Thermo Fisher Scientific) in PBS for 1 h
before 4 °C overnight incubation with primary antibodies targeting ubiquilin 2,
phosphorylated TDP-43 (pTDP-43), p62, and *C9orf72*-linked dipeptide repeat proteins
polyGA and polyGP (Supplementary Table 2). Following PBS washes, species- and
isotype-specific secondary antibodies and Hoechst 33342 nuclear stain (Supplementary
Table 3) were applied for 3 h in 1% normal donkey serum at room temperature. After final
3x 5-min PBS washes, sections were coverslipped with #1.5 coverslips (Menzel-Gläser)
using ProLong™ Gold Antifade Mountant (Thermo Fisher Scientific). All primary
antibodies used in this study have been previously validated in human brain tissue
[6,38,50,51], and secondary antibodies showed no non-specific binding or signal bleed-

through into adjacent channels (Supplementary Fig. 1) or cross-reactivity (Supplementary
Fig. 2).

Multiplex (five-label) images were acquired using a Zeiss Z2 Axioimager with a
MetaSystems VSlide slide scanning microscope (20x dry magnification lens, 0.9 NA) with
a Colibri 7 solid-state fluorescent light source. Sections were imaged with the same
exposure time and gain settings for each staining combination where possible. *UBQLN2*-
linked ALS/FTD case V:7-p.T487I showed consistently poor Hoechst immunoreactivity,
while *UBQLN2*-linked ALS/FTD case MN17-p.T487I showed poor immunoreactivity
overall, likely due to long-term fixation, therefore a longer exposure was used for both
cases when imaging. Single filters, as described and validated previously [49], were used
to excite fluorophores and detect the following wavelengths: Filter set #1 (LED 385; Em
447/60 nm), #3 (LED 475; Em 527/20 nm), #4 (LED 555; Em 580/23 nm), #5 (LED 590;
Em 628/32 nm), #6 (LED 630; Em 676/29 nm), and #7 (LED 735; Em 809/81 nm) and
visualised with MetaFer software (MetaSystems, v.3.12.1) equipped with a CoolCube 4m
TEC (monochrome) sCMOS digital camera. Image tiles were seamlessly stitched using
MetaCyte software, and stitched images were extracted using VSViewer software
(MetaSystems, v.1.1.106).

Double-label immunohistochemistry

For super-resolution stimulated emission depleted (STED) microscopy of *C9orf72*-linked
ALS case MN28, double-label immunohistochemistry was performed.
Immunohistochemistry was performed as above with primary antibody targeting ubiquilin
2 and detected by goat anti-mouse IgG_{2a} Alexa Fluor 594, but with primary antibody
targeting polyGA detected by goat anti-mouse IgG_1 Biotin (A10519, Thermo Fisher
Scientific), followed by 3x 5-min PBS washes, and an additional incubation with Star Red
neutravidin (STRED-0121, Abberior) for 1 hour at room temperature. After final 3x 5-min
PBS washes, sections were coverslipped with #1.5 coverslips using ProLong™ Glass
Antifade Mountant (P36984, Thermo Fisher Scientific).

STED images were acquired using an Abberior Facility STED microscope (60x
UPLXAPO oil immersion lens, 1.42 NA) using ImSpector Lightbox software (Specim,
v.16.3.13779). A 561-nm pulsed diode laser was used to excite Alexa Fluor 594, and a
640-nm diode laser was used to excite Alexa Fluor 647. For STED imaging, a pulsed 775-
nm laser was used for depletion of both fluorophores. After scanning, the images were
processed using the PureDenoise plugin [52] for ImageJ (National Institutes of Health,
USA v1.53f51).

Final figures of multiplex and double-label immunohistochemistry were compiled using
Adobe Photoshop CC (Adobe Systems Incorporated, v24.4.1). For clarity, multiplex labels
are shown in separate panels, either pTDP-43, ubiquilin 2, and p62, or ubiquilin 2, polyGA,
and polyGP, with negative labelling not shown.

Relatedness analysis

To further examine the p.T487I mutation in ALS/FTD, genome-wide genotype data from
the family of *UBQLN2*-linked ALS+FTD case MN17 (FALS5) and another Australian
family with an identical p.T487I mutation (FALS14) were analysed to determine whether
they inherited the mutation from a common ancestor (pedigree in Supplementary Fig. 3,
sample information in Supplementary Table 4). These two families were previously
reported to share a haplotype identical-by-state over the *UBQLN2* locus [17], but genealogy
analysis had been unable to link the pedigrees. Three individuals from FALS5 (all affected,
pedigree IDs III:8, IV:9, IV:18 (MN17)) and two individuals from FALS14 (pedigree IDs
II:1 (unaffected) and III:2 (affected)) underwent SNP genotyping using the Illumina
Infinium Global Screening Array v2.0. Identity-by-descent (IBD) analysis was performed
using XIBD software [53] with the combined HapMap Phase II and III European (CEU)
cohort as a reference dataset. SNPs in high linkage disequilibrium ($r^2 > 0.95$) or SNPs with
a low minor allele frequency (MAF < 0.01) were removed from analysis, as well as SNPs
with missing genotype calls in two or more samples. 39,002 SNPs remained for analysis
and 211 IBD segments greater than 3 cM were identified. The degree of relationship was
estimated for each pair of samples using the lengths of inferred IBD segments as in
Estimation of Recent Shared Ancestry [54].

Results

Ubiquilin 2 labelling in previous studies failed to discriminate between *UBQLN2*-linked and other genotypes of ALS/FTD

Systematic review of the literature describing ubiquilin 2 neuropathology in human ALS/FTD identified 137 results, of which 7 articles met the criteria for full review. The hippocampal neuropathology of ubiquilin 2 with respect to five other ALS/FTD-linked proteins is summarised in Fig. 1. Ubiquilin 2 aggregate deposition in the hippocampal molecular layer was most frequently reported upon, being found in *UBQLN2*-linked and *C9orf72*-linked ALS/FTD, sometimes together with ubiquitin or p62 [6]. Ubiquilin 2 aggregate deposition in the granule cell layer of the hippocampal dentate gyrus (GCL) was inconsistent between cases of *UBQLN2*-linked ALS/FTD but was found in all cases of "ALS-dementia" in [1] (all later confirmed to be *C9orf72*-positive) and in a single sporadic ALS/FTD case [15]. Cornu ammonis (CA) ubiquilin 2 pathology had been reported in both *UBQLN2*-linked and *C9orf72*-linked ALS/FTD [1,38,39]. Overall, neither ubiquilin 2 staining alone nor in combinations previously tested could distinguish mutant ubiquilin 2 aggregation in *UBQLN2*-linked ALS/FTD from wildtype ubiquilin 2 aggregation in other ALS/FTD genotypes.

Ubiquilin 2 and other pathology in ALS/FTD of various genotypes

Control and familial ALS cases had no hippocampal ubiquilin 2 aggregates but certain cases had sparse p62-only aggregates

Five of the six non-neurodegenerative disease control cases included in this study showed no deposition of ubiquilin 2, pTDP-43, p62, or DPR protein aggregates in any region of the hippocampus (not shown). The remaining control case (H230) showed sparse p62 aggregates in the GCL and CA pyramidal cell layer (CA – pyr) (Fig. $1A_{1,2}$).

All five familial ALS cases also showed sparse p62 aggregates in the GCL and CA – pyr regions; three cases were of unknown genotype, represented by case MN14 (Fig. $1B_{1,2}$), one harboured a *SOD1* p.E101G mutation (Fig. $1C_{1,2}$), and one harboured a *FUS* p.P525L

mutation (Fig. 1D$_{1,2}$). No other pathology was observed in familial cases in hippocampal

regions.

Sporadic ALS cases had no hippocampal ubiquilin 2 aggregates but certain cases had

pTDP-43-p62 aggregates

None of the 20 sporadic ALS cases (one with co-morbid FTD) showed hippocampal

ubiquilin 2 or DPR aggregates (not shown). Fourteen cases (70%) were devoid of either

ubiquilin 2 or pTDP-43 in the GCL, ML, CA-pyr, and CA – lm/rad layers (not shown),

however 4 cases (20%) had compact p62-positive NCIs in the GCL and CA – pyr regions

that were negative for all other markers (Supplementary Fig. 4). Six cases (30%; including

representative cases MN15 with ALS+FTD, and MN30 with ALS) had compact or net-like

perinuclear pTDP-43-positive neuronal cytoplasmic inclusions (NCIs) in the GCL and CA

– pyr regions ranging in density from sparse to frequent, suggestive of stage 4 pTDP-43

proteinopathy [55]. These pTDP-43 aggregates were almost always p62-positive and

ubiquilin 2-negative (Fig. 2E$_{1,2}$ and F$_{1,2}$, white arrows), with the exception of 1-2

perinuclear aggregates that were not p62-labelled (Fig. 2F$_1$, yellow arrow).

C9orf72 cases had molecular layer ubiquilin 2-only aggregates, granule cell layer

ubiquilin 2-DPR-p62 aggregates, and one case had granule cell layer ubiquilin 2-DPR-

pTDP-43-p62 aggregates

Four of the five *C9orf72*-linked ALS cases showed identical hippocampal pathology

(represented by case MN18), while the fifth case (MN28) differed only by having pTDP-

43 aggregates in the GCL and CA – pyr, described below.

All five cases shared a signature of numerous ubiquilin 2 aggregates in the ML, GCL and

CA – l-m/rad regions. In the ML and CA – l-m/rad, ubiquilin 2 aggregates were punctate

or skein-like, dendritic, and very rarely co-localised with p62 (Fig. 3A$_{1,2}$ and B$_{1,2}$, green

arrowheads) and never with DPR proteins (Fig. 3C$_{1,2}$ and D$_{1,2}$, green arrowheads). In the

CA – pyr, aggregates did not contain ubiquilin 2 (Fig. 3A$_3$-D$_3$) but were positive for p62

(Fig. 3A$_3$, B$_3$, pink arrows), polyGA, and polyGP (Fig 3C$_3$, D$_3$, orange arrows and insets),

with pTDP-43 in case MN28 also seen rarely (Fig. 3B$_3$, yellow arrow).

In the GCL, ubiquilin 2 aggregates were perinuclear NCIs that were negative for pTDP-43

except rare aggregates in case MN28 (Fig. 3B$_4$, yellow arrow), but always positive for p62

(Fig. 3A$_1$, B$_1$, white arrows and yellow arrow) and poly(glycine-arginine) (polyGA), but

rarely polyGP (Fig 3C$_1$, D$_1$, yellow arrows). Ubiquilin 2 aggregates positive for pTDP-43

were rare (1-2 GCL cells per section of MN28) and comprised a pTDP-43 'shell'

surrounding polyGA either with polyGP (Supplementary Fig. 5A, white arrow) or without

polyGP (Supplementary Fig. 5A, purple arrow). This supports a previous report that

ubiquilin 2 in *C9orf72*-linked cases rarely co-localises with GCL pTDP-43, doing so only

when DPR proteins are present in the aggregate [56]. Therefore, wildtype ubiquilin 2

aggregation may be 'seeded' in the GCL by polyGA rather than by pTDP-43.

UBQLN2 cases show molecular layer ubiquilin 2-p62 aggregates and certain cases had

granule cell layer pTDP-43 aggregates

All four *UBQLN2*-linked ALS/FTD cases showed signature ubiquilin 2-positive but pTDP-

43-negative punctate aggregates in the hippocampal ML and CA – lm/rad regions, albeit

at variable loads (Fig. 4A$_{2,3}$, B$_{2,3}$, C$_{2,3}$, D$_{2,3}$, white arrows). In the ML, ubiquilin 2

aggregates were punctate and appeared to be localised to the dendritic spines of the granule

cells, as reported in *UBQLN2*-linked ALS/FTD and mutant *UBQLN2* rodent models [1,57].

In contrast to *C9orf72*-linked cases, these ML ubiquilin 2 aggregates almost always co-

localised with p62. The PSP case with a *UBQLN2* p.S222G variant of unknown

significance did not share this signature of ubiquilin 2- and p62-positive aggregates in the

ML and CA – lm/rad, instead having p62-only aggregates in the CA – lm/rad (Fig. 4E$_{2,3}$,

pink arrowheads).

Confocal microscopy of the ML and CA – lm/rad regions confirmed that the p62-positive

mutant ubiquilin 2 aggregates in *UBQLN2*-linked ALS/FTD cases (Supplementary Fig.

5A, representative p.P506S case shown) differed from wildtype ubiquilin 2 aggregates in

those regions in *C9orf72*-linked ALS, which were predominantly p62 negative

(Supplementary Fig. 5B, representative case MN28). Further, while mutant ubiquilin 2 in

UBQLN2-linked cases formed small compact aggregates (Supplementary Fig. 6A)

wildtype ubiquilin 2 in *C9orf72*-linked cases formed small compact aggregates and wispy

skein-like structures (Supplementary Fig. 6B). Therefore, mutant but not wildtype

ubiquilin 2 in the hippocampal ML and CA – lm/rad regions may promote or scaffold the

co-aggregation of p62, and there may be structural differences between mutant and

wildtype.

In the GCL, two of the four *UBQLN2*-linked ALS/FTD cases were devoid of pTDP-43,

ubiquilin 2, and p62 aggregates, consistent with previous findings [1,38] (case P497H, Fig.

4A$_1$; and MN17 – T487I, Fig. 4B$_1$). In contrast, case V:7 – T487I showed very sparse

pTDP-43 cytoplasmic aggregates in the GCL that were negative for ubiquilin 2 but weakly

positive for p62 (Fig. 4C$_1$, white hollow arrowhead). Consistent with Gkazi and colleagues

[15] and with the known spectrum of pTDP-43 loads in the GCL in ALS/FTD [55], the

UBQLN2 p.P506S case had abundant p62-positive pTDP-43 cytoplasmic aggregates in the

GCL, that either co-localised with ubiquilin 2 (Fig. 4D$_1$, white filled arrowheads) or were

independent of ubiquilin 2 (Fig. 4D$_1$, white hollow arrowheads). Mutant ubiquilin 2

aggregation in the GCL was therefore likely scaffolded or 'seeded' by pTDP-43 when

present. The *UBQLN2* p.S222G PSP case showed sparse p62-only aggregates in the GCL

that were not seen in *UBQLN2*-linked ALS/FTD cases (Fig. 4E$_1$, pink arrowheads).

UBQLN2-linked ALS/FTD case P506S also had sparse pTDP-43 aggregates in the CA –

pyr cells, which were positive for both ubiquilin 2 and p62 (Fig. 4D$_3$, white filled

arrowhead). All *UBQLN2* variant and mutation cases were negative for DPRs (not shown).

Combined neuropathological signatures discriminate between *UBQLN2*-linked,

***C9orf72*-linked, sporadic, and familial cases**

Integration of all neuropathological findings (Fig. 5) revealed a characteristic hippocampal

neuropathological signature for *UBQLN2*-linked ALS/FTD, that was distinct from that in

other forms of ALS/FTD. Sporadic ALS cases were wholly devoid of hippocampal

ubiquilin 2 or DPR protein pathology, with a minority of cases showing pTDP-43

aggregates in the granule cells that were ubiquilin 2 negative. Thus, wildtype ubiquilin 2

is not seeded/ scaffolded to aggregate by pTDP-43 aggregation. *C9orf72*-linked cases

showed ML ubiquilin 2 aggregates that were wispy or punctate and predominantly p62

negative, while *UBQLN2*-linked cases showed ML ubiquilin 2 aggregates that were

punctate and predominantly p62 positive. Mutant ubiquilin 2 aggregates thus promote the

co-aggregation of p62, and this may relate to their conformational differences. Overall,

mutant ubiquilin 2 causes unique neuropathology that is shared by p.P506S, p.P497H, and

p.T487I ubiquilin 2 (Fig. 6). The absence of this signature in a case harbouring a p.S222G

variant manifesting with PSP suggests this neuropathology can be used to discriminate

pathological ALS/FTD-causing *UBQLN2* mutations.

Ubiquilin 2 hippocampal pathology was present in both *UBQLN2*-linked ALS/FTD and

C9orf72-linked ALS, but these genotypes could be discriminated when ubiquilin 2 was co-

labelled with pTDP-43, DPR proteins, or p62. *C9orf72*-linked cases always showed GCL

ubiquilin 2 that co-localised with DPR aggregates, but rarely pTDP-43. This supports the

lack of seeding of wildtype ubiquilin 2 aggregation by pTDP-43 but suggests that wildtype

ubiquilin 2 can be seeded by polyGA. In contrast, *UBQLN2*-linked cases showed GCL

ubiquilin 2 if co-localised pTDP-43 was present. Therefore, mutant ubiquilin 2 is more

aggregation-prone than wildtype, being seeded by pTDP-43.

UBQLN2 p.T487I mutation in ALS/FTD families FALS5 and FALS14 was inherited

from a common ancestor

Since the initial report by Williams et al. [17] of an identical *UBQLN2* p.T487I mutation

in ALS families FALS5 and FALS14, we report here that cases MN17 (IV:18) and V:7

from family FALS5 developed ALS+FTD, indicating that FTD is part of the clinical

phenotype in that family. To further examine relatedness between the families and confirm

that the *UBQLN2* p.T487I mutation arose in a common founder, we performed identity-

by-descent (IBD) analysis. IBD segments were identified over the *UBQLN2* locus between

all four affected individuals from both families (Table 1), while there were no IBD

segments inferred over *UBQLN2* between the affected individuals from FALS5 and the

unaffected individual from FALS14 who did not carry the *UBQLN2* p.T487I mutation. The

interval shared by all four affected individuals spanned rs952836 to rs6423133 and is 68

cM in length. This confirms a founder effect of *UBQLN2* p.T487I in FALS5 and FALS14.

The genotyped individuals across these families are estimated to be 4th- to 5th-degree

relatives (1st cousins once removed - 2nd cousins) (Table 2), now confirming segregation

of the *UBQLN2* p.T487I mutation in 17 individuals from the proposed combined Australia-

New Zealand pedigree and providing further strong genetic evidence, in addition to the

neuropathological evidence, that *UBQLN2* p.T487I is pathogenic for ALS/FTD.

Discussion

ALS shows considerable clinical, pathological, and genetic heterogeneity [58–60]. While TDP-43 proteinopathy is seen in 97% of cases [40,43,61], ALS/FTD-causing genetic mutations can cause deposition of the encoded mutant protein leading to additional pathological aggregate signatures. We previously exploited this to infer *C9orf72* repeat expansion genotypes [62,63] through neuropathological screening of our New Zealand ALS cases [38]. Mutant *SOD1* [64,65], *FUS* [44], and certain other genotypes [66,67] can also be inferred from neuropathology. However, for many ALS/FTD genes — particularly those such as *TARDBP*, *SQSTM1*, and *UBQLN2* that encode proteins already within the hallmark TDP-43 inclusions — no completely discriminating neuropathology has been reported [42,66,68–72]. This has hampered the validation of pathogenicity of novel variants in these genes, in turn obscuring understanding of protein domains and molecular processes important to ALS/FTD pathogenesis. However, we report here a unique neuropathological signature for mutant ubiquilin 2 in the hippocampus that discriminates *UBQLN2*-linked ALS/FTD cases from all others tested.

pTDP-43 in *UBQLN2*-linked cases: An independent pathology

Before discussing the unique pattern of hippocampal ubiquilin 2 pathology shared by *UBQLN2*-linked cases, it must first be noted that pTDP-43 deposition was variable. Two of the four *UBQLN2*-linked cases were devoid of GCL pTDP-43 aggregates (p.P497H (ALS) and MN17-p.T487I (ALS+FTD)), one case had very sparse GCL pTDP-43 aggregates that were ubiquilin 2-negative (V:7-p.T487I (ALS+FTD)), and one case showed frequent GCL pTDP-43 aggregates that were mostly ubiquilin 2-positive (p.P506S (ALS+FTD)). Although pTDP-43 aggregates are nearly ubiquitous in the ALS spinal cord and motor cortex, only in ~15-30% of ALS cases are they found in the hippocampus [55,73]. Our findings suggest that variable regional pTDP-43 deposition occurs in the context of *UBQLN2*-linked ALS/FTD just as it does in sporadic and *C9orf72*-linked ALS/FTD.

Wildtype ubiquilin 2 pathology in *C9orf72*-linked cases

Hippocampal ubiquilin 2 deposition is a known and striking feature of *C9orf72*-linked ALS/FTD [39]. In our *C9orf72*-linked ALS cases, large stellate GCL ubiquilin 2 was found to preferentially co-localise with the aggregation-prone polyGA DPR protein compared to polyGP. Furthermore, polyGP aggregates were very rarely independent of polyGA. These findings support the emerging consensus that polyGA aggregates 'seed' polyGP protein aggregation [74–76]. Similar observations were made by Mackenzie and colleagues [56] of aggregates with a core of polyGA surrounded by an aggregated TDP-43 shell; a finding recapitulated here and supported by *in vitro* work showing polyGA aggregation preceding TDP-43 accumulation [77]. The ability of DPR proteins to seed ubiquilin 2 however, appears more complex. We previously showed ubiquilin 2 at the core of polyGP-positive aggregates, but we did not co-label for polyGA [38]. STED imaging in the current study shows that ubiquilin 2 may either surround polyGA or be enmeshed with it, suggesting that in *C9orf72*-linked ALS pathogenesis, the interaction between aggregating polyGA and ubiquilin 2 may be an early event.

In contrast to ubiquilin 2 co-aggregation with large stellate DPR proteins in the GCL, small neuritic ubiquilin 2 aggregates in *C9orf72*-linked cases punctuated the ML, and CA – lm/rad regions seemingly independent of a nucleating protein. The neurites in which these small aggregates are found likely derive from the granule or pyramidal cells, respectively [78,79], so DPR protein inclusions in the somata may promote the aggregation of ubiquilin 2 in the dendrites of the same cell. DPR aggregates can sequester proteasome components [80–82], and loss of C9orf72 protein function in cells expressing the DPR-encoding variant 2 [83] can impair autophagy [84–89], which may underpin wildtype ubiquilin 2 aggregation in the ML, and CA – lm/rad layers in *C9orf72*-linked cases.

Mutant ubiquilin 2 pathology in *UBQLN2*-linked cases: A neuropathological signature

Even in its wildtype state, ubiquilin 2 intrinsically self-assembles [3] but there is now ample biophysical evidence demonstrating that mutations to ubiquilin 2 confer an increased propensity to oligomerise and undergo aberrant LLPS, forming insoluble aggregates within

the cell [8,9,12,90–92]. Here we confirm that mutant ubiquilin 2 in the human hippocampal

ML (granule cell dendrites) is aggregated under less permissive conditions than wildtype

ubiquilin 2, appearing to require no aggregated protein scaffold, or protein aggregation

event in the GCL (granule cell soma).

Our study additionally found that p62 labelling was required, while pTDP-43 labelling was

dispensable, to confirm cases as being *UBQLN2*-linked. P62 co-localises with pTDP-43

aggregates and DPR proteins in ALS/FTD [62,93,94], or with hyperphosphorylated tau in

a range of tauopathies [95–97], or with mutant α-synuclein in synucleinopathies [97,98].

Given this promiscuity for substrates, discrimination between mutant and wildtype

ubiquilin 2 by p62 suggests that there are unique structural or biochemical features of

mutant ubiquilin 2 aggregates. This leads to the intriguing possibility that mutant ubiquilin

2 would be selectively druggable.

In addition to mechanistic insights, the mutant ubiquilin 2 neuropathological signature we

describe will enable classification of *UBQLN2* variants of uncertain significance, clarifying

the implications of a positive genetic result in a patient. Currently, only four missense

mutations in *UBQLN2* are designated by ClinVar as pathogenic or likely pathogenic

(resulting   in   p.M392V,   p.Q425R,   p.P497H,   p.P497S,   p.P506T,

https://www.ncbi.nlm.nih.gov/clinvar/, accessed July 11 2023 [99]). However, 62 other

UBQLN2 missense changes listed in ClinVar are classified as benign or of uncertain

significance, including c.1516C>T resulting in p.P506S. Indeed, neither *UBQLN2* p.T487I

nor p.S222G are even listed in ClinVar, leaving individual diagnostics labs to perform

classification for patient reporting. We now confirm that two ALS/FTD cases with p.T487I,

which segregates with disease in a large kindred and shows early onset in males consistent

with X-linked disease, shared our newly identified mutant ubiquilin 2 neuropathological

signature, supporting its classification as pathogenic. Conversely, a male PSP case with

p.S222G who lived to 93 y did not share this signature, suggesting this variant is benign.

We encourage uptake of this hippocampal ubiquilin 2 neuropathology signature, by other

labs or in collaboration with the authors hereof, as a tool to explore *UBQLN2* variant

pathogenicity.

Conclusion

Ubiquilin 2 aggregates were seen in the hippocampus of ALS/FTD cases across a range of genotypes. Wildtype ubiquilin 2 *in vitro* is known to be aggregation-prone; in human brain it co-aggregated with polyGA, but not pTDP-43 and with little co-aggregation of p62. Mutant ubiquilin 2 *in vitro* is known to be more aggregation-prone than wildtype; in brain it was either co-aggregated with pTDP-43 or aggregated independently of a known scaffold and appeared to promote the co-aggregation of p62. This hippocampal ubiquilin 2 neuropathology signature demonstrates that ubiquilin 2 aggregation is likely to play a mechanistic role in *C9orf72*-linked and *UBQLN2*-linked ALS/FTD and provides a definitive framework for exploring the biological implications of *UBQLN2* genetic variation.

List of abbreviations

524		
525	ALS	Amyotrophic lateral sclerosis
526	bvFTD	Behavioural variant FTD
527	CA-lm/rad	Cornu ammonis lacunosum-molecular and radiatum layers
528	CA-pyr	Cornu ammonis pyramidal layer
529	C9orf72	Chromosome 9 open reading frame 72
530	DPR	Dipeptide repeat
531	FFPE	Formalin-fixed paraffin-embedded
532	FTD	Frontotemporal dementia
533	GCL	Granule cell layer
534	IBD	Identity by descent
535	LLPS	Liquid-liquid phase separation
536	ML	Molecular layer
537	NCI	Neuronal cytoplasmic inclusions
538	PBS	Phosphate-buffered saline
539	PolyGA	Poly(glycine-arginine)
540	PolyGP	Poly(glycine-proline)
541	pTDP-43	Phosphorylated TDP-43
542	SOD1	Superoxide dismutase
543	STED	Stimulated emission depletion
544	SQSTM1	Sequestosome 1
545	TDP-43	Transactive response DNA binding protein 43 kDa

Disclosures and declarations

Data transparency

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Compliance with ethical standards

Conflicts of interest

The authors declare that they have no conflicts of interest.

Research involving human participants

All protocols were approved by the University of Auckland Human Participants Ethics Committee (New Zealand) and carried out as per approved guidelines. This study was also approved by the Human Research Ethics Committee of Macquarie University (520211013428875).

Informed consent

Informed donor consent and ethical approvals were obtained at each site as described previously [1,15,17,38].

References

1. Deng H-XX, Chen W, Hong S-TT, Boycott KM, Gorrie GH, Siddique N, et al. Mutations in UBQLN2 cause dominant X-linked juvenile and adult-onset ALS and ALS/dementia. Nature. 2011;477(7363):211–5. doi:10.1038/nature10353

2. Marín I. The ubiquilin gene family: Evolutionary patterns and functional insights. BMC Evol Biol. 2014;14(1):1–22. 10.1186/1471-2148-14-63

3. Hjerpe R, Bett JS, Keuss MJ, Solovyova A, McWilliams TG, Johnson C, et al. UBQLN2 Mediates Autophagy-Independent Protein Aggregate Clearance by the Proteasome. Cell. 2016;166(4):935–49. 10.1016/j.cell.2016.07.001

4. Itakura E, Zavodszky E, Shao S, Wohlever ML, Keenan RJ, Hegde RS. Ubiquilins Chaperone and Triage Mitochondrial Membrane Proteins for Degradation. Mol Cell. 2016;63(1):21–33. 10.1016/j.molcel.2016.05.020

5. Protter DSW, Parker R. Principles and Properties of Stress Granules. Trends Cell Biol. 2016;26(9):668–79. 10.1016/j.tcb.2016.05.004

6. Wu JJ, Cai A, Greenslade JE, Higgins NR, Fan C, Le NTT, et al. ALS/FTD mutations in UBQLN2 impede autophagy by reducing autophagosome acidification through loss of function. Proceedings of the National Academy of Sciences. 2020;2(Line 356):201917371.

7. Alexander EJ, Niaki AG, Zhang T, Sarkar J, Liu Y, Nirujogi RS, et al. Ubiquilin 2 modulates ALS/FTD-linked FUS–RNA complex dynamics and stress granule formation. Proc Natl Acad Sci U S A. 2018;115(49):E11485–94.

8. Dao TP, Kolaitis RM, Kim HJ, O'Donovan K, Martyniak B, Colicino E, et al. Ubiquitin Modulates Liquid-Liquid Phase Separation of UBQLN2 via Disruption of Multivalent Interactions. Mol Cell. 2018;69(6):965-978.e6.

9. Sharkey LM, Safren N, Pithadia AS, Gerson JE, Dulchavsky M, Fischer S, et al. Mutant UBQLN2 promotes toxicity by modulating intrinsic self-assembly. Proceedings of the National Academy of Sciences. 2018 Oct 30;115(44):E10495–504. 10.1073/pnas.1810522115

10. Subudhi I, Shorter J. Ubiquilin 2: Shuttling Clients Out of Phase? Mol Cell. 2018;69(6):919–21. 10.1016/j.molcel.2018.02.030

11.   Chang L, Monteiro MJ. Defective proteasome delivery of polyubiquitinated
proteins by ubiquilin-2 proteins containing ALS mutations. PLoS One.
2015;10(6):1–15. 10.1371/journal.pone.0130162

12.   Dao TP, Martyniak B, Canning AJ, Lei Y, Colicino EG, Cosgrove MS, et al.
ALS-Linked Mutations Affect UBQLN2 Oligomerization and Phase
Separation in a Position- and Amino Acid-Dependent Manner. Structure.
2019;27(6):937-951.e5.

13.   Gilpin KM, Chang L, Monteiro MJ. ALS-linked mutations in ubiquilin-2 or
hnRNPA1 reduce interaction between ubiquilin-2 and hnRNPA1. Hum Mol
Genet. 2015 May 1;24(9):2565–77. 10.1093/hmg/ddv020

14.   Higgins N, Lin B, Monteiro MJ. Lou Gehrig's Disease (ALS): UBQLN2
Mutations Strike Out of Phase. Structure. 2019;27(6):879–81.
10.1016/j.str.2019.05.006

15.   Gkazi SA, Troakes C, Topp S, Miller JW, Vance CA, Sreedharan J, et al.
Striking phenotypic variation in a family with the P506S UBQLN2 mutation
including amyotrophic lateral sclerosis, spastic paraplegia, and frontotemporal
dementia. Neurobiol Aging. 2019;73:229.e5-229.e9.
10.1016/j.neurobiolaging.2018.08.015

16.   McCann EP, Williams KL, Fifita JA, Tarr IS, O'Connor J, Rowe DB, et al. The
genotype-phenotype landscape of familial amyotrophic lateral sclerosis in
Australia. Clin Genet. 2017 Sep;92(3):259–66. 10.1111/cge.12973

17.   Williams KL, Warraich ST, Yang S, Solski JA, Fernando R, Rouleau GA, et
al. UBQLN2/ubiquilin 2 mutation and pathology in familial amyotrophic
lateral sclerosis. Neurobiol Aging. 2012;33(10):2527.e3-2527.e10.
10.1016/j.neurobiolaging.2012.05.008

18.   Chen J, Liu X, Xu Y, Fan D. Association between rare UBQLN2 variants and
amyotrophic lateral sclerosis in Chinese population. National Medical Journal
of China. 2021;101(12):846–50.

19.   Kim HJ, Kwon MJ, Choi WJ, Oh KW, Oh S il, Ki CS, et al. Mutations in
UBQLN2 and SIGMAR1 genes are rare in Korean patients with amyotrophic

lateral sclerosis. Neurobiol Aging. 2014;35(8):1957.e7-1957.e8.
10.1016/j.neurobiolaging.2014.03.001

20. Baglivo M, Manara E, Capodicasa N, Maltese PE, Stuppia L, Michelini S, et
al. Early-onset of Frontotemporal Dementia and Amyotrophic Lateral Sclerosis
in an Albanian Patient with a c.1319C>T Variant in the UBQLN2 Gene. Open
Med J. 2020;7(1):25–31. 10.2174/1874220302007010025

21. Daoud H, Suhail H, Szuto A, Camu W, Salachas F, Meininger V, et al.
UBQLN2 mutations are rare in French and French-Canadian amyotrophic
lateral sclerosis. Neurobiol Aging. 2012;33(9):2230.e1-2230.e5.
10.1016/j.neurobiolaging.2012.03.015

22. Dillen L, van Langenhove T, Engelborghs S, Vandenbulcke M, Sarafov S,
Tournev I, et al. Explorative genetic study of UBQLN2 and PFN1 in an
extended Flanders-Belgian cohort of frontotemporal lobar degeneration
patients. Neurobiol Aging. 2013;34(6):1711.e1-1711.e5.

23. Fahed AC, McDonough B, Gouvion CM, Newell KL, Dure LS, Bebin M, et al.
UBQLN2 mutation causing heterogeneous X-linked dominant
neurodegeneration. Ann Neurol. 2014;75(5):793–8.

24. Gellera C, Tiloca C, del Bo R, Corrado L, Pensato V, Agostini J, et al. Ubiquilin
2 mutations in Italian patients with amyotrophic lateral sclerosis and
frontotemporal dementia. J Neurol Neurosurg Psychiatry. 2013;84(2):183–7.
10.1136/jnnp-2012-303433

25. Huang X, Shen S, Fan D. No evidence for pathogenic role of UBQLN2
mutations in sporadic amyotrophic lateral sclerosis in the mainland Chinese
population. Le W, editor. PLoS One. 2017;12(1):1–5.
10.1371/journal.pone.0170943

26. Lattante S, le Ber I, Camuzat A, Pariente J, Brice A, Kabashi E. Screening
UBQLN-2 in French frontotemporal lobar degeneration and frontotemporal
lobar degeneration-amyotrophic lateral sclerosis patients. Neurobiol Aging.
2013;34(8):2078.e5-2078.e6.

27. Luukkainen L, Helisalmi S, Kytövuori L, Ahmasalo R, Solje E, Haapasalo A,
et al. Mutation Analysis of the Genes Linked to Early Onset Alzheimer's

654		Disease and Frontotemporal Lobar Degeneration. Journal of Alzheimer's
655		Disease. 2019 Jun 4;69(3):775–82. 10.3233/JAD-181256
656	28.	McLaughlin RL, Kenna KP, Vajda A, Byrne S, Bradley DG, Hardiman O, et
657		al. UBQLN2 mutations are not a frequent cause of amyotrophic lateral sclerosis
658		in ireland. Neurobiol Aging. 2014;35(1):267.e9-267.e11.
659		10.1016/j.neurobiolaging.2013.07.023
660	29.	Millecamps S, Corcia P, Cazeneuve C, Boillée S, Seilhean D, Danel-Brunaud
661		V, et al. Mutations in UBQLN2 are rare in French amyotrophic lateral sclerosis.
662		Neurobiol Aging. 2012;33(4):839.e1-839.e3.
663		10.1016/j.neurobiolaging.2011.11.010
664	30.	Morgan S, Shatunov A, Sproviero W, Jones AR, Shoai M, Hughes D, et al. A
665		comprehensive analysis of rare genetic variation in amyotrophic lateral
666		sclerosis in the UK. Brain. 2017 Jun 1;140(6):1611–8. 10.1093/brain/awx082
667	31.	Özoğuz A, Uyan Ö, Birdal G, Iskender C, Kartal E, Lahut S, et al. The distinct
668		genetic pattern of ALS in Turkey and novel mutations. Neurobiol Aging.
669		2015;36(4):1764.e9-1764.e18. 10.1016/j.neurobiolaging.2014.12.032
670	32.	Synofzik M, Maetzler W, Grehl T, Prudlo J, vom Hagen JM, Haack T, et al.
671		Screening in ALS and FTD patients reveals 3 novel UBQLN2 mutations
672		outside the PXX domain and a pure FTD phenotype. Neurobiol Aging.
673		2012;33(12):2949.e13-2949.e17. 10.1016/j.neurobiolaging.2012.07.002
674	33.	Teyssou E, Chartier L, Amador M-D-M, Lam R, Lautrette G, Nicol M, et al.
675		Novel UBQLN2 mutations linked to amyotrophic lateral sclerosis and atypical
676		hereditary spastic paraplegia phenotype through defective HSP70-mediated
677		proteolysis. Neurobiol Aging. 2017 Oct;58(March):239.e11-239.e20.
678		10.1016/j.neurobiolaging.2017.06.018
679	34.	Tripolszki K, Gampawar P, Schmidt H, Nagy ZF, Nagy D, Klivényi P, et al.
680		Comprehensive Genetic Analysis of a Hungarian Amyotrophic Lateral
681		Sclerosis Cohort. Front Genet. 2019 Aug 16;10(JUL).
682		10.3389/fgene.2019.00732
683	35.	Ugwu F, Rollinson S, Harris J, Gerhard A, Richardson A, Jones M, et al. A
684		UBQLN2 variant of unknown significance in frontotemporal lobar

685  degeneration. Neurobiol Aging. 2015 Jan;36(1):546.e15-546.e16.
686  10.1016/j.neurobiolaging.2014.08.002

687 36. Vengoechea J, David MP, Yaghi SR, Carpenter L, Rudnicki SA. Clinical
688  variability and female penetrance in X-linked familial FTD/ALS caused by a
689  P506S mutation in UBQLN2. Amyotroph Lateral Scler Frontotemporal
690  Degener. 2013 Dec 14;14(7–8):615–9. 10.3109/21678421.2013.824001

691 37. Vildan C, Sule D, Turker B, Hilmi U, Sibel KB. Genetic alterations of C9orf72,
692  SOD1, TARDBP, FUS, and UBQLN2 genes in patients with Amyotrophic
693  Lateral Sclerosis. Cogent Med. 2019;6(1):1–10.
694  10.1080/2331205x.2019.1582400

695 38. Scotter EL, Smyth L, Bailey JWT, Wong C-H, de Majo M, Vance CA, et al.
696  C9ORF72 and UBQLN2 mutations are causes of amyotrophic lateral sclerosis
697  in New Zealand: a genetic and pathologic study using banked human brain
698  tissue. Neurobiol Aging. 2017 Jan;49:214.e1-214.e5.
699  10.1016/j.neurobiolaging.2016.06.019

700 39. Brettschneider J, Van Deerlin VM, Robinson JL, Kwong L, Lee EB, Ali YO,
701  et al. Pattern of ubiquilin pathology in ALS and FTLD indicates presence of
702  C9ORF72 hexanucleotide expansion. Acta Neuropathol. 2012 Jun
703  18;123(6):825–39. 10.1007/s00401-012-0970-z

704 40. Arai T, Hasegawa M, Akiyama H, Ikeda K, Nonaka T, Mori H, et al. TDP-43
705  is a component of ubiquitin-positive tau-negative inclusions in frontotemporal
706  lobar degeneration and amyotrophic lateral sclerosis. Biochem Biophys Res
707  Commun. 2006;351(3):602–11. 10.1016/j.bbrc.2006.10.093

708 41. Kwiatkowski TJ, Bosco DA, LeClerc AL, Tamrazian E, Vanderburg CR, Russ
709  C, et al. Mutations in the FUS/TLS gene on chromosome 16 cause familial
710  amyotrophic lateral sclerosis. Science (1979). 2009;323(5918):1205–8.
711  10.1126/science.1166066

712 42. Maruyama H, Morino H, Ito H, Izumi Y, Kato H, Watanabe Y, et al. Mutations
713  of optineurin in amyotrophic lateral sclerosis. Nature. 2010;465(7295):223–6.

714 43. Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT,
715  et al. Ubiquitinated TDP-43 in Frontotemporal Lobar Degeneration and

Amyotrophic Lateral Sclerosis. Science (1979). 2006 Oct 6;314(5796):130–3.

10.1126/science.1134108

44. Vance C, Rogelj B, Hortobagyi T, de Vos KJ, Nishimura AL, Sreedharan J, et

al. Mutations in FUS, an RNA Processing Protein, Cause Familial

Amyotrophic Lateral Sclerosis Type 6. Science (1979). 2009 Feb

27;323(5918):1208–11. 10.1126/science.1165942

45. Waldvogel HJ, Bullock JY, Synek BJ, Curtis MA, van Roon-Mom WMC,

Faull RLM. The collection and processing of human brain tissue for research.

Cell Tissue Bank. 2008 Sep 21;9(3):169–79. 10.1007/s10561-008-9068-1

46. Keogh MJ, Wei W, Wilson I, Coxhead J, Ryan S, Rollinson S, et al. Genetic

compendium of 1511 human brains available through the UK Medical

Research Council Brain Banks Network Resource. Genome Res.

2017;27(1):165–73.

47. Murray HC, Johson K, Sedlock A, Highet B, Dieriks BV, Anekal PV, et al.

Lamina-specific immunohistochemical signatures in the olfactory bulb of

healthy, Alzheimer's and Parkinson's disease patients. Commun Biol.

2021;5(1):88. 10.1038/s42003-022-03032-5

48. Waldvogel HJ, Curtis MA, Baer K, Rees MI, Faull RLM.

Immunohistochemical staining of post-mortem adult human brain sections. Nat

Protoc. 2007;1(6):2719–32. 10.1038/nprot.2006.354

49. Maric D, Jahanipour J, Li XR, Singh A, Mobiny A, Van Nguyen H, et al.

Whole-brain tissue mapping toolkit using large-scale highly multiplexed

immunofluorescence imaging and deep neural networks. Nat Commun.

2021;12(1):1550. 10.1038/s41467-021-21735-x

50. Liu F, Morderer D, Wren MC, Vettleson-Trutza SA, Wang Y, Rabichow BE,

et al. Proximity proteomics of C9orf72 dipeptide repeat proteins identifies

molecular chaperones as modifiers of poly-GA aggregation. Acta Neuropathol

Commun. 2022;10(1):1–18. 10.1186/s40478-022-01322-x

51. Neumann M, Kwong LK, Lee EB, Kremmer E, Flatley A, Xu Y, et al.

Phosphorylation of S409/410 of TDP-43 is a consistent feature in all sporadic

and familial forms of TDP-43 proteinopathies. Acta Neuropathol.
2009;117(2):137–49.

52.   Luisier F, Vonesch C, Blu T, Unser M. Fast interscale wavelet denoising of
Poisson-corrupted images. Signal Processing. 2010;90(2):415–27.
10.1016/j.sigpro.2009.07.009

53.   Henden L, Wakeham D, Bahlo M. XIBD: software for inferring pairwise
identity by descent on the X chromosome. Bioinformatics. 2016 Aug
1;32(15):2389–91. 10.1093/bioinformatics/btw124

54.   Li H, Glusman G, Hu H, Shankaracharya, Caballero J, Hubley R, et al.
Relationship Estimation from Whole-Genome Sequence Data. Visscher PM,
editor. PLoS Genet. 2014 Jan 30;10(1):e1004144.
10.1371/journal.pgen.1004144

55.   Brettschneider J, Del Tredici K, Toledo JB, Robinson JL, Irwin DJ, Grossman
M, et al. Stages of pTDP-43 pathology in amyotrophic lateral sclerosis. Ann
Neurol. 2013 Jul;74(1):20–38. 10.1002/ana.23937

56.   Mackenzie IRA, Arzberger T, Kremmer E, Troost D, Lorenzl S, Mori K, et al.
Dipeptide repeat protein pathology in C9ORF72 mutation cases: clinico-
pathological correlations. Acta Neuropathol. 2013 Dec 6;126(6):859–79.
10.1007/s00401-013-1181-y

57.   Gorrie GH, Fecto F, Radzicki D, Weiss C, Shi Y, Dong H, et al. Dendritic
spinopathy in transgenic mice expressing ALS/dementia-linked mutant
UBQLN2. Proceedings of the National Academy of Sciences. 2014 Oct
7;111(40):14524–9. 10.1073/pnas.1405741111

58.   Ravits J, Appel S, Baloh RH, Barohn R, Rix Brooks B, Elman L, et al.
Deciphering amyotrophic lateral sclerosis: What phenotype, neuropathology
and genetics are telling us about pathogenesis. Amyotroph Lateral Scler
Frontotemporal Degener. 2013 May 16;14(sup1):5–18.
10.3109/21678421.2013.778548

59.   Nguyen HP, Van Broeckhoven C, van der Zee J. ALS Genes in the Genomic
Era and their Implications for FTD. Trends in Genetics. 2018;34(6):404–23.
10.1016/j.tig.2018.03.001

60.   Taylor JP, Brown RH, Cleveland DW. Decoding ALS: from genes to
778   mechanism. Nature. 2016 Nov 9;539(7628):197–206. 10.1038/nature20413

779   61.   Ling S, Polymenidou M, Cleveland DW. Converging Mechanisms in ALS and
780   FTD: Disrupted RNA and Protein Homeostasis. Neuron. 2013 Aug;79(3):416–
781   38. 10.1016/j.neuron.2013.07.033

782   62.   Mori K, Weng S-M, Arzberger T, May S, Rentzsch K, Kremmer E, et al. The
783   C9orf72 GGGGCC Repeat Is Translated into Aggregating Dipeptide-Repeat
784   Proteins in FTLD/ALS. Science (1979). 2013 Mar 15;339(6125):1335–8.
785   10.1126/science.1232927

786   63.   Mori K, Arzberger T, Grässer FA, Gijselinck I, May S, Rentzsch K, et al.
787   Bidirectional transcripts of the expanded C9orf72 hexanucleotide repeat are
788   translated into aggregating dipeptide repeat proteins. Acta Neuropathol. 2013
789   Dec 17;126(6):881–93. 10.1007/s00401-013-1189-3

790   64.   Shibata N, Hirano A, Kobayashi M, Siddique T, Deng H-X, Hung W-Y, et al.
791   Intense Superoxide Dismutase-1 Immunoreactivity in Intracytoplasmic
792   Hyaline Inclusions of Familial Amyotrophic Lateral Sclerosis with Posterior
793   Column Involvement. J Neuropathol Exp Neurol. 1996 Apr;55(4):481–90.
794   10.1097/00005072-199604000-00011

795   65.   Bruijn LI, Houseweart MK, Kato S, Anderson KL, Anderson SD, Ohama E, et
796   al. Aggregation and Motor Neuron Toxicity of an ALS-Linked SOD1 Mutant
797   Independent from Wild-Type SOD1. Science (1979). 1998 Sep
798   18;281(5384):1851–4. 10.1126/science.281.5384.1851

799   66.   Keith JL, Swinkin E, Gao A, Alminawi S, Zhang M, McGoldrick P, et al.
800   Neuropathologic description of CHCHD10 mutated amyotrophic lateral
801   sclerosis. Neurol Genet. 2020 Feb 13;6(1):e394.
802   10.1212/NXG.0000000000000394

803   67.   Smith BN, Topp SD, Fallini C, Shibata H, Chen H-J, Troakes C, et al.
804   Mutations in the vesicular trafficking protein annexin A11 are associated with
805   amyotrophic lateral sclerosis. Sci Transl Med. 2017 May 3;9(388):eaad9157.
806   10.1126/scitranslmed.aad9157

807   68.   Smith BN, Vance C, Scotter EL, Troakes C, Wong CH, Topp S, et al. Novel
808   mutations support a role for Profilin 1 in the pathogenesis of ALS. Neurobiol
809   Aging. 2015;36(3):1602.e17-1602.e27. 10.1016/j.neurobiolaging.2014.10.032

810   69.   Teyssou E, Takeda T, Lebon V, Boillée S, Doukouré B, Bataillon G, et al.
811   Mutations in SQSTM1 encoding p62 in amyotrophic lateral sclerosis: genetics
812   and neuropathology. Acta Neuropathol. 2013 Apr 17;125(4):511–22.
813   10.1007/s00401-013-1090-0

814   70.   Hirsch-Reinshagen V, Pottier C, Nicholson AM, Baker M, Hsiung G-YR,
815   Krieger C, et al. Clinical and neuropathological features of ALS/FTD with
816   TIA1 mutations. Acta Neuropathol Commun. 2017 Dec 7;5(1):96.
817   10.1186/s40478-017-0493-x

818   71.   Tada M, Doi H, Koyano S, Kubota S, Fukai R, Hashiguchi S, et al. Matrin 3 Is
819   a Component of Neuronal Cytoplasmic Inclusions of Motor Neurons in
820   Sporadic Amyotrophic Lateral Sclerosis. American Journal of Pathology.
821   2018;188(2):507–14. 10.1016/j.ajpath.2017.10.007

822   72.   van Deerlin VM, Leverenz JB, Bekris LM, Bird TD, Yuan W, Elman LB, et al.
823   TARDBP mutations in amyotrophic lateral sclerosis with TDP-43
824   neuropathology: a genetic and histopathological analysis. Lancet Neurol. 2008
825   May;7(5):409–16. 10.1016/S1474-4422(08)70071-1

826   73.   Tan RH, Kril JJ, Fatima M, McGeachie A, McCann H, Shepherd C, et al. TDP-
827   43 proteinopathies: pathological identification of brain regions differentiating
828   clinical phenotypes. Brain. 2015 Oct;138(10):3110–22. 10.1093/brain/awv220

829   74.   McEachin ZT, Gendron TF, Raj N, García-Murias M, Banerjee A, Purcell RH,
830   et al. Chimeric Peptide Species Contribute to Divergent Dipeptide Repeat
831   Pathology in c9ALS/FTD and SCA36. Neuron. 2020 Jul;107(2):292-305.e6.
832   10.1016/j.neuron.2020.04.011

833   75.   Freibaum BD, Taylor JP. The Role of Dipeptide Repeats in C9ORF72-Related
834   ALS-FTD. Front Mol Neurosci. 2017 Feb 13;10(February):1–9.
835   10.3389/fnmol.2017.00035

836   76.   Lee Y-B, Baskaran P, Gomez-Deza J, Chen H-J, Nishimura AL, Smith BN, et
837   al. C9orf72 poly GA RAN-translated protein plays a key role in amyotrophic

lateral sclerosis via aggregation and toxicity. Hum Mol Genet. 2017 Dec
15;26(24):4765–77. 10.1093/hmg/ddx350

77. Nonaka T, Masuda-Suzukake M, Hosokawa M, Shimozawa A, Hirai S, Okado
H, et al. C9ORF72 dipeptide repeat poly-GA inclusions promote intracellular
aggregation of phosphorylated TDP-43. Hum Mol Genet. 2018 Aug
1;27(15):2658–70. 10.1093/hmg/ddy174

78. Lindsay RD, Scheibel AB. Quantitative analysis of dendritic branching pattern
of granular cells from human dentate gyrus. Exp Neurol. 1976 Aug;52(2):295–
310. 10.1016/0014-4886(76)90173-4

79. Amaral DG, Scharfman HE, Lavenex P. The dentate gyrus: fundamental
neuroanatomical organization (dentate gyrus for dummies). In: Prog Brain Res.
2007. p. 3–790. 10.1016/S0079-6123(07)63001-5

80. May S, Hornburg D, Schludi MH, Arzberger T, Rentzsch K, Schwenk BM, et
al. C9orf72 FTLD/ALS-associated Gly-Ala dipeptide repeat proteins cause
neuronal toxicity and Unc119 sequestration. Acta Neuropathol. 2014 Oct
14;128(4):485–503. 10.1007/s00401-014-1329-4

81. Gupta R, Lan M, Mojsilovic-Petrovic J, Choi WH, Safren N, Barmada S, et al.
The Proline/Arginine Dipeptide from Hexanucleotide Repeat Expanded
C9ORF72 Inhibits the Proteasome. eNeuro. 2017 Jan;4(1):ENEURO.0249-
16.2017. 10.1523/ENEURO.0249-16.2017

82. Zhang Y-J, Gendron TF, Grima JC, Sasaguri H, Jansen-West K, Xu Y-F, et al.
C9ORF72 poly(GA) aggregates sequester and impair HR23 and
nucleocytoplasmic transport proteins. Nat Neurosci. 2016 May 21;19(5):668–
77. 10.1038/nn.4272

83. van Blitterswijk M, Gendron TF, Baker MC, DeJesus-Hernandez M, Finch
NA, Brown PH, et al. Novel clinical associations with specific C9ORF72
transcripts in patients with repeat expansions in C9ORF72. Acta Neuropathol.
2015 Dec 5;130(6):863–76. 10.1007/s00401-015-1480-6

84. Webster CP, Smith EF, Bauer CS, Moller A, Hautbergue GM, Ferraiuolo L, et
al. The C9orf72 protein interacts with Rab1a and the <scp>ULK</scp> 1

complex to regulate initiation of autophagy. EMBO J. 2016 Aug
22;35(15):1656–76. 10.15252/embj.201694401

85. Wang M, Wang H, Tao Z, Xia Q, Hao Z, Prehn JHM, et al. C9orf72 associates
with inactive Rag GTPases and regulates mTORC1-mediated autophagosomal
and lysosomal biogenesis. Aging Cell. 2020 Apr 25;19(4):1–14.
10.1111/acel.13126

86. Boivin M, Pfister V, Gaucherot A, Ruffenach F, Negroni L, Sellier C, et al.
Reduced autophagy upon C9ORF72 loss synergizes with dipeptide repeat
protein toxicity in G4C2 repeat expansion disorders. EMBO J. 2020 Feb
17;39(4):1–15. 10.15252/embj.2018100574

87. Zhu Q, Jiang J, Gendron TF, McAlonis-Downes M, Jiang L, Taylor A, et al.
Reduced C9ORF72 function exacerbates gain of toxicity from ALS/FTD-
causing repeat expansion in C9orf72. Nat Neurosci. 2020;23(May).
10.1038/s41593-020-0619-5

88. Beckers J, Tharkeshwar AK, van Damme P. C9orf72 ALS-FTD: recent
evidence for dysregulation of the autophagy-lysosome pathway at multiple
levels. Autophagy. 2021;00(00):1–17. 10.1080/15548627.2021.1872189

89. Sellier C, Campanari M, Julie Corbier C, Gaucherot A, Kolb-Cheynel I, Oulad-
Abdelghani M, et al. Loss of C9ORF72 impairs autophagy and synergizes with
polyQ Ataxin-2 to induce motor neuron dysfunction and cell death. EMBO J.
2016 Jun 15;35(12):1276–97. 10.15252/embj.201593350

90. Renaud L, Picher-Martel V, Codron P, Julien J. Key role of UBQLN2 in
pathogenesis of amyotrophic lateral sclerosis and frontotemporal dementia.
Acta Neuropathol Commun. 2019 Dec 18;7(1):103. 10.1186/s40478-019-
0758-7

91. Zhang KY, Yang S, Warraich ST, Blair IP. Ubiquilin 2: A component of the
ubiquitin-proteasome system with an emerging role in neurodegeneration.
International Journal of Biochemistry and Cell Biology. 2014;50(1):123–6.
10.1016/j.biocel.2014.02.018

92. Sharkey LM, Sandoval-Pistorius SS, Moore SJ, Gerson JE, Komlo R, Fischer
S, et al. Modeling UBQLN2-mediated neurodegenerative disease in mice:

Shared and divergent properties of wild type and mutant UBQLN2 in phase
separation, subcellular localization, altered proteostasis pathways, and
selective cytotoxicity. Neurobiol Dis. 2020;143(June).

93.   Mann DM, Rollinson S, Robinson A, Bennion Callister J, Thompson JC,
Snowden JS, et al. Dipeptide repeat proteins are present in the p62 positive
inclusions in patients with frontotemporal lobar degeneration and motor
neurone disease associated with expansions in C9ORF72. Acta Neuropathol
Commun. 2013 Dec 14;1(1):68. 10.1186/2051-5960-1-68

94.   Al-Sarraj S, King A, Troakes C, Smith B, Maekawa S, Bodi I, et al. p62
positive, TDP-43 negative, neuronal cytoplasmic and intranuclear inclusions in
the cerebellum and hippocampus define the pathology of C9orf72-linked
FTLD and MND/ALS. Acta Neuropathol. 2011 Dec 19;122(6):691–702.
10.1007/s00401-011-0911-2

95.   Piras A, Collin L, Grüninger F, Graff C, Rönnbäck A. Autophagic and
lysosomal defects in human tauopathies: analysis of post-mortem brain from
patients with familial Alzheimer disease, corticobasal degeneration and
progressive supranuclear palsy. Acta Neuropathol Commun. 2016;4:22.
10.1186/s40478-016-0292-9

96.   Babu JR, Geetha T, Wooten MW. Sequestosome 1/p62 shuttles
polyubiquitinated tau for proteasomal degradation. J Neurochem. 2005
Jul;94(1):192–203. 10.1111/j.1471-4159.2005.03181.x

97.   Zatloukal K, Stumptner C, Fuchsbichler A, Heid H, Schnoelzer M, Kenner L,
et al. P62 Is a Common Component of Cytoplasmic Inclusions in Protein
Aggregation Diseases. American Journal of Pathology. 2002;160(1):255–63.
10.1016/S0002-9440(10)64369-6

98.   Kuusisto E, Salminen A, Alafuzoff I. Ubiquitin-binding protein p62 is present
in neuronal and glial inclusions in human tauopathies and synucleinopathies.
Neuroreport. 2001 Jul;12(10):2085–90. 10.1097/00001756-200107200-00009

99.   Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards
and guidelines for the interpretation of sequence variants: a joint consensus
recommendation of the American College of Medical Genetics and Genomics

and the Association for Molecular Pathology. Genetics in Medicine. 2015

May;17(5):405–24. 10.1038/gim.2015.30

100. Ramstetter MD, Dyer TD, Lehman DM, Curran JE, Duggirala R, Blangero J,

et al. Benchmarking relatedness inference methods with genome-wide data

from thousands of relatives. Genetics. 2017;207(1):75–82.

Tables

Table 1. IBD segments inferred over *UBQLN2* on the X chromosome (chromosome 23).

Individual 1		Individual 2					
Family ID	**Pedigree ID**	**Family ID**	**Individual ID**	**StartSNP**	**EndSNP**	**Start Pos(bp)**	**End Pos(bp)**
FALS5	III:8	FALS5	IV:9	rs17330993	rs11156600	2779749	154440161
FALS5	III:8	FALS5	IV:18[a]	rs7062445	rs6423133	19650411	123608292
FALS5	III:8	FALS14	III:2	rs952836	rs17315029	40344087	129605268
FALS5	IV:9	FALS5	IV:18	rs6628597	rs6423133	31382037	123608292
FALS5	IV:9	FALS14	III:2	rs952836	rs17277770	40344087	132774456
FALS5	IV:18	FALS14	III:2	rs952836	rs6423133	40344087	123608292

940 [a] FALS5 IV:18 is referred to as MN17 in this report.
941

942 **Table 2.** Estimated degree of relatedness and fraction of genome with zero, one, and two

943 alleles identical-by-descent (IBD) between all samples in *UBQLN2* p.T487I-linked

944 families FALS5 and FALS14.

		Individual 1		Individual 2					
		Family ID	Pedigree ID	Family ID	Individual ID	Fraction IBD=0	Fraction IBD=1	Fraction IBD=2	Degree[a]
Intra-family		FALS5	III:8	FALS5	IV:9	0.529	0.471	0	2
		FALS5	III:8	FALS5	IV:18[b]	0.529	0.471	0	2
		FALS5	IV:9	FALS5	IV:18	0.751	0.249	0	3
		FALS14	II:1	FALS14	III:2	0.515	0.485	0	2
Inter-family		FALS5	III:8	FALS14	II:1	0.877	0.123	0	4
		FALS5	III:8	FALS14	III:2	0.927	0.073	0	5
		FALS5	IV:9	FALS14	II:1	0.942	0.058	0	5
		FALS5	IV:9	FALS14	III:2	0.948	0.052	0	5
		FALS5	IV:18	FALS14	II:1	0.926	0.074	0	5
		FALS5	IV:18	FALS14	III:2	0.948	0.052	0	5

[a] Degrees of relatedness as per [100]:

2 – Grandparent-grandchild, Avuncular, Half-sibling

3 – First cousin, Great-grandparent, Grand-avuncular

4 – First cousin once removed, GG-grandparent

5 – Second cousin, First cousin twice removed

[b] FALS5 IV:18 is referred to as MN17 in this report.

	Figure legends

Figure 1. Ubiquilin 2 labelling in previous studies fails to discriminate between
***UBQLN2*-linked and other genotypes of ALS/FTD.** Previously published
immunohistochemical analyses of hippocampus molecular layer (ML), granule cell layer
(GCL), and cornus ammonus (CA) regions fail to discriminate between *UBQLN2*-linked
ALS/FTD and ALS/FTD with other genotypes. Key shown within figure. Hippocampal
ubiquilin 2 (green) pathology is reported when ubiquilin 2 is either mutant or wildtype;
being present in the molecular layer in *UBQLN2*-linked and *C9orf72*-linked ALS/FTD
(including "ALS-dementia") but not in sALS; and in the granule cell layer in *C9orf72*-
linked ALS/FTD (including "ALS-dementia"). Hippocampal molecular layer ubiquilin 2
(green) is frequently reported and found together with ubiquitin (orange) or p62 (beige),
but not TDP-43 (blue). Hippocampal granule cell layer ubiquilin 2 pathology is found
together with dipeptide repeats (DPRs, pink) with or without TDP-43; but is variably
present in *UBQLN2*-linked ALS/FTD and sALS. In the hippocampal CA regions, ubiquilin
2 is present in *UBQLN2*-linked and in *C9orf72*-linked ALS/FTD (including "ALS-
dementia") where it is found together with DPRs, ubiquitin and p62.

Figure 2. Pathology in the hippocampal dentate gyrus and cornu ammonis regions of
control, familial, and sporadic cases. The majority of control tissue had no pTDP-43 or
ubiquilin 2 pathology, but one case had sparse p62 immunoreactivity in the GCL and CA-
pyr layers (**A** H230, pink arrows). Such punctate p62-positive inclusions were also found
in the GCL (**B$_1$-D$_1$**), and CA-pyr layers (**B$_2$-D$_2$**) in MN14, an ALS case of unknown
genotypic cause (**B**), MN24, a *SOD1* p.E101G case (**C**), and P525L, a *FUS* p.P525L case
(**D**). P62-co-labelled perinuclear pTDP-43 aggregates negative for ubiquilin 2 were found
in the GCL and CA-pyr regions of sporadic case ALS/FTD case MN15 (**E$_1$** and **E$_2$**, white
arrows), and sporadic ALS case MN30 (**F$_1$** and **F$_2$**, white arrows). Rare pTDP-43
aggregates negative for p62 are indicated by the yellow arrow in **F$_2$**. Scale bar A-D, 50 µm;
E-F, 25 µm. Abbreviations: CA – pyr, cornu ammonis – pyramidal cells; GCL, granule cell
layer.

Figure 3. Pathology in the hippocampal dentate gyrus and cornu ammonis regions of _C9orf72_-linked cases. Wispy ubiquilin 2 aggregates in the ML and CA – l-m-rad layers were predominantly p62 negative (**A,B₁** and **A,B₂**, green arrows), with larger aggregates also co-labelling with p62 in the ML (**B₁**, white arrow). In MN18, the CA – pyr cells harboured stellate p62 aggregates (**A₃**, pink arrow) while the GCL showed stellate p62 aggregates either with ubiquilin 2 (**A₄**, white arrows) or without (**A₄**, pink arrow). Case MN28 shared this pattern of p62 and ubiquilin 2 pathology with the additional presence of perinuclear pTDP-43 inclusions, often found without other co-labelling in the CA-pyr region (**B₃**, yellow arrow) and GCL (**B₄**, yellow arrow). Ubiquilin 2 aggregates in the ML and CA – l-m/rad layers were negative for dipeptide repeat proteins (**C₁-C₂**, **D₁-D₂**, green arrows). The dominant pathology in CA – pyr regions was the presence of polyGA and polyGP inclusions negative for ubiquilin 2 (**C₃** and **D₃**, orange arrowheads). All GCL ubiquilin 2 was positive for at least one dipeptide repeat protein (**C₄** and **D₄**, purple arrows). Ubiquilin 2-negative aggregates were comprised of both dipeptide repeat proteins (**C₄** and **D₄**, orange arrows) or most commonly, polyGA without other co-labelling (**C₄** and **D₄**, red arrows). Super-resolution STED microscopy of co-localised ubiquilin 2-polyGA GCL aggregates in case MN28 demonstrated polyGA enmeshed with and encircling ubiquilin 2 (**E**) or ubiquilin 2 labelling around a core of polyGA (**F**). Scale bar in main images A-D, 50 µm; zooms in C₄ and D₄, 25 µm; in E-F, 1 µm. Abbreviations: CA – lm/rad, cornu ammonis – lacunosum-molecular and radiatum layers; CA – pyr, cornu ammonis – pyramidal cells; GCL, granule cell layer.

Figure 4. Ubiquilin 2, pTDP-43, and p62 pathology in the hippocampal dentate gyrus and cornu ammonis regions of _UBQLN2_-linked cases. All _UBQLN2_-linked ALS/FTD cases showed numerous p62-positive ubiquilin 2 aggregates, indicated by white arrows, in the ML (**A₁-D₁**) and CA – l-m/rad layers (**A₂-D₂**). Case P506S had the additional presence of aggregates positive for pTDP-43, ubiquilin 2, and p62 in the CA – pyr layer (**D₂**, filled white arrowhead). Cases P497H and MN17-T487I showed no aggregate labelling in the GCL (**A₃**, **B₃**). ALS/FTD cases V:7 – T487I and P506S showed rare GCL p62-positive pTDP-43 aggregates negative for ubiquilin 2 (**C₃**, **D₃**, hollow white arrowheads). Only in P506S were the numerous compact pTDP-43 aggregates found in the GCL co-localised

with ubiquilin 2 and p62 (**D₃**, filled white arrowheads). A PSP case harbouring a *UBQLN2*
p.S222G variant of uncertain significance (VOUS) (**E**) had pathology in the ML (**E₁**), and
sparse p62 positivity (pink arrows) in the CA – lm/rad layers (**E₂**) and GCL (**E₃**). Scale bar
in main images, 25 μm; zooms, 5 μm. Abbreviations: CA – lm/rad, cornu ammonis –
lacunosum-molecular and radiatum layers; CA – pyr, cornu ammonis – pyramidal cells;
GCL, granule cell layer; PSP, progressive supranuclear palsy.

1019

Figure 5. Combined pTDP-43, ubiquilin 2, p62, and dipeptide repeat protein
immunohistochemical staining discriminated between sporadic, *C9orf72*-linked, and
***UBQLN2*-linked ALS/FTD.** Key shown within figure. P62 aggregates ('+' symbol on
brown box) without other co-labelling were observed in CA – pyr and GCL cells in a subset
of neurologically normal controls (n=1 of 5), unrelated familial ALS (n=5 of 5), and
sporadic ALS cases (n=4 of 20). In contrast, pTDP-43 pathology (boxed '+' symbol on
blue box) was present in these layers in a separate subset of sALS cases (n=6 of 20), co-
localising with p62. Ubiquilin 2 pathology ('+' symbol on green box) was present in the
hippocampus when ubiquilin 2 was wildtype (*C9orf72*-linked ALS) or mutant (*UBQLN2*-
linked ALS/FTD). Wildtype ubiquilin 2 in *C9orf72*-linked ALS ML and CA – lm/rad
regions was p62-negative, and in the CA – pyr layer was associated with polyGA, polyGP,
and p62 aggregates. PolyGA and polyGP aggregates were found in the GCL with or
without ubiquilin 2, all colocalising with p62. CA – pyr and GCL aggregates had additional
pTDP-43 pathology in only one case, and even then, only rarely. Mutant ubiquilin 2
aggregates in *UBQLN2*-linked ALS/FTD ML, and CA – lm/rad regions were p62-positive
and associated with GCL pTDP-43 aggregates only in some cases and either with or
without ubiquilin 2 co-labelling. Blue outlines indicate unique aggregation features of
C9orf72-linked ALS and *UBQLN2*-linked ALS/FTD hippocampal pathology.
Abbreviations: CA – lm/rad, cornu ammonis – lacunosum-molecular and radiatum layers;
CA – pyr, cornu ammonis – pyramidal cells; GCL, granule cell layer.

1040

Figure 6. Schematic representation of the hippocampal neuropathological signature
defining *UBQLN2*-linked ALS/FTD. In *UBQLN2*-linked ALS/FTD cases without
hippocampal pTDP-43 proteinopathy, mutant ubiquilin 2 is punctate, p62-positive, and

forms aggregates exclusively in the CA l-m/rad layer and ML (cases p.P497H and MN17

– p.T487I). In *UBQLN2*-linked ALS/FTD cases with hippocampal pTDP-43

proteinopathy, in addition to punctate p62-positive ubiquilin 2 aggregates in the CA l-

m/rad, and molecular layers, there are granule cell layer pTDP-43 aggregates that are

either; frequent and ubiquilin 2-labelled (case p.P506S), or rare and ubiquilin 2-negative

(case V7-p.T487I), suggesting a pathological cascade in which granule cell layer pTDP-43

aggregates provide a scaffold around which mutant ubiquilin 2 can aggregate.

Abbreviations: CA – lm/rad, cornu ammonis – lacunosum-molecular and radiatum layers;

CA – pyr, cornu ammonis – pyramidal cells; GCL, granule cell layer. Image created in

Biorender.com

Control

Familial ALS

| H230 | MN14 | *SOD1* p.E101G | *FUS* p.P525L |

GCL

A₁ — pTDP-43 / Ubiquilin 2 / p62

B₁

C₁

D₁

CA - pyr

A₂ B₂ C₂ D₂

Sporadic ALS

| MN15 (ALS/FTD) | MN30 (ALS) |

GCL

E₁ — pTDP-43 / Ubiquilin 2 / p62

F₁

CA - pyr

E₂ F₂

C9orf72-linked ALS

UBQLN2-linked ALS/FTD

UBQLN2 VOUS

| P497H | MN17 - T487I | V:7 - T487I | P506S | S222G |

ML

A₁ *ALS*

pTDP-43
Ubiquilin 2
p62

B₁ *ALS/FTD*

C₁ *ALS/FTD*

D₁ *ALS/FTD*

E₁ *PSP*

CA - l-m/rad

A₂

B₂

C₂

D₂ CA - l-m/rad

CA - pyr

E₂

GCL

A₃

B₃

C₃

D₃

E₃

Genetic classification	Case	Diagnosis	Aggregate deposition			
			ML	CA - l-m/rad	CA - pyr	GCL
Control (n=5)		Neurologically normal				
Control (n=1)	H230	Neurologically normal				
Familial ALS (n=5)						
Sporadic ALS (n=10)						
Sporadic ALS (n=4)						
Sporadic ALS (n=6)	MN27	MND-ALS/LBD				
	MN19	MND-ALS				
	MN13	MND-ALS				
	MN29	MND-ALS				
	MN15	MND-ALS/FTD				
	MN30	MND-ALS				
C9orf72-linked ALS (n=5)	MN2	MND-ALS				
	MN18	MND-ALS				
	MN23	MND-ALS				
	MN33	MND-ALS				
	MN28	MND-ALS				
UBQLN2-linked VOUS (n=1)	S222G	PSP				
UBQLN2-linked ALS/FTD (n=4)	P497H	MND-ALS				
	MN17 - T487I	MND-ALS/FTD				
	V:7 - T487I	MND-ALS/FTD				
	P506S	MND-ALS/FTD				

Key

ML Molecular layer **GCL** Granular cell layer **CA - l-m/rad** Cornu ammonis lacunosum-molecular and radiatum layer **CA - pyr** Cornu ammonis pyramidal layer

pTDP-43 Ubiquilin 2 p62 polyGA polyGP **+** Presence of pathology **–** Absence of pathology ☐ Stage 4 ALS ☐ Protein signature

LBD Lewy body dementia **FTD** Frontotemporal dementia **PSP** Progressive supranuclear palsy **VOUS** Variant of uncertain significance

UBQLN2-linked ALS/FTD

P497H	MN17 - T487I	V:7 - T487I	P506S

CA l-m/rad
CA pyr
ML
GCL

Hippocampal pTDP-43   −   +

Ubiquilin 2   p62   pTDP-43