An analysis of LysM domain function in LytE when fulfilling the D,L-endopeptidase requirement for viability in *Bacillus subtilis*

Laura Uelze[1]

Smurfit Institute of Genetics, Trinity College Dublin, Dublin 2. Ireland.

[1] present address: German Federal Institute for Risk Assessment, Berlin, Germany.

Keywords: LytE D,L-endopeptidase, *Bacillus subtilis*, LysM domain function

Abstract

The D,L-endopeptidase requirement states that *Bacillus subtilis* requires either the

activity of the LytE or the CwlO enzyme for viability, therefore proving that these two

enzymes can complement for each other despite their very different N-terminal domains.

Here, we show that another D,L-endopeptidase, LytF, can also fulfill the D,L-

endopeptidase requirement for viability, when expressed from the *cwlO* promoter. Both

LytE and LytF contain N-terminally located LysM domains, three and five respectively.

However, cells expressing another very similar D,L-endopeptidase CwlS, with four LysM

domains were not viable. This led us to investigate whether a LytE protein with any one

of its three LysM domains permuted can fulfill the D,L-endopeptidase requirement for

viability. We found that the three LysM domains are not functionally equivalent and that

the N-terminally located LysM domain plays a greater role for functioning of the LytE

enzyme than the subsequent domains. Based on an investigation of orthologous

enzymes in 19 *B. subtilis* species we propose an evolutionary model describing the

development of the LytE-, CwlS- and LytF-type D,L-endopeptidases and their LysM

domain repeats. In summary, these results show that the LytE enzyme has been

optimized to fulfill the D,L-endopeptidase requirement for cell viability of *B. subtilis* with

regard to the number and properties of LysM domains that mediate peptidoglycan-

binding.

INTRODUCTION:

Peptidoglycan and teichoic acids are the main structural components of the *Bacillus*

subtilis cell wall (Bhavsar and Brown, 2006; Barreteau *et al.*, 2008; Sauvage *et al.*,

2008; Vollmer *et al.*, 2008; Sekiguchi and Yamamoto, 2012). Peptidoglycan is

composed of glycan chains formed by polymerization of a disaccharide of *N*-

acetylglucosamine (Glc*N*Ac) and *N*-acetylmuramic acid (Mur*N*Ac). Stem pentapeptides

composed (proximal to distal order) of amino acids L-alanine (position 1), D-glutamate

(position 2), *meso*-diaminopimelate (position 3), D-alanine (position 4) and D-alanine

(position 5) extend from the lactyl group of Mur*N*Ac. The glycan strands are linked by

formation of a covalent peptide bond between the carboxyl group of D-alanine (position

4) of one strand [with concomitant loss of D-alanine (position 5)] and the amino group of

meso-diaminopimelate (position 3) of a different strand. Glycan strands can be up to 200

disaccharides in length with a stem peptide linkage index of up to ~33%. Thus

peptidoglycan forms a mesh outside the cell to which wall-teichoic acids composed of

poly(ribitol- or glycerol-phosphate) are covalently attached (Weidenmaier and Peschel,

2008; Swoboda et al., 2010; Kawai et al., 2011). This rigid mesh structure confers shape

on the cell and is capable of withstanding turgor pressures of up to 5 atmospheres

(Sekiguchi and Yamamoto, 2012). Interference with peptidoglycan metabolism, either

through inhibition of its synthesis by antibiotics (*e.g.* penicillin and vancomycin) or an

imbalance between synthesis and turnover leads to aberrant cell structure, cell division

and often cell death. Thus maintenance of cell wall integrity is a prerequisite for normal

cell viability.

The dramatic changes that occur in cell wall structure during growth and division require

exquisite control and coordination of cell wall synthesis and turnover. Autolysins play a

key role in sculpting the cell wall during these processes by cleavage of glycosidic

linkages in the glycan chains and peptide linkages in the stem peptides (Smith *et al*.,

2000; Sekiguchi and Yamamoto, 2012). *B. subtilis* encodes a cohort of 35 enzymes with

glucosaminidase, muraminidase, amidase and endopeptidase activities that confer the

capability to cleave all covalent linkages of peptidoglycan thereby allowing it to be

sculpted, remodeled and turned over during cell growth and division (Smith *et al*., 2000;

Sekiguchi and Yamamoto, 2012). A notable feature of the autolysin cohort of *B. subtilis*

is the degree of redundancy of individual enzymatic activities (Smith *et al*., 2000;

Sekiguchi and Yamamoto, 2012). For example, there are seven enzymes (CwlO, CwlS,

LytE, LytF, PgdS, YkfC, YqgT) with D,L-endopeptidase activity encoded in the *B. subtilis*

genome, each of which cleaves the stem peptide between D-glutamate (position 2) and

meso-diaminopimelate (position 3). However these enzymes differ in the type and/or

number of domains that interact with cell wall components. Thus the LytE, CwlS and

LytF D,L-endopeptidases have 3, 4 and 5 LysM domains respectively located at the

amino terminus of the protein, a domain that functions to bind proteins to peptidoglycan

(Mesnage *et al*., 2014). In contrast, the N-terminal domain of CwlO functions to bind the

enzyme to the FtsEX ATP-type transporter that is located in the cell membrane

(Yamaguchi *et al.*, 2004; Dominguez-Cuevas *et al.*, 2013; Meisner *et al.*, 2013). Thus

variance in these proteins regions allows differential localization and binding affinity of

these enzymes to elements located within the cell envelope.

While none of the seven D,L-endopeptidases are essential, *lytE* and *cwlO* are

synthetically lethal, signifying that the cell requires one or the other of these enzymes for

viability (Bisicchia *et al.*, 2007). This indicates that they perform an overlapping essential

function for the cell, probably in elongation of the cell wall during growth (Bisicchia *et al.*,

2007; Hashimoto *et al.* 2012). Expression of both CwlO and LytE are positively

regulated by the WalRK two-component signal transduction system (TCS). In addition,

LytE expression is controlled by the SigI/RsgI sigma factor, which is induced by heat

and other cell wall stresses (Tseng *et al.*, 2011; Liu *et al.,* 2017). Therefore the the

WalRK and SigI/RsgI regulatory systems combine to fulfill the D,L-endopeptidase

requirement during normal growth and stress conditions respectively.

The synthetic lethality of the CwlO and LytE enzymes is interesting on two accounts

(BIsicchia *et* al., 2012). Firstly, these enzymes bind to different regions and structures

within the cell envelope. The three LysM domains direct binding of LytE to peptidoglycan

in the cell wall (Ishikawa *et al.*, 1998; Buist *et al.*, 2008; Mesnage *et al*., 2014). In

contrast, the amino terminal region of CwlO binds to the FtsEX protein complex resulting

in tethering of the active enzyme to the outer face of the cell membrane (Dominguez-

Cuevas *et al.*, 2013; Meisner *et al.*, 2013). The second feature is that the high level of

redundancy among the cohort of autolysins in *B. subtilis* has hampered assigning

specific functions to individual enzymes. However, the requirement for D,L-

endopeptidase activity evident from the synthetic lethality of CwlO and LytE permits

investigation of the ability of individual enzymes with this activity to fulfill that

requirement.

In this study, the ability of individual enzymes to fulfill the D,L-endopeptidase

requirement for viability when individually expressed using the *cwlO* promoter is

investigated. In addition, we tested the LysM domains of LytE to establish which, and

how many, are needed to fulfill the D,L-endopeptidase requirement for viability. Our

results show that both LytE (3 LysM domains) and LytF (5 LysM domains), but not CwlS

(4 LysM domains) can fulfill the D,L-endopeptidase requirement for viability. In addition

we show that the three LysM domains of LytE are not functionally equivalent, but that

each alone is sufficient to fulfill the D,L-endopeptidase requirement for viability.

MATERIAL AND METHODS

Bioinformatic analysis

To investigate LysM domain-containing proteins, all entries containing the term 'LysM'

were extracted from UniProt, (55,014 hits in January 2016). These data were then

screened to identify *bona fide* LysM domains using the following criteria: (i) they must be

from a bacterial species and be specific to an individual bacterial strain (*e.g.*

Bacillus subtilis 168); (ii) putative domains must have a range of 40 – 70 amino acids

(Buist *et al.*, 2008; Desvaux *et al.*, 2018); (iiii) each protein must have a unique amino

acid. The resulting dataset contained 38,576 LysM domains located in 26,078 proteins

that were encoded by 6,971 different bacterial strains in 68 phyla.

A total of 150 LysM domain-containing proteins in this dataset were identified in

nineteen closely related strains of the *B. subtilis* species group (NCBI taxonomy).

Cluster analysis using ClustalOmega with default settings was performed to identify the

subset of proteins with a high level of identify in their catalytic domain, thereby

identifying orthologues of CwlS, LytE and LytF D,L-endopeptidases orthologs in these

B. subtilis 168 strains (Sievers *et al.*, 2011; Bodenhofer *et al.*, 2015). The alignment was

then used for metric multidimensional scaling conducted with the bios2mds package

(Pele *et al.*, 2012) to identify clusters of orthologs. Similarly, the relationships among the

185 LysM domains contained in the cohort of identified orthologs was established based

on an alignment of their amino acid sequence (ClustalOmega, substitution matrix =

BLOSUM80), followed by construction of a phylogenetic tree using Maximum Likelihood

via the phangorn package (Schliep, 2011). The best fitting amino acid substitution

models were identified (LG+G+I), the different model parameters subsequently

optimized and bootstrapping performed (100 replicates), before the circular phylogenetic

tree was visualized with ggtree (Yu *et al.*, 2016).

The phylogeny of the 19 *Bacillus subtilis* strains was inferred through alignment-free

genome comparison with feature frequency profiles (FFP) (Sims *et al.*, 2009). Genome

sequences (excluding plasmid sequences) of all species were retrieved from GenBank.

E. coli BL21(DE3) was included as outgroup. FPP was performed essentially as

described by Wang and Ash (2015). Neighbour Joining (NJ) trees were constructed

based on the Jensen-Shannon divergence matrices. It was found that varying lengths of

l between 20 and 25 resulted in highly similar trees. Accounting for the higher degree of

relatedness between species investigated in this study compared to Wang and Ash

(2015), length was adjusted to l = 24 in the final analysis and the NJ tree visualized with

ggtree (Yu *et al.*, 2016). The robustness of the tree was verified with bootstrapping and

bootstrapping values of 100 out of 100 were achieved for all branches.

Strains, plasmids and growth conditions

Bacterial strains and plasmids used in this study are listed in Supporting Information

Tables S2 and S3 respectively. Oligonucleotides were obtained from IDT, (Belgium) and

are listed in the supporting information Table S4. *Escherichia coli* strain TG-1 (Sambrook

et al., 1989) was used for propagating plasmids. *E. coli* strain BL21(DE3) was used for

protein overexpression and purification. Strains were grown in Luria-Bertani (LB)

medium and in salt-free buffered LB (Boylan *et al.*, 1993), to which NaCl was added up

to the desired concentration. The high phosphate defined medium (HPDM) and low

phosphate defined medium (LPDM) used, are described in Botella *et al*., (2011).

Antibiotics were added at concentrations described by Noone *et al*., (2014). OD_{600} was

measured using a UVmini-1240 UV-VIS spectrophotometer (Shimadzu Scientific

Instruments). *B. subtilis* 168 was transformed according to the procedure of

Anagnostopoulos and Spizizen (1960). Growth studies were conducted with the BioTek

Synergy™2 system (BioTek Instruments) at 37°C with slow shaking. Three technical

repeats were performed for each growth curve.

Strain construction

A *lytE* null mutant strain (LU1) was constructed by replacing the *lytE* gene with a

spectinomycin resistance gene (*spc*) The *spc* gene was amplified from plasmid

pDG1727 using primer pairs Spec_Fwd / Spec_Rev. Chromosomal regions located

upstream and downstream of *lytE* were amplified by PCR using primer pairs

LytE_Up_Fwd / LytE_Up_Rev_Spec and LytE_Do_Fwd_Spec / LytE_Do_Rev

respectively. These DNA fragments were joined to the fragment encoding the *spc* gene

by strand overlap PCR (Horton *et al.*, 1989). This linear DNA fragment was then

transformed into *B. subtilis* strain 168 with selection for spectinomycin resistance,

thereby generating strain LU1 in which the *lytE* gene is replaced by the *spc* gene.

A second *lytE* null mutant strain (LU331) was generated in a similar manner. The strain

differentiates from LC01, in that part of the *lytE* signal sequence was preserved in this

mutant. The *spc* gene was amplified as described previously. Chromosomal regions

located upstream and downstream of *lytE* were amplified by PCR using primer

pairs LytE_Up_Fwd_2 / Signal_LytE_Rev_Spec and LytE_Do_Fwd_Spec /

LytE_Do_Rev. The DNA fragments were joined to the fragment encoding the *spc* gene

by strand overlap PCR and the linear DNA fragment was transformed into WT168, with

selection for spectinomycin resistance, to generate strain LU331.

Strains were constructed that express the LytE protein from the *cwlO* promoter (LU17,

LU21, LU25). These strains were generated through transcriptional fusion between the

cwlO promoter and the *lytE* gene. To generate strain LU17, four PCR fragments were

amplified. First, the DNA fragments located upstream and downstream of *cwlO* locus

were amplified with the primer pairs CwlO_UF_Sma / CwlO_UR and QE93_Fwd_Ery /

QE_94_Rev respectively. Then the *lytE* open reading frame was amplified with the

primer pair LytE_DF and LytE_DR_Ery. Lastly, the erythromycin resistance gene *erm*

was amplified with primer pair Mls_Fwd / Mls _Rev from plasmid pDG6464. All PCR

fragments were joined together by strand overlap extension PCR and transformed into

strain EB35 with selection for erythromycin resistance. The resulting strain LU17

encodes two copies of the *lytE* gene (one at the *cwlO* locus and one at the *lytE* locus).

Subsequently, the *lytE* gene at the *lytE* locus was replaced with a spectinomycin

resistance gene (*spc*). For this purpose, the spc gene was amplified from LU1 with the

primer pair LytE_Up_Fwd / LytE_Do_Rev. The PCR fragment was then transformed into

LU17 with selection for spectinomycin resistance. The resulting strain LU21 encodes

one copy of the *lytE* gene at the *cwlO* locus.

Another similar strain (LU25) was generated with the additional feature that the leader

region of *cwlO* was deleted, thereby stabilizing the *lytE* mRNA and yielding higher LytE

protein expression (Noone *et al*., 2014). For this strain, a DNA fragment upstream of

cwlO from strain which has a 141 bp markerless deletion of the leader region (Noone *et*

al., 2014) was amplified from strain DN3000. This PCR fragment was transformed into

LU21 with selection for   erythromycin resistance to generate strain LU25.

In order to investigate LytE protein expression levels, the LytE protein was joined to a

HA-tag which can be detected by western analysis. Strains LU45, LU47, LU61 and

LU63 are derivatives of strains WT168, LS103, LU21 and LU25 in which the *lytE* gene is

fused with a C-terminal HA-tag. These strains were constructed by transforming plasmid

pLU2 into WT168, LS103, LU21 and LU25 with selection for kanamycin resistance. The

successful integration of the plasmid into a *cwlO* mutant strain (LS103) served as proof

for the functionality of the HA-tag, given that a failure would lead to a combined *cwlO* /

lytE mutant, which is synthetically lethal (Bisicchia *et al.*, 2007).

A markerless null *lytF* mutant was generated by the method of Bloor and Cranenburgh,

2006. The DNA regions upstream and downstream of *lytF* were amplified with primers

LytF_Up_Fwd/ ytF_Up_Rev and LytF_Do_Fwd/LytF_Do_Rev respectively. The *cat* gene

was amplified using primers Cat_Diff_Fwd/Cat_Diff_Rev and plasmid pDG1661 as

template. A *dif* site for recombination was introduced into the 5' end of each primer

(Table S4). The three DNA fragments were then ligated and transformed into WT168

with selection for chloramphenicol resistance. Transformants were then grown without

for 24 hours in LB medium without selection, plated onto LB agar plates and screened

for loss of chloramphenicol resistance. The null mutation in *lytF* in the resultant strain

LU74 (Δ*lytF*) was confirmed by PCR.

In order to generate a strain that expresses the LytF protein from the *cwlO* promoter

(with *cwlO* leader – LC111, without *cwlO* leader – LC125) a similar approach as for LU21

and LU25 was followed. Downstream fragments and the erythromycin resistance

cassette were amplified as described previously. The upstream fragment was amplified

with the primers CwlO_UF and CwlO_UR_LytF from WT168 (containing the *cwlO* leader

for LU111) or from DN3000 (no *cwlO* leader for LU125). A middle fragment (containing

the *lytF* open reading frame) was amplified with the primers LytF_Fwd / LytF_Rev_Ery.

The fragments were joined together by strand overlap extension PCR and the PCR

product was transformed into LU74 with selection for erythromycin. The resulting strains

LU111 and LU125 encode a copy of the *lytF* gene from the *cwlO* promoter (with / without

cwlO leader) and the *lytE* gene at the *lytE* locus.

Subsequently, the *lytE* gene at the *lytE* locus was replaced with a spectinomycin

resistance gene (*spc*). For this purpose, the spc gene was with the primer pair

LytE_Up_Fwd / LytE_Do_Rev from chromosomal DNA of LU. The PCR fragment was

then transformed into the parental strains LU111 and LU125 with selection for

spectinomycin resistance. The resulting strains LU116 and LU134 encode one copy of

the *lytF* gene from the *cwlO* promoter (with / without *cwlO* leader) and are knock-outs for

cwlO and *lytE*.

In order to investigate LytE / LytF protein expression levels, the LytE / LytF protein were

joined to a HA-tag which can be detected by western analysis. A C-terminal fusion

between either the *lytE* or the *lytF* gene and a HA-tag was achieved by transforming the

pLU2 plasmid into strain LU17 to generate strain LU143 and plasmid pLU6 into WT168,

LU111, LU116, LU125 and LU134 to generate strains LU147, LU151, LU153, LU155 and

LU157 respectively, with selection for kanamycin.

To generate strains containing a variant LytE expressed from the *cwlO* promoter in a

background with endogenous *lytE* gene, the plasmids pLU16, pLU15, pLU14, pLU17,

pLU18, pLU19, pLU20, pLU21, pLU22, pLU23, pLU24, pLU25, pLU26, pLU27, pLU28,

pLU30, pLU31, pLU29, pLU37, pLU38 and pLU40, were transformed by double cross-

over into WT168 with selection for kanamycin to generate the strains: LU239, LU207,

LU212, LU216, LU220, LU223, LU227, LU232, LU235, LU261, LU265, LU269, LU272,

LU276, LU280, LU286, LU288, LU312, LU314, LU315 and LU318.

To generate strains containing a variant LytE expressed from the *cwlO* promoter in a *lytE*

null mutant background, the plasmids pLU16, pLU15, pLU14, pLU17, pLU18, pLU19,

pLU20, pLU21, pLU22, pLU23, pLU24, pLU25, pLU26, pLU27, pLU28, pLU30, pLU31,

pLU29, pLU38, and pLU40 were transformed by double cross-over into LS089 with

selection for kanamycin to generate the strains: LU205, LU209, LU213, LU217, LU222,

LU226, LU229, LU233, LU238, LU262, LU266, LU270, LU273, LU277, LU281, LU287,

LU289, LU313, LU316 and LU320.

Strains were constructed that express a variant LytE from the *lytE* promoter (with intact

cwlO gene). For the construction of these strains four PCR fragments were amplified.

The upstream and the downstream fragments flanking the *lytE* locus were amplified with

the primer pairs LytE_Up_Fwd_2 / Signal_LytE_Rev and LytE_Do_Fwd_Ery /

LytE_Do_Rev from chromosomal DNA of WT168. The downstream fragment was joined

to an erythromycin resistance cassette, which was amplified from pDG646 with the

primers Mls_Fwd / Mls_Rev. The upstream fragment was joined to fragments of the

variant *lytE* genes, which were amplified with the primers Signal_LytE_Fwd /

Catalytic_LytE_Rev_Ery from the plasmids pLU40, pLU23, pLU24, pLU25, pLU20,

pLU27, pLU37, pLU16, pLU17, pLU14 and pLU29. After joining of the upstream

fragment (upstream homology to the *lytE* locus plus variant *lytE* gene sequence) to the

downstream fragment (erythromycin resistance cassette plus downstream homology to

the *lytE* locus) by strand overlap extension PCR, the linear PCR fragments were then

transformed into strain LU331 with selection for erythromycin resistance, to create the

strains LU335, LU337, LU339, LU341, LU372, LU374, LU398, LU399, LU401, LU403

and LU405 respectively.

In order to disrupt the *cwlO* gene in a series of strains, expressing a variant LytE from

the *lytE* promoter, a fragment encompassing the *cwlO* locus was amplified with the

primer pair CwlO_UF / QE94_Rev from chromosomal DNA of the *cwlO* null mutant

strain EB35. The PCR fragment was then transformed by double cross-over into the

strains LU335, LU337, LU339, LU341, LU372, LU374, LU399, LU401, LU403 and

LU405 with selection for kanamycin resistance to obtain strains LU343, LU350, LU352,

LU354, LU383, LU385, LU407, LU409, LU411 and LU413.

In order to investigate variant LytE protein expression levels, the variant LytE proteins

were joined to a HA-tag which can be detected by western analysis. To enable a C-

terminal fusion of the variant LytE proteins with a HA-tag, plasmid pLU46 with homology

to the *lytE* catalytic domain, followed by the HA-tag and was integrated by single cross-

over at the *lytE* locus in the strains LU205, LU213, LU217, LU229, LU262, LU266,

LU320, LU343, LU385, LU407, LU411, LU313, LU350, LU270, LU352 and LU413 to

generate the strains LU414, LU415, LU416, LU417, LU418, LU419, LU421, LU422,

LU424, LU425, LU427, LU432, LU435, LU440, LU442 and LU443. Strains were

selected for spectinomycin resistance.

Plasmid construction

Plasmid pLU2, pLU6 and pLU46 enable a C-terminal transcriptional fusion between the

either the *lytE* (pLU2, pLU6) or the *lytF* (pLU46) open reading frame and a HA-tag.

Plasmids pLU2 and pLU6 were constructed by cloning a fragment, with homologous

region to the C-terminus of the *lytE* or the *lytF* gene with the HA-tag sequence inserted

before the stop codon, into the parental vector pDG780. The fragments was amplified

with the following primer pair LytE_Tag_F_BamHI / LytE_Tag_R_HA_EcoRI;

LytF_Tag_F_BamHI / LytF_Tag_R_HA_EcoRI and cloned into pDG780 by use of the

BamHI and EcoRI restriction sites. Plasmid pLU46 is a derivative from pLU2 with a

spectinomycin resistance cassette. Exchange of the antibiotic resistance cassette was

achieved through amplifying the spectinomycin resistance gene from LU1 with

Spec_Fwd_EcoRI and Spec_Rev_SalI and cloning it into pLU2 via the SalI and EcoRI

restriction sites.

A parental plasmid (pLU10) was constructed for convenient cloning of variant *lytE* open

reading frames into a cassette that would allow integration of variant *lytE* genes at the

cwlO locus, thereby exchanging the *cwlO* against the *lytE* open reading frame. Plasmid

pLU10 was generated in a two-step process: First, upstream homology (containing the

cwlO leader and promoter and part of the *lytE* signal sequence, with a silent BamHI

restriction site) was amplified with CwlO_UF_LysM and CwlO_UR_LysM and cloned into

pdG780 via the NotI and BamHI restriction sites to create plasmid pLU9. Secondly,

downstream homology was amplified with the primer pair CwlO_DF_LysM and

CwlO_DR _LysM and cloned into the vector pLU9 via the XhoI and KpnI restriction sites

resulting in plasmid pLU10.

To obtain plasmid pLU37, harbouring a variant *lytE* open reading frame without a LysM

domain, a fragment spanning the signal sequence, the spacer and the sequence of the

catalytic domain was amplified with the primers LytE_Spacer_Fwd_Sig /

Catalytic_LytE_Rev_XmaI and cloned into pLU10 via the XmaI and BamHI sites.

Three plasmids, that harbour a variant *lytE* open reading frame with a single LysM

domain each, were obtained in a similar way, following a two-step process. (i) First, the

three LysM domains sequences were amplified with the primer pairs

L1_Fwd_Sig_BamHI / L1_Rev_Spac, L2_Fwd_Sig_BamHI / L2_Rev_Spac and

L3_Fwd_Sig_BamHI / L3_Rev_Spac. (ii) Secondly the *lytE* catalytic domain was

amplified with the primer pair Catalytic_LytE_fwd_spac / Catalytic_LytE_rev_XmaI. Then

the fragments generated in (i) were joined individually to the PCR fragment (ii), utilizing

the spacer sequence as overlapping region. The final fragments were then cloned into

pLU10 after digestion of the vector and the fragments with XmaI and BamHI to generate

the plasmids pLU23 (*lytE[L1cat]*), pLU24 (*lytE[L2cat]*) and pLU25 (*lytE[L3cat]*).

Nine plasmids were generated, harbouring a variant *lytE* open reading frame with two

LysM domains each. To do so, the three LysM domains of *lytE* were amplified with two

primer pairs each (creating six PCR fragments in total). One primer pair generated part

of the *lytE* signal sequence, followed by the LysM domain, followed by the linker

sequence (L1_fwd_sig_BamHI / L1_rev_link, L2_fwd_sig_BamHI / L2_rev_link,

L3_fwd_sig_BamHI / L3_rev_link). The second primer pair generated the linker

sequence followed by the LysM domain, followed by parts of the spacer

sequence (L1_fwd_link / L1_rev_spac, L2_fwd_link / L2_rev_spac, L3_fwd_link /

L3_rev_spac). For cloning purposes, a silent BamHI restriction site was introduced into

the signal sequence and a silent SpeI restriction site was integrated in the linker

sequence. The fragments were then combined and joined to the *lytE* catalytic sequence

as described before and cloned into the parental vector pLU10, digested with BamHI

and XmaI. This process resulted in the plasmids: pLU14 (*lytE[L13cat]*),

pLU15 (*lytE[L12cat]*), pLU16 (*lytE[L11cat]*), pLU17 (*lytE[L21cat]*), pLU18 (*lytE[L22cat]*),

pLU19 (*lytE[L23cat]*), pLU20 (*lytE[L31cat]*), pLU21 (*lytE[L32cat]*) and

pLU22 (*lytE[L33cat]*)

Plasmids for the generation of variant LytE with three LysM domains were obtained by

cloning PCR fragments of single LysM domains into vectors already harbouring a *lytE*

open reading frame with two LysM domains. The three individual LysM domain

sequences of *lytE* were amplified with the following primer pairs: L1_fwd_ link / L1_rev_

link (L1), L2_fwd_link / L2_rev_link (L2), L3_fwd_ link / L3_rev_link (L3). The individual

LysM domains could then be cloned into the parental vectors after digestion of the

vectors and the fragments with SpeI. The number and orientation of integrated domains

was verified by a screening PCR and sequencing. Cloning the L1 fragment in pLU20,

pLU19, pLU18 and pLU14 resulted in the plasmids pLU26, pLU29, pLU31 and pLU38.

Cloning the L2 fragment in pLU18 and pLU14 resulted in the plasmid pLU30 and pLU40.

Cloning the L3 fragment in pLU20 and pLU19 resulted in the plasmids pLU27 and

pLU28.

To create plasmids that would allow the overexpression and purification of variant LytE

proteins in *E. coli*, the respective protein coding sequences (excluding the start codon

and the *lytE* signal sequence) were amplified with the forward primers Exp_L1_Fwd,

Exp_L3_Fwd, Exp_L1_Fwd, Exp_L1_Fwd, Exp_L3_Fwd and Exp_Spac_Fwd from the

plasmids pLU40, pLU20, pLU14, pLU23, pLU25 and pLU37 to generate the plasmids

pLU47, pLU49, pLU48, pLU50, pLU51 and pLU52. The same reverse primer

(Exp_Cat_Rev) was used in all reactions. PCR fragments were cloned into the

NdeI/XhoI sites of the vector pET21b+ thereby re-generating the start codon and

generating a C-terminal fusion between the protein and a 6xHis-tag.

RNA extraction and RT-qPCR

RNA was extracted and transcripts were quantified using RT-qPCR as previously

described by Salzberg *et* al., 2013). The following primer were used to amplify the

transcripts: 16s (16S_Bsu_F / 16S_Bsu_R), *cwlO* (YvcE_5´_qPCR_orf*2 / YvcE_3

´_qPCR_orf*2), *lytE* (LytE_qPCR_F2 / LytE_qPCR_R2), *lytF* (LytF_qPCR_fwd /

LytF_qPCR_rev), *mreBH* (MreBH_qPCR_F / MreBH_qPCR_R) and *sigI*

(SigI_Chip_fwd / SigI_Chip_rev).

Western analysis

Western analysis was performed as described in Noone *et al.* (2012).

Immunofluorescence microscopy

Immunofluorescence microscopy was performed as described previously (Noone *et al.*, 2012).

Protein expression and purification

The variant LytE proteins were expressed in *E. coli* strain BL21(DE3) which harboured plasmids pLU47 to pLU53. Protein expression and purification was performed as described previously (Howell *et al.*, 2006).

Insoluble peptidoglycan pull down assay

To determine the binding capacity of variant LytE proteins to peptidoglycan an insoluble peptidoglycan pull down assay was performed as described by Wong *et al.* (2014). Purified cell wall of *B. subtilis* 168 was prepared as described in Salzberg and Helmann (2007) (the number of washes was increased to 15). Forty-four pmoles of variant LytE protein were mixed with purified cell wall of *B. subtilis* in assay buffer (50 mM Tris/HCl, pH 8, 0.05 M NaCl, 5 mM ß-mercaptoethanol, 1 % TWEEN20) in a 25 µl reaction volume. Purified cell wall was added until OD600 measurements of the mixture reached a final absorbance of 10. For a negative control, 44 pmole of the wild type LytE protein were mixed with the assay buffer, without added cell wall material. All mixtures were incubated for 10 minutes at 10°C and centrifuged for 10 minutes at 10500 g. Both, the supernatant and the pellet were collected. The pellet was resuspended in 15 µl assay buffer.

The level of LytE protein that bound to the cell wall, and that which remained in the

supernated was established on a 12.5 % SDS / PAGE gel. The proteins were visualized

by coomassie blue staining and the image was quantified with the ImageJ software

(Schneider *et al.*, 2012). To test the influence of salt on the binding of variant LytE

proteins to cell walls, NaCl were added to the assay buffer at final concentrations

of 0.05 M, 0.1 M, 0.2 M, 0.3 M, 0.4 M, 1 M. Two replicates of the pull down assay at 1 M

NaCl were conducted to ensure the reproducibility of the assay. A negative control was

included in all experiments.

RESULTS:

Bioinformatic analysis of the LysM domains

There are seven D,L-endopeptidases encoded in the *B. subtilis* genome, three of which

contain LysM domains (*cwlS*, *lytE*, *lytF*). These three enzymes are structurally very

similarly and their C-terminally located catalytic domains exhibit a high sequence

similarity. However, they distinguished by the number of LysM domains with LytE, CwlS

and LytF having three, four and five N-terminally located domains respectively. Although

the high sequence similarity of their catalytic domains indicates a close evolutionary

relationship, the manner in which these three enzymes evolved to have different

numbers of LysM domains is unknown. Therefore, we sought to establish the

evolutionary relationship between the CwlS, LytE and LytF D,L-endopeptidases and to

investigate the relationships between their LysM domains, by identifying orthologs of

these enzymes in closely related bacterial species and comparing their LysM domains

through cluster analysis.

Orthologs of CwlS, LytE and LytF were identified in 19 *Bacillus* species by the levels of

amino acid identity shown by their catalytic domains (the complete list of proteins is

shown in Table S1 of Supplementary Material). Results showed that all strains

possessed an ortholog to LytE and LytF. However, only 13 of the 19 Bacillus species

encoded a CwlS ortholog. Interestingly, the number of LysM motifs within these

orthologs ranged from two to five. By these characteristics the investigated strains could

be divided into groups that agreed with their respective phylogenetic lineages.

Group (A) is comprised mainly of *Bacillus licheniformis* and *Bacillus paralicheniformis*

strains. Members do not possess a CwlS ortholog, but encode a LytE and a LytF

orthologs with 2 and 4 LysM domains respectively. Group (B) is comprised mainly of

Bacillus amyloliquefaciens strains, each having a CwlS ortholog with 4 LysM domains,

and LytE and LytF orthologs with 2 and 3 LysM domains respectively. Group C is

completely comprised of *Bacillus subtilis* strains each encoding a CwlS (4 LysM

domains), LytE (3 LysM domains) and LytF orthologs (5 LysM domains). Interestingly,

Bacillus subtilis strain XF-1 is unusual, in that it has a CwlS orthologs with 3 instead of 4

LysM domains (group X). The relationships between the 185 LysM domains in this

cohort of LytE, LytF and CwlS enzymes was established based on an alignment of their

amino acid sequence, followed by estimation of a phylogenetic tree with Maximum

Likelihood method. The results (Figure 1), supported by high bootstrap values show that

(i) individual LysM domains cluster according to their position in the protein relative to

the catalytic domain; (ii) the N-terminal LysM domain is significantly diverged from those

at the other positions; (iii) LysM domains group according to the enzyme type of their

respective catalytic domains (i.e. LytE, LytF, CwlS) and (iv) the LysM domains of LytF

and CwlS are more similar to each other than to those of LytE. The three LysM domains

of the CwlS orthologue of *Bacillus subtilis* XF-1 (M4KSM2) are in that each clusters with

the LysM domains at positions 2, 3, and 4 respectively, rather than with those at

positions 1, 2, and 3. Since M4KSM2 is the only CwlS ortholog identified with three

instead of four LysM domains, it is likely that this enzyme has undergone a deletion,

losing the N-terminal LysM domain.

To further investigate LysM domain duplication, the Uniprot database was interrogated

to identify the occurrence of identical LysM domains. Of ~39,000 LysM domains

identified, 46 occurred more than once, involving a total of 107 individual domains.

However the duplicated domains were located in the same protein in all instances, and

were adjacent to each other in 97 of the 107 instances discovered. This suggests that

duplication of LysM domains occur within open reading frames with no evidence of

horizontal domain transfer between proteins.

These results indicate that LysM domain expansion and reduction contributes to gene

divergence and that LysM domains are transferred through stable vertical inheritance.

LytE can fulfil the D,L-endopeptidase requirement for cell viability when

expressed from the *cwlO* promoter

As discovered by Bisicchia et al. (2007), LytE and CwlO are synthetically lethal, i.e. the

cell requires one or other of these enzymes for viability. It is important to note that

although LytE and CwlO possess similar C-terminal domains, they have different N-

terminal domains and are expressed under different circumstances. The regulation of

CwlO expression and activity is well described by Noone et al. (2014). In short, the TCS

WalRK is the sole activator for CwlO expression, which is therefore only expressed

during exponential growth, when the WalRK system is active (Salzberg *et al.*, 2013).

LytE expression is also activated by the WalRK system, but LytE expression can also be

triggered by active SigI under stress conditions (Tseng and Shaw, 2008; Tseng *et al.*,

2011; Hashimoto *et al.*, 2012).

We sought to investigate whether LytE when expressed from the *cwlO* promoter can

complement expression of CwlO. Since it was shown that the *cwlO* leader sequence

destabilizes the *cwlO* transcript (Noone *et al.*, 2014) we were also interested in seeing if

the same effect applied to the *lytE* transcript.

We decided to express LytE under the control of the *cwlO* promoter in a background

with a disrupted endogenous *lytE* gene [Figure 2, a].

In comparison to the wild type, *lytE* transcript levels in such a strain were 20 to 30-fold

elevated in exponentially growing cells as determined by RT-qPCR [Figure 2, c].

Furthermore a slight increase (~ 2 to 5-fold) in *sigI* and *mreBH* transcript levels was

observed. The removal of the *cwlO* leader sequence caused a further increase in *lytE*

transcript and a decrease in *sigI* transcript level (~ 2-fold). We confirmed that this can be

attributed to the increased longevity of the *lytE* transcript through a mRNA half-life

experiment [data not shown]. Elevated *lytE* transcript levels matched increased LytE

protein levels [Figure 2, d]. High levels of LytE protein were found both cell-associated

and secreted into the medium. However, the LytE protein appeared to be degraded in

the medium as cultures approached stationary phase.

Although removal of the *cwlO* leader sequence led to a drop in SigI levels, we observed

that overexpressing LytE at this high level affects cell morphology with a fraction of cells

showing bulges and thickening [Figure 2, b]. In short, both *lytE* null mutants as well as

strains overexpressing LytE, exhibit difficulties to maintain cell shape and longitudinal

growth.

In summary, this results confirm that the LytE D,L-endopeptidase can complement CwlO

when expressed using *cwlO* expression signals - both transcriptional (i.e. promoter) and

post-transcriptional (i.e. *cwlO* leader sequence). It is clear however that the two

enzymes are not equivalent since the SigI regulon (signalling cell wall stress), is induced

when *lytE* is expressed using the normal *cwlO* expression signals (i.e. promoter and

leader sequence), but to a lesser extent when the *cwlO* leader sequence is removed.

LytF can also fulfil the D,L-endopeptidase requirement

The synthetic lethality of the CwlO / LytE autolysins allows further investigation of

whether any other D,L-endopeptidase encoded in the genome can complement their

activity. Thus, it can be investigated which D,L-endopeptidase can support growth and

viability when expressed under the control of the *cwlO* promoter.

In view of these findings, we sought to establish if the D,L-endopeptidases LytF and

CwlS, - which we previously showed to be closely related to LytE (LysM domains), as

well as another autolytic enzyme, - YqgT (no LysM domain), can fulfil the D,L-

endopeptidase requirement when expressed using the *cwlO* promoter.

To establish whether the CwlS, LytF or YqgT D,L-endopeptidases when expressed from

the *cwlO* promoter can fulfill the D,L-endopeptidase requirement, we constructed strains

in which the *cwlS, lytF* and *yqgT* genes were individually expressed at the *cwlO* locus.

The level of transcript of each gene was varied by expressing each gene both with the

cwlO leader sequence (giving an unstable transcript) and without the leader

sequence (giving a stable transcript). The ability of each enzyme to fulfil the D,L-

endopeptidase requirement for viability was then established by attempting to disrupt the

lytE gene in each of the strains, which was successful for *lytF* but not for *cwlS* or *yqgT*.

LytF levels in the strains expressing LytF from the *cwlO* promoter were highly elevated

when compared to that observed in strains, in which the enzyme was expressed from its

natural promoter [Figure 3, b]. Despite the high level of cellular *lytF* mRNA transcript,

only a very low level of full-length LytF protein was cell associated, with a high fraction

found to be processed [Figure 3, c]. HA-tagged LytF protein was predominantly localized

at the cell septa and cell poles, but was also present along the lateral cell cylinder

[Figure 3, d]. Based on the *sigI* transcript levels, it seems that significant cell wall stress

is not generated in these strains [data not shown]. However, strains overexpressing LytF

from the *cwlO* promoter displayed major growth defects (delayed lag phase) and cell

morphology defects [Figure 3, a].

Thus, while cell viability can be achieved by replacing the CwlO and LytE D,L-

endopeptidases with expression of LytF, cells are not robust.

The individual LysM domains of LytE are not functionally equivalent

We could show that LytE (3 LysM domains) and LytF (5 LysM domains), but not CwlS (4

LysM domains) can support growth of *B. subtilis*, when expressed from the *cwlO*

promoter. Interestingly although the majority of LysM domain containing proteins in the

dataset constructed in section one possess only a single LysM domains (~ 67 % of

18,246 proteins), all identified orthologs to LytE, CwlS and LytF feature at least 2 LysM

domains.

There is little known about how individual LysM domains bind to peptidoglycan and the

significance of the fact that the identified D,L-endopeptidase orthologs possess a

minimum of 2 LysM domains.

Therefore we sought to investigate the properties of the individual LysM domains of LytE

by designing LytE enzymes with a single LysM domain. We then tested the

characteristics of these LytE variants proteins in strains which viability was depended on

the permutated autolysin. This was achieved by expressing the variant LytE containing

either L1, L2 or L3 in single copy at the *cwlO* or the *lytE* locus and attempting to disrupt

either the endogenous *lytE* or the *cwlO* gene [Supplementary Figure 2]. For control

purposes, we also included a variant LytE without any LysM domain as well as a

reconstituted wild type LytE with the three LysM domains in the wild type combination

(L1-L2-L3) in the further analysis.

We found that all constructed strains were viable, demonstrating that any single LysM

domain of LytE is capable of inferring functionality to the LytE enzyme. A control strain

with a LytE protein without a LysM domain could not be constructed, demonstrating that

the LytE enzyme requires at least one LysM domain for functionality.

Despite the fact that any individual LysM domain enabled a functional LytE enzyme, we

detected a significant change in cell morphology with shorter and thicker cells for the

variant strains. Interestingly this effect was greater for strains containing a LytE enzyme

with L2 or L3 than L1 [Figure 4, a].

We suggest that the extent to which a variant LytE protein fulfils the D,L-endopeptidase

requirement can be assessed by determining the extent to which the SigI regulon is

induced. This was determined by establishing the mRNA transcript levels of the SigI

regulon genes *sigI* and *mreBH* by RT-qPCR. We observed that transcript levels were

elevated if a strain contained a variant LytE with a single LysM domain with the

combination L2, L3, however no elevation occurred with L1 or with the reconstituted wild

type protein [Figure 4, b].

We further decided to test the robustness of these strains by determining their sensitivity

to salt stress. As expected, all controls grew normally at NaCl concentrations up to

0.4 M. In contrast, although all variant LytE strains grew comparable to wild type under

the absence of salt or at a salt concentration of 0.1 M, growth inhibitions were observed

at a salt concentration of 0.2 M and greater [Figure 4, c]. The extent, to which salt stress

caused a change in growth behaviour, correlated with the type of LysM domains as well

as the expression level (*cwlO* vs *lytE* promoter). Higher expression levels (*cwlO*

promoter) improved the resistance of strains to salt stress. Salt tolerance was also

increased when the LytE enzyme contained the first LysM domain (L1) as compared to

L2 and L3. Since sensitivity to salt stress was not altered in the control strains which

were in possession of an intact copy of the endogenous *lytE* or *cwlO* gene, we conclude

that it is not the presence of the variant LytE that causes the increased sensitivity to salt

stress, but rather a lack of D,L-endopeptidase enzymatic activity.

Our results indicate that although a variant LytE protein with any individual LysM domain

is functional, it does not fulfil the D,L-endopeptidase requirement to the same extent as

the wild type LytE protein. Cells that express a variant LytE with a single LysM domain,

under the absence of the endogenous *cwlO* and *lytE* gene display growth and

morphology defects. Furthermore our experiments show that the three LysM domains of

LytE are not functionally equivalent and that L1 is capable of conferring on the catalytic

domains the ability to fulfil the D,L-endopeptidase requirement to a greater extent than

L2 and L3 as can be seen from the induction of the *sigI* regulon.

LysM domain requirement of LytE

We observed that the three LysM domains are not equal in their ability to confer

functionality to the LytE enzyme with the N-terminally located LysM domain L1 being

superior to L2 and L3. It is not clear what properties distinguish the first LysM domain

from the subsequent LysM domains, other than its location adjacent to the signal

sequence domain.

We therefore sought to investigate the effect of the order as well as the number of LysM

domains on the functionality of the LytE enzyme.

This was achieved by expressing variant LytE enzymes with two or three LysM domains

in a permutated order from either the *cwlO* or the *lytE* promoter [Supplementary

Figure 2]. Functionality of these enzymes was then tested by disrupting the endogenous

lytE or *cwlO* gene. We could show that any LytE variant with two LysM domains in any

of the nine possible combinations resulted in a functional enzyme. The same was true

for two variant LytE with three LysM domains in the combinations L3-L3-L1 and L2-L1-

L3.

Although all strains were viable, we observed that strains expressing variant LytE

proteins with three LysM domains produced bigger colonies than those that expressed

variant LytE proteins with two LysM domains [Figure 5, a]. Strains containing a variant

LytE protein with a single LysM domain displayed the smallest colonies, which also

exhibited strong lysis over the course of the experiment. In general, colonies were

bigger and experienced less lysis when they were expressed using the stronger *cwlO*

promoter rather than the weaker *lytE* promoter. Interestingly the order of LysM domains

also mattered. Those strains that expressed a variant LytE with L1 at the first (N-

terminal) position displayed larger colonies, than those in which either L2 or L3 were

located adjacent to the signal sequence domain. This correlation was especially strong

for LytE variants with a set of 2 LysM domains, as can be seen when comparing L1-L3

to L3-L1.

To address whether the type of variant LytE protein has an effect on the cellular protein

level, the level of HA-tagged variant LytE protein was determined by western blot

analysis. All variant LytE-HA proteins could be detected and bands migrated according

to the predicted MW [Figure 5, b]. The level of variant LytE protein varied significantly in

the strains tested. Since the number of LysM domains is proposed to correlate with

binding affinity, we expected to find more cell-associated protein in variant proteins with

a greater number of LysM domains. Indeed, the level of cell-associated variant LytE

protein was higher for proteins with two or more LysM domains, than for proteins with a

single LysM domain. If it is assumed that the lower level of cell-associated LytE is

caused by lowered binding to the cell wall, and not lowered expression, then we predict

the level of variant LytE protein found in the medium to have the reverse profile of that

for cell-associated protein. To verify this, we compared the levels of cell-associated

variant LytE proteins with those in the culture supernatant. A strong inverse correlation

in the level of variant LytE was observed. Variant LytE proteins with one LysM domain

were predominantly found in the culture supernatant, while variant LytE proteins with

three LysM domains were predominantly found in association with the cell wall.

Moreover the distribution of variant LytE proteins with two LysM domains takes an

intermediate between that found for those with three and one LysM domains. We

confirmed these results by immunofluorescence microcopy, which demonstrated that HA

tagged variant LytE with more LysM domains gave a stronger fluorescence signal than

those with fewer LysM domains [data not shown]. Interestingly, the pattern of

localization along the cell cylinder was not affected.

We further tested binding affinity of variant LytE protein in vitro in an insoluble

peptidoglycan pull down assay. To investigate the effect of salt on the peptidoglycan

binding affinity we added increasing amounts of NaCl to the assay buffer (0.05 M - 1

M NaCl) [Figure 5, c]. At a low salt concentration of 0.05 M NaCl, all variant LytE

proteins showed an equally strong binding capacity of approximately 100 % (proportion

of protein found in the pellet). However, no binding could be detected for the variant

LytE protein without LysM domain. Furthermore our results clearly show that the

peptidoglycan-binding capacity is negatively correlated to the salt concentration. With an

increase in salt concentration (≥ 0.1 M NaCl) binding capacity for the variant LytE

proteins is decreased. The rate at which binding capacity is declined varies between

proteins and is in negative correlation to the number of LysM domains present. Proteins

with more LysM domains exhibited a greater robustness of binding affinity than proteins

with a single LysM domain.

In summary, this study demonstrates that the number and order of LysM domains are of

great relevance for the functionality of the LytE enzyme. Although a LytE enzyme with

any one of its three LysM domains is functional and can fulfil the D,L-endopeptidase

requirement, increasing the number of LysM domains to two or three has a positive

effect on normal colony morphology. The order in which the domains are arranged is

also of importance in that variant LytE proteins in which the first LysM domain (L1) is

localized in its natural position adjacent to the signal peptide fulfil the D,L-endopeptidase

requirement to a greater extent than L2 and L3.

Furthermore, the number of LysM domains of a LytE variant protein is positively

correlated with peptidoglycan binding affinity and a good determinant of the extent to

which the protein is cell-associated or shed into the culture media. We could show that

an increase of the number of LysM domains leads to increased binding affinity both *in*

vivo and *in vitro*. We further demonstrated that binding affinity is negatively affected by

salt *in vitro*.

DISCUSSION:

The WalRK and SigIRsgI systems work in tandem to ensure adequate expression

of D,L-endopeptidase activity

Expression of D,L-endopeptidase activity is required for normal growth and viability of

B. subtilis as reported by Bisicchia et al. (2007). Under normal growth condition this

activity is provided by CwlO and under stress conditions by the LytE enzyme. We

demonstrated in this study that the level of *lytE* transcript and LytE protein is significantly

increased when the *cwlO* gene is deleted, probably to compensate for the loss in D,L-

endopeptidase activity. Since *lytE* is part of the SigI regulon (Tseng *et al.*, 2011), we

suggest that this is achieved through activation of the alternative SigI factor as proposed

by Salzberg and colleagues (2013). A similar response was observed by Kasahara and

colleagues (2016) who reported that *lytE* transcription was considerably increased in

strains with defects in teichoic acid synthesis, notably in a *ltaS* null mutant and who

suggest that upregulation is achieved in a SigI dependent manner. Thus, the WalRK and

SigIRsgI regulators form a regulatory network that functions to ensure that there is an

adequate level of D,L-endopeptidase enzyme present under normal growth (CwlO) and

cell wall stress (LytE) conditions.

It was of interest to establish whether LytE can fulfil the D,L-endopeptidase requirement

for viability when expressed from the *cwlO* promoter in the absence of the endogenous

lytE gene (thus removing SigI coupling). It was found that such a strain is viable,

signifying that changing the regulation of LytE expression from its natural promoter to

the *cwlO* promoter is compatible with viability and indeed normal growth. We observed

that LytE expression was higher when driven by the *cwlO* promoter as compared to the

endogenous *lytE* promoter (un-induced or induced). LytE expression (stabilized

transcript and increased protein level) was further increased by deletion of the *cwlO*

leader sequence due to the destabilizing effect of the leader transcript (Noone *et al.*,

2014). Surprisingly, we detected a slight increase in *mreBH* and *sigI* when *lytE* was

expressed from the *cwlO* promoter with the *cwlO* leader sequence. Based on the

premise that *sigI* is released under stress, removal of the *cwlO* leader sequence is

beneficial for the cell, since it results in a decrease of *sigI* levels. However, at the same

time removal of the *cwlO* leader sequence caused minor deficiencies in cell morphology,

indicating that such an elevated level of autolytic activity is detrimental to the cell. The

time point at which these phenotypic defects were observed, correlates with the high

level of WalRK activity during these growth phases (Prunty *et al.*, 2018)

In summary, these data suggest that although expression of LytE from the *cwlO*

promoter can fulfil the D,L-endopeptidase requirement for cell viability, it does not

complement CwlO function completely and that further elevation of LytE protein levels

are required (i.e. by removal of the leader sequence) to achieve a state without a cell

wall stress (i.e. no elevation of *sigI* expression).

Expression levels and localization of D,L-endopeptidases within the cell envelope

are critical

We investigated four of the seven D,L-endopeptidases encoded in the *B. subtilis*

genome (LytE, LytF, CwlS, YqgT) and found that CwlS and YqgT cannot complement

CwlO / LytE activity when expressed from the *cwlO* promoter. LytE naturally fulfils the

D,L-endopeptidase requirement in a *cwlO* mutant, but can also do so when expressed

from the *cwlO* locus. LytF cannot complement CwlO / LytE activity when expressed from

its natural *lytF* promoter, but gives partial complementation when expressed from the

cwlO promoter. The fact that LytF is unable to fulfil the D,L-endopeptidase requirement

for viability when expressed from the *lytF* promoter, but can do so when expressed from

the *cwlO* promoter, indicates that complementation requires either greater than normal

LytF protein levels and / or LytF expression at a different time point throughout the

growth cycle. However, although a viable strain could be constructed, severe growth

and morphology defects were observed. Exacerbation of the aberrant phenotype in

strains without *cwlO* leader sequence (causing a further increase in LytF levels)

suggests that the level of LytF in this background may be too high for normal growth.

We do not know why neither *cwlS* nor *yqgT* were able to fulfil the D,L-endopeptidase

requirement. However, we ruled out the possibility of a synthetic lethality between these

enzymes by constructing viable double mutants.

The fact that the LytE, LytF and CwlS enzymes are highly similar in their C-terminal D,L-

endopeptidase catalytic domains, as well as in their N-terminally located LysM domains

poses the question as to what causes the inability of CwlS to fulfil the D,L-

endopeptidase requirement. We confirmed that there is indeed a high level of CwlS

expressed and associated with the cell, when expressed from the *cwlO* promoter [data

not shown]. However, it may be of relevance that the CwlS protein is primarily located at

the cell septa (Hashimoto *et al.*, 2012), while both LytE and LytF have some presence

along the cell cylinder (Hashimoto *et al.*, 2012; Kiriyama *et al.*, 2014; Kasahara *et al.*,

2016). This is significant since it has been established that the D,L-endopeptidase

requirement is associated with facilitating cell cylinder elongation (Bisicchia *et al.*, 2007).

Thus, there may not be a sufficient level of CwlS located along the cylinder to facilitate

cell elongation. This is supported by the observations that cells expressing *cwlS* from

the *cwlO* promoter appear to be shorter than wild-type cells and to have unusual

structures at the poles of the cell (data not shown).

Although both LytE and LytF show some binding to the lateral cell wall, LytF is primarily

bound to the cell septa and poles, which potentially explains why LytF achieves only

partial complementation in comparison to the LytE enzyme. In summary, these results

show that beside the level of D,L-endopeptidase expression, their proper subcellular

localization are paramount for normal growth.

The individual LysM domains of LytE are functionally and structurally distinct

We could show that strains expressing a LytE variants with any single LysM domain are

viable. However, we observed cell morphology, colony morphology and growth defects

in such strains. These phenotypic defects were more pronounced in LytE variants with a

single copy of the second (L2) or third (L3) LysM domains as opposed to the N-terminal

located L1, indicating that the individual LysM domains are not functionally equivalent.

The increased effectiveness of the first LysM domain (L1) is probably due to the fact that

it has evolved to be more suitable for interaction with the adjacent signal sequence

domain, which perhaps causes the enzyme to be expressed or exported more

efficiently.

This finding was confirmed in later experiments when we could show that in strains that

express a variant LytE proteins with two or three LysM domains, the order of the

domains was relevant for proper colony growth. Variant LytE proteins that contained the

first LysM domain (L1) at the N-terminal position resulted in strains that displayed

healthier colonies which experienced less lysis than variant LytE in which either L2 or L3

were located adjacent to the signal peptide. A possible mechanism by which a LytE

protein with impaired functionality could cause colony growth defects was investigated

by Mamou and colleagues (2017), who report that LytE is involved in proper colony

development through influencing Y arm extension and colony thickening patterns.

The fact that the first LysM domain is structurally different from the subsequent domains

was further supported through bioinformatic analysis. By comparing the amino acid

sequence of the 185 LysM domains contained in the LytE, CwlS and LytF orthologs of

19 closely related *B. subtilis* species we found that the N-terminal located LysM domains

were the most distinct. This may indicate that due its interaction with the signal peptide

there is a greater selection pressure on the N-terminal LysM domain than on

subsequent domains.

LysM domains contribute to peptidoglycan binding in vivo and in vitro

We could show that a variant LytE without LysM domains exhibited no binding affinity to

purified cell wall *in vitro* and that it could not sustain cell growth. Thus, LysM domains

are essential for cell wall binding and therefore functionality of the LytE enzyme. This

observation is supported by several previous studies that have shown that a total loss of

LysM domains greatly decreases the peptidoglycan-binding capacity in vitro (Steen *et*

al., 2005; Eckert *et al.*, 2006; Wong *et al.*, 2014). Increasing the number of LysM

domains from one to three, led to increased binding capacity both *in vivo* and *in vitro*,

suggesting that individual LysM domains contribute to binding in a accumulative

manner, as was proposed previously (Mesnage *et al.*, 2014; Wong *et al.*, 2014).

Surprisingly, strains expressing a LytE variant with any single LysM domain are viable,

suggesting that the peptidoglycan-binding affinity of the wild type LytE protein with three

LysM domains is excessive. However, our experiments were performed under

laboratory conditions and we observed that such strains have elevated stress sensitivity,

especially to salt stress. We demonstrated *in vitro* that salt causes a decrease in binding

capacity of variant LytE proteins, especially those with fewer than three LysM domains.

Therefore, growth defects in the presence of salt are probably due to insufficient binding

of the variant LytE D,L-endopeptidase under this circumstances. Given that *B. subtilis* is

a soil-dwelling organism, its capability to withstand salt stress is essential for its survival.

Therefore, cells that possess a LytE enzyme with less than three LysM domains would

not be able to survive natural fluctuations in the salt concentrations in the soil.

Expansion of the LysM domains and evolution of the CwlS, LytE and LytF

orthologs

We observed that the LysM domains of the CwlS, LytE and LytF orthologs cluster

primarily by position, with the N-terminally located LysM domains forming a remarkably

distinct cluster. This finding is contrary to a similar study conducted for fungal proteins.

In this study, Cen and colleagues (2017) report that the LysM domains of fungal

effectors proteins cluster independently of their sequential position within the parental

protein or protein structure family. We suggest that this difference stems from the fact

that the LytE, CwlS and LytF proteins have evolved fairly recent, leaving their LysM

domains little time to diverge.

We were further interested in the manner in which D,L-endopetidases acquire new LysM

domains. By screening 26.078 bacterial proteins for identical LysM domain amino acid

sequences we could show that LysM domains are primarily propagated through

duplication events that occur within a gene. Interestingly, the majority of duplicated LysM

domains were present within the phylum Firmicutes (79.4 %). This is a much higher

percentage than could be expected, provided that domain expansion occurred in all

phyla at the same rate (LysM domains of the Firmicutes phylum represent 32.8 % of the

total dataset), suggesting that LysM domain expansion is either a more recent or more

frequent event in species of the Firmicutes phylum. Interestingly, an uncharacterised

protein of *Bacillus subtilis* subsp. *natto* BEST195 (D4G4L0) was found to possess 6

LysM domains, that are of identical amino acid sequence and highly similar (although

not identical) on nucleotide level. This indicates that LysM domain duplication events

can create several repeats of a single domain at once, leading to the rapid expansion of

the domain repeats.

Based on these findings, we propose a model for the evolutionary development of the

three D,L-endopeptidases and their LysM domains. Our model is based on the premise

that LytE, CwlS and LytF share a common ancestor and that LysM domain repeats

expand independently within genes and are passed on through stable vertical

inheritance. In this model an ancestral gene with a single N-terminally located LysM

domain, undergoes a combination of gene and LysM domain duplication / loss events,

which lead to the development of the three orthologs and their observed LysM domain

combinations [Figure 6, a]. This model can also be used to infer basic phylogeny

between the investigated *Bacillus* species [Figure 6, b]*,* which is in line with their overall

phylogenetic relationships [Supplementary Figure 1].

Conclusion

Overall this study offers further insights into the D,L-endopeptidase requirement of

B. subtilis. To our knowledge this is the first work that has demonstrated that the D,L-

endopeptidase requirement of *B. subtilis* can be fulfilled by a D,L-endopeptidase other

than LytE or CwlO. It remains unknown why LytF, but not CwlS is able to complement

for CwlO, given their highly similar structure (also localization is a likely suspect).

This is also the first time that is has been investigated whether individual LysM domains

are of equal importance for the functionality of an autolytic enzyme in vivo. It was shown

that any single LysM domain is sufficient for functionality of the LytE enzyme, however

robust growth requires the three LysM domains present in the wild type enzyme. Among

these we demonstrated that the first LysM domain is superior over the subsequent

domains, especially when located N-terminally, which is probably due to some

interaction with the signaling peptide.

An investigation of bacterial LysM domains on a large scale led to some interesting

observations, such as that expansion of LysM domains occurs frequently in the

Firmicutes phylum and that LysM domains are unlikely to be transferred horizontally.

Accordingly, a putative model was developed describing the relationship between the

LytE, CwlS and LytF D,L-endopeptidases.

It is clear from our results that D,L-endopeptidases must have many features to be able

to fulfill the requirement for viability. Among those requirements are the level of enzyme

expressed, the stability of the protein when expressed and the localization of the protein

within the cell envelope.

Further investigating these and other aspects of autolytic enzymes and cell wall

metabolism will not only deepen our understanding of these processes, but also has the

potential to be utilized for the development of new antibiotics and for the optimization of

bacterial strains for enzyme and protein production. Furthermore an in-depth

understanding of LysM domains and conditions that affect their biding affinity is likely to

be of importance for the development of future binding assays and cell surface displays.

References

Anagnostopoulus, C., and Spizizen, J. (1960). Requirements for Transformation in
Bacillus subtilis. 81, 741–746. Available at: C:\Users\User\Documents\Citavi 6\
Projects\LytE\Citavi Attachments\Anagnostopoulus, Spizizen 1960 -
Requirements for Transformation in Bacillus.pdf.

Bisicchia, P., Noone, D., Lioliou, E., Howell, A., Quigley, S., Jensen, T., et al. (2007).
The essential YycFG two-component system controls cell wall metabolism in
Bacillus subtilis. *Mol. Microbiol.* 65, 180–200. doi:10.1111/j.1365-
2958.2007.05782.x.

Bloor, A. E., and Cranenburgh, R. M. (2006). An Efficient Method of Selectable Marker
Gene Excision by Xer Recombination for Gene Replacement in Bacterial
Chromosomes. *Appl. Environ. Microbiol.* 72, 2520–2525. Available at: C:\Users\

User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Bloor and Cranenburgh 2006.pdf.

Bodenhofer, U., Bonatesta, E., Horejs-Kainrath, C., and Hochreiter, S. (2015). msa: an R package for multiple sequence alignment. *Bioinformatics* 31, 3997–3999. doi:10.1093/bioinformatics/btv494.

Botella, E., Hübner, S., Hokamp, K., Hansen, A., Bisicchia, P., Noone, D., et al. (2011). Cell envelope gene expression in phosphate-limited Bacillus subtilis cells. *Microbiology* 157, 2470–2484. doi:10.1099/mic.0.049205-0.

Boylan, S. A., Redfield, A. R., Brody, M. S., and Price, C. W. (1993). Stress-induced activation of the sigma B transcription factor of Bacillus subtilis. *J. Bacteriol.* 175, 7931–7937. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Boylan, Redfield et al 1993 - Stress-induced activation of the sigma.pdf.

Buist, G., Steen, A., Kok, J., and Kuipers, O. P. (2008). LysM, a widely distributed protein motif for binding to (peptido)glycans. *Mol. Microbiol.* 68, 838–847. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\ Buist 2008.pdf.

Cen, K., Li, B., Lu, Y., Zhang, S., and Wang, C. (2017). Divergent LysM effectors contribute to the virulence of Beauveria bassiana by evasion of insect immune defenses. *PLOS Pathog.* 13, e1006604. doi:10.1371/journal.ppat.1006604.

Desvaux, M., Candela, T., and Serror, P. (2018). Surfaceome and Proteosurfaceome in Parietal Monoderm Bacteria: Focus on Protein Cell-Surface Display. *Front. Microbiol.* 9. doi:10.3389/fmicb.2018.00100.

Dominguez-Cuevas, P., Porcelli, I., Daniel, R., and Errington, J. (2013). Differentiated roles for MreB-actin isologues and autolytic enzymes in bacterial morphogenesis. *Mol. Microbiol.* 89, 1084–1098. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Dominguez-Cuevas, Porcelli et al 2013 - Differentiated roles for MreB-actin isologues.pdf.

Eckert, C., Lecerf, M., Dubost, L., Arthur, M., and Mesnage, S. (2006). Functional analysis of AtlA, the major N-acetylglucosaminidase of Enterococcus faecalis. *J. Bacteriol.* 188, 8513–8519. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Eckert, Lecerf et al 2006 - Functional analysis of AtlA.pdf.

Hashimoto, M., Ooiwa, S., and Sekiguchi, J. (2012). Synthetic Lethality of the lytE cwlO Genotype in Bacillus subtilis Is Caused by Lack of D,L-Endopeptidase Activity at the Lateral Cell Wall. *J. Bacteriol.* 194, 796–803. doi:10.1128/JB.05569-11.

Hayhurst, E. J., Kailas, L., Hobbs, J. K., and Foster, S. J. (2008). Cell wall peptidoglycan architecture in Bacillus subtilis. *Proc Natl Acad Sci* 105, 14603–14608. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Hayhurst et al 2008.pdf.

Horton, R. M., Hunt, H. D., Ho, S. N., Pullen, J. K., and Pease, L. R. (1989). Engineering

    hybrid genes without the use of restriction enzymes: gene splicing by overlap

    extension. *Gene* 77, 61–68.

Howell, A., Dubrac, S., Noone, D., Varughese, K. I., and Devine, K. M. (2006).

    Interactions between the YycFG and PhoPR two-component systems in Bacillus

    subtilis: the PhoR kinase phosphorylates the non-cognate YycF response

    regulator upon phosphate limitation. *Mol. Microbiol.* 59, 1199–1215.

    doi:10.1111/j.1365-2958.2005.05017.x.

Ishikawa, S., Hara, Y., Ohnishi, R., and Sekiguchi, J. (1998). Regulation of a New Cell

    Wall Hydrolase Gene, cwlF, which affects Cell Separation in Bacillus subtilis. *J.*

    Bacteriol. 1890, 2549–2555. Available at: C:\Users\User\Documents\Citavi 6\

    Projects\LytE\Citavi Attachments\Ishikawa 1998.pdf.

Kasahara, J., Kiriyama, Y., Miyashita, M., Kondo, T., Yamada, T., Yazawa, K., et al.

    (2016). Teichoic Acid Polymers Affect Expression and Localization of dl-

    Endopeptidase LytE Required for Lateral Cell Wall Hydrolysis in Bacillus subtilis.

    J. Bacteriol. 198, 1585–1594. doi:10.1128/JB.00003-16.

Kiriyama, Y., Yazawa, K., Tanaka, T., Yoshikawa, R., Yamane, H., Hashimoto, M., et al.

    (2014). Localization and expression of the Bacillus subtilis dl-endopeptidase LytF

    are influenced by mutations in LTA synthases and glycolipid anchor synthetic

    enzymes. *Microbiology* 160, 2639–2649. doi:10.1099/mic.0.080366-0.

Mamou, G., Fiyaksel, O., Sinai, L., and Ben-Yehuda, S. (2017). Deficiency in

    Lipoteichoic Acid Synthesis Causes a Failure in Executing the Colony

    Developmental Program in Bacillus subtilis. *Front. Microbiol.* 8.

    doi:10.3389/fmicb.2017.01991.

Meisner, J., Llopis, P. M., Sham, L.-T., Garner, E., Bernhardt, T. G., and Rudner, D. Z.

    (2013). FtsEX is required for CwlO peptidoglycan hydrolase activity during cell

    wall elongation in Bacillus subtilis. *Mol. Microbiol.* 89, 1069–1083. Available at:

    C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Jeffrey

    2013.pdf.

Mesnage, S., Dellarole, M., Baxter, N. J., Rouget, J.-B., Dimitrov, J. D., Wang, N., et al.

    (2014). Molecular basis for bacterial peptidoglycan recognition by LysM domains.

    Nat. Commun. 5, 1–11. doi:10.1038/ncomms5269.

Noone, D., Botella, E., Butler, C., Hansen, A., Jende, I., and Devine, K. M. (2012).

    Signal Perception by the Secretion Stress-Responsive CssRS Two-Component

    System in Bacillus subtilis. *J. Bacteriol.* 194, 1800–1814. Available at: C:\Users\

    User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Noone 2012.pdf.

Noone, D., Salzberg, L. I., Botella, E., Basell, K., Becher, D., Antelmann, H., et al.

    (2014). A Highly Unstable Transcript Makes CwlO D,L-Endopeptidase Expression

    Responsive to Growth Conditions in Bacillus subtilis. *J. Bacteriol.* 196, 237–247.

    Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\

    Noone 2014.pdf.

Pele, J., Becu, J.-M., Abdi, H., and Chabbert, M. (2012). Bios2mds: an R package for comparing orthologous protein families by metric multidimensional scaling. *BMC Bioinformatics* 13, 1–7. doi:10.1186/1471-2105-13-133.

Prunty, M. P., Noone, D., and Devine, K. M. (2018). The distinct PhoPR mediated responses to phosphate limitation in *Bacillus subtilis* subspecies *subtilis* and *spizizenii* stem from differences in wall teichoic acid composition and metabolism: PhoPR activation in *Bacillus subtilis* subspecies. *Mol. Microbiol.* 109, 23–40. doi:10.1111/mmi.13965.

Rajagopal, M., and Walker, S. (2017). Envelope Structures of Gram-Positive Bacteria. *Curr. Top. Microbiol. Immunol.* 404, 1–44. doi:10.1007/82_2015_5021.

Salamaga, B., Prajsnar, T. K., Jareño-Martinez, A., Willemse, J., Bewley, M. A., Chau, F., et al. (2017). Bacterial size matters: Multiple mechanisms controlling septum cleavage and diplococcus formation are critical for the virulence of the opportunistic pathogen Enterococcus faecalis. *PLoS Pathog.* 13. doi:10.1371/journal.ppat.1006526.

Salzberg, L. I., Powell, L., Hokamp, K., Botella, E., Noone, D., and Devine, K. M. (2013). The WalRK (YycFG) and σ I RsgI regulators cooperate to control CwlO and LytE expression in exponentially growing and stressed Bacillus subtilis cells. *Mol. Microbiol.* 87, 180–195. Available at: C:\Users\User\Documents\Citavi 6\Projects\ LytE\Citavi Attachments\Salzberg, Powell et al 2013 - The WalRK YycFG and σ.pdf.

Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989). *Molecular cloning*. 2. ed. Cold

    Spring Harbor, NY: Cold Spring Harbor Laboratory Press.

Schliep, K. P. (2011). phangorn: phylogenetic analysis in R. *Bioinforma. Oxf. Engl.* 27,

    592–593. doi:10.1093/bioinformatics/btq706.

Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH Image to ImageJ. *Nat.*

    Methods 9, 671–675. doi:10.1038/nmeth.2089.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., et al. (2011). Fast,

    scalable generation of high-quality protein multiple sequence alignments using

    Clustal Omega. *Mol. Syst. Biol.* 7, 1–6. doi:10.1038/msb.2011.75.

Sims, G. E., Jun, S.-R., Wu, G. A., and Kim, S.-H. (2009). Alignment-free genome

    comparison with feature frequency profiles (FFP) and optimal resolutions. *Proc.*

    Natl. Acad. Sci. U. S. A. 106, 2677–2682. doi:10.1073/pnas.0813249106.

Steen, A., Palumbo, E., Deghorain, M., Cocconcelli, P. S., Delcour, J., Kuipers, O. P., et

    al. (2005). Autolysis of Lactococcus lactis Is Increased upon d-Alanine Depletion

    of Peptidoglycan and Lipoteichoic Acids. *J. Bacteriol.* 187, 114–124.

    doi:10.1128/JB.187.1.114-124.2005.

Tseng, C.-L., Chen, J.-T., Lin, J.-H., Huang, W.-Z., and Shaw, G.-C. (2011). Genetic

    evidence for involvement of the alternative sigma factor SigI in controlling

    expression of the cell wall hydrolase gene lytE and contribution of LytE to heat

    survival of Bacillus subtilis. *Arch. Microbiol.* 193, 677–685. doi:10.1007/s00203-

    011-0710-0.

Tseng, C.-L., and Shaw, G.-C. (2008). Genetic Evidence for the Actin Homolog Gene mreBH and the Bacitracin Resistance Gene bcrC as Targets of the Alternative Sigma Factor SigI of Bacillus subtilis. *J. Bacteriol.* 190, 1561–1567. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Tseng 2008.pdf.

Wang, A., and Ash, G. J. (2015). Whole Genome Phylogeny of *Bacillus* by Feature Frequency Profiles (FFP). *Sci. Rep.* 5, 13644. doi:10.1038/srep13644.

Wong, J. E. M. M., Alsarraf, H. M. A. B., Kaspersen, J. D., Pedersen, J. S., Stougaard, J., Thirup, S., et al. (2014). Cooperative binding of LysM domains determines the carbohydrate affinity of a bacterial endopeptidase protein. *FEBS J.* 281, 1196–1208. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Wong, Jaslyn E M M, Alsarraf, Husam M A B et al 2013 - Cooperative binding of LysM domains.pdf.

Yamaguchi, H., Furuhata, K., Fukushima, T., Yamamoto, H., and Sekiguchi, J. (2004). Characterization of a New Bacillus subtilis Peptidoglycan Hydrolase Gene, yvcE (Named cwlO), and the Enzymatic Properties of Its Encoded Protein. *J. Biosci. Bioeng.* 98, 174–181. Available at: C:\Users\User\Documents\Citavi 6\Projects\LytE\Citavi Attachments\Yamaguchi 2004.pdf.

Yu, G., Smith, D. K., Zhu, H., Guan, Y., and Lam, T. T.-Y. (2016). ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* 8, 28–36. doi:10.1111/2041-210X.12628.

Figure 1:
phylogentic tree of the LysM motifs

position of LysM motif   ■ 1st   ■ 2nd   ■ 3rd   ■ 4st   ■ 5st

type by catalytic domain   ■ CwlS-type   ■ LytE-type   ■ LytF-type

example of LytF type enyzme   L1 L2 L3 L4 L5 cat

Figure 2:

a) strain construction

b) microscopy

c) RT-qPCR

d) western blot

Figure 3:

a) microscopy

intact *lytE*   Δ*lytE*

LU111   LU116

PcwlOleader

LU125   LU134

PcwlO

10 μm

b) RT-qPCR

lytF

fold change in expression

WT168 LS103 LS089 LU74 LU111 LU125 LU134

bacterial strain

c) western blot

strain / time point

LU147   LU153   LU157

kDa   t1   t2 | t1   t2 | t1   t2 | nc

35 —

25 —

10 —

cell associated

d) fluorescence microscopy

PH   FITC   overlay

WT168

LU151

10 μm

Figure 4:

a) microscopy

intact *cwlO*   Δ*cwlO*

L123cat   LU335   LU343

L1cat   LU337   LU350

L2cat   LU339   LU352

L3cat   LU341   LU354

10 µm

c) salt stress

WT168   EB35   LS089

NaCl
0 M
0.1 M
0.2 M
0.3 M
0.4 M

LU21   LU320   LU343

log OD$_{600}$

time in hours

L1cat   L2cat   L3cat

LU262   LU266   LU270 $P_{cwlOleader}$

LU350   LU352   LU354 P_{lytE}

log OD$_{600}$

time in hours

b) RT-qPCR

mreBH

fold change in expression

bacterial strain

sigI

fold change in expression

bacterial strain

Figure 5:

a) colony morphology

b) western blot

c) pull-down assay

Figure 6:

a) putative model of the evolutionary development of *lytE*, *lytF* and *cwlS*

b) proposed model for the evolutionary development of groups with different sets of paralogs

Supplementary Figure 1:

phylogeny of the *Bacillus* species

Supplementary Figure 2:

a) schematic structure of the wild type *lytE*

L1 L2 L3 cat   *lytE [L123cat]*

signal
LysM motif
linker
spacer
lytE catalytic domain

b) a variant *lytE* fragment with no LysM motif

cat   *lytE [cat]*

c) construction of variant lytE fragments with one LysM motif

L1 + cat → L1 cat   *lytE [L1cat]*

L2 + cat → L2 cat   *lytE [L2cat]*

L3 + cat → L3 cat   *lytE [L3cat]*

I   II   III

d) construction of variant *lytE* fragments with two LysM motifs

L1 L2 cat   *lytE [L11cat]*
L1 L2 cat   *lytE [L12cat]*
L1 L3 cat   *lytE [L13cat]*
L2 L1 cat   *lytE [L21cat]*
L2 L2 cat   *lytE [L22cat]*
L2 L3 cat   *lytE [L23cat]*
L3 L1 cat   *lytE [L31cat]*
L3 L2 cat   *lytE [L32cat]*
L3 L3 cat   *lytE [L33cat]*

IV

e) integration of a variant *lytE* fragment on either the *cwlO* or the *lytE* locus

Figure Legends

Figure 1

<u>Circular phylogenetic tree of the LysM motifs of the identified CwlS, LytE and LytF orthologs.</u>
The amino acid sequence of the LysM motifs of the CwlS, LytE and LytF orthologs were aligned and a phylogenetic tree estimated with Maximum Likelihood method. Bootstrap values are shown (100 replicates). Inner colours correspond to the position of the individual LysM motif within the respective protein starting at the N-terminus (red= first position, blue = second position, green = third position, purple = fourth position, orange = fifth position). Outer colours correspond to the type of the respective catalytic domain (blue=CwlS, pink=LytE, green=LytF). The three LysM motifs of the CwlS-type protein M4KSM2 are highlighted by arrows. The simplified structure of a LytF-type enzyme is shown.

Figure 2

<u>2a: Schematic illustration of strains that express LytE from the *cwlO* promoter.</u>
In both strains, the *cwlO* gene is replaced by the *lytE* open reading frame and the endogenous *lytE* gene is disrupted by a spectinomycin resistance cassette. In LU21 ($P_{cwlOleader}lytE$ Δ*cwlO* Δ*lytE*) the LytE D,L-endopeptidase is expressed from the *cwlO* promoter with *cwlO* leader sequence. In LU25 ($P_{cwlO}lytE$ Δ*cwlO* Δ*lytE*) the LytE enzyme is expressed from the *cwlO* promoter without *cwlO* leader sequence. Open reading frames for *lytE* and *cwlO* are not contiguous.

<u>2b: Typical cell morphology of strains in which LytE is expressed from the *cwlO* promoter</u>
Comparison of cell morphology of strains in which *lytE* is expressed from the *cwlO* promoter either with (LU17, LU21) or without the *cwlO* leader sequence (LU25) to the wild type (WT168). The presence or absence of the endogenous *lytE* gene is indicated. Cells were grown in LB to an OD600 of 0.1, heat-fixed and examined with an Olympus BX61 microscope. Magnification is 1000x. A scale bar is shown.

<u>2c: Transcript levels of *cwlO, lytE, sigI* and *mreBH* in strains that express LytE from the *cwlO* promoter.</u>
Transcript levels were determined by RT-qPCR for the following strains: WT168, LU21 ($P_{cwlOleader}lytE$ Δ*cwlO* Δ*lytE*) and LU25 ($P_{cwlO}lytE$ Δ*cwlO* Δ*lytE*). Cells were grown in HPDM and samples harvested at OD600=0.15, OD600=0.3 and OD600=3.00 respectively. RNA was prepared from the samples and transcribed into cDNA, which was analysed by RT-qPCR. Transcript levels were normalised relative to the level of transcript for the WT168 sample, which was assigned a value of 1. Error bars are shown.

<u>2d: Cell-associated and secreted LytE protein levels in strains in which LytE is expressed from the *cwlO* promoter.</u>
The level of LytE protein associated with the cell (left) and secreted in the medium (right) was established by western blot analysis in the following strains: LU45 (*lytE*-HA),

LU47 (*lytE*-HA Δ*cwlO*), LU61 (P*cwlOleader lytE*-HA Δ*cwlO* Δ*lytE*) and LU63 (P*cwlO lytE*-HA Δ*cwlO* Δ*lytE*). Cells were grown in HPDM and harvested at OD600=0.2 (t1) and OD600=1.0 (t2) respectively. Total cell protein was prepared from the samples and the HA-tagged LytE protein was detected by western analysis. Twenty μg of total cell protein extract was loaded onto each lane. The amount of protein loaded for medium samples was equivalent to 1 ml medium at an OD600 = 0.3. Molecular weight markers are shown (M, kDa) and the film was exposed for 1 minute. The arrowhead indicates the position of the full-length LytE protein (LytE, MW = 37 kDa).

Figure 3

3a: Typical cell morphology of strains that express LytF from the *cwlO* promoter.
Comparison of cell morphology of strains in which *lytF* is expressed from the *cwlO* promoter either with (LU111, LU116) or without the *cwlO* leader sequence (LU125, LU134). The presence or absence of the endogenous *lytE* gene is indicated. The endogenous *cwlO* and *lytF* genes are disrupted in all strains. Cells were grown in LB to an OD600 of 0.1, heat-fixed and examined with an Olympus BX61 microscope. Magnification is 1000x. A scale bar is shown.

3b: Transcript levels of *lytF*.
LytF transcript levels were determined by RT-qPCR, for the following strains: WT168, LS103 (Δ*cwlO*), LS089 (Δ*lytE*), LU74 (Δ*lytF*), LU111 (P*cwlOleader lytF* Δ*lytF* Δ*cwlO*), LU125 (P*cwlO lytF* Δ*lytF* Δ*cwlO*) and LU134 (P*cwlO lytF* Δ*lytF* Δ*cwlO* Δ*lytE*). Cells were grown in HPDM and samples harvested at OD600=0.15. RNA was prepared from the samples and transcribed into cDNA, which was analysed by RT-qPCR. Transcript levels were normalised relative to the level of transcript for the WT168 sample, which was assigned a value of 1. Error bars are shown.

3c: Cell-associated protein levels of LytF in WT168 and when expressed from the *cwlO* promoter with and without *cwlO* leader sequence.
Cell-associated protein samples of the following strains are shown: LU147 (*lytF*-HA), LU153 (Δ*lytF* P*cwlOleader lytF*-HA Δ*cwlO* Δ*lytE*) and LU157 (Δ*lytF* P*cwlO lytF*-HA Δ*cwlO* Δ*lytE*). A negative control (WT168 without HA-tag) is also displayed. Cells were grown in HPDM and harvested at OD600=0.2 (t1) and OD600=1.0 (t2). Total cell protein was prepared from the samples and the HA-tagged proteins were detected by western analysis. Ten μg total cell protein extract was deployed per lane. Molecular weight markers are shown (M, kDa). The film was exposed for one minute. The arrowhead indicates the degraded LytF protein (full length LytF, MW = 51 kDa).

3d: Localization of the LytF protein in a strain in which the protein is expressed from the *cwlO* promoter.
Strain LU151 (Δ*lytF* P*cwlOleader lytF*-HA Δ*cwlO*) was grown in buffered LB to an OD600 of 0.2 at which point cells were harvested, fixed in 2.6 % paraformaldehyde, immunostained and examined with an Olympus BX61 microscope. Immunostaining was achieved by endowing the LytF protein with a HA-tag and treating cells first with a primary anti-HA antibody followed by binding of the primary antibody with a fluorescein isothiocyanate (FITC)-

labelled secondary antibody. Proteins were visualized by phase contrast microscopy (PH) and by immunofluorescence microscopy (FITC). Magnification is 1000x. A negative control (nc) is displayed (WT168, no HA-tag). A scale bar is shown.

Figure 4

4a: Typical cell morphology of strains that express a variant LytE protein with one LysM motif from the *lytE* promoter either with or without the endogenous *cwlO* gene.
Comparison of cell morphology of strains in which a variant LytE with one LysM motifs is expressed from the *lytE* promoter in combination with an intact *cwlO* gene (LU337 (P_{lytE}*lytE[L1cat]* Δ*lytE*), LU339 (P_{lytE}*lytE[L2cat]* Δ*lytE*), LU341 (P_{lytE}*lytE[L3cat]* Δ*lytE*)), or in combination with a *cwlO* null mutant (LU350 (P_{lytE}*lytE[L1cat]* Δ*lytE* Δ*cwlO*), LU352 (P_{lytE}*lytE[L2cat]* Δ*lytE* Δ*cwlO*), LU354 (P_{lytE}*lytE[L3cat]* Δ*lytE* Δ*cwlO*)). A control with a reconstituted wild type *lytE* is shown (LU335 (P_{lytE}*lytE[L123cat]R* Δ*lytE*), LU343 (P_{lytE}*lytE[L123cat]R* Δ*lytE* Δ*cwlO*)). Cells were grown in LB to an OD600 of 0.1, heat-fixed and examined with an Olympus BX61 microscope. Magnification is 1000x. A scale bar is shown.

4b: *sigI* and *mreBH* transcript levels for strains that express a variant LytE protein from the *cwlO* promoter.
Transcript levels were established by RT-qPCR for the following strains: WT168, EB35 (Δ*cwlO*), LS089 (Δ*lytE*), LU320 ($P_{cwlOleader}$*lytE[L123cat]R* Δ*cwlO* Δ*lytE*), LU262 ($P_{cwlOleader}$*lytE[L1cat]* Δ*cwlO* Δ*lytE*), LU266 ($P_{cwlOleader}$*lytE[L2cat]* Δ*cwlO* Δ*lytE*) and LU270 ($P_{cwlOleader}$*lytE[L3cat]* Δ*cwlO* Δ*lytE*). Cells were grown in HPDM and samples harvested at OD600=0.15. RNA was prepared from the samples and transcribed into cDNA, which was analysed by RT-qPCR with specific primer sets. Transcript levels were normalised relative to the level of transcript for the WT168 sample, which was assigned a value of 1. Error bars are shown.

4c: Growth curves of strains that express a variant LytE protein with one LysM motif from the *cwlO* and the *lytE* promoter at different salt concentrations.
Growth curves were obtained for the following strains: WT168, EB35 (Δ*cwlO*), LS089 (Δ*lytE*), LU21 ($P_{cwlOleader}$*lytE* Δ*cwlO* Δ*lytE*), LU320 ($P_{cwlOleader}$*lytE[L123cat]R* Δ*cwlO* Δ*lytE*), LU343 (P_{lytE}*lytE[L123cat]R* Δ*cwlO* Δ*lytE*), LU262 ($P_{cwlOleader}$*lytE[L1cat]* Δ*cwlO* Δ*lytE*), LU266 ($P_{cwlOleader}$*lytE[L2cat]* Δ*cwlO* Δ*lytE*), LU270 ($P_{cwlOleader}$*lytE[L3cat]* Δ*cwlO* Δ*lytE*), LU350 (P_{lytE}*lytE[L1cat]* Δ*cwlO* Δ*lytE*), LU352 (P_{lytE}*lytE[L2cat]* Δ*cwlO* Δ*lytE*), LU354 (P_{lytE}*lytE[L3cat]* Δ*cwlO* Δ*lytE*). Growth curves were obtained in the following way: Overnight cultures of the strains were grown in salt-free buffered LB and used to inoculate 100 µl media in a CELLSTAR® 96 well plate to an OD600 of 0.03. NaCl was added to the final concentrations of 0.1 M, 0.2 M, 0.3 M and 0.4 M. The 96 well plate was incubated at 37°C with slow shaking and the OD600 was measured every 10 minutes over a time period of 10 hours. Three technical repeats for each strain were conducted. One representative example is shown.

Figure 5

5a: Colony morphology of strains that express a variant LytE protein from either the *cwlO* or the *lytE* promoter.

LB agar plates (no antibiotic) were inoculated with a toothpick from an overnight LB culture and incubated overnight at 37°C followed by incubation for 14 days at room temperature. Growth of each colony was documented after 1 day, 4 days, 7 days and 14 days. The following strains were characterized: LU262 ($P_{cwlOleader}$*lytE[L1cat] ΔcwlO ΔlytE*), LU266 ($P_{cwlOleader}$*lytE[L2cat] ΔcwlO ΔlytE*), LU270 ($P_{cwlOleader}$*lytE[L3cat] ΔcwlO ΔlytE*), LU205 ($P_{cwlOleader}$*lytE[L11cat] ΔcwlO ΔlytE*), LU217 ($P_{cwlOleader}$*lytE[L21cat] ΔcwlO ΔlytE*), LU229 ($P_{cwlOleader}$*lytE[L31cat] ΔcwlO ΔlytE*), LU213 ($P_{cwlOleader}$*lytE[L13cat] ΔcwlO ΔlytE*), LU277 ($P_{cwlOleader}$*lytE[L331cat] ΔcwlO ΔlytE*), LU313 ($P_{cwlOleader}$*lytE[L213cat] ΔcwlO ΔlytE*), LU320 ($P_{cwlOleader}$*lytE[L123cat]R ΔcwlO ΔlytE*), LU350 (P_{lytE}*lytE[L1cat] ΔlytE ΔcwlO*), LU352 (P_{lytE}*lytE[L2cat] ΔlytE ΔcwlO*), LU354 (P_{lytE}*lytE[L3cat] ΔlytE ΔcwlO*), LU407 (P_{lytE}*lytE[L11cat] ΔlytE ΔcwlO*), LU409 (P_{lytE}*lytE[L21cat] ΔlytE ΔcwlO*), LU383 (P_{lytE}*lytE[L31cat] ΔlytE ΔcwlO*), LU411 (P_{lytE}*lytE[L13cat] ΔlytE ΔcwlO*), LU385 (P_{lytE}*lytE[L331cat] ΔlytE ΔcwlO*), LU413 (P_{lytE}*lytE[L213cat] ΔlytE ΔcwlO*), LU343 (P_{lytE}*lytE[L123cat]R ΔlytE ΔcwlO*), WT168, LS089 (*ΔlytE*), EB35 (*ΔcwlO*).

5b: The levels of LytE variant proteins expressed from the *cwlO* promoter, for cell-associated proteins and secreted proteins.

Protein samples of the following strains are shown: LU61 (*ΔcwlO* $P_{cwlOleader}$*lytE*-HA *ΔlytE*), LU421 ($P_{cwlOleader}$*lytE*-HA *[L123cat]R ΔcwlO ΔlytE*), LU432 ($P_{cwlOleader}$*lytE*-HA *[L213cat] ΔcwlO ΔlytE*), LU418 ($P_{cwlOleader}$*lytE*-HA *[L1cat] ΔcwlO ΔlytE*), LU419 ($P_{cwlOleader}$*lytE*-HA *[L2cat] ΔcwlO ΔlytE*), LU440 ($P_{cwlOleader}$*lytE*-HA *[L3cat] ΔcwlO ΔlytE*), LU414 ($P_{cwlOleader}$*lytE*-HA *[L11cat] ΔcwlO ΔlytE*), LU415 ($P_{cwlOleader}$*lytE*-HA *[L13cat] ΔcwlO ΔlytE*) and LU417 ($P_{cwlOleader}$*lytE*-HA *[L31cat] ΔcwlO ΔlytE*). A negative control nc (WT168 without HA-tag) is also displayed. Cells were grown in salt-free buffered LB and harvested at an OD600 of 0.25. Total cell protein was prepared from the samples and the HA-tagged proteins were detected by western analysis. Five µg total cell protein extract was deployed per lane (left) and 0.15 OD equivalent media proteins were deployed per lane (right). Molecular weight markers are shown (M, kDa). Both films were exposed for seven minutes. The black arrowhead indicates the full length LytE protein (LytE, MW = 37 kDa), the grey arrowhead indicates a full length dimer protein (variant LytE with two LysM motifs, MW = 30 kDa) and the open arrowhead indicates a full length monomer protein (variant LytE with one LysM motif, MW = 23 kDa).

5c: *In vitro* binding affinity of purified variant LytE proteins to purified *B. subtilis* cell walls at different salt concentrations.

The diagram shows the results of peptidoglycan pull down assays, conducted for purified variant LytE proteins by proportion of protein found in the pellet. The following proteins were analysed: LytE[L123cat], LytE[cat], LytE[L13cat], LytE[L31cat], LytE[L1cat], LytE[L3cat]. Forty-four pmole of variant LytE protein were mixed with purified cell wall of *B. subtilis* to a final absorbance of OD600 = 10, in a 25 µl reaction volume. A negative

control (nc) with without cell wall was performed to ensure that proteins do not precipitate on their own. Pull-down assays were conducted at five different NaCl concentrations: 0.05 M, 0.1 M, 0.2 M, 0.3 M, 0.4 M and 1 M.

Figure 6

<u>6a: A putative model of the evolutionary development of the *cwlS*, *lytE* and *lytF* D,L-endopeptidases and their LysM motifs.</u>
The proposed model assumes a stable inheritance of genes and LysM motifs. The common ancestor is a gene that encodes for a D,L-endopeptidase type protein with one LysM motif. Through a combination of gene duplication (blue star), LysM motif duplication (red star) as well as LysM motif loss events (orange star), the model arrives at the set of genes observed in this study, with LytE orthologs (having either two or three LysM motifs), CwlS orthologs (four or three LysM motifs) and LytF orthologs (with four or five LysM motifs).

<u>6b: Proposed model for the evolutionary development of groups with different sets of paralogs.</u>
The model assumes a stable inheritance of genes and LysM motifs and describes how through a combination of LysM motif (red star) and gene (blue star) duplication events as well as loss of LysM motifs (orange star) different bacterial species gain different sets of paralogs.

Supplementary Figure 1

<u>Supplementary 1: Basic phylogeny of the *Bacillus* species.</u>
Phylogeny of the investigated 19 Bacillus species was inferred through alignment-free genome comparison with feature frequency profiles (FFP). Genome sequences (excluding plasmid sequences) of all species were retrieved from GenBank. FPP was performed with length l = 24. A neighbour Joining (NJ) tree was constructed based on the Jensen-Shannon divergence matrice. Bootstrapping was performed to verify the robustness of the tree and boostrapping values of 100 out of 100 were achieved for all branches. Letters (A, B, C, X) refer to the different groups of paralogs defined in Figure 2b.

Supplementary Figure 2:

<u>Supplementary 2a: schematic structure of the wild type *lytE*</u>
Schematic structure of the WT168 *lytE*, which features a signal sequence followed by three LysM motifs (L1, L2 and L3), separated by linker sequences and a catalytic domain (cat) which is separated from L3 by a spacer sequence.

<u>Supplementary 2b: schematic structure of a variant *lytE* fragment with no LysM motifs</u>
A linear fragment encoding a variant *lytE* without a LysM motif was generated by amplifying the catalytic domain with (forward) primers flanked by the signal sequence.

<u>Supplementary 2c: construction of variant *lytE* fragments with one LysM motifs</u>
Linear fragments encoding variant *lytE* with one LysM motif (III), are constructed by amplifying individual LysM motifs (I), with primers flanked by the signal and the spacer sequence, and joining them to the catalytic domain (II) by long-flanking homology PCR, thereby utilizing the spacer sequence for strand overlap.

<u>Supplementary 2d: construction of variant *lytE* fragments with two LysM motifs</u>
PCR fragments encoding variant *lytE* with two LysM motifs (IV), were constructed by amplifying individual LysM motifs with two set of primers (I and II) (one primer set is flanked by the signal and the linker sequence and the other by the linker and spacer sequence) and joining them alternating to the catalytic domain (III) by long-flanking homology PCR, thereby utilizing the linker and spacer sequence for strand overlap to generate all nine possible permutations.

<u>Supplementary 2e: integration of a variant *lytE* fragment on either the *cwlO* or the *lytE* locus</u>
Schematic representation of strains LU213 and LU411, in which a variant LytE open reading frame with 2 LysM motifs in the combination L1-L3 is integrated on the chromosome. In LU213 the variant LytE is expressed from the *cwlO* promoter with the *cwlO* leader sequence (replacing the endogenous *cwlO* gene) and in LU411 the variant *lytE* is expressed from from the *lytE* promoter (replacing the endogenous *lytE* gene). The endogenous *lytE* gene in LU213 is disrupted with a chloramphenicol resistance cassette, while the endogenous *cwlO* gene in LU411 is disrupted with kanamycin resistance cassette. Open reading frames for *lytE* and *cwlO* are not contiguous.

Suplementary Material

Table S1: List of the identified orthologs to LytE, CwlS and LytF with their UniProt identifier and NCBI taxonomic identifier

Organism	Taxonomic identifier	LytE ortholog	LytF ortholog	CwlS ortholog
Bacillus subtilis subsp. subtilis 168	224308	P54421	O07532	O31852
Bacillus subtilis subsp. inaquosorum KCTC 13429	1236548	L8PYS4	L8PUD5	L8PTT2
Bacillus subtilis subsp. natto BEST195	645657	D4G6X8	D4G6X3	D4FXM8
Bacillus subtilis subsp. spizizenii W23	655816	E0TX21	E0TX16	E0TZ07
Bacillus subtilis subsp. spizizenii TU-B-10	1052585	G4NUN9	G4NUN5	G4NQ36
Bacillus subtilis subsp. subtilis SC-8	1089443	G4EYF8	G4EYG3	G4EUZ6
Bacillus subtilis XF-1	1233100	M4KT97	M4KPW4	M4KSM2
Bacillus subtilis subsp. niger PCI 246	1495315	A0A080UQ42	A0A080UT85	A0A080UNC1
Bacillus amyloliquefaciens DSM 7	692420	E1UQ68	E1UQ63	E1UVA8
Bacillus amyloliquefaciens Y2	1126211	I2C2Z6	I2C2Z0	I2C6G3
Bacillus velezensis FZB42 / *Bacillus amyloliquefaciens subsp. Plantarum FZB42*	326423	A7Z2V6	A7Z2V1	A7Z5K3
Bacillus velezensis UCMB5033 / *Bacillus amyloliquefaciens subsp. Plantarum UCMB5033*	1338518	S6FIE6	S6FIE3	S6FL69
Bacillus atrophaeus 1942	720555	A0A0H3DZF9	A0A0H3DZF6	A0A0H3E2R9
Bacillus licheniformis DSM 13	279010	Q65LZ8	Q65IH0	none
Bacillus licheniformis CG-B52	1368424	T5HVY1	T5HPU6	none
Bacillus licheniformis S 16	1348578	W7RE25	W7RRW1	none
Bacillus licheniformis WX-02	1126218	A0A0H4X825	A0A0H4XYT8	none
Bacillus paralicheniformis ATCC 9945a	766760	R9TSA0	R9TUX7	none
Bacillus sonorensis L12	1274524	M5PGX3	M5PFH3	none

Table S2: List of strains used in this study

Strain	Genotype	Source
BL21(DE3)	*F ompT hsdSB (r_B $m_{B)}$ gal dcm (DE3)*	Studier and Moffatt, 1986
TG1	*supE hsdΔ5 thi Δ(laC-proAB) F´[traD36 proAB$^+$ lacI^q lacZΔM15]*	Sambrook et al., 1989
TP611	*recBC hsdRM cya610 pcn*	Glaser et al., 1993
EB35	*trpC2 ΔcwlO::kan^r*	unpublished, Eric Botella
DN3000	*trpC2 ΔcwlOleader*	Noone et al., 2014
LS089	*trpC2 ΔlytE::cm^r*	Salzberg et al. (2013)
LS103	*trpC2 ΔcwlO::cm^r*	Salzberg et al. (2013)
LU1	*trpC2 ΔlytE::spc^r*	LFH > WT168
LU17	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE erm^r*	LFH > WT168
LU21	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE erm^r ΔlytE::spc^r*	LFH > LU17
LU23	*trpC2 ΔcwlO::P$_{cwlo}$lytE erm^r*	LFH > WT168
LU25	*trpC2 ΔcwlO::P$_{cwlo}$lytE erm^r ΔlytE::spc^r*	LFH > LU23
LU45	*trpC2 lytE-HA kan^r*	pLU2 > WT168
LU47	*trpC2 lytE-HA kan^r ΔcwlO::cm^r*	pLU2 > LS103
LU61	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA kan^r erm^r ΔlytE::spc^r*	pLU2 > LU21
LU63	*trpC2 ΔcwlO::P$_{cwlo}$lytE-HA kan^r erm^r ΔlytE::spc^r*	pLU2 > LU25
LU72	*trpC2 ΔcwlS*	LFH dif > WT168
LU74	*trpC2 ΔlytF*	LFH dif > WT168
LU76	*trpC2 ΔyqgT*	LFH dif > WT168
LU106	*trpC2 ΔcwlO::P$_{cwlOleader}$cwlS erm^r ΔcwlS*	LFH > LU72

Strain	Genotype	Source
LU111	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytF* erm^r Δ*lytF*	LFH > LU74
LU112	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*yqgT* erm^r Δ*yqgT*	LFH > LU76
LU116	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytF* erm^r Δ*lytE*::spc^r Δ*lytF*	LFH > LU111
LU123	*trpC2* Δ*cwlO*::P_{cwlO}*cwlS* erm^r Δ*cwlS*	LFH > LU72
LU125	*trpC2* Δ*cwlO*::P_{cwlO}*lytF* erm^r Δ*lytF*	LFH > LU74
LU134	*trpC2* Δ*cwlO*::P_{cwlO}*lytF* erm^r Δ*lytE*::spc^r Δ*lytF*	LFH > LU125
LU145	*trpC2 cwlS-HA* kan^r	pLU7 > WT168
LU147	*trpC2 lytF-HA* kan^r	pLU6 > WT168
LU149	*trpC2 yqgT-HA* kan^r	pLU8 > WT168
LU151	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytF-HA* erm^r kan^r Δ*lytF*	pLU6 > LU111
LU153	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytF-HA* erm^r kan^r Δ*lytE*::spc^r Δ*lytF*	pLU6 > LU116
LU155	*trpC2* Δ*cwlO*::P_{cwlO}*lytF-HA* erm^r kan^r Δ*lytF*	pLU6 > LU125
LU157	*trpC2* Δ*cwlO*::P_{cwlO}*lytF-HA* erm^r kan^r Δ*lytE*::spc^r Δ*lytF*	pLU6 > LU134
LU159	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*cwlS-HA* erm^r kan^r Δ*cwlS*	pLU7 > LU106
LU161	*trpC2* Δ*cwlO*::P_{cwlO}*cwlS-HA* erm^r kan^r Δ*cwlS*	pLU7 > LU123
LU163	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*yqgT-HA* erm^r kan^r Δ*yqgT*	pLU8 > LU112
LU205	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L11cat] kan^r Δ*lytE*::cm^r	LFH > LU239
LU207	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L12cat] kan^r	Lin. pLU15 > WT168
LU209	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L12cat] kan^r Δ*lytE*::cm^r	LFH > LU207
LU212	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L13cat] kan^r	Lin. pLU14 > WT168
LU213	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L13cat] kan^r Δ*lytE*::cm^r	LFH > LU212
LU216	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L21cat] kan^r	Lin. pLU17 > WT168
LU217	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L21cat] kan^r Δ*lytE*::cm^r	LFH > LU216
LU220	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L22cat] kan^r	Lin. pLU18 > WT168
LU222	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L22cat] kan^r Δ*lytE*::cm^r	LFH > LU220
LU223	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L23cat] kan^r	Lin. pLU19 > WT168
LU226	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L23cat] kan^r Δ*lytE*::cm^r	LFH > LU223
LU227	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L31cat] kan^r	Lin. pLU20 > WT168
LU229	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L31cat] kan^r Δ*lytE*::cm^r	LFH > LU227
LU232	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L32cat] kan^r	Lin. pLU21 > WT168
LU233	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L32cat] kan^r Δ*lytE*::cm^r	LFH > LU232
LU235	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L33cat] kan^r	Lin. pLU22 > WT168
LU238	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L33cat] kan^r Δ*lytE*::cm^r	LFH > LU235
LU239	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L11cat] kan^r	Lin. pLU16 > WT168
LU261	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L1cat] kan^r	Lin. pLU23 > WT168
LU262	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L1cat] kan^r Δ*lytE*::cm^r	LFH > LU261
LU265	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L2cat] kan^r	Lin. pLU24 > WT168
LU266	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L2cat] kan^r Δ*lytE*::cm^r	LFH > LU265
LU269	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L3cat] kan^r	Lin. pLU25 > WT168
LU270	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L3cat] kan^r Δ*lytE*::cm^r	LFH > LU269
LU272	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L311cat] kan^r	Lin. pLU26 > 168
LU273	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L311cat] kan^r Δ*lytE*::cm^r	LFH > LU272
LU276	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L331cat] kan^r	Lin. pLU27 > WT168
LU277	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L331cat] kan^r Δ*lytE*::cm^r	LFH > LU276
LU280	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L233cat] kan^r	Lin. pLU28 > WT168
LU281	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L233cat] kan^r Δ*lytE*::cm^r	LFH > LU280
LU286	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L222cat] kan^r	Lin. pLU30 > WT168
LU287	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L222cat] kan^r Δ*lytE*::cm^r	LFH > LU286
LU288	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L212cat] kan^r	Lin. pLU31 > WT168
LU289	*trpC2* Δ*cwlO*::$P_{cwlOleader}$*lytE* [L212cat] kan^r Δ*lytE*::cm^r	LFH > LU288

Strain	Genotype	Source
LU312	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE [L213cat] kan^r*	Lin. pLU29 > WT168
LU313	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE [L213cat] kan^r ΔlytE::cm^r*	LFH > LU312
LU314	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE [cat] kan^r*	Lin. pLU37 > WT168
LU315	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE [L113cat] kan^r*	Lin. pLU38 > WT168
LU316	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE [L113cat] kan^r ΔlytE::cm^r*	LFH > LU316
LU318	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE [L123cat]^R kan^r*	Lin. pLU40 > WT168
LU320	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE [L123cat]^R kan^r ΔlytE::cm^r*	LFH > LU320
LU331	*trpC2 ΔlytE::spc^r*	LFH > WT168
LU335	*trpC2 ΔlytE::P$_{lytE}$lytE [L123cat]^R erm^r*	LFH > WT168
LU337	*trpC2 ΔlytE::P$_{lytE}$lytE [L1cat] erm^r*	LFH > WT168
LU339	*trpC2 ΔlytE::P$_{lytE}$lytE [L2cat] erm^r*	LFH > WT168
LU341	*trpC2 ΔlytE::P$_{lytE}$lytE [L3cat] erm^r*	LFH > WT168
LU343	*trpC2 ΔlytE::P$_{lytE}$lytE [L123cat]^R erm^r ΔcwlO::kan^r*	LFH > LU335
LU350	*trpC2 ΔlytE::P$_{lytE}$lytE [L1cat] erm^r ΔcwlO::kan^r*	LFH > LU337
LU352	*trpC2 ΔlytE::P$_{lytE}$lytE [L2cat] erm^r ΔcwlO::kan^r*	LFH > LU339
LU354	*trpC2 ΔlytE::P$_{lytE}$lytE [L3cat] erm^r ΔcwlO::kan^r*	LFH > LU341
LU372	*trpC2 ΔlytE::P$_{lytE}$lytE [L31cat] erm^r*	LFH > WT168
LU374	*trpC2 ΔlytE::P$_{lytE}$lytE [L331cat] erm^r*	LFH > WT168
LU383	*trpC2 ΔlytE::P$_{lytE}$lytE [L31cat] erm^r ΔcwlO::kan^r*	LFH > LU372
LU385	*trpC2 ΔlytE::P$_{lytE}$lytE [L331cat] erm^r ΔcwlO::kan^r*	LFH > LU374
LU398	*trpC2 ΔlytE::P$_{lytE}$lytE [cat] erm^r*	LFH > WT168
LU399	*trpC2 ΔlytE::P$_{lytE}$lytE [L11cat] erm^r*	LFH > WT168
LU401	*trpC2 ΔlytE::P$_{lytE}$lytE [L21cat] erm^r*	LFH > WT168
LU403	*trpC2 ΔlytE::P$_{lytE}$lytE [L13cat] erm^r*	LFH > WT168
LU405	*trpC2 ΔlytE::P$_{lytE}$lytE [L213cat] erm^r*	LFH > WT168
LU407	*trpC2 ΔlytE::P$_{lytE}$lytE [L11cat] erm^r ΔcwlO::kan^r*	LFH > LU399
LU409	*trpC2 ΔlytE::P$_{lytE}$lytE [L21cat] erm^r ΔcwlO::kan^r*	LFH > LU401
LU411	*trpC2 ΔlytE::P$_{lytE}$lytE [L13cat] erm^r ΔcwlO::kan^r*	LFH > LU403
LU413	*trpC2 ΔlytE::P$_{lytE}$lytE [L213cat] erm^r ΔcwlO::kan^r*	LFH > LU405
LU414	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L11cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU205
LU415	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L13cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU213
LU416	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L21cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU217
LU417	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L31cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU229
LU418	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L1cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU262
LU419	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L2cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU266
LU421	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L123cat]^R kan^r spc^r ΔlytE::cm^r*	pLU46 > LU320
LU422	*trpC2 ΔlytE::P$_{lytE}$lytE-HA [L123cat]^R erm^r spc^r ΔcwlO::kan^r*	pLU46 > LU343
LU424	*trpC2 ΔlytE::P$_{lytE}$lytE-HA [L331cat] erm^r spc^r ΔcwlO::kan^r*	pLU46 > LU385
LU425	*trpC2 ΔlytE::P$_{lytE}$lytE-HA [L11cat] erm^r spc^r ΔcwlO::kan^r*	pLU46 > LU407
LU427	*trpC2 ΔlytE::P$_{lytE}$lytE-HA [L13cat] erm^r spc^r ΔcwlO::kan^r*	pLU46 > LU411
LU432	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L213cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU313
LU435	*trpC2 ΔlytE::P$_{lytE}$lytE-HA [L1cat] erm^r spc^r ΔcwlO::kan^r*	pLU46 > LU350
LU440	*trpC2 ΔcwlO::P$_{cwlOleader}$lytE-HA [L3cat] kan^r spc^r ΔlytE::cm^r*	pLU46 > LU270
LU442	*trpC2 ΔlytE::P$_{lytE}$lytE-HA [L2cat] erm^r spc^r ΔcwlO::kan^r*	pLU46 > LU352
LU443	*trpC2 ΔlytE::P$_{lytE}$lytE-HA [L213cat] erm^r spc^r ΔcwlO::kan^r*	pLU46 > LU413
WT168	*trpC2*	Laboratory stock, Kunst et al., 1997

[LFH] long flanking homology fragment; [cm^r] chloramphenicol resistance gene; [emr^r] erythromycin resistance gene; [kan^r] kanamycin resistance gene; [spc^r] spectinomycin resistance gene

Table S3: List of plasmids used in this study

Plasmid	Genotype	Source
pDG780	plasmid with *kan*[r] cassette	Guerout-Fleury et al., 1995
pDG646	plasmid with *erm*[r] cassette	Guerout-Fleury et al., 1995
pDG1727	plasmid with *spc*[r] cassette	Guerout-Fleury et al., 1995
pDG1661	plasmid with *cm*[r] cassette	Guerout-Fleury et al., 1995
pETLytE	pET21b+ derivative, protein expression vector for expression of the wild-type LytE protein	unpublished, David Noone
pLU2	pDG780 derivative enabling C-terminal HA-tagging of *lytE* over single cross-over, *kan*[r] cassette	This work
pLU6	pDG780 derivative enabling C-terminal HA-tagging of *lytF* over single cross-over, *kan*[r] cassette	This work
pLU9	pDG780 derivative with upstream homology to the *cwlO* locus (including *cwlO* promoter and *cwlO* leader sequence)	This work
pLU10	pLU9 derivative with up- and downstream homology to the *cwlO* locus (including *cwlO* promoter and *cwlO* leader sequence)	This work
pLU14	pLU10 derivative enabling integration of variant *lytE [L13cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU15	pLU10 derivative enabling integration of variant *lytE [L12cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU16	pLU10 derivative enabling integration of variant *lytE [L11cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU17	pLU10 derivative enabling integration of variant *lytE [L21cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU18	pLU10 derivative enabling integration of variant *lytE [L22cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU19	pLU10 derivative enabling integration of variant *lytE [L23cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU20	pLU10 derivative enabling integration of variant *lytE [L31cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU21	pLU10 derivative enabling integration of variant *lytE [L32cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU22	pLU10 derivative enabling integration of variant *lytE [L33cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU23	pLU10 derivative enabling integration of variant *lytE [L1cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU24	pLU10 derivative enabling integration of variant *lytE [L2cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU25	pLU10 derivative enabling integration of variant *lytE [L3cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU26	pLU20 derivative enabling integration of variant *lytE [L311cat]* at the cwlO locus by double cross-over, *kan*[r] cassette	This work
pLU27	pLU20 derivative enabling integration of variant *lytE [L331cat]* at the cwlO locus by double cross-over, *kan*[r] cassette	This work
pLU28	pLU19 derivative enabling integration of variant *lytE [L233cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU29	pLU19 derivative enabling integration of variant *lytE [L213cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU30	PLU18 derivative enabling integration of variant *lytE [L222cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU31	pLU18 derivative enabling integration of variant *lytE [L212cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU37	pLU10 derivative enabling integration of variant *lytE [cat]* at the *cwlO* locus by double cross-over, *kan*[r] cassette	This work
pLU38	pLU14 derivative enabling integration of variant *lytE [L113cat]* at the cwlO locus by double cross-over, k*an*[r] cassette	This work
pLU40	pLU14 derivative enabling integration of variant *lytE [L123cat]*[R] at the cwlO locus by double cross-over, *kan*[r] cassette	This work
pLU46	pLU2 derivative enabling C-terminal HA-tagging of *lytE* over single cross-over, *spc*[r] cassette	This work
pLU47	pET21b+ derivative, protein expression vector for expression of the variant LytE [L123cat][R]	This work
pLU48	pET21b+ derivative, protein expression vector for expression of the variant LytE [L31cat]	This work
pLU49	pET21b+ derivative, protein expression vector for expression of the variant LytE [L13cat]	This work
pLU50	pET21b+ derivative, protein expression vector for expression of the variant LytE [L1cat]	This work
pLU51	pET21b+ derivative, protein expression vector for expression of the variant LytE [L3cat]	This work
pLU52	pET21b+ derivative, protein expression vector for expression of the variant LytE [cat]	This work

[cm[r]] chloramphenicol resistance gene; [erm[r]] erythromycin resistance gene; [kan[r]] kanamycin resistance gene; [spc[r]] spectinomycin resistance gene

Table S4: List of oligonucleotids used in this study

Primer name	Sequence
16S_Bsu_F	GTCAGCTCGTGTCGTGAGATG
16S_Bsu_R	CAACTGAATGCTGGCAACTAAGATC
Aaa_LysM_Fwd	cgcggatccacgttatttgcaggagcggcatctgcacaaGTAAAACCGGGTGAATCAG
Aaa_LysM_Rev	atgtagatgacgttttgctgcttgaagacgatgaagatgaagcaggttttgagctgcttgtgcttgttcc TACTTTTAGTACTTGGTTTG
Catalytic_LytE_Fwd_Spac	ttcaagcagcaaaacgtcatctacatcacttaatgtgagcaagctggtttctgatgcaaaagcgttagtc GGAACGCCATATAAATGGGG
Catalytic_LytE_Rev_Ery	attatgtctttgcgcagtcggcGAAAAAAACCCGTTCCAATGAACGG
Catalytic_LytE_Rev_Xmal	tcccccgggGAAAAAAACCCGTTCCAATGAACGG
CwlO_DF_LysM	ccgctcgagATTAACATATACCCTTTTCGAATA
CwlO_DR_LysM	cggggtaccCCTTCTTCATTCGTAATGCCCAGGT
CwlO_UF	CCTTCATCTGTGGAGCAG
CwlO_UF_LysM	aaggaaaaaagcggccgcATCATTATATAAAACCGGCTT
CwlO_UF_Smal	gcatcccgggCCTTCATCTGTGGAGCAG
CwlO_UR	cgtagctgtaatgatttgcttttcatTTTTTATATCCTCCCTTTTACTGTGAACC
CwlO_UR_CwlS	aagccggctacaatcttcttttcatTTTTTATATCCTCCCTTTTACTGTGAA
CwlO_UR_LysM	taacgtggatcctaaaacaactgctgtcgtagctgtaatgatttgcttttcat TTTTTATATCCTCCCTTTTAC
CwlO_UR_LytF	gccctgctgctaatttcttttcatTTTTTATATCCTCCCTTTTACTGTGAA
CwlO_UR_Yqgt	gtttcaccgtaataaacgccatcatTTTTTATATCCTCCCTTTTACTGTGAA
CwlS_Do_Fwd	tatattatgtaaactaatatgcaagAAAATGAGTATGACAAAAAGCC
CwlS_Do_Rev	TCTGGGGAAATAGGCTGCTCT
CwlS_Fwd	ATGAAAAAGAAGATTGTAGCCGGC
CwlS_LysM_Fwd	cgcggatccacgttatttgcaggagcggcatctgcacaaATAAAAGTAAAAAGCGGTGA
CwlS_LysM_Rev	atgtagatgacgttttgctgcttgaagacgatgaagatgaagcaggttttgagctgcttgtgcttgttcc AGGAACCTTAAGACTTTGTC
Cwls_Rev_Ery	tcgattatgtctttgcgcagtcggcTAAACAAAAAAGCCTATAAAAGGC
CwlS_Tag_F_BamHl	cgggatccGTCGGAGTGCCTTACCGATGGGGC
CwlS_Tag_R_HA_EcoRl	ggaattctaaacaaaaaagcctataaaaggctttttgtcatactcattttattatgcataatccg gaacatcatacggataAAAATAACTTCTTGCGCCCAAGTA
CwlS_Up_Fwd	CGGCAAGCATATTAAAACAAGTGAG
CwlS_Up_Rev	tatattctaggaagtaggtattcacTTCATTCAACCTCCTAAACTG
Exp_Cat_Rev	ccgctcgagGAATCTTTTCGCACCGAGGTAACG
Exp_L1_Fwd	ggaattccatatgAGCATTAAGGTGAAAAAAGGCGACA
Exp_L2_Fwd	ggaattccatatgACATACAAAGTAAAGAGCGGGGACA
Exp_L3_Fwd	ggaattccatatgACTTATAAAGTGAAGAGCGGAGACA
Exp_LytF_Fwd	ggaattccatatgATTAAGGTCAAAAGCGGAGATTCTTT
Exp_Spac_Fwd	ggaattccatatgGGAACAAGCACAAGCAGCTCAAAAC
Exp_XlyB_Fwd	ggaattccatatgTATCATGTAAAAAAAGGTGACACACTG
L1_Fwd_Link	ggcaaatctactagttcaaaaagcagcagttcttcttcctcttcttct AGCATTAAGGTGAAAAAAGGC
L1_Fwd_Sig_BamHl	cgcggatccacgttatttgcaggagcggcatctgcacaa AGCATTAAGGTGAAAAAAGGC
L1_Rev_Link	agaagaagaggaagaagaactgctgctttttgaactagtagatttgcc GTTAATCGATAAAGTTTGTCCC
L1_Rev_Spac	atgtagatgacgttttgctgcttgaagacgatgaagatgaagcaggttttgagctgcttgtgcttgttcc GTTAATCGATAAAGTTTGTC
L2_Fwd_Link	ggcaaatctactagttcaaaaagcagcagttcttcttcctcttcttct ACATACAAAGTAAAGAGCGGG
L2_Fwd_Sig_BamHl	cgcggatccacgttatttgcaggagcggcatctgcacaa ACATACAAAGTAAAGAGCGGG
L2_Rev_Link	agaagaagaggaagaagaactgctgctttttgaactagtagatttgcc TTTCAGTTTCAGGACTTGTCC
L2_Rev_Spac	atgtagatgacgttttgctgcttgaagacgatgaagatgaagcaggttttgagctgcttgtgcttgttcc TTTCAGTTTCAGGACTTGTC
L3_Fwd_Link	ggcaaatctactagttcaaaaagcagcagttcttcttcctcttcttct ACTTATAAAGTGAAGAGCGGA
L3_Fwd_Sig_BamHl	cgcggatccacgttatttgcaggagcggcatctgcacaa ACTTATAAAGTGAAGAGCGGA
L3_Rev_Link	agaagaagaggaagaagaactgctgctttttgaactagtagatttgcc TTTCACCTTCAATACTTGGTTG

Primer name	Sequence
L3_Rev_Spac	atgtagatgacgttttgctgcttgaagacgatgaagatgaagcaggttttgagctgcttgtgcttgttcc TTTCACCTTCAATACTTGGT
LytE_DF	ATGAAAAAGCAAATCATTACAGCTACG
LytE_Do_Fwd_Ery	ggttgccagagttaaaggatcATTAGACGGAGCACAGGCGTTTT
LytE_Do_Fwd_Spec	caataaacccttgccctcgctacgTACTGGAAGCCTCGTTACCTCGG
LytE_Do_Fwd_Spec	caataaacccttgccctcgctacgATTAGACGGAGCACAGGCGTTTT
LytE_Do_Rev	GTATTTGCCGTATGCGCTCAGGC
LytE_DR_Ery	attatgtcttttgcgcagtcggcGAAAAAAACCCGTTCCAATGAA
LytE_Probe_F	TATCGATTAACGGCAAATCTACAAGTT
LytE_Probe_T7_R	taatacgactcactatagggGCAGTGCTTGTTCTGCCCACACTT
LytE_qPCR_F2_	GGAGGAAAATATGAAAAAGCAAATCA
LytE_qPCR_R2	TCGCTCTCATACGTATCAACACTCT
LytE_Spacer_Fwd_Sig	cgcggatccacgttatttgcaggagcggcatctgcacaaGGAACAAGCACAAGCAGC
LytE_Tag_F_BamHI	cgggatccTTAGTCGGAACGCCATATAAATGG
LytE_Tag_R_HA_EcoRI	ggaattcgaaaaaaacccgttccaatgaacgggttttctctaaaaattattatgcata atccggaacatcatacggataGAATCTTTTCGCACCGAGGTAACG
LytE_Up_Fwd	AACGGTAGGCTCTTATGTAAGGC
LytE_Up_Fwd_2	AACGGTAGGCTCTTATGTAAGGC
LytE_Up_Rev_Spec	cgttacgttattagcgagccagtcCCTAAAACAACTGCTGTCGTAGC
LytF_Do_Fwd	tatattatgtaaactaatatgcaagTATTAGAGCCTAAAGACGTTGATGC
LytF_Do_Rev	CCGGGATTTGTTTATGTGAGAA
LytF_Fwd	ATGAAAAAGAAATTAGCAGCAGGG
LytF_LysM_Fwd	cgcggatccacgttatttgcaggagcggcatctgcacaaATTAAGGTCAAAAGCGGAG
LytF_LysM_Rev	atgtagatgacgttttgctgcttgaagacgatgaagatgaagcaggttttgagctgcttgtgcttgttcc TGGAATAGTAAGCGTCTGG
LytF_qPCR_Fwd	TGTCGGAGATTTCGTTTTCTTTTC
LytF_qPCR_fwd	TGTCGGAGATTTCGTTTTCTTTTC
LytF_qPCR_Rev	CCCGTTGCCTAGGTAAATTCC
LytF_qPCR_rev	CCCGTTGCCTAGGTAAATTCC
LytF_Rev_Ery	tcgattatgtcttttgcgcagtcggcGCATCAACGTCTTTAGGCTCTAATA
LytF_Tag_F_BamHl	cgggatccCTTGGGGGTTCCGTATCGCTGGGGA
LytF_Tag_R_HA_EcoRl	ggaattcgcatcaacgtctttaggctctaataaaaaactgtacagccgttaaggccgcacagtttctg tttttattatgcataatccggaacatcatacggataGAAATATCGTTTTGCACCGAGATA
LytF_Up_Fwd	CTTTTGGTTTATCCTTCGGCCTT
LytF_Up_Rev	tatattctaggaagtaggtattcacCTTAGCTCTCCTTTTTCCCCC
MreBH_qPCR_F	GCAAAACCCGCTTGAATCTG
MreBH_qPCR_R	TGCCTTTTGCAGCTTATCAATC
OE93_Fwd_Ery	ggttgccagagttaaaggatcGAGCAATCCTTACTGGAAAGCAGC
OE94_Rev	ATCGACAAGCGTTACGTTACCGAC
SigI_Chip_Fwd	AAATTCTCTGCTCGCATTTGC
SigI_Chip_Rev	GGAAAGCTCGGCTTCAATCAG
Signal_LytE_Fwd	ATGAAAAAGCAAATCATTACAGCTACG
Signal_LytE_Rev	TTGTGCAGATGCCGCTCCTGCAA
Signal_LytE_Rev_Spec	cgttacgttattagcgagccagtcTTGTGCAGATGCCGCTCCTGCAA
Spacer_Check_Rev	GATGTAGATGACGTTTTGC
Spec_Fwd_Ecorl	ccggaattccggCTAATAACGTAACGTGACTGGCA
Spec_Rev_Sall	ttccgcggccgctatggccgacgtcgacgcgtAGGGTTTATTGTTTTCTAAAATCTG
YocH_Northern_F	GCAGCAAACACAGCTACTGAGAAT
YocH_Northern_R	GCTCTTTTTGCACAGCCTCTT
YqgT_Do_Fwd	tatattatgtaaactaatatgcaagACTGAAACTTTTTCTTTTTTCTGC
YqgT_Do_Rev	TTGGAGCGGCCATTTCGCTGT
YqgT_Fwd	ATGATGGCGTTTATTACGGTG

Primer name	Sequence
YqgT_Rev_Ery	tcgattatgtcttttgcgcagtcggcTCTCATATTTGCTTCGGCAGAA
YqgT_Tag_F_BamHl	cgggatccACGGAGAATCCGCCTGACAGACTG
YqgT_Tag_R_HA_EcoRl	ggaattctctcatatttgcttcggcagaaaaagaaaaagtttcagttattatgcataatccgg aacatcatacggataCTCATGAAAAAATAACGCCCTCCA
YqgT_Up_Fwd	TTACTCGAGGCTGATTACACTG
YqgT_Up_Rev	tatattctaggaagtaggtattcacTCATCCCCCTTTTTTGTTACTATA
YvcE_3qPCR_Orf*2	GGCGGATCTCAAGGATACATAGATG
YvcE_5qPCR_Orf*2	GTCTGCGTCCACAATAGATGAAAC

*   lower case = 5´ overhang, upper case = annealing sequence