1   **Optimized Multicolour Immunofluorescence Panel for Cattle T Cell**

2   **Phenotyping by an 8-Colour, 10-Parameter Panel**

3   **Eduard O. Roos[1], William Mwangi[1], Wilhelm Gerner[1], Ryan Waters[1], John A.**
4   **Hammond[1*]**
5
6   [1] The Pirbright Institute, Ash Road, Pirbright, Woking, Surrey, GU24 0NF, UK

7   [*]Correspondence to: John Hammond; The Pirbright Institute, Ash Road, Pirbright,
8   Woking, Surrey, GU24 0N, UK. E-mail: john.hammond@pirbright.ac.uk
9

Purpose and appropriate sample types:

This multiplex staining panel was developed to differentiate cattle T cells into

conventional (CD4 and CD8) and unconventional ($\gamma\delta$-TCR) subsets as well as their

stage of differentiation and activation. The combination of CD45RO and CD62L

allows the identification of naïve ($T_{Naïve}$), central memory (T_{CM}), effector memory

(T_{EM}) and terminal effector (T_{TE}) T cells. Activated cattle T cells (T_{AV}) can be

identified by the cell surface expression of CD25. This panel was developed using

cryopreserved cattle peripheral blood mononuclear cells (PBMCs) and tested on

fresh as well as stimulated PBMCs. Therefore, this 8-colour, 10-parameter flow

cytometry panel simultaneously identifies cattle $T_{Naïve}$, T_{AV}, T_{CM}, T_{EM}, T_{TE} and $\gamma\delta$-

TCR cells. This panel will improve our ability to examine T cell response to

pathogens and vaccines in cattle including the potential to identify previously

undescribed subpopulations. Furthermore, this panel can be readily optimised for

other bovid species as many of these reagents are likely to cross react.

Key terms:

Flow cytometry; cattle PBMC; T cells; naïve T cells; effector memory T cells; central

memory T cells; activated T cells; γδ T cells; T cell subsets

Background:

Robust T cell responses are critical in the response to pathogen infection both for

clearance and the formation of strong and broad memory responses (1). Cattle, like

several other species, have a much higher proportion of γδ T cells compared to CD4

and CD8 (2–4). Consequently, it is important to study the entire T cell compartment

simultaneously to fully characterise how immune protection arises and persists.

Furthermore, as the research climate focusses on One Health approaches, the ability

to study the immune response at high resolution in species that underpin global

food security is essential.

Common to several non-model species, the first mAbs to study CD molecules on

cattle T cells were derived from mouse immunizations with whole cattle PBMC

populations or PBMC lysates. Antibodies were characterized in three international

workshops on ruminant antigens (2,5,6). Together with identification of cross-

reactive mAbs, this allowed the establishment of a basic toolbox to study cattle T

cells and various subsets within them (7). However, several limitations still exist for

the establishment of polychromatic flow cytometry staining panels. For example,

many of the current 371 human CD molecules do not have an antibody that cross

react with cattle. Another major limitation is the lack of useful mAbs that are

labelled to a wider range of fluorochromes. This makes it difficult to expand panels

beyond the three most common fluorochromes FITC (or AF488), PE and APC (or

AF647). By conjugating existing T cell markers in-house we were able to develop a

cattle T cell panel that utilises eight colours excluding PE and APC-conjugated

antibodies. This allows the addition of specific antibodies, such as for cytokines or

transcription factors, that maximises the broader utility of this panel for individual

research needs. Additionally, if more of the available mAbs would be conjugated to

fluorochromes that are excited by the violet laser, the panel can be further expanded.

This OMIP identifies all three main cattle T cell subsets (CD4, CD8 and $\gamma\delta$), as well

as their subsets that are activated (T_{AV}) or in the distinct differentiation states of

naïve ($T_{Naïve}$), central memory (T_{CM}), effector memory (T_{EM}) and terminal effector

(T_{TE}). The gating strategy we used initially identifies the two $\alpha\beta$ T cell subsets CD4

(mAb clones CC8/CC30) and CD8 (mAb clone CC63) as well as the $\gamma\delta$ T cells (mAb

clone GB21A) (2,8) (Fig. 1). Like in swine and chickens, $\gamma\delta$ T cells constitute a major

T cell subset in cattle blood and can comprise more than 50% of circulating T cells

(3,9). To identify activated T cells, CD25 (mAb clone IL-A111) can be used (8,10–12),

whereas the memory state of the cells can be defined using the CD45RO (mAb clone

IL-A116) and CD62L (mAb clone CC32) cell surface markers (6,8,13–15) (Fig. 1).

Using this gating strategy, the following known subsets can be identified for the

helper T cells, $T_{Naïve}$ (CD3$^+\gamma\delta$-TCR$^-$CD4$^+$CD25$^-$CD45RO$^-$CD62L$^+$), T_{CM} (CD3$^+\gamma\delta$-TCR$^-$

CD4$^+$CD25$^-$CD45RO$^+$CD62L$^+$), T_{EM} (CD3$^+\gamma\delta$-TCR$^-$CD4$^+$CD25$^-$CD45RO$^+$CD62L$^-$), T_{TE}

(CD3$^+\gamma\delta$-TCR$^-$CD4$^+$CD25$^-$CD45RO$^-$CD62L$^-$) and T_{AV} (CD3$^+\gamma\delta$-TCR$^-$CD4$^+$CD25$^+$).

Similarly, the cytotoxic T cells can be separated into $T_{Naïve}$ (CD3$^+\gamma\delta$-TCR$^-$

$CD8\alpha^+CD25^-CD45RO^-CD62L^+$), T_{CM} ($CD3^+\gamma\delta\text{-}TCR^-CD8\alpha^+CD25^-CD45RO^+CD62L^+$),

T_{EM} ($CD3^+\gamma\delta\text{-}TCR^-CD8\alpha^+CD25^-CD45RO^+CD62L^-$), T_{TE} ($CD3^+\gamma\delta\text{-}TCR^-CD8\alpha^+CD25^-$

$CD45RO^-CD62L^-$) and T_{AV} ($CD3^+\gamma\delta\text{-}TCR^-CD8\alpha^+CD25^+$). Furthermore, $\gamma\delta$ T cells can

also be identified by ($CD3^+\gamma\delta\text{-}TCR^+$) (Fig. 1; Online Table 3). A major improvement

by this panel is the simultaneous analysis of all these T cell subsets in a single

sample, hence reducing the variation between replicates and the number of samples

needed per animal.

The panel was designed and optimised on a BD LSRFortessa and was tested on a BD

Aria IIIU. The BD Aria IIIU allows for sorting of the T cell subsets. Further

adaptations to the panel are enabled by having both the PE and APC channel empty,

for which many antibodies are commercially available. If more reagents become

available in the violet channel, they can easily be added to the panel with only minor

influence on compensation requirements.

In conclusion, this cattle T cell panel will advance the understanding of the cattle

immune response as it allows the measurement of all major T cell subsets and their

differentiation stage within a single sample.

Similarity to published OMIPs:

None to date.

Acknowledgements:

The authors wish to acknowledge the valuable input of Dr Katy Moffat for technical

support on the instruments. This project received funding from the European

Union's Horizon 2020 research and innovation programme under the VetBioNet

grant agreement (731014). The authors would like to acknowledge the Pirbright

Flow Cytometry facility and the Immunological Toolbox unit supported by the

United Kingdom Research and Innovation- Biotechnology and Biological Sciences

Research Council awards (BBS/E/I/00007038 and BBS/E/I/00007039); JAH and

WG were further supported by BBS/E/I/00007030 and BBS/E/I/00007031.

104   Literature Cited

1. Laidlaw BJ, Craft JE, Kaech SM. The multifaceted role of CD4+ T cells in CD8+ T

cell memory. Nat. Rev. Immunol. 2016;16:102–111.

2. Davis WC, Brown WC, Hamilton MJ, Wyatt CR, Orden JA, Khalid AM, Naessens

J. Analysis of monoclonal antibodies specific for the $\gamma\delta$ TcR. Vet. Immunol.

Immunopathol. 1996;52:275–283.

3. Baldwin CL, Damani-Yokota P, Yirsaw A, Loonie K, Teixeira AF, Gillespie A.

Special features of $\gamma\delta$ T cells in ruminants. Mol. Immunol. 2021;134:161–169.

Available at: https://doi.org/10.1016/j.molimm.2021.02.028.

4. Telfer JC, Baldwin CL. Bovine gamma delta T cells and the function of gamma

delta T cell specific WC1 co-receptors. Cell. Immunol. 2015;296:76–86. Available at:

http://dx.doi.org/10.1016/j.cellimm.2015.05.003.

5. Howard CJ, Morrison WI, Bensaid A, Davis W, Eskra L, Gerdes J, Hadam M,

Hurley D, Leibold W, Letesson J-J, MacHugh N, Naessens J, O'Reilly K, Parsons KR,

Schlote D, Sopp P, Splitter G, Wilson R. Summary of workshop findings for

leukocyte antigens of cattle. Vet. Immunol. Immunopathol. 1991;27:21–27. Available

at: https://linkinghub.elsevier.com/retrieve/pii/016524279190072K.

6. Naessens J, Hopkins J. Introduction and summary of workshop findings. Vet.

Immunol. Immunopathol. 1996;52:213–235. Available at:

https://linkinghub.elsevier.com/retrieve/pii/0165242796055663.

7. Sopp P, Werling D, Baldwin C. Cross-reactivity of mAbs to human CD antigens

with cells from cattle. Vet. Immunol. Immunopathol. 2007;119:106–114. Available at:

https://linkinghub.elsevier.com/retrieve/pii/S0165242707002206.

8. Howard CJ, Naessens J. 4.1 Summary of workshop findings for cattle (Tables 1

and 2). Vet. Immunol. Immunopathol. 1993;39:25–47. Available at:

https://linkinghub.elsevier.com/retrieve/pii/016524279390161V.

9. Baldwin CL, Telfer JC. The bovine model for elucidating the role of $\gamma\delta$ T cells in

controlling infectious diseases of importance to cattle and humans. Mol. Immunol.

2015;66:35–47. Available at: http://dx.doi.org/10.1016/j.molimm.2014.10.024.

10. Coussens PM, Sipkovsky S, Murphy B, Roussey J, Colvin CJ. Regulatory T cells in

cattle and their potential role in bovine paratuberculosis. Comp. Immunol.

Microbiol. Infect. Dis. 2012;35:233–239. Available at:

http://dx.doi.org/10.1016/j.cimid.2012.01.004.

11. Caruso A, Licenziati S, Corulli M, Canaris AD, De Francesco MA, Fiorentini S,

Peroni L, Fallacara F, Dima F, Balsari A, Turano A. Flow cytometric analysis of

activation markers on stimulated T cells and their correlation with cell proliferation.

Cytometry 1997;27:71–6. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/9000587.

12. Waters WR, Rahner TE, Palmer M V., Cheng D, Nonnecke BJ, Whipple DL.

Expression of L-selectin (CD62L), CD44, and CD25 on activated bovine T cells.

Infect. Immun. 2003;71:317–326.

13. Golubovskaya V, Wu L. Different subsets of T cells, memory, effector functions,

and CAR-T immunotherapy. Cancers (Basel). 2016;8.

14. Bembridge GP, MacHugh ND, McKeever D, Awino E, Sopp P, Collins RA,

Gelder KI, Howard CJ. CD45RO expression on bovine T cells: relation to biological

function. Immunology 1995;86:537–44. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/8567018%0Ahttp://www.pubmedcentral.

nih.gov/articlerender.fcgi?artid=PMC1384052.

15. Whelan AO, Villarreal-Ramos B, Vordermeier HM, Hogarth PJ. Development of

an antibody to bovine IL-2 reveals multifunctional CD4 T EM cells in cattle naturally

infected with bovine tuberculosis. PLoS One 2011;6:2–9.

Figure 1 Gating strategy of the cattle T-cell panel into CD4$^+$, CD8$^+$ and γδ-TCR$^+$ subsets. A) Sub-gating of the CD3$^+$γδ-TCR$^+$ subset with CD45RO and CD62L. B) Sub-gating of the CD3$^+$ γδ-TCR$^-$CD8α$^+$subset with CD25. C) Sub-gating of CD3$^+$ γδ-TCR$^-$CD8α$^+$CD25$^-$ subset with CD45RO and CD62L. D) Sub-gating of CD3$^+$ γδ-TCR$^-$CD4$^+$ with CD25. E) Sub-gating of CD3$^+$ γδ-TCR$^-$CD4$^+$CD25$^-$ subset with CD45RO and CD62L. Sub-gating of the CD4 and CD8α with CD45RO and CD62L allows the identification of effector memory (T$_{EM}$), central memory (T$_{CM}$), terminal effector (T$_{TE}$) and naïve (T$_{Naïve}$) T cells. The CD25 identifies the activated T (T$_{AV}$) cells, that include the T regulatory cells.