**Age-Induced Changes in Mu Opioid Receptor Signaling in the Midbrain Periaqueductal Gray of Male and Female Rats**

Evan F. Fullerton[1], Mary C. Karom[1], John M. Streicher[2], Larry J. Young[3], and Anne Z. Murphy[1]

[1]Neuroscience Institute, Georgia State University, Atlanta, GA, 30303, USA

[2]Department of Pharmacology, College of Medicine, University of Arizona, Tucson, AZ 85724, USA.

[3]Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA, 30322 USA

**Corresponding author:**

Anne Z. Murphy, Ph.D.

Neuroscience Institute, Georgia State University

100 Piedmont Ave SE, Atlanta, Georgia, 30303, United States

Tel: 404/413-5332

Email: amurphy@gsu.edu

**Highlights:**

- Aged males and females (adult and aged) exhibit reduced vlPAG μ-opioid receptor binding potential compared to adult males.

- Aged males and females (adult and aged) exhibit reduced opioid-induced vlPAG G-protein activation compared to adult males.

- Aged males and females (adult and aged) exhibit reduced vlPAG MOR mediated cAMP inhibition compared to adult males.

- Aged rats (males and females) exhibit increased vlPAG mRNA expression of Regulator of G-Protein Signaling (RGS) proteins RGS4 and RGS9-2 compared to adult rats (males and females), which may explain the reduced receptor signaling observed in aged animals.

- These coordinate decreases in opioid receptor signaling may explain the previously reported reduced potency of opioids to produce pain relief in females and aged rats.

**Abstract: The analgesic effects of opioids are attenuated in aged rats.** Opioids such as morphine have

decreased analgesic potency (but not efficacy) in aged rodents compared to adults; however, the neural

mechanisms underlying this attenuated response are not yet known. The present study investigated the

impact of advanced age and biological sex on opioid signaling in the ventrolateral periaqueductal gray

(vlPAG) in the presence of chronic inflammatory pain. Assays measuring mu-opioid receptor (MOR)

radioligand binding, GTP$\gamma$S binding, receptor phosphorylation, cAMP inhibition, and regulator of G-

protein signaling (RGS) protein expression were performed on vlPAG tissue from adult (2-3mos) and

aged (16-18mos) male and female rats. Persistent inflammatory pain was induced by intraplantar

injection of Complete Freund's Adjuvant (CFA). Adult males exhibited the highest MOR binding potential

and the highest G-protein activation (activation efficiency ratio) in comparison to aged males and

females (adult and aged). No impact of advanced age or sex on MOR phosphorylation state was

observed. DAMGO-induced cAMP inhibition was highest in the vlPAG of adult males compared to aged

males and females (adult and aged). vlPAG levels of RGS4 and RGS9-2, critical for terminating G-protein

signaling, were assessed using RNAscope. Adult rats (both males and females) exhibited lower levels of

vlPAG RGS4 and RGS9-2 mRNA expression compared to aged males and females. The observed age-

related reductions in vlPAG MOR binding potential, G-protein activation efficiency, and cAMP inhibition,

along with the observed age-related increases in RGS4 and RGS9-2 vlPAG expression, provide potential

mechanisms whereby the potency of opioids is decreased in the aged population. These results have

significant implications for pain management in this population.

**Keywords:** chronic pain; opioid signaling; pharmacology; advanced age; sex differences; mu-opioid

receptor

## 1 Introduction

Chronic pain is a debilitating condition that contributes to mental health disorders, disability, and increased risk of premature death (Domenichiello and Ramsden, 2019; Patel et al., 2013). More than 50% of individuals aged 65 and over suffer from chronic pain, and as the United States population rapidly ages, the development of effective strategies for chronic pain management for this demographic is imperative (Dahlhamer, 2018). Opioids, including morphine and fentanyl, are commonly prescribed analgesics for the management of chronic pain (Manchikanti et al., 2010; Sullivan et al., 2010), but effective dosing strategies for the elderly remain ill-defined, due in large part to a dearth of knowledge regarding the impact of advanced age on opioid signaling and analgesia.

Clinical studies examining pain management in the elderly are challenging; co-morbid conditions such as diabetes and high blood pressure, and participant use of concomitant medications affect patient outcomes, contributing to difficulties in the interpretation of results (Naples et al., 2016; Prostran et al., 2016). Additionally, there exist age- and sex-related individual differences in the likelihood of reporting pain in a clinical setting which may lead to misrepresentations of analgesic efficacy (Dampier et al., 2013; Reddy et al., 2012). Particularly, aged populations are known to underreport pain due to fears of institutionalization and concerns with addiction and overdose (Reddy et al., 2012).

Using a preclinical model of persistent inflammatory pain (intraplantar Complete Freund's Adjuvant; (CFA)), we have previously reported a significant impact of age and sex on morphine potency. Specifically, we showed that aged rats (18mos) exhibit a decreased morphine potency compared to adults (2mos), with aged males requiring greater than 2x the concentration of morphine than their adult counterparts to produce equivalent analgesia (Fullerton et al., 2021). Similar results have been reported in prior preclinical studies, suggesting that aged rodents require higher doses of opioids to produce antinociception (Kavaliers

et al., 1983; Kramer and Bodnar, 1986; Webster et al., 1976). The mechanisms underlying the attenuation

of opioid potency in the aged population are currently unknown.

Morphine-induced analgesia is mediated primarily via binding to μ-opioid receptors (MOR), prototypical

G-protein coupled receptors (GPCRs) located predominantly on neuronal cell membranes (Martin, 1963;

Wolozin and Pasternak, 1981). Following agonist binding, coupled G-proteins undergo a conformational

change in which the guanosine diphosphate (GDP) bound to the inactive alpha subunit is replaced by

guanosine triphosphate (GTP), activating the subunit and promoting opioid signaling through interaction

with downstream effectors (Senese et al., 2020). MORs are coupled to a family of G-proteins called Gi/o

that signal via inhibition of adenylyl cyclase and subsequently decrease cyclic adenosine monophosphate

(cAMP) (Koehl et al., 2018; Bouchet et al., 2021). This G-protein signaling is downregulated or terminated

by **R**egulator of **G**-Protein **Si**gnaling (RGS) proteins, which act as GTPase activating proteins (GAPS),

hydrolyzing the active GTP back into GDP and terminating downstream signaling (Gerber et al., 2016). RGS

proteins, particularly RGS4 and RGS9-2, have been previously implicated in opioid-mediated G-protein

signaling, and have been shown to attenuate analgesia (Garnier et al., 2003; Psifogeorgou et al., 2007;

Avrampou et al., 2019; Senese et al., 2020). A primary target of MOR downstream signaling is the effector

adenylate cyclase (AC). AC enzymatically converts adenosine triphosphate (ATP) to the excitatory second

messenger cyclic adenosine monophosphate (cAMP). MOR binding inhibits AC, thereby reducing cAMP

expression, facilitating neuronal hyperpolarization, and promoting analgesia (Santhappan et al., 2015). Like

other G-protein coupled receptors, MOR signaling is subject to desensitization via the cellular mechanism

of receptor phosphorylation, which limits agonist binding and recruits arrestin proteins, and thus

attenuates opioid signaling. (Zhang et al., 1996; Yu et al., 1997; Groer et al., 2011).

The present studies test the hypothesis that the observed age-induced reduction in morphine potency in

the presence of chronic pain is mediated by changes in MOR signaling within the midbrain periaqueductal

gray, a CNS region critical for the opioid modulation of pain. (Basbaum et al., 1976; Behbehani and Fields,

1979; Morgan et al., 2006, 1992). The ventrolateral PAG (vlPAG) contains a large population of MOR+

neurons, and direct administration of MOR agonists into the PAG produces potent analgesia (Satoh et al.,

1983; Jensen and Yaksh, 1986; Bodnar et al., 1988; Loyd et al., 2008) In addition, intra-PAG administration

of MOR antagonists or lesions of PAG MOR significantly attenuate the analgesic effect of systemic

morphine, suggesting a critical role for PAG MOR in mediating morphine action (Ma and Han, 1991; Y.

Zhang et al., 1998; Loyd et al., 2008).

We have previously shown that aged rats exhibit reduced vlPAG MOR protein expression and reduced

MOR agonist binding in the vlPAG compared to adult rats (Fullerton et al., 2021), suggesting that

diminished levels of functioning MOR in the vlPAG contribute to the attenuated opioid potency seen in the

aged. The present studies build on these previous findings to assess age and sex differences in MOR

availability, ligand affinity, phosphorylation, G-protein activation, cAMP inhibition, and expression of RGS

proteins. We report that aged rats, and more specifically males, exhibit reduced MOR binding potential,

reduced G-protein activation, reduced cAMP inhibition, and increased RGS expression compared to adults,

with no change in MOR phosphorylation. Similar changes were observed in females regardless of age.

These findings identify several mechanisms whereby opioid signaling is attenuated in the aged rat, and

provide a framework for the development of novel pharmacological therapies to improve pain

management in the elderly.

**2 Materials and methods**

**2.1 Experimental subjects**

Adult (2-3mos) and aged (16-18mos) male and regularly cycling female Sprague–Dawley rats were used

in these experiments (Charles River Laboratories, Boston, MA). Rats were co-housed in same-sex pairs

on a 12:12 hour light/dark cycle (lights on at 08:00 am). Access to food and water was ad libitum

throughout the experiment, except during testing. All studies were approved by the Institutional Animal

Care and Use Committee at Georgia State University and performed in compliance with Ethical Issues of

the International Association for the Study of Pain and National Institutes of Health. All efforts were

made to reduce the number of rats used in these experiments and to minimize pain and suffering. All

assays were performed on separate cohorts of rats (n=8/sex/age; N=32), with the exception of the

radioligand binding and GTPγS assays, which were run simultaneously on PAG tissue from a single

cohort.

**2.2 Vaginal cytology**

Beginning ten days prior to testing, vaginal lavages were performed daily on adult and aged female rats

to confirm that all rats were cycling regularly and to keep daily records of the stage of estrous. Proestrus

was identified as a predominance of nucleated epithelial cells, and estrus was identified as a

predominance of cornified epithelial cells. Diestrus 1 was differentiated from diestrus 2 by the presence

of leukocytes. Rats that appeared between phases were noted as being in the more advanced stage

(Loyd et al., 2007).

**2.3 CFA-induced chronic pain treatment**

72 hours prior to experimentation, persistent inflammatory pain was induced by injection of complete

Freund's adjuvant (CFA; Mycobacterium tuberculosis; Sigma; 200μl), suspended in an oil/saline (1:1)

emulsion, into the plantar surface of the right hind paw. Edema was present within 24 hours of

injection, indicated by a >100% change in paw diameter, determined using calibrated calipers applied

midpoint across the plantar surface compared to handled paw.

**2.4 Membrane preparation for radioligand binding and GTPγS assays**

vlPAG membrane protein lysates were prepared from adult and aged, male and female, handled and

CFA treated rats to be used for radioligand binding and GTPγS assays. 72 hours post-CFA injection or

handling, rats were restrained using DecapiCones and decapitated. Brains were removed rapidly, flash-

frozen in 2-methyl butane on dry ice, and stored at −80°C. Ventrolateral PAG tissue from caudal PAG

(Bregma -7.5 through -8.5) was dissected from each brain using a straight edge razor at -20°C. On the

day of the assay, PAG sections were placed in ice-cold assay buffer (50 mM Tris-HCl, pH 7.4). Tissue was

homogenized with a glass dounce and centrifuged at 20,000g at 4°C for 30 min. The supernatant was

discarded and the pellet resuspended in assay buffer (Li et al., 2018; Shaqura et al., 2016b, 2016a;

Zollner et al., 2003). Membrane protein concentration was calculated using the Bradford Assay, and

lysates of vlPAG membrane protein from adult and aged, male and female, naïve and CFA treated rats

(n=4; N=32) were used immediately for radioligand binding and GTPγS assays.

**2.5 Saturation radioligand binding assay**

Saturation binding experiments were performed on vlPAG membranes using [$^3$H]DAMGO (specific

activity 50 Ci/mmol, American Radiolabeled Chemicals, Missouri). Briefly, 100 µg of membrane protein

was incubated with various concentrations of [$^3$H]DAMGO (0.5 to 5nM), in a total volume of 1ml of

binding buffer (50mM Tris-HCl pH 7.4). Nonspecific binding was defined as radioactivity remaining

bound in the presence of 10 µM unlabeled naloxone. At the end of the incubation period (1h at RT)

bound and free ligands were separated by rapid filtration over Whatman brand Grade GF/C glass filters

(Sigma-Aldrich) using a sampling vacuum manifold (MilliporeSigma). Filters were washed four times with

5ml of cold dH$_2$0. Bound radioactivity was determined by liquid scintillation spectrophotometry after

overnight extraction of the filters in 3ml of scintillation fluid. Specific activity was calculated by

subtracting the non-specific binding mean value at each concentration of [$^3$H]DAMGO. All experiments

were performed in triplicate. B$_{max}$ and K$_d$ values were determined by nonlinear regression analysis of

concentration-effect curve in GraphPad Prism 9.1.

**2.6 Agonist-stimulated [$^{35}$S]GTPγS binding**

DAMGO-stimulated [$^{35}$S]GTPγS binding to PAG membrane protein was assessed by incubating

membrane protein (100ug) in the presence or absence of [$^{35}$S]GTPγS (0.1nM) (specific activity 1250

Ci/mmol, American Radiolabeled Chemicals, Missouri) and various concentrations of DAMGO (2 to

30,000 nM) in assay buffer (20mM Tris, 10mM $MgCl_2$, 100mM NaCl, 0.2mM EGTA, pH 7.4) for 30min at

30°C. Stimulated [$^{35}$S]GTPγS binding was compared to unstimulated binding at each measurement point

and presented as percent basal binding. At the end of the incubation period, bound and free ligands

were separated by rapid filtration over Whatman brand Grade GF/B glass filters (Sigma-Aldrich) using a

sampling vacuum manifold (MilliporeSigma). Filters were washed four times with 5ml of cold buffer (50

mM Tris-HCl, pH 7.4). Bound radioactivity was determined by liquid scintillation spectrophotometry

after overnight extraction of the filters in 3ml of scintillation fluid. All experiments were performed in

duplicate. Efficacy (Emax) is defined as the maximum percent stimulation by an agonist; potency ($EC_{50}$) is

defined as the concentration of DAMGO required for half the maximal response. Emax and $EC_{50}$ values

were determined by nonlinear regression analysis of concentration-effect curves using Graph Pad Prism

9.1.

**2.7 Phosphorylated MOR analysis**

To assess the impact of advanced age and sex on MOR phosphorylation state, levels of MOR

phosphorylation were analyzed by Western blot. The lysis buffers used and general methods are the

same as reported in (Lei et al., 2017). Briefly, rat PAG samples were homogenized in ~500 µL of lysis

buffer using a glass dounce and rotated overnight at 4°C. The samples were then spun down at 13,000g

for 10 min at 4°C. The resulting lysates were quantified by DC Protein Assay Kit (Bio-Rad, #500-0111)

according to the manufacturer's instructions. 50-75 µg of soluble protein per sample (same for all

samples in a set) was run on 10% Bis-Tris Bolt PAGE gels (Fisher Scientific, #NW00100BOX), and wet-

transferred to nitrocellulose membrane (Protran 0.2 µm NC, #45-004-001 from Fisher Scientific) at 30V

at 4°C for 2-3 hours. Membranes were blocked using 5% nonfat dry milk in TBS, then blotted for primary

antibody target in 5% BSA in TBST overnight rocking at 4°C. Primary antibodies used were: pMOR

(1:1000, Bioss #bs-3724R), tMOR (1 µg/mL, R&D #MAB6866), and GAPDH (1:1000, Invitrogen #MA5-

15738). All 3 targets were always probed for on the same blot, using low pH stripping buffer between

each set. Secondary antibodies were used at 1:5000 each in 5% nonfat dry milk in TBST, rocking for ~1 hr

at room temperature. Secondary antibodies used were Goat-α-Mouse-IRDye800CW (Fisher Scientific

#NC9401841) and Goat-α-Mouse-IRDye680LT (Fisher Scientific #NC0046410). Target signal was acquired

using an Azure Sapphire imager using the near-infrared channels (658 and 784 nm). Band density was

quantified, and background subtracted using the onboard AzureSpot analysis software. Samples were

run with at least two representatives of each experimental group on the same gel. Adult and aged, male

and female, naïve and CFA treated rats (n=4; N=32) were used for these experiments. pMOR signal was

normalized to the same sample tMOR or GAPDH as indicated, and all samples were normalized to the

Adult Male average on each gel before combining data from different gels.

**2.8 Agonist-dependent cAMP inhibition**

DAMGO-dependent cAMP inhibition was analyzed using a cAMP assay. vlPAG membrane protein lysates

were prepared from adult and aged, male and female, handled and CFA treated rats. 72 hours post-CFA

injection or handling, rats were restrained using DecapiCones and decapitated. Brains were removed

rapidly, flash-frozen in 2-methyl butane on dry ice, and stored at −80°C. Ventrolateral PAG tissue from

caudal PAG (Bregma -7.5 through -8.5) was dissected from each brain using a straight edge razor at -

20°C. Briefly, vlPAG tissues were homogenized in ice-cold lysis buffer containing 0.25-M sucrose, 50-mM

Tris–HCl pH 7.5, 5-mM EGTA, 5-mM EDTA, 1-mM phenylmethylsulfonyl fluoride, 0.1-mM dithiothreitol,

and 10 µg/ml leupeptin. The homogenized tissues were then centrifuged at 1000×$g$ for 5 min (4 °C) and

the supernatant was centrifuged at 35000 ×$g$ for 10 min (Viganò et al., 2003). vlPAG membrane protein

samples (100ug) from adult and aged, male and female, naïve and CFA treated rats (n=4; N=32) were

incubated in 1uM forskolin (FSK) and in the presence or absence of 10uM DAMGO to stimulate

adenylate cyclase activity. cAMP levels in vlPAG tissue were determined using the LANCE *Ultra* cAMP

Detection Kit (Perkin Elmer) a time-resolved fluorescence resonance energy transfer (TR-FRET) cAMP

immunoassay. Data are plotted as % change in TR-FRET signal comparing FSK baseline measurements to

measurements following DAMGO-stimulated cAMP inhibition.

**2.9 Single-molecule fluorescence in situ hybridization**

Single-molecule fluorescent in situ hybridization (RNAscope) assays were used to determine mRNA

expression of OPRM1, RGS9-2, and RGS4 in the vlPAG. Rats were restrained using DecapiCones and

decapitated. Brains were removed rapidly, flash frozen in 2-methyl butane on dry ice, and stored RNase

free at −80 °C. Frozen tissue was sectioned in a 1:6 series of 20 μm coronal sections at −20 °C with a

Leica CM3050S cryostat. Sections were immediately mounted onto Superfrost slides (20 °C) and stored

at −80 °C until the time of the assay. vlPAG sections from adult and aged, male and female, naïve and

morphine treated rats were used (n=4; N=32). For morphine dosing paradigm see (Fullerton et al.,

2021). Tissue was processed for single-molecule fluorescence in situ hybridization (smFISH) according to

the RNAscope Multiplex Kit protocol (Advanced Cell Diagnostics) using probes for OPRM1, RGS4, and

RGS9-2. To facilitate cellular mRNA quantification within the vlPAG, sections were counterstained with

DAPI. mRNA puncta were visualized as fluorescent signal. Fluorescent images were captured on Zeiss

LSM 700 Confocal Microscope at 40x, and mRNA expression (target puncta/DAPI) was calculated using

Imaris software. To determine RGS4 and RGS9-2 expression levels in MOR+ neurons, quantification was

restricted to puncta located within 10μm of DAPI that co-expressed OPRM1 mRNA. mRNA expression

values were determined for the left and right ventrolateral subdivisions of each PAG image from two

representative levels of the mid-caudal PAG (Bregma −7.74 and −8.00). As there was no significant effect

of rostrocaudal level in the analyses, data were collapsed and presented as vlPAG and were averaged for

each rat. RGS4 and RGS9-2 mRNA values are expressed as the mean ± standard error of the mean (SEM).

All images were collected and analyzed by an experimenter blinded to the experimental condition.

**2.10 Statistical analysis and data presentation**

All values are reported as mean ± SEM. Data were assessed for normality and homogeneity of variance

using Shapiro-Wilk and Bartlett's tests. Significant main effects of sex, age, and treatment were assessed

using ANOVA; $p < 0.05$ was considered statistically significant. Tukey's post-hoc tests were conducted to

determine significant mean differences between groups that were a priori specified. Data are expressed

either as fmol/protein or nM DAMGO for radioligand binding assay, or Emax or EC50 for GTPγS assay.

**3 Results**

**3.1 Advanced age and sex impact vlPAG MOR binding properties**

To assess the impact of advanced age, biological sex, and pain on vlPAG MOR signaling, we first used

radioligand saturation binding assays to determine MOR binding parameters. The saturation binding

curves generated with [3H]DAMGO are shown in Fig. 1A. There was no significant impact of CFA

treatment on any measures, so CFA and handled groups were combined.

Kd values, indicative of receptor affinity, were determined using the concentration-effect curves

generated from each sample. Analysis of Kd values indicated no significant impact of age [$F_{(1,26)} = 0.770$,

p=0.388] or sex [$F_{(1,26)} = 1.241$, p=0.275], or a significant interaction [$F_{(1,26)} = 1.278$, p=0.269] (Fig. 1B).

Analyses of $B_{max}$ values, indicative of receptor availability, showed no significant impact of age [$F_{(1,28)} =$

2.661, p=0.114] or sex [$F_{(1,28)} = 0.002$, p=0.967], but a significant interaction [$F_{(1,28)} = 4.995$, p=0.034]. Post

hoc analysis showed a significant difference in vlPAG MOR availability between adult males and aged

males, as evidenced by reduced $B_{max}$ values in the aged males compared to their adult counterparts (p =

0.025)(Fig. 1C). $B_{max}$ values for aged males were markedly reduced compared to adult males, while adult

females and aged females exhibited smaller reductions in $B_{max}$ compared to their adult male

counterparts (%Δ -23.6 and -17.1, respectively. Fig. 1C).

Analyses of binding potential (BP) values, a measure that takes into consideration the density of

available receptors and the affinity of the receptor for its agonist, showed a significant impact of sex,

with males exhibiting greater binding potential than females [$F_{(1,28)}$ = 4.631, p=0.040] and a significant

interaction between age and sex [$F_{(1,28)}$ = 9.110, p=0.005]. No significant impact of age [$F_{(1,28)}$ = 3.312,

p=0.079] was observed. Post hoc analyses showed a significant difference in vlPAG MOR binding

potential between adult males and adult females (p = 0.006), and adult males and aged males (p =

0.010), indicating that adult males exhibit greater vlPAG MOR binding potential than their aged male

counterparts and their adult female counterparts (Fig. 1D). Adult females, aged males, and aged females

all exhibited marked reductions in MOR binding potential compared to their adult male counterparts

(%Δ -52, -48.7, and -40, respectively; Fig. 1D).

**Fig. 1.** Saturation binding curve of bound [3H]DAMGO (A). No significant effect of age or sex was observed in MOR affinity ($K_d$) (B). MOR availability ($B_{max}$) was significantly lower in aged males compared to adult males (C). Adult females, aged males, and aged females exhibited reduced MOR binding potential ($B_{max}/K_d$) compared to adult males, indicating attenuated agonist interaction at the receptor level (D). ns, not significant. *Significant difference between adult and aged males, or adult males and adult females; $p<0.05$ calculated by Tukey's post hoc test. Graphs indicate mean ± SEM. Values indicate % change from adult male.

**3.2 Advanced age and sex impact opioid-induced G-protein activation in the vlPAG**

We next conducted GTPγS binding assays to determine if advanced age or biological sex impacted MOR mediated G-protein activation. Preliminary studies revealed a significant main effect of chronic pain [$F_{(1, 23)}$ = 21.16, p = 0.0001]. To improve the translatability of these results to the target population of aged patients suffering from chronic pain, GTPγS experiments were performed exclusively on CFA-treated

rats. Concentration curves generated from [$^{35}$S]GTP$\gamma$S assays are shown in Fig. 2A. Analyses of $E_{max}$

values, a measure of G-protein availability combined with ligand efficacy, indicated no significant impact

of age [$F_{(1,24)}$ = 0.035, p = 0.8532] or sex [$F_{(1,25)}$ = 2.852, p = 0.1042], or a significant interaction [$F_{(1,24)}$ =

2.023, p = 0.1678] (Fig. 2B).

Analyses of $EC_{50}$ values, a measure of the concentration of DAMGO required for half-maximal G-protein

binding, indicated a significant impact of sex, with males exhibiting lower effective concentration values

than females, reflecting a higher potency of activation [$F_{(1,24)}$ = 5.177, p = 0.0321]. There was no

significant impact of age [$F_{(1,24)}$ = 2.339, p = 0.1393], and no significant interaction [$F_{(1,24)}$ = 2.473, p =

0.1289] (Fig. 2C). Compared to their adult male counterparts, aged males, adult females, and aged

females all exhibited marked increases in G-protein $EC_{50}$ as evidenced by their percent change values

(%$\Delta$ 60.5, 114.8, and 98.6, respectively), reflecting a lower potency of activation (Fig. 2C). Together,

these results suggest that opioid-induced G-protein signaling is impaired in the PAG of the aged rat and

the female rat.

Next, we calculated the coefficient ratio of analysis of $E_{max}$ / $EC_{50}$, which is indicative of overall G-protein

activation efficiency. A significant impact of age was noted, with adults exhibiting higher vlPAG G-

protein activation efficiency than aged rats [$F_{(1,24)}$ = 6.491, p = 0.0177]. There was also a significant

impact of sex, with males exhibiting higher vlPAG G-protein activation efficiency compared to females

[$F_{(1,24)}$ =10.39, p = 0.0036]. A significant interaction was also observed [$F_{(1,24)}$ = 13.93, p = 0.0010]. Post

hoc analyses showed a significant difference in vlPAG G-protein activation efficiency between adult

males and aged males (p = 0.0009), with adult males exhibiting greater vlPAG G-protein activation

efficiency than aged males. A significant difference in vlPAG G-protein activation efficiency between

adult males and adult females (p= 0.0003) was also observed, with adult males exhibiting greater vlPAG

G-protein activation efficiency than adult females (Fig. 2D). Aged males, adult females, and aged females

all exhibited marked reductions in G-protein activation efficiency compared to their adult male

counterparts (%Δ -40.0, -52.1, and -41.9, respectively; Fig. 2D).

**Fig 2.** [$^{35}$S]GTPγS binding curve of agonist-stimulated GTPγS binding (A). No significant effect of age or
sex was observed in $E_{max}$ (B). Aged males, adult females, and aged females exhibited increased $EC_{50}$
values compared to adult males, indicating that adult males have the greatest potency (C). Aged males,
adult females, and aged females exhibited reduced G-protein activation efficiency ($E_{max}/EC_{50}$) compared
to adult males, indicating attenuated G-protein signaling (D). ns, not significant. *Significant difference
between adult and aged males, or adult males and adult females; $p < 0.05$ calculated by Tukey's post hoc
test. Graphs indicate mean ± SEM. Values indicate % change from adult male.

**3.3 Advanced age and sex do not impact phosphorylated MOR in the vlPAG**

The results above showed significant reductions in MOR binding potential and G-protein activation

efficacy. Therefore, we next examined if there was an effect of age and sex on MOR phosphorylation, as

increased levels of pMOR would likely contribute to these reductions. Western blots were used to

determine the impact of advanced age on MOR phosphorylation at serine-375, a known site of

phosphorylation-mediated desensitization (Schulz et al., 2004). There was no significant impact of

treatment, so CFA and handled groups were combined. Although increased pMOR/tMOR was observed

in aged males compared to their adult counterparts (%Δ 34.0), no significant main effect of age [$F_{(1,41)}$ =

2.168, p = 0.1486], sex [$F_{(1,41)}$ = 1.807, p = 0.1863], or age x sex interaction [$F_{(1,41)}$ = 0.4615, p = 0.5008]

was observed (Fig. 3).

**Fig 3.** No significant impact of advanced age or biological sex on the ratio of phosphorylated MOR/ total
MOR. ns, not significant. Graph indicates mean ± SEM. Values indicate % change from adult male.

**3.4 Advanced age and sex impact opioid-induced cAMP inhibition in the vlPAG**

We next determined if advanced age, biological sex, or persistent pain impacted opioid-induced cAMP

inhibition in the vlPAG. Forskolin-induced cAMP release was used as a baseline measurement for each

group, while forskolin + DAMGO was used to assess the degree to which cAMP was inhibited by

DAMGO. Percent change from baseline was used to compare across treatment groups (Fig. 4A). There

was no significant impact of persistent pain on any measures, so CFA and handled groups were

combined. Our analyses revealed a significant main effect of age [$F_{(1,30)}$ = 9.314, p = 0.0047], a significant

main effect of sex [$F_{(1,30)}$ = 10.31, p = 0.0031], and a significant interaction [$F_{(1,30)}$ = 14.24, p = 0.0007].

Post hoc analyses showed a significant difference in cAMP inhibition between adult males and adult

females (p = 0.0002), adult males and aged males (p = 0.002), and adult males and aged females (p =

0.004), indicating that DAMGO elicits greater cAMP inhibition in adult males compared to aged males

and females (both adult and aged) (Fig. 4B).

**Fig 4.** TR-FRET immunoassay showing % change from FSK stimulated baseline and DAMGO inhibited cAMP (cAMP levels are inversely proportional to TR-FRET signal). Adult males exhibited the highest level of agonist-dependent % change in cAMP (A). Aged males and females exhibited reduced agonist-dependent cAMP inhibition compared to adult males, indicating attenuated downstream opioid signaling (B). *Significant difference between adult and aged males, or adult males and adult females; $p<0.05$ calculated by Tukey's post hoc test. Graphs indicate mean ± SEM. Values indicate % change from FSK baseline.

**328**    **3.5 Advanced age and sex impact RGS4 and RGS9-2 expression in the vlPAG**

**329**    Regulator of G-protein Signaling (RGS) proteins act as GAP accelerators to negatively modulate G-

**330**    protein signaling. RGS protein family members RGS4 and RGS9-2 are expressed in the vlPAG and have

**331**    both been shown to regulate opioid signaling by reversing G-protein activation. Therefore, we next used

**332**    smFISH to determine if RGS4 and RGS9-2 expression in the vlPAG was altered by advanced age and

**333**    biological sex. In these studies, following CFA administration, a cohort of rats was administered

**334**    morphine to examine the relationship between morphine $EC_{50}$ and RGS levels. However, no differences

**335**    were noted between the morphine and naive conditions, so these groups were combined. First, we

**336**    assessed total vlPAG expression of RGS4 and RGS9-2. These analyses revealed a significant main effect

**337**    of age on both RGS4 [$F_{(1,37)}$ = 18.15, p = 0.0001] and RGS9-2 [$F_{(1,41)}$ = 17.09, p < 0.0002], with aged rats

**338**    exhibiting increased levels of RGS4 and RGS9-2 mRNA compared to their adult counterparts (Fig. 5A &

**339**    B). No significant main effect of sex on RGS4 [$F_{(1,37)}$ = 0.0247, p = 0.8763] or RGS9-2 [$F_{(1,41)}$ = 0.4855, p =

**340**    0.4903], or significant interactions between age and sex for RGS4 [$F_{(1,37)}$ = 0.4994, p = 0.4842] or RGS9-2

**341**    [$F_{(1,41)}$ = 0.3246, p = 0.5723] were observed (Fig. 5A & B).

**342**    Following our assessment of total RGS4 and RGS9-2, we next restricted our analyses to RGS4 and RGS9-2

**343**    mRNA expressed on MOR+ neurons. Similar to what was noted above, a significant main effect of age on

**344**    expression of RGS4 [$F_{(1,41)}$ = 26.47, p < 0.0001] and RGS9-2 [$F_{(1,41)}$ = 21.69, p < 0.0001] was observed, with

**345**    aged rats exhibiting increased levels of RGS4 and RGS9-2 mRNA in MOR+ neurons compared to their

**346**    adult counterparts (Fig. 5C & D). No main effect of sex for RGS4 [$F_{(1,41)}$ = 0.7881, p = 0.3799] or RGS9-2

**347**    [$F_{(1,41)}$ = 0.0006, p = 0.9799], or significant interactions between age and sex for RGS4 [$F_{(1,41)}$ = 0.6583, p =

**348**    0.4219] or RGS9-2 [$F_{(1,41)}$ = 0.0075, p = 0.9314] were observed (Fig. 5C & D).

**349**

**350**

**Fig 5.** mRNA expression of RGS4 (A) and RGS9-2 (B) in the vlPAG was significantly greater in aged rats compared to adults. mRNA expression of RGS4 (C) and RGS9-2 (D) on MOR+ cells in the vlPAG was significantly greater in aged rats compared to adults. (E) Photomicrograph of RGS expression compared an adult male and an aged male PAG section. *Significant difference between adults and aged rats; $p<0.05$ calculated by 2x2 ANOVA. Graphs indicate mean ± SEM.

## 4 Discussion

The present studies are the first to show that advanced age results in a significant attenuation in MOR

signaling within the vlPAG of male and female rats. Specifically, aged males and females (regardless of

age) showed decreased MOR binding potential, decreased G-protein activation, and decreased agonist-

stimulated cAMP inhibition in comparison to adult males (Fig. 6). These changes, along with the

observed increase in RGS4 and RGS9-2 expression, provide a mechanism whereby morphine potency is

significantly reduced in aged rats (Fullerton et al., 2021). These changes also support our earlier

observed sex difference whereby adult females have a lower opioid analgesic potency than adult males,

equivalent to aged males and females.

Age-induced changes in PAG MOR signaling

|  | Morphine Potency | Binding Potential | G-Protein Activation | MOR Phosphor. | cAMP Inhibition | RGS mRNA |
|---|---|---|---|---|---|---|
| Adult ♂ | = | = | = | = | = | = |
| Adult ♀ | ↓ | ↓ | ↓ | = | ↓ | = |
| Aged ♂ | ↓ | ↓ | ↓ | = | ↓ | ↑ |
| Aged ♀ | ↓ | ↓ | ↓ | = | ↓ | ↑ |

**Fig 6.** Model and table of MOR signaling impairments in the vlPAG of the aged and female rat. Aged and female rats exhibit decreased MOR binding potential, decreased G-protein activation, and decreased cAMP inhibition, while aged rats alone demonstrate increased RGS expression. No impact of age or sex was found in phosphorylation of MOR at serine-375.

**Impact of advanced age and sex on MOR binding potential**

In our previous study, we reported that morphine potency to block persistent inflammatory pain is

reduced in aged and female rats, likely due to a reduction in DAMGO binding in the vlPAG (Fullerton et

al., 2021). Consistent with that finding, we report here that the vlPAG MOR binding potential in aged

males and females of either age is significantly reduced in comparison to adult male rats, providing

additional support that the decrease in opioid potency observed in these groups is driven, in part, by

reduced expression/availability of MOR in the vlPAG. Notably, we did not observe an impact of

advanced age or sex on the MOR's affinity for its ligand, as there were no significant effects of age or sex

on DAMGO $K_d$ values. However, aged males had reduced MOR availability as evidenced by lower $B_{max}$

values compared to their adult male counterparts. This result is consistent with our previous finding that

aged males exhibited reduced DAMGO binding in the vlPAG (assessed using autoradiography) (Fullerton

et al., 2021), and together, suggest an age-induced downregulation of PAG MOR in males matching a

consistently lower level of MOR expression in females of any age. This apparent decrease in MOR

expression may explain the reduced analgesic potency of morphine in females and aged males in our

earlier work (Fullerton et al., 2021).

Here we report no impact of persistent inflammatory pain on vlPAG MOR binding potential, consistent

with previous preclinical and clinical studies assessing the impact of chronic pain on MOR binding in the

CNS. For example, patients with chronic fibromyalgia pain exhibit no changes in MOR binding in the

PAG, despite having significant reductions in several brain regions, including the nucleus accumbens and

the amygdala (Harris et al., 2007). Similarly, no change in PAG MOR binding was noted as a result of

peripherally localized chronic pain generated from proximal nerve injury (Maarrawi et al., 2007).

Interestingly, centrally localized neuropathic pain (generated from supraspinal injury) was associated

with a significant reduction of PAG MOR binding, suggesting that the origin of chronic pain within the

CNS is critical in promoting alterations in neural MOR signaling (Maarrawi et al., 2007). Preclinical

studies in rodents also suggest that changes in MOR induced by persistent pain are CNS site-specific. For

example, persistent inflammatory pain induced by intraplantar CFA increased MOR binding in the dorsal

root ganglion at 24 and 96 hours post-CFA, (Mousa et al., 2001; Zollner et al., 2003), with no change in

MOR noted for the hypothalamus or spinal cord (Shaqura et al., 2004). Similarly, no change in PAG MOR

availability or expression was observed following sciatic nerve injury, although reductions were noted

for the insula, caudate putamen, and motor cortex (Thompson et al., 2018). Taken together, these

results suggest that chronic pain alters MOR signaling in a CNS region-specific manner that is dependent

on the site and modality of the pain.

**Impact of advanced age and sex on G-protein activation**

Agonist binding at MOR activates Gαi/o -proteins and downstream signaling cascades which are critical

for opioid-mediated hyperpolarization (Laugwitz et al., 1993; Connor and Christie, 1999; Koehl et al.,

2018). In the present study, we found that adult males exhibit greater G-protein activation efficiency

compared to aged males and adult and aged females, suggesting that the decreases in opioid potency

observed in the aged males and the females are driven, in part, by attenuated G-protein-MOR coupling

in the vlPAG. A significant impact of biological sex on G-protein potency ($EC_{50}$) was also observed, with

females and aged males showing a reduction in the potency of opioid-dependent G-protein activation,

as evidenced by increased $EC_{50}$ values. This reduction in vlPAG G-protein activation likely contributes to

previous studies reporting that females are less sensitive to the analgesic effects of opioids (Kest et al.,

2000; Loyd et al., 2008; Wang et al., 2014; Fullerton et al., 2018). Although the mechanism by which this

change in potency occurs is not known, the nature of the GTPγS assay is suggestive. The assay uses

membrane preparations without soluble signaling regulators like kinases present. Similarly, the GTPγS

molecule itself is not hydrolyzable, and is thus not subject to regulation by GAPs like the RGS proteins.

Considering these factors, the observed decrease in G protein activation potency may be due to the

observed decrease in opioid receptor expression/availability in the binding studies above.

Interestingly, this was the only assay where an impact of persistent inflammatory pain was observed.

Specifically, CFA treated rats had reduced $EC_{50}$ values, regardless of age or sex. Reductions in Gα subunit

expression have also been reported within both the rostral ventromedial medulla and spinal cord dorsal

horn following intraplantar CFA in adult male rats (Wattiez et al., 2017). The mechanism whereby

persistent inflammatory pain would impact G-protein activation, and not the other pharmacodynamics

of MOR, is not known. Also unknown are the mechanisms by which advanced age and sex alter G-

protein-MOR signaling. Previous studies have reported that advanced age results in a global

downregulation of Gαi/o, most notably in the prefrontal cortex (Young et al., 1991; Alemany et al., 2007;

de Oliveira et al., 2019), that is not associated with overall cell loss. Indeed, the Gαi/o subunit in

particular appears to be highly susceptible to the aging process, with estimates of reduced expression as

high as 65% in the frontal cortex, hippocampus, substantia nigra, and striatum (de Oliveira et al., 2019).

Although age-induced changes in Gαi/o expression were not assessed specifically in PAG, a widespread

reduction in Gαi/o as a function of advanced age would likely impact the PAG, thereby limiting MOR-G-

protein coupling and reducing both G-protein activation efficiency and opioid potency in the aged rat.

Alternatively, an uncoupling between MOR and Gαi/o and/or a switch in Gα subunit from Gαi/o to Gαs

cannot be ruled out (Gintzler and Chakrabarti, 2000, 2004, 2006). Indeed, a shift from Gαi/o to Gαs

would similarly increase adenylate cyclase activity, resulting in reduced hyperpolarization and decreased

morphine potency. Clearly, additional studies assessing the impact of advanced age on the expression of

Gα isoforms are warranted (Lamberts et al., 2011).

The present studies used the high-affinity MOR agonist DAMGO for both the radioligand binding and

GTPγS assay. As previous studies have reported that the pharmacodynamics of MOR may be agonist-

dependent, this possibility cannot be ruled out and future studies should consider incorporating at least

two agonists in all assays. Alternatively, a contribution by OPRM1 splice variants induced as a function of

age may also be present (Pan et al., 2005; Alvarez et al., 2002; Narayan et al., 2021). Indeed, specific

alternative MOR isoforms are known to promote ligand-dependent regulation of opioid signaling

(Pasternak et al., 2004; Oldfield et al., 2008). MOR-1B, a family of MOR isoforms expressed in the brain,

are indistinguishable from MOR-1 at the extracellular surface and the agonist binding pocket but elicit

alternative signaling properties that impact opioid-induced cell hyperpolarization, and dysregulation of

these MOR splice variants has been implicated in advanced age (Latorre and Harries, 2017; Li et al.,

2017; Wang et al., 2018). There are also reports of sex differences in the expression of MOR splice

variants in the PAG, with male rodents expressing higher levels of MOR-1B5, and females expressing

higher levels of MOR-1B1 and MOR-1B2 (Liu et al., 2018). The unique conformation of MOR isoforms

and subsequent alterations in receptor phosphorylation would directly influence the G-proteins to

which MORs couple, potentially resulting in biased agonism (Verzillo et al., 2014; Abrimian et al., 2021).

**Impact of advanced age and sex on MOR Phosphorylation**

The results of the radioligand binding and GTPγS assays suggested that aged and female rats exhibit

decreased receptor expression and activation potency. It has been previously shown that MOR can be

desensitized by phosphorylation in its basal state, thereby limiting agonist activation (Zhang et al., 1996;

Yu et al., 1997; Groer et al., 2011). Thus, we tested the hypothesis that the observed reductions in G-

protein activation efficiency were due to age- and sex-induced changes in basal MOR phosphorylation.

Although aged rats expressed higher levels of MOR phosphorylation at serine-375 compared to the

adults, this result was not significant, suggesting that the reductions in activation potency observed in

the vlPAG of aged and female rats is likely driven by an alternative mechanism.

MOR desensitization is mediated by G protein-coupled receptor kinase (GRK)-dependent

phosphorylation of the receptor, a mechanism by which β-arrestins bind to MOR, resulting in

endocytosis (J. Zhang et al., 1998; Schulz et al., 2004; Dang et al., 2009). Although our results suggest no

impact of advanced age or sex on GRK-mediated MOR phosphorylation, MOR signaling is also

desensitized via phosphorylation through the extracellular signal-regulated kinase 1 and 2 (ERK1/2)

pathway (Dang et al., 2009). ERK1/2 phosphorylation stimulates the activity of Gα-interacting protein

(GAIP), an RGS protein that acts as a GTPase activator to reduce opioid signaling at the level of G-protein

activation (Ogier-Denis et al., 2000). Indeed, pharmacological inhibition of ERK1/2 phosphorylation in a

rat model leads to improved morphine analgesia (Popiolek-Barczyk et al., 2014; Melkes et al., 2020).

Thus, the age- and sex-induced changes in morphine potency may be driven in part by age-induced

hyper-phosphorylation of ERK1/2 and result in the downregulation of G-protein signaling.

**Impact of advanced age and sex on cAMP inhibition**

Agonist binding at MOR elicits a conformational change in the receptor that allows the G-protein α and

βγ subunits to interact with downstream effectors. Notably, the Gα subunit binds to adenylate cyclase

(AC) and inhibits the conversion of adenosine triphosphate (ATP) to cAMP, thus limiting the activation of

cAMP-dependent protein kinase (PKA) and ultimately inducing higher levels of hyperpolarization

(Christie, 2008; Santhappan et al., 2015). In the present study, aged males and adult and aged females

exhibited significantly lower levels of DAMGO-induced cAMP inhibition compared to adult males,

suggesting an attenuated activity of the α subunit at the level of AC, or a weakened relationship

between AC and cAMP. This effect could easily be due to the observed decreases in MOR expression and

activation potency observed above. This attenuated cAMP inhibition in the aged and female rats likely

contributes to the observed reduction in opioid potency and may also be due, in part, to decreased

basal cAMP levels in the aged brain. Reductions in cAMP have been reported in several rodent brain

regions including the cortex, thalamus, hypothalamus, and midbrain (Puri and Volicer, 1981; Titus et al.,

2013; Kelly, 2018). Additional evidence suggests that aged rodents elicit widespread reductions in ATP

(Błaszczyk, 2020), which is required for cAMP synthesis. Not surprisingly, previous studies assessing the

impact of advanced age on cAMP expression and signaling were conducted exclusively in male rodents,

so it is not known whether advanced age impacts neural cAMP levels in a sex-specific manner.

Morphine binding to MOR results in the activation of Gai/o that primarily inhibits the activity of AC,

thereby reducing the ability to generate cAMP; in parallel, the βγ subunits activate G-protein inwardly

rectifying potassium (GIRK) channels to induce hyperpolarization and decrease excitability (Law et al.,

2000; Nobles et al., 2005; Yudin and Rohacs, 2018). Advanced age is concomitant with increased levels

of neuroinflammation and greater production of reactive oxygen species (ROS), both of which have been

implicated in age-induced potassium channel dysfunction (Sparkman and Johnson, 2008; Sesti, 2016;

Lieberman et al., 2020). Consistent with this, aged Sprague Dawley rats (26-30 mos) exhibit reduced

potassium currents in hippocampal neurons (Alshuaib et al., 2001). Further experimentation assessing

the impact of advanced age and sex on GIRK channel-mediated hyperpolarization, in addition to cAMP-

mediated hyperpolarization, will provide a more detailed description of the impact of advanced age and

sex on downstream opioid signaling in the PAG.

**Impact of advanced age and sex on RGS protein levels**

G-protein activation by MOR results in the exchange of GDP for GTP (Senese et al., 2020). GTP-bound Gα

proteins are inactivated by hydrolysis of the GTP back to GDP, catalyzed by the intrinsic GTPase activity

of Gα proteins. This inactivation mechanism is modulated by the activity of Regulator of G-protein

signaling (RGS) proteins. RGS proteins reduce the magnitude of GPCR signaling via enhancement of

GTPase activity of the α subunit of the G-protein complex. By promoting the hydrolysis of the alpha-

bound GTP during the active state, RGS proteins hasten the return of the α subunit to the GDP-bound

inactive state (Roman and Traynor, 2011). RGS proteins play a critical role in negatively modulating

opioid signaling, as morphine analgesia is increased in male mice lacking RGS9-2 (Garzón et al., 2001;

Zachariou et al., 2003) and overexpression of RGS4 attenuates MOR signaling in reconstituted MORs in

vitro (Ippolito et al., 2002). In the present study, we observed increased expression of RGS4 and RGS9-2

in the vlPAG of aged rats compared to adults, suggesting greater GTPase activity and thus reduced G-

protein signaling in the aged. Similar results were observed when the analysis was limited to MOR+ cells,

indicating that opioid-induced G-protein signaling is subjected to greater negative regulation in the

vlPAG of the aged rat. These results are consistent with Kim et al. (2005) who reported higher RGS9-2

protein levels in the PAG of 1-year old male rats compared to 3-week-old rats (Kim et al., 2005). In

humans, advanced age is associated with increased RGS4 expression in the prefrontal cortex (Rivero et

al., 2010), suggesting that the age-induced increase in RGS4 in our experiment may not be specific to the

vlPAG. These results provide a mechanism by which advanced age could decrease analgesic potency and

downstream cAMP signaling, although they do not provide a mechanism by which adult females

demonstrate lower potency and signaling than adult males.

**Summary and Conclusions**

The present studies are the first to show that sex and advanced age leads to an attenuation in vlPAG

opioid signaling compared to adult male rats. Taken together with our previous findings, these results

suggest that age- and sex-induced reductions in vlPAG MOR expression and binding, combined with

attenuated downstream MOR signaling, contribute to the diminished opioid potency reported in aged

and female rats (Fullerton et al., 2021). The results of our analyses demonstrate that aged and female

rats exhibit reductions in MOR expression/availability, G-protein activation, and cAMP inhibition, and

increased G-protein regulation by RGS proteins, each of which provides potential therapeutic targets for

improved pain management in the elderly.

The nature of the results observed also suggest potential mechanisms of action for sex and age to

impact analgesic potency. First, the normal ligand affinity (Kd) and lack of receptor phosphorylation

suggests that unit receptor performance is normal. When combined with the observed decrease in

expression/availability (Bmax), this suggests that one major mechanism is a decrease in receptor

expression instead of a change in unit receptor performance. When considering the G protein coupling

analysis, since soluble regulators (like RGS proteins) are absent in this assay and the $^{35}$S-GTPγS is non-

hydrolyzable, the decrease in G protein coupling potency is likely due to decreased receptor availability.

Downstream of the G proteins, we also observed a decrease in cAMP inhibition. This downstream effect

could be due to the observed increase in RGS expression, decreasing G protein activity, leading to less

cAMP inhibition. However, RGS expression was not increased in adult females. Overall then our results

suggest two major complementary mechanisms; 1) decreased receptor expression/availability in adult

females and aged males and females, and 2) increased RGS expression in aged males and females.

Further investigation could also uncover additional complementary sex- and age-related mechanisms

that help explain why opioids show lower analgesic potency in females and the aged.

**Funding:** This work was supported by the National Institutes of Health [Grant DA041529 AZM, Grant
UG3DA047717 JMS, Grant P50 MH100023 LJY]; and Georgia State University [Molecular Basis of Disease
Fellowship EFF, Provost's Dissertation Fellowship EFF].

**Author Contributions:** AZM and EFF designed the research and wrote the paper. MK, JMS, and LJY aided
in the design of the research. EFF, MK, and JMS performed the research, EFF, AZM, and JMS analyzed
the data.

## References

Abrimian, A., Kraft, T., Pan, Y.-X., 2021. Endogenous Opioid Peptides and Alternatively Spliced Mu Opioid Receptor Seven Transmembrane Carboxyl-Terminal Variants. Int. J. Mol. Sci. 22, 3779. https://doi.org/10.3390/ijms22073779

Alemany, R., Perona, J.S., Sánchez-Dominguez, J.M., Montero, E., Cañizares, J., Bressani, R., Escribá, P.V., Ruiz-Gutierrez, V., 2007. G protein-coupled receptor systems and their lipid environment in health disorders during aging. Biochim. Biophys. Acta BBA - Biomembr. 1768, 964–975. https://doi.org/10.1016/j.bbamem.2006.09.024

Alshuaib, W.B., Hasan, S.M., Cherian, S.P., Mathew, M.V., Hasan, M.Y., Fahim, M.A., 2001. Reduced potassium currents in old rat CA1 hippocampal neurons. J. Neurosci. Res. 63, 176–184. https://doi.org/10.1002/1097-4547(20010115)63:2<176::AID-JNR1009>3.0.CO;2-H

Alvarez, V.A., Arttamangkul, S., Dang, V., Salem, A., Whistler, J.L., von Zastrow, M., Grandy, D.K., Williams, J.T., 2002. μ-Opioid Receptors: Ligand-Dependent Activation of Potassium Conductance, Desensitization, and Internalization. J. Neurosci. 22, 5769–5776. https://doi.org/10.1523/JNEUROSCI.22-13-05769.2002

Avrampou, K., Pryce, K.D., Ramakrishnan, A., Sakloth, F., Gaspari, S., Serafini, R.A., Mitsi, V., Polizu, C., Swartz, C., Ligas, B., Richards, A., Shen, L., Carr, F.B., Zachariou, V., 2019. RGS4 Maintains Chronic Pain Symptoms in Rodent Models. J. Neurosci. 39, 8291–8304. https://doi.org/10.1523/JNEUROSCI.3154-18.2019

Basbaum, A.I., Clanton, C.H., Fields, H.L., 1976. Opiate and stimulus-produced analgesia: functional anatomy of a medullospinal pathway. Proc. Natl. Acad. Sci. 73, 4685–4688. https://doi.org/10.1073/pnas.73.12.4685

Behbehani, M.M., Fields, H.L., 1979. Evidence that an excitatory connection between the periaqueductal gray and nucleus raphe magnus mediates stimulation produced analgesia. Brain Res. 170, 85–93. https://doi.org/10.1016/0006-8993(79)90942-9

Błaszczyk, J.W., 2020. Energy Metabolism Decline in the Aging Brain—Pathogenesis of Neurodegenerative Disorders. Metabolites 10, 450. https://doi.org/10.3390/metabo10110450

Bodnar, R.J., Williams, C.L., Lee, S.J., Pasternak, G.W., 1988. Role of mu 1-opiate receptors in supraspinal opiate analgesia: a microinjection study. Brain Res. 447, 25–34. https://doi.org/10.1016/0006-8993(88)90962-6

Bouchet, C.A., McPherson, K.B., Li, M., Traynor, J.R., Ingram, S.L., 2021. Mice Expressing Regulators of G protein Signaling–insensitive Gαo Define Roles of $\mu$ Opioid Receptor G $\alpha$ o and G $\alpha$ i Subunit Coupling in Inhibition of Presynaptic GABA Release. Mol. Pharmacol. 100, 217–223. https://doi.org/10.1124/molpharm.121.000249

Christie, M.J., 2008. Cellular neuroadaptations to chronic opioids: tolerance, withdrawal and addiction. Br. J. Pharmacol. 154, 384–396. https://doi.org/10.1038/bjp.2008.100

Connor, M., Christie, M.D., 1999. Opioid receptor signalling mechanisms. Clin. Exp. Pharmacol. Physiol.
26, 493–499. https://doi.org/10.1046/j.1440-1681.1999.03049.x

Dahlhamer, J., 2018. Prevalence of Chronic Pain and High-Impact Chronic Pain Among Adults — United
States, 2016. MMWR Morb. Mortal. Wkly. Rep. 67. https://doi.org/10.15585/mmwr.mm6736a2

Dampier, C.D., Smith, W.R., Wager, C.G., Kim, H.-Y., Bell, M.C., Miller, S.T., Weiner, D.L., Minniti, C.P.,
Krishnamurti, L., Ataga, K.I., Eckman, J.R., Hsu, L.L., McClish, D., McKinlay, S.M., Molokie, R., Osunkwo, I.,
Smith-Whitley, K., Telen, M.J., Sickle Cell Disease Clinical Research Network (SCDCRN), 2013. IMPROVE
trial: a randomized controlled trial of patient-controlled analgesia for sickle cell painful episodes:
rationale, design challenges, initial experience, and recommendations for future studies. Clin. Trials
Lond. Engl. 10, 319–331. https://doi.org/10.1177/1740774513475850

Dang, V.C., Napier, I.A., Christie, M.J., 2009. Two Distinct Mechanisms Mediate Acute -Opioid Receptor
Desensitization in Native Neurons. J. Neurosci. 29, 3322–3327.
https://doi.org/10.1523/JNEUROSCI.4749-08.2009

de Oliveira, P.G., Ramos, M.L.S., Amaro, A.J., Dias, R.A., Vieira, S.I., 2019. Gi/o-Protein Coupled Receptors
in the Aging Brain. Front. Aging Neurosci. 11. https://doi.org/10.3389/fnagi.2019.00089

Domenichiello, A.F., Ramsden, C.E., 2019. The silent epidemic of chronic pain in older adults. Prog.
Neuropsychopharmacol. Biol. Psychiatry 93, 284–290. https://doi.org/10.1016/j.pnpbp.2019.04.006

Fullerton, E.F., Doyle, H.H., Murphy, A.Z., 2018. Impact of sex on pain and opioid analgesia: a review.
Curr. Opin. Behav. Sci. 23, 183–190. https://doi.org/10.1016/j.cobeha.2018.08.001

Fullerton, E.F., Rubaharan, M., Karom, M.C., Hanberry, R.I., Murphy, A.Z., 2021. Advanced age
attenuates the antihyperalgesic effect of morphine and decreases μ-opioid receptor expression and
binding in the rat midbrain periaqueductal gray in male and female rats. Neurobiol. Aging 98, 78–87.
https://doi.org/10.1016/j.neurobiolaging.2020.10.020

Garnier, M., Zaratin, P.F., Ficalora, G., Valente, M., Fontanella, L., Rhee, M.-H., Blumer, K.J., Scheideler,
M.A., 2003. Up-regulation of regulator of G protein signaling 4 expression in a model of neuropathic pain
and insensitivity to morphine. J. Pharmacol. Exp. Ther. 304, 1299–1306.
https://doi.org/10.1124/jpet.102.043471

Garzón, J., Rodríguez-Díaz, M., López-Fando, A., Sánchez-Blázquez, P., 2001. RGS9 proteins facilitate
acute tolerance to mu-opioid effects: RGS9 facilitates opioid tachyphylaxis. Eur. J. Neurosci. 13, 801–
811. https://doi.org/10.1046/j.0953-816x.2000.01444.x

Gerber, K.J., Squires, K.E., Hepler, J.R., 2016. Roles for Regulator of G Protein Signaling Proteins in
Synaptic Signaling and Plasticity. Mol. Pharmacol. 89, 273–286. https://doi.org/10.1124/mol.115.102210

Gintzler, A.R., Chakrabarti, S., 2006. Post-opioid receptor adaptations to chronic morphine; altered
functionality and associations of signaling molecules. Life Sci. 79, 717–722.
https://doi.org/10.1016/j.lfs.2006.02.016

Gintzler, A.R., Chakrabarti, S., 2004. Chronic morphine-induced plasticity among signalling molecules.
Novartis Found. Symp. 261, 167–176; discussion 176-180, 191–193.

Gintzler, A.R., Chakrabarti, S., 2000. Opioid tolerance and the emergence of new opioid receptor-
coupled signaling. Mol. Neurobiol. 21, 21–33. https://doi.org/10.1385/MN:21:1-2:021

Groer, C.E., Schmid, C.L., Jaeger, A.M., Bohn, L.M., 2011. Agonist-directed Interactions with Specific β-
Arrestins Determine μ-Opioid Receptor Trafficking, Ubiquitination, and Dephosphorylation. J. Biol.
Chem. 286, 31731–31741. https://doi.org/10.1074/jbc.M111.248310

Harris, R.E., Clauw, D.J., Scott, D.J., McLean, S.A., Gracely, R.H., Zubieta, J.-K., 2007. Decreased central
mu-opioid receptor availability in fibromyalgia. J. Neurosci. Off. J. Soc. Neurosci. 27, 10000–10006.
https://doi.org/10.1523/JNEUROSCI.2849-07.2007

Ippolito, D.L., Temkin, P.A., Rogalski, S.L., Chavkin, C., 2002. N-terminal Tyrosine Residues within the
Potassium Channel Kir3 Modulate GTPase Activity of Gαi. J. Biol. Chem. 277, 32692–32696.
https://doi.org/10.1074/jbc.M204407200

Jensen, T.S., Yaksh, T.L., 1986. III. Comparison of the antinociceptive action of Mu and delta opioid
receptor ligands in the periaqueductal gray matter, medial and paramedial ventral medulla in the rat as
studied by the microinjection technique. Brain Res. 372, 301–312. https://doi.org/10.1016/0006-
8993(86)91138-8

Kavaliers, M., Hirst, M., Teskey, G.C., 1983. Ageing, opioid analgesia and the pineal gland. Life Sci. 32,
2279–2287. https://doi.org/10.1016/0024-3205(83)90427-7

Kelly, M.P., 2018. Cyclic nucleotide signaling changes associated with normal aging and age-related
diseases of the brain. Cell. Signal. 42, 281–291. https://doi.org/10.1016/j.cellsig.2017.11.004

Kest, B., Sarton, E., Dahan, A., Fisher, D.M., 2000. Gender Differences in Opioid-mediated Analgesia.
Anesthesiology 93, 539–547. https://doi.org/10.1097/00000542-200008000-00034

Kim, K.J., Moriyama, K., Han, K.R., Sharma, M., Han, X., Xie, G., Palmer, P.P., 2005. Differential expression
of the regulator of G protein signaling RGS9 protein in nociceptive pathways of different age rats. Dev.
Brain Res. 160, 28–39. https://doi.org/10.1016/j.devbrainres.2005.08.003

Koehl, A., Hu, H., Maeda, S., Zhang, Y., Qu, Q., Paggi, J.M., Latorraca, N.R., Hilger, D., Dawson, R., Matile,
H., Schertler, G.F.X., Granier, S., Weis, W.I., Dror, R.O., Manglik, A., Skiniotis, G., Kobilka, B.K., 2018.
Structure of the μ-opioid receptor–G i protein complex. Nature 558, 547–552.
https://doi.org/10.1038/s41586-018-0219-7

Kramer, E., Bodnar, R.J., 1986. Age-related decrements in morphine analgesia: A parametric analysis.
Neurobiol. Aging 7, 185–191. https://doi.org/10.1016/0197-4580(86)90041-2

Lamberts, J.T., Jutkiewicz, E.M., Mortensen, R.M., Traynor, J.R., 2011. Mu-Opioid Receptor Coupling to
Gαo Plays an Important Role in Opioid Antinociception. Neuropsychopharmacology 36, 2041–2053.
https://doi.org/10.1038/npp.2011.91

Latorre, E., Harries, L.W., 2017. Splicing regulatory factors, ageing and age-related disease. Ageing Res.
Rev. 36, 165–170. https://doi.org/10.1016/j.arr.2017.04.004

Laugwitz, K.L., Offermanns, S., Spicher, K., Schultz, G., 1993. mu and delta opioid receptors differentially
couple to G protein subtypes in membranes of human neuroblastoma SH-SY5Y cells. Neuron 10, 233–
242. https://doi.org/10.1016/0896-6273(93)90314-h

Law, P.Y., Wong, Y.H., Loh, H.H., 2000. Molecular mechanisms and regulation of opioid receptor
signaling. Annu. Rev. Pharmacol. Toxicol. 40, 389–430.
https://doi.org/10.1146/annurev.pharmtox.40.1.389

Lei, W., Mullen, N., McCarthy, S., Brann, C., Richard, P., Cormier, J., Edwards, K., Bilsky, E.J., Streicher,
J.M., 2017. Heat-shock protein 90 (Hsp90) promotes opioid-induced anti-nociception by an ERK
mitogen-activated protein kinase (MAPK) mechanism in mouse brain. J. Biol. Chem. 292, 10414–10428.
https://doi.org/10.1074/jbc.M116.769489

Li, H., Wang, Z., Ma, T., Wei, G., Ni, T., 2017. Alternative splicing in aging and age-related diseases.
Transl. Med. Aging 1, 32–40. https://doi.org/10.1016/j.tma.2017.09.005

Li, X., Shaqura, M., Mohamed, D., Beyer, A., Yamada, S., Mousa, S.A., Schäfer, M., 2018. Pro- versus
Antinociceptive Nongenomic Effects of Neuronal Mineralocorticoid versus Glucocorticoid Receptors
during Rat Hind Paw Inflammation. Anesthesiology 128, 796–809.
https://doi.org/10.1097/ALN.0000000000002087

Lieberman, O.J., Bartolini, F., Miniaci, M.C., 2020. GIRK channels in Alzheimer's disease. Aging 12,
18793–18794. https://doi.org/10.18632/aging.104026

Liu, A., Zhang, H., Qin, F., Wang, Qisheng, Sun, Q., Xie, S., Wang, Qian, Tang, Z., Lu, Z., 2018. Sex
Associated Differential Expressions of the Alternatively Spliced Variants mRNA of OPRM1 in Brain
Regions of C57BL/6 Mouse. Cell. Physiol. Biochem. 50, 1441–1459. https://doi.org/10.1159/000494644

Loyd, D.R., Morgan, M.M., Murphy, A.Z., 2007. Morphine Preferentially Activates the Periaqueductal
Gray –Rostral Ventromedial Medullary Pathway in the Male Rat: A Potential Mechanism for Sex
Differences in Antinociception. Neuroscience 147, 456–468.
https://doi.org/10.1016/j.neuroscience.2007.03.053

Loyd, D.R., Wang, X., Murphy, A.Z., 2008. Sex Differences in μ-Opioid Receptor Expression in the Rat
Midbrain Periaqueductal Gray Are Essential for Eliciting Sex Differences in Morphine Analgesia. J.
Neurosci. 28, 14007–14017. https://doi.org/10.1523/JNEUROSCI.4123-08.2008

Ma, Q.P., Han, J.S., 1991. Naloxone blocks the release of opioid peptides in periaqueductal gray and N.
accumbens induced by intra-amygdaloid injection of morphine. Peptides 12, 1235–1238.
https://doi.org/10.1016/0196-9781(91)90200-9

Maarrawi, J., Peyron, R., Mertens, P., Costes, N., Magnin, M., Sindou, M., Laurent, B., Garcia-Larrea, L.,
2007. Differential brain opioid receptor availability in central and peripheral neuropathic pain. Pain 127,
183–194. https://doi.org/10.1016/j.pain.2006.10.013

Manchikanti, L., Fellows, B., Ailinani, H., Pampati, V., 2010. Therapeutic use, abuse, and nonmedical use
of opioids: a ten-year perspective. Pain Physician 13, 401–435.

Martin, W.R., 1963. Analgesic and antipyretic drugs in Physiological Pharmacology. Pharmacol Rev,
Academic Press, NY 275–312.

Melkes, B., Markova, V., Hejnova, L., Novotny, J., 2020. β-Arrestin 2 and ERK1/2 Are Important
Mediators Engaged in Close Cooperation between TRPV1 and μ-Opioid Receptors in the Plasma
Membrane. Int. J. Mol. Sci. 21, 4626. https://doi.org/10.3390/ijms21134626

Morgan, M.M., Fossum, E.N., Stalding, B.M., King, M.M., 2006. Morphine antinociceptive potency on
chemical, mechanical, and thermal nociceptive tests in the rat. J. Pain 7, 358–366.
https://doi.org/10.1016/j.jpain.2005.12.009

Morgan, M.M., Heinricher, M.M., Fields, H.L., 1992. Circuitry linking opioid-sensitive nociceptive
modulatory systems in periaqueductal gray and spinal cord with rostral ventromedial medulla.
Neuroscience 47, 863–871. https://doi.org/10.1016/0306-4522(92)90036-2

Mousa, S.A., Zhang, Q., Sitte, N., Ji, R., Stein, C., 2001. beta-Endorphin-containing memory-cells and mu-
opioid receptors undergo transport to peripheral inflamed tissue. J. Neuroimmunol. 115, 71–78.
https://doi.org/10.1016/s0165-5728(01)00271-5

Naples, J.G., Gellad, W.F., Hanlon, J.T., 2016. The Role of Opioid Analgesics in Geriatric Pain
Management. Clin. Geriatr. Med. 32, 725–735. https://doi.org/10.1016/j.cger.2016.06.006

Narayan, A., Hunkele, A., Xu, J., Bassoni, D.L., Pasternak, G.W., Pan, Y.-X., 2021. Mu Opioids Induce
Biased Signaling at the Full-Length Seven Transmembrane C-Terminal Splice Variants of the mu Opioid
Receptor Gene, Oprm1. Cell. Mol. Neurobiol. 41, 1059–1074. https://doi.org/10.1007/s10571-020-
00973-5

Nobles, M., Benians, A., Tinker, A., 2005. Heterotrimeric G proteins precouple with G protein-coupled
receptors in living cells. Proc. Natl. Acad. Sci. U. S. A. 102, 18706–18711.
https://doi.org/10.1073/pnas.0504778102

Ogier-Denis, E., Pattingre, S., El Benna, J., Codogno, P., 2000. Erk1/2-dependent phosphorylation of
Galpha-interacting protein stimulates its GTPase accelerating activity and autophagy in human colon
cancer cells. J. Biol. Chem. 275, 39090–39095. https://doi.org/10.1074/jbc.M006198200

Oldfield, S., Braksator, E., Rodriguez-Martin, I., Bailey, C.P., Donaldson, L.F., Henderson, G., Kelly, E.,
2008. C-terminal splice variants of the mu-opioid receptor: existence, distribution and functional
characteristics. J. Neurochem. 104, 937–945. https://doi.org/10.1111/j.1471-4159.2007.05057.x

Pan, Y.-X., Xu, J., Bolan, E., Moskowitz, H.S., Xu, M., Pasternak, G.W., 2005. Identification of four novel
exon 5 splice variants of the mouse mu-opioid receptor gene: functional consequences of C-terminal
splicing. Mol. Pharmacol. 68, 866–875. https://doi.org/10.1124/mol.105.011858

Pasternak, D.A., Pan, L., Xu, J., Yu, R., Xu, M.-M., Pasternak, G.W., Pan, Y.-X., 2004. Identification of three
new alternatively spliced variants of the rat mu opioid receptor gene: dissociation of affinity and
efficacy. J. Neurochem. 91, 881–890. https://doi.org/10.1111/j.1471-4159.2004.02767.x

Patel, K.V., Guralnik, J.M., Dansie, E.J., Turk, D.C., 2013. Prevalence and Impact of Pain among Older
Adults in the United States: Findings from the 2011 National Health and Aging Trends Study. Pain 154.
https://doi.org/10.1016/j.pain.2013.07.029

Popiolek-Barczyk, K., Makuch, W., Rojewska, E., Pilat, D., Mika, J., 2014. Inhibition of intracellular
signaling pathways NF-κB and MEK1/2 attenuates neuropathic pain development and enhances
morphine analgesia. Pharmacol. Rep. PR 66, 845–851. https://doi.org/10.1016/j.pharep.2014.05.001

Prostran, M., Vujović, K.S., Vučković, S., Medić, B., Srebro, D., Divac, N., Stojanović, R., Vujović, A.,
Jovanović, L., Jotić, A., Cerovac, N., 2016. Pharmacotherapy of Pain in the Older Population: The Place of
Opioids. Front. Aging Neurosci. 8. https://doi.org/10.3389/fnagi.2016.00144

Psifogeorgou, K., Papakosta, P., Russo, S.J., Neve, R.L., Kardassis, D., Gold, S.J., Zachariou, V., 2007. RGS9-
2 is a negative modulator of μ-opioid receptor function: RGS9-2 modulates μ-opioid receptor
responsiveness. J. Neurochem. 103, 617–625. https://doi.org/10.1111/j.1471-4159.2007.04812.x

Puri, S.K., Volicer, L., 1981. Age-related changes of cyclic nucleotide levels in rat brain regions. Mech.
Ageing Dev. 15, 239–242. https://doi.org/10.1016/0047-6374(81)90132-9

Reddy, K.S.K., Naidu, M.U.R., Rani, P.U., Rao, T.R.K., 2012. Human experimental pain models: A review of
standardized methods in drug development. J. Res. Med. Sci. Off. J. Isfahan Univ. Med. Sci. 17, 587–595.

Rivero, G., Gabilondo, A.M., García-Sevilla, J.A., La Harpe, R., Morentín, B., Javier Meana, J., 2010.
Characterization of regulators of G-protein signaling RGS4 and RGS10 proteins in the postmortem
human brain. Neurochem. Int. 57, 722–729. https://doi.org/10.1016/j.neuint.2010.08.008

Roman, D.L., Traynor, J.R., 2011. Regulators of G Protein Signaling (RGS) Proteins as Drug Targets:
Modulating G-Protein-Coupled Receptor (GPCR) Signal Transduction: Miniperspective. J. Med. Chem. 54,
7433–7440. https://doi.org/10.1021/jm101572n

Santhappan, R., Crowder, A.T., Gouty, S., Cox, B.M., Côté, T.E., 2015. Mu opioid receptor activation
enhances regulator of G protein signaling 4 association with the mu opioid receptor/G protein complex
in a GTP-dependent manner. J. Neurochem. 135, 76–87. https://doi.org/10.1111/jnc.13222

Satoh, M., Kubota, A., Iwama, T., Wada, T., Yasui, M., Fujibayashi, K., Takagi, H., 1983. Comparison of
analgesic potencies of mu, delta and kappa agonists locally applied to various CNS regions relevant to
analgesia in rats. Life Sci. 33 Suppl 1, 689–692. https://doi.org/10.1016/0024-3205(83)90596-9

Schulz, S., Mayer, D., Pfeiffer, M., Stumm, R., Koch, T., Höllt, V., 2004. Morphine induces terminal μ-
opioid receptor desensitization by sustained phosphorylation of serine-375. EMBO J. 23, 3282–3289.
https://doi.org/10.1038/sj.emboj.7600334

Senese, N.B., Kandasamy, R., Kochan, K.E., Traynor, J.R., 2020. Regulator of G-Protein Signaling (RGS)
Protein Modulation of Opioid Receptor Signaling as a Potential Target for Pain Management. Front. Mol.
Neurosci. 13. https://doi.org/10.3389/fnmol.2020.00005

Sesti, F., 2016. Oxidation of K+ Channels in Aging and Neurodegeneration. Aging Dis. 7, 130–135.
https://doi.org/10.14336/AD.2015.0901

Shaqura, M., Li, X., Al-Khrasani, M., Shakibaei, M., Tafelski, S., Fürst, S., Beyer, A., Kawata, M., Schäfer,
M., Mousa, S.A., 2016a. Membrane-bound glucocorticoid receptors on distinct nociceptive neurons as
potential targets for pain control through rapid non-genomic effects. Neuropharmacology 111, 1–13.
https://doi.org/10.1016/j.neuropharm.2016.08.019

Shaqura, M., Li, X., Al-Madol, M.A., Tafelski, S., Beyer-Koczorek, A., Mousa, S.A., Schäfer, M., 2016b.
Acute mechanical sensitization of peripheral nociceptors by aldosterone through non-genomic
activation of membrane bound mineralocorticoid receptors in naive rats. Neuropharmacology 107, 251–
261. https://doi.org/10.1016/j.neuropharm.2016.03.032

Shaqura, M.A., Zöllner, C., Mousa, S.A., Stein, C., Schäfer, M., 2004. Characterization of mu opioid
receptor binding and G protein coupling in rat hypothalamus, spinal cord, and primary afferent neurons
during inflammatory pain. J. Pharmacol. Exp. Ther. 308, 712–718.
https://doi.org/10.1124/jpet.103.057257

Sparkman, N.L., Johnson, R.W., 2008. Neuroinflammation Associated with Aging Sensitizes the Brain to
the Effects of Infection or Stress. Neuroimmunomodulation 15, 323–330.
https://doi.org/10.1159/000156474

Sullivan, M.D., Edlund, M.J., Fan, M.-Y., DeVries, A., Braden, J.B., Martin, B.C., 2010. Risks for possible
and probable opioid misuse among recipients of chronic opioid therapy in commercial and medicaid
insurance plans: The TROUP Study. Pain 150, 332–339. https://doi.org/10.1016/j.pain.2010.05.020

Thompson, S.J., Pitcher, M.H., Stone, L.S., Tarum, F., Niu, G., Chen, X., Kiesewetter, D.O., Schweinhardt,
P., Bushnell, M.C., 2018. Chronic neuropathic pain reduces opioid receptor availability with associated
anhedonia in rat. Pain 159, 1856–1866. https://doi.org/10.1097/j.pain.0000000000001282

Titus, D.J., Furones, C., Kang, Y., Atkins, C.M., 2013. Age-dependent alterations in cAMP signaling
contribute to synaptic plasticity deficits following traumatic brain injury. Neuroscience 231, 182–194.
https://doi.org/10.1016/j.neuroscience.2012.12.002

Verzillo, V., Madia, P.A., Liu, N.-J., Chakrabarti, S., Gintzler, A.R., 2014. Mu-opioid receptor splice
variants: sex-dependent regulation by chronic morphine. J. Neurochem. 130, 790–796.
https://doi.org/10.1111/jnc.12768

Viganò, D., Rubino, T., Di Chiara, G., Ascari, I., Massi, P., Parolaro, D., 2003. μ opioid receptor signaling in
morphine sensitization. Neuroscience 117, 921–929. https://doi.org/10.1016/S0306-4522(02)00825-4

Wang, K., Wu, D., Zhang, H., Das, A., Basu, M., Malin, J., Cao, K., Hannenhalli, S., 2018. Comprehensive
map of age-associated splicing changes across human tissues and their contributions to age-associated
diseases. Sci. Rep. 8, 10929. https://doi.org/10.1038/s41598-018-29086-2

Wang, Y.-J., Huang, P., Blendy, J.A., Liu-Chen, L.-Y., 2014. Brain region- and sex-specific alterations in
DAMGO-stimulated [ [35] S]GTPγS binding in mice with *Oprm1* A112G: A118G SNP on MOPR activation.
Addict. Biol. 19, 354–361. https://doi.org/10.1111/j.1369-1600.2012.00484.x

Wattiez, A.-S., Walder, R.Y., Sande, C.M., White, S.R., Hammond, D.L., 2017. Peripheral inflammatory
injury alters the relative abundance of Gα subunits in the dorsal horn of the spinal cord and in the rostral
ventromedial medulla of male rats. Mol. Pain 13, 1744806917715210.
https://doi.org/10.1177/1744806917715210

Webster, G.W., Shuster, L., Eleftheriou, B.E., 1976. Morphine analgesia in mice of different ages. Exp.
Aging Res. 2, 221–233. https://doi.org/10.1080/03610737608257178

Wolozin, B.L., Pasternak, G.W., 1981. Classification of multiple morphine and enkephalin binding sites in
the central nervous system. Proc. Natl. Acad. Sci. 78, 6181–6185.
https://doi.org/10.1073/pnas.78.10.6181

Young, L.T., Warsh, J.J., Li, P.P., Siu, K.P., Becker, L., Gilbert, J., Hornykiewicz, O., Kish, S.J., 1991.
Maturational and aging effects on guanine nucleotide binding protein immunoreactivity in human brain.
Brain Res. Dev. Brain Res. 61, 243–248. https://doi.org/10.1016/0165-3806(91)90137-8

Yu, Y., Zhang, L., Yin, X., Sun, H., Uhl, G.R., Wang, J.B., 1997. μ Opioid Receptor Phosphorylation,
Desensitization, and Ligand Efficacy*. J. Biol. Chem. 272, 28869–28874.
https://doi.org/10.1074/jbc.272.46.28869

Yudin, Y., Rohacs, T., 2018. Inhibitory Gi/O-coupled receptors in somatosensory neurons: Potential
therapeutic targets for novel analgesics. Mol. Pain 14, 1744806918763646.
https://doi.org/10.1177/1744806918763646

Zachariou, V., Georgescu, D., Sanchez, N., Rahman, Z., DiLeone, R., Berton, O., Neve, R.L., Sim-Selley, L.J.,
Selley, D.E., Gold, S.J., Nestler, E.J., 2003. Essential role for RGS9 in opiate action. Proc. Natl. Acad. Sci.
100, 13656–13661. https://doi.org/10.1073/pnas.2232594100

Zhang, J., Ferguson, S.S.G., Barak, L.S., Bodduluri, S.R., Laporte, S.A., Law, P.-Y., Caron, M.G., 1998. Role
for G protein-coupled receptor kinase in agonist-specific regulation of -opioid receptor responsiveness.
Proc. Natl. Acad. Sci. 95, 7157–7162. https://doi.org/10.1073/pnas.95.12.7157

Zhang, L., Yu, Y., Mackin, S., Weight, F.F., Uhl, G.R., Wang, J.B., 1996. Differential μ Opiate Receptor
Phosphorylation and Desensitization Induced by Agonists and Phorbol Esters (∗). J. Biol. Chem. 271,
11449–11454. https://doi.org/10.1074/jbc.271.19.11449

Zhang, Y., Du, L.N., Wu, G.C., Cao, X.D., 1998. Modulation of intrathecal morphine-induced
immunosuppression by microinjection of naloxone into periaqueductal gray. Zhongguo Yao Li Xue Bao
19, 519–522.

Zollner, C., Shaqura, M.A., Bopaiah, C.P., Mousa, S., Stein, C., Schafer, M., 2003. Painful inflammation-
induced increase in mu-opioid receptor binding and G-protein coupling in primary afferent neurons.
Mol. Pharmacol. 64, 202–210. https://doi.org/10.1124/mol.64.2.202

 **Figure Legends**

**Fig. 1.** Saturation binding curve of bound [3H]DAMGO (A). No significant effect of age or sex was
observed in MOR affinity ($K_d$) (B). MOR availability ($B_{max}$) was significantly lower in aged males compared
to adult males (C). Adult females, aged males, and aged females exhibited reduced MOR binding
potential ($B_{max}/K_d$) compared to adult males, indicating attenuated agonist interaction at the receptor
level (D). ns, not significant. *Significant difference between adult and aged males, or adult males and
adult females; $p < 0.05$ calculated by Tukey's post hoc test. Graphs indicate mean ± SEM. Values indicate
% change from adult male.

**Fig 2.** [$^{35}$S]GTPγS binding curve of agonist-stimulated GTPγS binding (A). No significant effect of age or
sex was observed in $E_{max}$ (B). Aged males, adult females, and aged females exhibited increased $EC_{50}$
values compared to adult males, indicating that adult males have the greatest potency (C). Aged males,
adult females, and aged females exhibited reduced G-protein activation efficiency ($E_{max}/EC_{50}$) compared
to adult males, indicating attenuated G-protein signaling (D). ns, not significant. *Significant difference
between adult and aged males, or adult males and adult females; $p < 0.05$ calculated by Tukey's post hoc
test. Graphs indicate mean ± SEM. Values indicate % change from adult male.

**Fig 3.** No significant impact of advanced age or biological sex on the ratio of phosphorylated MOR/ total
MOR. ns, not significant. Graph indicates mean ± SEM. Values indicate % change from adult male.

**Fig 4.** TR-FRET immunoassay showing % change from FSK stimulated baseline and DAMGO inhibited
cAMP (cAMP levels are inversely proportional to TR-FRET signal). Adult males exhibited the highest level
of agonist-dependent % change in cAMP (A). Aged males and females exhibited reduced agonist-
dependent cAMP inhibition compared to adult males, indicating attenuated downstream opioid
signaling (B). *Significant difference between adult and aged males, or adult males and adult females;
$p < 0.05$ calculated by Tukey's post hoc test. Graphs indicate mean ± SEM. Values indicate % change from
FSK baseline.

**Fig 5.** mRNA expression of RGS4 (A) and RGS9-2 (B) in the vlPAG was significantly greater in aged rats
compared to adults. mRNA expression of RGS4 (C) and RGS9-2 (D) on MOR+ cells in the vlPAG was
significantly greater in aged rats compared to adults. (E) Photomicrograph of RGS expression compared
an adult male and an aged male PAG section. *Significant difference between adults and aged rats;
$p < 0.05$ calculated by 2x2 ANOVA. Graphs indicate mean ± SEM.

**Fig 6.** Model and table of MOR signaling impairments in the vlPAG of the aged and female rat. Aged and
female rats exhibit decreased MOR binding potential, decreased G-protein activation, and decreased
cAMP inhibition, while aged rats alone demonstrate increased RGS expression. No impact of age or sex
was found in phosphorylation of MOR at serine-375

**A** [35S]GTPγS binding

- adult male
- adult female
- aged male
- aged female

% Basal Binding

nM DAMGO

**B** G-protein efficacy (Emax)

ns

% Basal Binding

male | female | male | female
adult | | aged |

**C** G-protein potency (EC50)

*

EC50 (nM DAMGO)

60.5 | 114.8 | 98.6

adult | aged | adult | aged
male | | female |

**D** G-protein activation efficiency

*

Emax / EC50

-40.0 | -52.1 | -41.9

adult | aged | adult | aged
male | | female |

# Phosphorylated MOR

**A**

## cAMP Assay

**B**

## Agonist-dependent cAMP inhibition

**A** vlPAG RGS4 expression

**B** vlPAG RGS9-2 expression

**C** MOR+ RGS4 expression

**D** MOR+ RGS9-2 expression

**E** DAPI + OPRM1 + RGS4 + RGS9-2

adult male

aged male

20µm

# Age-induced changes in PAG MOR signaling

| | Morphine Potency | Binding Potential | G-Protein Activation | MOR Phosphor. | cAMP Inhibition | RGS mRNA | |
|---|---|---|---|---|---|---|---|
| **Adult ♂** | = | = | = | = | = | = | |
| **Adult ♀** | ↓ | ↓ | ↓ | = | ↓ | = | |
| **Aged ♂** | ↓ | ↓ | ↓ | = | ↓ | ↑ | |
| **Aged ♀** | ↓ | ↓ | ↓ | = | ↓ | ↑ | |