Seasonality and competition select for variable germination behavior in perennials

Hanna ten Brink[*1,2,5], Thomas R. Haaland[*2,3] & Øystein H. Opedal[4]

1: Swiss Institute of Aquatic Research, EAWAG, Switzerland. hannatenbrink@gmail.com

2: Department of Evolutionary Biology and Environmental Studies, University of Zurich, Switzerland.

3: Centre for Biodiversity Dynamics, Department of Biology, Norwegian University of Science and Technology, Norway. trhaaland@gmail.com

4: Department of Biology, Biodiversity Unit, Lund University, Sweden. oystein.opedal@biol.lu.se

5: Department of Coastal Systems, Royal Netherlands Institute for Sea Research

* Co-first authorship and corresponding authors.

Keywords: dormancy, competition, bet-hedging, phenology, seed size, environmental variation

Abstract: The common occurrence of within-population variation in germination behavior and associated traits such as seed size has long fascinated evolutionary ecologists. In annuals, unpredictable environments are known to select for bet-hedging strategies causing variation in dormancy duration and germination strategies. Variation in germination timing and associated traits is also commonly observed in perennials, and often tracks gradients of environmental predictability. Although bet-hedging is thought to occur less frequently in long-lived organisms, these observations suggest a role of bet-hedging strategies in perennials occupying unpredictable environments. We use complementary analytical and evolutionary simulation models of within-individual variation in germination behavior in seasonal environments to show how bet-hedging interacts with fluctuating selection, life-history traits, and competitive asymmetries among germination strategies. We reveal substantial scope for bet-hedging to produce variation in germination behavior in long-lived plants, when "false starts" to the growing season results in either competitive advantages or increased mortality risk for alternative germination strategies. Additionally, we find that lowering adult survival may, in contrast to classical bet-hedging theory, result in less spreading of germination by decreasing density-dependent competition. These models extend insights from bet-hedging theory to perennials and explore how competitive communities may be affected by ongoing changes in climate and seasonality patterns.

Introduction

Accurate timing of seasonal phenology is key to population persistence in unpredictable environments. The timing of emergence can have strong and direct effects on individual fitness, and is expected to be subject to strong selection driving local adaptation (Donohue et al. 2010). Indeed, emergence behavior (e.g. patterns of germination or hatching) is often found to vary in predictable ways along environmental gradients (Meyer et al. 1995; Venable

2007; Wagmann et al. 2012; Simons 2014; Pinceel et al. 2017; Rubio de Casas et al. 2017;
Torres-Martínez et al. 2017; Scholl et al. 2020). In most plants and many animals, the timing
of emergence is controlled by dormancy (Vleeshouwers et al. 1995; Finch-Savage and
Leubner-Metzger 2006; Baskin and Baskin 2014), and the evolution of emergence behavior is
thus tightly linked to the evolution of dormancy mechanisms (Varpe 2017).

Theoretical models of seed dormancy and germination behavior have a long history (Cohen
1966; Venable and Lawlor 1980; Ellner 1985; Geritz et al. 2018; Hughes 2018; Kortessis and
Chesson 2019). Most of these, however, have focused on specific systems such as annual
plants in desert environments. This system provides a natural starting point because one
striking observation demands an explanation: a fraction of the seeds produced each year fail
to germinate the next year, and instead lie dormant for another year before germinating at the
beginning of the second growing season following their dispersal (Gremer and Venable
2014). While this reduction in number of seedlings may seem a waste of resources most years,
such a strategy has been identified as a risk-spreading adaptation to avoid complete
recruitment failure in the event of a bad year. Prolonged seed dormancy and variation in
germination timing have come to represent the archetype of a bet-hedging strategy, i.e. a
genotype-level strategy that sacrifices some short-term (arithmetic-mean) fitness gains in
order to lower its fitness variance over time (Levins 1962; Cohen 1966; Venable 2007). Such
bet-hedging strategies have received considerable empirical and theoretical attention as a
major mode of adaptation to unpredictable environments (Seger and Brockmann 1987;
Simons 2011; Starrfelt and Kokko 2012).

Empirical and theoretical studies of bet-hedging have yielded a good understanding of how
different types of risk-spreading adaptations co-evolve, interact, and cancel each other out.
For example, if phenotypic polymorphisms or continuous variation ensures that fitness
correlations among related individuals are sufficiently low, then there is less need for other
costly bet-hedging strategies at the individual level such as 'safer' offspring phenotypes that
are weaker competitors but better able to cope with environmental variation (Venable and
Brown 1988; Starrfelt and Kokko 2012; Haaland et al. 2020; Escobar et al. 2021). The scope
for bet-hedging is also lowered if the spatiotemporal scale of environmental variation (what is
often called environmental 'grain' in bet-hedging terms) already ensures that related
individuals experience different conditions and thus uncorrelated fitness returns. In a spatially
fine-grained environment relatives may be exposed to a wide range of different conditions at
any given time (as opposed to coarse-grained environments where all individuals experience
the same conditions), and in such cases, genotype-level diversification or other bet-hedging
strategies are not selectively favored (Levins 1962; Starrfelt and Kokko 2012). Furthermore,
longer lifespans with a larger number of selective events over which fitness can accumulate
(e.g. an individual experiencing multiple consecutive breeding seasons with variable
conditions) also reduces the need for any variance reduction if fitness payoffs among selective
events are uncorrelated (Haaland et al. 2019).

Given these insights, it is unsurprising that most models of bet-hedging, as well as many of
the best-documented empirical examples, consider rather specific types of short-lived
organisms with discrete generations occupying highly variable, coarse-grained environments,
such as germination strategies of desert annuals, or overwintering strategies for organisms
living in ephemeral ponds (Furness et al. 2015; García-Roger et al. 2016; Wang and Rogers
2018). However, this focus has also led to a knowledge gap regarding the evolution of

seemingly 'risk-spreading' traits in longer-lived organisms, as these may require other
explanations than bet-hedging in its strictest sense. Variation in seed germination behavior is
seen not only in annuals, and is at least sometimes associated with variation in seed size (Rees
1996; Simons and Johnston 2000; Susko and Lovett-Doust 2000; Norden et al. 2009; Harel et
al. 2011; Martins et al. 2019). A classic example of within-individual variation in seed size is
seed heteromorphism, as observed e.g. in many Asteraceae and Chenopodiaceae (reviewed in
Venable 1985; Imbert 2002), where distinct seed morphs differ in their dispersal abilities and
germination behavior (Venable et al. 1987; Brändel 2004; Fumanal et al. 2007; Yao et al.
2010; Wang et al. 2012). Continuous variation in seed size is also common in natural
populations, and may relate to variation in germination behavior (Pélabon et al. 2021). Within
species, larger seeds are generally more likely to germinate and/or germinate earlier than do
smaller ones (Biere 1991; Simons and Johnston 2000; Tremayne and Richards 2000;
Galloway 2001; Pélabon et al. 2005), but the opposite pattern of smaller seeds breaking
dormancy and germinating earlier is also observed (Martins et al. 2019).

Seed size variation is thus a common mechanism for producing seeds with variable duration
of dormancy and/or variable germination time both within individuals as well as within an
inflorescence (Pélabon et al. 2021). As argued, existing bet-hedging models alone are not able
to explain the ubiquity of seed heteromorphism or continuous seed size variation as observed
in perennials (Martins et al. 2019). Although several other non-mutually exclusive hypotheses
exist, including but not limited to variance-sensitivity due to asymmetrical fitness costs versus
benefits (Bednekoff 1996; Bårdsen et al. 2008), fluctuating selection or fluctuations in priority
effects (Chesson 2000; ten Brink et al. 2020), competitive asymmetries among offspring types
(Geritz 1995) and negative frequency-dependent selection (Metcalf et al. 2015; Poethke et al.
2016), these have seldom been analyzed jointly (but see Rees et al. 2004). In Scholl et al.'s
(2020) recent analysis of the flora of the south-east United States, annuals were not detectably
more likely to exhibit seed heteromorphism than were perennials, suggesting that selection for
within- and/or among-individual variation in germination behavior may be common also in
long-lived plants where bet-hedging is less strongly favored.

Here, we explore the evolution of individual variation in dormancy of long-lived plants in
seasonally varying environments, using both analytical and simulation models covering a
range of ecological scenarios. First, our analytical model derives general predictions for when
within- and/or among-plant variation in germination behavior can be favored as an adaptation
to seasonal environmental variation in perennials. To this end we use an adaptive dynamics
approach assuming that plants produce two discrete seed morphs, an "early" seed with a short
dormancy period, and a "late" seed with a long dormancy period (for example, "small" and
"large" seeds, respectively). This analysis examines the conditions favoring plants producing
seeds of variable dormancy duration (frequencies of seed morphs different from 0 or 1), and
identifies the shape of the function relating the (rainfall-dependent) benefits and costs of
producing early relative to late seeds as the key determinant of whether such mixed strategies
are stable. The potential for among-individual variation (presence of evolutionary branching
points) is also examined. We complement this analytical model with individual-based
evolutionary simulations of dormancy duration modelled as a near-continuous trait via freely-
evolving weekly germination probabilities. This model thus examines the predictions from the
analytical model without the restriction of only allowing two discrete seed morphs, and makes
explicit the mechanisms affecting benefits and costs of early and late seeds. Specifically, early
seeds germinating during the dry season gain a competitive advantage if they survive until

reproduction, but suffer high mortality if there is not enough rain between their germination and the beginning of the wet season ("false starts"). We let spatiotemporal rainfall patterns vary predictably and unpredictably within and among years, thus emulating the wet- and dry-season dynamics seen in large parts of the tropics where intermittent dry-season rains can trigger germination of certain seed types. However, we note that this setup applies equally well to other stochastic seasonal changes between harsh and favorable conditions, such as the onset of warmer spring weather in temperate regions where "false starts" of warm temperatures may still be followed by harmful frosts. In concert, our two models allow us to tease apart the relative effects of bet-hedging, competition and fluctuating selection on the evolution of variable germination strategies.

1. Selection for within-plant variation in germination behavior

We first present an analytical model where we assume that plants can produce two seed morphs, here "early" and "late" seeds, that may differ in e.g. size or shell thickness and hence the amount of rainfall required for germinating (Norden et al. 2009). Each year consists of a dry and a wet season, where the dry season is characterized by scarce intermittent rainfalls, affecting the germination and survival probability of early seeds. These early seedlings emerging during intermittent dry-season rains have a high probability of dying due to drought, and therefore only a small chance to recruit into the adult populations, but may persist if there is sufficient excess rain in the following weeks leading up to the wet season. Late seeds, on the other hand, remain dormant through any intermittent rains, and germinate at the beginning of the safe wet season. However, any early seedlings that survived until the beginning of the wet season may now have a competitive advantage over late seeds, as they have a head start to growth, and therefore a higher probability of recruitment into the adult population. Thus, the relative advantage b of being an early *vs.* a late seed in a given year depends on the amount of excess dry-season rainfall in that year, a random environmental variable we term X.

We designate as f the fraction of early seeds produced by a plant, so that a fraction $1–f$ are late seeds. We model a density-regulated population of long-lived plants, where adult plants survive with probability s_a, and all seeds compete for recruiting into the vacant space $1-s_a$ in the adult population. We incorporate asymmetric competition among early and late seeds in a 'lottery' for vacant space. Early seeds get the relative weight $b(x)f$ against the weight $1–f$ of all late seeds, such that the probability of drawing an early seed in the lottery for vacant space becomes

$$\frac{b(x)f}{b(x)f+1-f}.$$

(1)

Next, assuming that the population is sufficiently isolated (no immigration) and that the quality of environmental conditions X vary randomly from year to year without any autocorrelation, the relative fitness $w(f',f)$ of strategy f' (a rare mutant) against a resident population that has strategy f is given by the geometric average growth rate, which is the expectation over the log growth rates λ, i.e.

$$w(f',f) = \mathrm{E}[\log \lambda(f',f)] = \mathrm{E}[\log\left(s_a + (1-s_a)\frac{b(x)f'+1-f'}{b(x)f+1-f}\right)],$$

(2)

where E indicates the expectation over all environmental conditions $X=x$ (see appendix A for derivation of λ). The sign of $w(f',f)$ determines whether the mutant can invade the population

174 (positive sign) or not (negative sign) – examples are shown in Fig. 1a-b where expectations
175 are calculated over 10.000 realizations of $x\sim\text{Normal}(0,1)$.

Figure 1: Results from adaptive dynamics model of the evolution of variable seed germination timing (strategy f
defined as fraction of early-germinating seeds) in a long-lived plant ($s_a=0.9$). Results are shown when the sum of
the benefits and costs of early relative to late germination relate to environmental conditions x (excess dry-
season rainfall) with an exponential (left column, using $\beta=1$) or logistic (right column, using $\alpha=1$) function $b(x)$.
Top row: Pairwise invasibility plots showing fitness surfaces of invasion fitness (eq. 2) as a function of the
resident strategy f and the invader strategy f'. Dark shaded regions indicate regions where the mutant strategy
can invade. Bottom row: Selection gradients (eq. 4) on f' showing the direction and strength of evolution for
different f (x-axis) and values of adult survival s_a (line color). Values of f where the lines cross 0 (horizontal
dashed line) represent convergence stable ESSs (see eqs. 5&6).

We now use an adaptive dynamics approach to analyse the evolutionary consequences of
equation 2 (Kisdi and Meszéna 1995; Geritz et al. 1998). Assuming that $E[X]=0$ (the
expectation for "excess" rain is zero) and that the two seed types fare equally well in average
years when there is neither an excess nor deficit of rain between the early seeds germinating

and the end of the dry season (i.e., $b(0)=1$), we can examine the conditions leading to an
advantage for early seeds.

The selection gradient acting on a rare mutant with a germination strategy (fraction of early
seeds) f' slightly different from the resident population's strategy f, is obtained by
differentiating $w(f', f)$ with respect to f' and evaluating at $f'=f$. Differentiating eq. 2 yields:

196
$$\frac{\partial w}{\partial f'}(f', f) = \mathrm{E}\left[\frac{(1-s_\mathrm{a})\frac{b(x)-1}{b(x)f+1-f}}{s_\mathrm{a}+(1-s_\mathrm{a})\frac{b(x)f'+1-f'}{b(x)f+1-f}}\right],$$
(3)

197   such that the selection gradient becomes

198
$$\frac{\partial w}{\partial f'}\Big|_{f'=f} = (1-s_\mathrm{a})\mathrm{E}\left[\frac{b(x)-1}{b(x)f+1-f}\right].$$
(4)

Intuitively, the strength of selection decreases with increasing adult survival $0<s_\mathrm{a}<1$, as the
competition among seedlings for annual recruitment contributes a smaller part of the
population. Examples of selection gradients for two functions $b(x)$ with different s_a are shown
in Fig. 1c-d. Cases where $\frac{\partial w}{\partial f'}\Big|_{f'=f} = 0$ (colored lines cross the dashed black lines in Fig. 1c-
d) represent so-called singular strategies (Geritz et al. 1998). In order to determine whether
these are attractors or repellors we can examine the sign of the derivative of eq. 4 taken with
respect to the resident strategy f:

206
$$\frac{d}{df}\left(\frac{\partial w}{\partial f'}\Big|_{f'=f}\right) = -(1-s_a)\mathrm{E}\left[\left(\frac{b(x)-1}{b(x)f+1-f}\right)^2\right].$$
(5)

207   Because the squared term is positive, eq. 5 is always negative, indicating that all singular
208   strategies are attractors. Further, computing the second derivative of equation 2 with respect
209   to the mutant strategy f' and again evaluating at $f'=f$ allows us to determine whether the
210   singular strategies are evolutionarily stable strategies (negative sign) or branching points
211   (positive sign). Thus, the ESS criterion is:

212
$$\frac{\partial^2 w}{\partial f'^2}\Big|_{f'=f} = -(1-s_a)^2\mathrm{E}\left[\left(\frac{b(x)-1}{b(x)f+1-f}\right)^2\right] < 0,$$
(6)

which reveals already that $\frac{\partial^2 w}{\partial f'^2}(f', f)$ is always negative (as both squared terms are always
positive), thus there is no potential for evolutionary branching and any singular strategy is
necessarily a convergence stable ESS (Geritz et al. 1998). In other words, in this simple case
of two discrete seed morphs (early and late seeds), the evolution of their frequency f cannot
reach two distinct values at evolutionary equilibrium.

The need to compute expectations over realizations of the random variable X makes it
difficult to exactly identify ESSs in the current model. However, we can approximate
selection gradients using the delta method if we assume that X follows a sufficiently simple
distribution with all odd-numbered moments equal to 0 ($\mathrm{E}[X]=0$, symmetric distribution), a
small variance V, and even-numbered moments have an upper bound depending on V. Then,
recalling that $b(0)=1$ by assumption, we obtain the approximation

224
$$\frac{\partial w}{\partial f'}\Big|_{f'=f} \approx \frac{V}{2}(1-s_\mathrm{a})[b''(0) - 2b'(0)^2 f].$$
(7)

225   Setting this selection gradient equal to 0, we see that there can be one singular strategy only,

$$226 \qquad\qquad\qquad\qquad f \approx \frac{b''(0)}{2b'(0)^2}. \qquad\qquad\qquad\qquad (8)$$

In order for such an ESS to exist and represent a mixed strategy, it much lie between 0 and 1,
with conditions given by the shape of the function $b(x)$, which determines the advantage of
early relative to late seeds for a given level of excess dry-season rain x. Firstly, to ensure $f>0$,
$b''(0) > 0$ is demanded (eq. 8), i.e. $b(x)$ must be accelerating (convex) around 0. Secondly, to
ensure $f<1$, eq. 8 gives the condition $b''(0) < 2b'(0)^2$, i.e. it must not accelerate too fast (there
is an upper bound to how convex it can be, given by the steepness of the function). For
example (as shown in Fig. 1a,c), if $b(x) = e^{-\beta x}$, then we have an ESS for all $\beta>0$, since $b''(0) >$
0 and $b''(0) < 2b'(0)$ is fulfilled. The ESS becomes $f = b''(0) / 2b'(0) = 0.5$. In contrast, if b is
a logistic function $b(x)=2/[1+e^{-\alpha x}]$ (Fig. 1b,d), the conditions for an ESS are not fulfilled: $b'(0)$
$= 0$ and $b''(0)<0$ whenever $\alpha>0$.

Overall, this model reveals that there is ample scope for within-individual variation in
germination strategies when plants can evolve to produce fractions f and 1-f of distinct early
and late seeds, but that among-individual variation in f should not occur in such a scenario
because all ESSs are convergence stable. The evolution of a mixed strategy $0 < f < 1$ requires
accelerating rewards for germinating early with increasing amounts of excess rainfall x, but
only within a limited range. If rewards are not accelerating, then f evolves towards 0 (if
benefits of being early are not sufficient, there is no point in producing risky early-
germinating seeds), whereas if rewards accelerate too much, f evolves towards 1 and all seeds
germinate early. Interpreting $b(x)$ as the sum of the functions relating early seed dry-season
survival as a function of x and the competitive advantage of early seeds (given their survival)
over late-germinating seeds in the lottery for vacant space in the adult population, this
analysis provides a baseline expectation for our individual-based model with continuous
variation in germination time.

2. Individual-based model of mean and variance in dormancy

Model setup

Here, we build an individual-based simulation model that allows plants to produce a
continuous distribution of seeds (rather than just e.g. "early" and "late" seeds). We model a
population of asexually reproducing perennials occupying a wet-dry seasonal tropical
environment where the wet and dry seasons both last half the year (26 weeks). We assume
that seed germination is determined by the dormancy period d, which functions as an obligate
"biological clock" preventing germination until a certain time d has passed (see table 1 for an
overview of all mathematical notation and baseline parameter values used in the individual-
based model). Germination of a seed occurs with probability 1 as soon as its age (in weeks) is
greater than d_z. We set no constraints on the evolution of flexible dormancy strategies, as
probability of germination is determined by 27 haploid genes d_i, with i between 1 and 27,
each corresponding to a week in the dry half of the year, and the 27^{th} indicating germination at
the start of the wet season. Each gene determines the probability of a seed germinating in the
matching week, and therefore has values bounded by 0 and 1, with the condition that the sum
of all genes always equals 1. The actual phenotypic trait value (dormancy period d_z) of an
offspring j is randomly drawn from the probability distribution determined by its parent's
genetic values.

Population initialization. Each replicate simulation starts with N_0 adult individuals. Each of
these individuals are independently assigned a random combination of gene values, where
starting d_i values are uniformly distributed between 0 and 1. After assigning each gene a
random value, we standardize all values such that the sum of an individual's 27 genes equals
1. We start the simulations with ample standing genetic variation for computational purposes.

Abiotic conditions. We model time steps of one week and assign the first 26 weeks of a year
as the dry season and the last 26 weeks as the wet season. Dry-season rainfall varies
stochastically within years, with 26 weekly rainfalls randomly drawn at the start of the year
from a beta distribution bounded between 0 (no rain) and 1 (high rain). Weekly rainfall in the
wet season is always equal to 1. We parameterize the beta distribution in terms of its mode m,
such that the shape parameters are given by

$$\alpha = m + 1 \tag{9}$$

$$\beta = 2 - m.$$

The mode mainly affects the average rainfall (the higher the wetter), but note that it also
affects weekly rainfall variance within years $V \left(= \frac{1}{36}\frac{m+1}{2-m}\right)$, which is highest when $m=0.5$.

To tease apart effects of bet-hedging resulting from the spatiotemporal scale of environmental
stochasticity, we examine evolutionary outcomes under coarse-grained and fine-grained
rainfall variation. In coarse-grained environments, we assume that all seeds experience the
same sequence of weekly dry-season rainfalls R (a vector of 26 values) but R differs among
years. Under fine-grained environmental variation, different R vectors are assigned to
individual seeds within years at random. Specifically, we produce 1000 sequences of weekly
rainfall upfront and assign each seed a random R chosen among these. Thus, experienced
rainfall will vary a lot among individuals within a year in this scenario, but average rainfall
will vary less across years. In both scenarios, the weekly rainfall in the wet season (week 27–
52) is constant and equal to 1, which represents high rainfall.

Survival of seedlings and adults. After setting the weekly rainfalls of a particular year, we
determine the survival probability of seedlings and adults in that year. We assume that
survival increases both with rainfall and seedling age. The probability of survival for an
individual i with age $a_{i,t}$ in a given week t with rainfall R_t (or $R_{i,t}$ in the fine-grained scenario)
equals

$$S_{i,t}\left(a_{i,t}, R_t\right) = S_{\min}R_t + \frac{(S_{\max}-S_{\min}R_t)}{1+\exp(-\delta_1(a_{i,t}-a_{\mathrm{infl}}(1-R_t\delta_2)))}, \tag{10}$$

i.e. a sigmoid function of seedling age, the lower asymptote of which increases with
increasing rainfall (Fig. 2A, B). Note that rainfall also affects the horizontal position of the
function (how far "along" the x-axis the seed has come in terms of survival), and that the
strength of this effect increases with the parameter δ_2.

With the rainfall data of a particular year and the survival function (eq. 10), we can now
calculate each seedling's survival probability (Fig. 2C), which is given by the product of its
(age- and rain-dependent) survival probabilities of each week from germination time d_z until
the end of the year:

$$S_i(d_z, \mathbf{R}) = \prod_{t=d_z}^{52} S_{i,t}(a_{i,t}, R_t)$$

To determine whether a seedling survives its first year, we draw a random number from a
uniform distribution [0,1]. If this random number is lower than the survival probability of the
seedling, the seedling will survive, otherwise it is removed from the population. Likewise, we
can calculate which of the adults will be allowed to reproduce (see below) and survive to the
next year. We assume that weekly survival for plants of age $a_i \geq 27$ weeks, is independent of
age and rainfall and equal to s_{ad}, such that yearly survival becomes s_{ad}^{52} and expected lifespan
(in weeks) is $1/(1-s_{ad})$.

316
Fig. 2: Seedling survival functions. A and B: Weekly survival probability for different amounts of rainfall (blue
lines: $\mathbf{R}$=1; red lines: $\mathbf{R}$=0) and varying the inflection point a_{inf} of the juvenile survival function $S_{i,t}$ (panels). C:
Effect of week of germination (x-axis) and a_{inf} (colored lines) on the probability of a seedling surviving its first
year (52 weeks since germination), assuming a worst-case scenario with no rainfall ($\mathbf{R}$=0) in the dry season and
constant high rainfall ($\mathbf{R}$=1) in the wet season.

Recruitment and density dependence. We assume that only K adult plants can live in the
environment. At the end of the wet season, we determine which of the seedlings that
germinated in this year will recruit into the adult population. We first calculate the number of
available slots (K minus the number of surviving adults), and then fill the empty slots with
seedlings. Recruitment success depends on seedling age, with older seedlings having a higher
probability of recruiting. The relative recruitment probability of a seed is calculated as

329
$$(a_i - 25)^c, \tag{11}$$

where parameter c indicates the strength of competitive asymmetry. For $c = 0$, recruitment is
independent of age, while for high values of c, young seedlings will be outcompeted by older
seedlings. Because seeds germinate at the latest in week 27, the youngest seedling is 26 weeks
old at the time of recruitment. By subtracting 25 from the age of each seedling, the youngest
seedling has a weight of 1, and all other seedlings have weights exceeding 1.

Reproduction, inheritance and mutation. At the end of the wet season, all adult plants (i.e.
individuals older than 52 weeks, not the juveniles that were just recruited) reproduce
asexually. Individual seed production is Poisson distributed with a mean fecundity of γ.
Offspring inherit their parent's gene values for determining dormancy duration (d_i), but not
their actual phenotype d_z. Each locus has an independent probability μ of experiencing
mutation. Mutational effects are of constant effect, such that the new gene value is drawn
from a normal distribution with a mean equal to the previous value, and a standard deviation
of μ_σ. Mutations that would lead to negative values are absorbed at 0, ensuring that the new
gene value is non-negative. After mutation, the values of all genes of an individual are
normalized to sum to 1.

Table 1: Overview of variables and parameters of the individual-based model.

Parameter in text	Description	Default values and range
A) Abiotic environment parameters		
m	Mode of the beta distribution determining weekly dry-season rainfall (see eq. 9)	Default 0.5, we explore [0, 1]
v	Variance of the beta distribution	$\dfrac{1}{36}\dfrac{m+1}{2-m}$
R_t	Realized rainfall in a particular week t.	[0, 1].
B) Survival function parameters		
a	Age of seedling.	
S_{min}	Lower asymptote of the seedling survival function (survival probability given $a=1$ and $R_t=0$).	Default 0.6.
s_{max}	Upper asymptote of the seedling survival function (survival probability when a and R_t are large).	Default 0.997.
s_{ad}	Weekly survival probability of plants older than 26 weeks	Default 0.997, we also examine $s_{ad} = 0.993$
δ_1	Affects the steepness of the survival function.	Default 0.2.
δ_2	Enhances the effect of rain on the survival function.	Default 0.8.
a_{inf}	Age (weeks) at which the survival function increases most steeply.	Default 5. We also examine $a_{inf} = \{4, 7\}$.
C) Genes and trait values		
d_i	Genetic values determining weekly probabilities of breaking dormancy, $i \in \{1, 2, \ldots, 27\}$.	[0, 1]. Require $\Sigma_i d_i = 1$.
d_z	Realized individual dormancy period (in weeks).	
D) Demographic and evolutionary parameters		
K	Carrying capacity.	Default 5000.
γ	Yearly fecundity per adult plant.	Default 50, we also examine $\gamma = 130$
μ	Per-locus mutation rate.	Default 0.1.
μ_σ	Size of mutational effects (The standard deviation of the Gaussian distribution from which the new trait value is chosen).	Default 0.001.
N_0	Starting population size.	Default equal to K.
c	Strength of competitive asymmetries (competitive advantage to germinating earlier), see eq. 11.	Default 1. We also examine $c = \{0.5, 1.2\}$.

347   *Data generation and analysis*. Simulations run for 1.000.000 years. Results presented are
348   based on the populations at the end of each simulation. As there is hardly variation among
349   replicates, results are presented for only 3 independent replicate runs of each scenario. We
350   implemented the model in C++; the code will be made available online. Simulations were run
351   using GNU parallel (Tange 2021).

Results

Despite the added ecological complexity and evolutionary realism of the individual-based
simulation model, with its potential for a detailed flexible germination strategy, we overall
find that these results correspond well with those of the analytical model (section 1). Firstly,
despite the high potential for finely-tuned germination strategies, our simulations showed
remarkably little variation both among replicated simulations and within populations, in
agreement with our prediction that any ESS is convergence stable and so no evolutionary
branching should occur. Secondly, our simulations highlight the mechanisms modulating the
(rainfall-dependent) benefits and costs of early relative to late seeds – namely the strength of
age-dependent competitive asymmetries for recruitment and juvenile dry-season survival –
and corroborate the finding from our analytical model that mixed germination strategies only
arise at a certain balance point where the relative benefits of early seeds are neither too large
nor too small (Figs. 3, 4).

Additionally, expanding on the findings from the analytical model, our simulations
demonstrate that this within-individual dormancy variation evolves as an adaptation lowering
genotype-level fitness variance, confirming that such strategies do indeed represent bet-
hedging strategies (compare top vs. bottom rows of Figs. 3, 4). By comparing results obtained
in temporally coarse-grained environments (all individuals experience the same rainfall
patterns in a given year, resulting in high within-year fitness correlations and high among-
year fitness variation) with those obtained in fine-grained environments (rainfall patterns are
assigned independently for all individuals in a given year, resulting in low within-year fitness
correlations and low among-year fitness variation), we observe that any spreading of
germination disappears in fine-grained environments. Thus, the evolution of within-individual
seed size variation in perennials can here be attributed to the interaction between age-
dependent survival, competition for recruitment, and bet-hedging. This result also importantly
highlights that bet-hedging in germination strategies is not driven *per se* by weekly rainfall
variation within years, *V*, but also by the spatial scale of rainfall variation, which affects
correlations in environmental conditions experienced among individuals within populations.

When competitive asymmetries are weak, i.e. competition for recruitment only depends
weakly on seedling age, obligate dormancy evolves causing germination at the beginning of
the wet season (top row of Fig. 3a and S1a). This holds even under higher rainfall levels, for
which dry-season germination and seedling survival may often be possible, because there are
no competitive benefits to early germination and so even small risks are not worth taking.
Conversely, when competitive ability depends strongly on seedling age, minimal dormancy
periods evolve and seeds germinate as soon as possible (top row of Fig. 3c and S1c). These
seedlings then need to survive the entire risky dry-season, but this pays off as long as the dry
season is not too malign, because later-germinating seeds have little chance of recruiting if
there are even a few surviving early seeds. Accordingly, levels of within-individual variation
in dormancy period are low in both these scenarios. Bet-hedging has no effect on the
evolution of dormancy in this case, because there is no variance in expected fitness nor fitness

correlations among related individuals (top and bottom rows are qualitatively identical in Fig. 3a,c and S1a,c). Such fixed strategies and absence of evolutionary conflicts of interest in the short and long term are also predicted by our analytical model (eq. 8, no mixed strategies ($0 < f < 1$) can arise if the benefits b of germinating early do not increase sufficiently despite excess rain).

Figure 3: Evolved trait values (weekly germination probabilities) in three independent simulations (box colors) under increasing levels of competitive advantage to older seeds ("competitive asymmetries", columns). For clarity, results are shown summing the germination probabilities of 9 weeks together. Box plots show median (horizontal line), range of 50 % of variation (box limits), range of 95 % of variation (vertical lines) and any outliers (black points). Top row: Coarse-grained environment (all seedlings experience same rainfall patterns within years). Bottom row: Fine-grained environment (all seedlings experience different rainfall patterns within years). Strength of competitive asymmetry is determined by parameter c, which takes the values of 0.5 (left column), 1 (middle column) or 1.2 (right column). Other parameters as in table 1.

Similarly, only intermediate mortality risks select for mixed strategies. Because seedling mortality strongly affects the risks of early germination for a given rainfall level, any variation in germination timing occurs at intermediate levels of mortality risk in the dry season (Fig. 4). In the scenario shown in Fig. 4 we adjust the parameter a_{inf}, the inflection point of the survival function, i.e. the age at which survival rises steepest, such that higher (later) a_{inf} entails higher overall mortality (see Fig. 2). We find that low mortality (left column in Fig. 4) always selects for germinating as soon as possible (because of age-dependent competitive asymmetries). On the other hand, very high mortality (due to high a_{inf}; right column in Fig. 4) always selects for germinating as close as possible to the start of the wet season. Only intermediate mortality risks (Figs. 4 & S2, middle column) favor variable dormancy periods. Again, this general result corresponds well with our analytical model (Fig. 1).

Figure 4: Evolved trait values (weekly germination probabilities) in three independent simulations (box colors) under increasing mortality risks (columns). For clarity, results are shown summing the germination probabilities of 9 weeks together. Box plots show median (horizontal line), range of 50 % of variation (box limits), range of 95 % of variation (vertical lines) and any outliers (black points). Top row: Coarse-grained environment (all seedlings experience same rainfall patterns within years). Bottom row: Fine-grained environment (all seedlings experience different rainfall patterns within years). Mortality risk is modulated by adjusting the a_{inf} parameter, which takes the values 4 (left column), 5 (middle column) and 7 (right column).

A final observation in this model is that, in contrast with predictions from previous bet-hedging theory, shorter lifespans may select for less spreading of germination (Fig. 5, weekly results shown in Fig. S3). This effect arises because of our implementation of density-dependence, where lower adult survival means that more 'slots' are opened up every year for seedlings to recruit into. This decreases the intensity of age-dependent competition, leading to weaker selection for early germination the lower the adult survival (Fig. 5A shows results for high adult survival, versus Fig. 5B for low adult survival).

In situations where competition is strongly asymmetric (high value of *c*, bottom row of Fig. 5), lowering adult survival may select for more spreading of germination (Fig. 5D, shows results for high adult survival, versus Fig 5E for low adult survival). The reduction in density-dependence now intensifies selection for late germination, resulting in the evolution of more spread in germination timing. In the analytical model, adult survival affects the speed of evolution only, and not the evolutionary endpoint (equation 4). This discrepancy with the individual-based model is caused by the assumption in the analytical model that the strength of density-dependence depends on the relative benefit of germinating early and the frequency of this strategy. In the individual-based model, however, the strength of density-dependence is

also affected by the number of free spots and competing seedlings. If we reduce adult survival
and increase the average seed production at the same time, such that the strength of density-
dependence (expected number of seeds competing per open slot) remains the same, we no
longer see an effect of adult survival on the resulting strategies (Fig 5C and 5F show the same
pattern of uni- or bimodal germination as do Fig 5A and 5D).

As for Figs. 3 & 4, removing any effects of bet-hedging by implementing fine-grained rainfall
patterns (and thus removing environmentally-induced fitness correlations among related
individuals) leads to populations with only a single, fixed strategy in all scenarios depicted in
Fig. 5 (results not shown). Thus, within-individual variation in germination timing can arise
due to interactions between life-history traits, bet-hedging and intensity of intraspecific
competition.

Figure 5: Evolved trait values (weekly germination probabilities) in three independent simulations (box colors)
under different strengths of density dependence (columns) and levels of competitive asymmetry (rows). For
clarity, results are shown summing the germination probabilities of 9 weeks together. Box plots show median
(horizontal line), range of 50 % of variation (box limits), range of 95 % of variation (vertical lines) and any
outliers (black points). Top row: Intermediate competitive asymmetry ($c = 1$), bottom row: strong competitive
asymmetry ($c = 1.2$). Density-dependence is modulated by adjusting the adult survival rate s_a (0.997 in the left
panels, 0.993 in the middle and right panels) and the average seed production (50 in the left and middle panels,
130 in the right panels).

Discussion

Modelling results and insights

Our two complementary modelling approaches have shown how stochastic seasonal
environments can select for within-individual variation in germination behavior through
interactions between bet-hedging, life-history traits, and competitive asymmetries due to age-
dependent seedling recruitment. First, our analytical model considering discrete variation

(early- vs. late-germinating seeds) reveals ample scope for mixed germination strategies in
long-lived species. This model revealed that the key determinant of whether variable
germination evolves is the shape of the function relating how all benefits and costs of early vs
late seeds change depending on dry-season rainfall environmental conditions. Specifically,
both the risks of early germination if conditions are too dry before the beginning of the wet
season, and the benefits provided by competitive advantages over later-germinating seedlings
(who will be smaller at the time of competition for recruitment), are captured in such a
function, which is shown to have to be increasing and accelerating around 0 (expected dry-
season rainfall), but not too quickly, in order for a mixed strategy to evolve.

Inspired by these results, we next built an evolutionary simulation model of dormancy period,
where we included both these benefits and costs to allow them to vary independently, as well
as continuous within- and among-individual variation in seed dormancy. To allow
unconstrained evolution of flexible germination strategies, we modelled the joint evolution of
27 unlinked genes representing weekly germination probabilities of offspring. Despite this
flexibility, the results from our analytical model with only two discrete germination strategies
were largely recovered: Mixed strategies emerged only when the potential benefits
(competitive advantage) and costs (dry-season mortality) of earlier germination were neither
too strong nor too weak, but were still observed for a large range of parameter values.
Furthermore, intermediate dormancy durations emerged very rarely, with evolution rather
favoring mixed parental strategies producing some early- and some late-germinating
offspring. Through comparisons with evolutionary outcomes in fine-grained environments,
which never exhibited such mixed strategies, we conclude that this represents a diversifying
bet-hedging strategy evolving due to genotype-level fitness correlations. In coarse-grained
environments, phenotypically variable offspring ensure that fitnesses of siblings become
decoupled, thus ensuring low genotype-level variance in fitness (Starrfelt and Kokko 2012).

Our work demonstrates that explicit consideration of density-dependence is necessary for
understanding the adaptive value of within-individual variation in offspring phenotype in
long-lived organisms. Variation in offspring size within individuals is not only common in
perennial plants, but also in many other iteroparous organisms including arthropods (Fox and
Czesak 2000), birds (Amundsen and Slagsvold 1998), fish (Einum and Fleming 2004), and
marine invertebrates (Marshall and Keough 2007). While among-population variation is often
interpreted as adaptive, within-individual variation, on the other hand, is mostly viewed as
maladaptive (Fox and Czesak 2000). This interpretation is possibly driven by the focus on
optimality models (e.g. Smith and Fretwell 1974; Einum and Fleming 2004), which by
definition predict a single offspring size to maximize fitness (Metz et al. 2008). Approaches
that take frequency-dependent interactions into account, however, show that within-individual
variation in offspring size can be adaptive, even in the absence of environmental
unpredictability (e.g. Geritz 1995). Our work shows that such an ecological approach is also
necessary to understand evolution in unpredictable environments.

Germination strategies in seasonal environments

509
510   Our models were inspired by the wet-dry seasonal dynamics characterizing large parts of the
511   tropics (Feng et al. 2013). Yearly patterns of seedling emergence such as those described by
512   Garwood (1983) for Barro Colorado Island in Panamá suggest that germination during the
513   early part of the rainy season predominates, although there is considerable variation (see also
514   Escobar et al. 2018). Garwood's study also hinted at an important role of asymmetric

competition in that seedling emergence in highly competitive light gaps tended to occur earlier than did emergence in less competitive understory habitats. Our models provide a mechanistic explanation for these patterns, and also suggest that ongoing changes in the predictability and temporal variation in precipitation patterns in the tropics may select for altered germination strategies in tropical plants (Feng et al. 2013).

The adaptive challenges faced by plants of seasonal tropical forests are in some ways analogous to those of arctic-alpine plants facing the problem of 'false springs' followed by freezing events, which is thought to select for dormancy or other mechanisms to avoid detrimental fitness effects of premature germination (Schwienbacher et al. 2011; Mondoni et al. 2012). Arguing that within-individual variation in germination timing could be beneficial as a bet-hedging trait, Simons & Johnston (1997) used a model of such a system to illustrate their argument that developmental instability could be adaptive in annuals. During a time when developmental instability was largely seen as maladaptive (c.f. the debate surrounding fluctuating asymmetry), substantial work on bet-hedging strategies has largely resolved this controversy (Zhang and Hill 2005; Devaux and Lande 2010; Scheiner 2014; Tufto 2015), and the dynamics considered by Simons & Johnson (1997) are now a canonical example of bet-hedging in seasonal phenology.

A second source of inspiration for our models was the extensive variation in seed size often observed in angiosperms (Michaels et al. 1988), and the many studies reporting relationships between seed size and germination behavior (e.g. Simons and Johnston 2000; Norden et al. 2009; Martins et al. 2019). Although seed size is sometimes thought about as a highly canalized trait (because there should be an optimal balance between size and number of offspring; Smith and Fretwell 1974), both categorical (seed heteromorphism) and continuous variation in seed size is common (e.g. Susko and Lovett-Doust 2000). In a recent phylogenetic analysis, Scholl et al. (2020) analyzed the occurrence of seed heteromorphism in a large sample of the flora of southwestern North America, and detected weak associations between seed heteromorphism and several measures of environmental predictability (aridity and precipitation seasonality). While the tendency for seed heteromorphism to be more likely in more unpredictable environments is consistent with our simulation model, these authors explicitly excluded species with continuous variation in seed size from their analyses. Although our simulation model suggests that seed heteromorphism may be a more likely evolutionary outcome for dealing with environmental variability than continuous seed size variation, there is no reason why continuous seed size variation could not also represent a bet-hedging strategy in unpredictable environments (c.f. Simons & Johnston's (1997) suggestion of developmental instability causing adaptive variation in germination timing). Interestingly, Scholl et al. (2020) failed to detect a correlation between seed heteromorphism and the annual life cycle, contrary to their expectations from previous bet-hedging theory. Although seed heteromorphism was more common in annuals than in perennials, they attributed the lack of a detectable effect to the small overall proportion of seed-heteromorphic species. However, the common occurrence of seed heteromorphism also among perennials can further be seen as an indication that the processes interacting with bet-hedging in our model to produce variation in germination phenology may indeed be common and important in natural systems.

Empirical guidelines and predictions

Our models yield several testable predictions about the relationship between germination behavior, seed size, and environmental predictability. The simplest prediction arising from

our results is that within-individual variability in germination behavior and associated traits
(e.g. seed size) can be adaptive, and possibly more so in more unpredictable environments.
However, we also find that the distinctions between fine- vs. coarse-grained environmental
variation is important in driving germination patterns. This implies that caution is needed
when interpreting measurements of environmental variation and using them to predict
ecological gradients of germination strategies. For example, the measurement of rainfall
variation used in Scholl et al. (2020) was based on among-year variation (coefficient of
variation of annual rainfall measurements) and the smallest spatial grid used was 25 km^2,
which is too large to capture whether a given genotype may experience environmental
variation as fine- or coarse-grained. Thus, neither of these metrics provide information about
the different outcomes observed in our models, which might explain the lack of a relationship
between rainfall variation and the occurrence of seed heteromorphism. Another reason
highlighted by both our models could be that mixed germination strategies as adaptations to
unpredictable environments may only evolve under certain conditions of how fitness costs
and benefits vary among morphs across environments.

Recent studies of variation in seed size and germination behavior in the tropical vine
Dalechampia scandens have revealed predictable covariation between measures of
environmental predictability and germination behavior. Martins et al. (2019) found that seeds
from populations occupying more seasonal environments required longer periods of after-
ripening (duration of storage required before germination occurs in response to favorable wet
conditions) before germinating, and that within each population, smaller seeds required
shorter periods of after-ripening. In a follow-up study, Pélabon et al. (2021) considered
variation in seed size within individuals and detected a positive relationship between
environmental stochasticity and variation in seed size. Environmental unpredictability as
defined by Pélabon et al. more closely resembles our parameter *V* in describing stochastic
variation in rainfall patterns, and our model thus provides mechanistic support for the
hypothesis that the patterns of seed size variation and germination behavior in *Dalechampia*
represents a bet-hedging strategy in unpredictable environments.

While observational studies of germination timing and seed size across ecological gradients
may continue to yield insights into variation in these traits, experimental studies would be
highly valuable. Seed-sowing experiments with seed families of known descent would
specifically allow separating environmental and genetic causes of variation, and lead to
insights into the evolutionary potential of germination behavior (e.g. Simons & Johnson
2006). However, separating the presence of two distinct strategies from continuous variation
may continue to prove difficult, especially if the difference in mean behavior between
strategies is small, and variation around these means is large.

Limitations and caveats of the model

Our model results are broadly consistent with empirical patterns of seed size and germination
behavior, yet some caution is needed when interpreting the model output. In particular, the
specific genetic architecture we use (one freely evolving locus for each weekly germination
probability) is admittedly highly unrealistic. Although we hope that this flexibility has
allowed us to observe outcomes that explore all parts of unconstrained strategy space, some
potentially beneficial combinations of germination probabilities may have been difficult to
arrive at. For example, by linearly standardizing all loci of weekly germination probabilities

to sum to 1, loci with already large values could make the effect of potentially beneficial
mutations at other loci increasing the germination probability in other weeks very small.

Next, while this breach in realism aimed to limit bias and increase the robustness of our
evolutionary interpretation, other limitations may in fact introduce biases to our conclusions.
For example, it is difficult to say how our choice of asexual inheritance might have affected
the ability to evolve discrete vs. continuous strategies, or indeed the range of parameter space
favoring variable germination, and so including sexual reproduction could provide an
interesting next step for expanding our modelling work. Plant mating systems are highly
variable, and near-complete selfing is uncommon compared to mixed and predominantly
outcrossing systems (Moeller et al. 2017). If germination timing is indeed highly polygenic,
outcrossing among individuals producing predominantly early- and predominantly late-
germinating seeds might allow mixed strategies to evolve more easily than in our model,
where such mixed strategies instead are dependent on the right mutations occurring
sequentially. In addition to dormancy, germination timing is also driven by variation in
flowering and fruiting phenology as well as the duration of fruit maturation (Escobar et al.
2018). Although we chose to keep these processes fixed in our model, variation in flowering
time additionally leads to non-random mating and is therefore an interesting trait to add if
future studies consider sexual reproduction.

Modelling multilocus evolution in the absence of any constraints, as we do here, may in itself
introduce biases if evolution of optimal combinations is difficult to attain due to genetic
constrains, covariation with other traits, or trade-offs. Importantly, we did not incorporate a
trade-off between offspring quality and dormancy period, which could be expected if we
assume that seed size is the mechanism by which variation in germination strategy is achieved
(Stearns 1989). Furthermore, we did not include a cost to seed size variation (or canalization,
see Zhang and Hill 2005). However, while such additional assumptions might affect our
results (for example by narrowing the parameter space in which variable strategies are
selected for if adding a cost to variation, and widening it if adding a cost to canalization), they
also reduce generality and do not subtract from the general point. Variation in germination
strategies may arise from other sources than simply seed size variation resulting from per-
offspring parental investment (Simons and Johnston 2006; Baskin and Baskin 2014) and as
with costs of variation vs. canalization, we note that the pattern may also go either way with
regards to the seed size vs. germination strategy (offspring quantity vs quality) trade-off: In
many species, larger seeds germinate earlier (Biere 1991; Simons and Johnston 2000; Pélabon
et al. 2005), but the opposite pattern is also observed (Susko and Lovett-Doust 2000), and
among species time to germination is usually longer for larger seeds (Norden et al. 2009;
Harel et al. 2011). Thus, while adding some trade-off between seed set and germination
strategy might increase realism with respect to certain systems, generality would again
necessarily be lowered relative to our present model without such energetic constraints.

Conclusions
This study has demonstrated novel mechanisms by which variation in germination strategies
in perennial plants can evolve, shedding light on the prevalence of seed-size variation seen in
perennials as well as annuals. Seed size was historically considered a highly canalized trait, to
the point that the weight measurement 'carat' was based on the remarkably invariant seeds of
the carob tree (*Ceratonia siliqua*). However, it has since been discovered that substantial
variation in seed size is widespread, even in carob seeds (Turnbull et al. 2006), and its

universality begs an explanation. Bet-hedging in desert annuals has previously been the best-
studied empirical and theoretical example of variable germination strategies, but there are
arguably more perennial species occupying seasonal tropical or temperate environments than
there are desert annuals. Our modelling results, where bet-hedging in perennials interacts with
unpredictable seasonal variation and competitive asymmetries to produce within-individual
variation in germination strategies, therefore provide a much-needed extension to existing
theory on this topic. A better understanding of competitive dynamics in stochastic seasonal
environments can help improve the predictability of species and community responses to
ongoing changes in climate patterns and other anthropogenic challenges.

Acknowledgments:

The authors thank Hanna Kokko for helpful discussions, and an anonymous reviewer for a
great contribution to the analytical model. HtB and TRH were supported by the Swiss
National Science Foundation, on grant no. 310030B_182836 awarded to Hanna Kokko. TRH
was additionally supported by the Forschungskredit of the University of Zurich, grant no. FK-
21-122 awarded to TRH, and the Research Council of Norway, grant no. 313570 awarded to
Jane Reid.

Author contributions

666 All authors initiated the study. HtB and TRH developed and analyzed the models. HtB ran the
667 simulations. ØHO reviewed the empirical literature. TRH reviewed the theoretical literature
668 and drafted the manuscript with input from all authors.

Appendix A: Derivation of invasion fitness (eq. 2)

Following a standard adaptive dynamics approach (Geritz et al. 1998), we assume sequential
appearance of rare mutations of small phenotypic effect in populations of residents fixed for
the wild type, where each mutation is assumed to go to fixation (if its invasion fitness is
positive) before the next mutation occurs. Mutants are assumed to have negligible effect on
ecological dynamics, such that the number of mutants N' in a population of K residents can be
derived. As in the main text, we designate respectively the resident and mutant germination
strategies (fraction of seeds early) as f and f', and probability of recruitment for an early seed
is given by eq. 1. Then, the dynamics of number of mutants $N'(t + 1)$ is given by the number
of surviving mutant adults from the previous time step, $N'(t)s_a$, plus any mutant seedling
recruitment (eq. A1). As described in the main text, seedling recruitment is according to a
lottery model where all seeds compete for a limited number of spaces in the adult population
freed up after adult mortality; this number is $(1 - s_a)K$. Assuming equal yields of γ seeds per
mutant and wild type adult, the number of mutant seeds produced is $N'(t)\gamma$, and the number
of resident seeds is $K\gamma$, and their mean (averaged over small and large seeds) recruitment
probabilities are respectively $f'b + 1 - f'$ and $fb + 1 - f$. The number of mutants in the next
time step is given by

$$686 \qquad N'(t + 1) = N'(t)s_a + (1 - s_a)K \, \frac{N'(t)\gamma(f'b+1-f')}{K\gamma(fb+1-f)}. \qquad (A1)$$

687 Cancelling away $K\gamma$ and dividing by $N'(t)$ gives the growth rate λ of the mutant population
688 used in eq. 2:

689
$$\lambda = \frac{N\prime(t+1)}{N\prime(t)} = s_a + (1 - s_a) \, \frac{f\prime b + 1 - f\prime}{f b + 1 - f}. \tag{A2}$$

Data availability statement

691
692 All code and output is available upon request and will be uploaded to Dryad upon journal
693 acceptance.

References

Amundsen, T., and T. Slagsvold. 1998. Hatching asynchrony in great rits: A bet-hedging strategy?
Ecology 79:295–304.

Bårdsen, B.-J., P. Fauchald, T. Tveraa, K. Langeland, N. G. Yoccoz, and R. A. Ims. 2008. Experimental
evidence of a risk-sensitive reproductive allocation in a long-lived mammal. Ecology 89:829–
837.

Baskin, C. C., and J. M. Baskin. 2014. Seeds. Ecology, Biogeography, and Evolution of Dormancy and
Germination. 2nd ed. Academic Press, San Diego, US.

Bednekoff, P. A. 1996. Risk-sensitive foraging, fitness, and life histories: Where does reproduction fit
into the big picture? Integr. Comp. Biol. 36:471–483.

Biere, A. 1991. Parental effects in *Lychnis flos-cuculi*. I: Seed size, germination and seedling
performance in a controlled environment. J. Evol. Biol. 4:447–465.

Brändel, M. 2004. Dormancy and germination of heteromorphic achenes of *Bidens frondosa*. Flora
199:228–233.

Chesson, P. 2000. Mechanisms of maintenance of species diversity. Annu. Rev. Ecol. Syst. 31:343–
366.

Cohen, D. 1966. Optimizing reproduction in a randomly varying environment. J. Theor. Biol. 12:119–
129.

Crowley, P. H., S. M. Ehlman, E. Korn, and A. Sih. 2016. Dealing with stochastic environmental
variation in space and time: bet hedging by generalist, specialist, and diversified strategies.
Theor. Ecol. 9:149–161.

Devaux, C., and R. Lande. 2010. Selection on variance in flowering time within and among individuals.
Evolution (N. Y). 64:1311–1320.

Donohue, K., R. Rubio De Casas, L. Burghardt, K. Kovach, and C. G. Willis. 2010. Germination,
postgermination adaptation, and species ecological ranges. Annu. Rev. Ecol. Evol. Syst. 41:293–
319.

Einum, S., and I. A. Fleming. 2004. Environmental unpredictability and offspring size: Conservative
versus diversified bet-hedging. Evol. Ecol. Res. 6:443–455.

Ellner, S. 1985. ESS germination strategies in randomly varying environments. I. Logistic-type models.
Theor. Popul. Biol. 28:50–79.

Escobar, D. F. E., R. Rubio de Casas, and L. P. C. Morellato. 2021. Many roads to success: different
combinations of life-history traits provide accurate germination timing in seasonally dry
environments. Oikos 1–15.

Escobar, D. F. E., F. A. O. Silveira, and L. P. C. Morellato. 2018. Timing of seed dispersal and seed

dormancy in Brazilian savanna: Two solutions to face seasonality. Ann. Bot. 121:1197–1209.

Feng, X., A. Porporato, and I. Rodriguez-Iturbe. 2013. Changes in rainfall seasonality in the tropics.
Nat. Clim. Chang. 3:811–815. Nature Publishing Group.

Finch-Savage, W. E., and G. Leubner-Metzger. 2006. Seed dormancy and the control of germination.
New Phytol. 171:501–523.

Fox, C. W., and M. E. Czesak. 2000. Evolutionary ecology of progeny size in arthropods. Annu. Rev.
Entomol. 45:341–369.

Franch-Gras, L., E. M. García-Roger, M. Serra, and M. J. Carmona. 2017. Adaptation in response to
environmental unpredictability. Proc. R. Soc. B Biol. Sci. 284.

Fumanal, B., B. Chauvel, A. Sabatier, and F. Bretagnolle. 2007. Variability and cryptic heteromorphism
of *Ambrosia artemisiifolia* seeds: What consequences for its invasion in France? Ann. Bot.
100:305–313.

Furness, A. I., K. Lee, and D. N. Reznick. 2015. Adaptation in a variable environment: Phenotypic
plasticity and bet-hedging during egg diapause and hatching in an annual killifish. Evolution (N.
Y). 69:1461–1475.

Galloway, L. F. 2001. The effect of maternal and paternal environments on seed characters in the
herbaceous plant Campanula americana (Campanulaceae). Am. J. Bot. 88:832–840.

García-Roger, E. M., M. J. Carmona, and M. Serra. 2016. Modes, mechanisms and evidence of bet
hedging in rotifer diapause traits. Hydrobiologia, doi: 10.1007/s10750-016-2869-5.

Garwood, N. C. 1983. Seed Germination in a Seasonal Tropical Forest in Panama. Ecol. Monogr.
53:159–181.

Gerber, N., and H. Kokko. 2018. Abandoning the ship using sex, dispersal or dormancy: multiple
escape routes from challenging conditions. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 373.

Geritz, S. A. H. 1995. Evolutionarily stable seed polymorphism and small-scale spatial variation in
seedling density. Am. Nat. 146:685–707.

Geritz, S. A. H., É. Kisdi, G. Meszéna, and J. A. J. Metz. 1998. Evolutionarily singular strategies and the
adaptive growth and branching of the evolutionary tree. Evol. Ecol. 12:35–57.

Geritz, S., M. Gyllenberg, and J. Toivonen. 2018. Adaptive correlations between seed size and
germination time. J. Math. Biol. 77:1943–1968. Springer Berlin Heidelberg.

Gourbière, S., and F. Menu. 2009. Adaptive dynamics of dormancy duration variability: Evolutionary
trade-off and priority effect lead to suboptimal adaptation. Evolution (N. Y). 63:1879–1892.

Gremer, J. R., and D. L. Venable. 2014. Bet hedging in desert winter annual plants: Optimal
germination strategies in a variable environment. Ecol. Lett. 17:380–387.

Haaland, T. R., J. Wright, and I. I. Ratikainen. 2019. Bet-hedging across generations can affect the
evolution of variance-sensitive strategies within generations. Proc. R. Soc. B Biol. Sci. 286. Royal
Society Publishing.

Haaland, T. R., J. Wright, and I. I. Ratikainen. 2020. Generalists versus specialists in fluctuating
environments: a bet-hedging perspective. Oikos 1–12.

Harel, D., C. Holzapfel, and M. Sternberg. 2011. Seed mass and dormancy of annual plant populations
and communities decreases with aridity and rainfall predictability. Basic Appl. Ecol. 12:674–684.
Elsevier GmbH.

Hughes, P. W. 2018. Minimal-risk seed heteromorphism: Proportions of seed morphs for optimal
risk-averse heteromorphic strategies. Front. Plant Sci. 9:1–9.

Imbert, E. 2002. Ecological consequences and ontogeny of seed heteromorphism. Perspect. Plant
Ecol. Evol. Syst. 5:13–36.

Kisdi, É., and G. Meszéna. 1995. Life histories with lottery competition in a stochastic environment:
ESSs which do not prevail. Theor. Popul. Biol. 47:191–211.

Kortessis, N., and P. Chesson. 2019. Germination variation facilitates the evolution of seed dormancy
when coupled with seedling competition. Theor. Popul. Biol. 130:60–73. Elsevier Inc.

Levins, R. 1962. Theory of fitness in a heterogeneous environment. I. The fitness set and adaptive
function. Am. Nat. 96:361–373.

Marshall, D. J., and M. J. Keough. 2007. The evolutionary ecology of offspring size in marine
invertebrates. Adv. Mar. Biol. 53:1–60.

Martins, A. A., Ø. H. Opedal, W. S. Armbruster, and C. Pélabon. 2019. Rainfall seasonality predicts the
germination behavior of a tropical dry-forest vine. Ecol. Evol. ece3.5108.

Metcalf, C. J. E., L. T. Burghardt, and D. N. Koons. 2015. Avoiding the crowds: The evolution of plastic
responses to seasonal cues in a density-dependent world. J. Ecol. 103:819–828.

Metz, J. A. J., S. D. Mylius, and O. Diekmann. 2008. When does evolution optimize? Evol. Ecol. Res.
10:629–654.

Meyer, S. E., S. G. Kitchen, and S. L. Carlson. 1995. Seed germination timing patterns in intermountain
Penstemon (Scrophulariaceae). Am. J. Bot. 82:377–389.

Moeller, D. A., R. D. Briscoe Runquist, A. M. Moe, M. A. Geber, C. Goodwillie, P. O. Cheptou, C. G.
Eckert, E. Elle, M. O. Johnston, S. Kalisz, R. H. Ree, R. D. Sargent, M. Vallejo-Marin, and A. A.
Winn. 2017. Global biogeography of mating system variation in seed plants. Ecol. Lett. 20:375–
384.

Mondoni, A., G. Rossi, S. Orsenigo, and R. J. Probert. 2012. Climate warming could shift the timing of
seed germination in alpine plants. Ann. Bot. 110:155–164.

Norden, N., M. I. Daws, C. Antoine, M. A. Gonzalez, N. C. Garwood, and J. Chave. 2009. The
relationship between seed mass and mean time to germination for 1037 tree species across five
tropical forests. Funct. Ecol. 23:203–210.

Olofsson, H., J. Ripa, and N. Jonzén. 2009. Bet-hedging as an evolutionary game: the trade-off
between egg size and number. Proc. R. Soc. B Biol. Sci. 276:2963–2969.

Pélabon, C., M. L. Carlson, T. E. Hansen, and W. S. Armbruster. 2005. Effects of crossing distance on
offspring fitness and developmental stability in *Dalechampia scandens* (Euphorbiaceae). Am. J.
Bot. 92:842–851.

Pélabon, C., F. De Giorgi, Ø. H. Opedal, G. H. Bolstad, A. Raunsgard, and W. S. Armbruster. 2021. Is
there more to within-plant variation in seed size than developmental noise? Evol. Biol., doi:
10.1007/s11692-021-09544-y. Springer US.

Pinceel, T., B. Vanschoenwinkel, W. Hawinkel, K. Tuytens, and L. Brendonck. 2017. Aridity promotes
bet hedging via delayed hatching: a case study with two temporary pond crustaceans along a
latitudinal gradient. Oecologia 184:161–170. Springer Berlin Heidelberg.

Poethke, H. J., T. Hovestadt, and O. Mitesser. 2016. The evolution of optimal emergence times: bet
hedging and the quest for an ideal free temporal distribution of individuals. Oikos 125:1647–

1656.

Rees, M. 1996. Evolutionary ecology of seed dormancy and seed size. Philos. Trans. R. Soc. B Biol. Sci.
351:1299–1308.

Rees, M., D. Z. Childs, K. E. Rose, and P. J. Grubb. 2004. Evolution of size-dependent flowering in a
variable environment: Partitioning the effects of fluctuating selection. Proc. R. Soc. B Biol. Sci.
271:471–475.

Rubio de Casas, R., C. G. Willis, W. D. Pearse, C. C. Baskin, J. M. Baskin, and J. Cavender-Bares. 2017.
Global biogeography of seed dormancy is determined by seasonality and seed size: a case study
in the legumes. New Phytol. 214:1527–1536.

Scheiner, S. M. 2014. Bet-hedging as a complex interaction among developmental instability,
environmental heterogeneity, dispersal, and life-history strategy. Ecol. Evol. 4:505–515.

Scholl, J. P., L. Calle, N. Miller, and D. L. Venable. 2020. Offspring polymorphism and bet hedging: a
large-scale, phylogenetic analysis. Ecol. Lett. 23:1223–1231.

Schwienbacher, E., J. A. Navarro-Cano, G. Neuner, and B. Erschbamer. 2011. Seed dormancy in alpine
species. Flora Morphol. Distrib. Funct. Ecol. Plants 206:845–856. Elsevier GmbH.

Seger, J., and H. J. Brockmann. 1987. What is bet-hedging? Oxford Surv. Evol. Biol. 4:182–211.

Simons, A. M. 2011. Modes of response to environmental change and the elusive empirical evidence
for bet hedging. Proc. R. Soc. B Biol. Sci. 278:1601–1609.

Simons, A. M. 2014. Playing smart vs. playing safe: The joint expression of phenotypic plasticity and
potential bet hedging across and within thermal environments. J. Evol. Biol. 27:1047–1056.

Simons, A. M., and M. O. Johnston. 1997. Developmental instability as a bet-hedging strategy. Oikos
80:401–406.

Simons, A. M., and M. O. Johnston. 2006. Environmental and genetic sources of diversification in the
timing of seed germination: Implications for the evolution of bet hedging. Evolution (N. Y).
60:2280.

Simons, A. M., and M. O. Johnston. 2000. Variation in seed traits of *Lobelia inflata* (Campanulaceae):
Sources and fitness consequences. Am. J. Bot. 87:124–132.

Smith, C. C., and S. D. Fretwell. 1974. The optimal balance between size and number of offspring. Am.
Nat. 108:499–506.

Snyder, R. E. 2006. Multiple risk reduction mechanisms: Can dormancy substitute for dispersal? Ecol.
Lett. 9:1106–1114.

Starrfelt, J., and H. Kokko. 2012. Bet-hedging - a triple trade-off between means, variances and
correlations. Biol. Rev. 87:742–755.

Stearns, S. C. 1989. Trade-offs in life-history evolution. Funct. Ecol. 3:259–268.

Susko, D. J., and L. Lovett-Doust. 2000. Patterns of seed mass variation and their effects on seedling
traits in *Alliaria petiolata* (Brassicaceae). Am. J. Bot. 87:56–66.

Tange, O. 2021. GNU Parallel. Zenodo.

ten Brink, H., J. R. Gremer, and H. Kokko. 2020. Optimal germination timing in unpredictable
environments: the importance of dormancy for both among- and within-season variation. Ecol.
Lett., doi: 10.1111/ele.13461.

Torres-Martínez, L., P. Weldy, M. Levy, and N. C. Emery. 2017. Spatiotemporal heterogeneity in
precipitation patterns explain population-level germination strategies in an edaphic specialist.
Ann. Bot. 119:253–265.

Tremayne, M. A., and A. J. Richards. 2000. Seed weight and seed number affect subsequent fitness in
outcrossing and selfing *Primula* species. New Phytol. 148:127–142.

Tufto, J. 2015. Genetic evolution, plasticity, and bet-hedging as adaptive responses to temporally
autocorrelated fluctuating selection: A quantitative genetic model. Evolution (N. Y). 69:2034–
2049.

Turnbull, L. A., L. Santamaria, T. Martorell, J. Rallo, and A. Hector. 2006. Seed size variability: From
carob to carats. Biol. Lett. 2:397–400.

Varpe, Ø. 2017. Life history adaptations to seasonality. Integr. Comp. Biol. 57:943–960.

Venable, D. L. 2007. Bet hedging in a guild of desert annuals. Ecology 88:1086–1090.

Venable, D. L. 1985. The evolutionary ecology of seed heteromorphism. Am. Nat. 126:577–595.

Venable, D. L., and J. S. Brown. 1988. The selective interactions of dispersal, dormancy, and seed size
as adaptations for reducing risk in variable environments. Am. Nat. 131:360–384.

Venable, D. L., A. Búrquez, G. Corral, and E. Morales. 1987. The ecology of seed heteromorphism in
Heterosperma pinnatum in central Mexico. Ecology 68:65–76.

Venable, D. L., and L. Lawlor. 1980. Delayed germination and dispersal in desert annuals: Escape in
space and time. Oecologia 46:272–282.

Vleeshouwers, L. M., H. J. Bouwmeester, and C. M. Karssen. 1995. Redefining seed dormancy: An
attempt to integrate physiology and ecology. J. Ecol. 83:1031.

Wagmann, K., N. C. Hautekèete, Y. Piquot, C. Meunier, S. E. Schmitt, and H. Van Dijk. 2012. Seed
dormancy distribution: Explanatory ecological factors. Ann. Bot. 110:1205–1219.

Wang, C.-C., and D. C. Rogers. 2018. Bet hedging in stochastic habitats: an approach through large
branchiopods in a temporary wetland. Oecologia 188:1081–1093. Springer Berlin Heidelberg.

Wang, H. L., L. Wang, C. Y. Tian, and Z. Y. Huang. 2012. Germination dimorphism in *Suaeda*
acuminata: A new combination of dormancy types for heteromorphic seeds. South African J.
Bot. 78:270–275. Elsevier B.V.

Yao, S., H. Lan, and F. Zhang. 2010. Variation of seed heteromorphism in *Chenopodium album* and
the effect of salinity stress on the descendants. Ann. Bot. 105:1015–1025.

Zhang, X. S., and W. G. Hill. 2005. Evolution of the environmental component of the phenotypic
variance: Stabilizing selection in changing environments and the cost of homogeneity. Evolution
(N. Y). 59:1237–1244.

Supplementary material

Figure S1: As Figure 3, except with each weekly germination probability plotted individually rather than grouped by 9 weeks at the time. The dashed horizontal line indicates the probability 1/27, which would be the expected evolved value in the absence of selection on germination week.

Figure S2: As Figure 4, except with each weekly germination probability plotted individually rather than grouped by 9 weeks at the time. The dashed horizontal line indicates the probability 1/27, which would be the expected evolved value in the absence of selection on germination week.

899 **Figure S3:** As Figure 5, except with each weekly germination probability plotted individually rather than
900 grouped by 9 weeks at the time. The dashed horizontal line indicates the probability 1/27, which would be the
901 expected evolved value in the absence of selection on germination week.

903