Evidence of increased Cathepsin B/L and decreased TMPRSS2 usage for cell entry by the SARS-CoV-2 Omicron variant

Pranesh Padmanabhan[1,*], Narendra M. Dixit[2,3,*]

[1]Clem Jones Centre for Ageing Dementia Research, Queensland Brain Institute, The University of Queensland, Brisbane, Australia 4072

[2]Department of Chemical Engineering, Indian Institute of Science, Bangalore, India 560012

[3]Centre for Biosystems Science and Engineering, Indian Institute of Science, Bangalore, India 560012

[*]**Correspondence:**

Pranesh Padmanabhan, Narendra M. Dixit

Email: p.padmanabhan@uq.edu.au; narendra@iisc.ac.in

Letter details:

Main Text: ~1200 words; Figures: 1; References: 10

The SARS-CoV-2 Omicron variant appears to exhibit altered cell tropism and entry properties compared to other variants. Here, analysing recent data, we found a strong positive correlation between the entry efficiency of the Omicron variant and the relative usage of the host protease Cathepsin B/L by the original SARS-CoV-2 strain for cell entry. We developed a mathematical model to quantify entry efficiency in *in vitro* assays and found that the Omicron variant displayed >4-fold improved efficiency in using Cathepsin B/L for entry in 293T-ACE2 cells compared to the original strain. The preferential usage of Cathepsin B/L over TMPRSS2 for entry may explain the altered cell tropism of the Omicron variant and have implications for our understanding of its infectivity and transmissibility as well as for interventions.

The new SARS-CoV-2 variant Omicron (B.1.1.529), declared a variant of concern on 25th November 2021, is fast becoming the dominant strain in several countries. It harbours as many as 37 mutations in the spike protein compared to the original SARS-CoV-2 strain [1] and is able to evade neutralising antibodies generated by previously infected or vaccinated individuals [1-3], possibly leading to its increased transmissibility and rapid global spread. Emerging data indicates that the high number of mutations in the Omicron variant affects its viral entry properties and cell tropism [2-5], which in turn may influence its ability to establish infection post-exposure and the severity of the subsequent symptoms. In this letter, we analyse this new data and find that the Omicron variant has substantially altered preferences for cell entry pathways compared to the earlier variants. Further, using a mathematical model, we quantify the extent of the differential usage of the entry pathways by the Omicron variant.

Early studies showed that the original SARS-CoV-2 (Wuhan-Hu-1) strain displayed broad cell tropism, with viral entry a key determinant of the tropism both *in vitro* [6] and *in vivo* [7]. Intriguingly, multiple recent *in vitro* studies suggest that the Omicron variant's cell

tropism may be altered and its entry efficiency different from the original and other variants in

a cell line-dependent manner [2-5]. For instance, SARS-CoV-2 pseudotyped virus bearing

either the B.1 or the Delta variant spike protein showed higher entry efficiency than the

Omicron pseudotyped virus in Caco-2 (human, colon) and Calu-3 (human, lung) cells, whereas

the Omicron pseudotyped virus entered Vero (African green monkey, kidney) and 293T

(human, kidney) cells more efficiently than the other variants [3]. Similar trends were observed

in live SARS-CoV-2 virus infection assays [4, 5]; the Delta variant infection spread was

significantly greater than the Omicron variant in Calu-3 cells [4, 5], whereas the spread of the

two variants was similar in VeroE6 cells [4]. What causes the relative efficiency of Omicron

virus entry to be higher in some cells and lower in others than other variants?

The first step in SARS-CoV-2 entry into target cells is the binding of the viral spike

protein, S, with the host cell surface receptor angiotensin-converting enzyme 2 (ACE2). Cell

tropism is thus expected to be affected by ACE2 expression levels [6-8]. The Omicron spike

protein binds soluble human ACE2 strongly [1, 3]. Further, ACE2 is necessary for the entry of

all SARS-CoV-2 variants [2-6]. Thus, any variation in ACE2 expression across cell types is

likely to have similar effects on the entry efficiencies of all the variants. The cell type-

dependent variation in the entry efficiency of the Omicron variant is therefore unlikely to arise

from the variations in the ACE2 expression level across cell types. Omicron spike protein

incorporation into pseudotyped virus appears to be compromised compared to the Delta and

Wuhan D614G strains [5]. However, this reduction in spike protein density is likely to decrease

the Omicron entry efficiency across all cell types, thus ruling it out as a potential cause of the

differential entry efficiency observed.

Following ACE2 engagement, successful entry requires that the spike protein be

cleaved into subunits by host proteases, either by the transmembrane serine protease TMPRSS2

at the plasma membrane or the cysteine proteases Cathepsin B and Cathepsin L in the

endosomal vesicles. The proteases TMPRSS2 and Cathepsin B/L are thought to work

independently, facilitating SARS-CoV-2 entry through two independent pathways [8, 9] (Fig.

1A). We recently analysed SARS-CoV-2 pseudotyped virus infection of several cell lines [6]

using a mathematical model of SARS-CoV-2 entry and found that the relative usage of

TMPRSS2 and Cathepsin B/L entry pathways by the original SARS-CoV-2 strain varied

widely across cell lines [9]. For example, Vero cells predominantly admitted entry through the

Cathepsin B/L pathway, whereas Calu-3 cells allowed entry via the TMPRSS2 pathway. Vero

cells overexpressing TMPRSS2 permitted entry via both pathways. Importantly, the original

strain displayed the ability to use either pathway, with the preferred pathway possibly based on

the relative expression levels of the two proteases [8]. We reasoned that the Omicron variant

might use these pathways differently from the original (or the Delta) strain, possibly underlying

its different cell tropism.

To test this, we extended the above analysis to the recent experiments of Hoffmann *et*

al. [3] and found a correlation between the Omicron virus entry efficiency and the relative

usage of the two entry pathways by the original strain (Fig. 1B). With cell lines (Calu-3 and

Caco-2) where the usage of the TMPRSS2 entry pathway by the original strain was dominant,

the Omicron pseudotyped virus entry was significantly less efficient than the B.1 and delta

strains. By contrast, the Omicron virus entry was more efficient in cell lines (Calu-3 and Caco-

2) where the Cathepsin B/L entry pathway usage by the original strain was dominant (Fig. 1B).

One possible interpretation of this relationship is that the Omicron variant entry is relatively

less effective through the TMPRSS2 pathway and more effective via the Cathepsin B/L

pathway than the other strains. Consistent with this notion, camostat mesylate, a TMPRSS2

inhibitor, was less potent against the Omicron variant than the Delta variant in blocking live

virus infection of VeroE6 cells overexpressing TMPRSS2, which allows entry via both

pathways [4]. Moreover, syncytium formation in a cell-cell fusion assay, which requires

TMPRSS2 but not Cathepsins, was severely impaired for the Omicron variant compared to

both the Delta [4, 5] and the Wuhan-Hu-1 D614G [5] strains. The increased efficiency of

Cathepsin B/L usage together with the reduced efficiency of TMPRSS2 usage by the Omicron

variant may explain its altered cell tropism relative to the original strain or other variants.

To quantify the increased efficiency of the Cathepsin B/L pathway usage by the

Omicron variant, we developed a mathematical model and applied it to analyse data from

Garcia-Beltran *et al*. [2], who measured the fraction of 293T-ACE2 cells infected by

pseudotyped virus bearing the wild-type or the Omicron spike proteins at different viral

concentrations (Fig. 1C). We derived an analytical expression to describe infections in the

pseudotyped virus infection assay (Supplementary Information, Text S1). Accordingly, the

fraction of cells infected, f_a, is given by $f_a = 1 - e^{\frac{k\overline{S_{tc}}}{c_V}(e^{-c_V t_d}-1)V_0}$. Here, V_0 is the initial viral

concentration, and $\overline{S_{tc}}$ is the mean susceptibility of the cell population to virus entry,

accounting for the usage of both the pathways. (See Text S1 for the description of other model

parameters.) Because 293T-ACE2 cells predominantly permit original strain entry via the

Cathepsin B/L pathway [9] and given the relatively poor efficiency of TMPRSS2 usage by the

Omicron variant (Fig. 1B), we let $\overline{S_{tc}} = \overline{S_c}$, where $\overline{S_c}$ is the mean susceptibility through the

Cathepsin B/L pathway. Our model provided good fits to the data (Fig. 1C) and estimated ~4.6-

fold increased usage of the Cathepsin pathway by the Omicron variant in 293T-ACE2 cells

relative to the original strain (more details in Text S1).

The switch in the Omicron variant's preference for entry pathways may have clinical

implications. Drugs targeting TMPRSS2, such as camostat mesylate [6], and Cathepsin B/L

pathway, such as hydroxychloroquine [10], are in clinical development. The effectiveness of

TMPRSS2 inhibitors is likely to decrease and that of Cathepsin B/L inhibitors is likely to

increase against the Omicron strain. Furthermore, our mathematical model previously

predicted synergy between TMPRSS2 and Cathepsin B/L inhibitors against SARS-CoV-2

infection [9]. Because the Omicron variant's preference for the TMPRSS2 pathway is reduced but not eliminated, combination treatment with TMPRSS2 and Cathepsin B/L inhibitors may exhibit limited but possibly useful synergy against Omicron virus infection of cells that admit both entry pathways. Future studies may establish these implications quantitatively. We suggest that pseudotyped virus infection experiments at different viral concentrations together with inhibitors of TMPRSS2 and Cathepsin B/L pathways, analysed as described here and in Padmanabhan *et al*. [9], might help quantify the relative TMPRSS2 and Cathepsin B/L pathway usage by different strains across cell types including Omicron, potentially explaining their cell tropism and the efficacy of the above treatments.

In summary, our analyses support the notion that the Omicron variant has shifted to better usage of the Cathepsin B/L entry pathway at the expense of the TMPRSS2 pathway. It would be interesting to see how these changes in the entry pathway usage observed in *in vitro* conditions relate to disease severity and transmissibility in animal models and humans.

ACKNOWLEDGEMENTS

We thank Rajat Desikan for help with Figure 1A.

COMPETING INTERESTS

The authors declare that no conflicts of interests exist.

REFERENCES

1. Cameroni E, Bowen JE, Rosen LE, Saliba C, Zepeda SK, Kulap K, et al. Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic shift. Nature. 2021.

2. Garcia-Beltran WF, Denis KJS, Hoelzemer A, Lam EC, Nitido AD, Sheehan ML, et al. mRNA-based COVID-19 vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant. Cell. 2021.

3. Hoffmann M, et al. The Omicron variant is highly resistant against antibody-mediated
neutralization – implications for control of the COVID-19 pandemic. 2021.

4. Zhao H, Lu L, Peng Z, Chen LL, Meng X, Zhang C, et al. SARS-CoV-2 Omicron variant
shows less efficient replication and fusion activity when compared with delta variant in
TMPRSS2-expressed cells. Emerg Microbes Infect. 2021;1-18.

5. Meng B, Ferreira IATM, Abdullahi A, Saito A, Kimura I, al. e. SARS-CoV-2 Omicron spike
mediated immune escape, infectivity and cell-cell fusion. bioRxiv. 2021.

6. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-
CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven
protease inhibitor. Cell. 2020;181:271-80.e8.

7. Liu J, Li Y, Liu Q, Yao Q, Wang X, Zhang H, et al. SARS-CoV-2 cell tropism and
multiorgan infection. Cell Discov. 2021;7:17.

8. Koch J, Uckeley ZM, Doldan P, Stanifer M, Boulant S, Lozach PY. TMPRSS2 expression
dictates the entry route used by SARS-CoV-2 to infect host cells. EMBO J. 2021;40:e107821.

9. Padmanabhan P, Desikan R, Dixit NM. Targeting TMPRSS2 and Cathepsin B/L together
may be synergistic against SARS-CoV-2 infection. PLoS Comput Biol. 2020;16:e1008461.

10. Ou T, Mou H, Zhang L, Ojha A, Choe H, Farzan M. Hydroxychloroquine-mediated
inhibition of SARS-CoV-2 entry is attenuated by TMPRSS2. PLoS Pathog. 2021;17:e1009212.
164

Figure 1. Omicron variant entry efficiency relative to B.1 and Delta strains depends on the relative usage of the TMPRSS2 and Cathepsin B/L pathways. (A) Schematic of the two independent entry pathways accessible to SARS-CoV-2. **(B)** The Omicron entry efficiency relative to either the B.1 (blue) or the Delta (red) variant across cell types. Data was taken from Hoffmann *et al*. [3]. *Inset*, the relative usage of the entry pathways by the original strain. Data was taken from Padmanabhan *et al*. [9]. **(C)** Fits of model predictions (lines) to the

172   experimental data (filled circles) taken from Garcia-Beltran *et al*. [2] measuring the fraction of

173   cells infected by pseudotyped virus bearing either the wild-type or the Omicron spike proteins.

174   In B and C, experimental data was extracted using Engauge Digitizer 12.1. In C, Data was fit

175   using the tool NLINFIT in MATLAB R2017b.