1

Contrast-enhanced microCT evaluation of degeneration following partial and full width injuries to mouse lumbar intervertebral disc

+Remy E. Walk, MS[1,2]; +Hong Joo Moon, MD, PhD[2,3]; Simon Y. Tang, PhD, MSCI[1,2*]

Munish C. Gupta, MD[2]

[1] Dept of Biomedical Engineering, Washington University in St. Louis, St. Louis, MO, USA
[2] Dept of Orthopaedic Surgery, Washington University in St. Louis, St. Louis, MO, USA
[3] Dept of Neurosurgery, Korea University College of Medicine, Seoul, South Korea

+ These authors contributed equally.

Keywords: Lumbosacral surgical exposure; Intervertebral disc degeneration; annulus fibrosus injury; mouse model; contrast-enhanced microCT

Abstract

Study Design: Preclinical animal study

Objective: Evaluation of the degenerative progression resulting from either a partial- or full- width

injury to the mouse lumbar intervertebral disc (IVD) using contrast-enhanced micro-computed

tomography and histological analyses. We utilized a lateral-retroperitoneal surgical approach to access

the lumbar IVD, and the injuries to the IVD were induced by either incising one side of the annulus

fibrosus or puncturing both sides of the annulus fibrosus. The full-width injury caused dramatic reduction

in nucleus pulposus hydration and significant degeneration. A partial-width injury produces localized

deterioration around the annulus fibrosus that resulted in local tissue remodeling without gross

degeneration to the IVD.

Methods: Female C57BL/6J mice of 3-4 months age were used in this study. They were divided into

three groups to undergo partial-width, full-width, or sham injuries. The L5/L6 and L6/S1 lumbar IVDs

were surgically exposed using a lateral-retroperitoneal approach. The L6/S1 IVDs were injured using

either a surgical scalpel (partial-width) or a 33G needle (full-width), with the L5/L6 serving as an internal

control. These animals were allowed to recover and then sacrificed at 2-, 4-, or 8- weeks post-surgery.

The IVDs were assessed for degeneration using contrast-enhanced microCT (CEμCT) and histological

analysis.

Results: The high-resolution 3D evaluation of the IVD confirmed that the respective injuries localized

within one side of the annulus fibrosus or spanned the full width of the IVD. The full-width injury caused

deteriorations in the nucleus pulposus after 2 weeks and progressed to significant degeneration at 8 weeks,

while the partial width injury caused localized disruptions that remained limited to the annulus fibrosus.

Conclusion: The use of CEμCT revealed distinct IVD degeneration profiles resulting from partial- and

full- width injuries. The partial width injury may serve as an alternative for IVD degeneration resulting

from localized annulus fibrosus injuries in humans.

Introduction

Mouse models are often used for the preclinical validation of treatments and therapeutic

candidates. Mice offer particular advantages of easy maintenance, year-round breeding, short gestation

period, large litters, and inbred tolerance[1]. Furthermore, the ability to manipulate the mouse genome

enables mechanistic studies with greater biological precision than larger mammalian models[1]. Moreover,

the mouse lumbar intervertebral disc (IVD) provides close geometric and microstructural semblance to

the human lumbar IVD compared to other preclinical animal models[2]. While the mouse lumbar IVD

exhibits age-related degeneration[3,4], an injury is often utilized to reproduce the inflammatory conditions

and accelerate the IVD's degenerative cascade[5,6].

To create a targeted injury to the IVD, surgical exposure of the IVD is required. These

commonly involve the posterior-lateral, transperitoneal, and anterior/lateral retroperitoneal surgical

access. Of these, the retroperitoneal approach particularly advantageous in that it : (1) does minimal

damage to major organs, vessels, and musculature; (2) accesses multiple spinal levels with a relatively

small incision; and (3) enables direct visualization of the target tissue under microscopy. Moss et al.

demonstrated the efficacy and reproducibility of the retroperitoneal approach in rabbits [7]. Exposing the

lumbar IVD enables the ability to create a targeted, directed injury for the investigation of the subsequent

degenerative process [8–10]. Masuda et al. and Sobajima et al. described the rabbit annulus fibrosus injury

model where the injury was confined to the annulus fibrosus that caused slow progressive degeneration of

the IVD over 8 weeks. The phenotype and the progressive nature of this model recapitulates the human

disease, and may be better suited to explore therapies that leverage prevention or regeneration.   In

contrast, more damaging approaches that injures both the annulus fibrosus and nucleus pulposus produce

rapid and severe course of degeneration [7,11]. There are several studies describing IVD degeneration using

the lumbosacral IVD injury in the mouse model[12–14]. In the mouse IVD where the average disc height is

approximately 300-400 microns, an injury by needle puncture produces damage to 55-90% across the

height and 15-40% across the width of the IVD[2,11], representing significant trauma that is atypical in

human IVDs. Moreover, these procedures involve damage and injury to the nucleus pulposus, such as

with a complete puncture to the IVD, with no possibility to decouple the contributions of the annulus

fibrosus and the nucleus pulposus toward the ensuing degenerative cascade. Piazza and coauthors have

shown that unilateral (half-width) and bilateral (full-width) injuries in the tail IVD results in unique

degenerative trajectories, but this has not been investigated in the mouse lumbar spine. We thus sought to

compare the degenerative profiles of IVDs after partial- and full- width injuries in the mouse lumbar

spine. In order to evaluate the degeneration of the IVD in a spatially robust manner, we utilized contrast-

enhanced microCT[15,16], in addition to histological grading, to quantify the changes in structure and

composition at 2-, 4- and 8- weeks after surgery.

Materials and Methods

Animal Preparation

All procedures were performed following Washington University School of Medicine IACUC

approval. Female C57BL/6J mice of 3-4 months age were used (BW: 20 – 25 g). They were housed

under standard animal husbandry conditions (in a temperature-controlled [21 ± 1°C] room with normal

12-hr light/dark cycles). These animals were divided into three groups: Partial-width injury (PW), full-

width injury (FW), and Sham (n=15-18 per group) with all animals undergoing to retroperitoneal surgical

exposure of the lumbar IVD. The groups were cross-sectionally evaluated at 2-, 4-, and 8- week post-

surgery time points. The injury was delivered to the L6/S1 IVD with the L5/6 IVD used as the internal

control.

The partial-width injury mimics a localized injury to the annulus fibrosus, aka annular tear injury, in

humans[17]. To evaluate the feasibility of this injury, we first performed the partial-width injury on six

animals, and then they were euthanized shortly after recovery from anesthesia. The IVDs were harvested

and measured for the thickness of the annulus fibrosus and the depth of the injury using contrast-enhanced

microCT (CEμCT) and histological analysis.

The second set of animals were allowed to recover following PW, FW or Sham surgery for 2, 4 or 8

weeks (n = 5-6 per group) and then euthanized, and the IVD tissues were assessed for degeneration with

CEμCT and histological analyses.

Retroperitoneal approach to the intervertebral disc procedure

Mice were anesthetized with isoflurane gas in oxygen via a facemask (3–4% induction and 2–2.5%

maintenance at 1 L/min flow rate; Highland Medical Equipment) and were given a preoperative

intradermal injection of lidocaine (7 mg/kg; Hospira, Inc). The left flank was then shaved from the ventral

to the dorsal midlines, and the skin was sterilized. The skin was prepared for aseptic surgery via washing

with 70% ethanol and povidone iodine. Under microscopic guidance, retroperitoneal dissection was used

to expose the lateral aspect of the spine with a Penfield dissector. The pelvis could be rotated posteriorly

to expose a broad working space while holding the pelvic bone (Figure 2A). The Penfield dissector was

pulled out the peritoneal wall to expose the psoas muscle and to protect the abdominal organs (Figure

2B). The psoas muscle was stripped posteriorly to anteriorly by initially scraping out tissues attached to

the anterior surface of the pelvis (Figure 2C). The spinal column and the intervertebral disc were exposed

from the ventral midline to the posterior margin of the disc at the L5/6 and L6/S1 levels (Figure 2D).

Since the superior margin of the pelvic bone indicates the L6 vertebral body (Figure 2E), the location of

the L5/6 and L6/S1 IVD could be confirmed (Figure 2D and 2E). Extension to the cranial space beyond

the L5/6 was also easily achieved by blunt dissection. Additional details and images on the surgical

procedure can be found in Supplemental methods, Supplemental Figure S1, and Supplemental video.

Partial-width and full-width injuries

For the partial-width injury, a distance of 0.3 mm from the end of the No. 11-scalpel blade tip was

measured and marked with a micro-caliper under a microscope (Figure 1). The distance of 0.3 mm was

determined from our preliminary studies. The edge of the blade was allowed to insert into the IVD until

the 0.3 mm marking was no long visible under the microscope. The injury site was closely observed

under the microscope to confirm that there is no leakage of the NP.

For the full-width injury, a 33G needle was inserted bilaterally through the IVD lateral axis of the

IVD. In contrast to the partial-width injury, NP herniations were observed following the full-width

injury.

Contrast-enhanced microCT tomography (CEµCT)

Samples were incubated in a solution of 175 mg/mL solution diluted from a stock of OptiRay 350

(Guerbet, St. Louis) in PBS at 37°C. After 24 hrs of incubation, samples were scanned using a µCT40

(Scanco Medical, CH) at a 10-µm voxel size (45 kVp, 177 uA, high resolution, 300 ms integration).

CEμCT data was exported as a DICOM file for analysis in a custom MATLAB program. After an

initial median filter (sigma = 0.8, support = 5), functional spine units were isolated from surrounding soft

tissue not part of the IVD by drawing a contour around the outer edge every 5 transverse slices and

morphing using linear interpolation. The IVD was manually segmented from the vertebral bodies with the

same methodology as above. The remaining voxels were designated as the whole disc mask. The NP was

thresholded from the AF followed by a morphological close and morphological open to fill interior holes

and smooth the boundary. The volumes and average attenuations (intensity) were calculated from the

mask of the NP and whole disc. The volume was determined from the total number of voxels contained

within the mask and the attenuation was taken as the average 16-bit grayscale value of the voxels.

Visualizations of the microCT were obtained using the image processing application OsiriX (Pixmeo,

Geneva). AF thickness, partial-width injury depth and disc height index (DHI) were measured along the

mid-sagittal plane. DHI was calculated as the ratio of the IVD height to width. IVD height was taken as

the average at 5 equidistant points along the mid-sagittal plane. The ratio of NP intensity/disc intensity

(NI/DI), defined as the average attenuation of voxels in the NP mask divided by the full disc mask, is an

unbiased, fully three-dimensional measure that quantifies the relative size and hydration to inform the

relative changes in degeneration [18]. Samples where attenuation was saturated were not included in

analysis.

Histological preparation and evaluation

Following microCT, samples were fixed for 24 hours in 10% neutral buffered formalin followed by

3 days of decalcification in Immunocal (StatLab 1414-X). The samples were embedded in paraffin,

sectioned at a thickness of 10 µm, and then stained with Safranin-O and Fast Green. Histological

classification system recently developed by Melgoza et al. in 2021 was used to quantify the degeneration

of the injured level (L6/S1) which allows the evaluation of the nucleus pulposus, annulus fibrosus,

endplates and interface boundaries[19]. Morphology and NI/DI determined from CEμCT was used to further

inform the level of degeneration of the injured level (L6/S1) compared to the internal control (L5/6).

Measuring thickness of AF and the depth of partial-width injury

The CEμCT of the injured L6/S1 and histological analysis on uninjured L5/6 were used to measure

the thickness of the AF. The CEμCT on the injured L6/S1 was used to measure the depth of injury as

defined by the shortest perpendicular distance from the outer edge of annulus fibrosus to the visually

observable outline of the injury site.

Statistics

All statistics were examined for normality and nonparametric tests were used accordingly. A 2-

way ANOVA was used to assess the effects of injury and postsurgical time on histological grade,

attenuation and morphological parameters, with adjusted post hoc comparisons between groups (Tukey).

Comparisons and effects were considered significant when the p-value is less than 0.05 . All statistics

were run using GraphPad Prism 9.3.1 (GraphPad, San Diego, California).

Results

A total of 57 mice were subjected to the surgical procedure. The average surgery time was 15 min 38 sec

± 6 min 23 sec from incision to closure. There was no mortality or complications due to surgery. Caution

was taken to prevent injury of the lumbosacral plexus located posteriorly during the blunt dissection of

the psoas muscle from posterior to the anterior direction.

Annulus Fibrosus (AF) Thickness

CEµCT and histology measurements of AF thickness and IVD width were highly consistent.   The

anteroposterior IVD width measured with CEµCT was 1.27 ± 0.13 mm (mean ± standard deviation),

whereas that measured by histological analysis was 1.21 ± 0.11 mm.   AF thickness measured with

CEµCT was 0.38 ± 0.05 mm and that measured with histological analysis was 0.43 ± 0.10 mm (Figure 3).

The CEµCT measured values were statistically indistinguishable from ($p = 0.37$) and were highly

correlated ($r^2 = 0.96$) with the histologically measured values (Figure 2).

Depth of Partial-width Injury

All injuries except one were isolated to the AF (Figure 3B). In sample 5, the NP may have been injured

because CEµCT showed disruption of the AF/NP boundary (Figure 3C). The depth of injury measured

with CEµCT was 0.29 ± 0.05 mm. The ratio of the injury depth to the AF thickness was 0.80 ± 0.19

(Figure 3D and 3E).

Structural evaluation of the IVD

Neither the partial-width nor the full-width injuries resulted in changes in the Disc Height Index

across the two-, four-, and eight- week time points (Figure 4D). Similarly, the size of the nucleus

pulposus, characterized by the nucleus pulposus (NP) volume fraction, were not dramatically different

between groups and time points (Figure 4E). The relative hydration of the NP, quantified by the ratio of

NP attenuating intensity to that of the whole disc (NI/DI), revealed the full-width injury caused a dramatic

decrease in the attenuation of the NP, indicating a loss of NP hydration, at the two- and four- week time

points, with a trending change at eight- weeks.

The impact of the partial- and full- width injuries to the annulus fibrosus was observable at two

weeks, The relative hydration of the annulus fibrosus (AF) increases after an injury, and this is evident in

both the partial- and full- width groups. Two weeks after injury, both the partial- and full- injuries

showed an increase attenuation of the AF, with the full-width injury group sustaining this increase (Figure

4G).

Extent of IVD Degeneration

Quantitative histological analyses revealed a degenerative cascade to the IVD following full-

width injury as early as two weeks post injury but not with the partial-width injury. The histological

classification showed significant degeneration following FW injury at all timepoints ($p < 0.05$) in the NP,

AF, interface boundaries, and total IVD score but not endplate score compared to Sham while no

differences were detected between PW and Sham (Figure 5). No morphological differences between

timepoints or injury groups were observed.

Discussion

Animal models of IVD degeneration are imperative to elucidate the key molecular mechanisms of

the pathophysiology.   To date, rodents, rabbits, dogs, goats, sheep, and primates have been used as

models for intervertebral disc degeneration[20]. One main advantage of mouse models is the availability of

reagents and modifications that could be used concomitantly with surgically induced IVD degeneration.

However, the small size of the mouse requires a high degree of surgical precision, particularly with access

and exposure of the lumbar spine.   While many studies utilize the mouse tail for degenerative models[12,21–

23], the lumbar spine maintains the anatomical proximity to physiologically relevant structures such as the

dorsal root ganglions.

We describe here a novel procedure for mouse lumbosacral IVD injury with visual guidance via

microscopy and gross features. The retroperitoneal space of the spine can be located (Figure S1) with the

left pelvic bone with gluteus muscles and proximal thigh as landmarks (Figure 1C and 1D). Blunt-

dissection of the fat pad surrounding the thoracic cage anteriorly, the gluteus muscles superiorly, and the

thigh muscles posteriorly (Figure 1B) exposes these two landmarks (Figure 1C). The lateral approach to

the IVD allows for precise injury to the annulus fibrosus of the IVD (Figure 1). Exposure of the pelvic

bone from the gluteus muscle and rotation of the pelvic bone posteriorly is necessary to achieve ample

exposure of multiple IVDs in the surgical field. The psoas muscles can be easily scraped posterior-

anteriorly with a Penfield dissector. CEμCT allowed for a 3D spatially unbiased quantification of

morphology and composition[18] that confirmed the localization of injury across the cross-section time

points. CEμCT measurements of the IVD morphology were in excellent agreement with histological

measures. NI/DI and histological analysis indicated degeneration of injured IVDs starting 2 weeks post-

surgery.

The present study describes a surgical procedure to expose the lumbosacral spine of mice and to

induce a localized experimental injury with AF-limited depth and compared the degenerative response

with a more commonly used needle puncture injury. The technique described herein also focused on

achieving a consistent depth and size of injury. The marked tip of the No. 11-scalpel blade tip was

confirmed with CEμCT to provide AF-limited injury and resulted in no NP leakage under observation by

microscopy. While most injury models cause damage to the nucleus pulposus, the cartilaginous endplates,

and even the vertebral body[24–26], the partial-width injury here results in an isolated injury to the outer

annulus fibrosus. Surprisingly, we observed no degenerative changes in the endplate in the full-width

injury, suggesting that it is possible to induce an IVD only injury using a carefully applied 33G needle.

Following full-width injury, IVDs exhibited rapid degeneration as early as two weeks after injury, and

this is sustained through the four- and eight- week time points. Despite maintaining disc height, the FW

injury caused consistent degenerative changes in the nucleus pulposus, annulus fibrosus, and interface

boundaries which included the AF-endplate and NP-AF boundaries. The PW injury did not produce

statistically significant degenerative changes, but whether the innervation or vascularization changes at

the outer annulus fibrosus remains to be investigated.

CEμCT allowed for nondestructive visualization of the injury site following both partial- and full-

width injuries. The increased attenuation on CEμCT may indicate changes in composition and diffusion

properties resulting from the disruption in the annulus fibrosus with both injuries. The injured annulus

fibrosus appears to be highly hydrated, likely due to the increased interstitial fluids that localize around

the injury site (Figures 3-4). Since IVD degeneration in humans often starts as a tear in the annulus

fibrosus[17], this model will enable mechanistic investigations of how injuries of the annulus fibrosus

contribute to IVD degeneration and the subsequent development of low back pain. In contrast, the full-

width injury to the nucleus pulposus induces a more rapid and severe course of degeneration that is more

aligned with an acute trauma [7,20,21]. Unlike the increased attenuation of the injured annulus fibrosus, the

depressurized nucleus pulposus loses water and attenuates less than the healthy state (Figures 4).

Consistent with unilateral injuries of the tail IVD, the resultant degenerative cascade was nuanced[11], and

required high resolutions modalities to detect measurable changes. Future studies will evaluate the

partial-width injury as a slowly progressing IVD degeneration model following a clinically relevant

injury.

Acknowledgements

This work is in part supported by NIH R01AR074441, R01AR077678, K01AR069116, P30 AR007992,

and T32 DK108742. All authors contributed in the design, execution, and collection of data; as well as

writing and critical review of the manuscript.

References

1. Rao RD, Bagaria VB, Cooley BC. Posterolateral intertransverse lumbar fusion in a mouse model: surgical anatomy and operative technique. *Spine J*. 2007;7(1):61-67. doi:10.1016/J.SPINEE.2006.03.004

2. O'Connell GD, Vresilovic EJ, Elliott DM. Comparison of animals used in disc research to human lumbar disc geometry. *Spine (Phila Pa 1976)*. 2007;32(3):328-333. doi:10.1097/01.BRS.0000253961.40910.C1

3. Gruber HE, Sage EH, Norton HJ, Funk S, Ingram J, Hanley EN. Targeted deletion of the SPARC gene accelerates disc degeneration in the aging mouse. *J Histochem Cytochem*. 2005;53(9):1131-1138. doi:10.1369/JHC.5A6687.2005

4. Dahia CL, Mahoney EJ, Durrani AA, Wylie C. Postnatal growth, differentiation, and aging of the mouse intervertebral disc. *Spine (Phila Pa 1976)*. 2009;34(5):447-455. doi:10.1097/BRS.0B013E3181990C64

5. Lotz JC, Hsieh AH, Walsh AL, Palmer EI, Chin JR. Mechanobiology of the intervertebral disc. *Biochem Soc Trans*. 2002;30(Pt 6):853-858. doi:10.1042/BST0300853

6. Abraham AC, Liu JW, Tang SY. Longitudinal Changes in the Structure and Inflammatory Response of the Intervertebral Disc Due to Stab Injury in a Murine Organ Culture Model. *J Orthop Res*. 2016;34(8):1431. doi:10.1002/JOR.23325

7. Moss IL, Zhang Y, Shi P, Chee A, Piel MJ, An HS. Retroperitoneal approach to the intervertebral disc for the annular puncture model of intervertebral disc degeneration in the rabbit. *Spine J*. 2013;13(3):229-234. doi:10.1016/J.SPINEE.2012.02.028

8. Lotz JC. Animal models of intervertebral disc degeneration: lessons learned. *Spine (Phila Pa 1976)*. 2004;29(23):2742-2750. doi:10.1097/01.BRS.0000146498.04628.F9

9. Masuda K, Lotz JC. New challenges for intervertebral disc treatment using regenerative medicine. *Tissue Eng Part B Rev*. 2010;16(1):147-158. doi:10.1089/TEN.TEB.2009.0451

10. Vo N, Niedernhofer LJ, Nasto LA, et al. An overview of underlying causes and animal models for the study of age-related degenerative disorders of the spine and synovial joints. *J Orthop Res*. 2013;31(6):831-837. doi:10.1002/JOR.22204

11. Piazza M, Peck SH, Gullbrand SE, et al. Quantitative MRI correlates with histological grade in a percutaneous needle injury mouse model of disc degeneration. *J Orthop Res*. 2018;36(10):2771-2779. doi:10.1002/JOR.24028

12. Shi C, Das V, Li X, et al. Development of an in vivo mouse model of discogenic low back pain. *J Cell Physiol*. 2018;233(10):6589-6602. doi:10.1002/jcp.26280

13. Ohnishi T, Sudo H, Iwasaki K, Tsujimoto T, Ito YM, Iwasaki N. In Vivo Mouse Intervertebral Disc Degeneration Model Based on a New Histological Classification. *PLoS One*. 2016;11(8):e0160486. doi:10.1371/JOURNAL.PONE.0160486

14. Millecamps M, Stone LS. Delayed onset of persistent discogenic axial and radiating pain after a single-level lumbar intervertebral disc injury in mice. *Pain*. 2018;159(9):1. doi:10.1097/j.pain.0000000000001284

15. Lin KH, Wu Q, Leib DJ, Tang SY. A novel technique for the contrast-enhanced microCT imaging of murine intervertebral discs. *J Mech Behav Biomed Mater*. 2016;63:66-74. doi:10.1016/j.jmbbm.2016.06.003

16. Walk RE, Tang SY. In vivo contrast-enhanced microCT for the monitoring of mouse thoracic,

lumbar, and coccygeal intervertebral discs. *JOR spine*. 2019;2(2). doi:10.1002/JSP2.1058

17.   Berger-Roscher N, Galbusera F, Rasche V, Wilke HJ. Intervertebral disc lesions: visualisation
with ultra-high field MRI at 11.7 T. *Eur Spine J*. 2015;24(11):2488-2495. doi:10.1007/S00586-
015-4146-0

18.   Lin KH, Tang SY. The Quantitative Structural and Compositional Analyses of Degenerating
Intervertebral Discs Using Magnetic Resonance Imaging and Contrast-Enhanced Micro-Computed
Tomography. *Ann Biomed Eng*. 2017;45(11):2626-2634. doi:10.1007/S10439-017-1891-
8/FIGURES/6

19.   Melgoza IP, Chenna SS, Tessier S, et al. Development of a standardized histopathology scoring
system using machine learning algorithms for intervertebral disc degeneration in the mouse
model—An ORS spine section initiative. *JOR Spine*. 2021;4(2):e1164. doi:10.1002/JSP2.1164

20.   Masuda K, Aota Y, Muehleman C, et al. A novel rabbit model of mild, reproducible disc
degeneration by an anulus needle puncture: Correlation between the degree of disc injury and
radiological and histological appearances of disc degeneration. *Spine (Phila Pa 1976)*.
2005;30(1):5-14. doi:10.1097/01.BRS.0000148152.04401.20

21.   Sobajima S, Kompel JF, Kim JS, et al. A slowly progressive and reproducible animal model of
intervertebral disc degeneration characterized by MRI, X-ray, and histology. *Spine (Phila Pa
1976)*. 2005;30(1):15-24. doi:10.1097/01.BRS.0000148048.15348.9B

22.   Tian Z, Ma X, Yasen M, et al. Intervertebral Disc Degeneration in a Percutaneous Mouse Tail
Injury Model. *Am J Phys Med Rehabil*. 2018;97(3):170-177.
doi:10.1097/PHM.0000000000000818

23.   Yang F, Leung VYL, Luk KDK, Chan D, Cheung KMC. Injury-induced sequential transformation
of notochordal nucleus pulposus to chondrogenic and fibrocartilaginous phenotype in the mouse. *J
Pathol*. 2009;218(1):113-121. doi:10.1002/PATH.2519

24.   Jin L, Balian G, Li XJ. Animal Models for Disc Degeneration-an Update. *Histol Histopathol*.
2018;33(6):543. doi:10.14670/HH-11-910

25.   Cunha C, Lamas S, Gonçalves RM, Barbosa MA. Joint analysis of IVD herniation and
degeneration by rat caudal needle puncture model. *J Orthop Res*. 2017;35(2):258-268.
doi:10.1002/JOR.23114

26.   Elliott DM, Yerramalli CS, Beckstein JC, Boxberger JI, Johannessen W, Vresilovic EJ. The effect
of relative needle diameter in puncture and sham injection animal models of degeneration. *Spine
(Phila Pa 1976)*. 2008;33(6):588-596. doi:10.1097/BRS.0B013E318166E0A2

Figures

Figure 1. *Retroperitoneal dissection to expose the lumbosacral intervertebral disc using microscopic*

guidance

(A) The No. 11-scalpel blade indicates the left pelvic bone and forceps grip the gluteus muscles. **(B)** The

pelvis can be rotated posteriorly to increase working space by raising the gluteus muscles and rotating the

left pelvic bone. The abdominal wall and peritoneum are retracted anteriorly by a Penfield dissector to

expose the psoas muscles. (**C)** The psoas muscle can be stripped posteriorly-to-anteriorly by scraping out

muscles attached to the anterior surface of the pelvis with a Penfield dissector. (**D)** Superior margin of the

pelvic bone indicating the L6 vertebral body. (**E**) X-ray confirms the position of the pelvis relative to the

L5/6 and L6/S1 intervertebral discs.

mm

350   **Figure 2**. *Measuring the thickness of the intervertebral disc and the annulus fibrosus*

351   **(A-B)** Contrast-enhanced microCT (CEμCT) and **(C-D)** histology were used to measure the IVD widths

352   and anterior AF thicknesses. The CEμCT and histology measurements were highly correlated ($r^2 = 0.96$,

353    $p < 0.001$) and statistically indistinguishable (paired t-test, $p = 0.37$), confirming the fidelity of the non-

354   destructive evaluation of CEμCT.

Figure 3. *Depth of the partial-width injury measured by contrast-enhanced micro-computed tomography*

(A) The tip of the scalpel edge is marked at 0.3 mm to allow for visual confirmation during surgery that the partial-width injury is limited to the AF. **(B)** All of the injury depths were confirmed by CEμCT and histology to have depths that do not exceed of the AF in each injured IVD, confirmed by the ratio of injury depth to AF thickness which are less than 1 in all samples. **(C, D)** Representative CEμCT showing the range of depths achieved by this AF injury.

Figure 4. *Full-width (FW) injury induces a sustained loss of NP hydration compared to both the partial-width (PW) injury and Sham groups.*

(A-C) Contrast-enhanced microCT of injured level (L6/S1) where injury site is visible in both FW and PW groups. **(D)** Mean attenuation of the AF in the injured and uninjured groups. Paired t-test indicates significant increase in attenuation at injury site compared to rest of uninjured AF ($p < 10^{-5}$). **(E)** NI/DI of injured level (L6/S1) and uninjured control level (L5/6). While no changes in the whole IVD structure were detected between injured and uninjured levels with either FW or PW, the ratio of the nucleus pulposus intensity to the whole disc intensity (NI/DI – approximately 4% drop in intensity) revealed the loss hydration and proteoglycans in the nucleus pulposus of FW samples that is indicative of degeneration compared to the control IVDs.

Figure 5. *FW induces rapid and sustained IVD degeneration compared to both Sham and PW.*

(A) Safranin-O stained histological section of injured level (L6/S1) **(B)** Total IVD degenerative score of L6/S1. **(C)** Breakdown of NP, AF, endplate and interface boundary contributions to total IVD degenerative score. For both the PW and FW injuries, the site of injury were not observable on the histological sections after the 2-, 4- and 8- week post surgery. Significant degeneration was observed at all timepoints indicated by significant increases of degenerative scores in NP, AF, interface boundary and combined IVD degenerative grades in the FW group compared to Sham and PW (p < 0.05). While PW causes localized disruption to the AF, it does not appear to cause degeneration.

Supplemental data

Instrumentation

Instruments used in surgery included a size No. 11-scalpel blade (Henry Schein Medical), forceps

(Fisher Scientific), dissecting scissors (VWR), Penfield dissector (VWR), q-tips (Dukal Corporation), 4-0

nylon sutures (Ethicon), suture grabbers (Fisher Scientific), and microscope (Zeiss).

Methodology

Retroperitoneal approach to the intervertebral disc procedure: The left flank was shaved from the

ventral to the dorsal midlines using veterinary trimmers. Mice were then transferred to the operating suite

and positioned in the left lateral decubitus position. The skin was prepared for aseptic surgery via

washing/rinsing with povidone iodine scrub and alcohol rinse. The animals were draped to isolate the

prepped area. Isoflurane by inhalation was used to maintain surgical depth anesthesia throughout the

procedure.

There were two surface landmarks utilized for this approach: 1) the anterior margin of the thigh

(vertical) and bony prominence of the pelvis (horizontal) and 2) the fat pad reflected underneath the skin.

The fat pad located between the anterior margin of the thigh and abdominal wall was utilized as an access

point to the retroperitoneal space. A 1–1.5-cm horizontal incision was made along the bony prominence

of the pelvic bone line covering the entry and the anterior 1/3 of the thigh (Figure S1A). The vertical

caudal margin of the access point was retracted and resected by blunt dissection with Mosquito forceps

(Figure S1B and S1C). The gluteus muscles and left pelvic bone (a) were utilized as a grip to hold and

move the spine axis with forceps. The opening to the retroperitoneal space (c) between (a) and oblique

abdominal muscles (b) was carefully blunt-dissected until the psoas muscle was exposed (Figure S1D).

404
Supplemental Figure 1. *Lateral retroperitoneal approach to the mouse lumbosacral spine* **(A):** The

superficial anatomical landmarks are denoted by 1) the blue dotted line which outlines anterior margin of

the thigh and bony prominence of the pelvis, and 2) the red dotted line which outlines the fat pad located

between the thigh and abdominal wall. The black solid line is an approximately 1 cm incision along the

pelvis line that reveals the surgical access point adjacent to the anterior thigh. (B and C): The access point

to the retroperitoneal space lies between the blue and red dotted lines. The underlying fat pad and

anterior margin of the thigh can be easily separated and retracted to expose the retroperitoneal space by

blunt dissection. **(D):** The pelvis is rotated posteriorly to expose a broad working space by gripping (a)

with forceps. The area (c) between (a) and (b) is carefully separated by blunt dissection to expose the

psoas muscle: (a) gluteus muscles and left pelvic bone; (b) oblique abdominal muscles; (c) opening to the

retroperitoneal space.