1

Multisite Phosphorylation and Binding Alter Conformational Dynamics of the 4E-BP2 Protein

Spencer Smyth[1,2,*], Zhenfu Zhang[1,2,*], Alaji Bah[3,+], Thomas E. Tsangaris[2], Jennifer Dawson[3,$],

Julie D. Forman-Kay[3,4], and Claudiu C. Gradinaru[1,2,#]

[1]Department of Physics, University of Toronto, Toronto, Ontario, M5S 1A7, Canada;

[2]Department of Chemical & Physical Sciences, University of Toronto Mississauga, Mississauga,

Ontario, L5L 1C6, Canada; [3]Program in Molecular Medicine, The Hospital for Sick Children,

Toronto, ON M5G 0A4, Canada; [4]Department of Biochemistry, University of Toronto, Toronto,

ON M5S 1A8, Canada

[*] These authors contributed equally to this work

[+] Present address: Department of Biochemistry and Molecular Biology, SUNY Upstate Medical

University, Syracuse, NY 13210, United States

[$] Present address: Genomic Medicine Institute, Cleveland Clinic, Cleveland, OH 44195, United

States

[#]Author to whom correspondence may be addressed: claudiu.gradinaru@utoronto.ca

KEYWORDS: IDP, single-molecule fluorescence, anisotropy decay, PET-FCS, smFRET

ABSTRACT

Intrinsically disordered proteins (IDPs) play critical roles in regulatory protein interactions, but detailed structural/dynamics characterization of their ensembles remain challenging, both in isolation and they form dynamic 'fuzzy' complexes. Such is the case for mRNA cap-dependent translation initiation, which is regulated by the interaction of the predominantly folded eukaryotic initiation factor 4E (eIF4E) with the intrinsically disordered eIF4E binding proteins (4E-BPs) in a phosphorylation-dependent manner. Single-molecule Förster resonance energy transfer showed that the conformational changes of 4E-BP2 induced by binding to eIF4E are non-uniform along the sequence; while a central region containing both motifs that bind to eIF4E expands and becomes stiffer, the C-terminal region is less affected. Fluorescence anisotropy decay revealed a nonuniform segmental flexibility around six different labelling sites along the chain. Dynamic quenching of these fluorescent probes by intrinsic aromatic residues measured via fluorescence correlation spectroscopy report on transient intra- and inter-molecular contacts on ns-μs timescales. Upon hyperphosphorylation, which induces folding of ~40 residues in 4E-BP2, the quenching rates decreased at most labelling sites. The chain dynamics around sites in the C-terminal region far away from the two binding motifs significantly increased upon binding to eIF4E, suggesting that this region is also involved in the highly dynamic 4E-BP2:eIF4E complex. Our time-resolved fluorescence data paint a sequence-level rigidity map of three states of 4E-BP2 differing in phosphorylation or binding status and distinguish regions that form contacts with eIF4E. This study adds complementary structural and dynamics information to recent studies of 4E-BP2, and it constitutes an important step towards a mechanistic understanding of this important IDP via integrative modelling.

INTRODUCTION

Intrinsically disordered proteins (IDPs) are a class of proteins that lack well-defined three-dimensional structures while still carrying out biological functions (1-3). IDPs play a crucial role in mediating interactions with multiple partners and often function as protein interaction hubs (4, 5). IDPs within these protein complexes can undergo disorder-to-order transitions or remain dynamic (6). The lack of stable folded structures observed in IDPs leads to the simplistic assumption that IDPs resemble random coils. In fact, IDPs have transient secondary and tertiary structures and preferential backbone torsion angle propensities due to electrostatic and other interactions based on their amino acid sequence and exhibit a wide range of compactness (6-9).

Cap-dependent initiation of translation is regulated by the interaction of eukaryotic initiation factor 4E (eIF4E) with disordered eIF4E binding proteins (4E-BPs) in a phosphorylation-dependent manner (10-12). The eIF4E protein, which binds the 7-methyl guanosine cap structure of mRNA at the 5' end, has been shown to be an oncogene and be involved in the induction of cellular transformation (13, 14). The eIF4G, a scaffolding protein, plays a crucial role in docking and assembling several components of the translation initiation machinery at the 5' cap of mRNA to recruit the ribosome (15). The 4E-BP2 protein is involved in controlling cell growth and proliferation via regulating mRNA translation (16) and in immunity to viral infections (17). Neural 4E-BP2 also functions in regulating synaptic plasticity, playing an essential role in learning and memory, and has been implicated in autism spectrum disorder (18, 19).

The interaction between eIF4E and eIF4G is the fundamental step that initiates the translation process. This interaction involves the canonical binding helix-forming [54]YXXXXLϕ[60] (where Y is tyrosine, X is any amino acid, L is Leucine, and ϕ is a hydrophobic amino acid) motif of eIF4G, which is also found in all 4E-BPs, binding to the same convex interface of eIF4E. Thus,

the 4E-BPs regulate translation by competing with eIF4G to prevent the assembly of the eIF4F complex and the subsequent mRNA recruitment to the ribosome. Binding of IDPs often leads to ordering, the transient helical structure around the canonical [54]YXXXXLϕ[60] motif is stabilized upon eIF4E binding. However, the 4E-BP2:eIF4E complex has been shown by NMR to be highly dynamic with an exchanging bipartite interface (20), in which the secondary binding site [78]IPGTV[82] interacts with the lateral surface of eIF4E, as revealed by an X-ray crystal structure capturing a snapshot of the complex (21).

The 4E-BP2 protein is hierarchically phosphorylated. Modification of the first two sites T37 and T46 results in the hypo-phosphorylated state(22); this 2-site phosphorylation (2P) state induces formation of a 4-stranded β-sheet structure from residues 18-62, partially sequestering the canonical binding motif and weakening eIF4E binding but still enabling competition with eIF4G(12). Additional phosphorylation at S65, T70, and S83 leads to a 5-site phosphorylation (5P) state that stabilizes the fold (23), further decreasing the eIF4E affinity and allowing translation initiation to proceed (10). The disordered region C-terminal to the folded domain (C-IDR) remains disordered after phosphorylation and stabilizes the folded domain via long-range interactions (12, 23). However, important details of the structure and dynamics of 4E-BP2 and the eIF4E:4E-BP2 complex remain unknown, which prevent a clear mechanistic picture of the function of 4E-BP2 and its phosphoregulation of translation initiation.

Typically, IDPs have a wide range of interchanging conformations, therefore obtaining dynamic information is critically important for understanding their biological functions. Single-molecule fluorescence (SMF) techniques have been applied previously to measure the conformational heterogeneity, the global dimensions, and the dynamics of IDPs (24, 25). Here we applied a combination of ensemble and single-molecule techniques, i.e., fluorescence anisotropy

decay (FAD) and fluorescence correlation spectroscopy (FCS), to characterize global and local peptide chain motions in 4E-BP2 upon multisite phosphorylation and upon binding to eIF4E. FCS and FAD are sensitive to chain motions on the nanosecond-to-microsecond time scale, which is highly relevant for protein folding and for IDP dynamics (26). FAD measurements informed on the local, segmental flexibility of the peptide chain at various sites of the protein, leading to a coarse rigidity map of 4E-BP2 in the non-phosphorylated (NP), 5-site-phosphorylated (5P), and eIF4E-bound states. Using FCS, we resolved up to two different timescales of intra-molecular conformational dynamics in 4E-BP2 under NP, 5P, and denaturing conditions. In addition, we obtained kinetic information (amplitude and lifetime) of key inter-molecular contacts of the dynamic binding interface between NP 4E-BP2 and eIF4E.

Single-molecule Förster resonance energy transfer (smFRET) measurements of two different regions of the protein delineated changes in intramolecular distances and chain rigidity upon multisite phosphorylation, and upon binding to eIF4E. Surprisingly, smFRET showed an increase in the distance between residues 32 and 91 upon phosphorylation, despite folding-induced compaction of residues 18 to 62 (12). While in complex with eIF4E, a region of 4E-BP2 containing both canonical and secondary binding sites expands and stiffens considerably, while the C-terminal region expands only slightly but remains highly flexible. Our multifaceted spectroscopic characterization of the 4E-BP2 conformational dynamics is an important step towards understanding the interplay between folding and release of binding to eIF4E, and it provides valuable information for calculating conformational ensembles of this multistate IDP via integrative modelling.

 RESULTS AND DISCUSSION

 Local chain dynamics measured by FAD

The structural flexibility of IDPs is essential for regulating their interactions with other

proteins and their role in signaling processes (3). FAD measures the rotational dynamics of the

emission dipole of a fluorophore and is therefore a suitable reporter of the local (segmental) chain

flexibility around the labelling site. Inferred parameters from FAD analysis relate to the spatial

confinement and the friction experienced by the dye, the movement of a protein segment around

the labelling site, and the hydrodynamic radius of the protein segment (27). Probing local

conformational dynamics is particularly relevant for IDPs, for which motions of different segments

may be uncorrelated and obscured by the averaged global dynamics.

Six single-cysteine 4E-BP2 constructs were used for site-specific fluorescence labelling:

C0 (i.e., C0ins/C35S/C73S mutations), C14 (S14C/C35S/C73S), C35 (C73S), C73 (C35S), C91

(C35S/C73S/S91C), and C121 (C35S/C73S/C121ins) (**Fig. S1, Table S1**). While the fastest, sub-

ns FAD lifetime (ρ_{dye}) typically describes the rotation of the dye-linker around the labelling site,

in IDPs with transient structure the slowest FAD lifetime describes the rotational diffusion of a

protein segment (ρ_{seg}), where individual segments can rotate independently (28, 29). Alternatively,

for folded proteins the slowest FAD lifetime describes the rotational diffusion of the entire

protein.(30)

Fig. 1A shows a family of FAD curves of NP 4E-BP2 with the fluorophore attached to

each of the six different mutated cysteine sites. As dye-protein interactions are expected to be

negligible for Atto488 (31) (**Fig. S2**), the variations in the anisotropy decays indicate that the chain

flexibility is site/region dependent. FAD fitting parameters using **Eq. 1** (see Methods) are listed in

Table 1, for each labelling site and each sample condition. **Fig. 1C** shows a comparison between

segmental lifetimes (ρ_{seg}) around each 4E-BP2 labelling site in different phosphorylation states,

i.e., NP, 2P, and 5P. These values range from 1.4 ns to 3.4 ns and vary considerably with the

labelling site and with the phosphorylation state.

Figure 1. Fluorescence anisotropy decay data for 4E-BP2 fit to **Eq. 1.** The 4E-BP2 protein was labelled
with Atto 488 at positions 0, 14, 35, 73, 91 and 121 along the sequence, as described in the text. NP 4E-
BP2 fitted curves for all six labelling sites are shown in the absence (**A**) and in the presence of 1 µM of
eIF4E (**B**); the FAD of the free dye is shown in black in both panels, exemplar FAD data is shown for
positions 0 and 91. (**C**) The slowest rotational lifetime (ρ_{seg}) obtained by fitting FAD curves at each labelling
site for NP (**blue**) and 5P (**red**) 4E-BP2. (**D**) The slowest rotational lifetime (ρ_{seg}) obtained by fitting FAD
curves at each labelling site for NP 4E-BP2 in the absence (**blue**) and in the presence of (**red**) of eIF4E.
The concentration of 4E-BP2 used in these experiments was ~50 nM.

Many IDPs are more compact than an ideal statistical coil of the same length due to

transient intra-molecular contacts (7, 8, 32). Slower segmental dynamics in PBS buffer (pH 7.4,

140 mM NaCl) were observed than in chemical denaturant (6M GdmCl) at all labelling sites

(**Table S2**). The chemically denatured NP and 5P states of 4E-BP2 have similar segmental

flexibility signatures, with $\rho_{seg} = 0.8 - 1.0$ ns at the ends and $\rho_{seg} = 1.3 - 1.6$ ns at internal

sites. The values and the trend here match previous measurements for denatured proteins and the

expectations for a random coil state(27, 29). Additionally, the pattern of FAD lifetimes is

consistent overall with that of ^{15}N relaxation rates(20) and of ^{1}H-^{15}N nuclear Overhauser effect

(NOE) values (12). This suggests that FAD-measured segmental dynamics can probe transient

secondary structures and non-random chain contacts, and reports on the local degree of disorder

in IDPs.

Table 1. Anisotropy decay parameters for phosphorylated/bound 4E-BP2 states labelled at different sites [a].

	5P 4E-BP2		NP 4E-BP2		NP 4E-BP2 + elF4E	
	ρ_{dye} (ns)	ρ_{seg} (ns) (a)	ρ_{dye} (ns)	ρ_{seg} (ns) (a)	ρ_{dye} (ns)	ρ_{seg} (ns) (a)
C0	0.53	1.75 (0.72)	0.51	1.61 (0.66)	0.52	1.93 (0.70)
S14C	0.56	2.56 (0.51)	0.64	3.27 (0.42)	0.62	3.36 (0.49)
C35	0.55	3.13 (0.41)	0.64	2.48 (0.52)	0.62	2.65 (0.51)
C73	0.55	2.35 (0.54)	0.62	2.86 (0.47)	0.61	4.54 (0.46)
S91C	0.57	2.01 (0.51)	0.61	2.59 (0.50)	0.65	5.35 (0.43)
C121	0.52	1.43 (0.61)	0.53	1.90 (0.60)	0.59	3.34 (0.59)

162 [a] All data were fit to **Eq. 1** to estimate two rotational correlation lifetimes (dye & segment) and their fractions (*a* is the
163 fraction of the slowest component). Fitting error margins are on the order of ±0.1 ns for the lifetimes and ±0.05 for
164 amplitudes **(Table S2).**
165

Another contribution to the heterogeneous segmental flexibility is the amino acid

composition around each labelling site, with glycine and serine being the most flexible, and

proline, isoleucine, and valine the most rigid (33). With 4 serine and 2 glycine among the first 10

residues, the N-terminus is the most flexible region of NP 4E-BP2. Considering a 10-residue

window, the positions 14, 73 and 91 are flanked by several rigid residues (proline, isoleucine,

valine) and the protein appears to be much less flexible in these regions. The slowest rotational

lifetime was observed at position 14 ($\rho_{seg} = 3.27$ ns), which, in addition to two proline residues,

has two positively charged arginine residues in its proximity, which may rigidify the segment

further via electrostatic interactions.

In previous NMR studies, we have shown that a region of 4E-BP2 (residues 18 to 62) folds

upon phosphorylation, while the rest of the chain remains disordered (12). We used FAD to probe

the changes in segmental flexibility of 4E-BP2 that are induced by complete (5P) phosphorylation

(**Fig. 1C, Table 1**). The rotational lifetime (ρ_{seg}) increases at position 35 while it decreases at all

the other dye positions, which are outside the folded domain, indicating that the chain becomes

more flexible. This is consistent with the formation of the four-stranded beta-sheet fold between

residues 18-62 and with the C-terminal region remaining disordered after phosphorylation. From

an entropic perspective, increased conformational flexibility near the secondary binding site

(residues 78 to 82) may also contribute to decreasing the affinity for eIF4E.

FAD has been used previously to quantify and differentiate local binding constants of IDPs

in the context of multisite interactions (28) . **Fig. 1B** shows anisotropy decay data and fitted curves

for the NP 4E-BP2:eIF4E complex at each of the six labeling sites on 4E-BP2. In contrast with the

apo sample, these curves decay to significantly higher asymptotic values (r_{inf}, **Table S2**), indicating

that the local motions around each labeling site in 4E-BP2 are hindered after binding to eIF4E.

The largest changes in chain flexibility occur at three C-terminal sites while the changes observed

in the N-terminal sites are minor (**Fig. 1D**). At positions 73, 91 and 121, the segmental rotational

lifetime ρ_{seg} nearly doubles, from 2-3 ns in the apo state to 4-5 ns in the bound state. Similarly,

r_{inf} values significantly increase, with the largest changes at the C-terminal labelling sites (**Fig.**

S3).

Positions 73 and 91 are located near the secondary binding site while position 121 is far

from either binding site but the changes in lifetime are consistent with binding-induced changes to

NMR intensity ratios(20), which demonstrate complete broadening for residues 45-88, significant

broadening for residues 34-90, and broadening at residue 120. The data for positions 73 and 91

likely reflect favorable interactions at the secondary binding site, as well as potentially the

competitive interaction of the secondary binding region with the disordered N-terminus of eIF4E

that has been suggested to act as a negative regulator (34). The NMR broadening results for

position 120 together with these FAD data on position 121 support a picture of the 4E-BP2:eIF4E

dynamic complex involving a more extensive part of the C-terminus.

The results demonstrate that segmental motion parameters measured by FAD can be used

to disentangle the binding contributions of different regions in an intrinsically disordered protein

as it interacts with its binding partners. More specifically, the changes in r_{inf} are correlated with

the changes in ρ_{seg}, indicating that the rotational freedom of the segment and the segmental

dynamics of 4E-BP2 are hindered in the bound state.

Non-local chain dynamics in 4E-BP2 measured by PET-FCS

The same Atto488-labelled 4E-BP2 constructs used for FAD (**Fig. S1, Table S1**) were also

used for FCS experiments. FCS is sensitive to intensity fluctuations caused by the diffusion of the

labelled protein and by the photophysics of the fluorophore (**Eq. 2**). Notably, the fluorophore can

be dynamically quenched by aromatic amino acids via photoinduced electron transfer (PET)(35,

36). Tryptophan and tyrosine are the strongest PET quenchers, with a quenching range of 5-10 Å

for typical fluorophores (37, 38). The two tyrosines in the 4E-BP2 sequence (Y34 and Y54) are

most likely acting as quenchers in our FCS measurements. Given that the PET quenching effect in

our measurements is not produced by a single residue, we do not employ a specific kinetic model

and instead limit our interpretation to the PET induced exponential lifetimes and amplitudes

accounting for in our fitting model (**Eq. 2**).

Fig. 2A shows experimental FCS data, fitting curves, and residuals for a representative sample, 4E-BP2 labelled with Atto488 at position 73, in the NP 4E-BP2:eIF4E state. Models with one diffusion component and two or three (faster) kinetic components (**Eq. 2**) satisfactorily fit the experimental autocorrelation decays measured for all 4E-BP2 samples. The (sub-diffusion) kinetic components in FCS data are attributed to intrinsic triplet-state kinetics of the probe and to dynamic PET quenching induced by the protein environment.

The FCS lifetimes obtained by fitting the data to **Eq. 2** can be grouped into three timescales: $\tau_1 \approx$ 50-100 µs, $\tau_t \approx$ 7-8 µs, and $\tau_2 \approx$ 50-300 ns, respectively (**Fig. 2, Table 2, Fig. S4**). Conversely, the free dye exhibits a single kinetic component with a lifetime of ~3-10 µs (**Fig. S5**), which is similar to a previously reported triplet lifetime of Atto488.(39) Thus, for 4E-BP2, the τ_1 and τ_2 lifetimes are attributed to PET quenching of the dye due to conformational dynamics of the protein, while the τ_t lifetime was assigned to the intrinsic triplet lifetime of Atto488. Therefore the (τ_t) and (a_t) component were fit as shared parameters in each measurement condition, the diffusion coefficient is also a shared (see methods). Chain motions on the sub-µs (τ_2) timescale are typically associated with interconversion of states within the disordered conformational ensemble and with proximal loop formation.(40, 41). The slowest (τ_1) kinetics are on the same timescale as concerted motions of the protein chain, such as domain movements or transient tertiary structural contacts (43, 44).

Figure 2. FCS lifetimes of 4E-BP2 labelled with Atto488 at residue 0, 14, 35, 73, 91 or 121. Experimental
curves were fit to **Eq. 2**. Fitting of NP 4E-BP2 labeled at residue 73 in the presence of 0.5 μM eIF4E, the
kinetic decays and diffusion for the best fit plotted separately (**A**). FCS experiments were performed on 4E-
BP2 at a concentration of ~5 nM in different states: non-phosphorylated (**B**), hyper-phosphorylated (**C**), in
the presence of 0.5 μM of elF4E (**D**). Different decay timescales are shown in different colors: $\tau_1 \approx 100$ μs
(**blue**) and $\tau_2 = 0.05$-0.3 μs (**red**). The fitting error bars for each lifetime are shown in the figure. The full list
of fitting parameters is given in **Table S3**.

For NP 4E-BP2, one or two kinetic components were resolved in addition to the triplet

component at each labelling position, which follow the **τ_1** and **τ_2** timescales. The lifetimes for the

τ_1 components are site-dependent (**Fig. 2B**). The fastest component **τ_2** was only resolved at

positions C73 and S91C, this component was not resolved for the other positions within

experimental error limits. Similar PET-FCS analysis, with multiple lifetimes of intrachain

dynamics between ~100 ns and ~100 μs, was previously reported for other IDPs, e.g., the N-

terminal domain of p53-TAD (42) and the mouse prion protein moPrP (43).

Table 2. FCS decay parameters for five-phospho, non-phospho and elF4E-bound 4E-BP2 [a].

	5P-BP2	NP 4E-BP2		NP 4E-BP2 + elF4E	
	$\tau_1(\mu s)$	$\tau_1(\mu s)$	$\tau_2(\mu s)$	$\tau_1(\mu s)$	$\tau_2(\mu s)$
C0	83.9 *(0.23)*	57.1 *(0.13)*	-	96.3 *(0.28)*	-
S14C	70.1 *(0.17)*	59.7 *(0.13)*	-	101.8 *(0.22)*	-
C35	79.4 *(0.18)*	53.7 *(0.11)*	0.31 *(0.11)*	100.5 *(0.25)*	0.10 *(0.16)*
C73	64.9 *(0.14)*	55.5 *(0.11)*	0.20 *(0.11)*	99.9 *(0.19)*	0.04 *(0.35)*
S91C	65.2 *(0.16)*	67.2 *(0.17)*	-	101.0 *(0.26)*	0.14 *(0.19)*
C121	71.7 *(0.16)*	54.3 *(0.12)*	-	59.2 *(0.15)*	0.16 *(0.14)*

[a] All data were fit to **Eq. 2** Lifetimes are given in microseconds, with corresponding amplitudes in brackets. Fitting errors are given in SI (Table S3).

In contrast to the NP state, the 5P state exhibits only one FCS kinetic component (τ_1) in addition to the triplet component at each labelling position (**Fig. 2C**). In the 5P state, the addition of phosphate groups and the induced folding in the 18-62 region probably slows down the protein reconfiguration time sufficiently to be "absorbed" into the low μs triplet dynamics, as previously reported for p53-TAD (42). Upon phosphorylation, the τ_1 component slows down at all positions except 73 and 91 where no change is observed within the experiment error limits; the associated kinetic amplitude a_1 decreases at all sites except for position 91 (**Table 2, Fig. S6**), from which it can be inferred that the dominant effect of phosphorylation is to slow down intrachain contacts causing PET quenching. Similarly, slower conformational dynamics upon multisite phosphorylation has been observed for the disordered p53-TAD protein, even though phosphorylation does not lead to folding in this system (42). In 4E-BP2, phosphorylation stabilizes

the β-strand structure between residues 18 and 62, which is expected to limit the access of the

fluorophore to tyrosine quenchers (12).

Given the tight binding affinity of 4E-BP2 for eIF4E ($k_d = 3.2 \pm 0.6$ nM)(12), saturating

amounts of eIF4E (0.5 μM) were used to ensure that nearly all (>99%) 4E-BP2 molecules were in

the bound state. FCS measurements on Atto488-labelled 4E-BP2 with or without unlabeled eIF4E

(**Fig. S2**) showed a shift to longer diffusion times (larger R_H) in the presence of eIF4E, which is

consistent with the formation of the 4E-BP2:eIF4E complex. The R_H values estimated by fitting

the data to **Eq. 2** are 29.0 ± 0.5 Å for the NP 4E-BP2 (independent of labelling site) and $36.6 \pm$

0.8 Å for the 4E-BP2:eIF4E complex. As reference, FCS measurements of Atto488 (5 nM) in the

presence of excess eIF4E (1 μM) showed that the dye does not interact/bind with eIF4E (**Fig S2**).

When bound to elF4E, NP 4E-BP2 exhibits one to two FCS kinetic components in addition

to the triplet component which fall into the same τ_1 and τ_2 range as in the free (Apo) state, (**Fig.**

2D). eIF4E (PDB: 3AM7) contains 8 tryptophan and 6 tyrosine residues, of which four tryptophan

(W46, W56, W73 and W102) and three tyrosine (Y34, Y76, and Y 145) residues are largely surface

exposed (**Table S4**). Therefore, the observed changes in PET quenching dynamics are the

combined result of the new inter-molecular interactions between 4E-BP2 and the surface of eIF4E

and the changes in intra-molecular contacts within 4E-BP2.

Figure 3. The changes in lifetime and amplitude of the fastest and slowest FCS kinetic components (τ_2 –
circle, and τ_1– **triangle**) induced by binding of NP 4E-BP2 to elF4E. FCS measurements on NP 4E-BP2
labelled with Atto488 at six different sites were performed in the Apo state (**blue**) and in the presence of
0.5 μM elF4E (**red**), and the data were fit to **Eq. 2.** The fitting error bars for each parameter are indicated
in the figure; some error bars are too small to be seen, all fitting parameters are given in **Table S3**.

4E-BP2 interacts with eIF4E through dynamic interactions involving at least an α-helical

structure at the canonical ^{54}YXXXXLφ^{60} motif and the secondary ^{78}IPGVT^{82} site (20, 21).

Positions 35 and 73 show a decrease in lifetime accompanied by an increase in amplitude of the

fast component, positions 91 and 121 gain this component upon binding to eIF4E (**Fig. 3 C-F**). As

such, τ_2 decreases from ~200-300 ns to ~50-100 ns upon binding at positions 35 and 73, while τ_2

at positions 91 and 121 is ~150 ns when bound (**Table 2**). The appearance of the fast τ_2 component

at positions 91 and 121, as well as the decrease in τ_2 lifetime and increase in a_2 amplitude at position

73 are likely caused by dynamic exchange of the secondary binding site leading to quenching by

aromatic residues on the surface of eIF4E. The involvement of residues 73, 91 and 121 in eIF4E

binding is consistent with binding-induced NMR intensity changes reported previously(20). The

changes at residue 121 upon binding mirror those observed by FAD (**Fig. 1 D**), providing further

support for a dynamic complex involving more of the C-terminal portion of 4E-BP2. Taken

together, the results highlight the dynamic nature and range of timescales present in the bound

state and ability of PET-FCS to probe intermolecular dynamics within the bound state.

Chain dimensions and stiffness of 4E-BP2 assessed by smFRET

smFRET is exquisitely suited to delineate heterogeneous and dynamic states that are

inherent to IDPs. smFRET can resolve heterogeneous population distributions and kinetics, is

compatible with a wide range of solution conditions, and overcomes the ensemble averaging of

established structural techniques such as NMR and small-angle X-ray scattering (SAXS)(25). By

recording the arrival time, the color and the polarization of each detected photon, multiparameter

fluorescence (MPF) detection permits access to additional intrinsic properties of fluorescence that

can be related to properties of the conjugated molecule (46). The chain dimensions and stiffness

of 4E-BP2 were assessed using smFRET with MPF on two double-cysteine FRET constructs,

which were labelled stochastically with Alexa Fluor 488 (donor) and Alexa Fluor 647 (acceptor).

smFRET histograms for H32C/S91C in the NP, 5P, and NP+eIF4E states are shown in **Fig.**

4. Although each histogram was satisfactorily fit to a single Gaussian, a single FRET peak does

not necessarily reflect a homogenous distribution of states. Indeed, the underlying population is

likely in fast exchange compared to the burst duration (~1 ms), given the typical chain

reconfiguration times of IDPs (~100 ns) (51). The data suggests that the NP state is more compact

than a statistical coil ensemble, i.e., $\langle E^{NP}_{32-91} \rangle_{exp} = 0.63 \pm 0.02$ vs. $\langle E^{NP}_{32-91} \rangle_{coil} = 0.43 \pm 0.03$

(see SI section 2.3). This is consistent with the presence of transient secondary structure observed

by NMR(20), and with the multiple transient intrachain contacts observed by PET-FCS.

Figure 4. smFRET MPF results for the H32C/S91C 4E-BP2 construct labelled with Alexa Fluor 488 and Alexa Fluor 647 in the (**A**) non-phospho (NP), (**B**) five-phospho and (5P) (**C**) non-phospho eIF4E-bound (NP 4E-BP2 + eIF4E) states. (1st column) Cartoon depictions of conformations of 4E-BP2 in different states with representations of fluorophore accessible volume simulations (47). (2nd column) FRET efficiency histograms fit to a Gaussian function, with the dashed lines indicating the mean values. (3rd column) 2D histogram plots of the FRET efficiency (*E*) and the donor-only normalized fluorescence lifetime; τ_{DA} and τ_{D0} are donor lifetimes in the presence and absence of acceptor, respectively. The black line shows the relation expected for a static rigid molecule. The red line is a relation for a worm-like chain with a stiffness parameter (κ) that passes through the centroid of a 2D gaussian fit indicated by a yellow cross. The red shaded region shows the uncertainty range of κ.

Upon multi-site phosphorylation 4E-BP2 undergoes folding to a beta-sheet domain between residues 18 and 62 (12). Typically, when proteins fold, FRET efficiency increases, following the overall compaction of the structure and contraction of most (but not all!) inter-residue distances(48, 49). In this case however, the FRET efficiency of the 5P state is lower for the dye pairs at residues 32 and 91, $\langle E_{32-91}^{5P} \rangle_{exp} = 0.44 \pm 0.02$, not higher than the NP state, $\langle E_{32-91}^{NP} \rangle_{exp} = 0.63 \pm 0.02$ (**Table 3**). The dye at C32 is positioned in the long loop between

strands β1 and β2 which is proximal to the C-terminal end of the domain (PBD ID 2MX4), from

where the C-IDR, containing the dye at C91, extends. An alternate construct designed to

specifically probe the folded region (with both dyes within the folded domain boundaries from 18-

62) would likely lead to an increase in FRET efficiency upon phosphorylation. The H32C/S91C

construct, however, also partially probes the C-IDR, which expands when 4E-BP2 is

phosphorylated (**Fig. 5B**).

The FRET efficiency of the NP H32C/S91C construct exhibits an even larger decrease

upon binding to eIF4E, i.e., $E_{32-91}^{NP+4E} = 0.26 \pm 0.02$. This construct flanks both primary and

secondary binding sites and should be sensitive to 4E-BP2:eIF4E interactions and the 3D

arrangement of the complex. The large FRET decrease could result from a combination of more

extended conformations being compatible with the eIF4E bound state, in which 4E-BP2 wraps

around eIF4E (with the exception of the canonical helical element), and an excluded-volume effect

exerted by eIF4E on 4E-BP2. The conformations of 4E-BP2 bound to eIF4E are thought to be

defined by dynamic interactions of the canonical and secondary binding elements, and by

significant structural disorder elsewhere (20).

smFRET histograms of the C73/C121 construct for the same three 4E-BP2 states are shown

in **Fig. 5**. The average FRET efficiency for NP 4E-BP2, $\langle E_{73-121}^{NP} \rangle_{exp} = 0.58 \pm 0.02$, is also

higher than that expected from a statistical coil, $\langle E_{73-121}^{NP} \rangle_{coil} = 0.49 \pm 0.03$, suggesting that the

C-terminal disordered region contains transient intrachain contacts. The average C73/C121 FRET

efficiency decreases to $\langle E_{73-121}^{5P} \rangle = 0.44 \pm 0.02$ upon phosphorylation, but only to $\langle E_{73-121}^{NP+4E} \rangle =$

0.51 ± 0.02 upon binding to eIF4E. The electrostatic repulsion between the five phosphates may

contribute to the C-IDR expansion of 5P 4E-BP2. This expansion of the C-IDR has also been

observed from previously published NMR paramagnetic relaxation enhancement (PRE) data and

chemical shift-derived measures of secondary structure.(23)

Figure 5. smFRET MPF results for the C73/C121 4E-BP2 construct labelled with Alexa Fluor 488 and Alexa Fluor 647 in the (**A**) non-phospho (NP), (**B**) five-phospho (5P) and (**C**) non-phospho eIF4E bound states (NP 4E-BP2 + eIF4E). (1st column) Cartoon depictions of conformations of 4E-BP2 in different states with representations of fluorophore accessible volume simulations (47). (2nd column) FRET efficiency histograms fit to a Gaussian function, with the dashed lines indicating the mean values. (3rd column) 2D histograms plots of the FRET efficiency (E) and the donor-only normalized fluorescence lifetime; τ_{DA} and τ_{D0} are donor lifetimes in the presence and absence of acceptor, respectively. The black line shows the relation expected for a static rigid molecule. The red line is a relation for a worm-like chain with a stiffness parameter (κ) that passes through the centroid of a 2D gaussian fit indicated by a yellow cross. The red shaded region shows the uncertainty range of κ.

For the bound state, the reduction in FRET is much less for C73/C121 than for H32C/S91C, i.e.,

$\Delta E_{73-121}^{NP\rightarrow+4E} = -(0.07 \pm 0.03)$ *vs.* $\Delta E_{32-91}^{NP\rightarrow+4E} = -(0.37 \pm 0.03)$. This suggests that C-terminal

disordered conformations are less restricted in the NP 4E-BP2:eIF4E complex than those of the

32-91 region, as expected from the complete loss or very significant broadening of NMR

resonances upon binding from residues 34-90 (20).

The relation between the donor lifetime and the FRET efficiency can be used to infer

information about the dynamic exchange of the underlying states (50). A linear relation is expected

for a static structure, with conformations that are rigid or fluctuate on a timescale slower than ~100

µs; in contrast, a nonlinear relation is expected for IDPs, as the burst duration (~1 ms) is much

longer than typical chain reconfiguration times (~100 ns) (51). A family of dynamic τ vs. E lines

based on a worm-like chain (WLC) model with variable stiffness κ (or persistence length) was

generated using a method described by Barth et al. (52) The center of the experimental 2D FRET

histogram was best matched to a WLC curve to infer the average stiffness for different regions of

4E-BP2 in different phospho/binding states (**Table 3**).

Table 3. Mean smFRET efficiency and WLC stiffness values of the non-phospho (NP), five-phospho
(5P) and elF4E-bound 4E-BP2 (NP 4E-BP2 + eIF4E).

	NP 4E-BP2		5P-BP2		NP 4E-BP2 + elF4E	
	$<E>$	κ	$<E>$	κ	$<E>$	κ
H32C/ S91C	0.63 ± 0.02	0.05 ± 0.03	0.44 ± 0.02	0.25 ± 0.10	0.26 ± 0.02	0.66 ± 0.17
C73/ C121	0.58 ± 0.02	0.14 ± 0.08	0.44 ± 0.02	0.34 ± 0.10	0.51 ± 0.02	0.27 ± 0.08

The stiffness parameter of the 32-91 region, $\kappa_{32-91}^{NP} = 0.05 \pm 0.03$, corresponds to a

persistence length of $l_{p\,32-91}^{NP} = 1.0 \pm 0.6$ nm, which is similar to $l_p = 0.4 \pm 0.07$ nm reported for

a set of disordered and unfolded proteins.(53) The stiffness increases to $\kappa_{32-91}^{5P} = 0.25 \pm 0.10$

upon phosphorylation, and even more significantly, to $\kappa_{32-91}^{NP+4E} = 0.66 \pm 0.17$ upon binding to

eIF4E (**Fig. 4** and **Table 3**). The larger stiffness in the 5P state is consistent with the appearance

of a stable beta-fold between residues 18 and 62. In the bound state, the 2D FRET population lies

close to the static line; this is consistent with the decreased dynamics expected as residues 49-67

form a predominantly stable helix when 4E-BP2 is bound to eIF4E.(20, 54) At the same time, the

separation between the static and dynamic regimes is much reduced in the low E range, which

increases the uncertainty in estimating the chain stiffness.

The 73-121 region has a stiffness of $\kappa_{73-121}^{NP} = 0.14 \pm 0.08$ and $\kappa_{73-121}^{5P} = 0.34 \pm 0.10$

in the NP and 5P states, respectively. While slightly larger, these values mirror the changes

observed for the 32-91 region. The stiffening of the C-IDR upon phosphorylation can be attributed

to stabilizing interactions with the folded domain established previously (23). However, in contrast

to the 32-91 region, the stiffness of the C-terminal region shows only a moderate increase in the

eIF4E-bound state, to $\kappa_{73-121}^{NP+4E} = 0.27 \pm 0.08$, as the 2D FRET histogram remains well separated

from the static line (**Fig. 5**). Together with the FAD and PET-FCS results, this suggests that the

73-121 region remains disordered and interacts with eIF4E as part of a more extensive dynamic

complex than the current bipartite model (20).

CONCLUSIONS

Dynamics is likely the key factor in understanding how 4E-BP2 regulates cap-dependent

translation via interactions with the initiation factor eIF4E. Complementary to our previous NMR

studies of the 4E-BP2 protein (12, 20, 23), a suite of multiparameter fluorescence techniques was

used here to define local and global conformational dynamics of this intriguing IDP in its various

states. Polarization anisotropy (FAD) measurements showed evidence of heterogeneous chain

flexibility within the 4E-BP2 sequence, which correlates with variations in transient secondary

structure and local amino acid composition. Multisite phosphorylation decreased segmental

flexibility only in the region that undergoes folding (residues 18-62), while it made the rest of the

chain more flexible. As anticipated, in the presence of eIF4E, segmental dynamics near the two

binding sites was slowed down by binding interactions. Surprisingly however, the C-terminal

region adjacent to the secondary binding site experienced the largest changes, while the N-terminal

region exhibited much less change. These results implicate a much larger region of 4E-BP2

interacting with eIF4E than previously thought and showcase FAD as a sensitive method to probe

local binding affinity, at the level of individual protein segments, in the context of multivalent

interactions.

As seen previously for other IDPs (42), the quenching rates decreased at most labelling

sites. This likely arises from a combination of local steric and electrostatic effects of the phosphate

groups and the overall steric effect of the folded domain. In the presence of eIF4E, the quenching

rates near the two binding motifs in 4E-BP2 increased significantly, as expected from increased

contacts with several exposed tryptophan and tyrosine residues along the extensive binding

interface of eIF4E. Although contributions from intra- and inter-molecular contact could not be

distinguished, the significant changes observed at the C-terminus support the involvement of the

C-terminal region of 4E-BP2 in dynamic interactions with eIF4E.

smFRET measurements informed on changes in 4E-BP2 chain dimensions and stiffness of

central and C-terminal regions of the protein following phosphorylation or binding. In the NP state,

as a consequence of transient intrachain secondary and tertiary contacts, the donor-acceptor

distance turned out smaller (higher FRET efficiency) than the random coil prediction for both

FRET constructs. In the 5P state, the separation between residues 32 and 91, which partially

contains the folding domain, increased, instead of the decrease typically expected for a folding

transition. When bound to eIF4E, the chain expands and stiffens considerably around the canonical

and secondary binding sites, while the C-terminal region remains highly flexible. The canonical

binding motif and secondary binding site provide specificity, while the dynamic nature of the

complex which retains significant chain flexibility minimizes the entropic penalty. All

phosphorylation sites except S65, which is phosphorylated last, are in regions that remain highly

dynamic in the 4E-BP2:eIF4E complex. This facilitates the access of large kinases and allows a

rapid response to cellular conditions. Our fluorescence-based characterization of the 4E-BP2

conformational dynamics is an important step towards understanding the interplay between folding

and release of binding to eIF4E and its regulatory function, and it provides a foundation for future

studies of IDP conformational and binding equilibria.

MATERIALS AND METHODS

Materials. The fluorescent probes used for labelling the 4E-BP2 protein were: Alexa Fluor 488 (A488) maleimide, Alexa Fluor 647 (A647) maleimide (ThermoFisher Scientific, Canada) and, Atto488 (At488) maleimide (ATTO-TEC GmbH, Germany). Guanidinium chloride (GdmCl) (G9284, Sigma Aldrich) was used for protein denaturation. All samples were diluted in phosphate-buffered saline (PBS) containing 137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, and 1.8 mM KH_2PO_4 at pH 7.4. GdmCl solutions were adjusted to pH 7.4 for all the denaturation experiments.

Protein expression and purification. Small ubiquitin-like modifier (SUMO) fusion constructs of both the 4E-BP2 protein (residues 1 through 120) and eIF4E protein (residues 1 through 217) were expressed and purified as described previously[23]. Briefly, the proteins were expressed in BL21-codonplus (DE3)-RIPL competent *E. coli* cells (Agilent Technologies) in Lysogeny broth at 37 ℃ until OD_{600}~0.6-0.8, induced with isopropyl β-D-1-thiogalactopyranoside (IPTG), and expressed at 16 ℃ for ~16 h. Protein was purified from cell lysate with a nickel–nitrilotriacetic acid (Ni–NTA) column followed by cleavage of the SUMO solubility tag with ULP1 at 4 ℃ for ~16 h. The Sumo tag was separated using an Ni-NTA column followed by HiLoad Superdex 75 PG gel filtration column (28-9893-34, GE Healthcare) if the protein was not pure as assessed by SDS-PAGE. The molecular mass and the purity of protein samples were verified by electrospray ionization mass spectrometry (ESI-MS).

Phosphorylation of 4E-BP2 with activated Erk2 using a dialysis technique was performed as described previously[12, 23]. Briefly, 50 mL of 5 µM Erk2 and 20 µM of 4E-BP2 were dialyzed in a 3 kDa MWCO dialysis bag in 1 L of buffer. The dialysis buffer contained 50 mM Tris-HCl, pH 7.5, 1 mM EGTA, 5 mM BME, 20 mM $MgCl_2$, 10 mM EDTA, and 10 mM ATP, dialysis was

performed at 20 ℃ for 1-3 days. Phosphorylated 4E-BP2 was purified from Erk2 using a Ni-NTA

column. Purity and degree of phosphorylation of 4E-BP2 was confirmed by ESI-MS.

All single cysteine proteins (C0/C35S/C73S (cysteine insertion at 0 position),

S14C/C35S/C73S, C35/C73S, C73/C35S, C35S/C73S/S91C, C35S/C73S/C121 (cysteine

insertion at 121 position)) were labeled by adding the Atto488 maleimide fluorophore to a 50 µL

solution of 100 µM protein at a dye:protein molar ratio of 3:1. The double-cysteine mutants

(H32C/C35S/C73S/S91C) and (C35S/insC121) was labeled with Alexa Fluor 488 maleimide and

Alexa Fluor 647 maleimide by adding Alexa Fluor 488 and Alexa Fluor 647 to a 50 µL solution

of 100 µM protein at a A488:A647:protein molar ratio of 1.3:3:1. TCEP was added at a $10\times$ molar

excess to the protein in order to reduce the disulfide bonds. All the maleimide-cysteine coupling

reactions were performed in a PBS buffer at pH 7.4. Oxygen was removed by flushing the sample

with argon gas in a desiccator for 5 min. The vial was capped tightly and shaken gently for 3 hours

at room temperature. The excess dye was removed by size-exclusion chromatography using

Sephadex G-50 gel (G5080, Sigma Aldrich) in a BioLogic LP system (731-8300, Bio-Rad).

All samples were diluted to concentrations of 1–10 nM and 20–50 pM, which are most

suitable for FCS/FAD and smFRET burst experiments, respectively. For a typical experiment, a

sample solution of 30 µL was dropped on the surface of plasma-cleaned coverslip. Non-specific

protein adsorption to the coverslip was prevented by adding 0.005% (v/v) Tween-20 (P2287,

Sigma-Aldrich) to the solution, and bovine serum albumin (BSA) (15260-037, ThermoFisher

Scientific) was used to coat the clean coverslips. All experiments were performed at 20 °C.

Instrumentation. smFRET measurements were performed on a custom-built

multiparameter fluorescence microscope (32). The donor was excited at 480 nm by frequency

doubling the infrared output of a femtosecond laser (Tsunami HP, Spectra Physics), while the

acceptor was excited at 635-nm using a diode laser (WSTech, TECRL-25GC-635). Alternating-

laser excitation (ALEX) of the sample was performed by synchronous modulation of the two laser

sources to achieve alternating 50-μs periods of donor and acceptor fluorophore excitation,

respectively. Laser intensities of 10 kW/cm^2 and 3.6 kW/cm^2 at the sample were used for exciting

the donor and the acceptor fluorophores, respectively. FAD measurements were performed on the

same microscope, by exciting Atto488 at an average intensity of ~0.14 kW/cm^2 at the sample. FCS

measurements were performed on a separate custom-built fluorescence microscope described

elsewhere (55), where Atto488 was excited using a 488-nm diode laser (TECBL-488nm,

WorldStarTech) at an average intensity of ~5 kW/cm^2 at the sample.

FAD analysis. FAD monitors the rotation dynamics of the emission dipole of the dye. The

"wobble-in-a-cone" model (30) was used to fit the experimental FAD data:

$$r(t) = r_0\big[(1-a)e^{-t/\rho_{dye}} + a\big]e^{-t/\rho_{seg}} + r_{inf} \tag{1}$$

where ρ_{dye}, ρ_{seg} are rotational correlation lifetimes of the dye and the protein segment,

respectively, *a* is the fraction of the slower (segmental) component, *r₀* is initial (limiting)

anisotropy, and *r_inf* is the non-decaying (residual) anisotropy. The uncertainties of the fitted

parameters were estimated by taking the standard deviation of the fitting results of 3-7 data sets

collected consecutively; if any of the fitting derived errors were larger, then this value was

reported. The baseline offset r_{inf} accounts for the slow, global motion of the whole protein; this

is typically very small for IDPs due to their high backbone flexibility (29), but it can increase upon

binding to their targets.

FCS analysis. In FCS, the fluorescence autocorrelation function for free Brownian

diffusion of a single molecular species with multiple relaxation components is given by (57) :

522
$$G(\tau) = \frac{1}{N_{eff}}\left(1 + \frac{\tau}{\tau_d}\right)^{-1}\left(1 + \frac{\tau}{s^2\tau_d}\right)^{-0.5}\left(1 + a_t e^{\frac{-\tau}{\tau_t}} + \sum_i a_i e^{\frac{-\tau}{\tau_i}}\right)$$
(2)

In equation (3), N_{eff} is the average number of molecules in the detection volume, s is the

ratio between the axial and the lateral radii of the detection volume ($s = z_0/w_0$), and τ_d is the

average diffusion time, which is related to the diffusion coefficient ($w_0^2 = 4D\tau_d$) and to the

hydrodynamic radius R_H of the molecule via the Stokes-Einstein equation(58); a_t and τ_t are the

amplitude and lifetime of the triplet component and a_i and τ_i are the amplitude and the lifetime of

the i^{th} PET relaxation components, respectively. Global fitting was performed simultaneously on

each labelling position (C0, C14, C35, C73, C91, C121), a separate global fit was performed for

each for each measurement condition (NP 4E-BP2, 5P 4E-BP2 and NP 4E-BP2 + eIF4E), a total

of three independent global fits performed on six FCS curves simultaneously. In the global analysis

the diffusion time (τ_d), the triplet amplitude (a_t) and the triplet lifetime (τ_t) were shared

parameters, while the average number of molecules in the detection volume (N_{eff}), the PET

relaxation amplitudes (a_i) and lifetimes (τ_i) were fit as individual parameters. PET relaxation

components with amplitudes of 0.05 or less were considered insignificant and not reported. The

triplet lifetime (τ_t) were confined to a range of 5-10 µs based on characterization of the triplet state

of free Atto488 as a function of excitation intensity, see **Fig. S5**. FCS measurements performed

at ~5 kW/cm², returned a characteristic diffusion time (τ_d) for free Atto488 of 48 ± 1 µs. To

exclude the possibility that the slow relaxation component (τ_1) resulted from the diffusion of

unreacted free Atto488 remaining after size-exclusion chromatography purification, we performed

an additional reverse-phase high-performance liquid chromatography (RP-HPLC) purification of

542 the S91C NP 4E-BP2 construct. We did not detect unreacted Atto488 in the RP-HPLC

543 chromatogram (**Fig. S7**), we also observed the slow relaxation component (τ_1) in FCS

544 measurements of the S91C NP 4E-BP2 after RP-HPLC purification (**Fig. S8**). Fitting was

545 optimized by varying the number of relaxation components, the goodness of fit (χ^2) and Akaike

546 information criterion (AIC) were calculated for each fit. The addition of relaxation components

547 was accepted if: the χ^2 parameter decreased, the AIC decreased, and the fitting residuals were

548 featureless. The uncertainties of the fitted parameters were estimated using the Jacobian from

549 Levenberg-Marquardt least-squares fitting. Prior to each set of measurements, a calibration dye

550 (rhodamine 110) was used to estimate the *s* and w_0 parameters, typically ~8 and ~250 nm,

551 respectively.

552 **smFRET analysis.** A custom MATLAB script based on the 'MLT' algorithm was used to

553 identify fluorescence bursts and sort them into donor-only, acceptor-only and dual-labelled

554 (FRET) populations (32). The FRET efficiency for each burst was calculated based on the number

555 of detected photons in the donor (I_D) and acceptor (I_A) channels:

$$E = \frac{I_A}{I_A + \gamma I_D} \tag{3}$$

557 where γ is a correction factor for the difference in the detection efficiencies of donor and acceptor

558 channels, and the quantum yields of the dyes were determined as described in SI section 2.1. In

559 addition, corrections were applied on both I_D and I_A to subtract background, spectral cross talk, and

560 direct excitation of the acceptor. The background was obtained from a measurement of the sample

561 buffer while the corrections for cross talk and direct excitation were derived from donor-only and

562 acceptor-only bursts. A smFRET histogram was constructed from all bursts detected for a given

563 sample, each histogram was fit to a single Gaussian distribution.

To estimate the FRET efficiency if 4E-BP2 was a featureless statistical coil, 5000

conformers were generated using Trajectory Directed Ensemble Sampling (TraDES)(59) with

100% coil sampling in accordance with the sequences in **Table S1**. The FRET efficiency for each

conformer was calculated using the python library *LabelLib*(60) (see SI section **2.3**).

For each FRET construct, a family of τ_{DA}/τ *vs E* curves were generated based on a WLC

model with stiffness parameters κ between 0.01 and 1 using the *FRETlines* Python library(52).

The chain stiffness for given dataset was selected as the κ value of the τ_{DA}/τ *vs E* curve passing

through the centroid of a 2D Gaussian fit of the experimental τ_{DA}/τ *vs E* histogram. The confidence

interval was established from the range of WLC models that agree with the centroid from fitting

within the experimental accuracy of determining the FRET efficiency, i.e., 0.02 (32).

AUTHOR INFORMATION

Corresponding Authors

Email: claudiu.gradinaru@utoronto.ca

Author Contributions

 C.C.G. was responsible for overall project management and supervision. S.S. and Z.Z. expressed, purified and labelled proteins, performed measurements and analyzed data. A.B. and J.D. designed single- and double-cysteine 4E-BP2 mutants. T.T. performed smFRET simulations. S.S., Z.Z., and C.C.G. wrote the manuscript. J.D.F.-K. contributed to interpretation of results and critical revision of manuscript.

Competing Financial Interest

The authors report no competing financial interest.

ACKNOWLEDGEMENTS

This work has been supported by the Natural Sciences and Engineering Research Council of Canada (NSERC RGPIN 2017 – 06030 to C.C.G.), the Canadian Institutes of Health Research (CIHR FND-148375 to J.D.F.-K.) and National Institutes of Health (NIH 5R01GM127627-03 to J.D.F.-K.).

REFERENCES

1. P. E. Wright, H. J. Dyson, Intrinsically unstructured proteins: Re-assessing the protein structure-function paradigm. *Journal of Molecular Biology* **293**, 321-331 (1999).
2. V. N. Uversky, Intrinsically disordered proteins from A to Z. *The International Journal of Biochemistry & Cell Biology* **43**, 1090-1103 (2011).
3. P. E. Wright, H. J. Dyson, Intrinsically disordered proteins in cellular signalling and regulation. *Nature Reviews Molecular Cell Biology* **16**, 18-29 (2015).
4. C. Haynes *et al.*, Intrinsic disorder is a common feature of hub proteins from four eukaryotic interactomes. *PLoS Comput Biol* **2**, e100 (2006).
5. A. K. Dunker, I. Silman, V. N. Uversky, J. L. Sussman, Function and structure of inherently disordered proteins. *Current Opinion in Structural Biology* **18**, 756-764 (2008).
6. Julie D. Forman-Kay, T. Mittag, From Sequence and Forces to Structure, Function, and Evolution of Intrinsically Disordered Proteins. *Structure* **21**, 1492-1499 (2013).
7. V. N. Uversky, Natively unfolded proteins: A point where biology waits for physics. *Protein Science* **11**, 739-756 (2002).
8. J. A. Marsh, J. D. Forman-Kay, Sequence determinants of compaction in intrinsically disordered proteins. *Biophysical journal* **98**, 2383-2390 (2010).
9. A. H. Mao, S. L. Crick, A. Vitalis, C. L. Chicoine, R. V. Pappu, Net charge per residue modulates conformational ensembles of intrinsically disordered proteins. *Proceedings of the National Academy of Sciences* **107**, 8183-8188 (2010).
10. N. Sonenberg, A. G. Hinnebusch, Regulation of translation initiation in eukaryotes: mechanisms and biological targets. *Cell* **136**, 731-745 (2009).
11. S. Tait *et al.*, Local control of a disorder–order transition in 4E-BP1 underpins regulation of translation via eIF4E. *Proceedings of the National Academy of Sciences* **107**, 17627-17632 (2010).
12. A. Bah *et al.*, Folding of an intrinsically disordered protein by phosphorylation as a regulatory switch. *Nature* **519**, 106-109 (2015).
13. A. Lazaris-Karatzas, N. Sonenberg, The mRNA 5'cap-binding protein, eIF-4E, cooperates with v-myc or E1A in the transformation of primary rodent fibroblasts. *Molecular and cellular biology* **12**, 1234-1238 (1992).
14. A. Lazaris-Karatzas *et al.*, Ras mediates translation initiation factor 4E-induced malignant transformation. *Genes & development* **6**, 1631-1642 (1992).
15. A.-C. Gingras *et al.*, Regulation of 4E-BP1 phosphorylation: a novel two-step mechanism. *Genes & development* **13**, 1422-1437 (1999).
16. R. J. Dowling *et al.*, mTORC1-mediated cell proliferation, but not cell growth, controlled by the 4E-BPs. *Science* **328**, 1172-1176 (2010).
17. R. Colina *et al.*, Translational control of the innate immune response through IRF-7. *Nature* **452**, 323-328 (2008).
18. C. G. Gkogkas *et al.*, Autism-related deficits via dysregulated eIF4E-dependent translational control. *Nature* **493**, 371-377 (2013).
19. E. Klann, J. D. Sweatt, Altered protein synthesis is a trigger for long-term memory formation. *Neurobiology of learning and memory* **89**, 247-259 (2008).
20. S. Lukhele, A. Bah, H. Lin, N. Sonenberg, J. D. Forman-Kay, Interaction of the eukaryotic initiation factor 4E with 4E-BP2 at a dynamic bipartite interface. *Structure* **21**, 2186-2196 (2013).
21. D. Peter *et al.*, Molecular Architecture of 4E-BP Translational Inhibitors Bound to eIF4E. *Molecular Cell* **57**, 1074-1087 (2015).
22. G. J. Brunn *et al.*, Phosphorylation of the translational repressor PHAS-I by the mammalian target of rapamycin. *Science* **277**, 99-101 (1997).
23. J. E. Dawson *et al.*, Non-cooperative 4E-BP2 folding with exchange between eIF4E-binding and binding-incompatible states tunes cap-dependent translation inhibition. *Nat Commun* **11**, 3146 (2020).

24. G.-N. Gomes, C. C. Gradinaru, Insights into the conformations and dynamics of intrinsically disordered proteins using single-molecule fluorescence. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* **1865**, 1696-1706 (2017).

25. L. A. Metskas, E. Rhoades, Single-molecule FRET of intrinsically disordered proteins. *Annual review of physical chemistry* **71**, 391-414 (2020).

26. V. N. Uversky, "Biophysical Methods to Investigate Intrinsically Disordered Proteins: Avoiding an "Elephant and Blind Men" Situation" in Intrinsically Disordered Proteins Studied by NMR Spectroscopy. (Springer, 2015), pp. 215-260.

27. G. F. Schröder, U. Alexiev, H. Grubmüller, Simulation of fluorescence anisotropy experiments: probing protein dynamics. *Biophysical journal* **89**, 3757-3770 (2005).

28. S. Milles, E. A. Lemke, Mapping Multivalency and Differential Affinities within Large Intrinsically Disordered Protein Complexes with Segmental Motion Analysis. *Angewandte Chemie International Edition* **53**, 7364-7367 (2014).

29. N. Jain, M. Bhattacharya, S. Mukhopadhyay, Chain collapse of an amyloidogenic intrinsically disordered protein. *Biophysical journal* **101**, 1720-1729 (2011).

30. J. R. Lakowicz, *Principles of fluorescence spectroscopy* (Springer, New York, N.Y., ed. 3rd, 2006).

31. Z. Zhang, D. Yomo, C. Gradinaru, Choosing the right fluorophore for single-molecule fluorescence studies in a lipid environment. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1859**, 1242-1253 (2017).

32. G.-N. W. Gomes *et al.*, Conformational Ensembles of an Intrinsically Disordered Protein Consistent with NMR, SAXS, and Single-Molecule FRET. *Journal of the American Chemical Society* **142**, 15697-15710 (2020).

33. F. Huang, W. M. Nau, A conformational flexibility scale for amino acids in peptides. *Angewandte Chemie* **115**, 2371-2374 (2003).

34. Keum *et al.*, A conserved motif within the flexible C-terminus of the translational regulator 4E-BP is required for tight binding to the mRNA cap-binding protein eIF4E. *Biochemical Journal* **441**, 237-245 (2012).

35. A. P. De Silva, T. S. Moody, G. D. Wright, Fluorescent PET (Photoinduced Electron Transfer) sensors as potent analytical tools. *Analyst* **134**, 2385-2393 (2009).

36. S. Doose, H. Neuweiler, M. Sauer, Fluorescence quenching by photoinduced electron transfer: a reporter for conformational dynamics of macromolecules. *ChemPhysChem* **10**, 1389-1398 (2009).

37. H. Chen, E. Rhoades, J. S. Butler, S. N. Loh, W. W. Webb, Dynamics of equilibrium structural fluctuations of apomyoglobin measured by fluorescence correlation spectroscopy. *Proceedings of the National Academy of Sciences* **104**, 10459-10464 (2007).

38. S. Mukhopadhyay, R. Krishnan, E. A. Lemke, S. Lindquist, A. A. Deniz, A natively unfolded yeast prion monomer adopts an ensemble of collapsed and rapidly fluctuating structures. *Proceedings of the National Academy of Sciences* **104**, 2649-2654 (2007).

39. H. Blom, A. Chmyrov, K. Hassler, L. M. Davis, J. Widengren, Triplet-State Investigations of Fluorescent Dyes at Dielectric Interfaces Using Total Internal Reflection Fluorescence Correlation Spectroscopy. *The Journal of Physical Chemistry A* **113**, 5554-5566 (2009).

40. L. J. Lapidus, W. A. Eaton, J. Hofrichter, Measuring the rate of intramolecular contact formation in polypeptides. *Proceedings of the National Academy of Sciences* **97**, 7220-7225 (2000).

41. F. Krieger, B. Fierz, O. Bieri, M. Drewello, T. Kiefhaber, Dynamics of unfolded polypeptide chains as model for the earliest steps in protein folding. *Journal of molecular biology* **332**, 265-274 (2003).

42. J. K. Lum, H. Neuweiler, A. R. Fersht, Long-range modulation of chain motions within the intrinsically disordered transactivation domain of tumor suppressor p53. *Journal of the American Chemical Society* **134**, 1617-1622 (2012).

43. R. R. Goluguri, S. Sen, J. Udgaonkar, Microsecond sub-domain motions and the folding and misfolding of the mouse prion protein. *eLife* **8**, e44766 (2019).

44. S. Sen, H. Kumar, J. B. Udgaonkar, Microsecond Dynamics During the Binding-induced Folding of an Intrinsically Disordered Protein. *Journal of Molecular Biology* **433**, 167254 (2021).

704   45.   S. A. Waldauer, O. Bakajin, L. J. Lapidus, Extremely slow intramolecular diffusion in unfolded
705   protein L. *Proceedings of the National Academy of Sciences* **107**, 13713-13717 (2010).

706   46.   E. Sisamakis, A. Valeri, S. Kalinin, P. J. Rothwell, C. A. Seidel, Accurate single-molecule FRET
707   studies using multiparameter fluorescence detection. *Methods in enzymology* **475**, 455-514 (2010).

708   47.   S. Kalinin *et al.*, A toolkit and benchmark study for FRET-restrained high-precision structural
709   modeling. *Nature Methods* **9**, 1218-1225 (2012).

710   48.   B. Schuler, W. A. Eaton, Protein folding studied by single-molecule FRET. *Current opinion in
711   structural biology* **18**, 16-26 (2008).

712   49.   A. Borgia, P. M. Williams, J. Clarke, Single-molecule studies of protein folding. *Annu. Rev.
713   Biochem.* **77**, 101-125 (2008).

714   50.   I. V. Gopich, A. Szabo, Theory of the energy transfer efficiency and fluorescence lifetime
715   distribution in single-molecule FRET. *Proceedings of the National Academy of Sciences* **109**, 7747-
716   7752 (2012).

717   51.   B. Schuler, Perspective: Chain dynamics of unfolded and intrinsically disordered proteins from
718   nanosecond fluorescence correlation spectroscopy combined with single-molecule FRET. *The
719   Journal of Chemical Physics* **149**, 010901 (2018).

720   52.   A. Barth *et al.*, Unraveling multi-state molecular dynamics in single-molecule FRET experiments-
721   Part I: Theory of FRET-Lines. arxiv:2107.14770 (2021).

722   53.   H. Hofmann *et al.*, Polymer scaling laws of unfolded and intrinsically disordered proteins
723   quantified with single-molecule spectroscopy. *Proceedings of the National Academy of Sciences*
724   **109**, 16155-16160 (2012).

725   54.   C. Igreja, D. Peter, C. Weiler, E. Izaurralde, 4E-BPs require non-canonical 4E-binding motifs and
726   a lateral surface of eIF4E to repress translation. *Nature communications* **5**, 1-14 (2014).

727   55.   Y. Li, R. V. Shivnaraine, F. Huang, J. W. Wells, C. C. Gradinaru, Ligand-Induced Coupling
728   between Oligomers of the M_2 Receptor and the G_{i1} Protein in Live Cells. *Biophysical Journal* **115**,
729   881-895 (2018).

730   56.   D. Badali, C. C. Gradinaru, The effect of Brownian motion of fluorescent probes on measuring
731   nanoscale distances by Förster resonance energy transfer. *The Journal of Chemical Physics* **134**,
732   225102 (2011).

733   57.   E. Haustein, P. Schwille, Fluorescence Correlation Spectroscopy: Novel Variations of an
734   Established Technique. *Annual Review of Biophysics and Biomolecular Structure* **36**, 151-169
735   (2007).

736   58.   A. Mazouchi, B. Liu, A. Bahram, C. C. Gradinaru, On the performance of bioanalytical
737   fluorescence correlation spectroscopy measurements in a multiparameter photon-counting
738   microscope. *Anal Chim Acta* **688**, 61-69 (2011).

739   59.   H. J. Feldman, C. W. Hogue, Probabilistic sampling of protein conformations: new hope for brute
740   force? *Proteins: Structure, Function, and Bioinformatics* **46**, 8-23 (2002).

741   60.   M. Dimura *et al.*, Quantitative FRET studies and integrative modeling unravel the structure and
742   dynamics of biomolecular systems. *Current opinion in structural biology* **40**, 163-185 (2016).