Culture and genomic analysis of *Symbiopectobacterium purcellii*, gen.nov. sp. nov.,

isolated from the leafhopper *Empoasca decipiens*

Pol Nadal-Jimenez[1,*], Stefanos Siozios[1], Nigel Halliday[2], Miguel Cámara[2] & Gregory

D.D. Hurst[1]

Author affiliations: [1]Institute of Infection, Veterinary and Ecological Sciences,

University of Liverpool, Liverpool, UK; [2]The National Biofilms Innovation Centre,

School of Life Sciences, University of Nottingham Biodiscovery Institute, University of

Nottingham, UK.

*Correspondence: Pol Nadal-Jimenez, University of Liverpool, polnadal@gmail.com

ABSTRACT

Bacterial endosymbionts are found in multiple arthropod species, where they

play crucial roles as nutritional symbionts, defensive symbionts or reproductive

parasites. Recent work has highlighted a new clade of heritable microbes within the

gammaproteobacteria that enter into both obligate and facultative symbioses, with

an obligately required unculturable symbiont recently given the name *Cand.*

Symbiopectobacterium. In this study, we describe a culturable rod shaped non-

flagellated bacterial symbiont from this clade isolated from the leafhopper

Empoasca decipiens. The symbiont is related to the transovarially-transmitted 'BEV'

bacterium that was first isolated from the leafhopper *Euscelidius variegatus* by

Alexander Purcell*,* and we therefore name the symbiont *Symbiopectobacterium*

purcellii sp. nov. gen. nov. We further report the closed genome sequence for *S.*

purcellii. The genome is atypical for a heritable microbe, being large in size, without

profound AT bias and with little evidence of pseudogenization. The genome is

predicted to encode Type II, III and VI secretion systems and associated effectors and

a non-ribosomal peptide synthase array likely to produce bioactive small molecules.

Predicted metabolism is more complete than for other symbionts in the

Symbiopectobacterium clade, and the microbe is predicted to synthesize a range of B

vitamins. However, Biolog plate analysis indicate metabolism is depauperate

compared to the sister clade, represented by *Pectobacterium carotovorum*. A

quorum-sensing pathway related to that of *Pectobacterium* spp. (containing an

overlapping *expI-expR1* pair in opposite directions and a "solo" *expR2*) is evidenced,

and LC-MS/MS analysis reveals the presence of 3-hydroxy-C10-HSL as the sole *N*-

acylhomoserine lactone (AHL) in our strain. This AHL profile is profoundly divergent

from that of other *Erwinia* and *Pectobacterium* spp., which produce mostly 3-oxo-C6-

and 3-oxo-C8-HSL and could aid group identification. Thus, this microbe denotes one

that has lost certain pathways associated with a saprophytic lifestyle but represents

an important baseline against which to compare other members of the genus

Symbiopectobacterium that show more profound integration into host biology.

KEYWORDS

Symbiopectobacterium, *Empoasca*, symbiosis, leafhopper, quorum sensing

INTRODUCTION

It is now understood that microbes influence multiple aspects of animal biology (1). Symbiont contributions extend from involvement in the process of digestion in the gut, through anabolic activities and the supply of vitamins and amino acids, to protection against natural enemies and defence against prey/hosts (2). Conversely, other symbiotic microbes are pathogenic or parasitic, and many symbioses combine both parasitic and beneficial aspects. Levels of symbiont integration vary between symbioses (3). On the host axis, they vary from facultative relationships where the host does not require a particular symbiont, to obligate where the individual dies or becomes sterile in the absence of symbiosis. Likewise, symbionts vary in the degree to which they rely on a host – some only replicating within hosts with others having environmental replication. The process of symbiosis formation also varies – from arising within the host lifecycle through acquisition by the host or infection by the microbe, to being present through the host lifecycle, with symbiont transfer/transmission from parent to offspring.

Whilst arthropod-microbe symbioses are diverse in terms of the microbial partners, particular microbial taxa have established symbiosis with a number of host species, commonly establishing in new host species through a host switch event. Well-known 'heritable symbionts' found over a broad range of arthropods include *Wolbachia*, *Rickettsia*, *Spiroplasma*, *Cardinium* and *Arsenophonus* (4). The interactions found in these symbioses include obligate and facultative associations, and ones which are beneficial, parasitic or have a combination of features.

Recent research has added a new clade of insect symbionts, *Cand.* Symbiopectobacterium, to 'the big five' (5). The first member of this clade to be described was the BEV strain – an acronym for bacterium from *Euscelidius variegatus*, a planthopper host species. This strain was cultured (6), but never formally named. The symbiosis was characterized as one with vertical transmission, where the host's reproduction was negatively impacted by the microbe. In addition, there was also transmission to other insects on the plant – thus establishing the symbiosis as a pathogenic one maintained through mixed modes of transmission (7).

Experiments also suggested the symbiont facilitated the transmission of

phytoplasma from its bug host to plant (8). Later, the pest species *Cimex lectularius*

(common bedbug) was observed to carry a heritable symbiont related to BEV (9).

This symbiont has not been established in cell-free culture, and symbiosis is

facultative from the host perspective: the bedbug does not require the symbiont.

Following this, a third hemipteran – the bulrush bug *Chilacis* was observed to carry a

related vertically transmitted symbiont housed in a gut mycetome, in what appears

to be an obligate association, in which the host requires the symbiont (10). More

recently, symbioses involving members of this clade have extended beyond

Hemiptera hosts to nematodes, with Martinson et al (5) characterizing symbionts

related to BEV as obligate partners of *Howardula* nematodes. They named this

microbe *Cand.* Symbiopectobacterium, reflecting its symbiotic lifestyle and its sister

relationship to the well-characterized genus *Pectobacterium*.

Cand. Symbiopectobacterium has thus emerged as a potentially widespread

and significant symbiotic associate of invertebrates. The original culturable BEV

isolate, on which the genus could be formally described, was lost and genomic

information for this strain is partial (11). Recovering a model culturable member of

the genus is important, as it allows formal description of the microbe, completion of

a closed genome sequence against which reductive evolution in symbiosis can be

measured and presents a system in which gene function may be investigated. In this

paper, we report the isolation to pure culture of a member of this clade from the

planthopper *Empoasca decipiens*. We further present and analyse the complete

genome sequence of this microbe, assess its growth requirements compared to

Pectobacterium carotovorum and analyse its quorum sensing-signalling system.

MATERIALS AND METHODS

Symbiont isolation, morphology *in vitro* and identification through 16S rRNA sequence

Initial Cicadellidae samples with light green coloration were collected in Prince's Park, Liverpool in April 2018, scooping different plants with an insect net at a maximum height of 2m. Fresh specimens were transported alive to the lab and sacrificed by freezing at -20ºC for 15 min. The insect specimens were surface sterilized by immersion in 70% ethanol and washed with sterile water to remove the remaining alcohol. Insect legs were excised with a sterile surgical blade and stored at -20ºC for *post hoc* host species determination through DNA barcoding.

The remainder of the insect body was mechanically crushed and resuspended in 100 μl of sterile water. An aliquot of 10 μl was plated on brain heart infusion (BHI, Oxoid, UK) agar and grown at 30ºC for 6 d to allow the appearance of slow-growing bacterial colonies. Morphology was examined through Gram staining and scanning electron microscopy of overnight culture. To identify the bacterial species, we performed colony PCR of the 16S rRNA gene colonies emerging on the agar plates with primers 27F (AGAGTTTGATCMTGGCTCAG) (12) and 1492R(I) (GGTTACCTTGTTACGACTT) (13), and sequenced by Eurofins genomics, Germany. Sequences were manually curated and phylogenetic analysis performed based on the 16S rRNA gene sequences which included a large assemblage of members from the *Cand*. Symbiopectobacterium clade. To this end, 16S rRNA sequences were aligned using the SSU-ALIGN software (14). A Bayesian phylogeny was estimated with MrBayes v3.2.6 (15) by sampling across the GTR model space (nst=mixed, rates=gamma). Two independent runs were performed for 5,000,000 generations and sub-sampling every 500 generations using four Markov chains. The first 25% of the samples were discarded as burn-in.

The leafhopper host was identified through DNA barcoding using the COI sequence. To this end, insect legs were mechanically crushed and resuspended in 50 μl of sterile water and the genomic DNA (gDNA) was extracted using a Quick-DNA Universal kit (Zymo research, USA). 2 μl of the gDNA were added to a GoTaq® Green

Master Mix (Promega, USA) and used to amplify part of the mitochondrial

cytochrome oxidase 1 (CO1) with primers C1-J-1718

(GGAGGATTTGGAAATTGATTAGTTCC) and C1-N-2191

(CCCGGTAAAATTAAAATATAAACTTC) (16). The PCR program consisted of an initial

denaturation step at 95ºC for 5 min, followed by 30 cycles of DNA denaturation at

94ºC for 15s, primer annealing at 55ºC for 45 seconds, and primer extension at 72ºC

for 1 min. A final extension was carried out at 72ºC for 5 min. A few microliters of

each PCR product were run on an agarose gel to assess the success of the PCR

reaction and the remains cleaned through an Isolate II PCR and Gel kit (Bioline, USA)

and sent for sequencing with primer C1-N-2191. Identity was checked through

analysis against the Barcode of Life Database, BOLD.

In vitro Growth requirements

BIOLOG GEN III plates (Cat. No. 1030) were used to ascertain the

physiological and biochemical characteristics of *S. purcellii* SyEd1 *in vitro*, and these

were conducted alongside *Pectobacterium carotovorum* subsp. *carotovorum* LMG

02404^T for comparison. Within this, we also performed the assay in the

presence/absence of 0.4 % polygalacturonic acid PGA (Sigma, P3850), which is

commonly used to induce the expression of plant cell wall-degrading enzymes (for

preparation of PGA, see (17). For the BIOLOG GEN III assays, we used IF-A inoculating

fluid (Biolog, Cat. No. 72401) with or without PGA supplementation to a final

concentration of 0.4% PGA. Both bacterial species were grown overnight, diluted to

an OD_{600}= 0.4 and 50 μl of this aliquot were added to a tube containing IF-A fluid.

The aliquot in the IF-A tube was homogeneously mixed using a vortex and 100 μl of

this suspension was added to each of the 96 wells of the BIOLOG GEN III plate. The

plate was subsequently incubated at 30ºC without shaking.

Potato infection assays

Pectobacterium spp. are well-known plant pathogens causing soft-rot disease

in several plants including potatoes, carrots and cabbages. This damage is caused by

a series of secreted enzymes (cellulases, proteases, pectate lyases (Pel), pectin

lyases, and polygalacturonases) commonly referred to as plant cell wall-degrading

enzymes (PCWDEs). The presence of 15 putative PCWDEs and two copies of the KdgR

regulator (associated to their expression) in *Symbiopectobacterium purcellii* led us to

assess whether this symbiont retains the plant pathogenic activity of its sister clade,

Pectobacterium. To this aim, virulence was tested in potatoes, using a method

previously described in Nadal-Jimenez et al (18) with minor modifications. Briefly,

potatoes were bought at local stores, washed with tap water, dried and surface

sterilized with 70% ethanol. Slices about 0.5 cm thick were placed in sterile Petri

dishes. Overnight cultures of *S. purcellii* SyEd1 and *Pectobacterium carotovorum* LMG

02404^T were diluted to an OD600=0.4, and 20 µl were placed at the centre of the

potato slice. The same amount of sterile BHI medium was added to the negative

controls. The plates were sealed with parafilm to avoid moisture loss and incubated

at 25ºC in dark conditions. Tissue maceration was assessed visually 24, 48 and 72 h

after incubation.

Symbiont Genome sequencing, assembly and annotation

The genome of the symbiont was completed using a combination of short

(Illumina) and long (nanopore) reads by MicrobesNG (Birmingham, UK) using their

enhanced genome service. Briefly, Illumina sequencing was performed using the

Nextera XT library prep protocol on a HiSeq platform (Illumina, San Diego, CA, USA)

using a 250bp paired end protocol. Reads were adapter trimmed using Trimmomatic

0.30, with a sliding window quality cutoff of Q15 (19). Long read genomic DNA

libraries are prepared with Oxford Nanopore SQK-RBK004 kit (ONT, UK) using 400-

500ng of HMW DNA and sequenced in a FLO-MIN106 (R.9.4.1) flow cell in a GridION

(ONT, UK). Hybrid genome assembly of both short and long reads was performed

using Unicycler version 0.4.0 under the normal mode (20). The final assembly was

manually inspected for potential misassemblies by mapping the raw reads back to it.

Genome annotation was performed with the NCBI Prokaryotic Genome Annotation

Pipeline (21). Metabolic and functional assessment of the symbiont genome was

conducted using the Kyoto Encyclopaedia of Genes and Genomes (KEGG) database

(22). Identification of secondary metabolite biosynthesis gene clusters was

performed using the antiSMASH server (23). Finally, prophage regions were

predicted using the PHAge Search Tool Enhanced Release (PHASTER) web server

(24).

Phylogenomic analysis

The phylogenetic position of the symbiont was assessed based on the

concatenated analysis of 527 single copy core proteins identified among 56 publicly

available genomes. These include members of the closely related genera

Pectobacterium, *Brenneria*, *Dickeya*, *Lonsdalea*, *Sodalis* and the recently

characterized *Cand*. Symbiopectobacterium (5). Single copy orthologue protein

sequences were identified using OrthoFinder v2.3.11 (25). A maximum likelihood

phylogeny was inferred with IQ-TREE 2.0.3 [25] using the JTT+F+R3 substitution

model selected using ModelFinder according to the Bayesian information criterion

[26]. Clade support was assessed based on 1000 ultrafast bootstrap replicates [27].

Analysis of *N*-acyl homoserine lactone synthesis

S. purcellii SyEd1 cultures were grown in 5 ml of BHI medium at 30ºC for 24h

and 200 rpm. After incubation the cultures were centrifuged and the supernatant

collected and filtered through a 0.2 mm filter (SLGP033RS, Millipore). 500 µl of

acidified ethyl acetate was added to 1 ml of supernatant sample and the mixture was

vortexed for 1-2 min. Subsequently. The mixture was centrifuged for 1 min to allow

the formation of a clear interface between the aqueous and organic layer. The

organic (upper) layer was transferred using a pipette (without disturbing the

aqueous layer) to a new 2ml Eppendorf. The extraction process was repeated twice

more, combining the extracts for each sample into one of approximately 1.5 ml

extract. Upon completion, the samples were dried under vacuum in a centrifugal

evaporator.

Dried extract samples were reconstituted in 50 µl of methanol (MeOH) prior

to analysis. LC-MS/MS analysis of 5 µl sample injections were conducted using a

Qtrap 6500 hybrid triple-quadrupole linear ion trap mass spectrometer in tandem

with an Exion LC system (Sciex). The overall method was a modification of that

described by Ortori et al (26). Chromatography was achieved using a Phenomenex

Gemini C18 column (3.0 um, 50 x 3.0 mm) with a constant flow rate of 450 μl/min of

mobile phase A (0.1 % (v/v) formic acid) and mobile phase B (0.1 % (v/v) formic acid

in methanol). The LC gradient began at 10% B for 1.0 min, increased linearly to 50% B

over 0.5 min, then to 99% B over 4.0 min. The composition remained at 99% B for

1.5 min, decreased to 10% B over 0.1 min, and stayed at this composition for 2.9

min. Analyte detection was conducted with the MS operating in MRM (multiple

reaction monitoring) mode, screening the LC eluent for specific AHLs (unsubstituted,

3-oxo and 3-OH AHLs with even chain lengths from C4-C14).

Prevalence of symbiont in *Empoasca decipiens* leafhoppers

In order to assess the prevalence of this bacterium in *E. decipiens*, we performed a

PCR screening on various specimens. Additional insect collections were completed at

the same location in August 2019 and tested for SyEd1 by PCR assay. Using the full-

genome sequence of our cultured strain, we developed two set of specific PCR

primers to amplify part of the DNA gyrase subunit (*gyrB*) gene of this bacterium:

BEV_gyrB_F1 (CCGTGGTGTCGGTGAAAGTA) + BEV_gyrB_R1

(TGGTCTTCTGTCAGCGTGTC) and BEV_gyrB_F2 (CTCGTGAAATGACACGACGC) +

BEV_gyrB_R2 (CAGCAGTTCCACTTGTTCGC). The gDNA was extracted in the same

manner as for the leg samples and used as a template for the PCR reactions.

RESULTS

Symbiont isolation and identification

The bacterium grows under standard aerobic conditions in Brain heart

infusion (BHI) medium (CM1032, Oxoid), forming circular white colonies approx. 2-3

mm in diameter on BHI agar, and cultures emitted a pronounced plant-like odour.

The bacterium is Gram negative, and SEM revealed it to be a non-flagellated rod

shape, of length 1-1.5 mm (Figure 1). The bacterium will also grow in LB (Miller)

(110285, Millipore/Merck KGaA) although at a slower rate, and growth is inhibited

by light.

Phylogenetic analysis based on the 16S rRNA gene (Accession number
OK044380) placed the isolated microbe well within the recently characterized clade
Cand. *Symbiopectobacterium* (Figure 2), a group of microbes commonly associated
with arthropods and nematodes. Sequence of the CO-1 amplicons from the insect
host revealed the leafhopper host to be *Empoasca decipiens* (Hemiptera,
Cicadellidae), a common species of leafhopper in Europe. *Empoasca decipiens* has
been implicated in the transmission of various plant pathogens (27), and is
considered a pest in various crops (28).

In vitro Growth requirements

S. purcellii SyEd1 and *P. carotovorum* LMG 2404^T were grown at 30ºC in
BIOLOG GEN III plates. For *P. carotovorum* LMG 2404^T, the presence of the purple
tetrazolium dye as a result of growth and respiration in the wells where the strains
had grown was visible after 24 h, while the wells that did not supported the growth
of this strain remained colourless. In the case of *S. purcelliii* SyEd1, the plates had to
be incubated for a total of 72h to allow bacterial growth. This is not surprising since,
in our hands, *S. purcellii* SyEd1 grows slowly in BHI media (requiring up to 48h), and
even slower in less rich media. Analyses indicated *S. purcellii* was considerably more
fastidious than the comparator outgroup strain *P. carotovorum* LMG 2404^T in terms
of metabolites that supported growth (Table 1) but had broader resistance to
xenobiotics than this strain (Table 2). Growth conditions for *S. purcellii* on Biolog
analysis was only modestly altered by addition of PGA.

Potato infection assays

Potato slices infected with *S. purcellii* SyEd1 exhibited a complete absence of
infection/ tissue maceration at the different time points tested (72h time point
shown in Figure 3) in contrast to *P. carotovorum* LMG 02404^T, used as positive
control for infection. The assay was maintained for a week without any sign of
infection being visible in *S. purcellii* SyEd1 infected potatoes.

Genome sequence and assembly

The genome of the symbiont presented as a single circular chromosome of

circa 4.9 MB with an average GC content of 52.5% (Table 3). No plasmids were

identified. The complete predicted gene set consists of 4,494 protein-coding genes

(including 312 predicted pseudogenes), 7 ribosomal RNA operons (5S, 16S, 23S) and

76 tRNAs. The average length of the protein-coding genes is 948 bp accounting for a

coding density of about 86.2%. Pseudogenization rates were estimated to be circa

7% (312 predicted pseudogenes). The main chromosome was predicted to contain

six intact prophage regions and three additional incomplete fragments. The

complete genome assembly and the raw reads have been submitted to the

DDBJ/EMBL/GenBank database under the BioProject accession number

PRJNA756769 (genome accession number CP081864).

Phylogenomic and functional analysis

To confirm the phylogenetic position of the *E. decipiens* symbiont we conducted a

phylogenomic analysis base on the concatenated set of 527 single copy orthologue

proteins across 56 related genomes (Figure 4). These results further support the

placement of the symbiont in the *Cand.* Symbiopectobacterium clade.

The genome is predicted to encode type II, III and VI secretion systems alongside a

wide array of predicted secreted toxins, compatible with its likely status as a

symbiont of its insect host. Anti-SMASH predicted five genomic regions associated

with small molecule production. Notable amongst these is a non-ribosomal peptide

synthase (NRPS) region. It is unclear if the NRPS system produces siderophore

molecules that permit growth in iron poor host environments, or antimicrobial

compounds. In addition, anti-SMASH predicted putative gene clusters for the

biosynthesis of thiopeptide, an aryl-polyene potentially providing defence against

ROS, and betalactone synthesis. There is also a predicted homoserine lactone

synthesis island *expI/expR1* that may be involved in sensing of microbial titre (see

below); the genome encodes additional conserved elements of the Quorum sensing

system, *expR2, gacA (expA), gacS (expS), rsmA* and *rsmB,* and *kdgR*. Finally, the

genome encodes complete biosynthetic pathways for several B vitamins including

thiamine (B1), riboflavin (B2), pantothenate (B5), biotin (B7), pyridoxine (B6) and

folate (B9). A broad array of complete amino acid biosynthesis pathways was also
observed, including serine, threonine, cysteine, methionine, valine, leucine,
isoleucine, arginine, ornithine, arginine, proline, histidine, tryptophan, phenylalanine
and tyrosine. Vitamin and amino acid provision are common means through which
symbionts contribute to host function. There are also 5 *pel* genes predicted to
encoded pectate lyase enzymes. The failure of the strain to utilize pectin on Biolog
plates or on potato tubers may thus be context dependent.

Analysis of homoserine lactones

S. purcellii SyEd1 analysis using LC-MS/MS revealed the presence of a single
AHL that was characterised as *N*-(3-hydroxydecanoyl)-L-homoserine lactone (3-OH-
C10-HSL). Figure 5 shows the LC-MS/MS chromatogram obtained from the *S. purcellii*
SyEd1 sample compared to the 3-OH-C10-HSL standard and the uncultured BHI
medium. Members of the genus *Pectobacterium* have been reported to produce 3-
oxo-C6-HSL, 3-oxo-C8-HSL, C10-HSL (29), but, to the best of our knowledge, the
presence of 3-OH-C10-HSL as the sole AHL in *S. purcellii* is unreported in related
genera. This trait may help to identify novel members of this genus that we presume
may have been previously misidentified as *Pectobacterium* spp. associated to the
plants where the leafhoppers feed.

Prevalence of *S. purcellii* in *E. decipiens* samples.

Seven new *E. decipiens* specimens were collected and their identity
confirmed by CO-1 amplification and sequencing. All samples were confirmed to be
E. decipiens with >99.6% identity with previously deposited sequences in NCBI. The
same gDNA extract was used to screen for the presence of the bacterial symbiont by
PCR using our BEV_gyrB_F2 and BEV_gyrB_R2 primers. All seven samples produced
an amplicon for *S. purcellii* (10), and the identity of the amplicon was confirmed
through sequencing accounting for a 100% prevalence in the population tested
(95%CI: 64% - 100%).

Description of *Symbiopectobacterium purcelli gen. nov., sp. nov.*

Symbiopectobacterium purcellii . *Symbiopectobacterium* (L. n. *sym bio pecto*

bacterium) references the related *Cand.* Symbiopectobacterium that is an obligate symbiont

of nematode worms, this name reflecting the symbiotic habit of the microbe, and the

relationship of the genus as sister to *Pectobacterium*. The species name *purcellii* [pur.cell ii.

L. m. gen.] is given in reference to Alexander Purcell, who isolated the first member of this

clade, which he named the BEV symbiont (bacterium from *Euscelidius variegatus*).

Gram-negative rod-shaped bacterium. Grows optimally at 30 °C in BHI medium in

the dark forming colonies within 24-48 h. Using Biolog GENIII plates, *S. purcelli* responded

positively to the following carbon sources: D-glucose, D-mannose, D-fructose, glycerol, N-

acetyl glucosamine, L-rhamnose, and weaker to methyl pyruvate and acetic acid. Growth

was inhibited at pH5, by 4% and 8% NaCl, by D-serine, minocycline and sodium bromate.

Growth was not impaired by 1% sodium lactate, fusidic acid, Troleandomycin, Rifamycin S,

Lincomycin, Guanidine HCl, Vancomycin, Tetrazolium Violet, Tetrazolium blue, Potassium

tellurite, Nalidixic Acid, Lithium Chloride, Aztreonam, Sodium Butyrate. The microbe does

not cause macerations on potato slices.

Symbiopectobacterium purcellii gen. nov. sp. nov. form a cluster with a variety of

uncultured symbionts of insects and nematodes, as well as the previously cultured strain

BEV.

The type strain is SyEd1 (LMG 32449, CECT 30436) and was isolated from *Empoasca*

decipiens (Hexapoda: Hemiptera: Cicadellidae) from Liverpool UK (53.3868° N, 2.9565° W).

The genome consists of a single circular chromosome of size 4.9MB and DNA G+C content is

52.5 mol%. The 16S rRNA sequence of the type strain is available at accession OK044380.

The complete genome assembly and the raw reads have been submitted to the

DDBJ/EMBL/GenBank database under the BioProject accession number PRJNA756769

(genome accession number CP081864).

Acknowledgements - We are thankful to Alison Beckett (University of Liverpool, UK)

for SEM Electron microscopy imaging service; Sam Edwards (University of

Copenhagen, DK) for assistance in *E. decipiens* collection; Nigel Gotts (University of

Liverpool, UK) for assistance evaporating the AHL-containing extracts; and Rita

Valente (Instituto Gulbenkian de Ciência, PT) for useful discussions about the

Pectobacterium virulence assays. This work was funded by a BBSRC grant to GH

(grant BB/S017534/1). Miguel Cámara is partly funded by the National Biofilms
Innovation Centre (NBIC) which is an Innovation and Knowledge Centre funded by
the Biotechnology and Biological Sciences Research Council, Innovate UK and
Hartree Centre (Award Number BB/R012415/1).

Conflicts of interest

The authors declare that there are no conflicts of interest.

REFERENCES

1.   McFall-Ngai M, Hadfield MG, Bosch TCG, Carey H V., Domazet-Lošo T, Douglas
AE, et al. Animals in a bacterial world, a new imperative for the life sciences.
Vol. 110, Proceedings of the National Academy of Sciences of the United
States of America. 2013. p. 3229–36.
2.   Hurst GDD. Extended genomes: Symbiosis and evolution. Interface Focus
[Internet]. 2017 [cited 2021 Sep 22];7(5). Available from:
https://royalsocietypublishing.org/doi/abs/10.1098/rsfs.2017.0001
3.   Gerardo N, Hurst G. Q&A: Friends (but sometimes foes) within: The complex
evolutionary ecology of symbioses between host and microbes. BMC Biol
[Internet]. 2017 Dec 27 [cited 2021 Sep 22];15(1):1–6. Available from:
https://bmcbiol.biomedcentral.com/articles/10.1186/s12915-017-0455-6
4.   Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstädter J, et al. The
diversity of reproductive parasites among arthropods: Wolbachia do not walk
alone. BMC Biol. 2008;6.
5.   Martinson VG, Gawryluk RMR, Gowen BE, Curtis CI, Jaenike J, Perlman SJ.
Multiple origins of obligate nematode and insect symbionts by a clade of
bacteria closely related to plant pathogens. Proc Natl Acad Sci U S A [Internet].
2020 Dec 15 [cited 2021 Sep 22];117(50):31979–86. Available from:
https://www.pnas.org/content/117/50/31979
6.   Purcell AH, Steiner T, Mégraud F, Bové J. In vitro isolation of a transovarially
transmitted bacterium from the leafhopper Euscelidius variegatus (Hemiptera:
Cicadellidae). J Invertebr Pathol. 1986;48(1):66–73.
7.   Purcell AH, Suslow KG, Klein M. Transmission via plants of an insect
pathogenic bacterium that does not multiply or move in plants. Microb Ecol
[Internet]. 1994 Jan [cited 2021 Sep 22];27(1):19–26. Available from:
https://link.springer.com/article/10.1007/BF00170111
8.   Purcell AH, Suslow KG. Pathogenicity and effects on transmission of a
mycoplasmalike organism of a transovarially infective bacterium on the
leafhopper Euscelidius variegatus (Homoptera: Cicadellidae). J Invertebr
Pathol. 1987;50(3):285–90.
9.   Hypša V, Aksoy S. Phylogenetic characterization of two transovarially
transmitted endosymbionts of the bedbug Cimex lectularius (Heteroptera:

Cimicidae). Insect Mol Biol [Internet]. 1997 [cited 2021 Sep 22];6(3):301–4.
Available from: https://pubmed.ncbi.nlm.nih.gov/9272448/

10. Kuechler SM, Dettner K, Kehl S. Characterization of an obligate intracellular
bacterium in the midgut epithelium of the bulrush bug Chilacis typhae
(Heteroptera, Lygaeidae, Artheneinae). Appl Environ Microbiol.
2011;77(9):2869–76.

11. Degnan PH, Bittleston LS, Hansen AK, Sabree ZL, Moran NA, Almeida RPP.
Origin and examination of a leafhopper facultative endosymbiont. Curr
Microbiol [Internet]. 2011 May [cited 2021 Sep 22];62(5):1565–72. Available
from: /pmc/articles/PMC3069327/

12. Lane, D. J. (1991). 16S/23S rRNA sequencing, p. 115-175. *In* E. Stackebrandt
and M. Goodfellow (ed.), Nucleic acid techniques in bacterial systematics.
John Wiley & Sons, New York. [Internet]. [cited 2021 Sep 22]. Available from:
http://www.sciepub.com/reference/56892

13. Turner S, Pryer KM, Miao VPW, Palmer JD. Investigating deep phylogenetic
relationships among cyanobacteria and plastids by small subunit rRNA
sequence analysis. J Eukaryot Microbiol [Internet]. 1999 [cited 2021 Sep
22];46(4):327–38. Available from:
https://pubmed.ncbi.nlm.nih.gov/10461381/

14. Nawrocki EP. Structural RNA Homology Search and Alignment using
Covariance Models. PhD thesis [Internet]. 2009 Jan 1 [cited 2021 Sep 22];282.
Available from: http://openscholarship.wustl.edu/etd/256

15. Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, et al.
Mrbayes 3.2: Efficient bayesian phylogenetic inference and model choice
across a large model space. Syst Biol [Internet]. 2012 May [cited 2021 Sep
22];61(3):539–42. Available from:
https://pubmed.ncbi.nlm.nih.gov/22357727/

16. Simon C, Frati F, Beckenbach A, Crespi B, Liu H, Flook P. Evolution, weighting,
and phylogenetic utility of mitochondrial gene sequences and a compilation of
conserved polymerase chain reaction primers. Ann Entomol Soc Am [Internet].
1994 Nov 1 [cited 2021 Sep 22];87(6):651–701. Available from:
https://academic.oup.com/aesa/article/87/6/651/19291

17. Nadal-Jimenez P, Valente RS. Preparation of polygalacturonic acid (PGA) to
study virulence in Erwinia/Pectobacterium [Internet]. protocols.io. 2021.
Available from: https://doi.org/10.17504/protocols.io.bydyps7w

18. Jimenez PN, Koch G, Papaioannou E, Wahjudi M, Krzeslak J, Coenye T, et al.
Role of PvdQ in Pseudomonas aeruginosa virulence under iron-limiting
conditions. Microbiology. 2010;156(1):49–59.

19. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina
sequence data. Bioinformatics [Internet]. 2014 Aug 1 [cited 2021 Sep
22];30(15):2114–20. Available from:
https://academic.oup.com/bioinformatics/article/30/15/2114/2390096

20. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. PLoS Comput Biol
[Internet]. 2017 Jun 1 [cited 2021 Sep 22];13(6):e1005595. Available from:
https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.10055
95

21. Tatusova T, Dicuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L,
et al. NCBI prokaryotic genome annotation pipeline. Nucleic Acids Res. 2016
Aug 19;44(14):6614–24.
22. Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes.
Nucleic Acids Res [Internet]. 2000 Jan 1 [cited 2021 Sep 22];28(1):27–30.
Available from: /pmc/articles/PMC102409/
23. Blin K, Shaw S, Kloosterman AM, Charlop-Powers Z, Van Wezel GP, Medema
MH, et al. AntiSMASH 6.0: Improving cluster detection and comparison
capabilities. Nucleic Acids Res. 2021 Jul 2;49(W1):W29–35.
24. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better,
faster version of the PHAST phage search tool. Nucleic Acids Res [Internet].
2016 Jul 8 [cited 2021 Sep 22];44(W1):W16–21. Available from:
https://www.readcube.com/articles/10.1093%2Fnar%2Fgkw387
25. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome
Biol [Internet]. 2015 Aug 6 [cited 2021 Sep 22];16(1):1–14. Available from:
https://genomebiology.biomedcentral.com/articles/10.1186/s13059-015-
0721-2
26. Ortori CA, Halliday N, Cámara M, Williams P, Barrett DA. LC-MS/MS
quantitative analysis of quorum sensing signal molecules. Methods Mol Biol
[Internet]. 2014 [cited 2021 Sep 22];1149:255–70. Available from:
https://pubmed.ncbi.nlm.nih.gov/24818911/
27. Galetto L, Marzachì C, Demichelis S, Bosco D. Host Plant Determines the
Phytoplasma Transmission Competence of Empoasca decipiens (Hemiptera:
Cicadellidae). J Econ Entomol. 2011;104(2):360–6.
28. Fathi SAA, Nouri-Ganbalani G, Rafiee-Dastjerdi H. Life cycle parameters of
Empoasca decipiens Paoli (Hom.: Cicadellidae) on four potato cultivars
(Solarium tuberosum L.) in Iran. J Entomol. 2009;6(2):96–101.
29. Barnard AML, Salmond GPC. Quorum sensing in Erwinia species. Anal Bioanal
Chem. 2007;387(2):415–23.
529
530

Table 1: Utilization of carbon sources for Growth by *S. purcellii* in the presence and absence of PGA, with comparison to *Pectobacterium carotovorum* LMG 2404[T]. +++: strong growth; ++: medium growth; +: weak growth; and -: no growth.

	S. purcelli	*S. purcelli* + PGA	*P. carotovorum.* LMG 2404	*P. carotovorum* LMG 2404 + PGA
Dextrin	-	-	+++	+++
D-maltose	-	-	+++	+++
D-Trehalose	-	-	+++	+++
D-Cellobiose	-	-	+++	+++
Gentiobiose	-	-	+++	+++
Sucrose	-	-	+++	+++
D-Turanose	-	-	+++	+++
Stachyose	-	-	+++	+++
D-raffinose	-	-	+++	+++
α-D-lactose	-	-	+++	+++
D-melibiose	-	-	+++	+++
B-Methyl-D-Glucoside	-	-	+++	+++
D-Salicin	-	-	+++	+++
N-Acetyl-D-Glucosamine	+++	+++	+++	+++
N-Acetyl-B-D-Mannosamine	-	-	+++	+++
N-Acetyl-D-Galactosamine	-	-	+++	+++
N-Acetyl Neuraminic Acid	-	-	+++	+++
α-D-Glucose	+++	+++	+++	+++
D-mannose	+++	+++	+++	+++
D-Fructose	+++	+++	+++	+++
D-Galactose	-	-	+++	+++
3-Methyl Glucose	-	-	+++	+++
D-Fucose	-	-	+++	+++
L-Fucose	-	-	+++	+++
L-Rhamnose	+++	+	+++	+++
Inosine	-	-	+++	+++
D-sorbitol	-	-	+++	+++
D-mannitol	-	-	+++	+++
D-Arabitol	-	-	+++	+++
myo-inositol	-	-	+++	+++
Glycerol	+++	+++	+++	+++
D-Glucose-6-PO4	-	-	+++	+++
D-Fructose-6-PO4	-	-	+++	+++
D-Aspartic Acid	-	-	+++	+++
D-Serine	-	-	+++	+++
Gelatin	-	-	+++	+++
Glycyl-L-Proline	-	-	+++	+++

L-Alanine	-	-	+++	+++
L-Arginine	-	-	+++	+++
L-Aspartic Acid	-	-	+++	+++
L-Glutamic Acid	-	-	+++	+++
L-Histidine	-	-	-	-
L-Pyroglutamic Acid	-	-	-	-
L-Serine	-	-	+++	+++
Pectin	-	-	+++	+++
D-Galacturonic acid	-	-	+++	+++
L-Galactonic Acid Lactone	-	-	+++	+++
D-Gluconic Acid	-	-	+++	+++
D-Glucuronic Acid	-	-	+++	+++
Glucuronamide	-	-	+++	+++
Mucic Acid	-	-	+++	+++
Quinic Acid	-	-	-	-
D-Saccharic Acid	-	-	+++	+++
ρ-hydroxy phenyl acetic acid	-	-	-	-
Methyl Pyruvate	+	+	+++	+++
D-Lactic Acid Methyl Ester	-	-	-	-
L-Lactic acid	-	-	-	-
Citric Acid	-	-	+++	+++
α-Keto-Glutaric-Acid	-	-	-	-
D-Malic Acid	-	-	-	-
L-Malic Acid	-	-	+++	+++
Bromo-Succinic Acid	-	-	+++	+++
Tween-40	-	-	-	+++
Gamma-Amino-Butyric acid	-	-	-	+++
α-Hydroxy-Butyric Acid	-	-	-	+++
B-Hydroxy-D-L-Butyric Acid	-	-	-	+++
α-Keto-Butyric-Acid	-	++	-	+++
Acetoacetic Acid	-	-	+	+++
Propionic Acid	-	-	-	-
Acetic Acid	+	+	+++	+++
Formic Acid	-	-	+++	+++

539 Table 2: Impact of environmental and xenobiotic stress conditions on *S. purcellii*
540 growth on Biolog III plates compared to *P. carotovorum,* in the presence/absence of
541 PGA. +++: maintains full growth under condition stated, ++: medium growth; +: weak
542 growth; and - :no growth under condition stated.
543

	S. purcelli	*S. purcelli* + PGA	*P. carotovorum* LMG 2404[T]	*P. carotovorum* LMG 2404[T] + PGA
pH 6	+++	+++	+++	+++
pH 5	-	-	-	-
1% NaCl	+++	+++	+++	+++
4% NaCl	+	+	+++	+++
8% NaCl	-	-	-	-
1% Sodium lactate	+++	+++	+++	+++
Fusidic acid	+++	+++	+++	+++
D-Serine	-	+++	-	-
Troleandomycin	+++	+++	+++	+++
Rifamycin SV	+++	+++	+++	+++
Minocycline	-	+++	-	-
Lincomycin	++	++	+++	+++
Guanidine HCl	++	++	+++	+++
Niaproof 4	++	++	+++	+++
Vancomycin	+++	+++	+++	+++
Tetrazolium Violet	++	++	+++	+++
Tetrazolium Blue	+++	+++	+++	+++
Nalidixic Acid	++	++	-	-
Lithium Chloride	+++	+++	+++	+++
Potassium Tellurite	+++	+++	-	-
Aztreonam	+++	+++	-	-
Sodium Butyrate	++	++	+++	+++
Sodium Bromate	-	-	-	-

547 **Table 3.** Genome features of the *Symbiopectobacterium purcellii* strain SyEd1
548 isolated from the leafhopper *Empoasca decipiens*.
549

Chromosome size (bp)	4,942,431
Plasmids	no
GC contents (%)	52.5
Number of predicted CDS	4,494
Number of pseudogenes	312
Average CDS length (bp)	948
Coding density (%)	86.2
Number of rRNA operons	7 (5S, 16S, 23S)
Number of tRNAs	76

550
551

552 Figure 1: SEM of *S. purcellii* SyEd1.

Figure 2: Phylogenetic affiliation of the 16S rRNA of *S. purcellii* compared to other

strains, as estimated with Mr Bayes. Numbers on nodes represent posterior

probability.

560   Figure 3: Virulence assay in potatoes.

562
563

Figure 4: Affiliation of *S. purcellii* with other bacteria as estimated using IQTREE based on 527 shared single copy orthologues. Coloured dots on nodes represent bootstrap support.

570 Figure 5: LC-MS/MS of an extraction of *S. purcellii* SyEd1 grown in BHI medium (A), 3-

571 OH-C10-HSL standard (B) and negative control using a sterile BHI medium extract (C).

577
578