Divergent selection predating the Last Glacial Maximum mainly

acted on macro-phenotypes in Norway spruce

Tiret M.[1,2*], Olsson L.[3], Grahn T.[3], Karlsson B.[4], Milesi P.[1], Lascoux M.[1], Lundqvist S.-O.[5],

García-Gil M.R.[2]

[1] Program in Plant Ecology and Evolution, Department of Ecology and Genetics, EBC and

SciLife Lab, Uppsala University, Sweden.

[2] Department of Forest Genetics and Plant Physiology, SLU, Umeå Plant Science Centre

(UPSC), Umeå, Sweden.

[3] RISE Bioeconomy, Box 5604, 114 86 Stockholm, Sweden.

[4] Skogforsk, Ekebo, 2250 SE-268 90 Svalöv, Sweden.

[5] IIC, Rosenlundsgatan 48B, 11863 Stockholm, Sweden.

*: corresponding author: Mathieu Tiret, **Email**: mathieu.tiret@ebc.uu.se

Running head: Divergent selection on phenotypes in Norway spruce (50 characters)

Summary:

The current distribution and population structure of many species were, to a large extent, shaped by cycles of isolation in glacial refugia and subsequent population expansions. Isolation in, and post-glacial expansion through heterogeneous environments led to either neutral or adaptive divergence. Norway spruce is no exception, and its current distribution is the consequence of a constant interplay between evolutionary and demographic processes. We investigated population differentiation and adaptation of Norway spruce for juvenile growth, diameter of the stem, wood density and tracheid traits at breast height. Data from 4,461 phenotyped and genotyped Norway spruce from 396 half-sib families in two progeny tests were used to test for divergent selection in the framework of Q_{ST} vs F_{ST}. We show that the macroscopic resultant trait (stem diameter), unlike its microscopic components (tracheid dimensions) and juvenile growth, was under divergent selection that predated the Last Glacial Maximum. Altogether, the current variation in these phenotypic traits in Norway spruce is better explained by local adaptation to ancestral environments than to current ones, where populations were partly pre-adapted, mainly through growth-related traits.

181 words / 300 max.

main text: 5145 words / 6500 max.

2 figures.

1 table.

Keywords: Norway spruce, wood quality, population structure, Q_{ST} vs F_{ST}.

Introduction

During the Quaternary glacial cycles (up to 2.5 million years ago), many European animal and plant species were repeatedly confined to southern refugia. Those were often located in the Iberian, Italian and Balkan peninsulas, but surviving populations could also be distributed on less discrete spots and, instead, scattered along the southern edge of the ice sheet (Bennett, 1997; Binney et al., 2009; Kremer et al., 2010). Improvement of climatic conditions during post-glacial periods led to demographic expansions accompanied by population admixture (e.g., Petit et al., 2003), new selective pressures (e.g., Suarez & Tsuitsui, 2008), and inter and/or intraspecific competition (e.g., Savolainen et al., 2007). Although the present population structure is the result of successive cycles of glaciation events that led to a series of demographic contractions and expansions, the current distribution of many species has, at least in part, been shaped by the different ecological processes and selective pressures that acted during the period after the Last Glacial Maximum (e.g., Petit et al., 2002; Palmé et al., 2003).

New abiotic and biotic pressures encountered during the glacial cycles presumably led to new adaptations. In many tree species, current phenotypes partly reflect past selection (Von Gadow et al., 2021), and selection is known to play an important role in shaping species distribution (Case et al., 2005; Savolainen et al., 2007; Wisz et al., 2013; Li et al., 2022). Among others, wood is one of the main tissues undergoing selection, since radial growth of the woody stem provides the mechanical support for woody plants to grow tall and gain a competitive advantage (Givnish, 1982; Iwasa, 1985; Falster & Westoby, 2003; Falster & Westoby, 2005). However, adaptation for upright growth is often associated with modified wood properties and evolutionary trade-offs (Poorter et al., 2006; Chave et al., 2009). Wood density is a good example of such a growth-survival trade-off (Kitajima 1994; Poorter et al., 2010), where low wood density, resulting from wider lumina and, in hardwood species, wider vessels (e.g., Enquist et al., 1999; Preston et al., 2006; Coomes et al., 2008), is associated with fast growth. Conversely, high wood density provides higher biomechanical safety (e.g., Niklas, 1993; Van Gelder et al., 2006) and resistance against pathogens (Augspurger & Kelly, 1984), and is thus associated with higher survival rates. The growth-survival trade-off is a common ecological trade-off observed in both tropical (e.g., Faster & Westoby, 2005; Poorter, 2008) and boreal species (e.g., Rossi et al., 2015; Pretzsch et al., 2018).

Norway spruce (*Picea abies* (L.) H. Karst.) is a dominant boreal conifer species that thrives nowadays in a vast northern domain spanning from Norway to central Russia, as well as in smaller European domains, such as the Alps and the Carpathians (Vendramin et al., 2000; Tsuda et al., 2016; Li et al., 2022; Nota et al., 2022). Repeated demographic movements associated to climatic cycles and ensuing secondary contacts led to the division of Norway spruce into seven genetic clusters (Chen et al., 2019): Northern Fennoscandia (NFE), Central and South Sweden (CSE), Russia-Baltics (RBA), Northern Poland (NPL), Central Europe (CEU), Alpine (ALP), Carpathian (ROM) domains. In southern Sweden, recent hybrids between CSE and ALP clusters can be encountered (CSE-ALP). The demographic history leading to this population structure went along a history of successive adaptations to biotic and abiotic factors. Following the hypothesis that most adaptation is reflected by cluster specific allele frequencies (Kremer et al., 2012), we investigated past adaptive signature through the population structure of Norway spruce. Using the framework of Q_{ST} vs F_{ST}, we assessed whether a phenotype diverges more between the genetic clusters than expected under random genetic drift. The timing of the divergence thereby inferred was given by that of the population structure that long predated the Last Glacial Maximum (10 to 15 million years ago; Chen et al., 2019; Li et al., 2022). By doing so, we were able to infer the proportion of the current phenotypic variation that is due to adaptations predating the Last Glacial Maximum. For that, in order to account for possible evolutionary trade-offs between wood-related traits, we analyzed several phenotypes: stem diameter and ring widths, wood density and tracheid traits defining it. Since the nature of selection between trees is expected to change over ontogenetic stages (Miriti, 2006), we studied the development of these phenotypes with cambial age/ring number from increment cores sampled from two progeny trials of the Swedish breeding program. We have conducted the analyses on a large dataset of 4,461 twenty-year-old trees from 396 half-sib families, 118,145 SNPs from exome sequencing. We show that the current phenotypic variation is partly explained by divergent selection that predated the Last Glacial Maximum, which mainly acted on macro-phenotypes (stem diameter), rather than micro-phenotypes (tracheid dimensions).

Materials and Methods

Plant material. The study is based on previously presented data (Chen et al., 2014; Chen et al., 2016; Lundqvist et al., 2018), which are here analyzed with a special attention to population structure and selection. The data originates from two progeny trials of Norway spruce at two different sites: FF1146 (Höreda, Sweden; 57°60'N 15°04'E), and FF1147 (Erikstorp, Sweden; 56°80'N 13°90'E). Both trials were established by the Forestry Research Institute of Sweden (Skogforsk) in the spring of 1990 from open-pollinated seeds harvested on selected trees from 112 stands in southern Sweden. FF1146 comprised 1,373 half-sib families and FF1147, 1,375 half-sib families. In both trials, a randomized incomplete block design was implemented (20 blocks in FF1146, 23 in FF1147), with single-tree plots and a homogeneous spacing of 1.4 m x 1.4 m. Dead trees and trees of low breeding interest were removed in 2008. After filtering out (i) half-sib families absent in the established Skogforsk clonal archives, and (ii) stands with less than three half-sib families, the study included a total of 518 half-sib families with 5,644 progeny (2,983 from FF1146, and 2,661 from FF1147).

Population structure. Population structure was previously assessed at a fine scale using 399,801 unlinked putatively non-coding SNPs on 1,672 individuals (see details in Chen et al., 2019), categorizing families into the following genetic clusters (ordered by decreasing latitude): Central and South Sweden (CSE), Russia-Baltics (RBA), Northern Poland (NPL), Central Europe (CEU), Alps (ALP), Carpathians (ROM), and hybrids between CSE and ALP (CSE-ALP) (for tree geographical origins, see Fig. 1 in Milesi et al., 2019). The families examined in this study followed the same genetic clustering (Fig. 1A, 1B). Individuals from the ROM genetic cluster were removed, as our sample included only three such families, whereas all other clusters included at least ten families (Table S1).

Phenotype measurements. Increment cores of 12 mm diameter, extracted from the northern side of the stem, were collected in 2010 and 2011 at breast height (BH, 1.3 m above ground) from six trees per half-sib family per trial. The usual cutting age is 60-70 years or more, while the cores were extracted from 20- or 21-year-old trees: only juvenile stages of wood characteristic were then captured by the cores. The increment cores were analyzed by SilviScan (Evans, 1994; 2006) at Innventia, now RISE (Stockholm, Sweden), assessing high-resolution pith-to-bark radial variation in growth, tracheid and wood properties. In this study, we focused on the development with cambial age (ring number) of ring width (RW, in mm), and averages per ring of wood density (WD, in kg/m^3), tracheid

radial width (TRW, in μm), tracheid tangential width (TTW, in μm), tracheid wall thickness
(TWT, in μm), diameter at breast height (DBH, in mm) calculated from ring widths, and the
age when the trees reached breast height (ABH). Although ABH is related to upright growth,
it can be estimated accurately with high quality increment cores from old trees. The
measurements, identification of annual rings, and the importance of ABH are discussed in
Lundqvist et al. (2018).

The two innermost rings were removed as the extremely curved structure of the xylem does
not allow the same high data quality as for rings further out. In addition, only data up to the
13^{th} ring was kept for analyses, since only fast-growing offspring with low ABH are
represented at the highest cambial ages. To avoid the effect of the thinning performed, we
removed all annual rings formed after 2008. The study then covered six traits (RW, WD,
TRW, TTW, TWT, DBH) for 11 rings per offspring in most of the cases, and their ABH.

Exome capture and SNP filtering. The exome capture (target re-sequencing) of the mother
of each half-sib family is available on BioProject PRJNA511374 (Chen et al., 2019) and
BioProject PRJNA731384 (Chen et al., 2021). Total genomic data was extracted from the
buds of the 518 maternal trees (or needles when buds were not available) using the Qiagen
Plant DNA extraction kit (Qiagen, Germany). DNA concentration was assessed using the
Qubit® ds DNA Broad Range Assay Kit (Oregon, USA). The extracted DNA was sent to
RAPiD Genomics (Gainesville, USA) for library preparation and exome sequencing.
Sequence capture, with an average depth of 15X, was performed using the Illumina HiSeq
2500 (San Diego, USA), with 40,018 diploid probes (120bp-length) designed to capture
26,219 genes of *P. abies* (Vidalis et al., 2018). After mapping the raw reads against the
P. abies reference genome v1.0 (Nystedt et al., 2013) using BWA-MEM with default settings
(Li & Durbin, 2010), PCR duplicates were sorted and marked using SAMTOOLS v1.2 (Li et
al., 2009) and Picard v1.141 (http://broadinstitute.github.io/picard). INDELs were realigned
using GATK v3.5.0 (McKenna et al., 2010). Variant calling was performed using GATK
HaplotypeCaller v3.6, non-biallelic SNPs were filtered out using GATK SelectVariants v3.6
(option --restrict-alleles-to BIALLELIC), and quality filtering was carried out by the Variant
Quality Score Recalibration following the procedure of Van der Auwera et al. (2013).

As an alternative way to account for Minor Allele Frequency (MAF), we filtered out SNPs
with less than five individuals per genotype to avoid a leverage effect. We performed a loose
filtering for variant call rate (or missingness) with PLINK v1.9 (Chang et al., 2015), by

removing individuals with more than 70% SNP-missingness, resulting in 472 half-sib
families, and by removing SNPs with more than 70% individual-missingness, resulting in
118,145 SNPs (each mother is then genotyped for at least 35,443 SNPs). 70,277 SNPs
were successfully mapped on the consensus map of Bernhardsson et al. (2019).
180
181   **Statistical model.** The traits (DBH, WD, RW, TRW, TTW, and TWT) were normalized by
182   their respective overall means (across all trees and rings 3 to 13) to allow comparison
183   between traits. Then, they were analyzed at each cambial age separately, resulting in six
184   traits over 11 rings, a total of 66 traits in addition to ABH.
185
186   To study the family effect underlying the traits, we used the following mixed model:
187    $Y_{ijkl} = \mu + s_i + B_{ij} + F_k + \varepsilon_{ijkl},$ (1)
188
where 'Y_{ijkl}' is the trait of the l-th offspring of the k-th family in the j-th block at the i-th site, 'μ'
is the grand mean, 's_i' is the fixed effect of the i-th site, 'B_{ij}' and 'F_k' are uncorrelated random
effects of the j-th block nested in i-th site (with variance σ_B^2) and the k-th half-sib family (with
variance σ_F^2) respectively, and 'ε_{ijkl}' is the residual fitted to a Gaussian distribution of variance
σ^2. Heritability was estimated as $4 \times \sigma_F^2 / (\sigma_F^2 + \sigma_B^2 + \sigma^2)$ as an intraclass correlation
estimation of half-sib families (p.166, Falconer, 1989).
195
196   To investigate the effect of genetic clusters on traits jointly with the effect of ABH, we used
197   the following linear mixed model:
198
199    $Y_{ijkl} = \mu + c_l + s_i + g_k + (c \times g)_l + (c \times s)_l + B_{ij} + F_k + \varepsilon_{ijkl},$ (2)
200
where 'c_l' is the fixed effect of the l-th offspring's ABH, '$(c \times g)_l$' is the fixed interaction effect
between ABH and the genetic cluster of the l-th offspring, '$(c \times s)_l$' is the fixed interaction
effect between ABH and the site of the l-th offspring, and the other terms are as defined
above in models (1). Model (2) was applied on the 66 traits, as previously mentioned. Since
the interaction between ABH and the site effect was always significant, we split the dataset
per site, as recommended by Crawley (2007, pp. 323–386). The results are focusing on the
Höreda trial, but the conclusions were the same in the Erikstorp trial.
208
Finally, to disentangle the effect of divergent selection and neutral divergence, we estimated
Q_{ST} as defined by Spitze (1993) with the following mixed model:

$Y_{ijkl} = \mu + c_l + G_k + B_{ij} + F_k + \varepsilon_{ijkl}$,   (3)

where 'G_k' is the random effect of the k-th family's genetic cluster (with variance σ_A^2), and the

other terms are as defined above in model (1). Q_{ST} was estimated as $\sigma_A^2/(\sigma_A^2 + 2\sigma_F^2)$, and

averaged across rings. Following the recommendation of Whitlock (2008), we compared Q_{ST}

to the positive part of the distribution of F_{ST} (with VCFTools v0.1.16, option --weir-fst-pop;

Danecek et al., 2011) computed on SNPs in chromosomic regions identified as non-coding

by Nystedt et al. (2013). This set was not filtered for MAF or missingness, amounting then to

789,786 SNPs. If Q_{ST} of a trait is significantly larger than F_{ST}, the trait can be considered to

have diversified more than expected under neutrality in a heterogeneous environment. In

contrast, if Q_{ST} cannot be distinguished from F_{ST} values, we would have little evidence that

selection is responsible for phenotypic divergence.

All computations were done using R v4.0.4 (R core team, 2021), and all models were fitted

with the R package *lme4* (function *lmer*; Bates et al., 2015). Statistical significance of the

fixed effects was assessed using a type II Wald chi-square test (function *Anova* of the R

package *car*), and significance of a slope was assessed using a Student *t*-test (function *t.test*

of the R package *stats*). To account for the unbalance caused by consecutive filtering, we

only kept families with at least four offspring per site. Applying all the previous filters, we

ended up with 396 half-sib families and 4,461 offspring per trait. Multiple testing across the

66 traits were accounted for by considering a Bonferroni adjusted cut-off of $p = 7.6 \times 10^{-4}$.

Results

The genetic cluster effect was significant at all ages

At all cambial ages, phenotypic variation of all six phenotypes (DBH, WD, RW, TRW, TTW, and TWT) was associated with the geographical origin of the genetic cluster, as the genetic cluster effect was significant for all rings ($\chi^2 > 11.6$, degree of freedom or df = 5, p = 0.041), except for the 3rd ring TWT ($\chi^2 = 10.9$, df = 5, p = 0.053). Every trait of the southern populations was consistent with a higher investment in growth than in wood density (higher DBH, RW, TRW, and TTW; lower WD, and TWT).

Phenotypic variation exhibited a strong temporal pattern, an ontogenetic shift (Fig. 1C). For all rings, phenotypic variation also depended on the geographical origin of the trees. The ontogeny was quantitatively measurable, as it followed the second axis of the PCA on the six phenotypes (41.2% by PC1, 38.3% by PC2). As expected, the ontogeny was mainly driven by that of DBH, which alone explained 34% of PC2, and on average increased fourfold over the cambial ages investigated, while other phenotypes varied at most by 54% (for further details, see Lundqvist et al., 2018). As cambial age increased, ontogenetic shift was instead driven by other phenotypes, as the coefficient of variation of DBH was the only one decreasing over time (Student's t = -6.13, p < 0.001; Fig. S1). Overall, the ontogenetic shift resulted from a complex interplay of multiple phenotypes, all with heritabilities higher than 30% (Table 1), in accordance with previously reported values (Hannrup et al., 2004; Chen et al., 2014, 2021).

Accounting for early growth rate ruled out environmental causes from the analysis of ontogeny

By definition, ontogenic tempo is given by the ABH. It had a relatively higher environmental variance, and therefore a relatively lower heritability of 17.31% than other traits (Table 1). The lower heritability was not due to a lower genetic variance, in fact quite the opposite. ABH also depended on the geographical origin of the genetic cluster, as the genetic cluster effect was significant ($\chi^2 = 102.8$, df = 1, p < 10^{-5} in model (1)). As mentioned earlier, ABH was latitudinally structured, half-sib families of southern origin having low ABH reflecting higher early growth rates.

Despite evidence supporting the genetic control of ABH, the genetic signals in the model (2)
seems to have been absorbed by the genetic cluster and the family effects, and let the
feature "ABH" only reflect environmental factors out of a lesser correlation with the response
variable: when correcting for multiple testing, in model (2), the effects of ABH and of the
genetic clusters were significant for more than 70% of the phenotypes (47 out of 66 for both),
whereas their interaction was significant for less than 2% of the phenotypes (1 out of 66;
Table S2). In other words, ABH explained a large part of the phenotypic variation, while
being orthogonal to the family or genetic cluster effect - the variable "ABH" was then likely
only capturing the environmental part of ABH. In this particular statistical context, we will
assume that ABH reflects environmental factors – else than the block effect already
accounted for – impacting early growth and orthogonal to genetic factors (e.g., micro-
environmental heterogeneity of the field, supposedly orthogonal to genetic factors because
of the random block design). Under this assumption, it is possible to disentangle the effect of
genetic factors on the phenotype of a "mature" tree from environmental factors giving
advantages in early stages of life (i.e., acclimation). We will come back to this interpretation
in the Discussion.

Early favorable environment mainly impacted macro-phenotypes

The effect of acclimation, approximated by ABH, varied mainly with the cambial age,
following a consistent pattern for every trait (but RW): acclimation mostly impacted early (3rd
and 4th) or later (9th and higher) rings, and least impacted intermediate rings (5th to 8th). More
precisely, analysis of variance of model (2) showed that, for early rings, ABH significantly
impacted every trait but the 4th ring WD (χ^2 = 7.60, df = 1, p = 5.84 x 10^{-3}). For intermediate
rings, ABH significantly impacted only 6 out of 20 traits, namely the 5th ring DBH, the 6th ring
TTW, the 7th ring TTW, and the 8th ring DBH, WD, and TWT (χ^2 > 13.9, df = 1, p < 1.88 x 10^{-4}
). For later rings, ABH significantly impacted every trait but the 9th ring TTW (χ^2 = 3.30, df =
1, p = 6.92 x 10^{-2}), the 10th ring TTW (χ^2 = 4.70, df = 1, p = 3.02 x 10^{-2}), the 11th ring TWT (χ^2
= 5.65, df = 1, p = 1.75 x 10^{-2}) and the 12th ring TWT (χ^2 = 9.98 x 10^{-2}, df = 1, p = 0.752). Only
RW followed another pattern, for which ABH impacted mostly intermediate and later rings
(5th ring and beyond), as the significance of ABH followed a concave curve starting with an
increase of significance from the 5th ring (χ^2 = 40.70, df = 1, p < 10^{-5}) up until the 9th ring (χ^2 =
1199, df = 1, p < 10^{-5}), then a decrease until the 13th ring (χ^2 = 54.50, df = 1, p < 10^{-5}). This
can also be seen in Fig. 5a of Lundqvist et al. (2018), where similar age curves, shifting
sidewise with ABH, have their steepest phase during the intermediate cambial ages (see
also Supplemental Material, method A).

On later rings, trees with a low ABH, i.e., fast growers, ended up with a markedly larger

DBH, lower WD and TWT, and slightly higher TRW and TTW. Fast growers had increasingly

higher DBH over time, as the effect of ABH decreased from the 3^{rd} ring to the 13^{th} ring (from

5.36×10^{-3} to -2.87×10^{-2}). On the opposite, fast growers had increasingly lower WD, as the

effect of ABH increased from the 3^{rd} ring to the 13^{th} ring (from 4.93×10^{-3} to 3.1×10^{-2}). Fast

growers also had significantly larger 9^{th} rings, as the effect of ABH on RW followed a convex

curve matching its concave significance, reaching its largest negative value at the 9^{th} ring

(-5.82×10^{-1}). At all cambial ages, the effect of ABH was smaller for tracheid properties

(TRW, TTW, TWT) than for macro-phenotypes (DBH, WD) (Fig. S2). These results show

that environmental factors favoring early growth had an impact on all later stage phenotypes

(except for RW), albeit stronger on macro-phenotypes (DBH, WD) than on micro-phenotypes

(TWT, TRW, TTW).

312 Macro-phenotypes were also the ones showing signature of past
313 divergent selection

In addition to all phenotypes being, as shown above, significantly associated with

geographical location, the significance of this association increased over cambial ages

(Student's t > 4.36, p < 0.002), apart from TRW (Student's t = -0.764, p = 0.464). The

significance of the genetic cluster in model (2) on the 13^{th} ring was large for DBH and RW (χ^2

> 131.26, df = 5, p < 10^{-5}), intermediate for WD and TWT (χ^2 > 79.51, df = 5, p < 10^{-5}), and

low for TRW and TTW (χ^2 < 19.39, df = 5, p < 0.002). Presumably, the significance of TWT

might be due to its high correlation with the significant WD (ρ = 0.88). The variation of

growth-related macro-phenotypes, namely DBH and RW, therefore reflected the population

structure more pronouncedly than did tracheid dimensions or wood density (Fig. 2A).

Population structure was even more visible on phenotypes when correcting for

environmental factors, as the PCA on family effects clearly exhibited the population structure

(Fig. 2B): genetic clusters were disposed along PC1 (58.9% of variance explained) in

decreasing latitudinal order, except for CEU and ALP being reversed. PC1 was mostly

driven by DBH, RW, TRW and WD, while PC2 (17.0% of variance explained) was mostly

driven by TTW and TWT. That the traits are structured by both family and genetic cluster can

also be seen in the fact that intra-family (or intra-population) variance was significantly

increased when the family (or population) structure was shuffled (see Supplement, method

B). Not only did phenotypic variation reflect the underlying population structure, but it was

also quantitatively associated with latitude.

The quantitative association between environmental gradients and phenotypic variation was

partly due to selection (Fig. 2C), at least for macro-phenotypes. Selection as assessed by

Q_{ST} from model (3) was only significant for DBH (p = 0.0115), RW (p = 0.0086), and

marginally so for TTW (p = 0.0437). Despite the Q_{ST} of TTW being marginally significant,

what is striking is that TRW, which is a component of DBH and RW, is not (p = 0.2680).

Although interpreting the difference between Q_{ST} and F_{ST} can be fraught with difficulties, one

could note that whatever the distribution of F_{ST}, selection, if any, would primarily act upon

growth-related macroscopic traits. This strong association between population structure and

signatures of selection made impossible the inference of genetic variants underlying

phenotypic variation since correction for population structure in GWAS made all associations

non-significant (see Supplement, method C; Fig. S3-S4). Overall, phenotypic divergence in

growth-related macroscopic traits (DBH and RW) is partly due to divergent selection on top

of neutral divergence.

Discussion

The Swedish breeding program is a mixture of local and recently introduced trees from other European countries. As such, although incomplete, the progeny trials reflect the population structure due to glaciation events and constitute an important source of information for evolutionary biology (Chen et al., 2019; Milesi et al., 2019; Li et al., 2022). The breeding conditions constituted a unique opportunity to accurately correct phenotypes for environmental factors, by ruling out recruitment selection and considerably lowering interspecific competition. Furthermore, by correcting for population structure and using common garden experiments, we can rule out neutral and plastic processes so that the observed phenotypic differences can be interpreted as being the outcome of adaptation (Savolainen et al., 2013; de Villemereuil et al., 2020).

In our common garden experiments, growth exhibited a high level of population differentiation, as expected (Savolainen et al., 2007; Kremer et al., 2012). Population differentiation followed a latitudinal pattern, as previously reported (Rossi et al., 2015; Montwé et al., 2016; Milesi et al., 2019). The growth rate was generally differentiated among individual trees at early stages, supporting the fact that young ages are the most dynamic phases (Lundqvist et al., 2018). Differences in apparent growth rate were mainly due to a longer growth span (see discussion in Milesi et al., 2019), and despite its low heritability, its high genetic and environmental variances suggest that it could be related to fitness (Houle, 1992; Caballero, 2020, p.235). This is also supported by the fact that the effect of ABH (here only capturing the micro-environmental effect) was strongest on phenotypes on which signature of selection was also the strongest. Early advantages, possibly acquired by chance, seem to be re-invested by default on resultants – traits giving competitive advantages.

The quantitative association between phenotypes and population structure predates the Last Glacial Maximum

Phenotypic variation had a clear latitudinal component at both genetic and phenotypic levels, as previously reported in Norway spruce (e.g., Milesi et al., 2019) and in other species (e.g., Sang et al., 2019; Csillery et al., 2020). Since local adaptation matches the environment (Savolainen et al., 2013) and post-glacial recolonization routes followed environmental gradients (Gaggiotti et al., 2009; Li et al., 2022), population structure of Norway spruce is, as

a result, strongly confounded with local adaptation. Population structure is thus a good
predictor of selection in Norway spruce, as previously reported (Collignon et al. 2002;
Savolainen et al., 2007; Kremer et al. 2010). However, local adaptation is not necessarily
limited to recent population history. Some of the adaptations displayed by Norway spruce
were associated to ancestral environments, and some to current environments. If we
consider that adaptation can be dated by the population structure it is associated with,
knowing that the population structure of Norway spruce largely predates the Last Glacial
Maximum (10 to 15 million years ago; Chen et al., 2019; Li et al., 2022), the observed
phenotypic divergence is partly due to adaptations also predating the Last Glacial Maximum.
We posit that the quantitative association between phenotype, latitude, and population
structure would not be observed if there were no local adaptation to ancestral environments.
The last post-glacial expansion of Norway spruce was thus mainly shaped by adaptations
formerly acquired – not *newly* acquired. In other words, the phenotypic variation within the
refugia explains in part the latitudinal variation observed nowadays at large geographical
scale.
397
Because of the association between phenotypic variation and demographic processes,
assessing the genetic control underlying adaptation of Norway spruce is challenging: not
correcting for population structure in GWASs, will lead to a high false positive rate (Barton et
al., 2019; Lawson et al., 2019), and correcting for population structure, may remove true
association, i.e., lead to a high false negative rate. A comprehensive assessment of the
genetic variants underlying phenotypic variation is then not possible when there is such a
strong latitudinal component. Increasing the number of samples or the number of SNPs
remains inconclusive (Chen et al., 2021). A possible workaround suggested by Milesi et al.
(2019) is to consider phenotypes that are less latitudinally associated, such as the
longitudinally associated bud burst, and correct other phenotypes by the association with
population structure thereby inferred. Alternatively, we can consider several species jointly
as suggested by Alberto et al. (2013).

Divergent selection mostly affected resultants, not components

The strongest genetic divergence was observed for growth-related traits, and secondly for
wood density, suggesting that most of the adaptation is associated with radial or upright
growth. Divergent selection acting at early life stages most likely led to the accumulation of
competitive advantages, causing then an observable ontogenetic shift of the effect of
population structure. The average value of F_{ST} was slightly lower than previously reported

(0.0488 in our study, and ~0.116-0.145 in Table 1 of Chen et al., 2019), and the
interpretation of Q_{ST} *vs* F_{ST} remains difficult (e.g., Whitlock, 2008; Edelaar et al., 2011).
However, overall, correcting the estimates of F_{ST} would not change the conclusion of our
study: if there is selection, it would affect growth-related macroscopic traits more strongly
than wood density or its microscopic components. Consistently, the PCA on family effects
showed that the latitudinal component – and so, population structure – was mostly explained
by stem diameter, ring width and wood density. Neutral divergence could not be excluded for
early growth rate (ABH) in spite of its clear relation with growth. Either this trait is neutral,
which is highly unlikely as early-stage selection is known to be strong (e.g., Saleh et al.,
2022), or, more likely, selection on ABH is invariant among genetic clusters. In the latter
case, selection on early growth rate would be strong regardless of the heterogeneity of the
locations, where variability would not come from divergence but from, for instance, trade-offs
with other life-history traits (e.g., wood density).
429
The difference between Q_{ST} and F_{ST} is due to the covariance between effect sizes, and the
covariance between allelic frequencies (reviewed in Le Corre & Kremer, 2012). Altogether,
the resulting pattern of our analyses is that adaptation is more tightly associated to traits that
are resultants rather than to its components, in agreement with Caballero (2020, p. 253).
Indeed, components can have random variations that are invisible to selection – therefore
neutral, as long as the components contribute to a resultant that is fit. In other words,
components have a higher degree of freedom, and therefore are noisier. Since genetic
polymorphism depicted by SNPs are partly linked to selection, genomic selection on
resultant – such as stem diameter – should be much more efficient than on components –
such as tracheid traits. An alternative explanation is simply that components were not
associated with divergent selection predating the Last Glacial Maximum, and might be under
recent selection – not detectable with our approach.

Conclusion
As expected, phenotypic variation reflected the population structure of Norway spruce. The
fact that the phenotypes were latitudinally ordered suggests that the origin of population was
strongly confounded with adaptation. Teasing apart the demographic history and the
adaptation history is challenging, more so that it has been demonstrated that population
history had a strong impact on phenotypic divergence in Norway spruce (Lagercrantz &
Ryman, 1990; Milesi et al., 2019). Divergence of phenotypic traits might in part be the results
of past adaptations, which can be dated by the population structure it is associated with. In

accordance with previous studies (Savolainen et al., 2013; Milesi et al., 2019; Martinez-
Sancho et al., 2021), our results support the hypothesis that the current distribution of
Norway spruce reflects in part adaptations to ancestral environments, as some signatures of
selection were associated with population structure that predated the Last Glacial Maximum.
The divergent selection affected more growth-related macro-phenotypes than micro-
phenotypes at tracheid level – presumably invisible to selection.
456

Author contribution

459 RGG, SOL, and BK designed the original study; TG, LO and SOL managed the phenotypic
460 characterization of the samples and organized the data; PM, ML and RGG contributed
461 genomic data; MT and PM analyzed the data; MT, PM, ML, SOL, RGG drafted the
462 manuscript. All authors read and approved the final version of the manuscript.

Data Availability

464 The exome capture raw reads and the RNA-seq data have been deposited in NCBI's
465 sequence read archive (SRA) under accession number PRJNA511374 (Chen et al., 2019)
466 and PRJNA731384 (Chen et al., 2021). For question on availability of the raw phenotypic
467 data, contact RGG.

Acknowledgements

The authors acknowledge the Swedish research program Bio4Energy and Skogforsk for
access to phenotypic data, and the Swedish Foundation for Strategic Research (SSF), grant
number RBP14-0040 for exome capture data. This study was funded by the following
sources: Uppsala University, SLU Umeå, and the European Union's Horizon 2020 B4Est for
basic functioning and postdoc grant for MT. The authors would also like to thank Sophie
Karrenberg and Zhi-Qiang Chen for their useful feedback. The computations and data
handling were enabled by resources provided by the Swedish National Infrastructure for
Computing (SNIC 2017-7-296) at UppMax partially funded by the Swedish Research Council
through grant agreement no. 2018-05973.

478
479
480

481 References

483 Alberto, F. J., Aitken, S. N., Alía, R., et al. (2013). Potential for evolutionary responses to
484 climate change–evidence from tree populations. *Global change biology*, 19(6):1645–1661.
485

486 Augspurger, C. K., & Kelly, C. K. (1984). Pathogen mortality of tropical tree seedlings:
487 experimental studies of the effects of dispersal distance, seedling density, and light
488 conditions. *Oecologica*, 61:211–217.
489

490 Barton, N., Hermisson, J., & Nordborg, M. (2019). Population genetics: why structure
491 matters. *eLife*, 8:e45380
492

493 Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models
494 using lme4. *Journal of Statistical Software*, 67(1):1–48.
495

496 Bennett, K. D. (1997). *Evolution and ecology: the pace of life*. Cambridge University Press,
497 Cambridge, UK.
498

499 Bernhardsson, C., Vidalis, A., Wang, X., et al. (2019). An ultra-dense haploid genetic map
500 for evaluating the highly fragmented genome assembly of Norway spruce (*Picea abies*). *G3
501 Genes|Genomes|Genetics*, 9:1623–1632.
502

503 Biging, G. S., & Dobbertin, M. (1995). Evaluation of competition indices in individual tree
504 growth models. *Forest science*, 41(2):360–377.
505

506 Binney, H.A., Willis, K.J., Edwards, et al. (2009). The distribution of late-Quaternary woody
507 taxa in northern Eurasia: evidence from a new macrofossil database. *Quaternary Science
508 Reviews,* 28:2445–2464.
509

510 Burton, O. J., Phillips, B. L., & Travis, J. M. (2010). Trade-offs and the evolution of life-
511 histories during range expansion. *Ecology Letters*, 13:1210–1220.
512

513 Caballero, A. (2020). Quantitative Genetics (p. 235). Cambridge University Press.
514

Callaway, R. M. (1995). Positive interactions among plants. *The Botanical Review*,
61(4):306–349.

Case, T. J., Holt, R. D., McPeek, M. A., & Keitt, T. H. (2005). The community context of
species' borders: ecological and evolutionary perspectives. *Oikos*, 108:28–46.

Chang, C. C., Chow, C. C., Tellier, L. C. A. M., Vattikuti, S., Purcell, S. M., & Lee, J. J.
(2015). Second-generation PLINK: rising to the challenge of larger and richer datasets.
GigaScience, 4(1).

Chave, J., Coomes, D., Jansen, S., Lewis, S. L., Swenson, N. G., & Zanne, A. E. (2009).
Towards a worldwide wood economics spectrum. *Ecology Letters*, 12(4):351–366.

Chen, Z. Q., García-Gil, M. R., Karlsson, B., Lundqvist, S.-O., Olsson, L., & Wu, H. X.
(2014). Inheritance of growth and solid wood quality traits in a large Norway spruce
population tested at two locations in southern Sweden. *Tree Genetics & Genomes*,
10(5):1291–1303.

Chen, Z. Q., Karlsson, B., Mörling, T., et al. (2016). Genetic analysis of fiber dimensions and
their correlation with stem diameter and solid-wood properties in Norway spruce. *Tree
Genetics & Genomes*, 12(6):1–12.

Chen, J., Li, L., Milesi, P., et al. (2019). Genomic data provide new insights on the
demographic history and the extent of recent material transfers in Norway spruce.
Evolutionary Applications, 12:1539–1551.

Chen, Z.-Q., Zan, Y., Milesi, P., et al. (2021). Leveraging breeding programs and genomic
data in Norway spruce (*Picea abies* L. Karst) for GWAS analysis. *Genome Biology*, 22:179.

Collignon, A. M., Sype, H. V. D., & Favre, J. M. (2002). Geographical variation in random
amplified polymorphic DNA and quantitative traits in Norway spruce. *Canadian Journal of
Forest Research*, 32(2):266–282.

Coomes, D. A., Heathcote, S., Godfrey, E. R., Shepherd, J. J., & Sack, L. (2008). Scaling of
xylem vessels and veins within the leaves of oak species. *Biology Letters*, 4(3):302–306.

Crawley, M. J. (2007). Statistical modelling. In *The R book* (pp. 323–386). Chichester, UK.

Csilléry, K., Ovaskainen, O., Sperisen, C., Buchmann, N., Widmer, A., & Gugerli, F. (2020).

Adaptation to local climate in multi-trait space: evidence from silver fir (Abies alba Mill.)

populations across a heterogeneous environment. *Heredity*, 124(1):77–92.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, et al. (2011). The

variant call format and VCFtools. *Bioinformatics*, 27(15):2156–2158.

De Villemereuil, P., Gaggiotti, O. E., & Gouder, J. (2020). Common garden experiments to

study local adaptation need to account for population structure. *Journal of Ecology*, 00:1–5.

Edelaar, P. I. M., & Björklund, M. (2011). If FST does not measure neutral genetic

differentiation, then comparing it with QST is misleading. Or is it?. *Molecular Ecology*,

20(9):1805–1812.

Enquist, B. J., West, G. B., Charnov, E. L., & Brown, J. H. (1999). Allometric scaling of

production and life-history variation in vascular plants. *Nature*, 401:907–911.

Evans, R. (1994) Rapid measurement of the transverse dimensions of tracheids in radial

wood sections from Pinus Radiata. *Holz-forschung*, 48:168–172

Evans, R. (2006) Wood stiffness by X-ray diffractometry (pp 138–146). In: Stokke DD,

Groom HL (eds) Characterization of the cellulosic cell wall. Wiley, Hoboken.

Excoffier, L., Foll, M., & Petit, R. J. (2009). Genetic consequences of range expansions.

Annual Review of Ecology, Evolution, and Systematics, 40:481–501.

Falconer, D.S. (1989). *Introduction to quantitative genetics* (3rd edition, p. 116). Pearson

Education India.

Falster, D. S. & Westoby, M. (2003). Plant height and evolutionary games. *Trends in

Ecology & Evolution*, 18(7):337–343.

Falster, D. S. & Westoby, M. (2005). Alternative height strategies among 45 dicot rain forest
species from tropical Queensland, Australia. *Journal of Ecology*, 93(3):521–535.
587
588 Givnish, T. J. (1982). On the adaptive significance of leaf height in forest herbs. *The*
589 *American Naturalist*, 120(3):353–381.
590
591 Houle, D. (1992). Comparing evolvability and variability of quantitative traits. *Genetics*,
592 130(1):195–204.
593
594 Iwasa, Y., Cohen, D., and Léon, J. A. (1985). Tree height and crown shape, as results of
595 competitive games. *Journal of Theoretical Biology*, 112(2):279–297
596
597 Kitajima, K. (1994). Relative importance of photosynthetic traits and allocation patterns as
598 correlates of seedling shade tolerance of 13 tropical trees. *Oecologica*, 98:419–428.
599
600 Kremer, A., Le Corre, V., Petit, R. J., & Ducousso, A. (2010). Historical and contemporary
601 dynamics of adaptive differentiation in European oaks. *Molecular approaches in natural*
602 *resource conservation*, 101–122.
603
604 Kremer, A., Ronce, O., Robledo-Arnuncio, et al. (2012). Long-distance gene flow and
605 adaptation of forest trees to rapid climate change. *Ecology letters*, 15(4):378–392.
606
607 Lagercrantz, U., & Ryman, N. (1990). Genetic structure of Norway spruce (Picea abies):
608 concordance of morphological and allozymic variation. *Evolution*, 44(1):38–53.
609
610 Lawson, D. J., Davies, N. M., Haworth, et al. (2019). Is population structure in the genetic
611 biobank era irrelevant, a challenge, or an opportunity? *Human Genetics*, 139:23–41.
612
613 Le Corre, V., & Kremer, A. (2012). The genetic differentiation at quantitative trait loci under
614 local adaptation. *Molecular ecology*, 21(7):1548–1566.
615
616 Li, H., Handsaker, B., Wysoker, A., Fennell, T., et al. (2009). The sequence alignment/map
617 format and SAMtools. *Bioinformatics*, 25(16):2078–2079.
618

Li, H. & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows–Wheeler

transform. *Bioinformatics*, 26(5):589–595

Li, L., Milesi, P., Tiret, M., et al. (2022). Teasing apart the joint effect of demography and

natural selection in the birth of a contact zone. *New Phytologist* (in press).

Lundqvist, S.-O., Seifert, S., Grahn, T., et al. (2018). Age and weather effects on between

and within ring variations of number, width and coarseness of tracheids and radial growth of

young Norway spruce. *European Journal of Forest Research*, 137:719–743.

Martínez-Sancho, E., Rellstab, C., Guillaume, F., et al. (2021). Post-glacial re-colonization

and natural selection have shaped growth responses of silver fir across Europe. *Science of*

the Total Environment, 779:146393.

McKenna, A., Hanna, M., Banks, E., et al. (2010). The genome analysis toolkit: a

MapReduce framework for analyzing next-generation DNA sequencing data. *Genome*

Research, 20(9):1297–1303.

Milesi, P., Berlin, M., Chen, J., et al. (2019). Assessing the potential for assisted gene flow

using past introduction of Norway spruce in southern Sweden: local adaptation and genetic

basis of quantitative traits in trees. *Evolutionary Applications*, 12(10):1946–1959.

Miriti, M. N. (2006). Ontogenetic shift from facilitation to competition in a desert shrub.

Journal of Ecology, 94(5):973–979.

Montwé, D., Isaac-Renton, M., Hamann, A., & Spiecker, H. (2016). Drought tolerance and

growth in populations of a wide-ranging tree species indicate climate change risks for the

boreal north. *Global Change Biology*, 22(2):806–815.

Niklas, K. J. (1993). Influence of tissue density-specific mechanical properties on the scaling

of plant height. *Annals of Botany*, 72(2):173–179.

Nota, K., Klaminder, J., Milesi, P., Bindler, R., Nobile, A., van Steijn, T., et al. (2022). Norway

spruce postglacial recolonization of Fennoscandia. *Nature Communications*, 13(1):1–9.

Nystedt, B., Street, N. R., Wetterbom, A., et al. (2013). The norway spruce genome

sequence and conifer genome evolution. *Nature*, 497:579–584.

Orsucci, M., Milesi, P., Hansen, J., Girodolle, J., Glémin, S., & Lascoux, M. (2020). Shift in

ecological strategy helps marginal populations of shepherd's purse (Capsella bursa-pastoris)

to overcome a high genetic load. *Proceedings of the Royal Society B: Biological Sciences*,

287(1927):20200463.

Palmé, A. E., Su, Q., Rautenberg, A., Manni, F., & Lascoux, M. (2003). Postglacial

recolonization and cpdna variation of silver birch, *Betula pendula*. *Molecular Ecology*,

12:201–212.

Petit, R. J., Brewer, S., Bordács, S., et al. (2002). Identification of refugia and post-glacial

colonisation routes of European white oaks based on chloroplast DNA and fossil pollen

evidence. *Forest ecology and management*, 156(1-3):49–74.

Petit, R. J., Aguinagalde, I., De Beaulieu, J.-L., et al. (2003). Glacial refugia: hotspots but not

melting pots of genetic diversity. *Science*, 300(5625):1563–1565.

Poorter, L., Bongers, L., & Bongers, F. (2006). Architecture of 54 moist-forest tree species:

traits, trade-offs, and functional groups. *Ecology*, 87(5):1289–1301.

Poorter, L. (2008). The relationships of wood-, gas-and water fractions of tree stems to

performance and life history variation in tropical trees. *Annals of botany*, 102(3):367–375.

Poorter, L., McDonald, I., Alarcón, A., et al. (2010). The importance of wood traits and

hydraulic conductance for the performance and life history strategies of 42 rainforest tree

species. *New Phytologist*, 185(2):481–492.

Preston, K. A., Cornwell, W. K., & DeNoyer, J. L. (2006). Wood density and vessel traits as

distinct correlates of ecological strategy in 51 California coast range angiosperms. *New*

Phytologist, 170(4):807–818.

Pretzsch, H., Biber, P., Schütze, G., Kemmerer, J., & Uhl, E. (2018). Wood density reduced
while wood volume growth accelerated in Central European forests since 1870. *Forest*
Ecology and Management, 429:589–616.
690
691 R Core Team (2021). R: A language and environment for statistical computing. R
692 Foundation for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/.
693
694 Riis, T., Lambertini, C., Olesen, B., Clayton, J. S., Brix, H., & Sorrell, B. K. (2010). Invasion
695 strategies in clonal aquatic plants: are phenotypic differences caused by phenotypic
696 plasticity or local adaptation? *Annals of Botany*, 106(5):813–822
697
698 Rossi, S., Cairo, E., Krause, C., & Deslauriers, A. (2015). Growth and basic wood properties
699 of black spruce along an alti-latitudinal gradient in Quebec, Canada. *Annals of Forest*
700 *Science*, 72:77–87.
701
702 Saleh, D., Chen, J., Leplé, J. C., Leroy, T., Truffaut, L., Dencausse, et al. (2022). Genome-
703 wide evolutionary response of European oaks during the Anthropocene. *Evolution letters*,
704 6(1):4–20.
705
706 Sang, Z., Sebastian-Azcona, J., Hamann, A., Menzel, A., & Hacke, U. (2019). Adaptive
707 limitations of white spruce populations to drought imply vulnerability to climate change in its
708 western range. *Evolutionary Applications*, 12(9):1850–1860.
709
710 Savolainen, O., Pyhäjärvi, T., & Knürr, T. (2007). Gene flow and local adaptation in trees.
711 *Annu. Rev. Ecol. Evol. Syst.*, 38:595–619.
712
713 Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation.
714 Nature Reviews Genetics, 14:807–820.
715
716 Spitze, K. (1993). Population structure in Daphnia obtusa: quantitative genetic and allozymic
717 variation. *Genetics*, 135(2): 367–374.
718
719 Stephens, M. (2017). False discovery rates: a new deal. *Biostatistics*, 18(2):275–294.
720

Suarez, A. V., & Tsutsui, N.D. (2008). The evolutionary consequences of biological

invasions. *Molecular ecology,* 17(1):351–360.

Tiret, M., & Milesi, P. Statistical Peak Detection for GWAS. Zenodo.

https://doi.org/10.5281/zenodo.5079556. Deposited 7 July 2021.

Tomiolo, S., & Ward, D. (2018a). Species migrations and range shifts: A synthesis of causes

and consequences. *Perspectives in plant ecology, evolution and systematics*, 33:62–77.

Tomiolo, S., & Ward, D. (2018b). Soil properties and climate mediate the effects of biotic

interactions on the performance of a woody range expander. *Ecosphere*, 9(4):e02186.

Tsuda, Y., Chen, J., Stocks, M., et al. (2016). The extent and meaning of hybridization and

introgression between Siberian spruce (Picea obovata) and Norway spruce (*Picea abies*):

cryptic refugia as stepping stones to the west? *Molecular Ecology*, 25(12):2773–2789.

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., et al. (2013). From FastQ data to high-

confidence variant calls: the genome analysis toolkit best practices pipeline. *Current

Protocols in Bioinformatics*, 43:11.10.1–11.10.33.

Van Gelder, H. A., Poorter, L., & Sterck, F. J. (2006). Wood mechanics, allometry, and life-

history variation in a tropical rain forest tree community. *New Phytologist*, 171(2):367–378.

Van Valen, L. (1973). A new evolutionary law. *Evol theory*, *1*: 1–30.

VanWallendael, A., Hamann, E., & Franks, S. J. (2018). Evidence for plasticity, but not local

adaptation, in invasive Japanese knotweed (Reynoutria japonica) in North America.

Evolutionary Ecology, 32:395–410.

Vendramin, G. G., Anzidei, M., Madaghiele, A., Sperisen, C., & Bucci, G. (2000). Chloroplast

microsatellite analysis reveals the presence of population subdivision in Norway spruce

(*Picea abies* K.). *Genome*, 43(1): 68–78.

Vidalis, A., Scofield, D. G., Neves, L. G., Bernhardsson, C., García-Gil, M. R., & Ingvarsson,

P. K. (2018). Design and evaluation of a large sequence-capture probe set and associated

SNPs for diploid and haploid samples of Norway spruce (*Picea abies*). *BioRxiv*, 291716.

Von Gadow, K., Álvarez González, J. G., Zhang, C., Pukkala, T., & Zhao, X. (2021). The
Shape and Growth of Forest Trees. In *Sustaining Forest Ecosystems* (pp. 159-219).
Springer, Cham.

Whitlock, M. C. (2008). Evolutionary inference from QST. *Molecular ecology*, 17(8):1885–

1896.

Wisz, M.S., Pottier, J., Kissling, W.D., et al. (2013), The role of biotic interactions in shaping

distributions and realised assemblages of species: implications for species distribution

modelling. *Biological Reviews*, 88:15–30.

Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: a tool for genome-

wide complex trait analysis. *The American Journal of Human Genetics*, 88(1):76–82.

Figure 1. Overall description of phenotypes and genotypes of the sample of Norway spruce. (A) Principal Component Analysis of the genotypes of the half-sib families. Families are colored according to their genetic clusters (ordered by decreasing latitude in the legend): Central and South Sweden (CSE, dark red), Russia-Baltics (RBA, light green), Northern Poland (NPL, green), Central Europe (CEU, light blue), Alps (ALP, blue), and hybrids between CSE and ALP (CSE-ALP, pink). PC1 and PC2 explained 3.42% and 0.45% of the variance, respectively. **(B)** Admixture plot of the half-sib families, with K = 3. Individuals are grouped by population, and sorted according to their coancestry coefficient. **(C)** Principal Component Analysis of the phenotypes: Diameter at Breast Height (DBH), Ring Width (RW), Wood Density (WD), Tracheid Radial Width (TRW), Tracheid Tangential Width (TTW), and Tracheid Wall Thickness (TWT). Left panel only shows variable axes and 60% confidence ellipses. On the right panel, each data (dot) is colored according to its cambial age (3^{rd} year in light blue up until 13^{th} year in dark blue). PC1 and PC2 explained 41.2% and 38.3% of the variance, respectively.

Figure 2. Genetic control of wood properties in Norway spruce. (A) ANOVA of genetic clusters in model (2) as a function of cambial age (from 3 to 13), and of traits: DBH (plain circle), RW (plain triangle), WD (plain square), TRW (circle), TTW (triangle), and TWT (square). **(B)** Principal Component Analysis of the residuals of the phenotypes according to model (2): Diameter at Breast Height (DBH), Ring Width (RW), Wood Density (WD), Tracheid Radial Width (TRW), Tracheid Tangential Width (TTW), and Tracheid Wall Thickness (TWT). Left panel only shows variable axes and 10% confidence ellipses (zoomed x2.5), where clusters were ordered in decreasing latitude from right to left. On the right panel, each data (dot) is colored according to its genetic cluster (ordered by decreasing latitude in the legend): Central and South Sweden (CSE, dark red), Russia-Baltics (RBA, light green), Northern Poland (NPL, green), Central Europe (CEU, light blue), Alps (ALP, blue), and hybrids between CSE and ALP (CSE-ALP, pink). PC1 and PC2 explained 57.5% and 18.2% of the variance, respectively. **(C)** Distribution of pairwise F_{ST} across the genome, averaged among pairs of populations. The vertical solid black line is the average F_{ST} over the genome (0.0488), and vertical dotted lines are the Q_{ST} estimated for different traits (averaged across rings): Diameter at Breast Height (DBH, red, $Q_{ST} = 0.281$), Ring Width (RW, green, $Q_{ST} = 0.302$), Wood Density (WD, brown, $Q_{ST} = 0.121$), Tracheid Radial Width

806   (TRW, turquoise, Q_{ST} = 0.0579), Tracheid Tangential Width (TTW, blue, Q_{ST} = 0.177),

807   Tracheid Wall Thickness (TWT, purple, Q_{ST} = 0.112), and Age at Breast Height (ABH, pink,

808   Q_{ST} = 0.066).

	ABH (year)	DBH (mm)	RW (mm)	WD (kg.m^{-3})	TRW (µm)	TTW (µm)	TWT (µm)
σ_F^2	7.44 x 10^{-2}	3.03 x 10^{-2}	5.13 x 10^{-3}	2.77 x 10^{-3}	5.99 x 10^{-4}	4.73 x 10^{-4}	2.47 x 10^{-3}
$\sigma_B^2 + \sigma^2$	1.65	2.15 x 10^{-1}	5.63 x 10^{-2}	1.95 x 10^{-2}	6.04 x 10^{-3}	4.91 x 10^{-3}	1.88 x 10^{-2}
h^2 (%)	17.31	49.32	33.47	49.80	36.11	35.20	46.52

811 **Table 1.** Genetic variance (σ_F^2), environmental variance ($\sigma_B^2 + \sigma^2$), and heritability (h^2) were

812 estimated with model (1). Traits: Calendar age at Breast Height (ABH), Diameter at Breast

813 Height (DBH), Ring Width (RW), Wood Density (WD), Tracheid Radial Width (TRW),

814 Tracheid Tangential Width (TTW), and Tracheid Wall Thickness (TWT).