The Cif proteins from *Wolbachia* prophage WO modify sperm genome integrity to establish cytoplasmic incompatibility

Short title: The cytoplasmic incompatibility proteins modify sperm genome integrity

Authors: Rupinder Kaur[1,2,*], Brittany A. Leigh[1,2,*], Isabella T. Ritchie[1,2] and Seth R. Bordenstein[1,2,3,4,5]

Affiliations:

[1] Vanderbilt University, Department of Biological Sciences, Nashville, TN, USA

[2] Vanderbilt University, Vanderbilt Microbiome Innovation Center, Nashville, TN, USA

[3] Vanderbilt University Medical Center, Department of Pathology, Microbiology & Immunology, Nashville, TN, USA

[4] Vanderbilt University Medical Center, Vanderbilt Institute for Infection, Immunology and Inflammation, Nashville, TN, USA

[*] Co-first authors

[5] Correspondence to:

Seth R. Bordenstein, Nashville, TN, 37235, Phone 615.322.9087, e-mail

s.bordenstein@vanderbilt.edu

Abstract

Inherited microorganisms can selfishly manipulate host reproduction to drive through

populations. In *Drosophila melanogaster*, germline expression of the native *Wolbachia* prophage

WO proteins CifA and CifB cause cytoplasmic incompatibility (CI) in which embryos from

infected males and uninfected females suffer catastrophic mitotic defects and lethality; however,

in infected females, CifA expression rescues the embryonic lethality and thus imparts a fitness

advantage to the maternally-transmitted *Wolbachia*. Despite widespread relevance to sex

determination, evolution, and vector control, the mechanisms underlying when and how CI

impairs male reproduction remain unknown and a topic of debate. Here we use cytochemical,

microscopic, and transgenic assays in *D. melanogaster* to demonstrate that CifA and CifB

proteins of *w*Mel localize to nuclear DNA throughout the process of spermatogenesis. Cif

proteins cause abnormal histone retention in elongating spermatids and protamine deficiency in

mature sperms that travel to the female reproductive tract with Cif proteins. Notably, protamine

gene knockouts enhance wild type CI. In ovaries, CifA localizes to germ cell nuclei and

cytoplasm of early-stage egg chambers, however Cifs are absent in late-stage oocytes and

subsequently in fertilized embryos. Finally, CI and rescue are contingent upon a newly annotated

CifA bipartite nuclear localization sequence. Together, our results strongly support the Host

Modification model of CI in which Cifs initially modify the paternal and maternal gametes to

bestow CI-defining embryonic lethality and rescue.

Introduction

Numerous animal species harbor heritable microorganisms that alter host fitness in beneficial and harmful ways. The most common, maternally-inherited bacteria are *Wolbachia* that typically reside intracellularly in reproductive tissues of both male and female arthropods. Here, they induce reproductive modifications with sex specific effects such as cytoplasmic incompatibility (CI) that can selfishly drive the bacteria to high frequencies in host populations. CI also notably yields important consequences on arthropod speciation [1–3] and vector control strategies [4–9] by causing lethality of embryos from *Wolbachia*-infected males and uninfected females. As CI is rescued by *Wolbachia*-infected females with the same strain [10,11], the phenotype accordingly imparts a relative fitness advantage to infected females that transmit the bacteria [12].

Two genes, *cytoplasmic incompatibility factors cifA* and *cifB*, occur in *Wolbachia* prophage WO within the eukaryotic association module enriched for arthropod functions and homology [13–15]. We previously demonstrated that dual, transgenic expression of *cifA* and *cifB* from *w*Mel *Wolbachia* in *Drosophila melanogaster* males induces CI, while single expression of *cifA* in females rescues CI [16,17]. These results form the basis of the Two-by-One genetic model of CI for several, but not all, strains of *Wolbachia* [13,15,18,19]. At the cellular level, CI-defining lethality associates with chromatin defects and mitotic arrest within the first few hours of embryonic development. Normally after fertilization, the sperm-bound 'protamines' are removed in the embryo and replaced by maternally supplied 'histones', resulting in the rapid remodeling of the paternal chromatin [20]. However, during CI, there is a delay in the deposition of maternal histones onto the paternal chromatin, resulting in altered DNA replication, failed

chromosome condensation, and various mitotic defects that generate embryonic death [13,21–

27].

The incipient, pre-fertilization events in the reproductive tissues that establish CI and

rescue remain enigmatic and under recent debate, namely whether (i) Cifs modify the paternal

genome during spermatogenesis (Host-Modification model) or embryogenesis (Toxin-Antidote

model) and (ii) rescue occurs or does not occur by CifA binding CifB in the embryo [28,29].

These two key observations can conclusively differentiate the mechanistic models of CI, though

they have not been explicitly addressed to date. Notably, paternal transmission of the proteins

from sperm to embryo may occur under either mechanistic model [11,29]. Recent work proposed

the Toxin-Antidote model is operational using transgenic, heterologous expression of Cif

proteins from *w*Pip *Wolbachia* [30]. In this study, the CifB$_{w\text{Pip}}$ protein paternally transfers to the

fly embryo and associates with DNA replication stress of the paternal genome in the embryo.

However, without the ability to rescue this non-native, transgenic CI and thus visualize CifA-

CifB binding in the rescue embryo, these interesting results do not yet resolve the predictions of

the two models. Additionally, the paternal DNA replication defects observed in the embryos may

be established before fertilization by the Cif proteins, in concordance with Host Modification

model.

Here, we develop antibodies to localize Cif proteins from *w*Mel *Wolbachia* during *D.*

melanogaster gametogenesis and embryogenesis and then perform genome integrity

measurements of developing sperm across transgenic, mutant, and wild type treatment groups.

We describe the following cell biological and gametic chromatin events underpinning the Host

Modification model of CI and rescue: (i) CifA and CifB proteins localize to the developing

sperm nuclei from early spermatogonium stage to late elongating spermatids; (ii) In mature

sperm, CifA associates with sperm tails and occasionally occurs in the acrosome, whereas CifB

localizes to the acrosome in all mature sperms; (iii) Cifs increase histone retention in developing

spermatids and decrease protamine levels in mature sperms; (iv) Both CI and rescue are

dependent upon a newly annotated bipartite nuclear localization signal (bNLS) in CifA that

impacts nuclear localization and sperm protamine levels; (v) During copulation, both Cif

proteins transfer with the mature sperm exhibiting reduced protamine levels; importantly,

protamine mutant flies enhance wild type CI; (vi) In the ovaries, CifA is cytonuclear in germline

stem cells and colocalizes with *Wolbachia* in the nurse cell cytoplasm; (vi) CifA is absent in the

embryos, and thus rescue must be established in oogenesis independently of CifA's presence in

the embryo. Taken together, results demonstrate that prophage WO-encoded Cif proteins from

Wolbachia invade gametic nuclei to modify chromatin integrity at the histone to protamine

transition stage in males. At the mechanistic level, results support the Host Modification model

of CI and rescue whereby native Cifs wield their impacts prior to fertilization.

Results

CifA and CifB invade sperm nuclei during spermatogenesis and spermiogenesis

To evaluate the cellular localization of the Cif proteins, we generated monospecific

polyclonal antibodies for visualizing the proteins in reproductive tissues (Fig S1). In *D.*

melanogaster males, the sperm morphogenesis process is subdivided into two events (i)

Spermatogenesis including mitotic amplification and meiotic phases, and (ii) Spermiogenesis, a

post-meiotic phase. During spermatogenesis, the germline stem cell undergoes four rounds of

synchronous mitotic divisions to produce 16 precursor cells called spermatogonia. The

spermatogonia then grow and become spermatocytes [31]. After the growth phase, the

spermatocytes divide by meiosis and differentiate into 64 haploid round onion spermatids. Post-meiosis, the round sperm nuclei elongate to gradually change their shape accompanied by reorganization of the chromatin during the canoe stage [32]. This results in an individualization complex forming slim, needle-shaped sperm nuclei with reduced volume [31,32]. Elongation and individualization of the spermatids is the final stage of spermiogenesis, after which the mature sperms are transported to the seminal vesicle [31,33].

CifA, but not CifB, localizes in the germline stem cells at the apical end of testes in <8 hr old *cifA* and *cifB* transgene-expressing (Fig 1) and wild type *w*Mel+ males (Fig S2). CifA and CifB were both detected in the nuclei of mitotic spermatogonium and spermatocytes. CifA is more abundant than CifB in the spermatogonium stage (Fig S3A, Table S1). In the post-meiotic, round onion spermatids, clusters of both CifA and CifB are adjacent to the nuclei. In the elongating canoe-shaped spermatids, CifA and CifB localize apical to the sperm head nucleus, in what is likely the acrosome (Fig 1, Fig S2). CifB is present in all of the spermatid nuclei, whereas CifA is present on average in 39% of the elongating spermatids per sperm bundle (Fig S3B). During the elongating canoe stage, chromatin-bound histones are typically removed and replaced with protamines to yield compact nuclear packaging and chromatin organization of sperm DNA [32]. After nuclear compaction is complete, neither of the Cif proteins are detectable in late spermatid needle-shaped nuclei (Fig 1) and in the mature sperms from the seminal vesicle (Fig S4), indicating either the Cif proteins are fully stripped, or they might not be accessible by the antibodies when the chromatin is tightly compacted [34,35]. To evaluate Cif presence/absence in the mature sperm, we decondensed sperms after isolations from seminal vesicles of <8 hr old males (see methods) and stained them with the respective Cif antibodies. CifA is common along sperm tails in a speckled pattern (Fig 1, Fig S2) and infrequently present

in the acrosome region, on average in 45% or 0% of the mature sperm heads depending upon the

sampled seminal vesicles (Fig S3B). CifB is present in all of the acrosomal tips of the sperms

and not localized to the sperm tails (Fig 1, Fig S2).

During spermatogenesis, *Wolbachia* are stripped into the cytoplasmic waste bags, which

eliminate excess cytoplasmic material during the process of spermatid elongation [23]. Here, we

show that some CifA and CifB proteins are also stripped by the individualization complex into

the cytoplasmic waste bag (Fig S5). Since *Wolbachia* are not present in the mature sperms

[36,37], these data suggest that the Cif proteins exit *Wolbachia* cells during spermatogenesis to

possibly interact with and modify sperm DNA (see below). Taken together, these findings

demonstrate CifA and CifB proteins access *Drosophila* sperm nuclei throughout development.

Cifs cause abnormal histone retention and protamine deficiency

Since the Cifs localize to developing sperm nuclei during spermatogenesis, we

hypothesized that they may incipiently interact with nuclear DNA to impact sperm genome

integrity – a central prediction of the Host Modification model of CI [11,29]. At the histone-to-

protamine transition stage during spermatogenesis [38], histones normally undergo various post-

translational modifications (PTMs) for removal and replacement by smaller protamines for tight

chromatin reorganization [38–41]. Lack of PTMs can lead to histone-bound chromatin with

improper protamine deposition that causes paternal chromatin defects, male infertility, and

embryonic lethality [27,42,43]. Thus, the incipient defects initiated in the testes can lead to post-

fertilization catastrophes.

Utilizing a core histone antibody, we investigated histone abundance within spermatid

bundles at the late canoe stage in CI- and non-CI causing males. We detected significantly

increased histone retention in both *w*Mel+ and *cifAB*-expressing testes from <8 hr old males

compared to the negative controls (Fig 2A, Fig S6, Table S1). Single transgenic expression

showed significantly less histone-retaining bundles at this stage similar to *w*Mel- negative

controls (Fig S7A, Table S1). To detect if abnormal histone retention is linked with protamine

deficiency in mature sperms, we next used the fluorochrome chromomycin A3 (CMA3) stain

that fluoresces upon binding to protamine-deficient regions of DNA [44,45]. Mature sperms

isolated from wild-type seminal vesicles of young (high CI-inducing) *w*Mel+ males exhibit

increased protamine deficiency relative to *w*Mel- males (Fig 2B, Table S1).

To investigate if lack of protamines associates with CI, we isolated sperms from <8 hr old

males with a protamine A and B knockout mutant line. We show that protamine mutants, both in

the presence (ΔProt+) and absence (ΔProt-) of *Wolbachia,* also exhibit a significant increase in

fluorescence relative to *w*Mel- (Fig 3A, Table S1). Moreover, a key outcome of higher protamine

deficiency in ΔProt+ males with *Wolbachia* is an increase in CI compared to wild type *w*Mel+

CI, under the same experimental setup (Fig 3B, Table S2). These findings indicate an additive

effect by *Wolbachia* and ΔProt knockouts on the protamine deficiency and CI penetrance.

Notably, ΔProt- males do not recapitulate CI on their own; thus, the protamine deficiency is not

the sole cause of CI and must operate in conjunction with other CI modifications. Consistent

with these results, 7-day old *w*Mel+ males that express almost no CI exhibit a similarly weak

protamine deficiency level to *w*Mel- males, as expected (Fig S8A, Fig S8B and Table S2).

Moreover, transgene analyses specify both single and dual expression of CifA and CifB cause

protamine deficiencies at significantly higher levels than negative controls of *w*Mel- and a non-

CI transgene (Fig S7B, Table S1). Since singly expressed Cifs do not cause CI in *D.*

melanogaster [16] (Fig S7C, Table S2), additive effects on the protamine deficiency and/or

histone retention due to abnormal PTMs may be required to fully establish CI.

Both CI and rescue are dependent upon a CifA bipartite nuclear localization signal

Based on the cNLS mapping tool for nuclear localization signals [46], CifA amino acids

harbor a predicted bipartite nuclear localization sequence (bNLS) (Table S3) in the most

conserved region of the protein [13,47,48] that is under strong purifying selection [17]. As

nuclear localization signals bind to the extended surface groove of nuclear transport protein

importin-α, also known as karyopherin-α [49], we hypothesized that sperm nuclear localization

of CifA and CI and rescue are dependent on the bNLS. To test this hypothesis, we mutagenized

two bNLS sequences with alanine substitutions (aa189-190 for NLS1 (denoted *cifA$_{189}$*), aa224-

225 for NLS2 (denoted *cifA$_{224}$*)), and we additionally deleted the entire bNLS region (*cifA$_{\Delta bNLS}$*)

(Fig 4A). The bNLS deletion also corresponds to the weakly predicted catalase-rel domain in

CifA [47,48].

Each bNLS mutant, individually and together (*cifA$_{189;224}$*), was dually expressed in testes

with transgenic, intact *cifB* to assess CI and singly expressed in females to assess rescue.

Transgenic *cifA$_{189}$* expression significantly reduced CI and rescue as previously reported (Fig

4B,C and Table S2) [48]. Conversely, transgenic *cifA$_{224}$* expression showed no significant

difference from the controls in either CI or rescue, suggesting this region has little to no impact.

However, when both mutants are expressed in *cifA$_{189;224}$* or when the entire bNLS is deleted, CI

and rescue are strongly inhibited (Fig 4B,C and Table S2). These results highlight the importance

of the nuclear localization sequence in inducing CI as well as rescue. To determine if the lack of

CI induction is due to non-nuclear localization of CifA protein, we used the deletion mutant

cifA$_{\Delta bNLS}$ with wild type *cifB* to demonstrate that in contrast to its normal, nuclear localization

(Fig 1), it mislocalizes to the cytoplasm of onion stage spermatids rather than the nuclei (Fig

4D). Additionally, to test if deletion of the bNLS impacts sperm genomic integrity, we

performed CMA3 based protamine-deficiency assay (as described above) and found reduced

protamine deficiency levels in matures sperms due to non CI-causing *cifA$_{\Delta bNLS}$;B* line compared

to CI-causing *cifAB* (Fig 4E, Table S1), providing further evidence that protamine deficiency is

linked to CI. Overall, these data provide previously unknown findings that a functional nuclear

localization sequence and CifA nuclear localization impacts CI, rescue, and sperm protamine

levels.

Cifs cause a paternally-transferred protamine deficiency

To investigate if Cif proteins and/or the genome integrity modifications transfer

paternally to the female reproductive tract, single male:female pairwise matings were set up for

CI and rescue crosses. After 4 hrs of mating, we isolated the whole uterus (Fig 5A) including the

sperm storage organs (spermathecae (SP) and seminal receptacle (SR)). Following sperm

decondensation, antibody staining, and microscopy, we observed two key results. First, both

CifA and CifB proteins transfer with sperm tails and heads, respectively, to the sperm storage

organs of *Wolbachia*-free females (Fig 5B). Second, the CI-associated sperm protamine

deficiency induced by *w*Mel+ and *cifAB*-expressing males transfers and persists in the sperms

isolated from SP and SR of *Wolbachia*-free, mated females (Fig 5C,D and Table S1). These

findings connect a paternally-transferred sperm modification with the activity of *Wolbachia* and

the Cifs themselves. Results strongly support the Host Modification model of CI since a Cif-

induced sperm modification established in the testes transfers to the female reproductive tract.

CifA is present in early oogenesis and absent from mature eggs and rescue embryos

Expression of *cifA* alone in the ovaries rescues CI [16,17], yet how CifA protein mediates

rescue is unknown and central to further differentiating the mechanistic models of CI. For

instance, CifA in females may modify reproductive cell biology to nullify CI-inducing sperm

modifications in the embryo (Host-Modification), or alternatively, CifA may directly bind CifB

in the embryos to prevent its proposed CI toxicity (Toxin-Antitoxin). Using CifA antibodies, we

show in *w*Mel+ and *cifA* transgene-expressing ovaries that CifA protein is cytonuclear and

specifically localizes to cyst DNA in region 1 of the germarium (Fig 6A), indicative of nuclear

access in ovaries similar to that in testes (Fig 1). Cystoblast in the germarium undergoes rounds

of mitotic divisions to produce oocyte and nurse cells [50,51]. Along egg chamber stages 2-8 of

*w*Mel+ females, CifA colocalizes with *Wolbachia* in the nurse cells and oocyte cytoplasm (Fig

6A). While *Wolbachia* are abundant in the stage 10 egg chamber, CifA is notably absent. In

transgenic females, CifA is also primarily detected in the germarium and cytoplasm of the early

egg chambers and absent in late egg chambers (Fig 6A, Fig S9A). Presence of high levels of

CifA in *Wolbachia*-infected eggs is proposed to rescue CI; importantly, we did not detect CifA in

~30-60 min old rescue embryos during early mitotic divisions (Fig 6B). Moreover, CifA was not

detected in 1-2 hr old embryos (Fig S9B), whereas the positive control histone signals colocalize

with embryonic DNA at both embryonic developmental stages.

CifB from *w*Pip was recently shown to be paternally-transferred to the CI embryos [30].

We evaluated if *w*Mel CifB colocalizes with mitotic, embryonic DNA after fertilization. CifB is

absent from embryonic nuclei and the cytoplasm of CI embryos (Fig 6B), suggesting that CifB is

not inherited with the paternal DNA to the embryo. The absence of CifA and CifB in late wild

type and transgenic eggs and embryos indicates that host gametic changes prime the embryo for

CI and rescue before fertilization, as predicted by the Host Modification model of CI. This

inference is also consistent with previous studies where ovarian, rather than embryonic,

transgenic *cifA* expression rescues CI [13,15–17,52].

Discussion

236
At the genetic level, dual expression of *cifA and cifB* or single expression of *cifB* can

recapitulate CI; and *cifA* rescues CI [13,15–19,52]. However, the cellular and mechanistic bases

of CI and rescue remain unresolved and the subject of several questions: When and where do the

Cif proteins localize in testes to potentiate CI? Are the Cifs transferred to the embryo already

modified for CI-defining defects? Do CifA and CifB bind in the embryo to rescue lethality?

Here, we establish that both CifA and CifB proteins invade nuclei of developing spermatids and

modify paternal genome integrity by altering the normal Histone-Protamine transition process.

Specifically, dual CifA and CifB expression induces abnormal histone retention and protamine

deficiency in CI-causing male gametes to induce CI. Moreover, knocking out protamines

enhances wild type CI, and a nuclear localization sequence in CifA is essential for CI, rescue,

and protamine deficiency. Finally, binding of CifA and CifB in the embryo is not evident.

Sperm genome compaction is normally achieved during the post-meiotic canoe phase of

spermatogenesis, when histones are replaced by protamines [53]. This compaction process is

highly conserved from flies to humans [54–56] and plays a crucial role in successful fertilization

and embryonic development [57,58]. Histone marks are carriers of transgenerational epigenetic

information [59], and changes in the sperm epigenome can lead to detrimental consequences

including early embryonic lethality and birth defects [59,60]. Histones undergo various post-

translational modifications (PTMs) such as ubiquitination, methylation, phosphorylation and

acetylation before degradation and removal from the sperm chromatin [38]. Therefore,

abnormally-retained histones could result from aberrant PTMs in Cif-expressing flies. In a

protein interactome screen, both ubiquitin and histone H2B were determined as binding host

candidates to CifB [52]. Thus, it is possible that inhibition of the histone ubiquitination process

mediates abnormal histone retention. Additionally, histone acetylation during the canoe stage of

spermatogenesis is required for histone eviction in *Drosophila* [61]. Indeed, reduced acetylation

levels lead to abnormal histone retention and protamine deficiency, which causes embryonic

inviability in flies [61] and in mammals [40,62,63]. Future research investigating what specific

PTMs are altered at the Histone-Protamine transition stage will help elucidate the molecular

pathway(s) leading to CI.

The paternally-derived Cif proteins travel with the sperms to the female reproductive

tract, where CifB is present in the acrosomal region and CifA occurs along the tail. Notably, both

Cifs are not evident in the embryos. While the presence of paternally-transferred Cifs is

ambiguous to the mechanistic models of CI [11,29] and recently confirmred in another

transgenic study [30], it is the presence of CifA-CifB binding in the rescue embryo that would

support the Toxin-Antitoxin model. However, there is no evidence of this binding phenomenon

in the embryos to date. Thus, we conclude that Cifs act before fertilization to prime the sperm

chromatin and incipiently launch CI. Paternal effect proteins can modify sperm in various

systems to bestow embryonic defects, even though the proteins themselves do not transfer to the

embryos [60,64,65].

In *Drosophila*, the sperm enters the egg with an intact membrane [66]. Therefore,

absence of Cifs in the embryos raises the question at what point the presence of CifA along the

tail and CifB in the acrosome is lost before the sperm enters the egg. One possible explanation is

that the Cif proteins are released from the sperm upon exocytosis of the acrosome. The

acrosome, best known as a secretory vesicle, undergoes exocytosis and releases its contents to

facilitate sperm-egg binding in mammals [67,68]. Though not well characterized in insects,

studies in the house fly *Musca domestica* suggest loss of the sperm plasma membrane before

entry into the egg, followed by exocytosis of acrosomal contents during passage of the sperm

through the egg micropyle [69].

CifA is absent in *Wolbachia*-infected and transgenic embryos, which indicates that rescue

is established during oogenesis under the Host Modification model, and CifA thus does not bind

and nullify CifB from *w*Mel *Wolbachia* in the embryos. Indeed, a CifA mutant in the newly

annotated nuclear localization sequence ablates rescue, suggesting access to ovarian nuclei is

important for rescue. Interestingly, structures of CifA and CifB support binding of the two

proteins [70], yet in light of these results here, the CifA-CifB binding may be central to CI

induction instead of rescue. Moreover, mutating amino acid sites across the length of the CifA

protein, including binding and non-binding residues, generally ablates CI and rescue [48,70].

Thus, it is possible that CifA mutants that ablate rescue do so by altering the protein structure,

function, and/or location of ovarian targets to modify, rather than the embryonic binding of CifA

to CifB *in vivo*. Future work will be important to resolve how CifA primes oogenesis to alter

specific cell biological and biochemical events that underpin rescue.

Once fertilization occurs, protamines from the paternal chromatin must be removed and

replaced by maternal histones to decondense and activate the chromatin of the developing

embryo [71]. Interestingly, post-fertilization delays in maternal H3.3 histone deposition occur in

CI embryos [72]. The delay may in part be due to pre-loaded paternal histones or altered paternal

epigenome information leading to mistiming of maternal histone deposition, hence causing CI. Thus, we propose that a genome integrity network involving histones, protamines, and possibly other factors in the gametes may be a common and defining feature underpinning the onset of CI and rescue.

Altogether, discovery of nuclear-targeting Cif proteins in male and female gametes establishes new insights on the early cell biological and biochemical steps that underpin the CI drive system with relevance to arthropod speciation and pest control [11]. In addition to disentangling the reproductive events of the Cif proteins that control gametogenesis and embryogenesis, the evidence specifies that the Cif proteins modify sperm genomic integrity and transfer paternally, but they themselves do not enter and bind each other in the embryo to enable rescue. These findings are consistent with the Host Modification model of CI by *w*Mel *Wolbachia*. More generally, as there are no previous reports of prophage proteins invading animal gametic nuclei to impair the histone-protamine transition during spermatogenesis, the findings have implications for expanding an appreciation of prophage-bacteria-eukaryote interactions into the realm of animal reproduction.

Material and Methods

Cif proteins antibody development

Conserved amino acid regions of CifA and CifB proteins from *w*Mel *Wolbachia* were previously identified [47]. Using these regions, monospecific polyclonal antibodies were commercially generated by Pacific Immunology through injection of three synthesized and conserved short (20 aa) peptides for each protein into rabbits. Sequences of peptides were Cys-EYFYNQLEEKDKEKKLTE for CifA, and Cys-DENPPENLLSDQTRENFRR for CifB. The

resulting *α*-CifA and *α*-CifB antibodies were evaluated using an enzyme-linked immunosorbent

assay, and titers were determined to be higher than 1:500,000 for each antibody. Using standard

protocols of the Invitrogen WesternDot kit, antibody specificity to *w*Mel+ samples was verified

using western blots (1:1000-fold antibody dilution) on protein isolated from homogenates of 50

testes pairs (0-8 hour old males) and 10 ovary pairs (6 day old females) from *w*Mel+ (positive),

*w*Mel- (negative control), and *cifAB* transgenic (positive) flies. The correct size band was only

detected from *w*Mel+ and *cifAB* reproductive tissues (Fig S1). Because the antibodies were

generated in the same animal, all subsequent labeling was done with individual antibodies.

NLS identification

CifA amino acid sequences from known *Wolbachia* and close relatives were input into

the cNLS Mapper software [73] to identify putative NLS sequences within each protein (Table

S3). cNLS Mapper identifies sequences specific to the importin α/β pathway. A cut-off score of

4 was applied to all sequences. Higher scores indicate stronger NLS activities. Scores >8 indicate

exclusive localization to the nucleus, 7-8 indicate partial localization to the nucleus, 3-5 indicate

localization to both the nucleus and the cytoplasm, and score 1-2 indicate localization

exclusively to the cytoplasm. Predicted NLS sequences are divided into monopartite and

bipartite classes. Monopartite NLSs contain a single region of basic residues, and bipartite NLSs

contain two regions of basic residues separated by a linker region.

Development of transgenic lines

A *cifA* variant was synthesized de novo at GenScript and cloned into a pUC57 plasmid as

described previously [48]. Site-directed mutagenesis was performed by GenScript to produce the

mutants outlined in Figure 5. The $cifA_{189}$ variant was first described in Shropshire et al. [48] as

$cifA_2$. UAS transgenic *cifA* mutant flies were then generated using previously established

protocols [13]. Briefly, GenScript sub-cloned each gene into the pTIGER plasmid, a pUASp-

based vector designed for germline-specific expression. Transgenes were then integrated into

y^1 M{vas-int.Dm}ZH-2A w*; P{CaryP}attP40 attachment sites into the *D. melanogaster*

genome using PhiC31 integrase via embryonic injections by BestGene. At least 200 embryos

were injected per transgenic construct, and successful transformants were identified based on red

eye color gene included on the pTIGER plasmid containing the transgene. All sequences are

reported in Table S4.

Fly rearing and strains

D. melanogaster stocks y^1w* (BDSC 1495), *nos*-GAL4:VP16 (BDSC 4937), UAS

transgenic lines homozygous for *cifA*, *cifB*, *cifAB, WD0508* [13] and Protamine mutant

(w[1118]; ΔMst35B[floxed], Sco/CyO) [74] were maintained on a 12-hour light/dark cycle at

25^0C and 70% relative humidity on 50mL of standard media. Uninfected protamine mutant line

was generated by three generations of tetracycline treatment (20 μg/ml in 50 ml of fly media) as

described in previous studies [13], followed by two rounds of rearing on standard food media

before using in the experiments. Infection status for all lines was regularly confirmed by PCR

using Wolb_F and Wolb_R3 primers [75].

<u>*Hatch rates*</u>

Parental flies were either wild type uninfected (*w*Mel-) or infected (*w*Mel+) with

Wolbachia or transgene-expressing with no *Wolbachia* infection. Uninfected transgenic flies

were generated previously [13,17]. Paternal grandmother age was controlled to 9-11 days for

expression of naturally high penetrance of *w*Mel CI [76]. Parental transgenic males were

generated through crossing *nos*-Gal4:VP16 virgin females (aged 9-11 days) to UAS-*cif*

transgenic, uninfected males [76]. Mothers were aged 6-9 days before crossing, while father

males first emerged between 0-8 hours were used in hatch rates and tissue collections to control

for the younger brother effect associated with lower CI penetrance [13,77].

Hatch rates were set up as described previously [13,17]. Briefly, a male and female pair

was placed in an 8oz, round bottom, polypropylene *Drosophila* stock bottle with a grape juice-

agar plate containing a small amount of yeast placed at the base and secured with tape. These

bottles were then placed in a 25°C incubator overnight to allow for courting and mating. The

following day, these plates were discarded and replaced with new grape juice-agar plates with

fresh yeast. After an additional 24 hours, the plates were removed, and the embryos were

counted. The embryo plates were then incubated for 36 hours at 25°C before the total number of

unhatched embryos were counted. Any crosses with fewer than 25 embryos laid were discarded

from the analyses. Statistical significance (p<0.05) was determined by a Kruskal-Wallis test and

Dunn's multiple test correction in GraphPad Prism 7. All p-values are listed in Table S2.

Immunofluorescence: testes and seminal vesicles

Siblings from the hatch rate (males 0-8 hours) were collected for testes dissection in ice-

cold 1x PBS solution. Tissues were fixed in 4% formaldehyde diluted in 1x PBS for 30 min at

room temperature and washed in 1x PBS-T (1x PBS + 0.3% TritonX-100) three times for 5 min

each. Tissues were then blocked in 1% BSA in PBS-T for 1 hour at room temperature. They

were then incubated with 1° antibody (α-CifA 1:500 OR α-CifB 1:500) overnight at 4°C

rotating. After washing in 1x PBS-T three times for 5 min each at room temperature, they were

incubated with 1:1000 dilution Goat anti-rabbit Alexa Fluor 594 secondary antibody (Fisher

Scientific, Cat#A11037) for 4 hours at room temperature in the dark. Tissues were then washed

three times for 5 minutes each in 1X PBS-T and mounted on slides. To stain the nuclear DNA,

0.2mg/mL of DAPI was added to the mounting media before the coverslip was gently placed

over the tissue and excess liquid wiped away. Slides were allowed to dry overnight in the dark

before viewing on the Zeiss LSM 880 confocal microscope. All images were acquired with the

same parameters for each line and processed in ImageJ as described in [78].

Decondensation of Mature Sperm Nuclei

Squashed seminal vesicles collected from male flies (aged 0-8 hours) were treated with

10 mM DTT, 0.2% Triton X-100, and 400 U heparin in 1X PBS for 30 min [34]. The slides were

then washed quickly in 1X PBS before immunofluorescence staining (see above).

Immunofluorescence and quantification: Histones

Testes from male flies (aged 0-8 hours) were fixed and stained as described above for

testes. The tissues were stained with a core histone antibody (Millipore Sigma, Cat#MABE71)

(1:1000) and imaged on a Keyence BZ-800 microscope. Total late canoe stage sperm bundles

were quantified in each testis, and those that retained histones were determined. Ratios of late

canoe stage bundles containing histones relative to total bundles from each individual testis were

graphed in GraphPad Prism 7. Statistical significance ($p < 0.05$) were determined by pairwise

comparisons based on Kolmogorov-Smirnov test and multiple comparisons based on a Kruskal-

Wallis test and Dunn's multiple test correction in GraphPad Prism 7.

Sperm isolation and chromomycin A3 staining/quantification

Seminal vesicles were collected from male flies (aged 0-8 hours for 1-day old flies and 7-days for older flies) and placed on a microscope slide in ice-cold 1x PBS. Sperm was extracted on the slide using forceps and fixed in 3:1 vol/vol methanol:acetic acid at 4°C for 20 min. Excess solution was then removed, and the slide was air dried. Each slide was treated in the dark for 20 min with 0.25mg/mL of CMA3 in McIlvain's buffer, pH 7.0, with 10mM MgCl2. Sperm was then washed in 1x PBS, mounted, and imaged using a Keyence BZ-X700 Fluorescence microscope. All images were acquired with the same parameters for each line and did not undergo significant alteration. Fluorescence quantification was performed by scoring fluorescent pixels in arbitrary units (A.U.) within individual sperm heads using ImageJ as per the details described in [78], and calculated fluorescence intensity per sperm head was graphed. Statistical significance ($p < 0.05$) was determined by a Kruskal-Wallis test and Dunn's multiple test correction in GraphPad Prism 7. All of the experiments involving CMA3 staining were performed at 21^0C instead of 25^0C. CI hatch rate assays were run in parallel to ensure that CI and rescue phenotypes are not impacted due to changed temperature conditions.

Immunofluorescence: ovaries

Ovaries from females (6 days old) were dissected in 1x PBS on ice and processed as described previously [77,79]. Tissues were blocked in 1% BSA in PBS-T for 1 hour at room temperature and were first incubated with α-CifA (1:500) primary antibody at 4^0C overnight. After washing in 1x PBS-T three times for 5 min each at room temperature, they were incubated with 1:1000 dilution Alexa Fluor 488 secondary antibody (Thermo Fisher Scientific,

Cat#A11034) for 4 hours at room temperature in the dark. Samples were then rinsed properly

and blocked again before incubating with α-ftsZ (1:150) primary antibody (a kind gift from Dr.

Irene Newton) to stain *Wolbachia* at 4^0C overnight. After washing in 1x PBS-T three times,

samples were incubated with second secondary antibody (Alexa Fluor 594) for 4 hours in the

dark. Since both CifA and ftsZ antibodies were generated in the same animal, we used secondary

antibodies conjugated to two distant fluorophores to distinguish specific signals. Tissues were

then washed three times for 5 minutes each in 1X PBS, stained with DAPI to label nuclear DNA

and mounted on slides. Slides were allowed to dry overnight in the dark before viewing on the

Zeiss LSM 880 confocal microscope.

Immunofluorescence: embryos

After 24 hours of mating, plates were switched, and embryos were collected every 30

minutes. Embryos were collected in a 100μm mesh basket in embryo wash solution. To remove

the chorion, the basket was placed in 50% bleach for 3 min and then rinsed with 1x PBS. The

embryos were then transferred to 50:50 4% paraformaldehyde (PFA) and heptane in a

microcentrifuge tube and rotated for 20 min at room temperature. Tubes were then removed from

the rotator, and the heptane and PFA were allowed to separate before the bottom PFA phase was

carefully removed. Methanol was added to the remaining heptane, and the tube was shaken

vigorously for 20 seconds before the embryos settled to the bottom and solution was removed. A

new volume of methanol was added to the embryos, and they were allowed to settle to the

bottom of the tube. Methanol was removed, and all blocking, staining, and imaging steps were

carried out for testes and ovary tissues above.

Acknowledgments: The authors thank Jennifer Battle for her assistance in fly collections and

staining for sperm integrity assays, Alex Mansueto for his assistance in hatch rates, and Sarah

Bordenstein, Dylan Shropshire, and Luis Mendez for providing helpful feedback on the

manuscript. We thank Dr. Janna McLean for sending the Protamine mutant fly line for

conducting the experiments. We thank Dr. Irene Newton for sharing *Wolbachia* antibody and

providing useful feedback on the preprint version of this manuscript. We also thank the Cell

Imaging Shared Resource (CISR) at Vanderbilt for imaging assistance.

Funding: Digestive Disease Research Center Scholarships S1848284, S1848300, S1883559,

Vanderbilt-Ingram Cancer Center Scholarships S1871288, S1848887, S1848952, National

Institutes of Health Awards R01 AI132581 and AI143725 to S.R.B., F32 Ruth Kirschstein

Postdoctoral Fellowship to B.A.L., Vanderbilt Microbiome Initiative to S.R.B, and NIH grant

DK020593 to CISR.

Author contributions: Conceptualization, B.A.L. and S.R.B.; Methodology, B.A.L., R.K. and

S.R.B.; Investigation, B.A.L., R.K., and I.T.R.; Writing – Original Draft, B.A.L., R.K. and

S.R.B.; Writing – Review & Editing, B.A.L., R.K., I.T.R. and S.R.B.; Funding Acquisition,

B.A.L. and S.R.B.; Resources, S.R.B.; Supervision, S.R.B.

Competing interests: Authors declare no competing interests.

Data and materials availability: All data are available in the main text or the supplementary

materials. Unique biological materials will be available upon request.

References:
489
490   1.   Bordenstein SR, Ohara FP, Werren JH. Wolbachia-induced incompatibility precedes other

491   hybrid incompatibilities in Nasonia. Nature. 2001;409: 707–710. doi:10.1038/35055543

492   2.   Charlat S, Hurst GDD, Merçot H. Evolutionary consequences of Wolbachia infections.

493   Trends Genet. 2003;19: 217–223. doi:10.1016/S0168-9525(03)00024-6

494   3.   Jaenike J, Dyer KA, Cornish C, Minhas MS. Asymmetrical reinforcement and Wolbachia

495   infection in Drosophila. PLoS Biol. 2006;4: 1852–1862.

496   doi:10.1371/journal.pbio.0040325

497   4.   Bian G, Xu Y, Lu P, Xie Y, Xi Z. The Endosymbiotic Bacterium Wolbachia Induces

498   Resistance to Dengue Virus in Aedes aegypti. PLoS Pathog. 2010;6: e1000833.

499   doi:10.1371/journal.ppat.1000833

500   5.   Frentiu FD, Zakir T, Walker T, Popovici J, Pyke AT, van den Hurk A, et al. Limited

501   Dengue Virus Replication in Field-Collected Aedes aegypti Mosquitoes Infected with

502   Wolbachia. PLoS Negl Trop Dis. 2014;8. doi:10.1371/journal.pntd.0002688

503   6.   Walker T, Johnson PH, Moreira LA, Iturbe-Ormaetxe I, Frentiu FD, McMeniman CJ, et

504   al. The wMel Wolbachia strain blocks dengue and invades caged Aedes aegypti

505   populations. Nature. 2011;476: 450–455. doi:10.1038/nature10355

506   7.   Bourtzis K, Dobson SL, Xi Z, Rasgon JL, Calvitti M, Moreira LA, et al. Harnessing

507   mosquito-Wolbachia symbiosis for vector and disease control. Acta Trop. 2014;132.

508   doi:10.1016/j.actatropica.2013.11.004

509   8.   Hughes GL, Koga R, Xue P, Fukatsu T, Rasgon JL. Wolbachia infections are virulent and

510   inhibit the human malaria parasite Plasmodium falciparum in Anopheles gambiae. PLoS

511   Pathog. 2011;7: e1002043. doi:10.1371/journal.ppat.1002043

9.   O'Connor L, Plichart C, Sang AC, Brelsfoard CL, Bossin HC, Dobson SL. Open release

of male mosquitoes infected with a wolbachia biopesticide: field performance and

infection containment. PLoS Negl Trop Dis. 2012;6: e1797.

doi:10.1371/journal.pntd.0001797

10.   Taylor MJ, Bordenstein SR, Slatko B. Microbe profile: Wolbachia: A sex selector, a viral

protector and a target to treat filarial nematodes. Microbiol (United Kingdom). 2018.

doi:10.1099/mic.0.000724

11.   Shropshire JD, Leigh B, Bordenstein SR, Initiative M. Symbiont-mediated cytoplasmic

incompatibility□: What have we learned in 50 years? Elife. 2020; 1–51.

doi:10.20944/preprints202008.0350.v1

12.   Turelli M. Evolution of incompatibility-inducing microbes and their hosts. Evolution (N

Y). 1994;48: 1500–1513. doi:10.1111/j.1558-5646.1994.tb02192.x

13.   LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, et al.

Prophage WO genes recapitulate and enhance Wolbachia-induced cytoplasmic

incompatibility. Nature. 2017;543: 243–247. doi:10.1038/nature21391

14.   Bordenstein SR, Bordenstein SR. Eukaryotic association module in phage WO genomes

from Wolbachia. Nat Commun. 2016;7. doi:10.1038/ncomms13155

15.   Beckmann JF, Ronau JA, Hochstrasser M. A Wolbachia deubiquitylating enzyme induces

cytoplasmic incompatibility. Nat Microbiol. 2017;2. doi:10.1038/nmicrobiol.2017.7

16.   Shropshire JD, Bordenstein SR. Two-by-one model of cytoplasmic incompatibility:

Synthetic recapitulation by transgenic expression of cifa and cifb in drosophila. PLoS

Genet. 2019;15. doi:10.1371/journal.pgen.1008221

17.   Shropshire JD, On J, Layton EM, Zhou H, Bordenstein SR. One prophage WO gene

rescues cytoplasmic incompatibility in Drosophila melanogaster. Proc Natl Acad Sci U S

A. 2018;115: 4987–4991. doi:10.1073/pnas.1800650115

18.   Adams KL, Abernathy DG, Willett BC, Selland EK. Wolbachia cifB induces cytoplasmic

incompatibility in the malaria mosquito. 2021. doi:10.1101/2021.04.20.440637

19.   Sun G, Zhang M, Chen H, Hochstrasser M. The Wolbachia CinB Nuclease is Sufficient

for Induction of Cytoplasmic Incompatibility. bioRxiv. 2021; 2021.10.22.465375.

Available:

https://www.biorxiv.org/content/10.1101/2021.10.22.465375v1%0Ahttps://www.biorxiv.o

rg/content/10.1101/2021.10.22.465375v1.abstract

20.   Yang P, Wu W, Macfarlan TS. Maternal histone variants and their chaperones promote

paternal genome activation and boost somatic cell reprogramming. BioEssays. 2015;37:

52–59. doi:10.1002/bies.201400072

21.   Landmann F, Orsi GA, Loppin B, Sullivan W. Wolbachia-mediated cytoplasmic

incompatibility is associated with impaired histone deposition in the male pronucleus.

PLoS Pathog. 2009;5. doi:10.1371/journal.ppat.1000343

22.   Lassy CW, Karr TL. Cytological analysis of fertilization and early embryonic

development in incompatible crosses of Drosophila simulans. Mech Dev. 1996;57: 47–58.

doi:10.1016/0925-4773(96)00527-8

23.   Serbus LR, Casper-Lindley C, Landmann F, Sullivan W. The genetics and cell biology of

Wolbachia-host interactions. Annu Rev Genet. 2008;42: 683–707.

doi:10.1146/annurev.genet.41.110306.130354

24.   Tram U, Sullivan W. Role of delayed nuclear envelope breakdown and mitosis in

Wolbachia-induced cytoplasmic incompatibility. Science (80-). 2002;296: 1124–1126.

558   doi:10.1126/science.1070536

559   25.   Tram U, Fredrick K, Werren JH, Sullivan W. Paternal chromosome segregation during the

560   first mitotic division determines Wolbachia-induced cytoplasmic incompatibility

561   phenotype. J Cell Sci. 2006;119: 3655–3663. doi:10.1242/jcs.03095

562   26.   Callaini G, Riparbelli MG, Giordano R, Dallai R. Mitotic defects associated with

563   cytoplasmic incompatibility in Drosophila simulans. J Invertebr Pathol. 1996;67: 55–64.

564   doi:10.1006/jipa.1996.0009

565   27.   Callaini G, Dallai R, Riparbelli MG. Wolbachia-induced delay of paternal chromatin

566   condensation does not prevent maternal chromosomes from entering anaphase in

567   incompatible crosses of Drosophila simulans. J Cell Sci. 1997;110 (Pt 2: 271–280.

568   28.   Beckmann JF, Bonneau M, Chen H, Hochstrasser M, Poinsot D, Merçot H, et al. The

569   Toxin–Antidote Model of Cytoplasmic Incompatibility: Genetics and Evolutionary

570   Implications. Trends in Genetics. 2019. pp. 175–185. doi:10.1016/j.tig.2018.12.004

571   29.   Shropshire JD, Leigh B, Bordenstein SR, Duplouy A, Riegler M, Brownlie JC, et al.

572   Models and Nomenclature for Cytoplasmic Incompatibility: Caution over Premature

573   Conclusions – A Response to Beckmann et al. Trends in Genetics. 2019. pp. 397–399.

574   doi:10.1016/j.tig.2019.03.004

575   30.   Horard B, Terretaz K, Gosselin-Grenet A-S, Sobry H, Sicard M, Landmann F, et al.

576   Paternal transmission of the Wolbachia CidB toxin underlies cytoplasmic incompatibility.

577   Curr Biol. 2022. doi:10.1016/j.cub.2022.01.052

578   31.   Fuller M. The Development of Drosophila melanogaster. Cold Spring Harbor Laboratory

579   Press. Spermatogenesis. 1993; 71–147.

32.   Jayaramaiah Raja S, Renkawitz-Pohl R. Replacement by Drosophila melanogaster

Protamines and Mst77F of Histones during Chromatin Condensation in Late Spermatids

and Role of Sesame in the Removal of These Proteins from the Male Pronucleus . Mol

Cell Biol. 2005;25: 6165–6177. doi:10.1128/mcb.25.14.6165-6177.2005

33.   Fabian L, Brill JA. Drosophila spermiogenesis. Spermatogenesis. 2012;2: 197–212.

doi:10.4161/spmg.21798

34.   Eren-Ghiani Z, Rathke C, Theofel I, Renkawitz-Pohl R. Prtl99C Acts Together with

Protamines and Safeguards Male Fertility in Drosophila. Cell Rep. 2015;13: 2327–2335.

doi:10.1016/j.celrep.2015.11.023

35.   Bonnefoy E, Orsi GA, Couble P, Loppin B. The essential role of Drosophila HIRA for de

novo assembly of paternal chromatin at fertilization. PLoS Genet. 2007;3: 1991–2006.

doi:10.1371/journal.pgen.0030182

36.   Bressac C, Rousset F. The reproductive incompatibility system in Drosophila simulans:

Dapi-staining analysis of the Wolbachia symbionts in sperm cysts. J Invertebr Pathol.

1993;61: 226–230. doi:10.1006/jipa.1993.1044

37.   Clark ME, Veneti Z, Bourtzis K, Karr TL. The distribution and proliferation of the

intracellular bacteria Wolbachia during spermatogenesis in Drosophila. Mech Dev.

2002;111: 3–15. Available: http://www.ncbi.nlm.nih.gov/pubmed/11804774

38.   Rathke C, Baarends WM, Jayaramaiah-Raja S, Bartkuhn M, Renkawitz R, Renkawitz-

Pohl R. Transition from a nucleosome-based to a protamine-based chromatin

configuration during spermiogenesis in Drosophila. J Cell Sci. 2007;120: 1689–1700.

doi:10.1242/jcs.004663

39.   Brunner AM, Nanni P, Mansuy IM. Epigenetic marking of sperm by post-translational

modification of histones and protamines. Epigenetics and Chromatin. 2014;7.

doi:10.1186/1756-8935-7-2

40.   Luense LJ, Donahue G, Lin-Shiao E, Rangel R, Weller AH, Bartolomei MS, et al. Gcn5-

Mediated Histone Acetylation Governs Nucleosome Dynamics in Spermiogenesis. Dev

Cell. 2019;51: 745-758.e6. doi:10.1016/j.devcel.2019.10.024

41.   Goudarzi A, Shiota H, Rousseaux S, Khochbin S. Genome-scale acetylation-dependent

histone eviction during spermatogenesis. J Mol Biol. 2014;426: 3342–3349.

doi:10.1016/j.jmb.2014.02.023

42.   Fenic I, Sonnack V, Failing K, Bergmann M, Steger K. In vivo effects of histone-

deacetylase inhibitor trichostatin-A on murine spermatogenesis. J Androl. 2004;25: 811–

818. doi:10.1002/j.1939-4640.2004.tb02859.x

43.   Sonnack V, Failing K, Bergmann M, Steger K. Expression of hyperacetylated histone H4

during normal and impaired human spermatogenesis. Andrologia. 2002;34: 384–390.

doi:10.1046/j.1439-0272.2002.00524.x

44.   Lolis D, Georgiou I, Syrrou M, Zikopoulos K, Konstantelli M, Messinis I. Chromomycin

A3-staining as an indicator of protamine deficiency and fertilization. Int J Androl.

1996;19: 23–27. doi:10.1111/j.1365-2605.1996.tb00429.x

45.   Kazerooni T, Asadi N, Jadid L, Kazerooni M, Ghanadi A, Ghaffarpasand F, et al.

Evaluation of sperm's chromatin quality with acridine orange test, chromomycin A3 and

aniline blue staining in couples with unexplained recurrent abortion. J Assist Reprod

Genet. 2009;26: 591–596. doi:10.1007/s10815-009-9361-3

46.   Kosugi S, Hasebe M, Tomita M, Yanagawa H. Systematic identification of cell cycle-

dependent yeast nucleocytoplasmic shuttling proteins by prediction of composite motifs.

Proc Natl Acad Sci U S A. 2009;106: 10171–10176. doi:10.1073/pnas.0900604106

47.   Lindsey ARI, Rice DW, Bordenstein SR, Brooks AW, Bordenstein SR, Newton ILG.

628   Evolutionary Genetics of Cytoplasmic Incompatibility Genes cifA and cifB in Prophage

629   WO of Wolbachia. Genome Biol Evol. 2018;10: 434–451. doi:10.1093/gbe/evy012

630   48.   Shropshire JD, Kalra M, Bordenstein SR. Evolution-guided mutagenesis of the

631   cytoplasmic incompatibility proteins: Identifying CifA's complex functional repertoire

632   and new essential regions in CifB. PLoS Pathog. 2020;16: e1008794.

633   doi:10.1371/journal.ppat.1008794

634   49.   Kosugi S, Hasebe M, Matsumura N, Takashima H, Miyamoto-Sato E, Tomita M, et al.

635   Six classes of nuclear localization signals specific to different binding grooves of

636   importinα. J Biol Chem. 2009;284: 478–485. doi:10.1074/jbc.M807017200

637   50.   Ferree PM, Frydman HM, Li JM, Cao J, Wieschaus E, Sullivan W. Wolbachia utilizes

638   host microtubules and Dynein for anterior localization in the Drosophila oocyte. PLoS

639   Pathog. 2005;1: e14. doi:10.1371/journal.ppat.0010014

640   51.   Marlow FL. Maternal Control of Development in Vertebrates. Colloq Ser Dev Biol.

641   2010;1: 1–196. doi:10.4199/c00023ed1v01y201012deb005

642   52.   Beckmann JF, Sharma GD, Mendez L, Chen H, Hochstrasser M. The wolbachia

643   cytoplasmic incompatibility enzyme CIDB targets nuclear import and protamine-histone

644   exchange factors. Elife. 2019;8. doi:10.7554/eLife.50026

645   53.   Braun RE. Packaging paternal chromosomes with protamine. Nat Genet. 2001;28: 10–12.

646   doi:10.1038/ng0501-10

647   54.   Sassone-Corsi P. Editorial: Never enough - On the multiplicity and uniqueness of

648   transcriptional regulators in postmeiotic male germ cells. Endocrinology. 2002;143: 1575–

649   1577. doi:10.1210/endo.143.5.8874

650   55.   Gaucher J, Reynoird N, Montellier E, Boussouar F, Rousseaux S, Khochbin S. From

651   meiosis to postmeiotic events: The secrets of histone disappearance. FEBS J. 2010;277:

652   599–604. doi:10.1111/j.1742-4658.2009.07504.x

653   56.   Rathke C, Baarends WM, Awe S, Renkawitz-Pohl R. Chromatin dynamics during

654   spermiogenesis. Biochim Biophys Acta - Gene Regul Mech. 2014;1839: 155–168.

655   doi:10.1016/j.bbagrm.2013.08.004

656   57.   Schagdarsurengin U, Paradowska A, Steger K. Analysing the sperm epigenome: Roles in

657   early embryogenesis and assisted reproduction. Nat Rev Urol. 2012;9: 609–619.

658   doi:10.1038/nrurol.2012.183

659   58.   Carrell DT, Hammoud SS. The human sperm epigenome and its potential role in

660   embryonic development. Mol Hum Reprod. 2009;16: 37–47. doi:10.1093/molehr/gap090

661   59.   Kaneshiro KR, Rechtsteiner A, Strome S. Sperm-inherited H3K27me3 impacts offspring

662   transcription and development in C. elegans. Nat Commun. 2019;10. doi:10.1038/s41467-

663   019-09141-w

664   60.   Levine MT, Vander Wende HM, Malik HS. Mitotic fidelity requires transgenerational

665   action of a testis-restricted HP1. Elife. 2015;4: 1–20. doi:10.7554/eLife.07378

666   61.   Hundertmark T, Gärtner SMK, Rathke C, Renkawitz-Pohl R. Nejire/dCBP-mediated

667   histone H3 acetylation during spermatogenesis is essential for male fertility in Drosophila

668   melanogaster. PLoS One. 2018;13. doi:10.1371/journal.pone.0203622

669   62.   Aoki VW, Emery BR, Liu L, Carrell DT. Protamine levels vary between individual sperm

670   cells of infertile human males and correlate with viability and DNA integrity. J Androl.

671   2006;27: 890–898. doi:10.2164/jandrol.106.000703

63.   Cho C, Jung-Ha H, Willis WD, Goulding EH, Stein P, Xu Z, et al. Protamine 2 deficiency

leads to sperm DNA damage and embryo death in mice. Biol Reprod. 2003;69: 211–217.

doi:10.1095/biolreprod.102.015115

64.   Loppin B, Lepetit D, Dorus S, Couble P, Karr TL. Origin and neofunctionalization of a

Drosophila paternal effect gene essential for zygote viability. Curr Biol. 2005;15: 87–93.

doi:10.1016/j.cub.2004.12.071

65.   Clark ME, Heath BD, Anderson CL, Karr TL. Induced paternal effects mimic cytoplasmic

incompatibility in Drosophila. Genetics. 2006;173: 727–734.

doi:10.1534/genetics.105.052431

66.   Karr TL. Intracellular sperm/egg interactions in Drosophila: A three-dimensional

structural analysis of a paternal product in the developing egg. Mech Dev. 1991;34: 101–

111. doi:10.1016/0925-4773(91)90047-A

67.   Evans JP. The molecular basis of sperm-oocyte membrane interactions during mammalian

fertilization. Hum Reprod Update. 2002;8: 297–311. doi:10.1093/humupd/8.4.297

68.   Flesch FM, Gadella BM. Dynamics of the mammalian sperm plasma membrane in the

process of fertilization. Biochim Biophys Acta - Rev Biomembr. 2000;1469: 197–235.

doi:10.1016/S0304-4157(00)00018-6

69.   Degrugillier ME, Leopold RA. Ultrastructure of sperm penetration of house fly eggs. J

Ultrasructure Res. 1976;56: 312–325. doi:10.1016/S0022-5320(76)90006-X

70.   Xiao Y, Chen H, Wang H, Zhang M, Chen X, Berk JM, et al. Structural and mechanistic

insights into the complexes formed by Wolbachia cytoplasmic incompatibility factors .

Proc Natl Acad Sci. 2021;118: e2107699118. doi:10.1073/pnas.2107699118

71.   McLay DW, Clarke HJ. Remodelling the paternal chromatin at fertilization in mammals.

Reproduction. 2003;125: 625–633. doi:10.1530/rep.0.1250625

72. Landmann F, Orsi GA, Loppin B, Sullivan W. Wolbachia-mediated cytoplasmic

incompatibility is associated with impaired histone deposition in the male pronucleus.

PLoS Pathog. 2009;5. doi:10.1371/journal.ppat.1000343

73. Zhang B, Ye W, Ye Y, Zhou H, Saeed AFUH, Chen J, et al. Structural insights into

Cas13b-guided CRISPR RNA maturation and recognition. Cell Res. 2018;28: 1198–1201.

doi:10.1038/s41422-018-0109-4

74. Tirmarche S, Kimura S, Sapey-Triomphe L, Sullivan W, Landmann F, Loppin B.

Drosophila protamine-like Mst35ba and Mst35bb are required for proper sperm nuclear

morphology but are dispensable for male fertility. G3 Genes, Genomes, Genet. 2014;4:

2241–2245. doi:10.1534/g3.114.012724

75. Casiraghi M, Bordenstein SR, Baldo L, Lo N, Beninati T, Wernegreen JJ, et al. Phylogeny

of Wolbachia pipientis based on gltA , groEL and ftsZ gene sequences□: clustering of

arthropod and nematode symbionts in the F supergroup , and evidence for further diversity

in the Wolbachia tree. 2006; 4015–4022. doi:10.1099/mic.0.28313-0

76. Layton EM, On J, Perlmutter JI, Bordenstein SR, Shropshire JD. Paternal grandmother

age affects the strength of Wolbachia-induced cytoplasmic incompatibility in Drosophila

melanogaster. MBio. 2019;10. doi:10.1128/mBio.01879-19

77. Shropshire JD, Hamant E, Cooper BS. Male Age and Wolbachia Dynamics: Investigating

How Fast and Why Bacterial Densities and Cytoplasmic Incompatibility Strengths Vary.

MBio. 2021;12. doi:10.1128/mbio.02998-21

78. Kaur R, Martinez J, Rota-Stabelli O, Jiggins FM, Miller WJ. Age, tissue, genotype and

virus infection regulate Wolbachia levels in Drosophila. Mol Ecol. 2020;29: 2063–2079.

doi:10.1111/mec.15462

79.   Newton ILG, Savytskyy O, Sheehan KB. Wolbachia Utilize Host Actin for Efficient

720   Maternal Transmission in Drosophila melanogaster. PLoS Pathog. 2015;11.

721   doi:10.1371/journal.ppat.1004798

     Figure legends

Fig 1. CifA and CifB invade sperm nuclei during spermatogenesis and spermiogenesis.
Schematic representation of *Drosophila melanogaster* male reproductive system created by
Biorender is shown on the top. Testes (n=20) from <8 hrs old males expressing dual transgenes
cifAB were dissected and immunostained to visualize CifA (green) and CifB (red) during sperm
morphogenesis. DAPI stain (blue) was used to label nuclei. CifA, but not CifB, localizes in the
germline stem cells at the apical end of testes. Both CifA and CifB localize in the nuclei of
mitotic spermatogonium, spermatocytes, and round onion stage spermatids. In the later stages of
spermiogenesis, elongating spermatids harbor CifA and CifB at the acrosomal tip of the heads.
CifB is present in all canoe-stage spermatid nuclei, whereas CifA is present on average in 39%
of spermatids per bundle. Cifs are not accessible by the antibodies in the tightly compacted
spermatids at the needle-stage. After decondensing mature sperms isolated from seminal vesicles
(see methods), CifA and CifB are detectable in the acrosome regions at varying percentages.
CifA is common among sperm tails in a speckled pattern (white arrow) and either present on
average in 45% or 0% of the mature sperm heads depending upon the sampled seminal vesicles.
CifA's presence in the acrosome region is shown by solid white arrowheads and absence with
empty white arrowheads. CifB is present in acrosomal tips of all of the sperms (solid white
arrowheads) and does not occur with sperm tails. CifA and CifB localization patterns are similar
in wild type (*w*Mel+) line and signals are absent in *Wolbachia*-uninfected (*w*Mel-) negative
control line (Fig S2). Some of the CifA and CifB proteins are also stripped by the
individualization complex into the cytoplasmic waste bag (Fig S5). The experiment was repeated
in two biological replicates.

Fig 2. Cifs cause histone retention in late canoe spermatids and protamine deficiency in
mature sperms. (A) Testes (n=15) from <8 hrs old males of *w*Mel+, *w*Mel- and transgenic *cifAB*
lines were dissected and immunostained to visualize and quantify spermatid bundles with histone
retention (purple) during late canoe stage of spermatogenesis. DAPI stain (blue) was used to
label spermatid nuclei. Total spermatid bundles with DAPI signals and those with retained
Histones were manually counted and graphed. Compared to the negative control *w*Mel-, *w*Mel+
Wolbachia and dually-expressed *cifAB* transgenic lines show abnormal histone retention in the
late canoe stage. Vertical bars represent mean, and error bars represent standard deviation.
Letters indicate statistically significant (p<0.05) differences as determined by pairwise
comparisons based on Kolmogorov-Smirnov test. (B) Mature sperms isolated from seminal
vesicles (n=15) of <8 hr old males reared at 21°C were stained with fluorescent CMA3 (green)
for detection of protamine deficiency in each individual sperm nucleus. Individual sperm head
intensity was quantified in ImageJ (see methods) and graphed. *w*Mel+ and transgenic *cifAB* lines
show enhanced protamine deficiency levels compared to *w*Mel- control. Vertical bars represent
mean and error bars represent standard deviation. Letters indicate statistically significant
(p<0.05) differences as determined by multiple comparisons based on a Kruskal-Wallis test and

Dunn's multiple test correction. All of the P-values are reported in Table S1. The experiments were performed in two independent biological replicates and samples were blind-coded for the first run. Raw data underlying this figure can be found in S1 data file.

Fig 3. Protamine mutants enhance wild type CI and show significantly increased levels of protamine deficiency in mature sperms. (A) Sperms from the *Wolbachia*-infected (ΔProt+) and -uninfected (ΔProt-) protamine mutant (w[1118]; ΔMst35B[floxed], Sco/CyO) males exhibit significantly increased CMA3 fluorescence indicative of protamine deficiency compared to both wild type *w*Mel+ and *w*Mel-. Vertical bars represent mean, and error bars represent standard deviation. Letters indicate statistically significant (p<0.05) differences as determined by multiple comparisons based on a Kruskal-Wallis test and Dunn's multiple test correction. All of the P-values are reported in Table S1. (B) CI hatch rate analyses of male siblings used in CMA3 assays (panel A) validate that ΔProt+ males with increased sperm protamine deficiency causes stronger (rescuable) CI levels than *w*Mel+. ΔProt- males do not cause CI. Letters to the right indicate statistically significant (p<0.05) differences as determined by pairwise Mann-Whitney test and multiple comparisons calculated using a Kruskal-Wallis test and Dunn's multiple test correction. All of the P-values related to CI assay are reported in Table S2. Raw data underlying this figure can be found in S1 data file.

Fig 4. A nuclear localization signal in CifA is necessary for CI, rescue, and protamine levels. (A) Schematic representation of CifA annotation shows the annotated bipartite nuclear localization signal (bNLS) with engineered amino acid substitutions and deletions. (B, C) Hatch rate assays assessed both CI (B) and rescue (C) in flies expressing wild type, transgenic, and mutant *cifA*. Each dot represents the percent of embryos that hatched from a single male and female pair. Sample size is listed in parentheses. Horizontal bars represent the median. Letters to the right indicate significant differences determined by a Kruskal-Wallis test and Dunn's multiple comparison tests. All the P-values are reported in Table S2. (D) Antibody labeling (green) and DAPI staining of onion stage spermatids in the testes of the bNLS mutant line (*cifA~ΔbNLS~*) reveals that the deletion ablates CifA's localization to the nucleus, and CifA thus remains in the surrounding cytoplasm. The imaging experiment was conducted in parallel to *nos;cifAB* line shown in Fig 1. (E) Mature sperms isolated from seminal vesicles (n=15) of <8 hr old males of transgenic *cifA~Δbnls~B* line shows reduced fluorescence indicative of less Protamine deficiency compared to *cifAB*. To control for any background confounding effects of *nos-Gal4VP16* driver line, *w*Mel- fathers were prior crossed to *nos-* mothers to generate males with *nos;wMel-* genotype. CMA3 fluorescence levels of sperms isolated from *nos;wMel-* males were similar to *w*Mel- wild type lines used in previous assays in this study. Vertical bars represent mean and error bars represent standard deviation. Letters indicate statistically significant (p<0.05) differences as determined by multiple comparisons based on a Kruskal-Wallis test and Dunn's multiple test correction. All of the P-values are reported in Table S1 and raw data underlying this figure can be found in S1 data file.

Fig 5. CifA, CifB, and the protamine deficiency are transferred with the mature sperm to the female reproductive tract. (A) Schematic representation of *Drosophila melanogaster* female reproductive system. Mature oocytes leave the ovary (OV) and reach the uterus (UT), where they can be fertilized prior to being laid. Sperms from males are stored in specialized organs - spermathecae (SP) and seminal receptacle (SR) shown in the box, which open into the UT for fertilization to occur. Schematic is created with BioRender (B) Transgenic *cifAB*-expressing and *w*Mel- males were crossed to *w*Mel- females. 4 hrs post-fertilization, sperms isolated from females were decondensed and immunostained for localizing CifA (green) and CifB (red). DAPI stain (blue) was used to label nuclei. CifA is absent in sperm heads (empty arrowheads) and puctae are seen along the sperm tails (arrows). CifB is present in apical acrosomal tip of all of the sperm heads (solid arrowheads), with more distant signal in the more decondensed sperm nuclei. No Cifs are present in the sperms transferred from *w*Mel- negative control males. (C) Individual sperm intensity quantification shows that protamine deficiency of sperms from *w*Mel+ and transgenic *cifAB* males persists after transfer in the females compared to *w*Mel- males. Sperm protamine deficiency from transgenic *cifAB* males also persists in the reproductive tract of *w*Mel+ females. Vertical bars represent mean, and error bars represent standard deviation. Letters indicate statistically significant ($p<0.05$) differences as determined by multiple comparisons based on a Kruskal-Wallis test and Dunn's multiple test correction. All of the P-values are reported in Table S1 and the raw data underlying this panel can be found in S1 data file. (D) Representative images of CMA3-stained mature sperms (arrows) transferred from *w*Mel-, *w*Mel+ and transgenic *cifAB* males in *w*Mel- and *w*Mel+ female reproductive systems are shown.

Fig 6. CifA is present in early oogenesis and absent in late-stage egg chambers. Both CifA and CifB are absent in CI and rescue embryos. (A) Schematic representation of *Drosophila melanogaster* ovariole at the top illustrates the stages of oogenesis from left to right. Image was created with BioRender. Immunostaining assay indicates localization of CifA (green) to the cyst DNA (blue labeled with DAPI) in region 1 of the germarium of *Wolbachia*-uninfected transgenic *cifA* line. In *w*Mel+ line, CifA colocalizes with *Wolbachia* (red) in the germarium, nurse cells and oocyte cytoplasm along 2-8 stages of egg chambers. CifA is absent in stage 10 egg chamber, whereas *Wolbachia* signals persist. In the transgenic *cifA* line, we note the observed autofluorescence in green channel outlining the tissue morphology does not signify CifA signals. Ovariole images were manually adjusted in Affinity designer software to align egg chamber stages in the same plane. (B) Immunofluorescence of CifA (green) and CifB (red) in ~30-60 min old embryos obtained from rescue (*cifAB* male x *w*Mel+ female) and CI crosses (*nos;cifAB* male x *w*Mel- female). Histone antibody labeling core-histones (magenta) was used as a positive control. Histone signals were detected colocalizing with host DNA, labelled with DAPI (blue), whereas no CifA and CifB signals were detected. Dotted white embryonic periphery is drawn

843   around the embryo shape. White arrows indicate dividing nuclei post fertilization and
844   arrowheads indicate polar bodies.
845
846
847   **Supplementary figure legends.**
848
Fig S1. Western blots using Cif antibodies reveal proteins at the proper size. Western blots
were run on protein extracted from ovaries (n=10) of wild type infected *w*Mel+, uninfected
*w*Mel-, *cifA* transgenic (*cifA*), and dual *cifA;B* expressing transgenic lines. The expected size for
CifA is ~54 kD. Western blots were run using anti-CifB antibody on testes (n=15) of wild-type
infected (+), uninfected (-), and *cifA;B* transgenic (A;B) lines. Expected CifB size is ~133kD.
Cifs are absent in *w*Mel- control and present at accurate size in *w*Mel+ and *cif expressing* lines.
855
Fig S2. Cifs invade spermatid nuclei in wild type *w*Mel+ testes. Testes (n=20) from <8 hrs old
males of wild type *w*Mel+ and *w*Mel- lines were dissected and immunostained to visualize CifA
(green) and CifB (red) during sperm morphogenesis. DAPI stain (blue) was used to label nuclei.
CifA and CifB localization patterns in wild type lines are similar to that of transgenic *cifAB* (Fig
1) and signals are absent in *w*Mel- uninfected control line. The experiment was conducted in
parallel to the one shown in Fig 1.
862
Fig S3. CifA and CifB vary in abundance levels in spermatids and mature sperms. (A)
ImageJ-based signal intensity quantification indicates CifA (green) is more abundantly expressed
than CifB (red) in the spermatogonium stage of the spermatogenesis. Mean of individual data
points with standard deviation is plotted on the graph. Letters indicate statistically significant
($p<0.05$) differences as determined by pairwise comparisons based on a Mann-Whitney test. P-
values are reported in Table S1. (B) In the decondensed mature sperms isolated from seminal
vesicles, CifB is present in the acrosomal tip of canoe-shaped spermatids and mature sperm
heads, whereas CifA is present in only 40% and 20% of them, respectively. Quantification was
performed on the images obtained in Fig 1 data. Each dot represents percentage of Cifs present
in spermatids or mature sperms per testes examined. Raw data underlying this figure can be
found in S1 data file.
874
Fig S4. CifA and CifB are not detectable in the condensed mature sperms in seminal
vesicles due to technical limitations. Seminal vesicles (n=20) from <8 hrs old males of
transgenic *cifAB*, wild type *w*Mel+ and *w*Mel- lines were dissected and immunostained to
visualize CifA (green) and CifB (red) in the mature condensed sperms (indicated by white
arrows). DAPI stain (blue) was used to label nuclei. Absence of both CifA and CifB indicates
that proteins are not accessible to the antibodies when the sperm chromatin is condensed and
tightly packed.
882

Fig S5. CifA and CifB are also removed in the cytoplasmic waste bag. Testes (n=20) from <8 hrs old males of transgenic *cifAB*, and wild type *w*Mel- lines were dissected and immunostained to visualize CifA (green) and CifB (red) in the cytoplasmic waste bags (WB) that are present near the basal end of sperm tail bundles. Some of the Cif proteins strip down in the WB in *cifAB* line and absent in *w*Mel- control testes. Brightfield is shown to highlight the morphology of sperm tail bundles and waste bags, which are otherwise not visible using Cif antibodies and DAPI stain. The experiment was run in parallel to the ones shown in Fig 1 and Fig S2.

Fig S6. Full uncropped fluorescent images are shown related to Fig 2A.

Fig S7. Individual CifA- and CifB-expressing lines do not show abnormal histone retention but are protamine deficient. (A) Testes (n=15) from <8 hrs old males of single transgene-expressing lines *cifA*, *cifB* and a non CI-causing control gene *WD0508* were dissected to quantify spermatid bundles with histone retention (purple) during late canoe stage of spermatogenesis. DAPI stain (blue) was used to label spermatid nuclei. Total spermatid bundles with DAPI signals and those with retained Histones were manually counted and graphed. Single transgenic expressing lines showed significantly less histones similar to the negative control *w*Mel- at the late canoe stage. Vertical bars represent mean and error bars represent standard deviation. Letters indicate statistically significant ($p<0.05$) differences as determined by multiple comparisons based on a Kruskal-Wallis test and Dunn's multiple test correction. (B) Mature sperms isolated from seminal vesicles (n=15) of <8 hr old males reared at 21°C were stained with fluorescent CMA3 (green) for detection of protamine deficiency in each individual sperm nucleus. Individual sperm head intensity was quantified in ImageJ (see methods) and graphed. *cifA*- and *cifB*-expressing lines showed significantly higher fluorescence indicative of reduced levels of protamines compared to *w*Mel- and *WD0508* control lines. Vertical bars represent mean and error bars represent standard deviation. Letters indicate statistically significant ($p<0.05$) differences as determined by multiple comparisons based on a Kruskal-Wallis test and Dunn's multiple test correction. All of the P-values are reported in Table S1. The experiments were performed in parallel to the ones shown in Fig 2. (C) CI hatch rate analyses of transgenic male siblings used in CMA3 assays (Figure 2B, S6) validate that CI crosses (black circles) yielded significant less embryonic hatching compared to non CI-inducing ones, when reared at 21^0C. Letters to the right indicate statistically significant ($p<0.05$) differences as determined by multiple comparisons based on a Kruskal-Wallis test and Dunn's multiple test correction. All of the P-values are reported in Table S2. Raw data underlying this figure can be found in S1 data file.

Fig S8. 7 day-old males do not cause CI and are not protamine deficient. (A) Sperms from the 7 day-old wild type *Wolbachia*-infected (*w*Mel+) males show similar level of protamine levels as of *w*Mel-. Vertical bars represent mean, and error bars represent standard deviation. Letters indicate statistically significant ($p<0.05$) differences as determined by multiple

comparisons based on a Kruskal-Wallis test and Dunn's multiple test correction. All of the P-
values are reported in Table S1. (B) CI hatch rate analyses of male siblings used in CMA3 assays
(panel A) validate that 7d old *w*Mel+ do not induce CI that correlates with their normal levels of
sperm protamine levels. Letters to the right indicate statistically significant ($p<0.05$) differences
as determined by pairwise Mann-Whitney test and multiple comparisons calculated using a
Kruskal-Wallis test and Dunn's multiple test correction. All of the P-values are reported in Table
S2. Raw data underlying this figure can be found in S1 data file.
930
Fig S9. CifA is absent in late oocyte stages and the developing embryos from the rescue
cross. (A) In the transgenic *cifA* line, CifA (green) is absent in the late oocyte stages. Image was
acquired at 20x magnification to show mid and late oocytes in one plane. We note the
autofluorescence upon enhanced exposure in the green channel outlining the tissue morphology
of stage 15 egg chamber does not signify CifA signals. (B) Immunofluorescence of CifA (green)
and histones (magenta) in 1-2 hr old embryos (n=50) obtained from rescue cross (*cifAB* male x
*w*Mel+ female). Histone signals are detected in the developing embryos colocalizing with host
DNA, labelled with DAPI (blue), whereas CifA signals are absent.
939

	Apical tip	Spermatogonium	Spermatocytes	Round Onion spermatids	Canoe spermatids	Needle spermatids	Decondensed mature sperms

α-CifA DAPI

α-CifB DAPI

A

CifA | bNLS | | | DUF | | STE | |

$cifA_{189}$ $cifA_{224}$

F189A F224A
Y190A C225A

$cifA_{\Delta bNLS}$

B

$nos;cifAB$ ♂ x ♀   (30)   a

$nos;cifA_{\Delta bNLS};B$ ♂ x ♀   (31)   b

$nos;cifA_{189;224};B$ ♂ x ♀   (19)   c   CI inhibited

$nos;cifA_{189};B$ ♂ x ♀   (29)   b

$nos;cifA_{224};B$ ♂ x ♀   (12)   a

0 10 20 30 40 50 60 70 80 90 100
Embryos hatched (%)

C

$nos;cifAB$ ♂ x $nos;cifA$ ♀   (27)   a

$nos;cifAB$ ♂ x $nos;cifA_{\Delta bNLS}$ ♀   (28)   b

$nos;cifAB$ ♂ x $nos;cifA_{189;224}$ ♀   (30)   b   Rescue inhibited

$nos;cifAB$ ♂ x $nos;cifA_{189}$ ♀   (28)   c

$nos;cifAB$ ♂ x $nos;cifA_{224}$ ♀   (30)   a

0 10 20 30 40 50 60 70 80 90 100
Embryos hatched (%)

D

α-CifA   Merged with DAPI

$nos;cifA_{\Delta bNLS}B$
Round onion-stage spermatids

E

Protamine-deficient CMA3 fluorescence per sperm

2.5×10⁶
2.0×10⁶
1.5×10⁶
1.0×10⁶
5.0×10⁵
0.0

a   b   c

$nos;wMel-$ $nos;cifA_{\Delta bNLS}B$ $nos;cifAB$

A

OV SP SR UT

B

nos;cifAB ♂ x *wMel-* ♀ | *wMel-* ♂ x *wMel-* ♀

α-CifA DAPI

α-CifB DAPI

C

Protamine-deficient CMA3 fluorescence / sperm in female uterus

10^6

a   bc   b   c

10^5

10^4

10^3

♂ *w*Mel-
x
♀ *w*Mel-

*w*Mel+
x
*w*Mel-

nos;cifAB
x
*w*Mel-

nos;cifAB
x
*w*Mel+

D

wMel- ♂ x *wMel-* ♀ | *wMel+* ♂ x *wMel-* ♀

nos;cifAB ♂ x *wMel-* ♀ | *nos;cifAB* ♂ x *wMel+* ♀

A

Germarium   Egg chamber stages during oogenesis
                    2-8   10

Region 1   Nurse cells Oocyte   Mature egg

wMel+

α-CifA
α-*Wolb*
DAPI

nos;cifA

Region 1

α-CifA
α-*Wolb*
DAPI

B

α-CifA   DAPI   Merge

Rescue embryos

α-His

α-CifB

CI embryo