De novo analysis of bulk RNA-seq data at spatially resolved single-cell resolution

Jie Liao[1, 2, †], Jingyang Qian[1, †], Yin Fang[3, 4, †], Zhuo Chen[3, 4, †], Xiang Zhuang[3, 4, †], Ningyu Zhang[3, 4], Xin Shao[1, 2], Yining Hu[1], Penghui Yang[1], Junyun Cheng[1, 5], Yang Hu[1, 5], Lingqi Yu[1], Haihong Yang[3, 4], Jinlu Zhang[1, 3], Xiaoyan Lu[1, 5], Li Shao[6], Dan Wu[7], Yue Gao[8], Huajun Chen[3, 4] & Xiaohui Fan[1, 2, 5, 9]

[1] College of Pharmaceutical Sciences, Zhejiang University, Hangzhou 310058, China

[2] Future Health Laboratory, Innovation Center of Yangtze River Delta, Zhejiang University, Jiaxing 314100, China

[3] College of Computer Science and Technology, Zhejiang University, Hangzhou 310027, China

[4] Hangzhou Innovation Center, Zhejiang University, Hangzhou 310058, China

[5] Innovation Center in Zhejiang University, State Key Laboratory of Component-Based Chinese Medicine, Hangzhou 310058, China

[6] Institute of Translational Medicine, The Affiliated Hospital of Hangzhou Normal University, Hangzhou Normal University, Hangzhou 310015, China

[7] College of Biomedical Engineering and Instrument Science, Zhejiang University, Hangzhou 310013, China

[8] Department of Pharmaceutical Sciences, Beijing Institute of Radiation Medicine, Beijing, 100850, China

[9] Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, 310024, China.

Correspondence should be addressed to X.F. (e-mail: fanxh@zju.edu.cn), H.C. (e-mail: huajunsir@zju.edu.cn) and Y.G. (e-mail: gaoyue@bmi.ac.cn).

[†] These authors contributed equally to this work.

Abstract

Uncovering the tissue molecular architecture at single-cell resolution could help better understand organisms' biological and pathological processes. However, bulk RNA-seq can only measure gene expression in cell mixtures, without revealing the transcriptional heterogeneity and spatial patterns of single cells. Herein, we introduce Bulk2Space (https://github.com/ZJUFanLab/bulk2space), a deep learning framework-based spatial deconvolution algorithm that can simultaneously disclose the spatial and cellular heterogeneity of bulk RNA-seq data using existing single-cell and spatial transcriptomics references. The use of bulk transcriptomics to validate Bulk2Space unveils, in particular, the spatial variance of immune cells in different tumor regions, the molecular and spatial heterogeneity of tissues during inflammation-induced tumorigenesis, and spatial patterns of novel genes in different cell types. Moreover, Bulk2Space is utilized to perform spatial deconvolution analysis on bulk transcriptome data from two different mouse brain regions derived from our in-house developed sequencing approach termed Spatial-seq. We have not only reconstructed the hierarchical structure of the mouse isocortex but also further annotated cell types that were not identified by original methods in the mouse hypothalamus.

Introduction

Tissue complexity is portrayed by the spatial diversity and heterogeneity of cells[1]. Advances in spatially resolved transcriptomics[2-4] have made it possible to understand the cell composition, molecular architecture, and functional details of tissues at unanticipated spatial levels[5, 6]. State-of-the-art experimental technologies, including image-based methods[7-10], spatial barcoding RNA-seq methods[11-14], and laser capture microdissection-based methods[15, 16], have been developed to address either high throughput measuring of cells, unbiased detection of mRNA species, or single-cell resolution. To investigate the molecular variation during biological and pathological processes at a higher resolution, each sample is encouraged to be analyzed by spatially resolved single-cell transcriptomics, which is not yet fully achieved and is time-consuming, costly, and difficult to scale up[17]. Meanwhile, with investment for almost

two decades, RNA-seq has been extensively applied in transcriptome analysis[18], with

many large projects having been carried out, such as the Encyclopedia of DNA

Elements (ENCODE)[19], The Cancer Genome Atlas (TCGA)[20], and projects of the

International Cancer Genome Consortium (ICGC)[21]. A wealth of bulk RNA-seq data has

become a legacy for biological and clinical research[22]. Thus, reanalysis of the enormous

amount of bulk data to explain both cellular diversity and spatial expression patterns

is a challenging but consequential task.

In silico methods have great potential to predict spatial heterogeneity from bulk

RNA-seq data at single-cell resolution by integrating cutting-edge technologies. Several

approaches such as CPM[23], CIBERSORT[24], and MuSiC[25] can only extrapolate

proportions of cell types from bulk RNA-seq data and have failed to further generate

single-cell data, let alone map them to tissue coordinates. The emergence of an

approach that can efficiently decompose bulk RNA-seq data into spatially resolved

single-cell expression profiles is expected to reveal the cell diversity of complex tissue

and the spatial expression variation simultaneously.

Herein, we introduce Bulk2Space, a spatial deconvolution algorithm based on deep

learning frameworks, which generates spatially resolved single-cell expression profiles

from bulk transcriptomes using existing high-quality scRNA-seq data and spatial

transcriptomics as references. We hypothesize that the process of scRNA-seq is similar

to sampling cells from bulk tissue, and each selected cell is labeled with a unique

barcode. Consequently, bulk transcriptome data can be used as a weighted collection

of single-cell expression data in a defined clustering space of cells. Bulk2Space first

generates single-cell transcriptomic data within the clustering space to find a set of

cells whose aggregated data is proximate to the bulk data. Next, the generated single

cells were allocated to optimal spatial locations using a spatial transcriptome reference.

For this step, we chose as a spatial reference one of the two most commonly used

spatially resolved transcriptomics technologies. One is an image-based method with

limited target genes[7-9], and the other is a spatial barcoding method without single-cell

resolution[11-14]. Taken all, Bulk2Space showed a robust performance across multiple

datasets and conditions and is an open-access algorithm on GitHub

(https://github.com/ZJUFanLab/bulk2space).

Results

Design concept of Bulk2Space

The overall design of the Bulk2Space algorithm is illustrated in Fig. 1, which is divided
into two steps, deconvolution, and spatial mapping. Although bulk RNA-seq data are
obtained by sequencing a mixture of cells, whereas scRNA-seq labels individual cells
in advance, both methods share comparable cell types and states[23-26]. Thus, we
assume that bulk transcriptomics data can be decomposed into single-cell
transcriptomics data by a well-designed deconvolution algorithm (Step 1).
Subsequently, in the second step, the above-mentioned two prevailing spatially
resolved transcriptomics methods, though, failed in either achieving single-cell
resolution or delineating the whole transcriptome, can provide reference locations for
single cells generated in the first step of Bulk2Space based on the similarity of their
expression profiles (Step 2)[27, 28]. As a consequence, it could be rationally conceived
that the combination of single-cell profiles and spatial transcriptomics is expected to
overcome technical bottlenecks and accomplish the spatially resolved single-cell
deconvolution of bulk transcriptomes (Fig. 1a).

For the deconvolution of bulk transcriptomics data, we hypothesize that the process
of single-cell sequencing identifies with the process of sampling from the tissue, while
the ratio between different cell types is related to the zone of the tissue, the ways of
single-cell capturing, and the size of the cell itself. However, even if the ratio between
cell types changes, the state of each cell type still fluctuates within a relatively stable
high-dimensional space, which is namely the clustering space of cell types. This
hypothesis has been supported by several studies[29-31]. As shown in Fig. 1b, a single-
cell reference can be utilized to characterize the clustering space and average the gene
expression for each cell type[31, 32]. The expression vector of the bulk transcriptome is
taken as the input, which is equal to a product of the average gene expression matrix
of cell types and their abundance vector. By solving this nonlinear equation, the
proportion of each cell type is determined. Then, the solved proportion of each cell
type is taken as a control parameter to generate the corresponding number of single
cells. Specifically, a deep generative model, termed beta variational autoencoder (β-
VAE)[33], is employed to simulate a given number of single cells within the characterized
clustering space of each cell type. The simulation stops when the training loss no

longer reduces, thus deconvolves the heterogeneous bulk transcriptome into single-
cell transcriptome data. Instead of only calculating the abundance values of cell types,
as performed by other methods, such as Cell Population Mapping (CPM)[23],
CIBERSORT[24], and ImmuCC[26], Bulk2Space can generate biologically feasible single-cell
expression profiles.

For spatial mapping, the single-cell profiles generated in the first step are assigned
to the reference tissue section based on an optimization strategy. In the second step,
we consider two of the most commonly used spatially resolved transcriptomics
approaches as the spatial reference, namely, spatial barcoding-based RNA-seq
methods and image-based targeted methods.

Spatial barcoding-based methods employ arrayed reverse transcription primers
with unique positional barcodes to preserve the spatial information of mRNA via in
situ complementary DNA (cDNA) synthesis. Such methods could provide the whole
transcriptome as well as the location of each sequenced spot after decoding the spatial
barcodes. However, these methods are unable to achieve single-cell resolution,
because the spots are customized in shape and size, such as ST[14] (100 μm in diameter),
Visium (55 μm in diameter), Slide-seq[13] (10 μm in diameter), etc. As shown in Fig. 1c,
for spatial barcoding-based references, each barcoded spot is regarded as a mixture of
several cells. Similar to existing decomposition methods such as RCTD[34], SpatialDWLS[35],
stereoscope[36], and SPOTlight[37], we first calculate the cell-type composition of each
spot. The difference is that we then map the generated single cells into the
recommended spots based on the similarity of their cell expression profiles and ensure
that the proportions of cell types in each spot are consistent with the calculated results.
Since Bulk2Space generates spatially resolved single-cell transcriptomics data from
bulk RNA-seq, the spatial heterogeneity of individual cells can be analyzed, which
cannot be realized by RCTD, SpatialDWLS, stereoscope, and SPOTlight.

As illustrated in Fig. 1d, another strategy is to map the single-cell data generated in
the first step onto image-based targeted references capable of measuring hundreds to
thousands of RNA species[38], such as MERFISH[7], STARmap[9], seqFISH[10], etc. The pairwise
similarity between cells is calculated based on genes shared by both datasets, and
eventually, each generated single cell is robustly mapped to an optimized coordinate
of the target cell in the spatial reference. The resulting spatially resolved single-cell
RNA-seq data can provide unbiased transcriptomes of individual cells and improve the

gene coverage.

Taken all, unlike other computational methods that focus only on either of the two
spatially resolved transcriptomics approaches, Bulk2Space takes into account both
techniques to greatly increase the scalability, applicability, and reference data of the
algorithm.

Performance evaluation of Bulk2Space using simulated and biological

datasets

To demonstrate the robustness of the deconvolution step of Bulk2Space, a benchmark
test was performed using 30 paired simulations (both bulk and single-cell data were
generated from the same datasets) from 10 different high-quality single-cell datasets
across human blood, brain, kidney, liver, and lung, and mice brain, kidney, lung,
pancreas, and testis (Fig. S1a), and 12 unpaired simulations (both bulk and single-cell
data were generated from different datasets of mice pancreas) from 8 single-cell RNA-
seq data of human pancreas (Fig. S1b). All datasets used in this study were listed in
Supplementary Data 1.

First, for 30 paired simulations, each single-cell transcriptome dataset was divided
into two parts, one for reference and the other for the synthesis of bulk data. The
procedure for bulk data synthesis is shown in Fig. S1c. For each single-cell dataset, we
changed the proportion of cell types and synthesized 3 corresponding bulk
transcriptomics data with different cell compositions. Three generative deep learning
models, namely, the beta variational autoencoder (β-VAE)[33], generative adversarial
networks (GAN)[39], and conditional GAN (CGAN)[40], were introduced here for the
benchmarking of different methods using 30 paired single-cell datasets. The gene
expression correlation between the generated single-cell data and input bulk data was
calculated to evaluate and compare the three candidate algorithms. As illustrated in
Fig. 2a, β-VAE performed better than the other algorithms in both Pearson and
Spearman correlations with a lower root mean squared error (RMSE). The Pearson
correlation, Spearman correlation, and RMSE were calculated by combining the gene
expression across all cell types and averaging these metrics for each cell type. Ten
examples of different species and tissues are shown in Fig. S2. The results

demonstrated that the clustering space of the generated single-cell data was similar to that of the test data but varied in the proportions of different cell types (Fig. S2a and Fig. S2b). Furthermore, the correlation analysis of the expression of marker genes for each cell type between the generated and test data was performed for each method. The comparison of the correlation heatmaps between Bulk2Space, GAN, and CGAN showed that Bulk2Space had higher correlations across different simulations (Fig. S2c).

We then benchmarked the deconvolution step of Bulk2Space using 12 unpaired simulation data of mice pancreas. Because other deconvolution methods, such as CPM[23], CIBERSORT[24], and ImmuCC[26], can only predict cell-type proportions instead of gene expression of generated data, we compared Bulk2Space with GAN, CGAN, and a Bayesian deconvolution method termed bMIND[41]. As illustrated in Fig. 2b and Fig. S3a, Bulk2Space outperformed the other three methods with a higher Pearson correlation of gene expression and lower gene expression variation (RMSE). Although CGAN had a comparable performance as Bulk2Space in the single-cell generation, its computing speed was significantly lower than Bulk2Space. The detailed comparison of the cell-type-specific marker gene expression correlations of generated single-cell data and the ground truth between the four candidate methods were shown in Fig. S3b.

To further investigate the robustness of the single-cell generation of Bulk2Space, we introduced two noise mechanisms to test the performance of the algorithm (Fig. S4). One mechanism changes the expression values of certain genes in randomly selected cells (Fig. S4a), and the other alters the cell type labels of selected cells (Fig. S4b). Bulk2space showed robust performance and could effectively avoid the over-fitting phenomenon (Fig. S4c and Fig. S4d). We also demonstrated that Bulk2Space can deconvolve bulk RNA-seq data with an annotation-free single-cell reference in the supplementary information Fig. S5. Since the core generation model of Bulk2Space is β-VAE, a perturbation analysis was conducted to verify the robustness of single-cell generation by Bulk2Space. The perturbation test confirmed that better generation results can be obtained by assuming that the latent vectors follow the Gaussian distribution (Fig. S6). Finally, we validated Bulk2Space with paired biological bulk and single-cell data derived from different mice liver[42]. As shown in Fig. S7, the deconvolution results indicated that Bulk2Space could be well applied in biological scenarios.

In the second step, Bulk2Space assigns the generated single cells to spatial
coordinates based on a spatially resolved transcriptome reference (Fig. 1c and Fig. 1d).
There are two types of most common spatially resolved transcriptomic techniques.
One is the spatial barcoding-based RNA-seq technology, which includes 'spatial
transcriptomics' (ST)[14], high-definition spatial transcriptomics (HDST)[11], and Slide-
seq[12, 13], 10 Visium, etc. These methods are based on tissue microregions with certain
regular shapes (spots), and cannot achieve single-cell resolution. The other is image-
based in situ transcriptomics, including MERFISH[7], seqFISH[8, 10], and STARmap[9]. These
methods can only measure the expression of target genes but cannot cover the whole
transcriptome. To overcome the bottlenecks of the two approaches, we designed
corresponding spatial assignment strategies.

The first strategy was designed to map single cells to spots in spatially resolved
transcriptomics based on spatial barcoding. In this step, we first used paired synthetic
data to test the performance of Bulk2Space. The data synthesis procedure is illustrated
in Fig. S1c. Similar to the commonly used spatially resolved transcriptomics
technologies, such as ST, HDST, 10X Visium, and Slide-seq, we randomly chose 10 cells
from each scRNA-seq data and aggregated their gene expression profiles as a spot of
pseudo spatial transcriptomics data. A comprehensive comparison between
Bulk2Space (deep forest), six machine learning or deep learning approaches, and four
published methods, RCTD[34], SpatialDWLS[35], stereoscope[36], and SPOTlight[37] was then
conducted using 50 paired simulation datasets. These methods were benchmarked for
the optimization of cell-type composition and spatial gene expression patterns. As
shown in Fig. 2c, compared with these methods, Bulk2Space showed higher
correlations in gene expression and lower RMSEs than other methods. The cell
composition and spatial cell-type distribution predicted by Bulk2Space were highly
correlated to the ground truth (Fig. S8). Next, we benchmarked the spatial mapping
step of Bulk2Space using unpaired simulation data of mice pancreas. As shown in Fig.
2d, Bulk2Space showed the highest correlations in gene expression and the lowest
RMSEs. The comparison of the cell-type composition for each spot between
Bulk2Space and the spatial reference was shown in Fig. S9.

The second spatial mapping strategy uses image-based targeted methods. A
scRNA-seq   data   ("GSE113576
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE113576]") and MERFISH
data (image-based reference) of the mouse hypothalamus tissue from the same

experiment[43] were used as paired datasets to test the performance of the Bulk2Space

algorithm. Another scRNA-seq data[44] derived from a separated experiment was used

as unpaired single-cell data. We wanted to find the shortest path that could shift single

cells to optimal locations. First, a nonparametric empirical Bayes network was used to

eliminate the expression differences between the single-cell data and spatially

resolved transcriptome data. As illustrated in Fig. 2e, the clustering spaces of the two

datasets were very similar, with most cell types sharing the distribution in both

datasets. The two discrete datasets could be well merged in distribution, and the gene

expression of each cell type had a correlation of over 0.8 between the two datasets

(Fig. 2f). Of the 150 target genes identified in MERFISH, a leave-one-out test was

conducted for the evaluation of the spatial mapping step of Bulk2Space. Specifically,

149 were used as reference genes, and the rest one was used for validation. The results

showed that spatially, the gene expression pattern of the matched cells was highly

correlated with that of the reference, and the predicted spatial expression patterns

across all 150 targeted genes are strongly correlated with the ground truth with

averaged Pearson correlation coefficients (PCCs) over 0.9 for both paired and unpaired

datasets (Fig. 2g and Fig. S10).

We further verified Bulk2Space's spatial mapping using five-fold cross-validation

for image-based methods (Fig. S11). specifically, the 150 targeted genes in MERFISH

were split into five folds using R package caret (version 6.0-92), and 80% of genes were

used as the reference and the remaining was used for validation of predicted spatial

expression. As shown in Fig. S11, the spatial expressions of the top 25 genes predicted

by Bulk2Space for the paired and unpaired single-cell data were compared with

MERFISH data.

To validate the spatial mapping step of Bulk2Space in the real situation, two

biological datasets, a spatial reference sequenced by Slide-seq v2[12], and a single-cell

data[45] from separated experiments were used to reconstruct the structure of the

mouse hippocampus (Fig. S12). The spatial distribution of single cells predicted by

Bulk2Space was consistent with the real pattern of cell types in that region. Notably,

the refined subregions in the mouse hippocampus were successfully reconstructed by

Bulk2Space, which was confirmed by the spatial expression of cell-type-specific

marker genes (Fig. S12e).

Since Bulk2Space employed β-VAE to generate single-cell profiles within the

clustering space of cell types, we then investigated whether the randomness of β-VAE

could affect the generation of single-cell data and the spatial mapping results. We

conducted 100 repetitions on the deconvolution step of Bulk2Space to evaluate the

robustness of β-VAE using simulated bulk and single-cell reference[46]. The results

demonstrated that the single-cell generation remained highly robust across 100

repetitions (Fig. S13).

We then used simulated PDAC[47] and melanoma[48, 49] bulk data to evaluate the

robustness of the spatial mapping step of Bulk2Space using three repetitions of bulk

data deconvolution and spatial mapping. As shown in Fig. S14 and Fig. S15, the spatial

deconvolution results suggested a robust performance of Bulk2Space with 3

repetitions for PDAC and melanoma bulk data, respectively.

In conclusion, although the single-cell data generated by β-VAE were slightly

different each time, the overall prediction results showed robust performance in the

spatial distribution of cell types, the cell-type composition and proportion in spots,

and the spatial patterns of gene expression.

Validation of Bulk2Space using biological data

To further verify the performance of the Bulk2Space, two consecutive slices[49] from the same melanoma tissue were used to demonstrate the spatial deconvolution result. As shown in Fig. S16, one slice was used as the spatial reference and the other was used to synthesize the bulk data. The expression of cell-type-specific marker genes between the generated data and the reference were highly correlated and the spatial distribution pattern of generated single cells was consistent with the histological annotations of different regions. Moreover, we found a spatial heterogeneity of generated B cells from different tissue regions, which was associated with the biological functions in the lymph node area and the melanoma region. Because Bulk2Space can predict spatially resolved single-cell transcriptomics data from bulk RNA-seq or scRNA-seq, spatial heterogeneity of the same cell type can be discovered. This was difficult for other spatial deconvolution algorithms to achieve such as RCTD[34], SpatialDWLS[35], stereoscope[36], and SPOTlight[37].

Similarly, the same spatial deconvolution procedure of Bulk2Space was performed for another two discrete slices from different PDAC[47] tissues to reveal the spatial heterogeneity of bulk transcriptomics data. One slice was used as the spatial reference and the other was used to synthesize the bulk data. As shown in Fig. S17, similar results were retrieved as the validation of melanoma data. The expression of cell-type-specific marker genes between the generated data and the reference were highly correlated and the spatial distribution pattern of generated single cells was consistent with the histological feature of tissue regions.

Bulk2Space integrates spatial gene expression and histomorphology in PDAC

Based on the success in the spatial deconvolution of PDAC using simulated data and discrete slices, we further verified the performance of the Bulk2Space algorithm using biological bulk RNA-seq data. As shown in Fig. 3a, two bulk RNA-seq data[50] of the pancreatic adjacent (PA) and the PDAC tissues, one scRNA-seq data[47], and one spatial transcriptomics data[47] were derived from two individual experiments and three different technologies. The scRNA-seq data sequenced by inDrop-seq[51] were used as the single-cell reference for the deconvolution of PA tissue and PDAC bulk data, and the spatial barcoding-based ST data were used as the spatial reference.

The PA and PDAC bulk data were deconvolved into single-cell data with different cell-type proportions by Bulk2Space. We compared the cell-type proportions of single cells generated from PA and PDAC bulk data with Bulk2Space. As shown in Fig. 3b, the proportion of acinar cells in PA was significantly higher than that in PDAC, where the proportion of two cancer clone subtypes was higher. The clustering space of single cells generated from PDAC and PA tissue was exhibited in Fig. 3c. The detailed information for the deconvolution of PA tissue was shown in Fig. S19.

Here, we focused on the PDAC bulk data. The expression of cell-type-specific marker genes between generated single-cell profiles by Bulk2Sapce and the reference data was highly correlated (Fig. 3d). Then, each cell was mapped to coordinates based on the spatial reference (Fig. 3e). To further correlate the spatial expression pattern of the tissue with its histological feature, we investigated the molecular architecture of the annotated regions in the spatial reference (Fig. 3f). As shown, the spatial distribution of two cancer clones, acinar cells, and ductal cells predicted by Bulk2Space (Fig. 3e) was consistent with the histological annotation of cancer cells and desmoplasia, normal pancreatic tissues, and duct epithelium, as well as the spatially resolved transcriptomics of the cancer region, pancreatic tissue and stroma, and the duct epithelium, respectively. The spatial cell-type proportions predicted by Bulk2Space were illustrated in Fig. 3g and Fig. S19c. The spatial expression of cell-type-specific marker genes at spot (left) and single-cell (right) resolution for different cell types were shown in Fig. 3h and Fig. S19d. Compared with the spatial distribution of single cells generated by Bulk2Space, the spatial expression of cell-type-specific marker genes

353   exhibited consistent patterns.

Bulk2Space reconstructs the hierarchical structure of the mouse

isocortex region sequenced by Spatial-seq

In addition to linking the histomorphology and transcriptomics in pathological tissues,
another application scenario of Bulk2Space is to reconstruct the structure of tissues
with spatial patterns. For instance, the spatial organization of the isocortex region of
the mouse brain exhibits a layered pattern. Therefore, Bulk2Space was applied to
reconstruct the hierarchical structure of the mouse isocortex region through spatial
deconvolution of the bulk transcriptomics data. The bulk transcriptomics data of
mouse isocortex were sequenced by our in-house developed multiplexed RNA-seq
approach, termed Spatial-seq. The detailed information for Spatial-seq was described
in the 'Methods' section and illustrated in Fig. S20a. In short, the laser capture
microdissection (LCM) was used to isolate regions of interest from the tissue sections
and each isolated tissue was collected independently. Then, barcoded beads were
used to capture and label the mRNA from the collected samples. Finally, the captured
mRNA was pooled together for RNA-seq. The transcriptome data of each sample can
be obtained by identifying the sample-specific barcode sequence. Using Spatial-seq,
we isolated and sequenced 13 main brain regions from coronal and sagittal sections
across the entire mouse brain (Fig. S20b).

As shown in Fig. 4a, the bulk transcriptomics data of the mouse isocortex was
obtained by Spatial-seq. The mouse primary visual cortex regions in different coronal
sections from anterior to posterior were collected and sequenced by SMART-seq2[52].
The result scRNA-seq data were used as the single-cell reference. A sagittal section of
the mouse brain was divided into two parts and sequenced using 10X Visium to obtain
spatial transcriptomics data of the isocortex region. The spatial reference data were
downloaded from 10X datasets. The detailed information of datasets used in this study
was summarized in Supplementary Data 1.

The bulk data were first deconvolved into single-cell RNA-seq data (Fig. 4b).
Subsequently, the generated single cells were mapped to spatial locations by
Bulk2Space and the spatial distribution of cell types showed a distinct layered pattern
(Fig. 4c). The expression of cell-type-specific marker genes was highly correlated

between Bulk2Space results and the single-cell reference data (Fig. 4d). The spatial distribution of the cell-type proportion in each spot predicted by Bulk2Space was illustrated in Fig. 4e, which corresponded well with the hierarchical structure of the mouse isocortex. The spatial distributions of cell-type proportions predicted by Bulk2Space for seven cell types including Astro, L2/3 IT, L4, L5 IT, L5 PT, L6 CT, and L6 IT cells were shown in Fig. 4f. The results supported the ability of Bulk2Space to reconstruct the spatial organization of tissues. Moreover, the spatial expression patterns of cell-type-specific marker genes predicted by Bulk2Space at single-cell resolution were consistent with the proportion distributions of the corresponding cell types (Fig. 4g).

Different from traditional spatial deconvolution algorithms such as RCTD[34], SpatialDWLS[35], stereoscope[36], and SPOTlight[37], Bulk2Space can generate spatially resolved single-cell transcriptomics data from bulk RNA-seq. Besides, Bulk2Space allowed us to analyze the spatial heterogeneity of individual cells from the spatially resolved transcriptomics data without single-cell resolution.

Bulk2Space re-annotates ambiguous cells in the mouse hypothalamus

The Bulk2Space algorithm was also used to spatially deconvolve bulk transcriptome data derived from the hypothalamus region of the mouse brain using Spatial-seq, to explain the spatial distribution of single cells and gene expression. The resources of the bulk, single-cell, and spatial data was illustrated in Fig. 5a. The single-cell reference was obtained by profiling ~31000 cells using Drop-seq and the MERFISH data of the mouse hypothalamus were used as the spatial reference[43]. More details were summarized in Supplementary Data 1.

After deconvolution of the bulk RNA-seq data of the mouse hypothalamus by Bulk2Space, the clustering space of the generated single cells was close to that of the MERFISH data (Fig. 5b). The generated single cells were then mapped to spatial coordinates based on the image-based spatial reference (Fig. 5c). As shown in Fig. 5c, the cell-type distributions in MERFISH data and Bulk2Space results were comparable at single-cell resolution. Among the 10 cell types, immature oligodendrocytes were unique to MERFISH, and macrophages were predicted by Bulk2Space, but absent in MERFISH data. The heatmap suggested that 7 of the 8 shared cell types were strongly correlated in both MERFISH and Bulk2Space results with an average pairwise

correlation over 0.9. The poor correlation of mural cells between MERFISH data and
Bulk2Space results may be related to the few numbers of pericytes. The spatial
patterns of the MERFISH targeted genes were consistent with that of Bulk2Space
results (Fig. 5d).

Meanwhile, owing to the limited number of measured target genes, some
ambiguous cells in MERFISH data could not be further identified according to their
expression profiles. However, using Bulk2Space, the generated single cells with
unbiased gene expression were assigned to the spatial context, thus further clustering
these ambiguous cells using novel genes that were absent in MERFISH data (Fig. 5e).
Meanwhile, the spatial distribution of novel genes beyond the targeted RNA species
was predicted by Bulk2Space and was consistent with the spatial pattern of
corresponding cell types as illustrated in Fig. 5f and Fig. S21.

Bulk2Space uncovers spatial gene expression dynamics in different
stages of the inflammation-induced prostate cancer

Tumor development is linked to chronic infection, dietary factors, obesity, inhaled
pollutants, tobacco use, and autoimmunity. The unifying principle underlying these
processes is inflammation, which is an aberrantly prolonged form of a protective
response to a loss of tissue homeostasis[53]. To explore whether Bulk2Space could
identify the cellular and molecular pathways that coordinate the tumor-promoting
effects in inflammation-induced cancer, we deconvolved the bulk transcriptome data
of prostate cancer and mapped single cells generated by Bulk2Space to the
corresponding spatial references, including normal glands, inflammatory tissue, and
tumor sites. The resources of the bulk (from TCGA), single-cell[54], and spatial data[55]
were listed in Supplementary Data 1.

These tissue regions were isolated from the prostate, and three of the sections
were subjected to RNA-seq based on spatial barcoding (Fig. 6a). Next, Bulk2Space was
used to deconvolve the bulk data into single-cell profiles (Fig. 6b) and mapped the
generated single cells to spatial coordinates (Fig. 6c). Notably, the composition and
distribution of cells in different tissue regions were quite different and closely related
to the state of the tissue (Fig. 6d and 6e). A comparison of cell distribution between
normal, inflammatory, and cancerous tissues (Fig. 6f to 6h) showed that in normal

glands, cancer-associated fibroblasts (CAFs) were the most abundant, but in
inflammatory and cancer tissues, the proportions of CAFs were significantly lower
(52.4% in normal, 20.5% in inflammation, and 27.5% in cancer). In the areas of
inflammation, cells involved in the inflammatory response, such as immunocompetent
B cells, endothelial cells, monocytes, and NK cells (Fig. 6g), were more abundant than
in normal or tumor tissues (13.4% in inflammation, 2.0% in normal, and 1.7% in cancer).
In cancer with a Gleason score (Gs) of 3+4, the proportion of cancer epithelial cells was
significantly higher (53.0% in cancer, 6.3% in normal, and 4.3% in inflammation) than
that in normal and inflammatory areas (Fig. 6h), while the proportion of the
periventricular layer cells was significantly lower (17.9% in cancer, 39.3% in normal,
and 44.1% in inflammation).

During the progress of inflammation-induced prostate cancer, CAFs accumulated
in normal glands at the early stage, resulting in high expression of inflammatory factors
and promoting the occurrence of local inflammation, a fundamental innate immune
response to perturbed tissue homeostasis, thus leading to tissue cancerization[53].
Bulk2Space results were consistent with this phenomenon.

Discussion

Understanding the transcriptional heterogeneity within the tissue from the
perspective of single-cell spatial resolution and full gene coverage is an essential
direction for the development in the field of biological sciences[1, 56]. The spatial
locations of cells may determine their identity and how they interplay with each other
in the microenvironment[57, 58]. However, there are many technical challenges, and thus,
such a technology has not been fully realized yet. Traditional bulk transcriptome data
can provide gene expression patterns and reflect the overall status of tissue, which is
widely used in studies on the occurrence and development of diseases. Nevertheless,
disease-inherent cell composition and spatial distribution are difficult to analyze.
Although several attempts can infer the cell-type proportions from traditional RNA-
seq data, understanding bulk tissue at single-cell spatial resolution remains a pressing
need in the field. Cell identity can be characterized by the clustering space of single-
cell gene expression profiles[59], and expression features can be stably retained across
different conditions, technologies, and species[29]. Based on this, we used a deep

learning model, termed β-VAE, to generate single-cell profiles with biological
significance within the clustering space of each cell type and map them accurately to
tissue coordinates, thus deconvolving the bulk transcriptomics into spatially resolved
single-cell transcriptomics data.

As thoroughly considered the merits and demerits of the widespread use of two
spatially resolved transcriptomics technologies, we employed different spatial
mapping strategies for each method. For image-based in situ hybridization,
Bulk2Space could predict the spatial expression of mRNA species apart from the target
genes. Moreover, Bulk2Space can help annotate cells whose types are difficult to
further distinguish by targeted methods. For spatial barcoding-based RNA-seq, each
sequenced tissue region contains transcripts from different cells and can thus be
regarded as a mixture of cells, not allowing single-cell resolution. Bulk2Space can
assign single cells to optimal spots, and these mixtures are split into collections of
individual cells, thus providing a spatially resolved single-cell deconvolution strategy
for these spatial data. Our results showed that Bulk2Space had great biological and
clinical application prospects, including linking tissue molecular characteristics with
histological phenotypes, revealing the spatial specific variation of cells, discovering the
spatial expression pattern of novel genes, achieving more refined cell clustering, and
predicting the molecular mechanism underlying the progression of the disease.

Methods

Datasets

All scRNA-seq, spatially resolved transcriptomics, and bulk RNA-seq datasets used in
this study were collected from high-quality publications, Gene Expression Omnibus
(GEO), and The Cancer Genome Atlas (TCGA), wherein unannotated, ambiguous, or
low quality-cells were excluded. The detailed description of each dataset was
summarized in Supplementary Data 1.

Computing environment

The Bulk2Space was developed on two workstations, as listed below.

Workstation 1: Dell Precision Tower 7820 Workstation, CPU (Intel Xeon Gold 5118, 2.3 GHz ×
2), RAM (64 GB, 16 GB × 4, DDR4, 2933 MHz), Hard Drive (SSD, SATA Class 20, 512 GB; HDD,
7200 rpm, SATA), Graphics Card (NVIDIA, Quadro P4000, 8 GB), Operating System (Ubuntu
16.04), Running Environment (CUDA 11.6, Torch 1.12.1, Python 3.8.5, deep-forest 0.1.5,
easydict 1.9, numpy 1.19.2, pandas 1.1.3, scanpy 1.8.1, scikit-learn 1.0.1, scipy 1.5.2, tqdm
4.50.2, Unidecode 1.3.0).

Workstation 2: Dell Precision Tower 7920 Workstation, CPU (Intel Xeon Gold 6230, 2.1 GHz ×
2), RAM (192 GB, 16 GB × 12, DDR4, 2933 MHz), Hard Drive (SSD, SATA Class 20, 512 GB; HDD,
7200 rpm, SATA), Graphics Card (NVIDIA, RTX2080Ti VIDEO CARD V2 × 2, 22 GB), Operating
System (Ubuntu 18.04), Running Environment (CUDA 11.0, Torch 1.7.1, Python 3.8.5, deep-
forest 0.1.5, easydict 1.9, numpy 1.19.2, pandas 1.1.3, scanpy 1.8.1, scikit-learn 1.0.1, scipy
1.5.2, tqdm 4.50.2, Unidecode 1.3.0).

Data processing

All data were preprocessed using R (version 4.1.1). For mouse brain scRNA-seq data by
Moffitt[43] et al., we filtered 841 ambiguous cells and 88 unstable cells. For human
prostate cancer scRNA-seq data by Wu[54] et al., we filtered 183 unassigned cells. All
cells of other scRNA-seq as well as spatially resolved transcriptomics datasets were
retained. For mouse hypothalamus MERFISH data, we filtered 5 "blank" barcodes as
well as *the Fos* gene, whose expression value in all cells was 'NA'. For human prostate
cancer RNA-seq data, we used biomaRt (version 2.48.3) to transform ensemble id to
gene name. For all scRNA-seq and spatially resolved transcriptomics datasets, the raw
counts were normalized using the global-scaling normalization method "LogNormalize"
by Seurat (version 4.0.4)[60]. For the MERFISH dataset, the normalized data was
determined as the raw count per cell divided by the cell volume and scaled by 1000.

Step 1: Deconvolution of bulk transcriptome data.

Cell type proportions prediction. For a given bulk RNA-seq dataset, we aimed to
calculate its cell-type proportions firstly, and then generate single-cell gene expression
profiles based on the calculated proportions. Since cell type and gene expression of all
single cells can be accessed from the scRNA-seq data reference, we collected cells with

the same cell type and then average their gene expression. This average vector $\mathbf{c}_i \in$

$\mathbb{R}^N$ is defined as the gene expression of cell type i ($i \in \{1,2,\dots,C\}$, where C

denotes the total number of cell types). Given the gene expression vector $\mathbf{x} \in \mathbb{R}^N$ of

a bulk RNA-seq dataset, we aimed to predict the cell type proportion:

540
$$\sum_{i=1}^{C} p_i \mathbf{c}_i = \mathbf{x} \quad (1)$$

541   Here, p_i is the proportion of cell type i.

542   We applied least square estimation (LSE) to estimate p_i by minimizing the

543   squared discrepancies between observed data.

544
$$\min_{\boldsymbol{p}} \|\mathbf{x} - \mathbf{C}\boldsymbol{p}\|_2 \quad (2)$$

where $\mathbf{C} \in \mathbb{R}^{N \times C}$ denotes the cell type gene expression profile, and each column

represents a different cell type. $\boldsymbol{p} = [p_1, p_2, \dots, p_C]^T$ is the proportion vector to be

estimated.

Single-cell simulation by β-VAE. Let $\mathcal{D} = \{X, V, W\}$ be a set consisting of gene

expression vectors $x \in \mathbb{R}^N$ and two sets of ground truth data generative factors:

conditionally independent factors $v \in \mathbb{R}^K$, where $\log p(v|x) = \sum_k \log p(v_k|x)$; and

conditionally dependent factors $w \in \mathbb{R}^H$. We assume that x are generated by a true

world simulator S using the corresponding ground truth data generative factors:

$p_\theta(x|v,w) = S(v,w)$, where θ is the generative model parameter.

We wanted to develop an unsupervised deep generative model that, using samples

from X only, can learn the joint distribution of the data x and a set of generative

latent factors z ($z \in \mathbb{R}^M$, where $M \geq K$) that can be used to generate the observed

data x; i.e., $p_\theta(x|z) \approx p(x|v,w) = S(v,w)$. However, since the integral of the

marginal likelihood $p_\theta(x) = \int p_\theta(\mathbf{z}) p_\theta(\mathbf{x}|\mathbf{z}) \, d\mathbf{z}$ is intractable (so we cannot

evaluate or differentiate the marginal likelihood), the true posterior density

$p_\theta(\mathbf{z}|x) = p_\theta(\mathbf{x}|\mathbf{z}) p_\theta(\mathbf{z}) / p_\theta(\mathbf{x})$ is intractable.

To solve this problem, for a given observation x, we described the inferred

posterior configurations of the latent factors $\mathbf{z}$ through a probability distribution

$q_\phi(\mathbf{z}|\mathbf{x})$: an approximation to the intractable true posterior $p_\theta(x|z)$. We aimed to

ensure that the inferred latent factors $q_\phi(\mathbf{z}|\mathbf{x})$ capture the generative factors v in a

disentangled manner. A disentangled representation can be defined as one where

single latent units are sensitive to changes in single generative factors, while being

relatively invariant to changes in other factors. In a disentangled representation,

knowledge about one factor can generalize to novel configurations of other factors.

The conditionally dependent data generative factors w can remain entangled in a

separate subset of $\mathbf{z}$ that is not used for representing v.

An intuitive approach was to minimize the KL divergence between the approximate

and the true posterior:

573
$$\mathcal{D}_{KL}\left(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z}|x)\right) = -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{z}|x)}{q_\phi(\mathbf{z}|\mathbf{x})}\right)$$

574
$$= -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{\frac{p_\theta(x,\mathbf{z})}{p_\theta(x)}}{q_\phi(\mathbf{z}|\mathbf{x})}\right)$$

575
$$= -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \left[\log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right) - \log(p_\theta(x))\right]$$

576
$$= -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right) + \log(p_\theta(x))$$

577
$$= -\mathcal{L}(\theta, \phi; \mathbf{x}) + \log(p_\theta(x)) \quad (3)$$

578 Here, $\mathcal{L}(\theta, \phi; \mathbf{x})$ is called the variational lower bound, and can be written as:

579
$$\mathcal{L}(\theta, \phi; \mathbf{x}) = \sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right)$$

580
$$= \sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})p_\theta(\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right)$$

581
$$= \sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \left[\log(p_\theta(\mathbf{x}|\mathbf{z})) + \log\left(\frac{p_\theta(\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right)\right]$$

$$582 \qquad = \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}\big[\log(p_\theta(\mathbf{x}|\mathbf{z}))\big] - \mathcal{D}_{KL}\Big(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\Big) \quad (4)$$

To encourage this disentangling property in the inferred $q_\phi(\mathbf{z}|\mathbf{x})$, we introduced

a constraint over it by trying to match it to a prior $p(\mathbf{z})$ that can both control the

capacity of the latent information bottleneck, and embodies the desiderata of

statistical independence mentioned above. We set the prior to be an isotropic unit

Gaussian $p(z) \sim \mathcal{N}(0, I)$, then the constrained optimization problem can be written

as:

$$589 \qquad \max_{\phi,\theta} \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}[\log(p_\theta(\mathbf{x}|\mathbf{z}))] \; subject \; to \; \mathcal{D}_{KL}\Big(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\Big) < \epsilon \quad (5)$$

590   where ϵ specifies the strength of the applied constraint. Re-writing the above

591   equation as a Lagrangian under the KKT conditions, we obtain:

$$592 \qquad \mathcal{F}(\theta, \phi, \beta; \mathbf{x}, \mathbf{z}) \; = \; \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}[\log(p_\theta(\mathbf{x}|\mathbf{z}))] - \beta\Big(\mathcal{D}_{KL}\Big(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\Big) - \epsilon\Big) \quad (6)$$

593   where the KKT multiplier β is the regularization coefficient that constrains the

594   capacity of the latent information channel $\mathbf{z}$ and puts implicit independence pressure

595   on the learnt posterior due to the isotropic nature of the Gaussian prior $p_\theta(\mathbf{z})$. Since

596      $\beta, \epsilon \geq 0$, according to the complementary slackness KKT condition, the equation can

597   be re-written as:

$$598 \qquad \mathcal{F}(\theta, \phi, \beta; \mathbf{x}, \mathbf{z}) \geq \mathcal{L}(\theta, \phi; \mathbf{x}, \mathbf{z}, \beta) = \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}[\log(p_\theta(\mathbf{x}|\mathbf{z}))] -$$

$$599 \qquad \beta\Big(\mathcal{D}_{KL}\Big(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\Big)\Big) \quad (7)$$

which is the β-VAE formulation with the addition of the β coefficient.

Here, different β will change the degree of applied learning pressure during

training, thereby encouraging different learned representations. We postulated that

to learn disentangled representations of the conditionally independent data

generative factors v, it is important to set $\beta > 1$ to impose a stronger constraint on

latent bottleneck than the original VAE. These constraints limit the capacity of $\mathbf{z}$,

coupled with the pressure to maximize the log likelihood of the training data $\mathbf{x}$ under

the model, encouraging the model to learn the most efficient representation of the

data. The additional pressure from high β values may create a trade-off between

reconstruction fidelity and the quality of disentanglement within the learned latent

representations. When the appropriate balance is found between information

preservation (reconstruction cost as regularization) and latent channel capacity

restriction ($\beta > 1$), disentangled representations emerge.

Model Configuration. In step1, we simulate single-cell by β-VAE. Both the encoder and

decoder apply a four-layer perceptron, where each layer is followed by a RELU

activation except the last layer of the encoder. For the encoder, the number of neurons

in each layer is 2048, 1024, 512 and 512 respectively. While for the decoder, the

number of neutrons in each layer is 512, 1024, 2048 and k respectively, where k

represents the number of genes in the dataset. The dimensionality of the latent space

learned by the beta-VAE is 256. The relative weighting of the reconstruction loss and

regularization loss is 1:4. We apply the Adam with weight decay (AdamW) optimizer

with an initial learning rate as 1e-4, the decoupled weight decay as 5e-4 and Adam's β

parameters as 0.9 and 0.999. Moreover, the default running epoch is fixed to 3000,

but we use early stopping during the training phase, with which we stop training when

the training loss no longer reduces for 50 epochs.

Here we introduce the implementation details of our benchmark approaches. For GAN

and cGAN, the generator consists of two fully connected layers, followed by LeakyRELU

and RELU activations respectively. The discriminator is also made up of two fully

connected layers, where the first layer is followed by a LeakyRELU activation. We use

the Adam optimizer with the initial learning rate of 1e-4 and the betas parameters of

0.5 and 0.999. We train util the loss in the generative and discriminative phases is no

longer reduced for 50 epochs, and use the model obtained at this time as the final

model for prediction.

Step2: Mapping generated single cells to spatial locations

Spatial barcoding-based RNA-seq. After obtaining the generated data, we needed to

map each cell and the spot it belongs to. To improve the accuracy and reduce the

complexity of this process, we divided it into two steps. First, we calculated the cell

type proportion of each spot, and then predicted which cells are contained in each

spot based on this proportion.

1) Calculate cell type proportions. The procedure is identical to the Step 1
deconvolution, we repeated this process to obtain the cell type proportion of each
spot in the tissue.

2) Deep forest for spot recommendation. With cell type proportion, we next
predicted which cells are contained in each spot. We defined this task as a binary
classification problem. We designed a classifier, whose input is the gene expression
vectors of both single cell $\mathbf{s}$ and spot $\mathbf{x}$. When the cell belongs to the spot, the output
of the classifier is 1, otherwise the output is 0. Specifically, the input can be
represented as $\mathbf{v} = [\mathbf{s}; \mathbf{x}; \mathbf{s} - \mathbf{x}] \in \mathbb{R}^{N \times 3}$, where semicolon means concatenate
operation. $\mathbf{s} - \mathbf{x} \in \mathbb{R}^{N}$ can be regarded as an auxiliary signal, since when there is a
negative number occurring in this vector, the single cell must not belong to the spot.

We applied multi-Grained Cascade Forest (gcForest), a decision tree ensemble with
a cascade structure and further enhanced by multi-grained scanning, to build our
classifier. Each level of cascade receives feature information processed by its preceding
level and outputs its processing result to the next level. Each level is an ensemble of
decision tree forests, i.e., an ensemble of ensembles. We also included completely-
random tree forests) to encourage diversity. Given an instance, each forest counts the
percentage of different classes of training samples at the leaf node where the
concerned instance falls, and then average across all trees in the same forest to
produce an estimate of class distribution. The estimated class distribution forms a class
vector, which was generated by k-fold cross validation and then concatenated with
the original feature vector to be input to the next level of the cascade.

To enhance cascade forest, we introduced a procedure of multi-grained scanning.
We used sliding windows to scan the raw features. Specially, we used a window with
a fixed length of k to slide the input vector into k-dimensional feature vectors. We
then fed these vectors to n forests and finally produce n two-dimensional
vectors. By using multiple sizes k of sliding windows, differently grained feature
vectors were generated.

The we summarized the overall procedure of gcForest. For the original raw features,
three window sizes $\{\lfloor N/16 \rfloor, \lfloor N/8 \rfloor, \lfloor N/4 \rfloor\}$ were used for multi-grained scanning.
The generated data were used to train a completely-random tree forest containing 100
trees. We concatenated the output of these two forests as the transformed feature

vectors, which also acted as the input of cascade forest. The final model was a cascade

of cascades, where each cascade consisted of multiple levels each corresponding to a

grain of scanning. Each level consisted of 4 completely-random tree forests, each

containing 100 trees. In other words, the transformed feature vectors were

augmented with the class vector generated by the previous grade, and then were used

to train the current grade of cascade forests.

Given a test instance, it would obtain its corresponding transformed feature vector

through the multi-grained scanning procedure, and then go through the cascade till

the last level. The two-dimensional class vectors at the last level were aggregated, and

the class with the largest aggregate value was selected to obtain the final prediction

result.

This decision tree ensemble approach not only had much fewer hyper-parameters

than deep neural networks, but also retained the interpretability of tree models. Its

model complexity could be automatically determined in a data-dependent manner,

which made gcForest work well even on small-scale data.

Image-based in situ hybridization. We developed a simple method for predicting the

spatial distributions of genes not measured in spatial transcriptomic data which

produced by in situ RNA imaging-based technologies. To do this, Bulk2Space used the

spatial transcriptomic data as the reference, and generated scRNA-seq data were

mapped to the tissue space corresponding to this reference. To eliminate the

differences between scRNA-seq data and spatial transcriptomic data caused by

different experiment types, an empirical Bayes framework was used to remove the

batch effect, so that both sets of data are at the same scale level. Subsequently, the

cross-dataset k-nearest neighbor graph of spatial transcriptomic data in scRNA-seq

data was computed in the aligned space, and then the predicted whole gene

expression profile of each cell of spatial transcriptomic data could be calculated to the

mean of its k nearest neighbors in scRNA-seq data.

Model Configuration. Deep forest is applied for spot recommendation. Specifically, the

number of samples used to construct feature discrete bins is set to 200000. If the size

of training set is smaller than it, then all training samples will be used. The type of

binner used to bin feature values into integer-valued bins is "percentile", which means

each bin will have approximately the same number of distinct feature values. We set

the maximum number of cascade layers in the deep forest as 20, and apply 2 estimator
in each cascade layer. Gini impurity is used to measure the quality of a split. We have
no constraints on the maximum depth of each tree. The training process terminates
when the validation performance on the training set does not improve compared
against the best validation performance achieved so far to 2 tolerant rounds. And the
counting on tolerant rounds is triggered if the performance of a fitted cascade layer
does not improve by 1e-5 compared against the best validation performance achieved
so far.

We also introduce the implementation details of our benchmark approaches. (1) For
Logistic Regression (LR), we implement L2 regularization as the additional penalty term
to solve the problem of overfitting. We use L-BFGS algorithm as the solver, which uses
the Hessian matrix to iteratively optimize the loss function. We fit the model according
to the given training data and return the probability estimates on testing set. (2) For
Decision Tree (DT), we use the Gini impurity to measure the quality of a split and
choose the best split at each node. We don't set any maximum depth of the tree, so
nodes are expanded until all leaves are pure or until all leaves contain less than 2
samples. We build the decision tree classifier from the training set, and then predict
class probabilities of the input samples. (3) For Gradient Boosting Decision Tree(GBDT),
we set the learning rate as 0.1 and the number of boosting stages to perform as 100.
The loss function to be optimized is log loss function, and we choose Friedman MSE to
measure the quality of a split. The maximum depth of the individual regression
estimators is set to 3. We fit the gradient boosting model on the training set and use
the trained model to make predictions on testing set. (4) For Multilayer
Perceptron(MLP), we apply a two-layer perceptron, where each layer is followed by a
batch normalization, a RELU activation, and a dropout layer with the probability of
element zeroing as 0.1. We use the Adam optimizer with the initial learning rate of 1e-
4 and the betas parameters of 0.9 and 0.999. We train until the loss is no longer
reduced for 30 epochs, and use the model obtained at this time as the final model for
prediction. (5) For DeepGBM, we set the number of tree groups to 100. The dimension
of leaf embedding for a tree group is set to 20. The structure of the distilled NN model
is a fully connected network with "100-100-100-50" hidden layers. We adopt the
feature selection in each tree group, where we first sort the features according to the
information gain, and the top 128 of them are selected as the inputs of distilled NN
model. We use the Adam with weight decay (AdamW) optimizer with an initial learning
rate as 2e-3, the decoupled weight decay as 1e-6, and Adam's betas parameters as 0.9

and 0.999. We train until the model reaches the highest ROC-AUC score, and use it for
prediction.

Performance evaluation of Bulk2Space (Simulated datasets)

Benchmarking the deconvolution step of Bulk2Space on simulated datasets

1. Simulated bulk data and the single-cell reference were generated from same
datasets (paired simulations)

Datasets. Ten scRNA-seq datasets were applied to benchmark our method, including
human and mouse primary tissues. Five human scRNA-seq datasets including
peripheral blood[46] ("GSE92495
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE92495]"), brain[61]
("GSE103723 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE103723]"),
kidney[62] ("GSE121862
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE121862]"), liver[63]
("GSE124395 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE124395]"),
and lung[64] ("GSE130148
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130148]"). Five mice
scRNA-seq datasets including brain[65] ("GSE60361
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE60361]"), kidney[66]
("GSE119531 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE119531]"),
lung[67] ("GSE127465
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE127465]"), pancreas[68]
("GSE84133 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE84133]"), and
testis[69] ("GSE112393
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE112393]").

Data simulations. The simulated data were derived from the above 10 scRNA-seq
datasets. We randomly divided the scRNA-seq data into two parts, one was treated as
the single-cell reference (data_2) and the other (data_1) for the construction of bulk
transcriptome data via aggregating all the single-cell gene expression profiles.
Considering the cell composition of bulk RNA-seq data varies greatly in the natural
situation, for each data_1, we further changed the cell-type proportion and

synthesized 3 corresponding bulk transcriptome data with different cell compositions. In total, 30 paired simulated data were synthesized in this study. The detailed description of the experimental design was summarized in Supplementary Data 2.

Compared methods. **a**, Generative adversarial networks (GAN), **b**, conditional generative adversarial networks (CGAN). The generator consists of two fully connected layers, followed by LeakyRELU and RELU activations respectively. The discriminator is also made up of two fully connected layers, where the first layer is followed by a LeakyRELU activation. We use the Adam optimizer with the initial learning rate of 1e-4 and the betas parameters of 0.5 and 0.999. We train until the loss in the generative and discriminative phases is no longer reduced for 50 epochs, and use the model obtained at this time as the final model for prediction.

Benchmark metrics. Pearson correlation coefficient (PCC), Spearman's rank correlation coefficient (SRCC), and root mean squared error (RMSE) were used to assess the similarity of the gene expression profile between scRNA-seq data reference and synthetic bulk transcriptome data.

2. Simulated bulk data and the single-cell reference were generated from different datasets (unpaired simulations)

Datasets. Eight human pancreas scRNA-seq datasets from different resources were also applied to benchmark our method, including 1 CelSeq[70] ("GSE81076 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE81076]"), 1 CelSeq2[71] ("GSE85241 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE85241]"), 1 Fluidigm C1[72] ("GSE86469 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE86469]"), 4 inDrops[68] ("GSE84133 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE84133]"), and 1 SMART-Seq2[73] (E-MTAB-5061).

Data simulations. The simulated data were derived from the above 8 scRNA-seq datasets. We randomly selected one as the single-cell reference and another dataset from different resources as bulk transcriptome data. The detailed description of the experimental design was summarized in Supplementary Data 3.

Compared methods. **a**, Generative adversarial networks (GAN). **b**, conditional

generative adversarial networks (CGAN). The model details were described above. **c**,

bMIND[41]. We followed the guidelines on the bMIND GitHub repository:

https://github.com/randel/MIND. We set the cell type proportion parameter "frac" to

the output value of Bulk2Space to ensure the generated gene expression profile

comparability.

Benchmark metrics. The benchmark metrics were described above.

Evaluating the robustness of the deconvolution step of Bulk2Space

we evaluated the robustness of Bulk2space by introducing two kinds of noise: cell type

noise and gene expression noise. For cell type noise, we randomly altered the type of

cells with the ratio of 0.01, 0.02, 0.04, 0.08, 0.1, 0.2, 0.3, 0.4, 0.5, 0.7, and 0.9. For gene

expression noise, we firstly constructed a noise expression profile C,

$$C \sim \mathrm{U}(-x, x) \quad (8)$$

Here, x is the maximum gene expression value of scRNA-seq data reference.

We next constructed synthetic expression matrix by combining noise expression

profile with scRNA-seq data reference, and the ratio of gene expression-altered cells

remains 0.01, 0.02, 0.04, 0.08, 0.1, 0.2, 0.3, 0.4, 0.5, 0.7, and 0.9. PCC, SRCC and RMSE

between the synthetic expression matrix and constructed bulk RNA-seq data were

used to evaluate performance.

Benchmarking the spatial mapping step of Bulk2Space on simulated datasets

1. Spatial barcoding-based data

Datasets. Eighteen datasets metioned above were applied to benchmark our method.

10 datasets were applied to construct single-cell data and spatial reference from same

datasets, and other 8 datasets were applied to construct single-cell data and spatial

reference from different datasets.

Data simulations. Starting from scRNA-seq data reference, we randomly chose 10 cells

from it and aggregated their transcriptomic profiles as a spot of pseudo spatial

transcriptomic data. The spot with over 25000 UMI counts would be sampled down to

20000 UMI counts to better meet the true situation. We also constructed pseudo

spatial transcriptomic data with 100, 200, 500, 1000, and 5000 spot numbers to

simulate true spatial transcriptomics data produced by different spatial barcoding

technologies. Besides, the simulation that the scRNA-seq and spatial datasets are from

different sources was also considered. The detailed description of the experimental

design was summarized in Supplementary Data 4 (paired simulation) and

Supplementary Data 5 (unpaired simulation).

Compared methods. **a**, Logistic Regression (LR). We implement L2 regularization as

the additional penalty term to solve the problem of overfitting. We use L-BFGS

algorithm as the solver, which uses the Hessian matrix to iteratively optimize the loss

function. We fit the model according to the given training data and return the

probability estimates on testing set. **b**, Decision Tree (DT). We use the Gini impurity to

measure the quality of a split and choose the best split at each node. We don't set any

maximum depth of the tree, so nodes are expanded until all leaves are pure or until all

leaves contain less than 2 samples. We build the decision tree classifier from the

training set, and then predict class probabilities of the input samples. **c**, Gradient

Boosting Decision Tree (GBDT). We set the learning rate as 0.1 and the number of

boosting stages to perform as 100. The loss function to be optimized is log loss function,

and we choose Friedman MSE to measure the quality of a split. The maximum depth

of the individual regression estimators is set to 3. We fit the gradient boosting model

on the training set and use the trained model to make prediction on testing set. **d**,

Multilayer Perceptron (MLP). We apply a two-layer perceptron, where each layer is

followed by a batch normalization, a RELU activation, and a dropout layer with the

probability of element zeroing as 0.1. We use the Adam optimizer with the initial

learning rate of 1e-4 and the betas parameters of 0.9 and 0.999. We train until the loss

is no longer reduced for 30 epochs, and use the model obtained at this time as the

final model for prediction. **e**, DeepGBM. We set the number of tree groups to 100. The

dimension of leaf embeddin for a tree group is set to 20. The structure of the distilled

NN model is a fully connected networks with "100-100-100-50" hidden layers. We

adopt the feature selection in each tree group, where we first sort the features

according to the information gain, and the top 128 of them are selected as the inputs

of distilled NN model. We use the Adam with weight decay (AdamW) optimizer with

an initial learning rate as 2e-3, the decoupled weight decay as 1e-6, and Adam's betas

parameters as 0.9 and 0.999. We train until the model reach the highest ROC-AUC

score, and use it for prediction. **f**, RCTD[34]. We followed the guidelines on the RCTD

GitHub repository: https://github.com/dmcable/spacexr. We set the parameter doublet_mode = 'full'. **g**, spatialDWLS[35]. We followed the guidelines on the spatialDWLS GitHub repository: https://github.com/rdong08/spatialDWLS_dataset/tree/main/codes. We set the parameter n_cell = 20. **h**, stereoscope[36]. We followed the guidelines on the stereoscope GitHub repository: https://github.com/almaan/stereoscope. We set the parameter sc epochs = 10000, st epochs = 10000. **i**, SPOTlight[37]. We followed the guidelines on the SPOTlight GitHub repository: https://github.com/MarcElosua/SPOTlight. We set the parameter n_cells = 75.

Benchmark metrics. Pearson correlation coefficient (PCC), Spearman's rank correlation coefficient (SRCC), and root mean squared error (RMSE) were used to assess the similarity of the gene expression profile per spot between the predict result and ground truth. Due to RCTD, spatialDWLS, stereoscope, and SPOTlight could not obtain gene expression profiles for each spot, we calculated gene signature matrix of each cell type based on single-cell reference, and the product of the predicted proportions of each cell type and the gene signature matrix of each cell type was regarded as the gene expression profile of each spot.

2. Image-based in situ hybridization data

Datasets. A mouse hypothalamus scRNA-seq data ("GSE113576 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE113576]") and MERFISH data (Bregma +0.26) (image-based reference) from the same laboratory[43] were applied to test the performance of the Bulk2Space algorithm for the second spatial mapping strategy. We also applied another mouse hypothalamus scRNA-seq data[44] ("GSE87544 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE87544]") from different source as single-cell reference to confirm the robustness of Bulk2Space.

Benchmark metrics. Pearson correlation coefficient (PCC) was used to assess the similarity of the gene expression profile between the predict result and ground truth. We took two strategies, five-fold cross validation and leave-one-out cross validation, to validate performance of Bulk2Space. In five-fold cross validation, we random split 150 target genes identified in MERFISH into five folds using R package caret (version

891 6.0-92), a split of 80% for reference and 20% for validation. In leave-one-out cross

892 validation, 149 were used as reference genes and and one was used for validation.

Performance evaluation of Bulk2Space using biological datasets

Deconvolution performance evaluation using paired bulk and single-cell datasets

895 Three paired mouse liver bulk and single-cell datasets[74] ("GSE119340

896 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE119340]") were applied to

897 evaluate the performance of the deconvolution step of Bulk2Space.

Reconstruction of mouse hippocampus subregions at single-cell resolution

The mouse hippocampus Slide-seq v2[12] and scRNA-seq data[45] were downloaded from

the Seurat website: https://satijalab.org/seurat/articles/spatial_vignette.html were

applied to evaluate the performance of the spatial mapping step of Bulk2Space on real

single-cell data and real spatial data. We down sampled Slide-seq v2 data to 5000 spots

to speed the spatial mapping step. For Bulk2Space, we set the parameter k = 2 and

top_marker_num = 100, all other parameter followed the default values. For Seurat,

we followed the standard analysis workflow on the Seurat website and set the

parameter resolution = 0.3 to perform unsupervised clustering.

Robustness evaluation of Bulk2Space using repeated data

We conducted 100-time repetitions on the deconvolution step of Bulk2Space to

evaluate the robustness of β-VAE using the human peripheral blood[46] ("GSE92495

[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE92495]") scRNA-seq data.

The construction of the single-cell reference and bulk transcriptome data was the

same as mentioned above. We further evaluated the spatial mapping results using the

pancreatic ductal adenocarcinoma (PDAC) dataset[47] and the human melanoma

dataset[48, 49] with 3-time repetitions. For Bulk2Space, all parameters followed the

default values.

Deconvolution of bulk RNA-seq data using annotation-free single-cell reference by Bulk2Space

We further evaluated the performance of Bulk2Space without providing cell type

information. We randomly divided the human peripheral blood[46] ("GSE92495

[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE92495]") scRNA-seq data

into two parts, one was treated as the single-cell reference and the other for the

construction of bulk transcriptome data via aggregating all the single-cell gene

expression profiles. For the single-cell reference, we followed the scRNA-seq data

analysis   workflow   on   the   Seurat   website:

https://satijalab.org/seurat/articles/pbmc3k_tutorial.html to get 5 "clustering spaces"

using "FindClusters" function with the parameter resolution = 0.2. For Bulk2Space, all

parameters followed the default values.

Revealing spatial, molecular, and functional heterogeneity of B cells in melanoma

(consecutive slices)

The   scRNA-seq   data   ("GSE72056

[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE72056]") by Tirosh[48] et. al.

was used as the single-cell reference, and two consecutive slices of the ST data by

Thrane[49] et. al. were used as spatial reference (slice 1) and synthesized bulk data (slice

2) (via aggregating all the spots gene expression profiles), respectively. For Bulk2Space,

we set the parameter epoch_num = 3500, k = 10, and top_marker_num = 500, all other

parameters followed the default values.

We firstly evaluated the expression correlation of marker genes for five major cell

types between generated single-cell expression profiles from synthesized bulk data

(slice 2) and ground truth. We applied "FindAllMarkers" function of Seurat (version

4.0.4)[60] to calculate the marker genes of each cell type with the parameter

logfc.threshold = 0.5. We also evaluated the expression correlation of marker genes

for five major cell types between spatial mapping result of Bulk2Space and spatial

reference (slice 1).

For B cells spatial heterogeneity analysis, we determined two spatial areas (lymphoid

area and tumor area) based on histological annotation firstly. Then, differentially

expressed genes analysis between B cells from different areas was applied by the

"FindAllMarkers" function with the parameter logfc.threshold = 0.25. For these

differentially expressed genes, we further performed the pathway enrichment analysis

using the Metascape (https://metascape.org) to investigate the biological functions.

Linking histomorphology and transcriptomics in PDAC (disrete slices)

Bulk2Space was performed for another PDAC dataset[47]. We selected PDAC-B scRNA-seq data as the single-cell reference, and two inconsecutive slices of the ST data were used as spatial reference (PDAC-B ST1, slice 1) and synthesized bulk data (PDAC-B ST2, slice 2) (via aggregating all the spots gene expression profiles), respectively. For Bulk2Space, we set the parameter epoch_num = 3500, k = 10, and top_marker_num = 500, all other parameters followed the default values. The evaluation procedure of the marker gene expression correlation between Bulk2Space results and slices was same as it in the human melanoma dataset analysis.

Application of Bulk2Space

Bulk2Space integrates spatial gene expression and histomorphology in PDAC

The pancreatic adjacent tissues and pancreatic cancer tissues bulk data by Wu[50] et.al. were downloaded from "GSE171485 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE171485]". We firstly applied the PDAC-A scRNA-seq data by Moncada[47] et. al. as the single-cell reference to deconvolute pancreatic adjacent tissues bulk data and pancreatic cancer tissues bulk data respectively. We then applied the PDAC-A ST1 by Moncada[47] et. al. as the spatial reference and performed spatial mapping step for the generated single-cell gene profiles from pancreatic cancer tissues bulk data. For Bulk2Space, we set the parameter epoch_num = 3500, k = 10, and top_marker_num = 200, all other parameters followed the default values.

Reconstruction of mouse isocortex layers at single-cell resolution using Spatial-seq data

The mouse isocortex bulk data was produced by Spatial-seq (see Methods). We applied a mouse primary visual cortex scRNA-seq data[52] and two tissue sections (anterior section 1 and posterior section 1) sequenced by 10X Visium as the single-cell reference and spatial references, respectively. For Bulk2Space, we set the parameter epoch_num = 3500 and top_marker_num = 300, all other parameters followed the default values.

Reconstruction of mouse hypothalamus structure at single-cell resolution using Spatial-seq data

The mouse hypothalamus bulk data was produced by Spatial-seq (see Methods). We applied a mouse hypothalamus scRNA-seq data ("GSE113576 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE113576]") and MERFISH data (Bregma -0.04) by Moffitt[43] et.al. as the single-cell reference and spatial reference, respectively. For Bulk2Space, we set the parameter epoch_num = 3500 and top_marker_num = 500, all other parameters followed the default values. For those ambiguous cells in MERFISH data, we re-annotated their cell type based on the whole gene expression profiles predicted by Bulk2Space.

Uncovering spatial gene expression dynamics in inflammation-induced prostate cancer

The human prostate cancer bulk data was downloaded from TCGA (https://portal.gdc.cancer.gov). We selected the human prostate cancer scRNA-seq data by Wu[54] et.al. as the single-cell reference, and three ST data from different tissue section location as spatial references (p1.1 Normal glands, p2.3 Inflammation, p3.3 Cancer Gs 3+4). For Bulk2Space, we set the parameter epoch_num = 3500, k = 10, and top_marker_num = 500, all other parameters followed the default values. We then compared the distribution of various cell types in these three different tissue regions.

Spatial-seq protocol

Multiplexed spatial barcoding. A virtual 48×48×48 three-dimensional well array (barcoding array) was constructed in X, Y, and Z directions, with 48×48 wells in each layer. 384-well plates (24×16 wells) were selected as the basic units (2×3 plates) of each layer. Each well in the barcoding array can be labeled as barcode $(\{X_i, Y_j, Z_k\}, i, j, k \in N, [1, 48])$. Magnetic beads coated with carboxyl group (Cat. # 40200, purchased from BEAVER Biomedical Engineering Co., Ltd.) were distributed into each well for barcode synthesis. There were three rounds of barcode extension reactions. In the first round, 5' amino-modified barcoded oligonucleotides (barcode X_i) were conjugated to the beads. In the second round of extension reaction, barcoded oligonucleotides (barcode Y_j) which had a sequence of bases at the 5' end complementary to the 3' end of barcode X_i were linked to the end of barcode X_i by

PCR. Similarly, in the third round of reaction, barcode sequences containing a unique

molecular identifier (UMI) and a polyT tail (barcode Z_k) were extended to barcode Y_j

by PCR. All oligonucleotides were purchased from Sangon Biotech (Shanghai) Co., Ltd.

This method only needs 144 (48×3) different barcodes to theoretically generate

magnetic beads with 48^3 distinct kinds of magnetic beads with known sequences.

More importantly, the entire synthesis process was serialized and multiplexed, which

can be accomplished with the help of arrayed pipettes or liquid workstations, rather

than synthesizing the magnetic beads well by well. The spatially barcoded beads can

be used in many areas, such as multiplexed labeling of tissue samples or even single

cells.

Collection of mouse brain regions. Three wild-type adult C57BL/6J mice (SPF, male, 20-

25 g) aged 8-10 weeks were used for Spatial-seq experiments. The environmental

conditions in the mouse facility were: 12 h light and 12 h dark cycle (light on from 8:00

a.m. to 8:00 p.m.), light intensity range of 15-20 lux, temperature range of 22–26 °C,

humidity range of 40–70%, and free access to food and water. The use and care of the

mice were in accordance with the guidelines of the Animal Advisory Committee of

Zhejiang University and the US National Institutes of Health Guidelines for the Care

and Use of Laboratory Animals. All procedures were approved by the Animal Advisory

Committee of Zhejiang University. The mouse brain was sliced into 14-μm sections

from the coronal and sagittal directions. Each tissue slice was registered to a reference

brain template provided by the Allen Brain Atlas (https://portal.brain-map.org/). After

spatial registration, the anatomical regions of the mouse brain were delineated and

annotated. The outlines, dissection sequences, and collectors of the brain regions

were specified from the annotated data and then imported to an LCM instrument

(Laser Microdissection System equipped with a DM68 microscope, a Leica LMD6 laser

cutter, and a single-cell capture collector LMT350, Leica Microsystems, Germany).LCM

is a microscope-guided powerful cutting system incorporating UV light for contact- and

contamination-free isolation of areas of interest from tissue sections[15]. Brain regions

were dissected from the tissue by LCM according to the imported files and fell into the

collector loaded with barcoded beads in advance. Spatial-seq has the potential to

achieve spatially resolved whole transcriptome sequencing of a large number of single

cells.

Multiplex RNA-seq. The tissue was lysed in the barcoded well, allowing mRNA to be

captured by polyT tail on the surface of the magnetic beads. The lysis buffer was

prepared with Tris-HCl (120 μL, pH 7.5, Cat. # T1140, purchased from Beijing Solarbio

Science & Technology Co., Ltd.), LiCl (80 μL, Cat. # AM9480, purchased from Invitrogen),

10% SDS solution (120 μL, RNase-free, Cat. # AM9823, purchased from Invitrogen),

EDTA (16 μL, Cat. # ST066, Shanghai beyotime Biological Co.,Ltd.), 0.5 M DTT solution

(16 μL, DNase, RNase & Protease free, Cat. # ST041, Shanghai beyotime Biological

Co.,Ltd.), and water (852 μL, nuclease-free, Cat. # AM9930, Ambion). Each well was

loaded with lysis buffer. The tissue was lysed on ice for 12 minutes. The beads were

then washed and transferred for reverse transcription (RT), The captured mRNA was

reverse transcribed using PrimeScript II Reverse Transcriptase (Cat. # 2690A, Takara).to

construct cDNA libraries following the guidance of Illumina Nextera XT DNA Library

Preparation Kits. A paired-end sequencing was conducted to decode the spatial

barcode in the 3' end and detect RNA species in the 5' end of the cDNA.

Sequence alignment. FastQC was utilized for quality control of the RNA-seq data

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). The derived RNA-seq

data were fragmented into lots of files based on the spatial barcodes. The digital gene

expression matrix for each barcode was retrieved follow the Drop-seq sequence

alignment cookbook.

Data normalization. FPKM, RPKM, RPM, and TPM were calculated from the gene

expression matrix according to the following formulas.

$$1064 \quad \text{FPKM} = \frac{ExonMappedFragments*10^9}{TotalMappedFragments*ExonLength} \quad (9)$$

$$1065 \quad \text{RPKM} = \frac{(ExonMappedRead/TotalMappedReads*10^6)}{ExonLength} * 10^3 \quad (10)$$

$$1066 \quad \text{RPM} = \frac{ExonMappedReads*10^6}{TotalMappedReads} \quad (11)$$

$$1067 \quad \text{TPM} = \frac{N_i/L_i*10^6}{sum(N_1/L_1+N_2/L_2+\cdots+N_n/L_n)} \quad (12)$$

The data were normalized using the global-scaling normalization method

"LogNormalize" in Seurat. A Bayesian network-based method termed ComBat was

used for batch effect removal of data derived from different cDNA libraries. The

expression matrix of each sample was normalized using the "Deconvolution

Normalization" algorithm to align the median gene expression.

Ethical statement. All experiments were approved by and conducted in accordance
with the ethical guidelines of the Zhejiang University Animal Experimentation
Committee (Protocol number, 14875).

1076 Data Availability

The original data used in this paper can be accessed through the following links: (1)
single-cell RNA-seq data of the human blood: GEO accession: "GSE92495
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE92495]"[46]; (2) single-cell
RNA-seq data of the human brain: GEO accession: "GSE103723
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE103723]"[61]; (3) single-cell
RNA-seq data of the human kidney: GEO accession: "GSE121862
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE121862]"[62]; (4) single-cell
RNA-seq data of the human liver: GEO accession: "GSE124395
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE124395]"[63]; (5) single-cell
RNA-seq data of the human lung: GEO accession: "GSE130148
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130148]"[64]; (6) single-cell
RNA-seq data of the mouse brain: GEO accession: "GSE60361
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE60361]"[65]; (7) single-cell
RNA-seq data of the mouse kidney: GEO accession: "GSE119531
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE119531]"[66]; (8) single-cell
RNA-seq data of the mouse lung: GEO accession: "GSE127465
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE127465]"[67]; (9) single-cell
RNA-seq data of the mouse pancreas: GEO accession: "GSE84133
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE84133]"[68]; (10) single-cell
RNA-seq data of the mouse testis: GEO accession: "GSE112393
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE112393]"[69]; (11) single-
cell RNA-seq data of the human pancreas with 975 cells: GEO accession: "GSE81076
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE81076]"[70]; (12) single-cell
RNA-seq data of the human pancreas with 2133 cells: GEO accession: "GSE85241
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE85241]"[71]; (13) single-cell
RNA-seq data of the human pancreas with 597 cells: GEO accession: "GSE86469
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE86469]"[72]; (14) four sets
of single-cell RNA-seq data of the human pancreas with 1635, 1562, 3330, 1230 cells,

respectively: GEO accession: "GSE84133
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE84133]"[68]; (15) single-cell
RNA-seq data of the human pancreas with 2288 cells[73]
(https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-5061/); (16) single-cell
RNA-seq data of the mouse hypothalamus using 10X Genomics: GEO accession:
"GSE113576 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE113576]"[43];
(17) single-cell RNA-seq data of the mouse hypothalamus using Drop-seq: "GSE87544
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE87544]"[44]; (18) three sets
of single-cell RNA-seq data of the mice liver: GEO accession: "GSE119340
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE119340]"[42]; (19) single-
cell RNA-seq data of the mouse hippocampus region[45]
(https://www.dropbox.com/s/cs6pii5my4p3ke3/mouse_hippocampus_reference.rds
?dl=0); (20) two sets of single-cell RNA-seq data of the human pancreatic ductal
adenocarcinoma (PDAC): GEO accession: "GSE111672
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE111672]"[47]; (21) single-
cell RNA-seq data of the human melanoma: GEO accession: "GSE72056
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE72056]"; (22) single-cell
RNA-seq data of the mouse cortex region[52]
(https://www.dropbox.com/s/dl/cuowvm4vrf65pvq/allen_cortex.rds); (23) single-cell
RNA-seq data of the human prostate cancer[54]
(https://singlecell.broadinstitute.org/single_cell/study/SCP1415); (24) MERFISH data
of the mouse hypothalamic preoptic region at bregma 0.26[43]
(https://datadryad.org/stash/dataset/doi:10.5061/dryad.8t8s248); (25) slide-seq v2
data of the mouse hippocampus region[12]
(https://singlecell.broadinstitute.org/single_cell/study/SCP815/highly-sensitive-
spatial-transcriptomics-at-near-cellular-resolution-with-slide-seqv2); (26) three sets of
spatially resolved transcriptomics data of the human PDAC using "Spatial
Transcriptomics": GEO accession: "GSE111672
[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE111672]"[47]; (27) two sets
of spatially resolved transcriptomics data of the human melanoma using "Spatial
Transcriptomics"[49] (https://www.spatialresearch.org/resources-published-
datasets/doi-10-1158-0008-5472-can-18-0747/); (28) two sets of 10X Visium data of
the mouse anterior cortex and posterior cortex regions
(https://www.10xgenomics.com/cn/resources/datasets/mouse-brain-serial-section-
1-sagittal-anterior-1-standard-1-1-0); (29) two sets of spatially resolved

transcriptomics data of the human prostate cancer using "Spatial Transcriptomics"[55] (https://www.spatialresearch.org/resources-published-datasets/10-1038-s41467-018-04724-5/); (30) three sets of bulk RNA-seq data of the mice liver: GEO accession: "GSE119340 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE119340]"[42]; (31) two sets of bulk RNA-seq data of the human PDAC: GEO accession: "GSE171485 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE171485]"[50]; (32) bulk RNA-seq data of the human prostate cancer (https://portal.gdc.cancer.gov).

The bulk RNA-seq data of the mouse cortex and mouse hypothalamus regions reported in this manuscript using our in-house developed Spatial-seq have been deposited to the Gene Expression Omnibus under accession number "GSE192999 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE192999]".All other relevant data supporting the key findings of this study are available within the article and its Sup-plementary Information files or from the corresponding author upon reasonable request. All source data are provided with this paper.

Statistics & Reproducibility

In this study, 152 simulated datasets (the data points of paired simulations for deconvolution, unpaired simulations for deconvolution, paired simulations for spatial mapping, and unpaired simulations for spatial mapping are 30, 12, 50, and 60, respectively), 13 biological datasets, and 2 experimental datasets were used to evaluate the Bulk2Space algorithm. Unannotated, ambiguous, or low quality-cells were excluded from the analysis. Pearson correlation coefficient, Spearman's rank correlation coefficient, and root mean squared error were used to compare the performance between different methods. For paired data simulations, single cells were randomly selected to synthesize the reference and the test set. For unpaired data simulations, single-cell profiles were randomly selected to synthesize the reference and the test set. The Investigators were not blinded to allocation during experiments and outcome assessment.

Code Availability

Bulk2Space is available as a python package and the source code is deposited at the GitHub (https://github.com/ZJUFanLab/bulk2space)[75].

1170

References

1171

1. Liao, J., Lu, X., Shao, X., Zhu, L. & Fan, X. Uncovering an Organ's Molecular Architecture at Single-Cell Resolution by Spatially Resolved Transcriptomics. *Trends Biotechnol* **39**, 43-58 (2021).

2. Wang, Y.S. & Guo, J. Multiplexed Single-Cell in situ RNA Profiling. *Front Mol Biosci* **8**, 775410 (2021).

3. Srivatsan, S.R. et al. Embryo-scale, single-cell spatial transcriptomics. *Science* **373**, 111-117 (2021).

4. Waylen, L.N., Nim, H.T., Martelotto, L.G. & Ramialison, M. From whole-mount to single-cell spatial assessment of gene expression in 3D. *Commun Biol* **3**, 602 (2020).

5. Qian, X. et al. Probabilistic cell typing enables fine mapping of closely related cell types in situ. *Nat Methods* **17**, 101-106 (2020).

6. Larsson, L., Frisen, J. & Lundeberg, J. Spatially resolved transcriptomics adds a new dimension to genomics. *Nat Methods* **18**, 15-18 (2021).

7. Chen, K.H., Boettiger, A.N., Moffitt, J.R., Wang, S. & Zhuang, X. RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* **348**, aaa6090 (2015).

8. Lubeck, E., Coskun, A.F., Zhiyentayev, T., Ahmad, M. & Cai, L. Single-cell in situ RNA profiling by sequential hybridization. *Nat Methods* **11**, 360-361 (2014).

9. Wang, X. et al. Three-dimensional intact-tissue sequencing of single-cell transcriptional states. *Science* **361** (2018).

10. Eng, C.L. et al. Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH. *Nature* **568**, 235-239 (2019).

11. Vickovic, S. et al. High-definition spatial transcriptomics for in situ tissue profiling. *Nat Methods* **16**, 987-990 (2019).

12. Stickels, R.R. et al. Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nat Biotechnol* **39**, 313-319 (2021).

13. Rodriques, S.G. et al. Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. *Science* **363**, 1463-1467 (2019).

14. Stahl, P.L. et al. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78-82 (2016).

15. Nichterwitz, S. et al. Laser capture microscopy coupled with Smart-seq2 for precise spatial transcriptomic profiling. *Nat Commun* **7**, 12139 (2016).

16. Chen, J. et al. Spatial transcriptomic analysis of cryosectioned tissue samples with Geo-seq. *Nat Protoc* **12**, 566-580 (2017).

17. Cheng, J., Liao, J., Shao, X., Lu, X. & Fan, X. Multiplexing Methods for Simultaneous Large-Scale Transcriptomic Profiling of Samples at Single-Cell Resolution. *Adv Sci (Weinh)* **8**, e2101229 (2021).

18. Liu, Y., Ai, N., Liao, J. & Fan, X. Transcriptomics: a sword to cut the Gordian knot of traditional Chinese medicine. *Biomark Med* **9**, 1201-1213 (2015).

19. Djebali, S. et al. Landscape of transcription in human cells. *Nature* **489**, 101-108 (2012).

20. Cancer Genome Atlas Research, N. Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature* **455**, 1061-1068 (2008).

21. International Cancer Genome, C. et al. International network of cancer genome projects. *Nature* **464**, 993-998 (2010).

22. Consortium, S.M.-I. A comprehensive assessment of RNA-seq accuracy, reproducibility and information content by the Sequencing Quality Control Consortium. *Nat Biotechnol* **32**, 903-914 (2014).

23. Frishberg, A. et al. Cell composition analysis of bulk genomics using single-cell data. *Nat Methods* **16**, 327-332 (2019).

24. Newman, A.M. et al. Robust enumeration of cell subsets from tissue expression profiles. *Nat Methods* **12**, 453-457 (2015).

25. Wang, X., Park, J., Susztak, K., Zhang, N.R. & Li, M. Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. *Nat Commun* **10**, 380 (2019).

26. Chen, Z. et al. Inference of immune cell composition on the expression profiles of mouse tissue. *Sci Rep* **7**, 40508 (2017).

27. Gangalum, R.K. et al. Spatial Analysis of Single Fiber Cells of the Developing Ocular Lens Reveals Regulated Heterogeneity of Gene Expression. *iScience* **10**, 66-79 (2018).

28. Ma, A., McDermaid, A., Xu, J., Chang, Y. & Ma, Q. Integrative Methods and Practical Challenges for Single-Cell Multi-omics. *Trends Biotechnol* **38**, 1007-1022 (2020).

29. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018).

30. Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019).

31. Shao, X. et al. scCATCH: Automatic Annotation on Cell Types of Clusters from Single-Cell RNA Sequencing Data. *iScience* **23**, 100882 (2020).

32. Shao, X. et al. scDeepSort: a pre-trained cell-type annotation method for single-cell transcriptomics using deep learning with a weighted graph neural network. *Nucleic Acids Res* **49**, e122 (2021).

33. Higgins, I. et al. beta-vae: Learning basic visual concepts with a constrained variational framework. *International Conference on Learning Representations*, 1-22 (2017).

34. Cable, D.M. et al. Robust decomposition of cell type mixtures in spatial transcriptomics. *Nat Biotechnol* **40**, 517-526 (2022).

35. Dong, R. & Yuan, G.C. SpatialDWLS: accurate deconvolution of spatial transcriptomic data. *Genome Biol* **22**, 145 (2021).

36. Andersson, A. et al. Single-cell and spatial transcriptomics enables probabilistic inference of cell type
topography. *Commun Biol* **3**, 565 (2020).

37. Elosua-Bayes, M., Nieto, P., Mereu, E., Gut, I. & Heyn, H. SPOTlight: seeded NMF regression to
deconvolute spatial transcriptomics spots with single-cell transcriptomes. *Nucleic Acids Res* **49**, e50
(2021).

38. Li, H., Wei, W. & Xu, H. Drug discovery is an eternal challenge for the biomedical sciences. *Acta
Materia Medica* **1**, 1-3 (2022).

39. Goodfellow, I.J. et al. Generative Adversarial Networks. *arXiv*, 1-9. (2014).
https://doi.org/10.48550/arXiv.1406.2661.

40. Mirza, M. & Osindero, S.J.A. Conditional Generative Adversarial Nets. arXiv, 1-7. (2014).
https://doi.org/10.48550/arXiv.1411.1784.

41. Wang, J., Roeder, K. & Devlin, B. Bayesian estimation of cell type-specific gene expression with prior
derived from single-cell data. *Genome Res* **31**, 1807-1818 (2021).

42. Xiong, X. et al. Landscape of Intercellular Crosstalk in Healthy and NASH Liver Revealed by Single-
Cell Secretome Gene Analysis. *Mol Cell* **75**, 644-660 e645 (2019).

43. Moffitt, J.R. et al. Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic
region. *Science* **362** (2018).

44. Chen, R., Wu, X., Jiang, L. & Zhang, Y. Single-Cell RNA-Seq Reveals Hypothalamic Cell Diversity. *Cell
Rep* **18**, 3227-3241 (2017).

45. Saunders, A. et al. Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain.
Cell **174**, 1015-1030 e1016 (2018).

46. Gierahn, T.M. et al. Seq-Well: portable, low-cost RNA sequencing of single cells at high throughput.
Nat Methods **14**, 395-398 (2017).

47. Moncada, R. et al. Integrating microarray-based spatial transcriptomics and single-cell RNA-seq
reveals tissue architecture in pancreatic ductal adenocarcinomas. *Nat Biotechnol* **38**, 333-342 (2020).

48. Tirosh, I. et al. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-
seq. *Science* **352**, 189-196 (2016).

49. Thrane, K., Eriksson, H., Maaskola, J., Hansson, J. & Lundeberg, J. Spatially Resolved Transcriptomics
Enables Dissection of Genetic Heterogeneity in Stage III Cutaneous Malignant Melanoma. *Cancer
Res* **78**, 5970-5979 (2018).

50. Wu, H. et al. Identification and functional analysis of novel oncogene DDX60L in pancreatic ductal
adenocarcinoma. *BMC Genomics* **22**, 833 (2021).

51. Macosko, E.Z. et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using
Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).

52. Tasic, B. et al. Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat
Neurosci* **19**, 335-346 (2016).

53. Elinav, E. et al. Inflammation-induced cancer: crosstalk between tumours, immune cells and
microorganisms. *Nat Rev Cancer* **13**, 759-771 (2013).

54. Wu, S.Z. et al. Cryopreservation of human cancers conserves tumour heterogeneity for single-cell
multi-omics analysis. *Genome Med* **13**, 81 (2021).

55. Berglund, E. et al. Spatial maps of prostate cancer transcriptomes reveal an unexplored landscape
of heterogeneity. *Nat Commun* **9**, 2419 (2018).

56. Shao, X., Lu, X., Liao, J., Chen, H. & Fan, X. New avenues for systematically inferring cell-cell
communication: through single-cell transcriptomics data. *Protein Cell* **11**, 866-880 (2020).

57. Shao, X. et al. CellTalkDB: a manually curated database of ligand-receptor interactions in humans
and mice. *Brief Bioinform* **22** (2021).

58. Shao, X. et al. Knowledge-graph-based cell-cell communication inference for spatially resolved
transcriptomic data with SpaTalk. *Nat Commun* **13**, 4429 (2022).

59. Shao, L. et al. Identify differential genes and cell subclusters from time-series scRNA-seq data using
scTITANS. *Comput Struct Biotechnol J* **19**, 4132-4141 (2021).

60. Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529 (2021).

61. Fan, X. et al. Spatial transcriptomic survey of human embryonic cerebral cortex by single-cell RNA-
seq analysis. *Cell Res* **28**, 730-745 (2018).

62. Lake, B.B. et al. A single-nucleus RNA-sequencing pipeline to decipher the molecular anatomy and
pathophysiology of human kidneys. *Nat Commun* **10**, 2832 (2019).

63. Aizarani, N. et al. A human liver cell atlas reveals heterogeneity and epithelial progenitors. *Nature*
572, 199-204 (2019).

64. Vieira Braga, F.A. et al. A cellular census of human lungs identifies novel cell states in health and in
asthma. *Nat Med* **25**, 1153-1163 (2019).

65. Zeisel, A. et al. Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-
cell RNA-seq. *Science* **347**, 1138-1142 (2015).

66. Wu, H., Kirita, Y., Donnelly, E.L. & Humphreys, B.D. Advantages of Single-Nucleus over Single-Cell
RNA Sequencing of Adult Kidney: Rare Cell Types and Novel Cell States Revealed in Fibrosis. *J Am Soc*
Nephrol **30**, 23-32 (2019).

67. Zilionis, R. et al. Single-Cell Transcriptomics of Human and Mouse Lung Cancers Reveals Conserved
Myeloid Populations across Individuals and Species. *Immunity* **50**, 1317-1334 e1310 (2019).

68. Baron, M. et al. A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter-
and Intra-cell Population Structure. *Cell Syst* **3**, 346-360 e344 (2016).

69. Green, C.D. et al. A Comprehensive Roadmap of Murine Spermatogenesis Defined by Single-Cell
RNA-Seq. *Dev Cell* **46**, 651-667 e610 (2018).

70. Grun, D. et al. De Novo Prediction of Stem Cell Identity using Single-Cell Transcriptome Data. *Cell*
Stem Cell **19**, 266-277 (2016).

71. Muraro, M.J. et al. A Single-Cell Transcriptome Atlas of the Human Pancreas. *Cell Syst* **3**, 385-394
e383 (2016).

72. Lawlor, N. et al. Single-cell transcriptomes identify human islet cell signatures and reveal cell-type-
specific expression changes in type 2 diabetes. *Genome Res* **27**, 208-222 (2017).

73. Segerstolpe, A. et al. Single-Cell Transcriptome Profiling of Human Pancreatic Islets in Health and
Type 2 Diabetes. *Cell Metab* **24**, 593-607 (2016).

74. Xiong, X. et al. Mapping the molecular signatures of diet-induced NASH and its regulation by the
hepatokine Tsukushi. *Mol Metab* **20**, 128-137 (2019).

75. Liao, J. et al. De novo analysis of bulk RNA-seq data at spatially resolved single-cell resolution.
ZJUFanLab/bulk2space: Bulk2Space v1.0.0 (v1.0.0), doi: 10.5281/zenodo.7134575 (2022).

1321

Acknowledgments

This work is supported by the National Natural Science Foundation of China (81973701, X.F.; 91846204, H.C.; and U19B2027, H.C.), the Natural Science Foundation of Zhejiang Province (LZ20H290002, X.F.), the Innovation Team and Talents Cultivation Program of the National Administration of Traditional Chinese Medicine (ZYYCXTD-D-202002, X.F.; ZYYCXTD-D-202207, Y.G.), the Westlake Laboratory (Westlake Laboratory of Life Sciences and Biomedicine), and the Alibaba Cloud (Alibaba Cloud Computing Co. Ltd.).

Author Contributions Statement

X.F., H.C., Y.G., and J.L. conceived the study. J.L. and J.Q. collected datasets involved in this article, benchmarked all methods, and participated in the development of the Bulk2Space algorithm. Y.F., Z.C., and X.Z. implemented the code and interface to the Bulk2Space algorithm. N.Z., X.S., H.Y., P.Y., J.C., L.S., and J.Z. provided a lot of advice on algorithm implementation and biological applications. J.L., Y.H. (Yang Hu), Y.H.(Yining Hu), L.Y., and X.L. conducted the multiplexed RNA-seq experiment using Spatial-seq. D.W. provided important advice on brain tissue isolation, spatial registration, and annotation. All authors wrote the manuscript, and read and approved the final manuscript.

Competing Interests Statement

The authors declare no competing interests.

Figure Legends

Fig. 1 Workflow of Bulk2Space. a, Overview of the design concept of Bulk2Space. Bulk transcriptome data is taken as the input, and a single-cell profile is used as the reference for characterizing the clustering space of the heterogeneous tissue. After deconvolution, input bulk data is deconvolved into single-cell transcriptomics data. Then, either of the two spatially resolved transcriptomics, spatial barcoding-based RNA-seq or image-based in-situ hybridization, is used as the spatial reference. Generated single cells are assigned to the corresponding coordinates based on the spatial reference. The output is a set of generated single-cell profiles with specified x and y spatial coordinates. **b,** Detailed deconvolution procedure. The input vector of bulk tissue is equal to the production of the expression matrix of cell types and the proportion vector of each cell type. The calculated proportion of all cell types is employed for the subsequent single-cell generation. The single-cell reference is used to characterize the clustering space of the tissue, and a deep learning model generates single-cell profiles within the clustering space of each cell type. **c** and **d** demonstrate the strategies for spatial mapping based on the two mainly used spatially resolved transcriptomics approaches. **c,** For spatial barcoding-based reference, each generated single cell is assigned to the spot with the highest gene expression correlation until the aggregation of cells within the spot is close enough to the exact expression value. **d,** For image-based reference, each generated cell is assigned to the location where the cell on tissue has the highest similarity with the given cell.

Fig. 2. Benchmark test for Bulk2Space. a, Gene expression correlation (Pearson correlations, left, and Spearman correlations, middle) and gene expression variation (RMSE, right) of three methods, namely, Bulk2Space (β-VAE), GAN, and CGAN, using paired simulation data (n=30). Each point represented a simulated dataset. Data are presented as boxplots (minima, 25th percentile, median, 75th percentile, and maxima). **b**, Gene expression correlation and gene expression variation of three methods with unpaired simulation data (n=12). Data are presented as boxplots (minima, 25th percentile, median, 75th percentile, and maxima). **c**, Benchmark test for 10 spatial mapping methods (from left to right, Bulk2Space (deep forest), gradient boosting decision tree, decision tree, multi-layer perceptron, DeepGBM, linear regression, RCTD, spatialDWLS, stereoscope, and SPOTlight) using paired simulation data (n=50). Data are presented as boxplots (minima, 25th percentile, median, 75th percentile, and maxima). Left, Pearson correlation of gene expression, Middle, Spearman correlation of gene expression, Right, gene expression variation (RMSE). **d**, Benchmark test for 10 spatial mapping methods using paired simulation data (n=60). Data are presented as

boxplots (minima, 25th percentile, median, 75th percentile, and maxima). **e**, The UMAP layout of the MERFISH data (image-based spatially resolved transcriptomics) and the single-cell data. Share cell types were colored with corresponding colors. **f**, Expression correlation heatmap of the marker genes of all shared cell types in MERFISH data and scRNA-seq data. **g**, Spatial expression of genes in the ground truth and predicted by Bulk2Space using paired and unpaired single-cell RNA-seq data. PCC for each gene was shown. Source data are provided as a Source Data file.

Fig. 3. Spatially resolved analysis of PDAC by Bulk2Space. a, Collection of pancreatic adjacent (PA) and PDAC bulk transcriptome data, PDAC scRNA-seq data, and PDAC spatial transcriptomics data. Two bulk data were sequenced by bulk RNA-seq. The scRNA-seq data sequenced by inDrop-seq[51] is used as the single-cell reference. One slice of the sectioned PDAC tissue sequenced by ST[14] was employed as the spatial reference. **b**, The cell-type proportions of single cells generated from PDAC and PA tissues by Bulk2Space. Purple box, the proportion of the cell type was higher in PA tissue than that in PDAC tissue. Black box, the proportion of the cell type was higher in PDAC tissue than that in PA tissue. Teal, the PDAC tissue. Red, PA tissue. **c**, The clustering space of single cells generated from the PDAC (Top) and PA (Bottom) bulk data by Bulk2Space using UMAP layout. Different colors represented distinct cell types. **d**, Pairwise expression correlation of cell-type-specific marker genes between single cells generated by Bulk2Space and the single-cell reference for PDAC. Marker genes were found by 'FindAllMarkers' function in Seurat. *P* value was calculated with the Wilcoxon Rank Sum test. **e**, Spatial mapping of single cells generated from PDAC bulk data. Scale bar, 1 mm. **f**, Top, Histological annotation for cancer region (red), duct epithelium (yellow), and normal pancreatic tissue (light blue). Middle, clusters of cancer region (yellow), pancreatic tissue (blue), duct epithelium (green), and stroma (dark grey) from the spatial transcriptomics data. Bottom, spatial deconvolution result of the bulk PDAC data by Bulk2Space with each spot displaying the composition of cell types in a pie chart. Scale bar, 1 mm. **g**, The spatial abundance of acinar cells, cancer clone A cells, cancer clone B cells, and ductal cells in each spot on the tissue section predicted by Bulk2Space. Scale bar, 1 mm. **h**, The spatial expression of the marker genes in acinar cells, cancer clone A cells, cancer clone B cells, and ductal cells predicted by Bulk2Space at spot (left) and cellular (right) resolution. Source data are provided as a Source Data file.

Fig. 4. Spatially resolved single-cell analysis of the mouse isocortex bulk data by Bulk2Space. a, The resources of the bulk transcriptome data, single-cell reference, and spatial reference. The bulk RNA-seq data of the mouse isocortex was sequenced using our in-house developed Spatial-seq which combines LCM and spatial barcoding strategies. The single-cell reference data is derived from scRNA-seq of a collection of consecutive mouse primary visual cortex sections from anterior to posterior. The spatial reference data is obtained from an open-access database using the 10X Visium approach. **b**, The deconvolution results of the mouse isocortex bulk data by Bulk2Space. The UMAP layout showed the clustering space of the generated single cells. **c**, The spatial mapping of the single cells generated from the mouse isocortex bulk data by Bulk2Space. Scale bar, 1 mm. **d**, The pairwise expression correlation of cell-type-specific marker genes between single cells generated by Bulk2Space and single cells in the reference data. **e**, The spatial distribution of the cell-type proportion predicted by Bulk2Space. Scale bar, 1 mm. **f**, The spatial cell-type abundance of seven cell types with layered structure predicted by Bulk2Space. Scale bar, 1 mm. **g**, The spatial expression distribution of cell-type-specific marker genes predicted by Bulk2Space at single-cell resolution. Scale bar, 1 mm. Source data are provided as a Source Data file.

Fig. 5. Spatially resolved analysis of the mouse hypothalamus bulk data by Bulk2Space. a, The resources of the bulk transcriptome data, single-cell reference, and spatial reference. The bulk RNA-seq data of the mouse hypothalamus was profiled using our in-house developed Spatial-seq. The single-cell reference data was derived from Drop-seq. The spatial reference data is obtained from MERFISH. **b**, The UMAP layout of single-cell profiles in MERFISH data and generated from bulk tissue by Bulk2Space. Top, clustering of all cells in both datasets. Two heterogeneous sets of data were aligned through a joint analysis. Each cell type was represented by a unique color. **c**, The spatial distribution of distinct cells in MERFISH and Bulk2Space results. The stacked bar chart showed the cell number of each cell type (grey) and the number of cells predicted by Bulk2Space to the corresponding cell type at each coordinate (blue). Bottom, Pearson correlation of gene expression between cell types in MERFISH data and Bulk2Space results. The correlation of shared cell types was in red border and the expression correlation between MERFISH ambiguous cells and Bulk2Space predicted cell types was in green border. **d**, The spatial expression of genes predicted by Bulk2Space and in the ground truth (MERFISH). **e**, Re-annotation of ambiguous cells in MERFISH using Bulk2Space results. Left, clustering of the ambiguous cells. Right, spatial assignment of cells that were colored according to their predicted cell types. **f**, Predicted spatial expression patterns of novel marker genes for different cell types. Source data are provided as a Source Data file.

Fig. 6. Spatial gene expression dynamics analysis of inflammation-induced prostate

cancer by Bulk2Space. a, Experimental design of the bulk transcriptome data. The

tissue was segmented into distinct regions, namely, normal glands (pink),

inflammation (blue), and several cancer regions (red, green, purple, and yellow). Three

tissue regions (1.1 normal, 2.3 inflammation, and 3.3 cancer region) were isolated to

conduct spatially resolved transcriptomics. **b**, The t-SNE layout of the generated single-

cell profiles by Bulk2Space. Each cell type was represented by a unique color. **c**, The

proportions of generated single cells that were mapped to normal, inflammation, and

cancer region by Bulk2Space, respectively. The color of each cell type was consistent

with **b**. **d**, The proportion of cells assigned to normal (pink), inflammation (blue), and

cancer (green) regions for each cell type. **e**, The predicted proportion of distinct cell

types for each tissue region by Bulk2Space. **f**, The proportions of cancer-associated

fibroblasts (CAFs) in normal (left), inflammation (middle), and cancer regions (right). **g**,

The proportions of immune cells in normal, inflammation, and cancer regions. **h**, The

proportions of cancer epithelial cells in normal, inflammation, and cancer regions.

Source data are provided as a Source Data file.

a

Input

Bulk RNA-seq

Cell types

Genes

Deconvolution

Single-cell reference

Generated

Cells

Genes

Single cells

(1) Spatial barcoding

Cell

Spots

80% 15% 4%

Spatial mapping

(2) Image-based

Cell

Cells

✗ ✗ ✓ ✗ ✗

Full-gene coverage

Single-cell resolution

y

Coordinates

x

Output

Cells

x
y

Genes

Spatially resolved
transcriptomics

Deconvolution (Step 1)

Spatial mapping (Step 2)

b Generate single cells within the clustering space from bulk data

Input

Bulk

Genes

Single-cell reference

Cell types

Genes

×

Prediction

Abundance

Cell types	
5%	
40%	
10%	
45%	

The clustering space of each cell type

C2 C1
C3 C4

β-VAE architecture

Encoder   Mean   Decoder

Single-cells

Variance

Single-cells

Generated SC data

40% 5%
C2   C1
C3   C4
10% 45%

c Map generated single cells to reference based on spatial barcoding

Generated
SC data

Spot
assignment

80% 10%
5%   4%

Spot
deconvolution

Output
single-cell resolution

d Map generated single cells to image-based reference

Generated
SC data

Similarity
calculation

Cell
assignment

Output
Full-gene coverage

a Paired simulation (bulk and reference were generated from same SC datasets)

Gene expression correlation (Pearson)

Gene expression correlation (Spearman)

Gene expression variation (RMSE)

b Unpaired simulation (bulk and reference were from different SC datasets)

Gene expression correlation (Pearson)

Gene expression correlation (Spearman)

Gene expression variation (RMSE)

c Paired simulation (single-cell and spatial transcriptomics data were generated from same datasets)

Gene expression correlation (Pearson)

Gene expression correlation (Spearman)

Gene expression variation (RMSE)

d Unpaired simulation (single-cell and spatial transcriptomics data were generated from different datasets)

Gene expression correlation (Pearson)

Gene expression correlation (Spearman)

Gene expression variation (RMSE)

- ● Bulk2Space (deep forest)
- ● Gradient boosting decision tree
- ● Decision tree
- ● Multi-layer perceptron
- ● DeepGBM
- ● Linear regression
- ● RCTD
- ● spatialDWLS
- ● stereoscope
- ● SPOTlight

e Shared clustering space of MERFISH and scRNA-seq data

MERFISH

scRNA-seq

UMAP2

UMAP2

UMAP1

UMAP1

OD immature
Mural/Pericyte
Endothelial
Astrocyte
Microglia
OD mature
Excitatory
Inhibitory

f The gene expression correlation between scRNA-seq and MERFISH

MERFISH

scRNA-seq

Astrocyte
Endothelial
Ependymal
Excitatory
Inhibitory
Microglia
OD Immature
OD Mature
Mural/Pericyte

Astrocyte, Endothelial, Ependymal, Excitatory, OD Immature, Inhibitory, Microglia, OD Mature, Mural/Pericyte

g Spatial distribution of the gene expression in MERFISH and Bulk2Space prediction

Sgk1 *Selplg* *Ttyh2* *Ndrg1* *Gad1*

MERFISH

Single-cell and MERFISH data were collected from the same experiment

PCC = 0.981 PCC = 0.961 PCC = 0.960 PCC = 0.958 PCC = 0.945

Bulk2Space

Single-cell and MERFISH data were collected from separate experiments

PCC = 0.958 PCC = 0.932 PCC = 0.936 PCC = 0.943 PCC = 0.941

Bulk2Space

a Collection of bulk RNA-seq data, single-cell (SC) RNA-seq data, and spatially resolved transcriptomics (ST) data

Pancreatic adjacent tissue (PA tissue)

Pancreatic ductal adenocarcinoma (PDAC tissue)

Separated experiments

Bulk RNA-seq

PA tissue   PDAC tissue

Genes   Genes

Bulk transcriptomes
GSE171485

inDrop-seq

Cell

Barcodes   RT mix

Oil   Oil

SC reference

ST Technology

Slide

Spatial barcoding

ST reference
GSE111672

b Comparison of the cell-type proportions generated from PDAC and PA tissues

Proportion

PDAC tissue
PA tissue

Acinar cells
Fibroblasts
mDCs
Endocrine cells
Endothelial cells
Cancer clone B
Cancer clone A
Ductal
Mast cells
pDCs
Monocytes
Macrophages

Less than 5%

More abundant in PA tissue
More abundant in PDAC tissue
PDAC tissue
PA tissue

c Deconvolution of bulk transcriptomics from PDAC and PA tissues by Bulk2Space

PDAC tissue

UMAP2

UMAP1

PA tissue

UMAP2

UMAP1

• Acinar cells
• Cancer clone A
• Cancer clone B
• Ductal
• Endocrine cells
• Endothelial cells
• Fibroblasts
• Mast cells
• mDCs
• pDCs
• RBCs

d Expression correlation of marker genes between SC reference and deconvolved PDAC bulk data

log(Fold Change) > 0.5, P value<0.05

Bulk2Space

single-cell reference

Acinar cells
Cancer clone A
Cancer clone B
Ductal
Endocrine cells
Endothelial cells
Fibroblasts
Macrophages
Mast cells
mDCs
Monocytes
pDCs
RBCs

Acinar cells
Cancer clone A
Cancer clone B
Ductal
Endocrine cells
Endothelial cells
Fibroblasts
Macrophages
Mast cells
mDCs
Monocytes
pDCs
RBCs

1
0.75
0.5
0.25
0

e Spatial mapping of generated cells from the PDAC tissue by Bulk2Space

y coord

x coord

• Acinar cells
• Cancer clone A
• Cancer clone B
• Ductal
• Endocrine cells
• Fibroblasts
• Macrophages
• mDCs
• Monocytes
• pDCs
• RBCs

f Mapping the histological annotations and spatial cluster assignments with the spatial deconvolution results from PDAC tissue by Bulk2Space

Histological annotations

— Cancer cells and desmoplasia
— Duct epithelium
— Normal pancreatic tissue

Cancer region
Pancreatic tissue
Duct epithelium
Stroma

Cluster assignments

Acinar cells
Cancer clone A
Cancer clone B
Ductal cells
Endothelial cells
Fibroblasts
Macrophages
mDCs
Monocytes
pDCs
RBCs

Bulk2Space result

g Bulk2Space predicted cell-type proportions in spatial distribution

Acinar cells   Cancer clone A

Ductal cells   Cancer clone B

Proportion
1.00
0.75
0.50
0.25
0.00

Scale bar

h Spatial expression of cell-type-specific maker genes at spot and single-cell resolution

low   high

Acinar cells   Cancer clone A   Cancer clone B   Ductal cells

Spots Cells   Spots Cells   Spots Cells   Spots Cells

CPA1   TM4SF1   SLC20A1   CRISP3

CTRB1   GABRP   KRT16   DMBT1

REG1A   CLDN1   KRT17   SPP1

a Bulk RNA-seq data of Mouse isocortex are obtained by Spatial-seq

Bulk data

Genes

Decoding

Isocortex (Barcode X,Y,Z)

Isolation of bulk tissue

LCM

Collection

Spatial barcoding

RNA-seq

Barcodes   mRNA

scRNA-seq data of the mouse primary visual cortex by SMART-seq2

Microdissection

Posterior

Anterior

SMART-seq2

Single-cell reference

Spatial transcriptomics of the mouse isocortex region by 10X Visium

mRNA

Barcodes

Spatial reference

b Deconvolution of mouse isocortex bulk RNA-seq data by Bulk2Space

L5 PT   L5 IT   L2/3 IT
L6 CT   L6 IT   Astro
Sst   NP   Lamp5
Pvalb   Vip

UMAP2 / UMAP1

Cell types

Astro, L2/3 IT, L4, L5 IT, L5 PT, L6 CT, L6 IT, Lamp5, NP, Pvalb, Sst, Vip

c Spatial mapping of generated single cells by Bulk2Space

Anterior

Posterior

Astro, L2/3 IT, L4, L5 IT, L5 PT, L6 CT, L6 IT, Lamp5, NP, Pvalb, Sst, Vip

d Expression correlation of marker genes between Bulk2Space and SC reference

Bulk2Space / scRNA-seq

Astro, L2/3 IT, L4, L5 IT, L5 PT, L6 CT, L6 IT, Lamp5, NP, Pvalb, Sst, Vip

e Spatial distribution of the predicted cell-type compositions in the anterior and posterior of the mouse isocortex region by Bulk2Space

Astro, L2/3 IT, L4, L5 IT, L5 PT, L6 CT, L6 IT, Lamp5, NP, Pvalb, Sst, Vip

f Spatial distribution of predicted cell-type proportions in the anterior and posterior of the mouse isocortex region by Bulk2Space

Astro, L2/3 IT, L4, L5 IT, L5 PT, L6 CT, L6 IT

g Spatial distribution of Bulk2Space predicted expression of cell-type-specific marker genes at single-cell resolution

Astro (*Osr1*)   L2/3 IT (*Myh7*)

L4 (*Rspo1*)   L5 IT (*Dkkl1*)

L5 PT (*Fam84b*)   L6 CT (*Foxp2*)

L6 IT (*Osr1*)

a Bulk RNA-seq data of mouse hypothalamus region are obtained by Spatial-seq

Bulk data

Hypothalamus (Barcode X,Y,Z)

Isolation of bulk tissue

LCM

Collection

Spatial barcoding

Decoding

Barcodes   mRNA

RNA-seq

scRNA-seq data of the mouse hypothalamus are derived by Drop-seq

Barcoded beads

Cells

Single-cell reference (GSE113576)

Spatially resolved transcriptomics of the mouse hypothalamic preoptic region by MERFISH

Spatial reference

b Merged clustering

- Astrocytes
- Endothelial cells
- Ependymal cells
- Excitatory neurons
- Inhibitory neurons
- Microglias
- Mural/Pericytes
- Oligodendrocytes (OD) Mature
- OD Immature
- Macrophages

Bulk2Space   MERFISH

c Spatial mapping of generated single cells

MERFISH

- Astrocytes
- Endothelial cells
- Ependymal cells
- Excitatory neurons
- Inhibitory neurons
- Microglias
- Mural/Pericytes
- OD Mature
- OD Immature
- Macrophages

Scale bar

Bulk2Space

No. of matched cells

Gene expression correlation between cell types

MERFISH

Bulk2Space

Astrocytes
Endothelial cells
Ependymal cells
Excitatory neurons
Inhibitory neurons
Microglias
Mural/Pericytes
OD Mature

Ambiguous

d Prediction of the spatial gene expression

Bulk2Space   MERFISH

Syt4

Lpar1

Prlr

Necab1

ermn

low   high

e Prediction of the spatial gene expression

Clustering Space

Inhibitory neurons
Excitatory neurons
Mural/Pericytes
Macrophages
Endothelial cells
OD Mature
Astrocytes

Spatial distribution

Coord y

Coord x

f Spatial distribution of novel marker genes predicted by Bulk2Space

Astrocytes *Ntsr2*
Endothelial cells *Cldn5*
Ependymal cells *S100a6*

Excitatory neurons *Resp18*
Inhibitory neurons *Syt1*
Microglias *Ctss*

Mural/Pericytes *Tagln*
OD Mature *Mag*
Macrophages *Lyz2*

low   high