Spatially resolved single-cell deconvolution of bulk RNA-seq using Bulk2Space

Jie Liao[1,4,†], Jingyang Qian[1,4,†], Yin Fang[2,3,†], Zhuo Chen[2,3,†], Xiang Zhuang[2,3,†], Ningyu Zhang[2,3], Xin Shao[1,4], Haihong Yang[2,3], Penghui Yang[1,4], Junyun Cheng[1,4], Yang Hu[1,4], Yining Hu[1], Lingqi Yu[1,4], Jinlu Zhang[1,2], Xiaoyan Lu[1,4], Li Shao[8], Dan Wu[9], Yue Gao[10,*], Huajun Chen[2,3,6,7,*], Xiaohui Fan[1,4,5*]

[1]College of Pharmaceutical Sciences, Zhejiang University, Hangzhou 310058, China

[2]College of Computer Science and Technology, Zhejiang University, Hangzhou 310027, China

[3]Hangzhou Innovation Center, Zhejiang University, Hangzhou 310058, China

[4]Innovation Center in Zhejiang University, State Key Laboratory of Component-Based Chinese Medicine, Hangzhou 310058, China

[5]Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou 310058, China

[6]Alibaba-ZJU Frontier Technology Research Center, Hangzhou 310023, China

[7]The First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou 310003, China

[8]Institute of Translational Medicine, The Affiliated Hospital of Hangzhou Normal University, Hangzhou Normal University, Hangzhou 310015, China

[9]College of Biomedical Engineering and Instrument Science, Zhejiang University, Hangzhou 310013, China

[10]Department of Pharmaceutical Sciences, Beijing Institute of Radiation Medicine, Beijing, 100852, China.

*Corresponding Authors: Prof. Xiaohui Fan, e-mail: fanxh@zju.edu.cn, Prof. Huajun Chen, e-mail: huajunsir@zju.edu.cn, and Prof. Yue Gao, e-mail: gaoyue@bmi.ac.cn.

[†]These authors contribute equally to this work.

Abstract

Uncovering the tissue molecular architecture at single-cell resolution could help better understand the biological and pathological processes in organisms. However, bulk RNA-seq can only measure gene expression in cell mixtures, without revealing molecular heterogeneity and spatial patterns of single cells. Herein, we introduce Bulk2Space (https://github.com/ZJUFanLab/bulk2space), a deep learning framework-based spatial deconvolution algorithm that can simultaneously reveal the spatial and cellular heterogeneity of bulk RNA-seq data. The use of bulk transcriptomics to validate Bulk2Space revealed, in particular, the spatial variance of immune cells in different tumor regions, the molecular and spatial heterogeneity of tissues during inflammation-induced tumorigenesis, and spatial patterns of novel genes in different cell types. Moreover, Bulk2Space further annotated cell types that cannot be identified by original methods when it was used to spatially deconvolve the bulk data derived from our in-house developed sequencing approach Spatial-seq. Bulk2Space is distributed via an open-source Python software package.

Keywords

RNA-seq, Spatially resolved transcriptomics, Spatial deconvolution, Spatial-seq, scRNA-seq, Deep learning.

Main

Tissue complexity is portrayed by the spatial diversity and heterogeneity of cells[1]. Advances in spatially resolved transcriptomics[2-4] have made it possible to understand the cell composition, molecular architecture, and functional details of tissues at unanticipated spatial levels[5, 6]. State-of-the-art experimental technologies, including image-based methods[7-10], spatial barcoding RNA-seq methods[11-14], and laser capture microdissection-based methods[15, 16], have been developed to address either high throuput measuring of cells, unbiased detection of mRNA species, or single-cell resolution. To investigate the molecular variation during biological and pathological processes at a higher resolution, each sample is encouraged to be analyzed by spatially resolved single-cell transcriptomics, which is not yet fully achieved and is time-consuming, costly, and difficult to scale up[17]. Meanwhile, with investment for almost two decades, RNA-seq has been extensively applied in transcriptome analysis[18], with many large projects having been carried out, such as the Encyclopedia of DNA Elements (ENCODE)[19], The Cancer Genome Atlas (TCGA)[20], and projects of the International Cancer Genome Consortium (ICGC)[21]. A wealth of bulk RNA-seq data has become a legacy for biological and clinical research[22]. Thus, reanalysis of the enormous amount of bulk data to explain both cellular diversity and spatial expression patterns is a challenging but consequential task.

The *in silico* method has great potential to predict spatial heterogeneity from bulk RNA-seq data at single-cell resolution by integrating cutting-edge technologies. Several approaches such as CPM[23], CIBERSORT[24], and MuSiC[25] stopped short of extrapolating the proportion of cell types from bulk RNA-seq data yet failed to obtain single-cell data, let alone map them to tissue coordinates. The emerging of an approach that can efficiently convert bulk RNA-seq data into spatially resolved single-cell expression profiles is expected to reveal the cell diversity of complex tissue and the spatial expression variation simultaneously.

Herein, we introduce Bulk2Space, a spatial deconvolution method based on deep learning frameworks, which converts bulk transcriptomes into spatially resolved single-cell expression profiles using existing high-quality scRNA-seq data and spatial transcriptomics as references. We hypothesize that the process of scRNA-seq is similar to sampling cells from bulk tissue, and each

selected cell is labeled with a unique barcode. Consequently, bulk transcriptome data can be used

as a weighted collection of single-cell expression data in a defined clustering space of cells.

Bulk2Space first generates single-cell transcriptomic data within the clustering space to find a set

of cells whose aggregated data are close to the bulk data. Next, the generated single cells were

allocated to optimal spatial locations using a spatial transcriptome reference. For this step, we

chose as a spatial reference one of the two most commonly used spatially resolved transcriptomics

technologies. One is an image-based method with limited target genes[7-9], and the other is a spatial

barcoding method without single-cell resolution[11-14]. Bulk2Space showed a robust performance

across multiple datasets and conditions and is an open-access algorithm in GitHub

(https://github.com/ZJUFanLab/bulk2space).

Results

Design of Bulk2Space

The overall design of the Bulk2Space algorithm is illustrated in Fig. 1, which is divided into

deconvolution and spatial mapping. Although bulk data are obtained by sequencing a mixture of

cells, whereas scRNA-seq labels individual cells, both methods share comparable cell types and

states[23, 24, 26]. Meanwhile, the prevailing spatially resolved transcriptomics methods can provide

reference coordinates for single cells based on their expression profiles[27, 28]. As a consequence, it

could rationally conceive that the combination of single-cell profiles and spatial transcriptomics is

expected to overcome technical bottlenecks and accomplish the spatially resolved single-cell

deconvolution of bulk transcriptomes. We hypothesize that the process of single-cell sequencing

is the process of sampling from tissue and that the ratio between different cell types is related to

the zone of tissue, the ways of single-cell capture, and the size of the cell itself. However, even if

the ratio of cell types changes, the state of each cell still fluctuates within a relatively stable high-

dimensional space, which is namely the clustering space of cells. Several studies have backed up

this hypothesis[29-31].

Fig. 1 Workflow of Bulk2Space. (a) Overview of the design concept of Bulk2Space. Bulk transcriptome data is taken as input, and a single-cell profile is used as the reference for characterizing the clustering space of the heterogeneous tissue. After deconvolution, input bulk data is transformed into single-cell transcriptomics. Then, either of the two spatially resolved transcriptomics, spatial barcoding RNA-seq or image-based in-situ hybridization, is used as the spatial reference. Generated single cells are assigned to the corresponding coordinates based on the spatial reference. The output is a set of deconvolved single-cell profiles with specified x and y spatial coordinates. **(b)** Detailed deconvolution procedure. The input vector of bulk tissue is a production of the expression matrix of cell types and the proportion vector of each cell type. The calculated proportion of all cell types is employed for the subsequent single cells generation. The single-cell reference is used to characterize the clustering space of the tissue, and a deep learning model generates single-cell profiles within the clustering space of each cell type. **(c)** and **(d)** demonstrate the strategies for spatial mapping based on the two mainly used spatially resolved transcriptomics approaches. **(c)** For spatial barcoding reference, each generated single cell is assigned to the spot with the highest gene expression correlation until the aggregation of cells within the spot is close enough to the exact expression value. **(d)** For image-based reference, each generated cell is assigned to the location where the cell on tissue has the highest similarity with the given cell.

For single-cell deconvolution, single-cell reference profiles can be used to characterize the clustering space and the average gene expression in each cell type[31, 32]. The expression vector of bulk transcriptome is taken as an input, which is a product of the average gene expression matrix

of each cell type and its abundance vector. By solving this nonlinear equation, the proportion of

each cell type is determined. Using the percentage of each cell type as a control parameter to

generate the number of single cells, a deep learning model, termed the constrained variational

autoencoder (β-VAE)[33], is utilized to randomly simulate a given number of single cells within the

clustering space of each cell type, thus transforming the heterogeneous bulk transcriptome into

single-cell transcriptome data (Fig. 1b).

For spatial mapping, the generated single-cell profiles are assigned to the reference tissue

section map based on either of the two most commonly used spatially resolved transcriptomics

approaches, namely, spatial barcoding or image-based targeted method. As shown in Fig. 1c, for

spatial barcoding references, each barcoded spot was regarded as a mixture of several cells. We

calculated the probability of each generated single cell in each spot based on its gene expression

and assigned this cell to the most likely spot until the recommended spot was saturated. Finally,

the cell composition of each spot was determined. For image-based references, simulated single-

cell data were mapped to the coordinates of the target cell according to the similarity in their gene

expression (Fig. 1d). Instead of only calculating the abundance values of cell types, as performed

by other methods, such as Cell Population Mapping (CPM)[23], CIBERSORT[24], and ImmuCC[26],

Bulk2Space can generate biologically feasible single-cell expression profiles and assign them

robustly to tissue coordinates, thus achieving spatially resolved single-cell deconvolution of the

bulk transcriptomics.

Bulk2Space robustly deconvolves bulk transcriptomes

To demonstrate the robustness of the Bulk2Space algorithm, a benchmark test was performed

using 30 single-cell transcriptome datasets (Supplementary Table 1) obtained from different tissues.

First, each single-cell transcriptome dataset was divided into two parts, one for reference and the

other for the synthesis of bulk data. The procedure for bulk data synthesis is shown in Fig. 2a. The

composition ratio of each cell type was first assumed, and then single-cell data were randomly

selected from the given dataset according to this ratio. This newly created set of cells can be

employed as the ground truth for subsequent evaluation and is aggregated to synthesize the input

bulk transcriptome data. Three generative deep learning models, namely, the constrained

variational autoencoder (β-VAE)[33], generative adversarial networks (GAN)[34], and conditional GAN

(CGAN)[35], were introduced here for the benchmarking of the 30 single-cell datasets. The gene

expression correlation between the generated single-cell data and input bulk data was calculated

to evaluate and compare the three candidate algorithms. As illustrated in Fig. 2b, β-VAE performed

better than the other algorithms in both Pearson and Spearman correlations with a lower root

mean squared error (RMSE). Two examples are shown in Fig. 2c. The first includes single-cell data

for human peripheral blood[36], which consists of four cell types. Bulk2Space was used to deconvolve

the synthetic data, and the results demonstrated that the clustering space of the generated single-

cell data was similar to that of the test data but varied in the proportions of different cell types.

Furthermore, the correlation analysis of the expression of marker genes for each cell type showed

that the generated and test data had a correlation over 0.9. Similar results were obtained for single-

cell data collected generated from the human lung[37] (Fig. 2c) and other datasets (Supplementary

Fig. 1). To further investigate the robustness of Bulk2Space, we introduced two noise mechanisms

to test the performance of the algorithm (Supplementary Fig. 2). One mechanism changes the

expression values of certain genes in randomly selected cells, and the other alters the cell type

labels of selected cells. Bulk2space showed robust performance and could effectively avoid the

over-fitting phenomenon.

Bulk2Space maps single cells to spatial locations

In the second step, Bulk2Space assigns the generated single cells to spatial coordinates based on a

spatially resolved transcriptome reference (Fig.2 d-i). There are two most common spatially

resolved transcriptomic techniques. One is the spatial barcoding technology, which includes

'spatial transcriptomics' (ST)[14], high-definition spatial transcriptomics (HDST)[11], and Slide-seq[12, 13].

These methods are based on tissue microregions with certain regular shapes (spots), and cannot

achieve single-cell resolution. The other is image-based *in situ* transcriptomics, including MERFISH[7],

seqFISH[8, 10], and STARmap[9]. These methods can only measure the expression of target genes but

cannot cover the whole transcriptome. To overcome the bottlenecks of the two approaches, we

designed corresponding spatial assignment strategies.

The first strategy was designed to map single cells to spots in spatially resolved transcriptomics

based on spatial barcoding. In this step, we used the synthesized data to test the performance of

Bulk2Space. The data synthesis procedure is illustrated in Fig. 2d. Similar to the commonly used

spatially resolved transcriptomics technologies, such as ST, HDST, 10X Visium, and Slide-seq, we

randomly selected certain single cells from the scRNA-seq data to constitute a spot and assigned it

a defined spatial coordinate in a virtual tissue with a reference system. The gene expression value

of a spot is the sum of the gene expression values of all cells within it. A comprehensive comparison

of six machine learning or deep learning approaches was then conducted using 40 datasets. These

methods were benchmarked for the optimization of cell type composition and spatial gene

expression patterns. As shown in Fig. 2e, compared with these methods, Bulk2Space showed a

higher correlation with gene expression and a lower RMSE. The cell composition and spatial

expression patterns predicted by Bulk2Space were highly correlated to the ground truth (Fig. 2f).

A scRNA-seq data (GSE113576) and MERFISH data (image-based reference) from the same

tissue[38] were used to test the performance of the Bulk2Space algorithm for the second spatial

mapping strategy. We wanted to find the shortest path that could shift single cells to optimal

locations. First, a nonparametric empirical Bayes network was used to eliminate the expression

differences between the single-cell data and spatially resolved transcriptome data. As illustrated in

Fig. 2g, the clustering spaces of the two datasets were very similar. Most cell types were matched

in both datasets. The two discrete datasets could be well merged in distribution, and the gene

expression of each cell type had a strong correlation between the two datasets (Fig. 2h). Of the

150 target genes identified in MERFISH, 149 were used as reference genes, and one was used for

validation. The results showed that spatially, the gene expression pattern of the matched cells was

highly correlated with that of the reference (Fig. 2i), and the predicted spatial expression patterns

across all 150 targeted genes are strongly correlated with the ground truth (Supplementary Fig. 3).

Fig. 2. Benchmark test for Bulk2Space. (a) Synthesis of the bulk transcriptome data. The color bar donated the abundance of distinct cell types. For a given single-cell profile and the specified composition proportions of cell types, single cells were randomly selected from each cell type to form a new dataset. The input bulk transcriptomics was retrieved by aggregating the generated single-cell profiles, which can be used as the ground truth for the performance evaluation of various methods. **(b)** Gene expression correlation (Pearson correlations, left, and Spearman correlations, middle) and root mean squared error (RMSE, right) of three methods, namely, Bulk2Space (β-VAE), GAN, and CGAN. Each point represented a data set. **(c)** Illustration of the deconvolution examples by Bulk2Space on two datasets, human peripheral blood (GSE92495) on the top and human lung (GSE130148) on the bottom. Left, distribution of generated (yellow) and test (grey) data. Middle, t-SNE layout of the datasets. Different colors represented distinct cell types. Right, Pearson expression correlation heatmap of different cell types for generated and test data. **(d)** Synthesis of the spatially resolved transcriptome for spatial barcoding methods. Single cells were randomly distributed on the virtual tissue section. Arrayed regions were uniformly segmented from the virtual tissue at a certain diameter (regarding the spatial resolution of

different technologies) to simulate existing spatial barcoding-based RNA-seq data, e.g., ST, HDST,

slide-seq, and 10X Visium, etc. Therefore, the specific cell composition of the generated spatial

reference was known. **(e)** Benchmark test for 6 spatial mapping methods (red, Bulk2Space (deep

forest); yellow, multi-layer perceptron; green, linear regression; blue, DeepGBM; indigo, decision

tree; and purple, gradient boosting decision tree). Left, box plot of gene expression correlation and

RMSE across 40 datasets for each method. **(f)** The layout of the cell compositions that were

predicted by Bulk2Space and the ground truth. Each pie chart exhibited the cell type proportions

in each spot. **(g)** The t-SNE layout of the MERFISH data (image-based spatially resolved

transcriptomics) and the single-cell data. Share cell types were colored with corresponding colors.

(h) Expression correlation heatmap of the marker genes of all shared cell types in MERFISH data

and scRNA-seq data. **(i)** Spatial expression patterns of genes in the ground truth and predicted by

Bulk2Space.

Bulk2Space integrates spatial gene expression and histomorphology in pancreatic ductal adenocarcinoma (PDAC)

To further verify the performance of the Bulk2Space algorithm, it was applied to reveal the spatial

heterogeneity in bulk PDAC data. Moncada[39] et al. performed scRNA-seq and ST on PDAC tissues.

As shown in Fig. 3a, the scRNA-seq data were used as a deconvolution reference, and one piece of

a tissue section was used as a spatial reference (Slice 1). An additional slice (Slice 2) was used to

synthesize bulk data. After deconvolution, the bulk data were transformed into single-cell data.

There was a good correlation between the generated single-cell profiles and Slice 2 on the

expression of marker genes corresponding to each cell type (Fig. 3b). The deconvolved single cells

were then mapped to Slice 1 by Bulk2Space (Fig. 3c). To further correlate the spatial expression

pattern of the tissue with its histological feature, we investigated the molecular architecture of the

annotated regions in Slice 1 (Fig. 3d). As shown, the spatial distribution of cancer, endothelial, and

ductal cells predicted by Bulk2Space (Fig. 3e) was consistent with the histological annotation and

the spatially resolved transcriptomics of cancer, interstitium, and duct epithelium regions. This was

confirmed by the spatial expression of marker genes in these three histologically associated cell

types (Fig. 3f). The gene expression correlation of all six cell types between the predicted spatial

data and Slice 1 was exhibited in Fig. 3g.

Fig. 3. Spatially resolved analysis of PDAC by Bulk2Space. (a) Synthesis of the bulk transcriptome
data. The scRNA-seq data is used as the deconvolution reference, one slice of the sectioned tissue
is employed as the spatial reference (Slice 1), and another slice (Slice 2) is used to synthesize the
input bulk data. **(b)** The expression correlation of the marker genes in six cell types between the
Bulk2Space result and Slice 2. **(c)** Single-cell deconvolution of the bulk data. Top, t-SNE plot showed
the clustering of generated single cells and the rose chart showed the proportion of each predicted
cell type. Bottom, distribution of predicted single cells. Different colors represented distinct cell
types. **(d)** The top, Histological annotation for cancer region (red), interstitium (green), and duct
epithelium (yellow). Middle, clusters of cancer region (yellow), interstitium (blue), and duct
epithelium (green) from the ST data. Bottom, single-cell decomposition result of each spot in the
ST data by Bulk2Space. **(e)** The spatial abundance of different cell types in each spot on the tissue
section is predicted by Bulk2Space. **(f)** The spatial expression of the marker genes in cancer (left),
endothelial (middle), and ductal (right) cells in predicted data. **(g)** The expression correlation of the
marker genes in six cell types between the Bulk2Space result and Slice 1.

Bulk2Space reveals spatial, molecular, and functional heterogeneity of B cells in melanoma

Similarly, the same spatial deconvolution procedure of Bulk2Space was performed for another dataset deriving from Tirosh[40] et al., the melanoma bulk transcriptome data. The scRNA-seq data were used as a single-cell reference, one slice of tissue section was used as a spatial reference (Slice 1), and another (Slice 2) was used for bulk data synthesis (Fig. 4a). The expression correlation of marker genes for five cell types between Bulk2Space result and Slice 2 was shown in Fig. 4b. After deconvolution, single cells were assigned to Slice 1 by Bulk2Space (Fig. 4c). As illustrated in Fig. 4d, the composition of cell types in each spot of the tissue showed a strong spatial pattern, which was consistent with the histological annotation of the tissue section. The spatial distribution of each cell type is shown in Fig. 4e. The predicted results for B cells, cancer-associated fibroblasts (CAFs), macrophages, and malignant cells showed distinct spatial heterogeneity. Next, we compared the expression of marker genes in both predicted and reference tissues (Fig. 4f) and found that the spatial pattern of the marker genes in the predicted tissue was highly correlated with that in the spatial reference (Fig. 4g). Surprisingly, B cells assigned to the melanoma area were quite different from those assigned to the lymph node area (Fig. 4h), which suggested that there was heterogeneity in B cells from different tissue regions. However, this spatial difference cannot be detected by other deconvolution methods that only predict the composition of cell types in each spot. The differential expression analysis (Fig. 4i) and pathway enrichment (Fig. 4j) were carried out to further investigate the molecular and functional heterogeneity of B cells. The result indicated that B cells in the lymph node area were related to immunoregulation such as lymphocyte activation, T cell activation, and immune effector process, etc. However, in the melanoma region, many pathways, such as regulation of growth, cell-substrate junction, cell-substrate adhesion, and other pathways associated with cancer cell growth, adhesion, and diffusion were enriched.

289
Fig. 4. Spatially resolved analysis of Melanoma by Bulk2Space. (a) Synthesis of the bulk
transcriptome data. The scRNA-seq data is used as deconvolution reference, one slice of the
sectioned tissue is employed as a spatial reference, and another slice is used to synthesize the input
bulk data. **(b)** Expression correlation of marker genes for five predicted cell types between
Bulk2Space result and Slice 2. **(c)** Single-cell deconvolution of the bulk data. Top, t-SNE plot showed
the clustering of generated single cells and the rose chart showed the proportion of each predicted
cell type. Bottom, distribution of predicted single cells. Different colors represented distinct cell
types. **(d)** The top, Histological annotation for stroma (red), melanoma (black), and lymphoid region
(yellow). Bottom, single-cell decomposition result of each spot in the ST data by Bulk2Space. **(e)**
The spatial abundance of different cell types in each spot on the tissue section is predicted by
Bulk2Space. **(f)** The expression correlation of the marker genes in six cell types between the

Bulk2Space result and Slice 1. **(g)** The spatial expression of the marker genes in B cells, CAFs, macrophages, and malignant cells in predicted data, respectively. **(h)** Spatial distribution of B cells. Segmented area, lymphoid tissue (purple), and melanoma region (black). **(i)** Differentially expressed genes for lymphoid and melanoma tissue. **(j)** Pathway enrichment analysis for lymphoid and melanoma tissue.

Bulk2Space re-annotates ambiguous cells in the mouse hypothalamus

The Bulk2Space algorithm was also used to spatially deconvolve bulk transcriptome data derived from the hypothalamus region of the mouse brain using our in-house developed multiplexed barcoding RNA-seq method, termed Spatial-seq, to explain the spatial distribution of single cells and gene expression. The Spatial-seq workflow is shown in Fig. 5a, laser capture microdissection (LCM) was used to isolate brain regions and a barcoding strategy was utilized for multiplexed RNA-seq (see 'methods' section). After deconvolution by Bulk2Space, the resulting single cells had a clustering space close to the reference scRNA-seq data and MERFISH data (Fig. 5b). The generated single cells were mapped to spatial coordinates based on a spatial reference (Figure 5c). As shown in Fig. 5c, the MERFISH and Bulk2Space results were very close in single-cell spatial distribution. Among the 10 cell types, immature oligodendrocytes were unique to MERFISH, and macrophages were predicted by Bulk2Space, but absent in MERFISH. The heatmap suggested 7 of the 8 shared cell types were strongly correlated in both MERFISH and Bulk2Space results. The low correlation of mural cells may be related to the low number of pericytes. The spatial patterns of the MERFISH targeted genes were quite similar to that of Bulk2Space results (Fig. 5d). Meanwhile, owing to the limited number of measured target genes, some ambiguous cells in MERFISH could not be defined according to their expression profiles. Using Bulk2Space, the resulting single cells were assigned to the spatial context, thus further clustering these ambiguous cells (Fig. 5e). Meanwhile, the spatial distribution of novel genes beyond the targeted RNA species was predicted by Bulk2Space and was highly consistent with the spatial pattern of corresponding cell types. (Fig. 5f).

Fig. 5. Spatially resolved analysis of mouse hypothalamus by Bulk2Space. (a) Spatial-seq workflow. The mouse brain was sectioned continuously from coronal and sagittal directions. For each slice, spatial registration was carried out to annotate each brain region to be isolated. Brain regions are dissected by LCM and labeled with a unique barcode and then followed with RNA-seq. **(b)** The t-SNE layout of single-cell profiles in MERFISH data and generated from bulk tissue by Bulk2Space. The top, clustering of all cells in both datasets. Two heterogeneous sets of data were aligned through a joint analysis. Each cell type was represented by a unique color. **(c)** The spatial distribution of distinct cells in MERFISH and Bulk2Space results. The stacked bar chart showed the cell number of each cell type (grey) and the number of cells predicted by Bulk2Space to the corresponding cell type at each coordinate (blue). Bottom, Pearson correlation of gene expression for different cell types in MERFISH and Bulk2Space data. Correlation for shared cell types was in the red border and the expression correlation between MERFISH ambiguous cells and Bulk2Space predicted cell types was in the green border. **(d)** The spatial expression of the Bulk2Space predicted genes and the ground truth (MERFISH). **(e)** Re-annotation of ambiguous cells in MERFISH using Bulk2Space results. Left, clustering of the ambiguous cells. Right, spatial assignment of cells that

343 were colored according to their predicted cell types. **(f)** Predicted spatial expression patterns of

344 novel marker genes for different cell types.

Bulk2Space uncovers spatial gene expression dynamics in inflammation-induced

prostate cancer

Tumor development is linked to chronic infection, dietary factors, obesity, inhaled pollutants,

tobacco use, and autoimmunity. The unifying principle underlying these processes is inflammation,

which is an aberrantly prolonged form of a protective response to a loss of tissue homeostasis[41-43].

To explore whether Bulk2Space could identify the cellular and molecular pathways that coordinate

the tumor-promoting effects in inflammation-induced cancer, we deconvolved the bulk

transcriptome data of prostate cancer[44] and mapped them to the corresponding spatial references,

including normal glands, inflammatory tissue, and tumor sites. These tissue regions were isolated

from the prostate, and three of the sections were subjected to RNA-seq based on spatial barcoding

(Fig. 6a). Next, Bulk2Space was used to deconvolve the bulk data into single-cell profiles (Fig. 6b)

and mapped the generated single cells to spatial coordinates (Fig. 6c). Interestingly, the

composition and distribution of cells in different tissue regions were quite different and closely

related to the state of the tissue (Fig. 6d and 6e). Comparison of cell distribution among normal,

inflammatory, and cancerous tissues (Fig. 6f to 6h) showed that in normal glands, cancer-

associated fibroblasts (CAFs) were the most abundant, but in inflammatory and cancer tissues, the

proportions of CAFs were significantly lower (52.4% in normal, 20.5% in inflammation, and 27.5%

in cancer). In the areas of inflammation, cells involved in the inflammatory response, such as

immunocompetent B cells, endothelial cells, monocytes, and NK cells (Fig. 6g), were more

abundant than in normal or tumor tissues (13.4% in inflammation, 2.0% in normal, and 1.7% in

cancer). In cancer with a Gleason score (Gs) of 3+4, the proportion of cancer epithelial cells was

significantly higher (53.0% in cancer, 6.3% in normal, and 4.3% in inflammation) than that in normal

and inflammatory areas (Fig. 6h), while the proportion of the periventricular layer cells was

significantly lower (17.9% in cancer, 39.3% in normal, and 44.1% in inflammation). The Bulk2Space

results indicated that, during the progress of inflammation-induced prostate cancer, CAFs

accumulated in normal glands at the early stage, resulting in high expression of inflammatory

factors and promoting the occurrence of local inflammation, a fundamental innate immune

response to perturbed tissue homeostasis, thus leading to tissue cancerization.

Fig. 6. Spatial gene expression dynamics analysis of inflammation-induced prostate cancer by Bulk2Space. (a) Experimental design of the bulk transcriptome data. The tissue was segmented into distinct regions, namely, normal glands (pink), inflammation (blue), and several cancer regions (red, green, purple, and yellow). Three tissue regions (1.1 normal, 2.3 inflammation, and 3.3 cancer region) were isolated to conduct spatially resolved transcriptomics. **(b)** The t-SNE layout of the deconvolved single-cell profiles by Bulk2Space. Each cell type was represented by a unique color. **(c)** The proportions of generated single cells that were mapped to normal, inflammation, and cancer region by Bulk2Space, respectively. The color of each cell type was consistent with (b). **(d)** The proportion of cells assigned to normal (pink), inflammation (blue), and cancer (green) regions for each cell type. **(e)** The predicted proportion of distinct cell types for each tissue region by Bulk2Space. **(f)** The proportions of CAFs in normal (left), inflammation (middle), and cancer regions (right). **(g)** The proportions of immune cells in normal, inflammation, and cancer regions. **(h)** The proportions of cancer epithelial cells in normal, inflammation, and cancer regions.

Discussion

Understanding transcriptional heterogeneity within the tissue from the perspective of single-cell spatial resolution and full gene coverage is an essential direction for the development in the field of biological sciences[1]. However, there are many technical challenges, and thus, such a technology has not been fully realized yet. Traditional bulk transcriptome data can provide gene expression patterns and reflect the overall status of tissue, which is widely used in studies on the occurrence and development of diseases. Nevertheless, disease-inherent cell composition and spatial distribution are difficult to analyze. Although several attempts can infer the cell-type proportions from traditional RNA-seq data, understanding bulk tissue at single-cell spatial resolution remains a pressing need in the field. Cell identity can be characterized by the clustering space of single-cell gene expression profiles[45], and expression features can be stably retained across different conditions, technologies, and species[29]. Based on this, we used a deep learning model, termed β-VAE, to generate single-cell profiles with biological significance within the clustering space of each cell type and map them accurately to tissue coordinates, thus transforming bulk transcriptomics into spatially resolved single-cell transcriptome data. As thoroughly considered the merits and demerits of the widespread used two spatially resolved transcriptomics technologies, we employed different spatial mapping strategies for each method. For image-based *in situ* hybridization, Bulk2Space could predict the spatial expression of novel mRNA species apart from the target genes. Moreover, Bulk2Space can help annotate cells whose types are difficult to further distinguish by targeted methods. For spatial barcoding-based RNA-seq, each sequenced tissue region contains transcripts from different cells and can thus be regarded as a mixture of cells, not allowing single-cell resolution. Bulk2Space can assign single cells to optimal spots, and these mixtures are split into collections of individual cells, thus achieving single-cell resolution. Our results showed that Bulk2Space had great biological and clinical application prospects, including linking tissue molecular characteristics with histological phenotypes, revealing the spatial specific variation of cells, discovering the spatial expression pattern of novel genes, achieving more refined cell clustering, and uncovering the molecular mechanism underlying the progression of the disease.

Acknowledgment

This work is supported by the National Natural Science Foundation of China (81973701, 91846204, and U19B2027), the Natural Science Foundation of Zhejiang Province (LZ20H290002), the Innovation Team and Talents Cultivation Program of the National Administration of Traditional Chinese Medicine (ZYYCXTD-D-202002), the Westlake Laboratory (Westlake Laboratory of Life Sciences and Biomedicine), and the Alibaba Cloud (Alibaba Cloud Computing Co. Ltd.).

Author contributions

X.F., H.C., Y. G., and J. L. conceived the study. J. L. and J. Q. collected datasets involved in this article, benchmarked all methods, and participated in the development of the Bulk2Space algorithm. Y. F., Z. C., and X. Z. implemented the code and interface to the Bulk2Space algorithm. N. Z., X. S., H. Y., P. Y., J. C., L. S., and J. Z. provided a lot of advice on algorithm implementation and biological applications. J. L., Y. H., Y. H., L. Y., and X. L. conducted the multiplexed RNA-seq experiment using Spatial-seq. D. W. provided important advice on the brain tissue isolation, spatial registration, and annotation. All authors wrote the manuscript, read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Methods

Datasets

All scRNA-seq, spatially resolved transcriptomics, and bulk RNA-seq datasets used for

benchmarking of Bulk2Space were retrieved from high-quality publications, Gene Expression

Omnibus (GEO), and The Cancer Genome Atlas (TCGA), wherein unannotated, ambiguous, or low

quality-cells were excluded.

scRNA-seq datasets. Ten scRNA-seq were applied to benchmark the Bulk2Space algorithm,

including five human datasets, peripheral blood, brain, kidney, liver, and lung tissues, and five

mouse datasets, brain, kidney, lung, pancreas, and testis tissues. A dataset from the hypothalamic

preoptic region in mouse brain containing 30370 cells of 12 cell types is used for spatial mapping

using image-based methods. The human pancreatic ductal adenocarcinomas (PDAC), human

melanoma, and human prostate cancer data were used for the validation of Bulk2Space.

Spatially resolved transcriptome data. For image-based spatial mapping, the mouse

hypothalamus MERFISH data were obtained from Dryad (doi:10.5061/dryad.8t8s248), and the

Bregma +0.26 slice of mouse #1 was chosen to further analysis. For spatial barcoding-based

methods, the human PDAC, human melanoma, and human prostate cancer data were used for

reference and validation.

Bulk transcriptome data. Two bulk RNA-seq datasets were used to illustrate the applications of

Bulk2Space. The mouse hypothalamus data were derived from experiments of our in-house

developed Spatial-seq protocols. The human prostate cancer RNA-seq data were downloaded from

TCGA.

Data processing

All data were preprocessed using R (version 4.1.1). For mouse brain scRNA-seq data by Moffitt[38] et

al., we filtered 841 ambiguous cells and 88 unstable cells. For human prostate cancer scRNA-seq

data by Wu[46] et al., we filtered 183 unassigned cells. All cells of other scRNA-seq as well as spatially

resolved transcriptomics datasets were retained. For mouse hypothalamus MERFISH data, we

filtered 5 "blank" barcodes as well as *Fos* gene, whose expression value in all cells was 'NA'. For

human prostate cancer RNA-seq data, we used biomaRt (version 2.48.3) to transform ensemble id

to gene name. For all scRNA-seq and spatially resolved transcriptomics datasets, the raw counts

were normalized using the global-scaling normalization method "LogNormalize". For MERFISH

dataset, the normalized data was determined as the raw count per cell divided by the cell volume

and scaled by 1000.

Step 1: Deconvolution of bulk transcriptome data.

Cell type proportions prediction. For a given bulk RNA-seq dataset, we aimed to calculate its cell

type proportions firstly, and then generate single-cell gene expression profiles based on the

calculated proportions. Since cell type and gene expression of all single cells can be accessed from

the scRNA-seq data reference, we collected cells with the same cell type and then average their

gene expression. This average vector $\mathbf{c}_i \in \mathbb{R}^N$ is defined as the gene expression of cell type i ($i \in$

$\{1,2,\dots,C\}$, where C denotes the total number of cell types). Given the gene expression vector

$\mathbf{x} \in \mathbb{R}^N$ of a bulk RNA-seq dataset, we aimed to predict the cell type proportion:

477
$$\sum_{i=1}^{C} p_i \mathbf{c}_i = \mathbf{x}$$

478 Here, p_i is the proportion of cell type i.

479 We applied least square estimation (LSE) to estimate p_i by minimizing the squared

480 discrepancies between observed data.

481
$$\min_{\boldsymbol{p}} \|\mathbf{x} - \boldsymbol{Cp}\|_2$$

where $\mathbf{C} \in \mathbb{R}^{N \times C}$ denotes the cell type gene expression profile, and each column represents

a different cell type. $\boldsymbol{p} = [p_1, p_2, \dots, p_C]^T$ is the proportion vector to be estimated.

Notably, we only chose top 500 marker genes of each cell type in this step, to reduce the noise

effect.

Single-cell simulation by β-VAE. Let $\mathcal{D} = \{X, V, W\}$ be a set consisting of gene expression vectors

$x \in \mathbb{R}^N$ and two sets of ground truth data generative factors: conditionally independent factors

$v \in \mathbb{R}^K$, where $\log p(v|x) = \sum_k \log p(v_k|x)$; and conditionally dependent factors $w \in \mathbb{R}^H$. We

assume that x are generated by a true world simulator S using the corresponding ground truth

data generative factors: $p_\theta(x|v, w) = S(v, w)$, where θ is the generative model parameter.

We wanted to develop an unsupervised deep generative model that, using samples from X

only, can learn the joint distribution of the data x and a set of generative latent factors z ($z \in$

$\mathbb{R}^M$, where $M \geq K$) that can be used to generate the observed data x; i.e., $p_\theta(x|z) \approx$

$p(x|v, w) = S(v, w)$. However, since the integral of the marginal likelihood $p_\theta(x) =$

$\int p_\theta(\mathbf{z})p_\theta(\mathbf{x}|\mathbf{z}) \, d\mathbf{z}$ is intractable (so we cannot evaluate or differentiate the marginal likelihood),

the true posterior density $p_\theta(\mathbf{z}|x) = p_\theta(\mathbf{x}|\mathbf{z})p_\theta(\mathbf{z})/p_\theta(\mathbf{x})$ is intractable.

To solve this problem, for a given observation x, we described the inferred posterior

configurations of the latent factors $\mathbf{z}$ through a probability distribution $q_\phi(\mathbf{z}|\mathbf{x})$: an

approximation to the intractable true posterior $p_\theta(x|z)$. We aimed to ensure that the inferred

latent factors $q_\phi(\mathbf{z}|\mathbf{x})$ capture the generative factors v in a disentangled manner. A

disentangled representation can be defined as one where single latent units are sensitive to

changes in single generative factors, while being relatively invariant to changes in other factors. In

a disentangled representation, knowledge about one factor can generalize to novel configurations

of other factors. The conditionally dependent data generative factors w can remain entangled in

a separate subset of $\mathbf{z}$ that is not used for representing v.

An intuitive approach was to minimize the KL divergence between the approximate and the

true posterior:

$$\mathcal{D}_{KL}\left(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z}|x)\right) = -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{z}|x)}{q_\phi(\mathbf{z}|\mathbf{x})}\right) = -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{\frac{p_\theta(x, \mathbf{z})}{p_\theta(x)}}{q_\phi(\mathbf{z}|\mathbf{x})}\right)$$

$$= -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \left[\log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right) - \log(p_\theta(x))\right]$$

$$= -\sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right) + \log(p_\theta(x)) = -\mathcal{L}(\theta, \phi; \mathbf{x}) + \log(p_\theta(x))$$

Here, $\mathcal{L}(\theta, \phi; \mathbf{x})$ is called the variational lower bound, and can be written as:

512
$$\mathcal{L}(\theta, \phi; \mathbf{x}) = \sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right) = \sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \log\left(\frac{p_\theta(\mathbf{x}|\mathbf{z})p_\theta(\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right)$$

513
$$= \sum_{\mathbf{z}} q_\phi(\mathbf{z}|\mathbf{x}) \left[\log(p_\theta(\mathbf{x}|\mathbf{z})) + \log\left(\frac{p_\theta(\mathbf{z})}{q_\phi(\mathbf{z}|\mathbf{x})}\right)\right]$$

514
$$= \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}\left[\log(p_\theta(\mathbf{x}|\mathbf{z}))\right] - \mathcal{D}_{KL}\left(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\right)$$

515 To encourage this disentangling property in the inferred $q_\phi(\mathbf{z}|\mathbf{x})$, we introduced a constraint

516 over it by trying to match it to a prior $p(\mathbf{z})$ that can both control the capacity of the latent

517 information bottleneck, and embodies the desiderata of statistical independence mentioned

518 above. We set the prior to be an isotropic unit Gaussian $p(z) \sim \mathcal{N}(0, I)$, then the constrained

519 optimization problem can be written as:

520
$$\max_{\phi,\theta} \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}[\log(p_\theta(\mathbf{x}|\mathbf{z}))] \; subject \; to \; \mathcal{D}_{KL}\left(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\right) < \epsilon$$

521 where ϵ specifies the strength of the applied constraint. Re-writing the above equation as a

522 Lagrangian under the KKT conditions, we obtain:

523
$$\mathcal{F}(\theta, \phi, \beta; \mathbf{x}, \mathbf{z}) = \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}[\log(p_\theta(\mathbf{x}|\mathbf{z}))] - \beta\left(\mathcal{D}_{KL}\left(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\right) - \epsilon\right)$$

524 where the KKT multiplier β is the regularization coefficient that constrains the capacity of the

525 latent information channel $\mathbf{z}$ and puts implicit independence pressure on the learnt posterior due

526 to the isotropic nature of the Gaussian prior $p_\theta(\mathbf{z})$. Since $\beta, \epsilon \geq 0$, according to the

527 complementary slackness KKT condition, the equation can be re-written as:

528
$$\mathcal{F}(\theta, \phi, \beta; \mathbf{x}, \mathbf{z}) \geq \mathcal{L}(\theta, \phi; \mathbf{x}, \mathbf{z}, \beta) = \mathbb{E}_{q_\phi(\mathbf{z}|\mathbf{x})}[\log(p_\theta(\mathbf{x}|\mathbf{z}))] - \beta\left(\mathcal{D}_{KL}\left(q_\phi(\mathbf{z}|\mathbf{x})||p_\theta(\mathbf{z})\right)\right)$$

which is the β-VAE formulation with the addition of the β coefficient.

Here, different β will change the degree of applied learning pressure during training, thereby

encouraging different learnt representations. We postulated that to learn disentangled

representations of the conditionally independent data generative factors $\boldsymbol{v}$, it is important to set

$\beta > 1$ to impose a stronger constraint on latent bottleneck than the original VAE. These

constraints limit the capacity of $\mathbf{z}$, coupled with the pressure to maximize the log likelihood of the

training data $\mathbf{x}$ under the model, encouraging the model to learn the most efficient

representation of the data. The additional pressure from high β values may create a trade-off

between reconstruction fidelity and the quality of disentanglement within the learnt latent

representations. When the appropriate balance is found between information preservation

(reconstruction cost as regularization) and latent channel capacity restriction ($\beta > 1$), disentangled

representations emerge.

Step2: Mapping generated single cells to spatial locations

Spatial barcoding-based RNA-seq. After obtaining the generated data, we needed to map each cell

and the spot it belongs to. To improve the accuracy and reduce the complexity of this process, we

divided it into two steps. First, we calculated the cell type proportion of each spot, and then

predicted which cells are contained in each spot based on this proportion.

1) Calculate cell type proportions. The procedure is identical to the Step 1 deconvolution, we

repeated this process to obtain the cell type proportion of each spot in the tissue.

2) Deep forest for spot recommendation. With cell type proportion, we next predicted which

cells are contained in each spot. We defined this task as a binary classification problem. We

designed a classifier, whose input is the gene expression vectors of both single cell $\mathbf{s}$ and spot $\mathbf{x}$.

When the cell belongs to the spot, the output of the classifier is 1, otherwise the output is 0.

Specifically, the input can be represented as $\mathbf{v} = [\mathbf{s}; \mathbf{x}; \mathbf{s} - \mathbf{x}] \in \mathbb{R}^{N \times 3}$, where semicolon means

concatenate operation. $\mathbf{s} - \mathbf{x} \in \mathbb{R}^N$ can be regarded as an auxiliary signal, since when there is a

negative number occurring in this vector, the single cell must not belong to the spot.

We applied multi-Grained Cascade Forest (gcForest), a decision tree ensemble with a cascade

structure and further enhanced by multi-grained scanning, to build our classifier. Each level of

cascade receives feature information processed by its preceding level and outputs its processing

result to the next level. Each level is an ensemble of decision tree forests, i.e., an ensemble of

ensembles. We also included completely-random tree forests) to encourage diversity. Given an

instance, each forest counts the percentage of different classes of training samples at the leaf node where the concerned instance falls, and then average across all trees in the same forest to produce an estimate of class distribution. The estimated class distribution forms a class vector, which was generated by k-fold cross validation and then concatenated with the original feature vector to be input to the next level of the cascade.

To enhance cascade forest, we introduced a procedure of multi-grained scanning. We used sliding windows to scan the raw features. Specially, we used a window with a fixed length of k to slide the input vector into k-dimensional feature vectors. We then feeded these vectors to n forests and finally produce n two-dimensional vectors. By using multiple sizes k of sliding windows, differently grained feature vectors were generated.

The we summarized the overall procedure of gcForest. For the original raw features, three window sizes $\{\lfloor N/16 \rfloor, \lfloor N/8 \rfloor, \lfloor N/4 \rfloor\}$ were used for multi-grained scanning. The generated data were used to train a completely-random tree forest containing 500 trees. We concatenated the output of these two forests as the transformed feature vectors, which also acted as the input of cascade forest. The final model was a cascade of cascades, where each cascade consisted of multiple levels each corresponding to a grain of scanning. Each level consisted of 4 completely-random tree forests, each containing 500 trees. In other words, the transformed feature vectors were augmented with the class vector generated by the previous grade, and then were used to train the current grade of cascade forests.

Given a test instance, it would obtain its corresponding transformed feature vector through the multi-grained scanning procedure, and then go through the cascade till the last level. The two-dimensional class vectors at the last level were aggregated, and the class with the largest aggregate value was selected to obtain the final prediction result.

This decision tree ensemble approach not only had much fewer hyper-parameters than deep neural networks, but also retained the interpretability of tree models. Its model complexity could be automatically determined in a data-dependent manner, which made gcForest work well even on small-scale data.

Image-based in situ hybridization. We developed a simple method for predicting the spatial

distributions of genes not measured in spatial transcriptomic data which produced by *in situ* RNA

imaging-based technologies. To do this, Bulk2space used the spatial transcriptomic data as the

reference, and generated scRNA-seq data were mapped to the tissue space corresponding to this

reference. To eliminate the differences between scRNA-seq data and spatial transcriptomic data

caused by different experiment types, an empirical Bayes framework was used to remove the batch

effect, so that both sets of data are at the same scale level. Subsequently, the cross-dataset k-

nearest neighbor graph of spatial transcriptomic data in scRNA-seq data was computed in the

aligned space, and then the predicted whole gene expression profile of each cell of spatial

transcriptomic data could be calculated to the mean of its k nearest neighbors in scRNA-seq data.

Data simulations

Using several simulations, we evaluated the performance of Bulk2space. The first simulation

compared the generation of single-cell gene expression profiles by Bulk2Space with other methods,

GAN and CGAN. The simulated data were derived from 10 scRNA-seq data of benchmarking usage.

We randomly divided the scRNA-seq data into two parts, one was treated as the single-cell

reference and the other for the construction of bulk transcriptome data via aggregating all the

single-cell gene expression profiles. Considering the cell composition of bulk RNA-seq data varies

greatly in the natrual situation, we further changed the cell-type proportion and synthesized three

bulk transcriptome data with different cell compositions from the same part of scRNA-seq data. In

total, 40 paired simulated data were synthesized in this study. Performance was assessed using the

Pearson correlation coefficient (PCC), Spearman's rank correlation coefficient (SRCC), and root

mean squared error (RMSE) between the gene expression profile of scRNA-seq data reference and

synthetic bulk transcriptome data.

In the second simulation, we evaluated the robustness of Bulk2space. We introduced two kinds

of noise: cell type noise and gene expression noise. For cell type noise, we randomly altered the

type of cells with the ratio of 0.01, 0.02, 0.04, 0.08, 0.1, 0.2, 0.3, 0.4, 0.5, 0.7, and 0.9. For gene

expression noise, we firstly constructed a noise expression profile $\mathbf{C}$,

$$\mathbf{C} \sim \mathrm{U}(-x, x)$$

Here, x is the maximum gene expression value of scRNA-seq data reference.

We next constructed synthetic expression matrix by combining noise expression profile with

scRNA-seq data reference, and the ratio of gene expression-altered cells remains 0.01, 0.02, 0.04,

0.08, 0.1, 0.2, 0.3, 0.4, 0.5, 0.7, and 0.9. PCC, SRCC and RMSE between the synthetic expression

matrix and constructed bulk RNA-seq data were used to evaluate performance.

In the third simulation, we simulated spatial barcoding data to evaluate the spatial

reconstruction capacity of Bulk2space with other five classical classification models: Logistic

Regression (LR), Decision Tree (DT), Gradient Boosting Decision Tree (GBDT), Multilayer Perceptron

(MLP) and DeepGBM. Starting from scRNA-seq data reference, we randomly chose 10 cells from it

and aggregated their transcriptomic profiles as a spot of pseudo spatial transcriptomic data. The

spot with over 25000 UMI counts would be sampled down to 20000 UMI counts to better meet

the true situation. We also constructed pseudo spatial transcriptomic data with 100, 200, 500 and

1000 spot numbers to simulate true ST data produced by different spatial barcoding technologies.

PCC, SRCC, and RMSE of cell type proportion and gene expression profile between model

prediction results and ground truth were used to evaluate performance.

In the fourth and fifth simulations, we systematically evaluated the performance of Bulk2space.

We used Bulk2space into 2 types of cancer data, pancreatic ductal adenocarcinomas (PDAC) and

melanoma. The bulk RNA-seq data were simulated from the biological replicates of selected ST

slices.

Spatial-seq protocol

Multiplexed spatial barcoding. A virtual 48×48×48 three-dimensional well array (barcoding array)

was constructed in X, Y, and Z directions, with 48×48 wells in each layer. 384-well plates (24×16

wells) were selected as the basic units (2×3 plates) of each layer. Each well in the barcoding array

can be labeled as barcode $\left(\{X_i, Y_j, Z_k\}, i, j, k \in N, [1, 48]\right)$. Magnetic beads coated with carboxyl

group were distributed into each well for barcode synthesis. There were three rounds of barcode

extension reactions. In the first round, 5' amino-modified barcoded oligonucleotides (barcode X_i)

were conjugated to the beads. In the second round of extension reaction, barcoded

oligonucleotides (barcode Y_j) which had a sequence of bases at the 5' end complementary to the

3' end of barcode X_i were linked to the end of barcode X_i by PCR. Similarly, in the third round

of reaction, barcode sequences containing a unique molecular identifier (UMI) and a polyT tail

(barcode Z_k) were extended to barcode Y_j by PCR. This method only needs 144 (48×3) different

barcodes to theoretically generate magnetic beads with 48^3 distinct kinds of magnetic beads with

known sequences. More importantly, the entire synthesis process was serialized and multiplexed,

which can be accomplished with the help of arrayed pipettes or liquid workstations, rather than

synthesizing the magnetic beads well by well. The spatially barcoded beads can be used in many

areas, such as multiplexed labeling of tissue samples or even single cells.

Collection of mouse brain regions. The mouse brain was sliced into 14-μm sections from the coronal

and sagittal directions. Each tissue slice was registered to a reference brain template provided by

the Allen Brain Atlas (https://portal.brain-map.org/). After spatial registration, the anatomical

regions of the mouse brain were delineated and annotated. The outlines, dissection sequences,

and collectors of the brain regions were specified from the annotated data and then imported to

an LCM instrument (Laser Microdissection System equipped with a DM68 microscope, a Leica

LMD6 laser cutter, and a single-cell capture collector LMT350, Leica Microsystems, Germany).LCM

is a microscope-guided powerful cutting system incorporating UV light for contact- and

contamination-free isolation of areas of interest from tissue sections[15]. Brain regions were

dissected from the tissue by LCM according to the imported files and fell into the collector loaded

with barcoded beads in advance. Spatial-seq has the potential to achieve spatially resolved whole

transcriptome sequencing of a large number of single cells.

Multiplex RNA-seq. The tissue was lysed in the barcoded well, allowing mRNA to be captured by

polyT tail on the surface of the magnetic beads. Each well was loaded with lysis buffer. The tissue

was lysed on ice for 12 minutes. The beads were then washed and transferred for reverse

transcription (RT). The captured mRNA was reverse transcribed to construct cDNA libraries

following the guidance of Illumina Nextera XT DNA Library Preparation Kits. A paired-end

sequencing was conducted to decode the spatial barcode in the 3' end and detect RNA species in

the 5' end of the cDNA.

Sequence alignment. FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) was

utilized for quality control of the RNA-seq data. The derived RNA-seq data were fragmented into

lots of files based on the spatial barcodes. The digital gene expression matrix for each barcode was

retrieved follow the Drop-seq sequence alignment cookbook.

Data normalization. FPKM, RPKM, RPM, and TPM were calculated from the gene expression matrix

according to the following formulas.

677
$$\text{FPKM} = \frac{ExonMappedFragments * 10^9}{TotalMappedFragments * ExonLength}$$

678
$$\text{RPKM} = \frac{(ExonMappedRead/TotalMappedReads * 10^6)}{ExonLength} * 10^3$$

679
$$\text{RPM} = \frac{ExonMappedReads * 10^6}{TotalMappedReads}$$

680
$$\text{TPM} = \frac{N_i/L_i * 10^6}{sum(N_1/L_1 + N_2/L_2 + \cdots + N_n/L_n)}$$

681   The data were normalized using the global-scaling normalization method "LogNormalize" in

682   Seurat. A Bayesian network-based method termed ComBat was used for batch effect removal of

683   data derived from different cDNA libraries. The expression matrix of each sample was normalized

684   using the "Deconvolution Normalization" algorithm to align the median gene expression.

Data availability

687   The original and processed mouse brain data used in this study were uploaded to GEO (GSE192999).

Code availability

Bulk2Space is available as a python package and the source code is deposited at the GitHub (https://github.com/ZJUFanLab/bulk2space).

Statistics

Python (version 3.8.5) and R (version 4.1.1) were used for the statistical analysis.

References

1. Liao, J., Lu, X., Shao, X., Zhu, L. & Fan, X. Uncovering an Organ's Molecular Architecture at Single-Cell Resolution by Spatially Resolved Transcriptomics. *Trends Biotechnol* **39**, 43-58 (2021).

2. Wang, Y.S. & Guo, J. Multiplexed Single-Cell in situ RNA Profiling. *Front Mol Biosci* **8**, 775410 (2021).

3. Srivatsan, S.R. et al. Embryo-scale, single-cell spatial transcriptomics. *Science* **373**, 111-117 (2021).

4. Waylen, L.N., Nim, H.T., Martelotto, L.G. & Ramialison, M. From whole-mount to single-cell spatial assessment of gene expression in 3D. *Commun Biol* **3**, 602 (2020).

5. Qian, X. et al. Probabilistic cell typing enables fine mapping of closely related cell types in situ. *Nat Methods* **17**, 101-106 (2020).

6. Larsson, L., Frisen, J. & Lundeberg, J. Spatially resolved transcriptomics adds a new dimension to genomics. *Nat Methods* **18**, 15-18 (2021).

7. Chen, K.H., Boettiger, A.N., Moffitt, J.R., Wang, S. & Zhuang, X. RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* **348**, aaa6090 (2015).

8. Lubeck, E., Coskun, A.F., Zhiyentayev, T., Ahmad, M. & Cai, L. Single-cell in situ RNA profiling by sequential hybridization. *Nat Methods* **11**, 360-361 (2014).

9. Wang, X. et al. Three-dimensional intact-tissue sequencing of single-cell transcriptional states. *Science* **361** (2018).

10. Eng, C.L. et al. Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH. *Nature* **568**, 235-239 (2019).

11. Vickovic, S. et al. High-definition spatial transcriptomics for in situ tissue profiling. *Nat Methods* **16**, 987-990 (2019).

12. Stickels, R.R. et al. Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nat Biotechnol* **39**, 313-319 (2021).

13. Rodriques, S.G. et al. Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. *Science* **363**, 1463-1467 (2019).

14. Stahl, P.L. et al. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78-82 (2016).

15. Nichterwitz, S. et al. Laser capture microscopy coupled with Smart-seq2 for precise spatial transcriptomic profiling. *Nat Commun* **7**, 12139 (2016).

16. Chen, J. et al. Spatial transcriptomic analysis of cryosectioned tissue samples with Geo-seq. *Nat Protoc* **12**, 566-580 (2017).

17. Cheng, J., Liao, J., Shao, X., Lu, X. & Fan, X. Multiplexing Methods for Simultaneous Large-Scale Transcriptomic Profiling of Samples at Single-Cell Resolution. *Adv Sci (Weinh)* **8**, e2101229 (2021).

18. Liu, Y., Ai, N., Liao, J. & Fan, X. Transcriptomics: a sword to cut the Gordian knot of traditional Chinese

medicine. *Biomark Med* **9**, 1201-1213 (2015).

19. Djebali, S. et al. Landscape of transcription in human cells. *Nature* **489**, 101-108 (2012).

20. Cancer Genome Atlas Research, N. Comprehensive genomic characterization defines human
glioblastoma genes and core pathways. *Nature* **455**, 1061-1068 (2008).

21. International Cancer Genome, C. et al. International network of cancer genome projects. *Nature* **464**,
993-998 (2010).

22. Consortium, S.M.-I. A comprehensive assessment of RNA-seq accuracy, reproducibility and
information content by the Sequencing Quality Control Consortium. *Nat Biotechnol* **32**, 903-914
(2014).

23. Frishberg, A. et al. Cell composition analysis of bulk genomics using single-cell data. *Nat Methods*
16, 327-332 (2019).

24. Newman, A.M. et al. Robust enumeration of cell subsets from tissue expression profiles. *Nat*
Methods **12**, 453-457 (2015).

25. Wang, X., Park, J., Susztak, K., Zhang, N.R. & Li, M. Bulk tissue cell type deconvolution with multi-
subject single-cell expression reference. *Nat Commun* **10**, 380 (2019).

26. Chen, Z. et al. Inference of immune cell composition on the expression profiles of mouse tissue. *Sci*
Rep **7**, 40508 (2017).

27. Gangalum, R.K. et al. Spatial Analysis of Single Fiber Cells of the Developing Ocular Lens Reveals
Regulated Heterogeneity of Gene Expression. *iScience* **10**, 66-79 (2018).

28. Ma, A., McDermaid, A., Xu, J., Chang, Y. & Ma, Q. Integrative Methods and Practical Challenges for
Single-Cell Multi-omics. *Trends Biotechnol* **38**, 1007-1022 (2020).

29. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data
across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018).

30. Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019).

31. Shao, X. et al. scCATCH: Automatic Annotation on Cell Types of Clusters from Single-Cell RNA
Sequencing Data. *iScience* **23**, 100882 (2020).

32. Shao, X. et al. scDeepSort: a pre-trained cell-type annotation method for single-cell transcriptomics
using deep learning with a weighted graph neural network. *Nucleic Acids Res* **49**, e122 (2021).

33. Higgins, I. et al. in ICLR (2017).

34. Goodfellow, I.J. et al. in NIPS (2014).

35. Mirza, M. & Osindero, S.J.A. Conditional Generative Adversarial Nets.   **abs/1411.1784** (2014).

36. Gierahn, T.M. et al. Seq-Well: portable, low-cost RNA sequencing of single cells at high throughput.
Nat Methods **14**, 395-398 (2017).

37. Vieira Braga, F.A. et al. A cellular census of human lungs identifies novel cell states in health and in
asthma. *Nat Med* **25**, 1153-1163 (2019).

38. Moffitt, J.R. et al. Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic
region. *Science* **362** (2018).

39. Moncada, R. et al. Integrating microarray-based spatial transcriptomics and single-cell RNA-seq
reveals tissue architecture in pancreatic ductal adenocarcinomas. *Nat Biotechnol* **38**, 333-342 (2020).

40. Tirosh, I. et al. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-
seq. *Science* **352**, 189-196 (2016).

41. Elinav, E. et al. Inflammation-induced cancer: crosstalk between tumours, immune cells and
microorganisms. *Nat Rev Cancer* **13**, 759-771 (2013).

42. Shao, X., Lu, X., Liao, J., Chen, H. & Fan, X. New avenues for systematically inferring cell-cell
communication: through single-cell transcriptomics data. *Protein Cell* **11**, 866-880 (2020).

43. Shao, X. et al. CellTalkDB: a manually curated database of ligand-receptor interactions in humans
and mice. *Brief Bioinform* **22** (2021).

44. Berglund, E. et al. Spatial maps of prostate cancer transcriptomes reveal an unexplored landscape
of heterogeneity. *Nat Commun* **9**, 2419 (2018).

45. Shao, L. et al. Identify differential genes and cell subclusters from time-series scRNA-seq data using
scTITANS. *Comput Struct Biotechnol J* **19**, 4132-4141 (2021).

46. Wu, S.Z. et al. Cryopreservation of human cancers conserves tumour heterogeneity for single-cell
multi-omics analysis. *Genome Med* **13**, 81 (2021).