Title: Neuron-specific protein network mapping of autism risk genes identifies shared biological mechanisms and disease relevant pathologies

Authors: *Nadeem Murtaza[1,2], Annie A. Cheng[2], Chad O. Brown[1,2], Durga Praveen Meka[3], Shuai Hong[3], Jarryll A. Uy[2,4], Joelle El-Hajjar[1], Neta Pipko[1], Brianna K. Unda[2], Birgit Schwanke[3], Sansi Xing[1], Bhooma Thiruvahindrapuram[5], Worrawat Engchuan[5], Brett Trost[5], Eric Deneault[6], Froylan Calderon de Anda[3], Bradley W. Doble[7], James Ellis[5,8], Evdokia Anagnostou[9,10], Gary D. Bader[8], Stephen W. Scherer[5,8], Yu Lu[1], Karun K. Singh[2,4]**

Affiliations:
[1]Department of Biochemistry and Biomedical Sciences, Faculty of Health Sciences, McMaster University, Hamilton, ON, Canada
[2]Krembil Research Institute, University Health Network, Toronto, ON, Canada
[3]Center for Molecular Neurobiology Hamburg (ZMNH), Research Group Neuronal Development, University Medical Center Hamburg-Eppendorf (UKE), Hamburg, Germany
[4]Department of Laboratory Medicine and Pathobiology, Faculty of Medicine, University of Toronto, Toronto, ON, Canada
[5]The Centre for Applied Genomics, The Hospital for Sick Children, Toronto, ON, Canada
[6]Centre for Biologics Evaluation, Biologic and Radiopharmaceutical Drugs Directorate, Health Products and Food Branch, Health Canada, Ottawa, Ontario, Canada
[7]Department of Paediatrics, Department of Biochemistry and Medical Genetics, University of Manitoba, Winnipeg, MB, Canada
[8]Department of Molecular Genetics, Faculty of Medicine, University of Toronto, Toronto, ON, Canada
[9]Department of Neurology, The Hospital for Sick Children, Toronto, ON, Canada.
[10]Bloorview Research Institute, Holland Bloorview Kids Rehabilitation Hospital, Toronto, ON, Canada.

*Corresponding author and Lead Contact: karun.singh@uhnresearch.ca

Acknowledgements

We thank Courtney Irwin. and Paolo de Guzman for proof reading the manuscript. We also thank Kristin J. Brennand and Michael B. Fernandez for providing the NRXN1 cDNA. Graphical abstract and flowcharts were created with BioRender.com (BU24JATEEX, SD235B8ORF, KW235KT7TM, RZ235KTA0S).

Funding

The study was supported by the Canadian Institute of Health Research (CIHR), the Ontario Brain Institute (OBI), the National Science and Engineering Research Council (NSERC) of Canada, the Network for European Funding for Neuroscience Research (NEURON ERA-NET), and the Donald K. Johnson Eye Institute at the Krembil Research Institute.

Author contributions

N.M. and K.K.S. conceived the project. N.M. and K.K.S. wrote the paper with input from A.A.C., B.T., W.E., B.T., and B.W.D. A.A.C and B.K.U created the BioID2 lentiviral construct. N.M. generated all subsequent DNA constructs and performed all experiments and data analysis unless otherwise specified. N.M and A.A.C. generated all lentiviruses. S.X. and Y.L. ran mass spectrometer samples and data acquisition. C.O.B. performed electrophysiology recordings. J.A.U, J.E.H., and N.P. assisted with western blots. D.P.M., S.H., B.S., and F.C.dA. performed and analysed mitochondrial experiments. E.D., J.E, and S.W.S helped to create the human *TAOK2* KO and A135P iPSC lines. E.A. advised on clinical score analysis and G.D.B. advised on pathway analyses. K.S.S supervised the project.

Declaration of Interests

The authors declare no competing interests.

Inclusion and Diversity

We support inclusive, diverse, and equitable conduct of research.

Manuscript summary

There are hundreds of risk genes associated with autism spectrum disorder

(ASD), but signaling networks at the protein level remain unexplored. We use neuron-

specific proximity-labeling proteomics (BioID) to identify protein-protein interaction (PPI)

networks for 41 ASD-risk genes. Neuron-specific PPI networks, including synaptic

transmission proteins, are disrupted by *de novo* missense variants. The PPI network

map reveals convergent pathways, including mitochondrial/metabolic processes, Wnt

signaling, and MAPK signaling. CRISPR knockout reveal an association between

mitochondrial activity and ASD-risk genes. The PPI network shows an enrichment of

112 additional ASD-risk genes and differentially expressed genes from post-mortem

ASD patients. Clustering of risk genes based on PPI networks identifies gene groups

corresponding to clinical behavior score severity. Our data reveal that cell type-specific

PPI networks can identify individual and convergent ASD signaling networks, provide a

method to assess patient variants, and reveal biological insight into disease

mechanisms and sub-cohorts of ASD.

Introduction

The risk of developing ASD has a strong genetic basis, including common and rare genetic variants (Gaugler et al., 2014; Iossifov et al., 2014; Robinson et al., 2016; Weiner et al., 2017). Large scale sequencing studies have identified hundreds of genes associated with ASD risk (Feliciano et al., 2019; Glessner et al., 2009; Ruzzo et al., 2019; Samocha et al., 2014; Satterstrom et al., 2020; Wilfert et al., 2021; Yuen et al., 2017). One hypothesis is multiple ASD risk genes converge in developmental brain signaling networks.

The majority of known convergent ASD-associated pathways are based on sequencing, transcriptomics, and gene co-expression analyses (Cederquist et al., 2020; Jin et al., 2020; Parikshak et al., 2013; Ramaswami et al., 2020; Sanders et al., 2015; Satterstrom et al., 2020; Willsey et al., 2013, 2021; Yuen et al., 2017). These studies have implicated synaptic transmission, translation, transcription, chromatin remodeling and gene splicing (Chang et al., 2015; O'Roak et al., 2012a; Ramaswami et al., 2020; Velmeshev et al., 2019; Voineagu et al., 2011). However, the majority of autism risk genes encode proteins, and protein-protein interactions (PPIs) are an essential mechanism of signaling (Neale et al., 2012; O'Roak et al., 2012b). Given that a large proportion of ASD genes have functions that do not occur in the nucleus (De Rubeis et al., 2014; Satterstrom et al., 2020), PPI networks provide an unbiased approach to gain insights into unknown convergent ASD disease processes (Kuzmanov and Emili, 2013; Murtaza et al., 2020). Previous studies have shown that ASD genes form PPI networks that can be shared (Chang et al., 2015; Chen et al., 2020; Corominas et al., 2014; Li et al., 2015; Neale et al., 2012; O'Roak et al., 2012a; Sakai et al., 2011). However, these data do not represent brain-specific networks (Lage, 2014). The lack of ASD risk-gene PPI networks represents a missing link towards understanding biological mechanisms in ASD.

Multiple proteomic techniques can be used to identify PPIs (reviewed in Richards et al., 2021). Further, many brain-expressed genes are large in size, including ASD-risk genes, which limits possible expression systems (Casanova et al., 2019). We balanced gene size limitations to capture strong and transient interactions, as well as proteins in

close proximity, to build BioID PPI networks for ASD risk genes. We developed an *in*
vitro proximity-labeling proteomics (BioID2) system with mouse primary neurons.
Proximity-labeling proteomics can capture physiologically relevant interactomes in
neural cell-types (Hamdan et al., 2020; Loh et al., 2016; Spence et al., 2019; Uezu et
al., 2016) or to map cellular compartments (Go et al., 2021; Hung et al., 2017;
Markmiller et al., 2018; Youn et al., 2018). We captured PPI networks from mouse
cortical neurons given their role in ASD pathology, while growing them with glial cells to
promote maturation (Barres, 2008; Stogsdill and Eroglu, 2017; Wilton et al., 2019).

In the current study, we screened the interactome of 41 ASD-risk proteins in
neurons using BIOID. We targeted non-nuclear proteins (e.g., cytosolic proteins,
receptors, kinases, and intracellular signaling proteins) and identified 1770 protein-level
connections and neighbourhood proteins, which was approximately 50-times that
reported in the STRING database (Snel et al., 2000). Convergent protein networks
included synaptic transmission, mitochondrial/metabolic processes and Wnt signaling.
Further investigation revealed that a subset of ASD genes regulates mitochondrial
function. We also determined that *de novo* missense variants in synaptic or poorly
characterized ASD-risk genes disrupt PPIs that lead to synaptic deficits. The shared
ASD-risk gene network revealed an enrichment of an additional 112 ASD risk genes,
and is enriched with ASD-associated brain cell types (Feliciano et al., 2019; Ruzzo et
al., 2019; Sanders et al., 2015; Satterstrom et al., 2020; Yuen et al., 2017). We also
found that individuals with variants within the 41-risk genes, with a high degree of
shared interactions, had similar adaptive behavior scores using human clinical data
from the MSSNG database (Trost et al., 2020; Yuen et al., 2017).

Taken together, we demonstrate that neuron-specific PPI networks provide a
scalable approach to reveal individual and convergent disease mechanisms in ASD.
This PPI network resource and screening system can be applied more broadly to
additional autism risk genes to identify disease mechanisms that are not captured with
current approaches.

Results

Development of a neuronal proximity-based proteomic system to identify PPI networks

We used mouse cortical neurons and glia co-cultures infected with neuron-specific human Synapsin1 promoter-driven lentiviral constructs expressing BioID2 fusion proteins (pLV-hSyn-tGFP-P2A-POI-13xLinker-BioID2-3xFLAG) (Figure 1A and Figure S1A). A 13x Gly-Ser linker sequence attached to the proteins-of-interest (POIs) increased the range of biotinylation to include direct/indirect interactors and proteins in close proximity. Embryonic age 16-17 (E16-17) mouse cortical neurons were infected at days *in vitro* (DIV) 14 (Figure 1A). Biotin was added at DIV17 for 18 hours, followed by processing for mass spectrometry. We used a Luciferase-P2A-BioID2-3xFLAG construct as a negative control (Figure S1B). To promote high efficiency infections, we optimized lentiviral production for small and large risk genes (Figure 1B).

To validate the BioID2 screening system, we used the well characterized excitatory synapse protein DLG4 (PSD95). Neurons expressing PSD95-BioID2 displayed punctate localization of BioID2-3xFLAG fusion proteins and biotinylated proteins around the dendrites, suggesting appropriate synaptic expression and biotinylation (Figure S1C). Synaptic punctate localization was also seen for TAOK2β-BioID2 as expected (Figure S1C) (Richter et al., 2019). The Luciferase-P2A-BioID2 control showed non-specific localization and biotinylation throughout the neuron as expected (Figure S1C). We identified 74 proteins that interact with PSD95, and Reactome pathway enrichment revealed neurotransmitter receptors and glutamatergic synapses (Figure S1D and Table S1). Comparison of our PSD95 PPI network with previously published data revealed 25 shared proteins (Fernández et al., 2009; Uezu et al., 2016) (Figure S1E), highlighting that our system captures relevant networks, acknowledging there are differences between methods and approaches.

Cortical neurons are a major cell type associated with ASD (Satterstrom et al., 2020); however, BioID approaches have typically been done primarily in cell lines (Go et al., 2021). To determine the importance of using neurons for the screen, we performed

BioID2 in HEK293 cells (Figure S2 and Table S2). The PSD95 PPI network from
HEK293 cells showed an absence of synaptic pathways (Figure S2A). Furthermore,
BioID2 of six ASD risk proteins in HEK293 cells revealed a significant loss of protein
interactions localized in neuron-specific compartments, and large differences in the PPI
network compared to mouse neurons (Figure S2B-S2G, and Table S3).

To further validate the neuron-specific BioID2 screening system, we targeted
compartment proteins (Fazal et al., 2019), including the microtubules (MAP2C),
endoplasmic reticulum network (CANX), plasma membrane (PDGFR transmembrane
domain), trans-Golgi apparatus (TGOLN), the presynaptic terminal (SNCA), and
nucleus (MECP2). Compartment analysis of each PPI network revealed enrichment of
the compartments expected for each bait protein (Figure S3, Figure S4D, Table S1 and
Table S4). SNCA did not have a strong enrichment of presynaptic compartments;
however, it did identify pathways involving axons, growth cones and the synapse
(Figure S3E). BioID2 of MECP2, a nuclear protein, indicated localization to the nucleus
(Figure S4A) and interaction with proteins enriched in nucleus-specific pathways (Figure
S4B). The MECP2 PPI network was different in mouse neurons compared to HEK293
cells (Figure S4C and S4D), suggesting that mouse neurons have differing MECP2
interactions localized to the nucleus. Further, the PPI network of MECP2 did not include
some of the known protein interactions in mouse neurons (e.g., ATRX, CREB1, SIN3A,
NCor, and TET1), possibly due to the presence of highly biotinylated endogenous
nuclear proteins or experimental artifacts. The enrichment of proteins specific to each
compartment analyzed provides validation that the BioID2 screen in mouse cortical
neurons can provide relevant PPI networks.

Identification of a shared PPI network map and common pathways of 41 ASD-risk genes in mouse cortical neurons

To develop a shared PPI network map for ASD risk genes, we selected 41 ASD-
risk genes that encode proteins with a range of molecular functions (Figure 1B). These
genes were chosen from a combined list of ASD-risk genes from the SPARK, SFARI

category 1, 2, and syndromic gene lists and previous sequencing studies (Feliciano et
al., 2019; Ruzzo et al., 2019; Sanders et al., 2015; Satterstrom et al., 2020; Wilfert et
al., 2021; Yuen et al., 2017). The final list was filtered for size limitations of the
lentivirus, however we included some large genes (>4kb) such as SHANK3 and
SYNGAP1 by optimizing lentivirus production. All genes chosen for the screen have a
cytoplasmic function or localization (Satterstrom et al., 2020). For each gene, the
human cDNA was cloned into a BioID2 lentiviral backbone and protein expression was
confirmed (Figure S5A). We identified the individual PPI networks and enriched
Reactome pathways, biological processes and cellular compartments for each gene in
mouse cortical neurons (Table S1 and Table S5). Validation of BioID protein hits verified
the interactions between Taok2 and Fbxo41, similar to another publication (Pennemann
et al., 2021) (Figure S5B). We also verified interactions between Fmrp and Stxbp1,
Taok2 and Syngap1, and the mitochondrial citrate synthase protein (Cs) and Syngap1
(Figure S5C-S5E). Furthermore, we found that Psd95 and Taok2β, and to a lesser
degree, Shank2 and Taok2β, co-localized in mouse cortical neurons (Figure S5F).

The 41 ASD-risk gene PPI network consisted of 1109 proteins (41 ASD bait
proteins and 1068 prey proteins) and 2349 connections. Every ASD bait protein shared
at least 4 shared prey proteins with one other ASD bait protein (Figure 2A). Reciprocal
identification was observed between DLG4 and CDKL5, GRIA1, GRIA2, or SYNGAP1
and between GRIA1 and GRIA2. BioID2 of CDKL5, DLG4, LRRC4C, SYNGAP1, and
TAOK2 identified the most ASD bait proteins, suggesting high connectivity between a
subset of ASD bait proteins. We identified 3 groups of highly connected ASD-risk genes
from the individual PPI networks of the 41 ASD-risk genes (Group 1, 2 and 3), based on
the correlation between individual PPI networks (Figure 2B). Groups 1 and 2 showed
high connectivity between the ASD-risk genes within each group, whereas connectivity
was lower in Group 3.

We compared our results with physical interactions between the 41 ASD bait
proteins extracted from the STRING database (greater than or equal to medium
confidence, 0.4). Our neuron-generated 41 ASD-risk gene PPI network had a near 50-
fold increase in the number of connections compared to the STRING-db network

(Figure 2A and S6A). STRING is primarily derived from non-neuronal sources using gene co-expression or direct interaction data (Lage, 2014), which do not capture proteins within close proximity to the POI. However, the PPI networks we identified include both direct interacting proteins, shared interacting proteins and close proximity proteins, highlighting potential connections missed by traditional methods.

The most significant pathways in the shared 41 ASD-risk gene PPI network involve synaptic transmission (Figure 2C and Table S6). Other enriched pathways included TCA cycle and mitochondrial activity, Wnt signaling, potassium channel activity, and MAPK signaling (Figure 2C and S6B, and Table S6). The majority of the shared ASD-risk PPI network localized to specific cellular compartments including axons, dendrites and synapses (Figure S6C and Table S6), while the majority of biological processes involve synaptic signaling and organization, and protein transport (Figure S6D and Table S6). Shared pathways in the ASD-risk gene PPI network reflect the major role of synaptic dysfunction in ASD, but also highlight that less well-studied mechanisms are important in ASD pathology.

The shared PPI network map identifies the tricarboxylic acid (TCA) cycle and pyruvate metabolism as a common signaling pathway in ASD

The 41-gene shared PPI network enriched for the TCA cycle and pyruvate metabolism proteins, implicating dysregulation in mitochondrial function and cellular metabolism. This pathway has been associated with some ASD associated genes such as *Fmr1* and *Mecp2*, but it is unknown whether this extends to other ASD risk genes (Bülow et al., 2021; Licznerski et al., 2020; Shulyakova et al., 2017). Previous clinical studies have identified abnormal mitochondrial function in ASD patient lymphoblastoid cells (Frye, 2020; Rose et al., 2014, 2018; Shen et al., 2019a). The TCA cycle and respiratory electron transfer chain protein Reactome pathway was highly enriched in the shared ASD-risk gene PPI network map (adj. p-value = 3.14×10^{-12}), even without the PPI network for the mitochondrial protein ETFB (adj. p-value = 3.21×10^{-6}) (Table S6). 28 out of 41 ASD-risk genes associated with at least one TCA cycle and pyruvate

metabolism associated protein (Figure 3A). Citrate synthase (CS), which is involved in turning acetyl-CoA into citrate early in the TCA cycle, interacted with eight ASD bait proteins (ERBIN, MET, NRXN1, SHANK3, SPAST, STXBP1, SYNGAP1, TAOK2). The TCA cycle and pyruvate metabolism are essential for proper cellular respiration (adj p-value = 1.64×10^{-10}) (Table S6). Therefore, we investigated this finding using a gene not previously associated with mitochondrial and metabolic processes, *TAOK2*, a gene in the 16p11.2 deletion/duplication region associated with ASD (Calderon de Anda et al., 2012; Richter et al., 2019; Ultanir et al., 2014; Yadav et al., 2017). We measured cellular respiration using live-cell metabolic assays in mouse cortical neurons. *Taok2* heterozygous knockout (Het) mouse cortical neurons showed an increase in maximal respiration, proton leak, non-mitochondrial respiration, spare respiratory capacity, and a decrease in ATP coupling efficiency (Figure 3B and 3C, and Figure S7A-S7D) compared to wildtype (WT) neurons. These changes indicated less functional mitochondria, which was corroborated by proteomic analysis of post-synaptic density (PSD) fractions from *Taok2* WT and homozygous knockout (KO) mouse cortices (Figure S7E). *Taok2* KO mice PSD fractions had downregulation of proteins involved in synaptic function and activity, and ETC complex proteins (Figure S7F and Table S7). Analysis at the transcriptome level also revealed reduced mRNA levels of mitochondrial membrane proteins in *Taok2* KO mouse cortices (Figure S7G and S7H, and Table S7), coinciding with the reduced protein levels of mitochondrial proteins (Figure S7F). Further investigation revealed that *Taok2* Het and KO neurons have a reduction in active Teramethylrhodamine (TMRM)-stained mitochondria (Figure 3D and 3E, and Figure S7I). There was also an increase in mitochondria size and number using the marker TOMM20 (Figure 3F and 3G). We examined the morphology of mitochondria *in vivo* from electron microscopy (EM) images from WT and *Taok2* KO mouse brains (Richter et al., 2019). *Taok2* KO mouse neurons had altered mitochondrial morphology with a reduction in category 1 and 3 mitochondria (typical morphology), and an increase in category 2 mitochondria at synapses (Figure 3H and 3I) (Da Costa et al., 2018). Category 2 mitochondria indicate enlarged non-contiguous mitochondrial cristae, which can reduce oxidative phosphorylation and cause aberrant mitochondrial protein translation (John et al., 2005; Schmidt et al., 2010). We extended these studies to

human induced pluripotent stem cell (iPSC)-derived NGN2-neurons. We generated isogenic *TAOK2* KO and heterozygous knock-in *TAOK2 A135P* iPSC lines. A135P is a *de novo* missense variant which renders TAOK2 as kinase dead (Richter et al., 2019). We generated human neurons through direct differentiation of iPSCs via NGN2 overexpression and found altered cellular respiration in *TAOK2* KO neurons, similar to mouse neurons, and increases in the spare respiratory capacity of *TAOK2* KO and *A135P* neurons (Figure S7J-S7L). *TAOK2* KO and *A135P* human neurons transfected with Mito7-DsRed also displayed an increase in mitochondrial puncta size, suggesting an increase in the number or size of the mitochondria (Figure S7M and S7N). To determine if these changes were due to long-term deficits caused by loss of TAOK2 function, we used acute *Taok2* shRNA knock-down and found a decrease in mitochondrial membrane potential (Figure S8A-S8C). Taken together, using *TAOK2* as a validation gene, we determined that mouse and human models with disruption of *TAOK2* have altered cellular respiration, likely caused by altered activity, size and number of mitochondria.

To determine if other ASD risk genes converging on the mitochondrial/metabolic network regulate cellular respiration, we used the CRISPR/Cas9 system to knock out *Syngap1*, *Spast*, and *Taok2*. We also targeted *Etfb* and *Rheb*, both ASD risk genes that localize to the mitochondrion or regulate neuronal mitochondrial function (Yang et al., 2021). Combined gRNAs against BFP and Luciferase were used as a negative control (Hart et al., 2015; Richardson et al., 2016), and we used and validated 1-3 gRNAs targeting each gene (Figure S8D and S8E). CRISPR/Cas9 knockout of Etfb showed increased basal and maximal respiration, proton leakage, and no change in ATP synthase-dependent cellular respiration (Figure 3J and 3K, and Figure S8F and S8G). Mouse neurons with CRISPR knockout of *Taok2*, *Syngap1*, and *Rheb* also showed changes in many aspects of cellular respiration (Figure 3J and 3K, and Figure S8F-S8I). CRISPR KO of *Spast* did not cause changes in cellular respiration; however, Spast could regulate other aspects of the TCA cycle and pyruvate metabolism. The increase in basal respiration in *Taok2*, *Syngap1*, and *Etfb* KO neurons (Figure S8F) may be indicative of cellular respiration not yet reaching homeostasis (Ruggiero et al., 2021). We further validated the BioID data by using CRISPR/Cas9 to knockout Ap2s1, Gria1,

Ppp1r9b, and Ppp2r5d; genes which did not have mitochondrial or metabolic proteins in

their PPI network. Seahorse assays on these genes showed no changes in cellular

respiration (Figure S8J-S8L). These findings suggest that a subset of ASD risk genes

regulate cellular respiration in neurons, and highlight the relevance of TCA cycle and

pyruvate metabolism pathways in ASD.

PPI networks identify differences in signaling between missense variants in ASD

risk genes

Next, we hypothesized that PPI networks could be used to identify differences in

pathogenic mechanisms of *de novo* ASD-linked missense variants, in particular variants

of unknown significance (VUS). Missense variants have been suggested to impact

protein stability and protein-protein interaction networks (Chen et al., 2020); however,

these data were imputed from databases using primarily non-neuronal datasets. Due to

the strong link between synaptic functional deficits and ASD pathophysiology, we chose

two known synaptic genes (TAOK2β, the synapse-specific isoform of TAOK2, and

GRIA1) and a less well-characterized risk gene with no known cellular localization

(PPP2R5D) (Figure 4A-4C, Table S8, and Table S9).

We determined the change in the TAOK2β PPI network due to the A135P *de*

novo missense variant. The TAOK2β A135P PPI network had reduced numbers of

proteins associated with the synaptic compartment, and increased dendritic and

ribosomal proteins (Figure 4D, Table S8 and Table S9). The latter changes may be due

to the loss of PPIs in dendritic spines where TAOK2β localizes, causing an increase in

dendritic and ribosome translation complex PPIs for the TAOK2β A135P (Figure 4E),

combined with decreased expression of the A135P mutant (Figure S9A). To corroborate

the possible synaptic deficits caused by the A135P variant, we performed patch-clamp

electrophysiology on isogenic iPSC-derived NGN2-neurons (Figure S9B-S9C)

(Deneault et al., 2018, 2019; Zhang et al., 2013). *TAOK2* KO and *TAOK2* A135P

neurons had decreases in frequency and amplitude of spontaneous excitatory post-

synaptic currents (sEPSCs) (Figure 4F and 4G). This data coincides with the reduced

biotinylation of synaptic proteins for TAOK2 A135P-BioID2 (Figure 4D). The lack of

change in the intrinsic firing properties or Synapsin1-positive punctae in *TAOK2* A135P

neurons, as opposed to the *TAOK2* KO neurons (Figure S9B-S9E), suggest that the

shift in interaction of proteins and localization for the heterozygous A135P line has

dissimilar phenotypes compared to the *TAOK2* KO. In fact, *TAOK2* A135P neurons

displayed increased size of Synapsin1 punctae, suggesting possible changes in

synapse structure (Figure S9D and S9E). Taken together, these data demonstrate that

changes in PPI networks can help to support functional deficits.

We also asked whether PPI networks can distinguish the impact of missense

variants based on their location within functional domains of a gene. We investigated

GRIA1 and two ASD-linked *de novo* missense variants, R208H and A636T (Geisheker

et al., 2017; Iossifov et al., 2014; de Ligt et al., 2012), located in the extra-cellular ligand

binding domain and the transmembrane domain, respectively (Fig. 4B). The *GRIA1*

variants showed differential effects in their enriched cellular compartments (Figure 4H)

and the number of shared interacting proteins compared to wildtype (Figure 4I). GRIA1

R208H had a loss of proteins localizing to the AMPA receptor and post-synaptic density.

GRIA1 A636T had a less severe impact, with small increases in compartment-specific

PPIs, including membrane junction and ER proteins (Figure 4H, Table S8, and Table

S9). There was no difference between GluA1 variant expression (Figure S9F) or in

GluA2 subunit expression (Figure S9G). We identified electrophysiological changes that

coincide with the changes in PPI networks, revealing a trend towards decreased sEPSC

frequency in neurons expressing the R208H variant, but not the A636T variant (Figure

4J and 4K). Although the A636T mutant had no change in sEPSCs, we did observe

large sEPSC bursts (Figure 4J), which may be indicative of altered trafficking of AMPA

receptors (Pick and Ziff, 2018; Schwenk et al., 2019). These data demonstrate the

potential use of PPI networks to identify functional differences in missense variants for

receptor proteins.

Finally, we used BioID2 to test missense variants in the ASD risk gene

PPP2R5D, a regulatory subunit of phosphatase-2A (Shang et al., 2016). We selected

three *de novo* variants, P53S, E198K, and E420K (Houge et al., 2015; Shang et al.,

2016). The PPI networks for the variants had both common and dissimilar effects
(Figure 4C, 4L, and S9H), with all three variants reducing PPIs with synaptic and
dendritic proteins (Figure 4L). All of the variants caused a loss and gain of diverse
interactions (Figure S9H), with no change in expression levels (Figure S9I).
Interestingly, both the E198K and the E420K variants gained trans-Golgi compartment
proteins (Figure 4L, Table S8, and Table S9), suggesting altered localization to the
Golgi apparatus. Previous studies have linked overactive AKT signaling to the
PPP2R5D E420K variant (Papke et al., 2021); however, we found no difference in
phospho-AKT levels (Figure S9J). To probe the E198K and E420K variants, we stained
infected mouse neurons for a trans-Golgi apparatus-specific protein, GALNT2 (Figure
4M). In WT neurons, the GALTN2 expression is primarily located in the cell body
(Figure 4M). However, GALTN2 staining in the E198K variant was smaller and spread
throughout the primary dendrite (Figure 4L and 4M, and Table S8 and S9). Neurons
expressing the E410K variants showed no change in size; however, GALTN2
expression was also spread into the apical dendrites (Figure 4L and 4M, and Table S8
and S9). As a control, we measured cell size and found that overexpression of
PPP2R5D variants had no effect (Figure S9K). These data suggest that ASD-linked
PPP2R5D variants may impart pathogenicity by altering Golgi-related functions in
neurons.

The 41 ASD-risk gene PPI network map enriches for additional ASD risk genes,
human disease cell types, and correlates with human behavioral phenotypes from
clinical datasets

To demonstrate the utility of the 41 ASD-risk gene PPI network map resource,
we used enrichment analysis to determine relevance to human ASD. We found a
significant enrichment of 112 additional ASD-risk genes (Fisher's Exact test p =
2.69×10^{-30}, OR = 3.45), highlighting the functional connectivity between ASD-risk genes
at the protein level (Figure 5A). We also found that gene lists from individual sequencing
studies were enriched in our PPI network, especially when examining cytoplasmic (non-
nuclear) proteins (Figure S10A). Gene lists with only nuclear proteins were not enriched

(Figure S10B), providing evidence for less interaction between proteins in the nucleus
and the cytoplasm. Of the 153 ASD-risk proteins in the network, 69 are interacting with
2 or more ASD bait proteins.

While the PPI network from 41 ASD-risk genes was generated using human
genes, it was obtained using mouse cortical neurons; therefore, it is unknown whether
this network map is applicable to human brain cell types or differentially expressed
genes (DEGs) implicated in ASD pathology. We examined the enrichment of specific
cell types based on their single cell RNA-sequencing profiles (Nowakowski et al., 2017;
Velmeshev et al., 2019). We found the 41 ASD-risk gene PPI network map enriches for
excitatory and inhibitory neuron cell types, and neural progenitor cells, astrocytes and
microglia (Figure S10C), which have been associated with ASD (Parikshak et al., 2013;
Tang et al., 2014; Velmeshev et al., 2019; Willsey et al., 2013; Xu et al., 2020). When
examining the ASD-specific DEGs of cell types from human post-mortem brains
(Velmeshev et al., 2019), the shared PPI network was enriched for DEGs in layer 2/3
and 4 neurons, parvalbumin and VIP interneurons, and protoplasmic astrocytes (Figure
5B). The enrichment of ASD DEGs of specific cell types suggests the human disease
relevance of the 41 ASD-risk gene PPI network map.

Finally, we hypothesized a potential relationship between highly connected
genes and human ASD behavioral phenotypes. We obtained clinical data of individuals
with rare variants in the 41 ASD-risk genes from the MSSNG database and grouped
them based on the correlation of individual ASD-risk gene PPI networks (Figure 2B and
5C) (Yuen et al., 2017). We obtained the adaptive behavior and socialization scores
from between 112-879 individuals, which was dependant on data availability, and the
number of individuals per group with at least one rare missense/splicing/LoF variant in
the 41 ASD-risk genes (data-explorer.mss.ng). Remarkably, we found that individuals
with missense variants in Group 1 genes had lower adaptive behavior standard scores
compared to Groups 2 or 3, suggesting that missense variants impact Group 1 genes in
regards to adaptive behavior (Figure 5D and Figure S10D). However, individuals with
variants impacting mRNA splicing in Groups 1 had significantly higher standard
adaptive behavior and socialization scores compared to Group 2 or 3 (Figure 5E and

5F). Interestingly, the NRXN1 gene that is part of group 2 has been found to have
alternative splicing in individuals with neuropsychiatric disorders (Flaherty et al., 2019).
No significant differences were seen between individuals with frame shift or stop gain
variants (Figure S10E and S10F). Group 1 genes were found to have the largest
enrichment of ASD-risk genes (Figure S10G), suggesting that the highly interconnected
PPI networks and shared pathways for this group of genes may be linked to the clinical
phenotypes.

Discussion and Conclusion

A long-standing question in ASD research is whether there are convergent
signaling mechanisms between different risk genes. Specific disease cell types or
signaling pathways have been proposed as convergent mechanisms, but these data are
based on RNA expression, not protein levels (Chang et al., 2015; O'Roak et al., 2012b;
Ramaswami et al., 2020; Velmeshev et al., 2019; Voineagu et al., 2011). To address
this, we devised a neuron-specific proteomic screen to study 41 ASD-risk genes, and
identified links between risk genes and convergent pathways. In addition, PPI network
mapping could predict the impact of disease-associated missense variants. Finally, the
PPI network map revealed an enrichment of additional ASD risk genes and cell types
implicated in ASD pathology. Cross-referencing the PPI network with human clinical
data showed correlation between highly interacting ASD-risk genes and ASD diagnostic
behavioral severity.

A main finding was the identification of shared interactions with the TCA cycle
and mitochondrial proteins in 28 ASD-risk genes, where dysfunction is indirectly
associated with neurodevelopmental disorders (Iwata et al., 2020) (Figure 3A). Clinical
studies have found mitochondrial and metabolic dysfunction or changes in metabolites
in primary lymphocytes or brain tissue in individuals with ASD, but whether this is direct
or indirect is not known (Anitha et al., 2012; Frye and Rossignol, 2011; Frye et al., 2013;
Kurochkin et al., 2019; Rose et al., 2014, 2018; Rossignol and Frye, 2012; Wang et al.,
2016b). Some ASD-associated syndromic disorders, co-morbid disorders and genetic

ASD models have shown deficits in mitochondrial and metabolic processes, however the specific proteins involved were unknown (Bülow et al., 2021; Ebrahimi-fakhari et al., 2016; Fernandez et al., 2019; Jagtap et al., 2019; Kanellopoulos et al., 2020; Li et al., 2019; Licznerski et al., 2020; Madison et al., 2021; Menzies et al., 2021; Shen et al., 2019b; Shulyakova et al., 2017). Our findings indicate that TCA cycle and mitochondrial activity proteins are interacting with multiple ASD-risk genes. In addition, we identified two ASD-risk genes (*RHEB* and *BCKDK*) that were previously implicated in metabolic processes (Heinemann-Yerushalmi et al., 2021; Yang et al., 2021). This highlights that our screen can identify relevant protein interactions and may indicate a more direct connection between mitochondrial/metabolic processes and some ASD genes (Jagtap et al., 2019; Madison et al., 2021).

Our CRISPR/Cas9 KO studies revealed that multiple ASD-risk genes are important for proper cellular respiration (Figure 3A), suggesting that cross-talk may exist between ASD-risk genes and TCA cycle function. Deficits in the TCA cycle can cause overreliance on glutaminolysis to produce energy and cause a decrease in synaptic vesicle glutamate levels, possibly explaining the synaptic transmission deficits caused by *Syngap1* and *Taok2* KO (Divakaruni et al., 2018; Fendt and Verstreken, 2017; Namba et al., 2021).

ASD-associated *de novo* missense mutations are enriched in hub genes of known protein interaction networks (Chen et al., 2018, 2020). However, few studies have used proximity-labeling to study disease-relevant mutations on PPI networks (Chou et al., 2018; Pintacuda et al., 2021; Tracy et al., 2021). Our screen provides functional evidence of the impact of *de novo* missense variants on the PPI network of three ASD-risk genes as examples. PPP2R5D is the regulatory subunit of PP2A, which was previously associated with Golgi assembly through the beta subunit (Lowe et al., 2000; Schmitz et al., 2010). We found PPP2R5D variants disrupted the Golgi apparatus morphology and localization, highlighting a potential pathogenic mechanism in neurons. These studies are an example of how PPI networks can be used to discover unknown disease mechanisms. Using neuron-specific PPI networks allows the use of a relevant

cell type, and the potential to study variants of unknown significance to provide
important information on severity.

The enrichment of 112 ASD risk genes in the shared ASD-risk gene PPI network
map, and the enrichment of the network in ASD-associated cell types further
emphasizes the interconnectedness of ASD risk proteins. Mid-fetal deep cortical
projection neurons and superficial cortical glutamatergic neurons are enriched for ASD-
risk genes and are associated with autism pathology (Parikshak et al., 2013; Willsey et
al., 2013). The ASD-shared PPI network was highly enriched for genes expressed in
excitatory and inhibitory neurons, and for DEGs in individuals with ASD specific to
layers 2/3 excitatory neurons and VIP interneurons. Future studies will need to
distinguish which ASD PPI networks are specific to each cell type, or possible
subpopulations, to understand the subtle network changes that impact disease
mechanisms.

The grouping of the 41 ASD-risk genes based on their PPI network, and
correlation to clinical behavioral scores is an emerging area. Although we focused on
missense/LoF variants, we found that the type of variant in each group of genes
influenced the average score of the individuals within the group. However, the overall
effect size of the analysis was small, suggesting that further BioID of ASD-risk genes, or
methods that consider the genetic background of individuals, is required. To work
through the complexity, it may be important to combine our analysis with other methods
of categorizing mutations (e.g., gnomAD pLI, Polyphen-2) as higher or lower impact,
which would reduce the number of people shared between groups (Adzhubei et al.,
2010; Lek et al., 2016).

In conclusion, our neuron-specific 41 ASD-risk gene PPI network map
demonstrates that interacting and close proximity proteins, and their associated
pathways and sub-cellular compartments are relevant to ASD disease pathology, which
is missing from transcriptome-based approaches. This resource containing the
individual PPI networks of 41 ASD-risk genes will be valuable for future in-depth study
of the genes, and has the potential to grow larger with PPI networks of additional risk
genes.

Limitations of the study

540   In our screen, we excluded nuclear genes due to the functional separation with cytoplasmic genes (Satterstrom et al., 2020). Therefore, we did not identify many disease-relevant convergent pathways associated with gene regulation and chromatin modification. Another limitation is the variability between BioID runs that necessitated reduced stringency in our comparisons. Therefore, functional validation is important to filter out changes caused by background noise. Finally, we used single canonical isoforms and C-terminal tagged BioID (no N-terminal tag) of each gene; therefore some PPI networks may not encompass the full scope of possible interactions in the neuron.

References

Adzhubei, I.A., Schmidt, S., Peshkin, L., Ramensky, V.E., Gerasimova, A., Bork, P., Kondrashov, A.S., and Sunyaev, S.R. (2010). A method and server for predicting damaging missense mutations. Nat. Methods *7*, 248–249. https://doi.org/10.1038/nmeth0410-248.

Afgan, E., Baker, D., Batut, B., Van Den Beek, M., Bouvier, D., Ech, M., Chilton, J., Clements, D., Coraor, N., Grüning, B.A., et al. (2018). The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res. *46*, W537–W544. https://doi.org/10.1093/nar/gky379.

Alford, S.C., Ding, Y., Simmen, T., and Campbell, R.E. (2012). Dimerization-dependent green and yellow fluorescent proteins. ACS Synth. Biol. *1*, 569–575. https://doi.org/10.1021/sb300050j.

Anitha, A., Nakamura, K., Thanseem, I., Yamada, K., Iwayama, Y., Toyota, T., Matsuzaki, H., Miyachi, T., Yamada, S., Tsujii, M., et al. (2012). Brain region-specific altered expression and association of mitochondria-related genes in autism. Mol. Autism *3*, 1–12. https://doi.org/10.1186/2040-2392-3-12.

Bader, G.D., Isserlin, R., Merico, D., and Voisin, V. (2014). Enrichment Map - a Cytoscape app to visualize and explore OMICs pathway enrichment results. F1000Research *3*, 1–8. https://doi.org/10.12688/f1000research.4536.1.

Barger, C.J., Branick, C., Chee, L., and Karpf, A.R. (2019). Pan-cancer analyses reveal genomic features of FOXM1 overexpression in cancer. Cancers (Basel). *11*, 251. https://doi.org/10.3390/cancers11020251.

Barres, B.A. (2008). The Mystery and Magic of Glia: A Perspective on Their Roles in Health and Disease. Neuron *60*, 430–440. https://doi.org/10.1016/j.neuron.2008.10.013.

Bateman, A., Martin, M.J., Orchard, S., Magrane, M., Agivetova, R., Ahmad, S., Alpi, E., Bowler-Barnett, E.H., Britto, R., Bursteinas, B., et al. (2021). UniProt: The universal protein knowledgebase in 2021. Nucleic Acids Res. *49*, D480–D489. https://doi.org/10.1093/nar/gkaa1100.

Braganza, A., Li, J., Zeng, X., Yates, N.A., Dey, N.B., Andrews, J., Clark, J., Zamani, L., Wang, X.H., St Croix, C., et al. (2017). UBE3B is a calmodulin-regulated,

mitochondrion-associated E3 ubiquitin ligase. J. Biol. Chem. *292*, 2470–2484. https://doi.org/10.1074/jbc.M116.766824.

Braun, J.E., Huntzinger, E., Fauser, M., and Izaurralde, E. (2011). GW182 proteins directly recruit cytoplasmic deadenylase complexes to miRNA targets. Mol. Cell *44*, 120–133. https://doi.org/10.1016/j.molcel.2011.09.007.

Bülow, P., Zlatic, S.A., Wenner, P.A., Bassell, G.J., and Faundez, V. (2021). FMRP attenuates activity dependent modifications in the mitochondrial proteome. Mol. Brain *14*, 1–11. https://doi.org/10.1186/s13041-021-00783-w.

Butko, M.T., Yang, J., Geng, Y., Kim, H.J., Jeon, N.L., Shu, X., MacKey, M.R., Ellisman, M.H., Tsien, R.Y., and Lin, M.Z. (2012). Fluorescent and photo-oxidizing TimeSTAMP tags track protein fates in light and electron microscopy. Nat. Neurosci. *15*, 1742–1751. https://doi.org/10.1038/nn.3246.

Calderon de Anda, F., Rosario, A.L., Durak, O., Tran, T., Gräff, J., Meletis, K., Rei, D., Soda, T., Madabhushi, R., Ginty, D.D., et al. (2012). Autism spectrum disorder susceptibility gene TAOK2 affects basal dendrite formation in the neocortex. Nat. Neurosci. *15*, 1022–1031. https://doi.org/10.1038/nn.3141.

Casanova, E.L., Switala, A.E., Dandamudi, S., Hickman, A.R., Vandenbrink, J., Sharp, J.L., Feltus, F.A., and Casanova, M.F. (2019). Autism risk genes are evolutionarily ancient and maintain a unique feature landscape that echoes their function. Autism Res. *12*, 860–869. https://doi.org/10.1002/aur.2112.

Cederquist, G.Y., Tchieu, J., Callahan, S.J., Ramnarine, K., Ryan, S., Zhang, C., Rittenhouse, C., Zeltner, N., Chung, S.Y., Zhou, T., et al. (2020). A Multiplex Human Pluripotent Stem Cell Platform Defines Molecular and Functional Subclasses of Autism-Related Genes. Cell Stem Cell *27*, 35-49.e6. https://doi.org/10.1016/j.stem.2020.06.004.

Chang, J., Gilman, S.R., Chiang, A.H., Sanders, S.J., and Vitkup, D. (2015). Genotype to phenotype relationships in autism spectrum disorders. Nat. Neurosci. *18*, 191–198. https://doi.org/10.1038/nn.3907.

Chen, S., Fragoza, R., Klei, L., Liu, Y., Wang, J., Roeder, K., Devlin, B., and Yu, H. (2018). An interactome perturbation framework prioritizes damaging missense mutations for developmental disorders. Nat. Genet. *50*, 1032–1040.

https://doi.org/10.1038/s41588-018-0130-z.

Chen, S., Wang, J., Cicek, E., Roeder, K., Yu, H., and Devlin, B. (2020). De novo missense variants disrupting protein-protein interactions affect risk for autism through gene co-expression and protein networks in neuronal cell types. Mol. Autism *11*, 1–16. https://doi.org/10.1186/s13229-020-00386-7.

Chou, C., Zhang, Y., Umoh, M.E., Vaughan, S.W., Lorenzini, I., Liu, F., Sayegh, M., Donlin-asp, P.G., Chen, Y.H., Duong, D.M., et al. (2018). TDP-43 pathology disrupts nuclear pore complexes and nucleocytoplasmic transport in ALS/FTD. Nat. Neurosci. *21*. https://doi.org/10.1038/s41593-017-0047-3.

Corominas, R., Yang, X., Lin, G.N., Kang, S., Shen, Y., Ghamsari, L., Broly, M., Rodriguez, M., Tam, S., Trigg, S.A., et al. (2014). Protein interaction network of alternatively spliced isoforms from brain links genetic risk factors for autism. Nat. Commun. *5*, 1–12. https://doi.org/10.1038/ncomms4650.

Da Costa, R.M., Karmirian, K., and Rehen, S.K. (2018). Deformation of mitochondrial cristae in human neural progenitor cells exposed to valproic acid. An. Acad. Bras. Cienc. *90*, 2223–2232. https://doi.org/10.1590/0001-3765201820170762.

Cox, J., and Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat. Biotechnol. *26*, 1367–1372. https://doi.org/10.1038/nbt.1511.

Cummins, J.M., and Vogelstein, B. (2004). HAUSP is required for p53 destabilization. Cell Cycle *3*, 687–690. https://doi.org/10.4161/cc.3.6.924.

Daly, M.J., Patterson, N., Mesirov, J.P., Golub, T.R., Tamayo, P., and Spiegelman, B. (2003). PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat. Genet. *34*, 267–273. .

Deneault, E., White, S.H., Rodrigues, D.C., Ross, P.J., Faheem, M., Zaslavsky, K., Wang, Z., Alexandrova, R., Pellecchia, G., Wei, W., et al. (2018). Complete Disruption of Autism-Susceptibility Genes by Gene Editing Predominantly Reduces Functional Connectivity of Isogenic Human Neurons. Stem Cell Reports *11*, 1211–1225. https://doi.org/10.1016/j.stemcr.2018.10.003.

Deneault, E., Faheem, M., White, S.H., Rodrigues, D.C., Sun, S., Wei, W., Piekna, A., Thompson, T., Howe, J.L., Chalil, L., et al. (2019). CNTN5-/+or EHMT2-/+human

iPSC-derived neurons from individuals with autism develop hyperactive neuronal
networks. Elife *8*, 1–26. https://doi.org/10.7554/elife.40092.

Divakaruni, S.S., Dyke, A.M. Van, Chandra, R., Lobo, M.K., Thompson, S.M., Blanpied,
T.A., Divakaruni, S.S., Dyke, A.M. Van, Chandra, R., Legates, T.A., et al. (2018).
Long-Term Potentiation Requires a Rapid Burst of Dendritic Mitochondrial Fission
during Induction. Neuron *100*, 1–16. https://doi.org/10.1016/j.neuron.2018.09.025.

Ebrahimi-fakhari, D., Saffari, A., Wahlster, L., Nardo, A. Di, Turner, D., Lewis, T.L.,
Hoffmann, G.F., Han, M.J., Di Nardo, A., Turner, D., et al. (2016). Impaired
Mitochondrial Dynamics and Mitophagy in Neuronal Models of Tuberous Sclerosis
Complex. Cell Rep. *17*, 1053–1070. https://doi.org/10.1016/j.celrep.2016.09.054.

Fazal, F.M., Han, S., Parker, K.R., Kaewsapsak, P., Xu, J., Boettiger, A.N., Chang,
H.Y., and Ting, A.Y. (2019). Atlas of Subcellular RNA Localization Revealed by
APEX-Seq. Cell *178*, 1–18. https://doi.org/10.1016/j.cell.2019.05.027.

Feliciano, P., Zhou, X., Astrovskaya, I., Turner, T.N., Muzny, D.M., Sabo, A.,
Consortium, T.S., Gibbs, R.A., and Evan, E. (2019). Exome sequencing of 457
autism families recruited online provides evidence for novel ASD genes. Genomic
Med. *4*, https://doi.org/10.1038/s41525-019-0093-8. .

Fendt, S.M., and Verstreken, P. (2017). Neurons eat glutamate to stay alive. J. Cell Biol.
216, 863–865. https://doi.org/10.1083/jcb.201702003.

Fernandez, A., Meechan, D.W., Karpinski, B.A., Rothblat, L.A., Maynard, T.M.,
Lamantia, A., Fernandez, A., Meechan, D.W., Karpinski, B.A., Paronett, E.M., et al.
(2019). Mitochondrial Dysfunction Leads to Cortical Under- Connectivity and
Cognitive Impairment 22q11. Neuron *102*, 1127-1142.e3.
https://doi.org/10.1016/j.neuron.2019.04.013.

Fernández, E., Collins, M.O., Uren, R.T., Kopanitsa, M. V., Komiyama, N.H., Croning,
M.D.R.R., Zografos, L., Armstrong, J.D., Choudhary, J.S., Grant, S.G.N.N., et al.
(2009). Targeted tandem affinity purification of PSD-95 recovers core postsynaptic
complexes and schizophrenia susceptibility proteins. Mol. Syst. Biol. *5*.
https://doi.org/10.1038/msb.2009.27.

Flaherty, E., Zhu, S., Barretto, N., Cheng, E., Deans, P.J.M., Fernando, M.B., Schrode,
N., Francoeur, N., Antoine, A., Alganem, K., et al. (2019). Neuronal impact of

patient-specific aberrant NRXN1α splicing (Springer US).

Frye, R.E. (2020). Mitochondrial Dysfunction in Autism Spectrum Disorder: Unique

Abnormalities and Targeted Treatments. Semin. Pediatr. Neurol. *11*, 1071–9091.

https://doi.org/10.1016/j.spen.2020.100829.

Frye, R.E., and Rossignol, D.A. (2011). Mitochondrial dysfunction can connect the

diverse medical symptoms associated with autism spectrum disorders. Pediatr.

Res. *69*, 41–47. https://doi.org/10.1203/PDR.0b013e318212f16b.

Frye, R.E., Delatorre, R., Taylor, H., Slattery, J., Melnyk, S., Chowdhury, N., and

James, S.J. (2013). Redox metabolism abnormalities in autistic children associated

with mitochondrial disease. Transl. Psychiatry *3*, 1–12.

https://doi.org/10.1038/tp.2013.51.

Furlong, R.A., Narain, Y., Rankin, J., Wyttenbach, A., and Rubinsztein, D.C. (2000). α-

Synuclein overexpression promotes aggregation of mutant huntingtin. Biochem. J.

346, 577–581. https://doi.org/10.1042/0264-6021:3460577.

Gaugler, T., Klei, L., Sanders, S.J., Bodea, C. a, Goldberg, A.P., Lee, A.B., Mahajan,

M., Manaa, D., Pawitan, Y., Reichert, J., et al. (2014). Most genetic risk for autism

resides with common variation. Nat. Genet. *46*, 881–885.

https://doi.org/10.1038/ng.3039.

Geisheker, M.R., Heymann, G., Wang, T., Coe, B.P., Turner, T.N., Stessman, H.A.F.,

Hoekzema, K., Kvarnung, M., Shaw, M., Friend, K., et al. (2017). Hotspots of

missense mutation identify neurodevelopmental disorder genes and functional

domains. Nat. Neurosci. *20*, 1043–1051. https://doi.org/10.1038/nn.4589.

Glessner, J.T., Wang, K., Cai, G., Korvatska, O., Kim, C.E., Wood, S., Zhang, H., Estes,

A., Brune, C.W., Bradfield, J.P., et al. (2009). Autism genome-wide copy number

variation reveals ubiquitin and neuronal genes. Nature *459*, 569–573.

https://doi.org/10.1038/nature07953.

Go, C.D., Knight, J.D.R., Rajasekharan, A., Rathod, B., Hesketh, G.G., Abe, K.T., Youn,

J., Samavarchi-tehrani, P., Zhang, H., Zhu, L.Y., et al. (2021). A proximity-

dependent biotinylation map of a human cell. Nature *595*.

https://doi.org/10.1038/s41586-021-03592-2.

Hamdan, H., Lim, B.C., Torii, T., Joshi, A., Konning, M., Smith, C., Palmer, D.J., Ng, P.,

Leterrier, C., Oses-Prieto, J.A., et al. (2020). Mapping axon initial segment structure and function by multiplexed proximity biotinylation. Nat. Commun. *11*. https://doi.org/10.1038/s41467-019-13658-5.

Hart, T., Chandrashekhar, M., Aregger, M., Steinhart, Z., Brown, K.R., MacLeod, G., Mis, M., Zimmermann, M., Fradet-Turcotte, A., Sun, S., et al. (2015). High-Resolution CRISPR Screens Reveal Fitness Genes and Genotype-Specific Cancer Liabilities. Cell *163*, 1515–1526. https://doi.org/10.1016/j.cell.2015.11.015.

Heinemann-Yerushalmi, L., Bentovim, L., Felsenthal, N., Vinestock, R.C., Michaeli, N., Krief, S., Silberman, A., Cohen, M., Ben-Dor, S., Brenner, O., et al. (2021). BCKDK regulates the TCA cycle through PDC in the absence of PDK family during embryonic development. Dev. Cell *56*, 1182-1194.e6. https://doi.org/10.1016/j.devcel.2021.03.007.

Hiday, A.C., Edler, M.C., Salek, A.B., Morris, C.W., Thang, M., Rentz, T.J., Rose, K.L., Jones, L.M., and Baucum, A.J. (2017). Mechanisms and Consequences of Dopamine Depletion-Induced Attenuation of the Spinophilin/Neurofilament Medium Interaction. Neural Plast. *2017*. https://doi.org/10.1155/2017/4153076.

Hockemeyer, D., Soldner, F., Cook, E.G., Gao, Q., Mitalipova, M., and Jaenisch, R. (2008). A Drug-Inducible System for Direct Reprogramming of Human Somatic Cells to Pluripotency. Cell Stem Cell *3*, 346–353. https://doi.org/10.1016/j.stem.2008.08.014.

Houge, G., Haesen, D., Vissers, L.E.L.M., Mehta, S., Parker, M.J., Wright, M., Vogt, J., McKee, S., Tolmie, J.L., Cordeiro, N., et al. (2015). B56δ-related protein phosphatase 2A dysfunction identified in patients with intellectual disability. J. Clin. Invest. *125*, 3051–3062. https://doi.org/10.1172/JCI79860.

Howarth, M., Takao, K., Hayashi, Y., and Ting, A.Y. (2005). Targeting quantum dots to surface proteins in living cells with biotin ligase. Proc. Natl. Acad. Sci. U. S. A. *102*, 7583–7588. https://doi.org/10.1073/pnas.0503125102.

Hung, V., Lam, S.S., Udeshi, N.D., Svinkina, T., Guzman, G., Mootha, V.K., Carr, S.A., and Ting, A.Y. (2017). Proteomic mapping of cytosol-facing outer mitochondrial and ER membranes in living human cells by proximity biotinylation. Elife *6*, e24463. .

Iossifov, I., Roak, B.J.O., Sanders, S.J., Ronemus, M., Krumm, N., Levy, D., Stessman,

H.A., Witherspoon, K.T., Vives, L., Patterson, K.E., et al. (2014). The contribution of de novo coding mutations to autism spectrum disorder. Nature *515*, 216–221. https://doi.org/10.1038/nature13908.

Iwata, R., Casimir, P., and Vanderhaeghen, P. (2020). Mitochondrial dynamics in postmitotic cells regulate neurogenesis. Science (80-.). *369*, 858–862. https://doi.org/10.1126/science.aba9760.

Jagtap, S., Thanos, J.M., Fu, T., Wang, J., Lalonde, J., Dial, T.O., Feiglin, A., Chen, J., Kohane, I., Lee, J.T., et al. (2019). Aberrant mitochondrial function in patient-derived neural cells from CDKL5 deficiency disorder and Rett syndrome. Hum. Mol. Genet. *28*, 3625–3636. https://doi.org/10.1093/hmg/ddz208.

Jin, X., Simmons, S.K., Guo, A.X., Shetty, A.S., Ko, M., Nguyen, L., Jokhi, V., Robinson, E., Oyler, P., Curry, N., et al. (2020). In vivo Perturb-Seq reveals neuronal and glial abnormalities associated with autism risk genes. Science (80-.). *370*, eaaz6063. https://doi.org/10.1126/science.aaz6063.

Johannessen, C.M., Boehm, J.S., Kim, S.Y., Thomas, S.R., Wardwell, L., Johnson, L.A., Emery, C.M., Stransky, N., Cogdill, A.P., Barretina, J., et al. (2010). COT drives resistance to RAF inhibition through MAP kinase pathway reactivation. Nature *468*, 968–972. https://doi.org/10.1038/nature09627.

John, G.B., Shang, Y., Li, L., Renken, C., Mannella, C.A., Selker, J.M.L., Rangell, L., Bennett, M.J., and Zha, J. (2005). The Mitochondrial Inner Membrane Protein Mitofilin Controls Cristae Morphology. Mol Biol Cell *16*, 1543–1554. https://doi.org/10.1091/mbc.E04.

Kanellopoulos, A.K., Mariano, V., Spinazzi, M., Fiala, A., Achsel, T., Kanellopoulos, A.K., Mariano, V., Spinazzi, M., Woo, Y.J., Mclean, C., et al. (2020). Aralar Sequesters GABA into Hyperactive Article Aralar Sequesters GABA into Hyperactive Mitochondria , Causing Social Behavior Deficits. Cell *180*, 1178-1197.e20. https://doi.org/10.1016/j.cell.2020.02.044.

Kapfhamer, D., Taylor, S., Zou, M.E., Lim, J.P., Kharazia, V., and Heberlein, U. (2013). *Taok2* controls behavioral response to ethanol in mice. Genes, Brain Behav. *12*, 87–97. https://doi.org/10.1111/j.1601-183X.2012.00834.x.

Kim, E., Ilic, N., Shrestha, Y., Zou, L., Kamburov, A., Zhu, C., Yang, X., Lubonja, R.,

Tran, N., Nguyen, C., et al. (2016). Systematic functional interrogation of rare

cancer variants identifies oncogenic alleles. Cancer Discov. *6*, 714–726.

https://doi.org/10.1158/2159-8290.CD-16-0160.

Kucera, M., Isserlin, R., Arkhangorodsky, A., and Bader, G.D. (2016). AutoAnnotate: A

Cytoscape app for summarizing networks with semantic annotations [version 1;

referees: 2 approved]. F1000Research *5*, 1–12.

https://doi.org/10.12688/F1000RESEARCH.9090.1.

Kurochkin, I., Khrameeva, E., Tkachev, A., Stepanova, V., Vanyushkina, A.,

Stekolshchikova, E., Li, Q., Zubkov, D., Shichkova, P., Halene, T., et al. (2019).

Metabolome signature of autism in the human prefrontal cortex. Commun. Biol. *2*,

1–10. https://doi.org/10.1038/s42003-019-0485-4.

Kuzmanov, U., and Emili, A. (2013). Protein-protein interaction networks: Probing

disease mechanisms using model systems. Genome Med. *5*, 1–12.

https://doi.org/10.1186/gm441.

Kwan, V., Meka, D.P., White, S.H., Hung, C.L., Holzapfel, N.T., Walker, S., Murtaza, N.,

Unda, B.K., Schwanke, B., Yuen, R.K.C.C., et al. (2016). DIXDC1 Phosphorylation

and Control of Dendritic Morphology Are Impaired by Rare Genetic Variants. Cell

Rep. *17*, 1892–1904. https://doi.org/10.1016/j.celrep.2016.10.047.

Lage, K. (2014). Protein-protein interactions and genetic diseases: The interactome.

Biochim. Biophys. Acta - Mol. Basis Dis. *1842*, 1971–1980.

https://doi.org/10.1016/j.bbadis.2014.05.028.

Lek, M., Karczewski, K.J., Minikel, E. V., Samocha, K.E., Banks, E., Fennell, T.,

O'Donnell-Luria, A.H., Ware, J.S., Hill, A.J., Cummings, B.B., et al. (2016). Analysis

of protein-coding genetic variation in 60,706 humans. Nature *536*, 285–291.

https://doi.org/10.1038/nature19057.

Li, J., Ma, Z., Shi, M., Hallmayer, J., Babu, M., Li, J., Ma, Z., Shi, M., Malty, R.H., Aoki,

H., et al. (2015). Identification of Human Neuronal Protein Complexes Reveals

Biochemical Activities and Convergent Mechanisms of Action in Autism Spectrum

Disorders. Cell Syst. *1*, 361–374. https://doi.org/10.1016/j.cels.2015.11.002.

Li, J., Ryan, S.K., Deboer, E., Cook, K., Fitzgerald, S., Lachman, H.M., Wallace, D.C.,

Goldberg, E.M., and Anderson, S.A. (2019). Mitochondrial deficits in human iPSC-

810   derived neurons from patients with 22q11.2 deletion syndrome and schizophrenia.

811   Transl. Psychiatry *9*. https://doi.org/10.1038/s41398-019-0643-y.

812 Licznerski, P., Park, H.A., Rolyan, H., Chen, R., Mnatsakanyan, N., Miranda, P.,

813   Graham, M., Wu, J., Cruz-reyes, N., Mehta, N., et al. (2020). ATP Synthase c-

814   Subunit Leak Causes Aberrant Cellular Metabolism in Fragile X Syndrome. Cell

815   *182*, 1170-1185.e9. https://doi.org/10.1016/j.cell.2020.07.008.

816 de Ligt, J., Willemsen, M.H., van Bon, B.W.M., Kleefstra, T., Yntema, H.G., Kroes, T.,

817   Vulto-van Silfhout, A.T., Koolen, D.A., de Vries, P., Gilissen, C., et al. (2012).

818   Diagnostic Exome Sequencing in Persons with Severe Intellectual Disability. N.

819   Engl. J. Med. *367*, 1921–1929. https://doi.org/10.1056/nejmoa1206524.

820 Loh, K.H., Stawski, P.S., Draycott, A.S., Udeshi, N.D., Lehrman, E.K., Wilton, D.K.,

821   Svinkina, T., Deerinck, T.J., Ellisman, M.H., Stevens, B., et al. (2016). Proteomic

822   Analysis of Unbounded Cellular Compartments : Synaptic Clefts Resource

823   Proteomic Analysis of Unbounded Cellular Compartments : Synaptic Clefts. Cell

824   *166*, 1295-1307.e21. https://doi.org/10.1016/j.cell.2016.07.041.

825 Lowe, M., Gonatas, N.K., and Warren, G. (2000). The mitotic phosphorylation cycle of

826   the cis-Golgi matrix protein GM130. J. Cell Biol. *149*, 341–356.

827   https://doi.org/10.1083/jcb.149.2.341.

828 Lu, G., Middleton, R.E., Sun, H., Naniong, M.V., Ott, C.J., Mitsiades, C.S., Wong, K.K.,

829   Bradner, J.E., and Kaelin, W.G. (2014). The myeloma drug lenalidomide promotes

830   the cereblon-dependent destruction of ikaros proteins. Science (80-.). *343*, 305–

831   309. https://doi.org/10.1126/science.1244917.

832 Madison, J.M., Duong, K., Vieux, E.F., Udeshi, N.D., Iqbal, S., Requadt, E.,

833   Fereshetian, S., Lewis, M.C., Gomes, A.S., Pierce, K.A., et al. (2021). Regulation of

834   purine metabolism connects KCTD13 to a metabolic disorder with autistic features.

835   IScience *24*, 101935. https://doi.org/10.1016/j.isci.2020.101935.

836 Małecki, J., Ho, A.Y.Y., Moen, A., Dahl, H.A., and Falnes, P. (2015). Human METTL20

837   is a mitochondrial lysine methyltransferase that targets the β subunit of electron

838   transfer flavoprotein (ETFβ) and modulates its activity. J. Biol. Chem. *290*, 423–

839   434. https://doi.org/10.1074/jbc.M114.614115.

840 Markmiller, S., Soltanieh, S., Server, K.L., Bennett, E.J., Yeo, G.W., Markmiller, S.,

Soltanieh, S., Server, K.L., Mak, R., Jin, W., et al. (2018). Context-Dependent and

Disease-Specific Diversity in Protein Interactions within Stress Granules. Cell *172*,

590–604. https://doi.org/10.1016/j.cell.2017.12.032.

Menzies, C., Naz, S., Patten, D., Lacoste, B., Alquier, T., and Bennett, B.M. (2021).

Distinct basal metabolism in three mouse models of neurodevelopmental disorders.

ENeuro *8*, 1–14. https://doi.org/10.1523/ENEURO.0292-20.2021.

Merico, D., Isserlin, R., Stueker, O., Emili, A., and Bader, G.D. (2010). Enrichment map:

A network-based method for gene-set enrichment visualization and interpretation.

PLoS One *5*, e13984. https://doi.org/10.1371/journal.pone.0013984.

Murtaza, N., Uy, J., and Singh, K.K. (2020). Emerging proteomic approaches to identify

the underlying pathophysiology of neurodevelopmental and neurodegenerative

disorders. Mol. Autism *11*, 27. https://doi.org/10.1186/s13229-020-00334-5.

Namba, T., Nardelli, J., Gressens, P., and Huttner, W.B. (2021). Perspective Metabolic

Regulation of Neocortical Expansion in Development and Evolution. Neuron *109*,

408–419. https://doi.org/10.1016/j.neuron.2020.11.014.

Nathanson, J.L., Yanagawa, Y., Obata, K., and Callaway, E.M. (2009). Preferential

labeling of inhibitory and excitatory cortical neurons by endogenous tropism of

adeno-associated virus and lentivirus vectors. nsc. NSC *161*, 441–450.

https://doi.org/10.1016/j.neuroscience.2009.03.032.

Neale, B.M., Kou, Y., Liu, L., Ma'ayan, A., Samocha, K.E., Sabo, A., Lin, C.-F., Stevens,

C., Wang, L.-S., Makarov, V., et al. (2012). Patterns and rates of exonic de novo

mutations in autism spectrum disorders. Nature *485*, 242–245.

https://doi.org/10.1038/nature11011.

Nowakowski, T.J., Bhaduri, A., Pollen, A.A., Alvarado, B., Mostajo-Radji, M.A., Di Lullo,

E., Haeussler, M., Sandoval-Espinosa, C., Liu, S.J., Velmeshev, D., et al. (2017).

Spatiotemporal gene expression trajectories reveal developmental hierarchies of

the human cortex. Science (80-.). *358*, 1318–1323.

https://doi.org/10.1126/science.aap8809.

O'Roak, B.J., Vives, L., Fu, W., Egertson, J.D., Stanaway, I.B., Phelps, I.G., Carvill, G.,

Kumar, A., Lee, C., Ankenman, K., et al. (2012a). Multiplex targeted sequencing

identifies recurrently mutated genes in autism spectrum disorders. Science (80-.).

338, 1619–1622. https://doi.org/10.1126/science.1227764.

O'Roak, B.J., Vives, L., Girirajan, S., Karakoc, E., Krumm, N., Coe, B.P., Levy, R., Ko, A., Lee, C., Smith, J.D., et al. (2012b). Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations. Nature *485*, 246–250. https://doi.org/10.1038/nature10989.

Ohno, H., Hirabayashi, S., Iizuka, T., Ohnishi, H., Fujita, T., and Hata, Y. (2002). Localization of p0071-interacting proteins, plakophilin-related armadillo-repeat protein-interacting protein (PAPIN) and ERBIN, in epithelial cells. Oncogene *21*, 7042–7049. https://doi.org/10.1038/sj.onc.1205852.

Papke, C.M., Smolen, K.A., Swingle, M.R., Cressey, L., Heng, R.A., Toporsian, M., Deng, L., Hagen, J., Shen, Y., Chung, W.K., et al. (2021). A disorder-related variant (E420K) of a PP2A-regulatory subunit (PPP2R5D) causes constitutively active AKT-mTOR signaling and uncoordinated cell growth. J. Biol. Chem. *296*, 100313. https://doi.org/10.1016/j.jbc.2021.100313.

Parikshak, N.N., Luo, R., Zhang, A., Won, H., Lowe, J.K., Chandran, V., Horvath, S., and Geschwind, D.H. (2013). Integrative functional genomic analyses implicate specific molecular pathways and circuits in autism. Cell *155*, 1008. https://doi.org/10.1016/j.cell.2013.10.031.

Pennemann, F.L., Mussabekova, A., Urban, C., Stukalov, A., Andersen, L.L., Grass, V., Lavacca, T.M., Holze, C., Oubraham, L., Benamrouche, Y., et al. (2021). Cross-species analysis of viral nucleic acid interacting proteins identifies TAOKs as innate immune regulators. Nat. Commun. *12*, 1–22. https://doi.org/10.1038/s41467-021-27192-w.

Pick, J.E., and Ziff, E.B. (2018). Regulation of AMPA receptor trafficking and exit from the endoplasmic reticulum. Mol. Cell. Neurosci. *91*, 3–9. https://doi.org/10.1016/j.mcn.2018.03.004.

Pintacuda, G., Hsu, Y.-H.H., Tsafou, K., Li, K.W., Martin, J.M., Risemann, J., Gonzalez-Lozano, M.A., Egri, S.B., Jaffe, J., Smit, A.B., et al. (2021). Interaction studies of risk proteins in human induced neurons reveal convergent biology and novel mechanisms underlying autism spectrum disorders. MedRxiv https://doi.org/10.1101/2021.10.07.21264575. .

Ramaswami, G., Won, H., Gandal, M.J., Haney, J., Wang, J.C., Wong, C.C.Y., Sun, W., Prabhakar, S., Mill, J., and Geschwind, D.H. (2020). Integrative genomics identifies a convergent molecular subtype that links epigenomic with transcriptomic differences in autism. Nat. Commun. *11*, 1–14. https://doi.org/10.1038/s41467-020-18526-1.

Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013). Genome engineering using the CRISPR-Cas9 system. Nat. Protoc. *8*, 2281–2308. https://doi.org/10.1038/nprot.2013.143.

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J. (2019). G:Profiler: A web server for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic Acids Res. *47*, W191–W198. https://doi.org/10.1093/nar/gkz369.

Richards, A.L., Eckhardt, M., and Krogan, N.J. (2021). Mass spectrometry-based protein–protein interaction networks for the study of human diseases. Mol. Syst. Biol. *17*, 1–18. https://doi.org/10.15252/msb.20188792.

Richardson, C.D., Ray, G.J., DeWitt, M.A., Curie, G.L., and Corn, J.E. (2016). Enhancing homology-directed genome editing by catalytically active and inactive CRISPR-Cas9 using asymmetric donor DNA. Nat. Biotechnol. *34*, 339–344. https://doi.org/10.1038/nbt.3481.

Richter, M., Murtaza, N., Scharrenberg, R., White, S.H., Johanns, O., Walker, S., Yuen, R.K.C.C., Schwanke, B., Bedürftig, B., Henis, M., et al. (2019). Altered TAOK2 activity causes autism-related neurodevelopmental and cognitive abnormalities through RhoA signaling. Mol. Psychiatry *24*, 1329–1350. https://doi.org/10.1038/s41380-018-0025-5.

Robinson, E.B., St Pourcain, B., Anttila, V., Kosmicki, J.A., Bulik-Sullivan, B., Grove, J., Maller, J., Samocha, K.E., Sanders, S.J., Ripke, S., et al. (2016). Genetic risk for autism spectrum disorders and neuropsychiatric variation in the general population. Nat. Genet. *48*, 552–555. https://doi.org/10.1038/ng.3529.

Rose, S., Frye, R.E., Slattery, J., Wynne, R., Tippett, M., Melnyk, S., and James, S.J. (2014). Oxidative stress induces mitochondrial dysfunction in a subset of autistic lymphoblastoid cell lines. Transl. Psychiatry *4*, 1–8.

https://doi.org/10.1038/tp.2014.15.

Rose, S., Bennuri, S.C., Davis, J.E., Wynne, R., Slattery, J.C., Tippett, M., Delhey, L.,

Melnyk, S., Kahler, S.G., MacFabe, D.F., et al. (2018). Butyrate enhances

mitochondrial function during oxidative stress in cell lines from boys with autism.

Transl. Psychiatry *8*. https://doi.org/10.1038/s41398-017-0089-z.

Rossignol, D.A., and Frye, R.E. (2012). Mitochondrial dysfunction in autism spectrum

disorders: A systematic review and meta-analysis. Mol. Psychiatry *17*, 290–314.

https://doi.org/10.1038/mp.2010.136.

De Rubeis, S., He, X., Goldberg, A.P., Poultney, C.S., Samocha, K., Ercument Cicek,

A., Kou, Y., Liu, L., Fromer, M., Walker, S., et al. (2014). Synaptic, transcriptional

and chromatin genes disrupted in autism. Nature *515*, 209–215.

https://doi.org/10.1038/nature13772.

Ruggiero, A., Katsenelson, M., and Slutsky, I. (2021). Trends in Mitochondria : new

players in homeostatic regulation of fi ring rate set points. Trends Neurosci. *44*,

605–618. https://doi.org/10.1016/j.tins.2021.03.002.

Ruzzo, E.K., Pe, L., Jung, J., Prober, D.A., Geschwind, D.H., Wall, D.P., Jung, J.,

Wang, L., and Kashef-haghighi, D. (2019). Inherited and De Novo Genetic Risk for

Autism Article Inherited and De Novo Genetic Risk for Autism Impacts Shared

Networks. Cell *178*, 850–866. https://doi.org/10.1016/j.cell.2019.07.015.

Sakai, Y., Shaw, C.A., Dawson, B.C., Dugas, D. V, Al-, Z., Hill, D.E., Zoghbi, H.Y., Al-

Mohtaseb, Z., Hill, D.E., Zoghbi, H.Y., et al. (2011). Protein interactome reveals

converging molecular pathways among autism disorders. Sci. Transl. Med. *3*,

86ra49. https://doi.org/10.1126/scitranslmed.3002166.

Samocha, K.E., Robinson, E.B., Sanders, S.J., Stevens, C., Sabo, A., McGrath, L.M.,

Kosmicki, J.A., Rehnström, K., Mallick, S., Kirby, A., et al. (2014). A framework for

the interpretation of de novo mutation in human disease. Nat. Genet. *46*, 944–950.

https://doi.org/10.1038/ng.3050.

Sanders, S.J., He, X., Willsey, A.J., Ercan-Sencicek, A.G., Samocha, K.E., Cicek, A.E.,

Murtha, M.T., Bal, V.H., Bishop, S.L., Dong, S., et al. (2015). Insights into Autism

Spectrum Disorder Genomic Architecture and Biology from 71 Risk Loci. Neuron

87, 1215–1233. https://doi.org/10.1016/j.neuron.2015.09.016.

Satterstrom, F.K., Kosmicki, J.A., Wang, J., Roeder, K., Daly, M.J., Buxbaum, J.D.,

Breen, M.S., De Rubeis, S., An, J.Y., Peng, M., et al. (2020). Large-Scale Exome

Sequencing Study Implicates Both Developmental and Functional Changes in the

Neurobiology of Autism. Cell *180*, 568-584.e23.

https://doi.org/10.1016/j.cell.2019.12.036.

Schmidt, O., Pfanner, N., and Meisinger, C. (2010). Mitochondrial protein import: From

proteomics to functional mechanisms. Nat. Rev. Mol. Cell Biol. *11*, 655–667.

https://doi.org/10.1038/nrm2959.

Schmitz, M.H.A., Held, M., Janssens, V., Hutchins, J.R.A., Hudecz, O., Ivanova, E.,

Goris, J., Trinkle-Mulcahy, L., Lamond, A.I., Poser, I., et al. (2010). Live-cell

imaging RNAi screen identifies PP2A-B55α and importin-β 21 as key mitotic exit

regulators in human cells. Nat. Cell Biol. *12*, 886–893.

https://doi.org/10.1038/ncb2092.

Schwenk, J., Boudkkazi, S., Kocylowski, M.K., Brechet, A., Zolles, G., Bus, T., Costa,

K., Kollewe, A., Jordan, J., Bank, J., et al. (2019). An ER Assembly Line of AMPA-

Receptors Controls Excitatory Neurotransmission and Its Plasticity. Neuron *104*,

680-692.e9. https://doi.org/10.1016/j.neuron.2019.08.033.

Seeling, J.M., Miller, J.R., Gil, R., Moon, R.T., White, R., and Virshup, D.M. (1999).

Regulation of β-catenin signaling by the B56 subunit of protein phosphatase 2A.

Science (80-.). *283*, 2089–2091. https://doi.org/10.1126/science.283.5410.2089.

Shang, L., Henderson, L.B., Cho, M.T., Petrey, D.S., Fong, C.T., Haude, K.M., Shur, N.,

Lundberg, J., Hauser, N., Carmichael, J., et al. (2016). De novo missense variants

in PPP2R5D are associated with intellectual disability, macrocephaly, hypotonia,

and autism. Neurogenetics *17*, 43–49. https://doi.org/10.1007/s10048-015-0466-9.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,

Schwikowski, B., and Ideker, T. (2003). Cytoscape: A Software Environment for

Integrated Models. Genome Res. *13*, 2498–2504.

https://doi.org/10.1101/gr.1239303.metabolite.

Sharma, K., Schmitt, S., Bergner, C.G., Tyanova, S., Kannaiyan, N., Manrique-Hoyos,

N., Kongi, K., Cantuti, L., Hanisch, U.K., Philips, M.A., et al. (2015). Cell type- and

brain region-resolved mouse brain proteome. Nat. Neurosci. *18*, 1819–1831.

https://doi.org/10.1038/nn.4160.

Shen, L., Feng, C., Zhang, K., Chen, Y., and Gao, Y. (2019a). Proteomics Study of
Peripheral Blood Mononuclear Cells (PBMCs) in Autistic Children. Front. Cell.
Neurosci. *13*, 1–16. https://doi.org/10.3389/fncel.2019.00105.

Shen, M., Wang, F., Li, M., Sah, N., Stockton, M.E., Tidei, J.J., Gao, Y., Korabelnikov,
T., Kannan, S., Vevea, J.D., et al. (2019b). Reduced mitochondrial fusion and
Huntingtin levels contribute to impaired dendritic maturation and behavioral deficits
in Fmr1-mutant mice. Nat. Neurosci. *22*, 386–400. https://doi.org/10.1038/s41593-
019-0338-y.

Shulyakova, N., Andreazza, A.C., Mills, L.R., and Eubanks, J.H. (2017). Mitochondrial
dysfunction in the pathogenesis of rett syndrome: Implications for mitochondria-
targeted therapies. Front. Cell. Neurosci. *11*, 1–9.
https://doi.org/10.3389/fncel.2017.00058.

Snel, B., Lehmann, G., Bork, P., and Huynen, M.A. (2000). String: A web-server to
retrieve and display the repeatedly occurring neighbourhood of a gene. Nucleic
Acids Res. *28*, 3442–3444. https://doi.org/10.1093/nar/28.18.3442.

Solowska, J.M., Garbern, J.Y., and Baas, P.W. (2010). Evaluation of loss of function as
an explanation for SPG4-based hereditary spastic paraplegia. Hum. Mol. Genet.
19, 2767–2779. https://doi.org/10.1093/hmg/ddq177.

Sowa, M.E., Bennett, E.J., Gygi, S.P., and Harper, J.W. (2009). Defining the Human
Deubiquitinating Enzyme Interaction Landscape. Cell *138*, 389–403.
https://doi.org/10.1016/j.cell.2009.04.042.

Spence, E.F., Dube, S., Uezu, A., Locke, M., Soderblom, E.J., and Soderling, S.H.
(2019). In vivo proximity proteomics of nascent synapses reveals a novel regulator
of cytoskeleton-mediated synaptic maturation. Nat. Commun. *10*, 1–16.
https://doi.org/10.1038/s41467-019-08288-w.

Stogsdill, J.A., and Eroglu, C. (2017). The interplay between neurons and glia in
synapse development and plasticity. Curr. Opin. Neurobiol. *42*, 1–8.
https://doi.org/10.1016/j.conb.2016.09.016.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set

enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proc. Natl. Acad. Sci. U. S. A. *102*, 15545–15550. https://doi.org/10.1073/pnas.0506580102.

Tang, G., Gudsnuk, K., Kuo, S.H., Cotrina, M.L., Rosoklija, G., Sosunov, A., Sonders, M.S., Kanter, E., Castagna, C., Yamamoto, A., et al. (2014). Loss of mTOR-Dependent Macroautophagy Causes Autistic-like Synaptic Pruning Deficits. Neuron *83*, 1131–1143. https://doi.org/10.1016/j.neuron.2014.07.040.

Tracy, T.E., Madero-pérez, J., Swaney, D., Chang, T.S., Konrad, C., Ward, M.E., Stevenson, E., Hüttenhain, R., Kauwe, G., Mercedes, M., et al. (2021). Tau interactome mapping reveals dynamic processes in synapes and mitochondria associated with neurodegenerative disease. BioRxiv https://doi.org/10.1101/2021.06.17.448349. .

Trost, B., Engchuan, W., Nguyen, C.M., Thiruvahindrapuram, B., Dolzhenko, E., Backstrom, I., Mirceta, M., Mojarad, B.A., Yin, Y., Dov, A., et al. (2020). Genome-wide detection of tandem DNA repeats that are expanded in autism. Nature *586*, 80–86. https://doi.org/10.1038/s41586-020-2579-z.

Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., and Cox, J. (2016). The Perseus computational platform for comprehensive analysis of (prote)omics data. Nat. Methods *13*, 731–740. https://doi.org/10.1038/nmeth.3901.

Uezu, A., Kanak, D.J., Bradshaw, T.W.A., Soderblom, E.J., Catavero, C.M., Burette, A.C., Weinberg, R.J., and Soderling, S.H. (2016). Identification of an elaborate complex mediating postsynaptic inhibition. Science (80-.). *353*, 1123–1129. https://doi.org/10.1126/science.aag0821.

Ultanir, S.K., Yadav, S., Hertz, N.T., Oses-Prieto, J.A., Claxton, S., Burlingame, A.L., Shokat, K.M., Jan, L.Y., and Jan, Y.-N. (2014). MST3 Kinase Phosphorylates TAO1/2 to Enable Myosin Va Function in Promoting Spine Synapse Development. Neuron *84*, 968–982. https://doi.org/10.1016/j.neuron.2014.10.025.

Urano, J., Sato, T., Matsuo, T., Otsubo, Y., Yamamoto, M., and Tamanoi, F. (2007). Point mutations in TOR confer Rheb-independent growth in fission yeast and nutrient-independent mammalian TOR signaling in mammalian cells. Proc. Natl. Acad. Sci. U. S. A. *104*, 3514–3519. https://doi.org/10.1073/pnas.0608510104.

Velmeshev, D., Schirmer, L., Jung, D., Haeussler, M., Perez, Y., Mayer, S., Bhaduri, A.,
Goyal, N., Rowitch, D.H., and Kriegstein, A.R. (2019). Single-cell genomics
identifies cell type–specific molecular changes in autism. Science (80-.). *364*, 685–
689. https://doi.org/10.1126/science.aav8130.

Voineagu, I., Wang, X., Johnston, P., Lowe, J.K., Tian, Y., Horvath, S., Mill, J., Cantor,
R.M., Blencowe, B.J., and Geschwind, D.H. (2011). Transcriptomic analysis of
autistic brain reveals convergent molecular pathology. Nature *474*, 380–384.
https://doi.org/10.1038/nature10110.

Wang, S., Tukachinsky, H., Romano, F.B., and Rapoport, T.A. (2016a). Cooperation of
the ER-shaping proteins atlastin, lunapark, and reticulons to generate a tubular
membrane network. Elife *5*, 1–29. https://doi.org/10.7554/eLife.18605.

Wang, Y., Picard, M., and Gu, Z. (2016b). Genetic Evidence for Elevated Pathogenicity
of Mitochondrial DNA Heteroplasmy in Autism Spectrum Disorder. PLoS Genet. *12*,
1–24. https://doi.org/10.1371/journal.pgen.1006391.

Weiner, D.J., Wigdor, E.M., Ripke, S., Walters, R.K., Kosmicki, J.A., Grove, J.,
Samocha, K.E., Goldstein, J.I., Okbay, A., Bybjerg-Grauholm, J., et al. (2017).
Polygenic transmission disequilibrium confirms that common and rare variation act
additively to create risk for autism spectrum disorders. Nat. Genet. *49*, 978–985.
https://doi.org/10.1038/ng.3863.

Wilfert, A.B., Turner, T.N., Murali, S.C., Hsieh, P.H., Sulovari, A., Wang, T., Coe, B.P.,
Guo, H., Hoekzema, K., Bakken, T.E., et al. (2021). Recent ultra-rare inherited
variants implicate new autism candidate risk genes. Nat. Genet. *53*, 1125–1134.
https://doi.org/10.1038/s41588-021-00899-8.

Willsey, A.J., Sanders, S.J., Li, M., Dong, S., Tebbenkamp, A.T., Muhle, R.A., Reilly,
S.K., Lin, L., Fertuzinhos, S., Miller, J.A., et al. (2013). Coexpression networks
implicate human midfetal deep cortical projection neurons in the pathogenesis of
autism. Cell *155*, 997. https://doi.org/10.1016/j.cell.2013.10.020.

Willsey, H.R., Exner, C.R.T., Xu, Y., Everitt, A., Sun, N., Wang, B., Dea, J., Schmunk,
G., Zaltsman, Y., Teerikorpi, N., et al. (2021). Parallel in vivo analysis of large-effect
autism genes implicates cortical neurogenesis and estrogen in risk and resilience.
Neuron *109*, 788–804. https://doi.org/10.1016/j.neuron.2021.01.002.

Wilton, D.K., Dissing-Olesen, L., and Stevens, B. (2019). Neuron-Glia Signaling in Synapse Elimination. Annu. Rev. Neurosci. *42*, 107–127. https://doi.org/10.1146/annurev-neuro-070918-050306.

Xu, Z.X., Kim, G.H., Tan, J.W., Riso, A.E., Sun, Y., Xu, E.Y., Liao, G.Y., Xu, H., Lee, S.H., Do, N.Y., et al. (2020). Elevated protein synthesis in microglia causes autism-like synaptic and behavioral aberrations. Nat. Commun. *11*. https://doi.org/10.1038/s41467-020-15530-3.

Yacoub, N. al, Romanowska, M., Haritonova, N., and Foerster, J. (2012). Optimized production and concentration of lentiviral vectors containing large inserts. J. Gene Med. *9*, 579–584. https://doi.org/10.1002/jgm.

Yadav, S., Oses-Prieto, J.A., Peters, C.J., Zhou, J., Pleasure, S.J., Burlingame, A.L., Jan, L.Y., Jan, Y.-N., Samuel, J., Burlingame, A.L., et al. (2017). TAOK2 Kinase Mediates PSD95 Stability and Dendritic Spine Maturation through Septin7 Phosphorylation. Neuron *93*, 1–15. https://doi.org/10.1016/j.neuron.2016.12.006.

Yang, W., Pang, D., Chen, M., Du, C., Jia, L., Wang, L., He, Y., Jiang, W., Luo, L., Yu, Z., et al. (2021). Rheb mediates neuronal-activity-induced mitochondrial energetics through mTORC1-independent PDH activation. Dev. Cell *56*, 811-825.e6. https://doi.org/10.1016/j.devcel.2021.02.022.

Youn, J.Y., Dunham, W.H., Hong, S.J., Knight, J.D.R., Bashkurov, M., Chen, G.I., Bagci, H., Rathod, B., MacLeod, G., Eng, S.W.M., et al. (2018). High-Density Proximity Mapping Reveals the Subcellular Organization of mRNA-Associated Granules and Bodies. Mol. Cell *69*, 517-532.e11. https://doi.org/10.1016/j.molcel.2017.12.020.

Yuen, R.K.C., Merico, D., Bookman, M., Howe, J.L., Thiruvahindrapuram, B., Patel, R. V., Whitney, J., Deflaux, N., Bingham, J., Wang, Z., et al. (2017). Whole genome sequencing resource identifies 18 new candidate genes for autism spectrum disorder. Nat. Neurosci. *20*, 13–17. https://doi.org/10.1038/nn.4524.

Zhang, Y., Pak, C.H., Han, Y., Ahlenius, H., Zhang, Z., Chanda, S., Marro, S., Patzke, C., Acuna, C., Covy, J., et al. (2013). Rapid single-step induction of functional neurons from human pluripotent stem cells. Neuron *78*, 785–798. https://doi.org/10.1016/j.neuron.2013.05.029.

Figure 1. Development of a proximity-based proteomics screen to identify neuronal PPI networks for ASD-risk genes. (A) Workflow of neuron-specific BioID2 screen for identification of ASD-risk gene PPI networks. (B) List of 41 ASD-risk gene used in the BioID2 screen

1137

**Figure 2. Neuron-specific PPI network map identifies convergent signaling
pathways between 41 ASD-risk genes.** (A) Shared PPI network map of 41 ASD-risk
bait genes. Large red nodes represent bait proteins. Color and increased thickness of
connecting lines represent number of interactions (direct or shared prey proteins)
between bait genes. (B) Correlation plot of 41 ASD-risk genes through individual PPI
networks. Genes were ordered by hierarchical clustering and clustered using kmeans (k
= 3). (C) Top 50 enriched Reactome pathways of the shared 41 ASD-risk gene PPI
network map. Individual pathways are grouped by functional similarity (g:Profiler,
Bejamini-Hochberg FDR adj. p<0.05).

1147

1148

1149

1150

1151

1152

1153

Figure 3. Regulation of cellular respiration and mitochondrial activity by ASD-risk

genes. (A) Network of TCA cycle and pyruvate metabolism proteins enriched in the

shared ASD-risk gene PPI network (g:Profiler, Benjamini-Hochberg FDR adj. p<0.05).

(B) Loss of Taok2 alters cellular respiration. WT = 118 wells and Het = 122 wells from

three separate cultures. Arrows indicate the addition of oligomycin (blue), FCCP

(green), and Antimycin A and Rotenone (red). (C) Loss of Taok2 increases maximal

respiration (*left*) and proton leakage (*right*). WT = 118 wells, Het = 122 wells from three

separate cultures. (D) Representative images of Taok2 WT, Het, and KO neurons

stained with MitoTrack (green) and TMRM (red). Arrows indicate Mitotracker-labeled

mitochondria with no TMRM staining. Scale bar is 5 µm. (E) Taok2 Het and KO neurons

have decreased active (TMRM) Mitotracker-labeled mitochondria. WT = 14, Het = 15,

KO = 15 neurons from 1-3 separate pups per genotype from one culture. (F)

Representative images of Taok2 WT, Het, and KO neurons stained with TOMM20 (red)

and Phalloidin (green). Scale bar 5 µm. (G) Taok2 Het and KO neurons have increased

TOMM20 staining. WT = 45, Het = 42, KO = 48 neurons from 1-4 separate pups per

genotype from two cultures. (H) Representative images of synaptic mitochondria

morphological categories. (I), Taok2 KO neurons have increased proportion of category

2 mitochondria with enlarged cristae. 19-25 images per animal and three animals per

genotype. (J) CRISPR/Cas9 KO of Taok2, Syngap1, Etfb, Rheb, and Spast alters

cellular respiration. Control = 45 wells, Taok2 KO = 51 wells, Syngap1 KO = 50 wells,

Etfb KO = 47 wells, Rheb KO = 48 wells, Spast KO = 45 wells from five separate

cultures. Arrows indicate the addition of oligomycin (blue), FCCP (green), and Antimycin

A and Rotenone (red). (K) Maximal respiration (*top*) and proton leakage (*bottom*) show

an increase in Syngap1 and Etfb KO neurons. Mean ± s.e.m. *$p < 0.05$, **$p < 0.01$,

***$p < 0.001$.

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

Figure 4. *De novo* missense variants alter the PPI networks of uncharacterized and established ASD-risk genes that correspond to functional deficits. (A-C), Diagram of TAOK2, GRIA1, and PPP2R5D and *de novo* missense variants. ATD: amino-terminal domain, LDB: ligand-binding domain, TM: transmembrane domain, CTD: carboxy-terminal domain, PM: plasma membrane. (D) Dot plot of top 15 cellular compartment gene sets and top 10 variant-specific gene sets in TAOK2 variants (g:Profiler, Benjamini-Hochberg FDR adj. p<0.05).. Size of dots indicate protein number and the color represents the significance. (E) Venn diagram of PPI network proteins of TAOK2 WT and A135P. (F) Representative traces of sEPSC recordings of DIV21 TAOK2 WT, KO, and A135P human iPSC-derived NGN2 neurons. (G) TAOK2 KO and A135P neurons show decreased sEPSC frequency (*left*) and amplitude (*right*). WT = 23, KO = 22, and A135P = 21 neurons from three transductions. (H) Dot plot of top 15 cellular compartment gene sets and top five variant-specific gene sets in GRIA1 variants (g:Profiler, Benjamini-Hochberg FDR adj. p<0.05). (I) Venn diagram of PPI network proteins of GRIA1 WT and variants. (J) Representative traces of sEPSC recordings of mouse neurons expressing GRIA1 or variants. (K) R208H variant shows

trend in decrease sEPSC frequency (*left*) and no change in amplitude (*right*). WT = 14,

R208H = 11, and A636T= 11 neurons from three transductions. (L) Dot plot of top 15

cellular compartment gene sets and top five variant-specific gene sets in PPP2R5D

variants (g:Profiler, Benjamini-Hochberg FDR adj. p<0.05). (M) Representative images

of PPP2R5D WT and variants show altered Golgi morphology and localization to the

dendrite (white arrow) and soma (blue arrow) (*right*) through GALNT2 staining. Inlet

shows magnified Golgi. Scale bar is 20μm. Neurons expressing the E198K variant have

decreased Golgi size, while both the E198K and E420K variants cause mislocalization

of the Golgi into the dendrite (*left*). 20 neurons from 5 separate infections per condition.

Mean ± s.e.m. *p<0.05, **p<0.01, ***p<0.001.

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

Figure 5. Shared ASD PPI network map network correlates to human brain development and disease pathology. (A) Network of ASD genes enriched in the shared ASD-risk gene PPI network. Large red nodes represent bait genes, smaller colored nodes represent sharedness of ASD-risk genes between bait genes, between 1 (yellow) and 10 (blue) shared bait genes (g:Profiler, Benjamini-Hochberg FDR adj. p<0.05). (B) Enrichment of ASD differentially expressed genes (DEGs) in cell types. (Fisher's exact test). Dashed lines represent nominal (p = 0.05, *left*) and Bonferroni corrected (p = 0.05/number of cell types, *right*) significance thresholds. EN = excitatory neurons, IN = inhibitory neurons, RGC = radial glial cells, MGE RGC = medial ganglionic eminence, IPC = intermediate progenitor cells, Astro. = astrocyte, OPC = oligodendrocyte progenitor cells, Micro. = microglia, Endo. = endothelial cells, CP = choroid plexus cells, C-F = cortico-fugal, C-C = cortico-cortico, PV = paravalbumin, SST = somatostatin, VIP = vasoactive intestinal peptide, FB = fibrous, PP = protoplasmic, Neu_NRGN = neurogranin-expressing. Light blue bars have nominal p-value significance, while dark blue bars have Bonferroni corrected significance. (C) Flow chart of clinical data extraction (D) Decrease in the average standard scores of individuals

1266   with ASD, who have rare inherited missense mutations in Group 1 genes compared to

1267   Groups 2 and 3 (Group 1 = 350, Group 2 = 113, and Group 3 = 416 probands).

1268   Individuals with splice site mutations in Cluster 1 have significantly higher adaptive

1269   behavior (E) and socialization standard (F) scores than Cluster 2 and 3. Group 1 = 32,

1270   Group 2 = 9, and Group 3 = 71 probands. Mean ± s.e.m. *p<0.05, **p<0.01.

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Dr. Karun K. Singh (karun.singh@uhnresearch.ca)

Material availability

All constructs and material are available by request from Dr. Karun K. Singh.

Data and code availability

- Mass spectrometry datasets consisting of raw files and results files with statistical analysis to identify PPI networks are available via ProteomeXchange with identifier PXD036946. DEPs are available via ProteomeXchange with identifier PXD036937. Individual PPI networks and shared ASD-risk gene PPI network map protein lists and enriched pathways can be found in Tables S1-S9. The Mouse_Human_Reactome and Mouse_GO_ALL_no_GO_iea gene sets used for overrepresentation and gene set enrichment analyses were downloaded on 13 August 2021 from http://download.baderlab.org/EM_Genesets/(Merico et al., 2010). RNA sequencing raw sequence files and results files with statistical analysis to identify significant DEGs are available through the Gene Expression Omnibus (Accession: GSE213899). ASD proband variant information and clinical scores are available through the MSSNG database (research.mss.ng)(Yuen et al., 2017) and the associated Metabase (data-explorer.mss.ng), respectively.
- This paper does not report original code.
- Any additional information required to reanalyze the data reported in this work paper is available from the Lead Contact upon request.

METHOD DETAILS

Antibodies

The following antibodies were used for immunostaining and immunoblotting experiments: rabbit anti-FLAG (IB 1:2,000, MilliporeSigma, F7425), mouse anti-FLAG (IF 1:1,000, IB: 1:2000, MilliporeSigma, F3165), rabbit anti-turboGFP (IF 1:1,000, IB 1:1,000, Fisher, PA5-22688), chicken anti-MAP2 (IF 1:1,000, Cedarlane, CLN182), rabbit anti-β-actin (IB 1:1,000, Cell Signaling, 8457S), mouse anti-β-actin (IB 1:5,000, MilliporeSigma, A5316), goat anti-TAOK2α/β (IB 1:1,000, Santa Cruz Biotechnology, sc-47447), rabbit anti-TAOK2β (IB 1:1,000, Synaptic Systems, 395 003), mouse anti-Synapsin1 (IF: 1:1000, Synaptic Systems, 106 001), mouse anti-TOMM20 (IF 1:100, US Biological, 134604), mouse anti-Shank2 (IF 1:500, Antibodies Inc), mouse anti-PSD95 (IF 1:1000, NeuroMab), rabbit anti-FMRP (IB 1:1000, Abcam), rabbit anti-STXBP1 (IB 1:1000, Proteintech), mouse anti-CS (IB 1:1000, Thermofisher), rabbit anti-CS (IP, Proteintech), rabbit anti-SYNGAP1 (IB 1:1000, Thermofisher). DAPI (IF 300mM, ThermoFisher, D21490), Hoechst (IF 1:10,000, Invitrogen, 1050083), Phalloidin-488 (IF 1:120, Cytoskeleton Inc., PHDG1), Anti-mouse-Cy3 (IF 1:500, Jackson Immunoresearch, 715-165-151), Alexa 488 anti-rabbit (IF 1:500, Jackson Immunoresearch, 711-545-152), Alexa 488 anti-chicken (IF 1:500, Jackson Immunoresearch, 703-545-155), 405 conjugated-streptavidin (IF 1:500, Jackson Immunoresearch, 016-470-084), 405 anti-chicken (IF 1:500, Jackson Immunoresearch, 703-475-155), Alexa 647 anti-mouse (IF 1:500, Jackson Immunoresearch, 715-605-150).

Generation of constructs

All cloning was accomplished using the In-Fusion HD cloning kit (Takara). To create the BioID2 fusion constructs, we obtained an expression construct containing a 198bp (13x "GGGGS" repeat) linker sequence upstream of a C-terminal 3xFLAG-tagged BioID2 sequence with BioID2 (Genscript). For lentiviral expression, 13xlinker-BioID2-3xFLAG was amplified and cloned into the lentiviral backbone pLV-hSYN-RFP (Addgene #22909)(Nathanson et al., 2009). For ease of visualization and to create a bicistronic construct, the RFP in the pLV-hSYN-RFP backbone was replaced with the TurboGFP(tGFP)-P2A from pCW57-GFP-2A-MCS (Addgene #71783)(Barger et al., 2019). NheI digest sites were added after the P2A sequence and before the 13xLinker

to allow easy insertion of ASD-risk bait genes. The final construct being pLV-hSyn-
tGFP-P2A-Bait-13xLinker-BioID2-3xFLAG (referred to as the BioID2 fusion construct).
For the control luciferase construct a second P2A was cloned in between the luciferase
ORF and the 13xLinker, creating the pLV-hSyn-tGFP-P2A-Luciferase-P2A-13xLinker-
BioID2-3xFLAG construct (referred to as the Luciferase control construct). ASD-risk
genes open reading frames (ORFs) were purchased from Addgene and Genscript or
amplified from human adult and fetal brain RNA (Takara) (see Table S10)(Alford et al.,
2012; Braganza et al., 2017; Braun et al., 2011; Butko et al., 2012; Cummins and
Vogelstein, 2004; Furlong et al., 2000; Hiday et al., 2017; Howarth et al., 2005;
Johannessen et al., 2010; Kim et al., 2016; Lu et al., 2014; Małecki et al., 2015; Ohno et
al., 2002; Seeling et al., 1999; Solowska et al., 2010; Sowa et al., 2009; Urano et al.,
2007; Wang et al., 2016a). For mouse electrophysiology experiments, the GRIA1,
GRIA1 R208H and GRIA1 A636T ORFs were inserted between the GFP-P2A and
3xFLAG. The pLV-CMV-Cas9-T2A-EGFP plasmid was made by replacing the UBC
promoter-rTetR in the FUW-M2rtTA plasmid (Addgene #20342)(Hockemeyer et al.,
2008) with CMV-Cas9-T2A-EGFP from PX458 (Addgene #48138)(Ran et al., 2013). All
generated constructs are available upon request.

1375

Experimental Models

Taok2 Het (*Taok2* +/-) and KO (*Taok2* -/-) mice were created by Kapfhamer *et*
al.(Kapfhamer et al., 2013). The E15-16 or E18 mouse embryo brains were used for
cortical neuronal cultures. P21-P23 mice were used for mass spectrometry or RNA
sequencing experiments. Animals housed at the Central Animal Facilities at McMaster
University were approved for experiments and procedures by the Animal Research
Ethics Board (AREB) at McMaster University. Animals housed at the University Medical
Center Hamburg-Eppendorf, Hamburg were approved for experiments and procedures
by local authorities of the city-state Hamburg (Behörde für Gesundheit und
Verbraucherschutz, Fachbereich Veterinärwesen) and the animal care committee of the
University Medical Center Hamburg-Eppendorf. All procedures were performed
according to the German and European Animal Welfare Act. Animals housed at the

Animal Resource Center at University Health Network were approved for experiments
and procedures by the University Health Network animal care committees.

All work with the human iPSCs was performed with the approval of the Hamilton
Integrated Research Ethics Board. iPSCs were previously generated as described in
(Deneault et al., 2018).

1393

Subject Details

1395   No subjects were directly tested in this study. Adaptive behavior and socialization
1396 standard scores of affected individuals was extracted from the MSSNG associated
1397 Metabase (data-explorer.mss.ng). These individuals are part of the MSSNG database
1398 (research.mss.ng)(Yuen et al., 2017), which has whole genome sequences of 4,258
1399 families and 5,102 ASD-affected individuals.

1400

Mouse Cortical Neuron Cultures

E15-E16 CD1 mice (Charles River) embryo cortices were harvested using a
dissecting microscope and kept in HBSS. Cortices were then digested in 300 µg/ml of
papain (Worthington) and 2 U/ml of DNase (Thermo) for 20 minutes at 37 °C. Cortices
were then washed three times with mouse plating media (Neurobasal media
supplemented with 2 mM GlutaMAX (Thermo), Pen-Strep (Thermo), and 10%
FBS(Gibco)). Digested cortices were triturated and put through 40 µm strainer. Cells
were counted, suspended in plating media, and plated at 600,000 cells per well of a 12-
well plate. Plates were coated with 100 µg/ml poly-D-lysine (mol wt > 300,000, Sigma)
and 3 µg/ml Laminin (Sigma). For immunostaining, 12 mm coverslips (Fisher) were
placed in the well prior to coating. The cells were incubated at 37 °C (with 5 % CO_2) for
one hour, after which plating media was removed and replaced with mouse culturing
media (Neurobasal media supplemented with 2 mM GlutaMAX, Pen-Strep, and B27).
Cells were grown at 37 °C (with 5 % CO_2) and half media changes were done on day 7
and every 3-4 days onwards.

1416

CRISPR/Cas9 editing of human induced pluripotent stem cells (iPSCs)

Human iPSCs were maintained on Matrigel (Corning) coated plates using

mTeSR1 media (Stem Cell Technologies) and passaged every 3-4 days using ReLeSR

(Stem Cell Technologies). Human iPSCs were edited for homozygous knockout of

TAOK2 or heterozygous knock-in of the A135P mutation as described in Deneault *et*

al.(Deneault et al., 2018). MGB probes were ordered from ThermoFisher scientific and

ssODN were designed on Benchling.com (Biology Software) and ordered from

Integrated DNA Technologies. For the A135P mutation a mutant and wildtype ssODN

containing the A135P (G to C) mutation and a PAM site mutation or just the PAM site

mutation, respectively, were used to create a heterozygous knock-in.

1427

1428   Human iPSC to neuron differentiation via NGN2 induction

Human iPSCs were cultured on Matrigel (Corning) coated plates using mTeSR1

media (Stem Cell Technologies) and passaged every 3-4 days using ReLeSR (Stem

Cell Technologies) until neural induction. A modified NGN2 induction protocol (Zhang *et*

al. 2013) was used to differentiate human iPSCs into excitatory NGN2 neurons(Zhang

et al., 2013). Human iPSCs were dual infected with pTet-O-NGN2-P2A-EGFP and

FUW-M2rtTA lentiviruses for dox-inducible expression and were titered for > 90%

infection efficiency. On Day -1 iPSCs were singularized using Accutase (Stem Cell

Technologies) and plated with mTeSR1 media (supplemented with 10 µM Y-27632) on

Matrigel at 400,000 cells per well in a 6-well plate. On Day 0, media exchanged and

supplemented with Doxycycline (1 µg/ml). On Day 1 and 2, media was replaced with

iNPC media (DMEM/F12 media (Gibco) supplemented with N2 (Gibco), MEM NEAA

(Thermo), 2mM GlutaMAX, and Pen-Strep) with Doxycycline and Puromycin (2µg/ml).

On Day 3, media was then replaced with iNi media (Neurobasal media with SM1 (Stem

Cell Technologies), 2mM GlutaMAX, Pen-Strep, 20 ng/ml BDNF, 20 ng/ml GDNF, and 1

µg/ml Laminin) with Doxycycline. On day 4, differentiated neurons were singularized

using Accutase and re-plated at 100,000 cells per well in a 24-well plate in only iNi

media. Plates were pre-coated with 20 µg/ml Laminin and 67 ug/ml Poly-ornithine

(Sigma). Mouse glial cells were plated on top of the differentiated neurons after 24

hours at a density of 50,000 cells per well. Half-media changes were carried out every

other day, and iNi media was supplemented with 2.5 % FBS on Day 9 and onwards. Neurons were grown until day 28 post NGN2-induction.

Generation of high-titer lentivirus

All viruses were made using the 2nd generation lentiviral packaging systems in Lenti-X HEK293 FTT cells (Takara). Lenti-X cells were passaged maximum 3 times before being used for virus production in HEK media (High glucose DMEM with 4 mM GlutaMAX, 1 mM Sodium Pyruvate, and 10 % FBS). Lenti-X cells were passaged once with 500µg/ml Gentamycin (Thermo) to increase T antigen expressing cells. Cells were plated into T150 flasks and each flask was transfected with the BioID2 lentiviral plasmid and the packaging plasmids, pMD2.G and pPAX2 (Addgene #12259 and #12260), using Lipofectamine 2000 in a 3:5 Opti-MEM: HEK media mix. Media was exchanged for fresh media after 5.5-6 hours. Media was harvested twice, first at 48 hours and then at 72 hours post-transection and spun at 100,000xG for 2 hours (maximum acceleration and deceleration). The virus was resuspended in PBS and kept at -80°C until they were used. Larger and unstable viruses were spun at 20,000xG for 4 hours in a table top centrifuge using a 20 % sucrose cushion(Yacoub et al., 2012). See nature exchange protocol for detailed procedure.

Infection of mouse cortical neurons for BioID2 screen

One 12-well plate of 7.2 million mouse cortical neurons was considered as one biological replicate. Each cortical neuron culture produced at least 5 plates for four separate BioID2 bait gene samples and one luciferase control sample. Three separate cultures were done in a 3 days span in one week to get 3 biological replicates per protein-of-interest (POI). On days *in vitro* (DIV) 14, the conditioned media from the mouse neuron cultures were removed, leaving only 0.5 ml of media per well. Extra wells with and without coverslips were infected at the same MOI for flow cytometry measurements of GFP positive neurons and immunostaining, respectively. On DIV14, lentivirus with BioID2 fusion constructs were added to each well with an average transduction efficiency of 70%, and on DIV17 each well was supplemented with 50 µM of Biotin. After 18-20 hours, cells for mass spectrometry were lysed with RIPA buffer (1

% NP40, 50 mM Tris-HCl, 150 mM NaCl, 0.1 % SDS, 0.5 % deoxycholic acid, and
protease inhibitor cocktail (PIC)) and flash frozen in liquid nitrogen. Cells for flow
cytometry were dissociated with 0.25 % Trypsin-EDTA (Fisher) and resuspended in
PEF media (PBS with 2 mM EDTA and 5 % FBS) (See flow cytometry section). Cells for
immunostaining were fixed with 4 % PFA for 20 minutes, washed with PBS, and kept at
4 °C for staining.

1485

Transfection of HEK293 FT cells for BioID2 screen

10 million HEK293 FT cells were plated in a 10 cm culture dish and transfected
24 hours later with the BioID2 fusion construct plasmids using Lipofectamine 2000.
Media was changed 6 hours after transfection and 50µM biotin was added 48 hours
post-transfection. Cells were lysed 72 hours post-transfection in RIPA buffer and flash
frozen in liquid nitrogen. Each individual plate was considered as biological replicate
and three plates were used for each gene and the luciferase control. An extra plate was
used for flow cytometry measurements of GFP positive cells.

1494

Processing of mouse cortical neuron and HEK293 FT cell BioID2 samples

Lysed cells were thawed and DNA was digested using benzonase (Sigma).
Lysates were than sonicated at high speed for 5 seconds and centrifuged at 20,000xG
for 30 minutes. The lysate supernatants were incubated with streptavidin Sepharose
beads (GE Healthcare) at 4 °C for 3 hours. Following the incubation, the supernatant
was spun down at 100xG for 2 minutes and the supernatant was removed. The beads
were than washed once with RIPA buffer, and then six times with 100 mM
triethylammonium bicarbonate (TEAB) with centrifugation between each wash. After the
final wash, the beads are then resuspended in 100 mM TEAB and sequencing-grade
trypsin (Promega) was added to digest the biotinylated proteins on the beads into
peptides. The beads were incubated at 37 °C for 16 hours while rotating, and additional
trypsin was added and incubated for a further 2 hours. The beads were then pelleted
and the supernatant was transferred to a new tube. The beads were washed twice with

100 mM TEAB and each wash was added to the supernatant. The supernatant was
then transferred to a 1.5 mL screw cap tub and speed vacuum dried. The dried peptides
were stored at 4 °C for TMT-labeling.

1511

Multiplex TMT-labeling of BioID2 samples

TMT10plex isobaric-labeling was used to combine at least 3 biological replicates
per gene. One additional technical TMT-labeling replicate of luciferase control sample
chosen at random was used to account for differences in labeling. Dried peptides were
resuspended in 100 mM TEAB. Each sample was TMT-labeled using the TMT 10plex
Isobaric Mass Tagging Kit (Thermo). The four genes (proteins-of-interest, POI) were
divided into two separate batches and the luciferase control samples were divided
between the batches. Each batch had three biological replicates of the two genes and
the luciferase control. One luciferase sample chosen at random was divided and labeled
with two different labels to determine variance due to labeling efficiencies. In brief, TMT-
label resuspended in acetonitrile was added to each sample and incubated at room
temperature for one hour. To stop the reaction, 5 % hydroxylamine was then added to
the samples and incubated for 15 minutes at room temperature. All ten samples were
combined into one tube and divided into two samples. Both samples were then speed
vacuum dried. One sample was kept at -80 °C for storage and the second sample was
kept at 4 °C to be run in the mass spectrometer.

1528

Identification of biotinylated proteins from BioID2 screen samples using LC-MS/MS

Peptide samples were resuspended in 0.1% Trifluoroacetic acid (TFA) and
loaded for liquid chromatography, which was conducted using a home-made trap-
column (5 cm x 200 µm inner diameter; POROS 10 µm 10R2 C18 resin) and a home-
made analytical column (50 cm x 50 µm inner diameter; Monitor 5 µm 100A C18
resin), running a 120min (label free) or 180min (TMT) reversed-phase gradient at
70nl/min on a Thermo Fisher Ultimate 3000 RSLCNano UPLC system coupled to a
Thermo QExactive HF quadrupole-Orbitrap mass spectrometer. A parent ion scan was
performed using a resolving power of 120,000 and then up to the 20 most intense peaks

were selected for MS/MS (minimum ion count of 1000 for activation), using higher energy collision induced dissociation (HCD) fragmentation. Dynamic exclusion was activated such that MS/MS of the same m/z (within a range of 10 ppm; exclusion list size = 500) detected twice within 5 seconds were excluded from analysis for 30 seconds. Data were analyzed using Proteome Discoverer 2.2 (Thermo). For protein identification, search was against the Swiss-Prot mouse proteome database (55,366 protein isoform entries)(Bateman et al., 2021), while the search parameters specified a parent ion mass tolerance of 10 ppm, and an MS/MS fragment ion tolerance of 0.02 Da, with up to two missed cleavages allowed for trypsin. Dynamic modification of +16@M was allowed.

Analysis for the identification of ASD-risk and cellular compartment protein PPI networks

Only proteins identified with two unique peptides were used for analysis. Two statistical cut-offs were used to identify positive hits for the PPI networks of each POI: Biotinylated proteins in the POI sample with 1) a significant increase in Log2 abundance compared to the luciferase control (Student's t-test, $p<0.05$)(Uezu et al., 2016) and 2) that were significant outliers when accounting for the overall protein abundance compared to the protein abundance ratio between the POI and control samples (SigB $p<0.05$)(Cox and Mann, 2008). Protein abundances were normalized between biological replicates based on the sample with the highest total protein abundance. To reduce variability between each viral transduction, flow cytometry was used to determine the total abundance of GFP in the infected neurons between samples. The protein abundance levels of samples that had less total GFP (area under the curve in GFP intensity histogram) than the luciferase control were normalized by the factor required to equalize the total GFP of the POI sample to the luciferase control sample. To account for false positive hits due to variability in TMT-labeling between samples, the ratio of protein abundances between the luciferase control technical TMT-labeling replicates were used as the minimal required ratio between POI and control sample abundances.

1569 Proteins that did not have abundance ratios (POI/Luciferase control) higher than this
1570 minimal ratio were considered false positives and removed from further analysis.

1571

Pathway enrichment analyses

All pathway enrichment analysis was done using the g:Profiler GOst functional
profiling tool (https://biit.cs.ut.ee/gprofiler/gost)(Raudvere et al., 2019). We used internal
sources without electronic GO annotations for GO biological processes and GO cellular
component (compartment), and curated Reactome pathway gene sets from the Bader
lab (http://download.baderlab.org/EM_Genesets/)(Merico et al., 2010). All three sources
were used for the shared ASD-risk gene network proteins. Only GO cellular component
enrichment was used for the HEK293 FT cell BioID2 PPI networks, neuron cellular
compartment BioID2 PPI networks, and *de novo* missense mutation network BioID2 PPI
network comparisons. We compared the protein lists against a custom statistical
domain of proteins identified through fractionated mass spectrometry of the mouse
brain(Sharma et al., 2015) and combined with any additional proteins identified in the
BioID2 screen. The final mouse brain proteome background had a total of 11992
proteins after removing multiple isoforms of the same protein. HEK293 BioID2 PPI
networks were compared to all annotated gene lists. The g:Profiler Benjamini-Hochberg
FDR multiple correction was used and only pathways with an adj. p-value < 0.05 were
considered significantly enriched. For cellular component enrichment for *de novo*
missense variant BioID2 PPI networks, the ggplot package in R was used to create the
dot plots. For *de novo* missense variant BioID2 bait genes, all proteins identified in the
wildtype samples were used for analysis, while for the shared PPI network map only
proteins found in all wildtype samples were used for pathway enrichment analysis.

1593

Virus titering and GFP normalization for BioID2 screen

Mouse cortical neurons were cultured as described above and infected on DIV3
at three dilutions of virus (1:100, 1:333, 1:1000). On DIV 5, infected mouse cortical
neurons were singularized using 0.25 % Trypsin-EDTA (Fisher) and resuspended in
PEF media (PBS with 2 mM EDTA and 5 % FBS). For GFP normalization, DIV18
mouse neurons infected with the BioID2 lentiviruses were dissociated with Trypsin and

resuspended in PEF media. CytoFLEX-LX or CANTO II flow cytometers (UHN
Research Flow Cytometry Facility – KDT Site) were used to measure the percentage of
GFP-positive cells with the 488 laser and 525/40 or 525/50 filters, respectively, using
CytExpert software (Beckman Coulter). Functional titers were calculated based on the
linear relationship between virus amount and percent of GFP positive cells. Mouse
cortical neurons were infected at an MOI of 0.7, where 70 percent of cells were
expected to be infected with the BioID2 lentiviral constructs. For normalization, the total
GFP per 20,000 GFP-positive cells were quantified by taking the area under the GFP
intensity histogram. GFP percentage and total amount was calculated using FlowJo
software.

1610

Western blots

HEK293 FT cells were transfected with the BioID2 constructs using
Lipofectamine 2000 (Invitrogen) in Opti-MEM: HEK media. Cells were harvested 48
hours post-transfection and lysed with RIPA buffer (with fresh PIC). Lysates were either
snap-frozen in liquid nitrogen or taken directly for western blot sample preparation.
Thawed or fresh lysates were sonicated at high frequency for 5 seconds and
centrifuged at 20,000xG for 5 minutes at 4 °C. Lysates were than quantified using the
Bio-Rad Bradford protein assay (Bio-Rad) by measuring absorbance with the
SPECTROstar Nano machine and MARS Data analysis software (BMG LABTECH) and
diluted to equal concentrations with RIPA buffer. 30-50 µg of protein were run on 8 % or
10 % SDS-PAGE Tris-Glycine gels (depending on the size of the proteins) at 100V for
initial stacking and then 140V for 1-1.25 hours in a Tris-Glycine running buffer. Proteins
were then transferred onto PVDF membrane using a Tris-Glycine buffered wet transfer
system for 2 hours at constant 200 mA. Blots were then blocked with 5% milk in TBS-T
(Tris buffered saline pH 7.4 with 0.1 % Tween). Blots were incubated with primary
antibodies overnight in 5 % milk/TBS-T. The next day, membranes were washed three
times with TBS-T for 5 minutes each and then incubated with secondary antibodies in 5
% milk/TBS-T for 1 hour. Blots were imaged by incubating them with the Amersham
ECL western blotting detection reagent (VWR) for 1 minute and then imaging every 10

seconds for 5 minutes on the ChemiDoc XRS+ machine (Bio-Rad). ImageLab (Bio-Rad)

was used for band intensity quantification.

1632

1633 Co-immunoprecipitation Western Blots

Cortices were harvested from 3-4 week old CD1 mice and homogenized in NP40

lysis buffer (150 mM NaCl, 1% NP40, 50 mM Tris-Cl pH 8.0). Magentic dynabeads

Protein G or Protein A (Invitrogen) were washed with 0.05% Triton X-100 in PBS, and

then incubated with 4 µg of the antibody, or IgG for the control (Santa Cruz). Antibody-

bead complexes were then washed 4 times with NT2 buffer (50 mM Tris-HCl, 150 mM

NaCl, 1 mM $MgCl_2$, 0.05% NP40). Lysates were quantified using the Bradford assay

and 1 mg of protein was incubated with the antibody-bead complexes for 16hrs at 4

degrees Celsius. The beads were then washed 5 times with NT2 buffer, resuspended in

2x Laemmli buffer, and boiled at 95 degrees Celsius for 10 minutes. The supernatant

was then stored at -80 degrees until they were used for western blots.

1644

1645 *In vitro* whole-cell patch clamp recordings of human iPSC-derived neurons and
1646 mouse cortical neurons

Human iPSC-derived neurons were used for electrophysiology experiments

between days 21-24 of the neural differentiation protocol. Whole-cell patch-clamp

recordings were performed at room temperature using Multiclamp 700B amplifier

(Molecular Devices) from borosilicate patch electrodes (P-97 puller and P-1000 puller;

Sutter Instruments) containing a potassium-based intracellular solution (123 mM K-

gluconate, 10 mM KCl, 10 mM HEPES; 1 mM EGTA, 1 mM MgCl2, 0.1 mM CaCl2, 1

mM Mg-ATP, and 0.2mM Na4GTP, pH 7.2). 0.06% sulpharhodamine dye was added to

the intracellular solution to confirm the selection of multipolar neurons. The extracellular

solution consisted of 140 mM NaCl, 5 mM KCl, 1.25 mM NaH_2PO_4, 1 mM $MgCl_2$, 2 mM

$CaCl_2$, 10 mM glucose, and 10 mM HEPES (pH 7.4). Data was digitized at 10 – 20 kHz

and low-pass filtered at 1 - 2 kHz. Recordings were omitted if access resistance was

>30 MΩ. Whole-cell recordings were clamped at -70 mV and corrected for a calculated -

10mV junction potential. Rheobase was determined by a step protocol with 5 pA

increments, where the injected current had 25 ms duration. Action potential waveform

parameters were all analyzed in reference to the threshold. Repetitive firing step protocols ranged from -20 pA to +50 pA with 5 pA increments. No more than two neurons per coverslip were used to reduce the variability. Data were analyzed using the Clampfit software (Molecular Devices), while phase-plane plots were generated in the OriginPro software (Origin Lab). For GRIA1 overexpression experiments, mouse neurons were infected with GRIA1, GRIA1 R208H, and GRIA1 A636T lentiviral constructs at DIV11 and recorded on DIV14-15. Mouse neurons for electrophysiology experiments were cultured in Neurobasal media (supplemented with an additional 0.3 % (w/v) glucose and 0.22 % (w/v) NaCl). The same intracellular solution was used as the human neuron recordings, with a mouse artificial cerebrospinal fluid extracellular solution (125 mM NaCl, 2.5 mM KCl, 2 mM CaCl2, 1 mM $MgCl_2$, 5 mM HEPES, 33 mM Glucose, pH 7.2). Whole-cell recordings of mouse neurons were clamped at -80 mV and corrected for a calculated -10mV junction potential.

Staining and imaging of mouse cortical neurons and human iPSC-derived neurons

Mouse cortical neurons and human iPSC-derived neurons on coverslips were fixed with 4% paraformaldehyde (PFA) for 10 minutes at room temperature, washed once with PBS, and stored in PBS at 4 °C protected from light. Fixed coverslips were then blocked and permeabilized in BP solution (PBS with 10% donkey serum and 0.3% Triton-X) for 45 minutes at room temperature. Coverslips were then incubated with primary antibodies at 4 °C overnight. The following day, coverslips were washed three times with PBS for eight minutes each. Coverslips were then incubated with secondary antibodies for one hour at room temperature, followed by three washes with PBS. For human iPSC-derived neuron Synapsin1 staining, coverslips were incubated with 300 mM of DAPI for 15 minutes, before the third wash with PBS. Excess liquid was then removed from the coverslips and they were mounted onto VistaVision glass microscope slides (VWR) with 10 µL of Prolong Gold Anti-Fade mounting medium (Life Technologies).

For TOMM20 staining, mouse neurons were fixed with 4% PFA at 37°C for 10 min and then permeabilized with 0.5 % Triton X-100 for 10 minutes. Non-specific

binding was blocked by incubation with 5 % donkey serum in PBS for 50 minutes at room temperature, followed by primary antibody incubation. The secondary antibody was added for 50 minutes at room temperature. Primary and secondary antibodies were diluted in PBS with 0.5 % BSA, 2.5 % Donkey-serum, and 0.15 % Triton X-100. After primary and secondary antibody incubation, three washing steps with PBS were performed. Then, coverslips were incubated with Phalloidin-488, for F-actin labeling, and Hoechst dye for 45 minutes at room temperature followed by three PBS washes. Coverslips were mounted onto slides using Fluoromount-G® (Southern Biotech) and were stored protected from light. Synapsin1 and BioID2 stained images were taken on the Zeiss LSM 700 confocal microscope with63x or 40x oil objective, respectively. Mito-dsRed images were taken on the Echo Revolve microscope with a 20x objective.

For Duolink Proximity Labeling assays, the manufacturers protocol was followed in detail (Sigma). DIV17 mouse neuron cultures were fixed in 4% PFA for 20 minutes, permeabilized and blocked as described above, and incubated with the primary antibody overnight at 4 degrees Celsius. Anti-Mouse MINUS and Anti-Rabbit PLUS probes were used with the FarRed In Situ detection reagents to identify co-localized proteins. Anti-MAP2 was co-stained on coverslips during the primary and secondary incubations. Coverslips were mounted with the DuoLink In Situe Mounting Medium with DAPI.

Synapsin1 puncta analysis in human iPSC-derived neurons

Synapsin1 stained images were processed and analyzed with ImageJ software. The Synapsin1 antibody was co-immunostained with MAP2 to determine dendrites with presynaptic puncta. Five biological replicates, which represent five separate neural inductions, were used for synaptic analysis. 5-10 neurons per genotype per replicate were used. Imaging settings were kept the same between images and synapsin1 images were analyzed at the same threshold. Dendrites were traced using ImageJ and the measure tool was used to quantify the number and size of the puncta within the traced region.

Proteomic profiling of *Taok2* KO mice cortical post-synaptic density fraction through LC-MS/MS

The right cortical lobes of three P21-23 *Taok2* KO mice and five P21-23 wildtype littermates were harvested and differential centrifugation was used to obtain the crude post-synaptic density fraction(Kwan et al., 2016). PSD fractionations were validated by western blot for PSD-95 and synaptophysin (data not shown). Final post-synaptic density pellets were resuspended using 8 M urea and 100 mM ammonium bicarbonate. Protein samples were then reduced with 10 mM Tris(2-carboxyethyl)phosphine for 45 min at 37 °C, alkylated with 20 mM iodoacetamide for 45 min at room temperature, and digested by trypsin (Promega) (1:50 enzyme-to-protein ratio) overnight at 37 °C. The peptides were desalted with the 10 mg SOLA C18 Plates (Thermo Scientific), dried, and labeled with Multiplex 10-plex TMT labels (Thermo) in 100 mM triethylammonium bicarbonate, and quenched with 5% hydroxylamine before combined. 40 µg of the pooled sample was separated into 60 fractions by high-pH reverse-phase liquid chromatography (RPLC) using a homemade C18 column (200 µm × 30 cm bed volume, Waters BEH 130 5 µm resin) running a 70 min gradient from 11 to 32% acetonitrile− 20 mM ammonium formate (pH 10) at a flow rate of 5 µL/min. Each fraction was then loaded onto a homemade trap column (200 µm × 5 cm bed volume) packed with POROS 10R2 10 µm resin (Applied Biosystems), followed by a homemade analytical column (50 µm × 50 cm bed volume) packed with Reprosil-Pur 120 C18-AQ 5 µm particles (Dr. Maisch) with an integrated Picofrit nanospray emitter (New Objective). LC-MS experiments were performed on a Thermo Fisher Ultimate 3000 RSLCNano UPLC system that ran a 3 h gradient (11− 38% acetonitrile−0.1% formic acid) at 70 nL/min coupled to a Thermo QExactive HF quadrupole-Orbitrap mass spectrometer. A parent ion scan was performed using a resolving power of 120 000; then, up to 30 of the most intense peaks were selected for MS/MS (minimum ion counts of 1000 for activation) using higher energy collision-induced dissociation (HCD) fragmentation. Dynamic exclusion was activated such that MS/MS of the same m/z (within a range of 10 ppm; exclusion list size = 500) detected twice within 5 seconds was excluded from the analysis for 30 seconds. Data were analyzed using Proteome Discoverer 2.2 (Thermo). For protein identification, search was against the SwissProt mouse proteome database

(55,366 protein isoform entries), while the search parameters specified a parent ion mass tolerance of 10ppm, and an MS/MS fragment ion tolerance of 0.02Da, with up to two missed cleavages allowed for trypsin. Dynamic modification of +16@M was allowed. Only proteins with two unique peptides were used for further analysis. Differentially expressed proteins (DEPs) were calculated through Significance B outlier test using the Perseus software(Tyanova et al., 2016), and only proteins that had adj. p-value < 0.05 were considered as DEPs.

Transcriptome profiling of *Taok2* KO mice cortices through RNA sequencing

Cortices from *Taok2* KO and wildtype littermates were also harvested for RNA at post-natal day 21-23, with 3 males and 3 females from each genotype. The RNA was extracted using Trizol and was sent for total RNA sequencing at the Center for Applied Genomics (TCAG). mRNA was purified using poly(A) selection to avoid contamination of ribosomal RNAs and miRNAs. All samples were run on one lane resulting in ~31-34 million of read pairs per sample. All analysis was carried out using the open-source platform Galaxy (usegalaxy.org)(Afgan et al., 2018). RNA reads were checked for good quality using the FastQC tool. The trimmomatic tool was used to identify and trim off known adaptors and remove any bases that have a Phred score of less than 20. FastQC was used again to ensure that adaptor sequences were removed and that the quality of the reads was not affected. We next used the HISAT2 alignment program for alignment of the RNA sequences to the mouse genome GRCm38 (NCBI). On average 85% of reads from mouse samples were aligned once and 5% were aligned more than once to distinct genome locations. Moving on, the featureCounts tool was used to count the number of reads per gene using the same reference genome as the HISAT2 tool. The DESeq2 tool was used to determine the significant differentially expressed genes (DEGs) between *Taok2* WT and KO mouse cortices. Genes were considered as DEGs if they had an adjusted p-value lower than 0.05.

Gene set enrichment analysis (GSEA) of *Taok2* KO mouse proteome and transcriptome

DEGs and DEPs were ranked based on the equation $-\log_{10}$(adj. p-value)*Ln(fold

change). GSEA 4.1.0 (Broad Institute)(Daly et al., 2003; Subramanian et al., 2005) was

used to run the GSEA preranked test. Tests were run with 1000 permutations, weighted

enrichment statistics, and excluding gene sets smaller than 15 and larger than 500

genes. All other settings were kept as default. All mouse GO term gene sets without

electronic GO annotations (http://download.baderlab.org/EM_Genesets/) were used for

the analysis(Merico et al., 2010). Visualization of the enriched gene sets was done on

Cytoscape 3.8.2 using the EnrichmentMap app and the AutoAnnotate app was used for

clustering similar gene sets(Bader et al., 2014; Kucera et al., 2016; Shannon et al.,

2003). All visualized gene sets had an FDR < 0.1.

1793

Seahorse assay of *in vitro* mouse and human iPSC-derived neurons

Mouse cortical neurons were cultured as described above at a density of 30,000

cells/well in the Seahorse XF96 cell culture microplate. CRISPR/Cas9 KO mouse

neurons were infected at DIV 7 and assayed at DIV 14. Human iPSC-derived NGN2

neurons were plated on day 4 of dox induction at a density of 50,000 cells per well in

the Seahorse XF96 cell culture microplates (Agilent), pre-coated with 20 µg/ml Laminin

and 67 µg/ml Poly-ornithine (Sigma). Mouse glia was plated on top of the neurons at a

density of 25,000 cells per well, 24 hours later. Cells were used for the Seahorse assay

on day 7. The day prior to the seahorse assay the Seahorse XFe96 sensor cartridge

was filled with Calibrant XF solution and incubated at 37 °C (without CO_2) overnight. On

the day of the assay the Seahorse XF96 microplates were washed twice with 200 µl per

well of pre-warmed MST media (Seahorse XF DMEM pH 7.4 media supplemented with

1 mM sodium pyruvate, 2.5 mM GlutaMAX, and 17.5 mM Glucose). The plate was then

filled with 180 µl per well MST media and incubated at 37 °C (without CO2) for 1 hour.

During the incubation, the mitochondrial stress test drugs were added to the XFe96

sensor cartridge (1 µM Oligomycin for mouse neurons and 3 µM Oligomycin for human

neurons, 1 µM FCCP, and 1 µM Rotenone/Antimycin A resuspended in MST media).

The cartridge plate with the drug compounds were then put in the Seahorse XFe96

analyzer for calibration. After calibration, the microplate was placed into the Seahorse

XFe96 analyzer for the pre-set mitochondrial stress test protocol. Oxygen consumption

rates (OCR) were recorded every seven minutes and the drug compounds were added
in 21-minute intervals. Oligomycin was used to inhibit ATP-synthase to measure ATP-
synthase dependant respiration, FCCP was added to decouple the inner membrane to
measure maximal respiration, and Rotenone and Antimycin A were added together to
measure non-mitochondrial respiration. After the assay, microplates were frozen at -
80°C overnight and cell content was measured using the Cyquant cell proliferation
assay (Thermo) by measuring fluorescence with the CLARIOStar machine and MARS
data analysis software (BMG LABTECH). Cellular respiration analysis was performed
using the Wave software (Agilent) and OCR values were normalized to the number of
cells per well.

1824

CRISPR/Cas9 knockout in mouse cortical neurons

Mouse cortical neurons were infected at DIV7 with the pLV CMV-Cas9-T2A-
EGFP (MOI 1) and pLV U6-sgRNA/EF1a-mCherry (MOI 3) lentiviruses. Cultures were
allowed to recover until DIV14 and were then taken for the seahorse assay. The
GeneArt genomic cleavage detection kit (Thermo) was used to detect insertions or
deletions in the targeted sites.

1831

Measuring mitochondrial activity in shRNA knockdown mouse neurons

Embryonic age E15 C57BL6/J mouse pups were *in utero* electroporated with
Taok2 shRNA and control shRNA. Electroporated mouse embryo cortices were than
harvested and cultured at E18. Mouse neuron cultures were imaged at DIV5 after
incubation with 2nM TMRM (Thermo). Images were analyzed on ImageJ. Soma regions
were delineated and integrated density in the soma (soma area x mean intensity) was
measured. For background correction, mean background intensity was obtained from
the neighbouring region.

1840

Measuring mitochondrial activity and content in mouse cortical neurons

DIV 6 mouse cortical neurons cultured from *Taok2* WT, Het, and KO mouse
embryos were incubated with 2 nM TMRM (Invitrogen, #T668) and/or 100 nM
MitoTracker Green (Cell Signaling Technology, #9074P) were directly added to the

conditioned medium, and incubated for 15 minutes. Cells were then imaged within 30

minutes after the incubation time. Images were loaded onto ImageJ, background mean

intensity was measured from the region without TMRM and MitoTracker signals inside

the cell, then the cell was delineated and the background was removed. After

background correction, using the JACoP plugin the TMRM-MitoTracker signal

colocalization was analyzed using Manders' correlation coefficients. For Manders'

correlation coefficients, threshold values for TMRM (red channel) and MitoTracker

(green channel) were set to 335±55 and 640±50 respectively. 16-bit wide field images

were taken on a Nikon EclipseTi2 inverted spinning disk microscope equipped with 60X

oil (NA 1.4) objective, an LED light source (Lumencor® from AHF analysentechnik AG,

Germany), and a digital CMOS camera (ORCA-Flash4.0 V3 C13440-20CU from

Hamamatsu) controlled with NIS-Elements software. The microscope imaging chamber

is equipped to maintain 37 °C temperature and 5 % CO_2. Illumination, exposure and

gain settings were kept the same across different conditions for imaging TMRM and

MitoTracker signals.

1860

TOMM20 staining analysis for mitochondria content in mouse cortical neurons

DIV 7-8 mouse cortical neurons cultured from *Taok2* WT, Het, and KO mouse

embryos were fixed and stained for TOMM20. 16-bit Z-series images with a step size of

300 nm Images were acquired on confocal spinning disk microscope using a 60X oil

(NA 1.4) objectives. Illumination, exposure and gain settings were kept the same across

the conditions. The images were loaded onto ImageJ and z-projection (sum slices) for

the entire cell in z-axis was performed on the confocal images. Using ImageJ, soma

region was carefully delineated and total intensity, also known as integrated density, in

the soma (soma area * mean intensity) was measured. For background correction,

mean intensity (background mean intensity) was obtained from the neighbouring region

(out of the cell). Using the following equation, we obtained the corrected values.

Corrected value = total intensity in the soma – (background mean intensity * soma

area).

1874

1875 **Electron microscopy of synaptic mitochondria from mouse brain cortices**

Coronal vibratome sections of the cingulate cortex (cg1 and cg2) and the prelimbic cortex (PL) of the PFC, the primary somatosensory regions S1HL, S1Fl, S1BF, and the intermediate HC were collected and prepared for electron microscopy as described in Richter *et al.*(Richter et al., 2019). Semithin sections (0.5 µm) were prepared for light microscopy mounted on glass slides and stained for 1 min with 1% Toluidine blue. Ultrathin sections (60 nm) were examined in an EM902 (Zeiss, Munich, Germany). Pictures were taken with a MegaViewIII digital camera (A. Tröndle, Moorenweis, Germany). EM images that were collected and analyzed for synapse formation on the dendritic spines or shafts from Richter *et al.* were reanalyzed for mitochondrial morphology. Mitochondria morphology from the EM images obtained from Taok2 Wt and Taok2 KO genotypes were analyzed manually using ImageJ. based on their morphology the mitochondria are and categorized to Category 1 - Normal mitochondria with well stacked Cristae, Category 2 - mitochondria with enlarged Cristae, Category 3 - mitochondria with condensed Cristae.

Mito7-dsRed puncta analysis in human iPSC-derived neurons

TAOK2 KO, A135P and wildtype human iPSC-derived NGN2 neurons were transfected with 0.8 µg of Mito7-dsRed (Addgene #55838) and 0.2 µg of pCAG-Venus at day 5, with 2 µl of Lipofectamine 2000 (Thermo). Venus was used to trace neuron projections.10 neurons per genotype from two separate neural inductions were used for analysis. Imaging settings were kept the same between images and Mito7-dsRed images were analyzed at the same threshold. Dendrites were traced using ImageJ and the measure tool was used to quantify the size of the puncta within the traced region

Correlation of 41 ASD-risk gene PPI networks

Corrplot (R package) was used to create the correlation plot. The normalized biotinylation score to the bait protein was used to calculate the correlation between ASD-risk gene PPI networks. The Silhouette and Within cluster sum of squares methods were used to calculate the optimal kmeans number for clustering. Genes were ordered by hierarchal clustering.

Cell type/DEG/ASD gene list enrichment analysis

Human cell type gene expression and ASD DEGs and ASD gene lists were obtained from their respective publications(Feliciano et al., 2019; Nowakowski et al., 2017; Ruzzo et al., 2019; Sanders et al., 2015; Satterstrom et al., 2020; Velmeshev et al., 2019; Wilfert et al., 2021; Yuen et al., 2017). For the enrichment analysis we used the Fisher exact test comparing each gene list with the shared ASD-risk gene PPI network in the mouse brain background protein list, which was used for pathway enrichment analysis. P-values and ODDs ratios were calculated for each comparison. To account for multiple comparisons, Bonferroni correction thresholds were calculated as p = 0.05 divided by the number of comparisons.

Clinical score analysis

Rare variants of individuals diagnosed with ASD were extracted from the MSSNG database (research.mss.ng)(Yuen et al., 2017). Only variants with estimated high or medium impact strengths were used for analysis, and variants were categorized into three groups (missense variants, splicing variants, and frame shift/premature stop codon variants). Adaptive behavior and socialization standard scores of affected individuals were extracted from the MSSNG associated Metabase (data-explorer.mss.ng). Individuals were grouped based on the presence of mutations in the 41 ASD-risk genes that were clustered into three groups. Individuals that had variants in genes between multiple groups were not included in the analysis. Separate analyses were carried out between individuals grouped by missense, splicing or frame shift/premature stop codon variants. Clinical data was considered as non-parametric and the Kruskal-Wallis ranked test with post hoc Dunn's test was used for comparison between the adaptive behavior and socialization standard scores of each group.

Data representation and figure generation

Networks and gene set enrichment maps were created on Cytoscape v3.8.2. Graphs were created on GraphPad Prism 7. Representative electrophysiology traces were extracted onto CorelDRAW. Microscopy images were prepared using ImageJ. Dot plots, correlation plots, and heat maps were created on R Studio. Graphical abstract and

1938 flowcharts were created on and exported from BioRender.com (BU24JATEEX,
1939 SD235B8ORF, KW235KT7TM, RZ235KTA0S). Final figures were organized and
1940 created using Adobe Illustrator CC.

1941

QUANTIFICATION AND STATISTICAL ANALYSIS

Data are expressed as mean ± s.e.m, except the clinical analysis which is shown
as a box and whisker plot showing the minimum, median, and maximum scores. A
minimum of three biological replicates were used for all experiments, where separate
HEK cell transfections, iPSC dox-inductions, mouse neuron cultures, or littermates are
considered as individual biological replicates. All statistical analysis was done on
GraphPad Prism 7. All comparisons were assumed to be parametric, except for the
clinical score analyses. ROUT's outlier test was used to identify possible outliers, with a
Q value of 0.1 %. For statistical analysis unpaired t-test, or One-Way ANOVA and Two-
Way ANOVA with *post hoc* Holm-Sidak tests were used to compare all experimental
conditions to the control condition. All unpaired t-tests were two-sided, except for the
one-sided t-test used for identification of BioID2 prey proteins. Clinical scores were
assumed to be non-parametric and the Kruskal-Wallis H test with post hoc Dunn's test
was used to compare all groups to each other. Any variation from the described
statistical analyses is described and explained in the figure legends. The p-values are
defined in the figure legends and $p < 0.05$ are considered statistically significant.

Supplementary Tables

Table S1. BioID2 PPI networks of 41 ASD-risk genes and cellular compartment genes. Related to Figure 2.

Table S2. Comparison of BioID2 PPI networks identified in HEK293 cells and mouse cortical neurons. Related to Figures 1, S2 and S4.

Table S3. Comparison of BioID2 PPI network enriched cellular components identified in HEK293 cells and mouse cortical neurons. Related to Figures 1, S3 and S4**.**

Table S4. BioID2 PPI network enriched cellular components of compartment specific genes. Related to Figure 2.

Table S5. BioID2 PPI network enriched pathways of 41 ASD-risk genes. Related to Figure 2.

Table S6. 41 ASD-risk gene PPI network map enriched pathways. Related to Figure 2.

Table S7. Differentially expressed genes and proteins and dysregulated pathways in *Taok2* KO mouse cortices. Related to Figures 3 and S7.

Table S8. Comparison of BioID2 PPI networks between ASD-risk genes and their variants. Related to Figure 4.

Table S9. BioID2 PPI network enriched pathways of ASD-risk genes and their variants. Related to Figure 4.

Table S10. List of sources for 41 ASD-risk genes and cellular compartment genes. Related to Figure 1.