1 **Title**

2 # Catch bond models may explain how force amplifies TCR signaling and
3 # antigen discrimination
4

5 **Authors**
6 Hyun-Kyu Choi[1,2], Kaitao Li[1,2], Peiwen Cong[1,2], Muaz Nik Rushdi[1,2,†], Chenghao Ge [1,2,‡], Wei
7 Chen[3], Jizhong Lou[4-6], and Cheng Zhu[1,2,7,*]
8
Affiliations

[1]Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of
Technology and Emory University, Atlanta, Georgia 30332, USA

[2]Parker H. Petit Institute for Bioengineering and Biosciences, Georgia Institute of
Technology, Atlanta, Georgia 30332, USA

[3]Department of Cell Biology and Department of Cardiology of the Second Affiliated
Hospital, Zhejiang University School of Medicine, Hangzhou 310058, China

[4]National Laboratory of Biomacromolecules, Institute of Biophysics, Chinese Academy of
Sciences, Beijing 100101, China

[5]Key Laboratory of RNA Biology, CAS Center for Excellence in Biomacromolecules,
Institute of Biophysics, Chinese Academy of Sciences, Beijing 100101, China

[6]University of the Chinese Academy of Sciences, Beijing 100049, China

[7]George W. Woodruff School of Mechanical Engineering, Georgia Institute of
Technology, Atlanta, Georgia 30332, USA

[*]Corresponding Author: Cheng Zhu, Email: cheng.zhu@bme.gatech.edu

[†]Present address: Medtronic CO., Minneapolis, MN, USA

[‡]Present address: Amgen Inc., One Amgen Center Dr., Thousand Oaks, CA, USA

Abstract

Central to T cell biology, the T cell receptor (TCR) integrates forces in its triggering process upon interaction with peptide-major histocompatibility complex (pMHC)[1-3]. Phenotypically, forces elicit TCR catch-slip bonds with strong pMHCs but slip-only bonds with weak pMHCs[4-9]. While such correlation is generally observed, the quantitative bond pattern and degree of "catchiness" vary. We developed two models based on the structure, elastic properties, and force-induced conformational changes of the TCR–pMHC-I/II complexes to derive from their bond characteristics more intrinsic parameters that underlie structural mechanisms, predict T cell signaling, and discriminate antigens. Applying the models to all published 50 datasets of 11 TCRs and their mutants interacting with corresponding pMHCs without coreceptor engagement revealed the ability for structural and physical parameters to quantitatively integrate and classify a broad range of bond behaviors and biological activities. Comparing to the generic two-state model for catch-slip bond that also fits the data, our models can distinguish class I from class II MHC systems and their best-fit parameters correlate with the TCR/pMHC potency to trigger T cell activation, which the generic model cannot. The extensive comparisons between theory and experiment provided strong validation of the models and testable hypothesis regarding specific conformational changes that control bond profiles, thereby suggesting structural mechanisms for the inner workings of the TCR mechanosensing machinery and plausible explanation of why and how force amplifies TCR signaling and antigen discrimination.

MAIN TEXT

Introduction

Antigen recognition via TCR–pMHC interactions is essential for T cell activation, differentiation, proliferation, and function[10]. Mechanical forces applied to TCR via engaged pMHC substantially increase antigen sensitivity and amplify antigen discrimination[4-9,11]. As a fundamental force-elicited characteristic, strong cognate pMHCs form catch-slip bonds with TCR where bond lifetimes increase with force until reaching a peak, and decrease as force increases further, whereas weak agonist and antagonist pMHCs form slip-only bonds with TCR where bond lifetimes decrease monotonically with increasing force[4-9]. However, the mechanism underlying the correlation between the force-lifetime pattern and the ability for force on TCR to induce T cell signaling remains unclear.

An intuitive hypothesis is that catch bonds prolong interactions, which allow the process of CD3 signal initiation to proceed sufficient number of phosphorylation steps to the threshold for downstream signal propagation, as proposed by the kinetic proofreading model[12]. However, this hypothesis faces two challenges upon scrutiny of multiple TCR–pMHC systems. First, since bond lifetime *vs* force profiles are monotonically decreasing for slip-only bonds but bell-shaped for catch-slip bonds, the type of bonds that would last longer depends on the force range, which may be catch-slip bonds in one force regime but slip-only bonds in another force regime. Second, TCR–pMHC interactions exhibiting catch-slip bonds often have longest lifetimes around 10-20 pN[4-8,13] and it has been reported that upon engaging pMHC, T cells would exert 12-19 pN endogenous forces on the TCR in a signaling-dependent fashion[14]. However, the relevance of this force range to T cell signaling remains incompletely understood. More perplexingly, some signal-inducing pMHCs form catch-slip bonds with TCRs but exhibit shorter lifetime than other pMHCs that do not induce signaling by, and form slip-only bonds with, the same TCRs even in the optimal force range[15]. These observations prompt the questions of what mechanism underlies the association of TCR–pMHC bond type with the T cell signaling capacity and what impact this specific force range

has on TCR mechanotransduction. To answer these questions requires an in-depth analysis of the

multiple datasets with mathematical models, which was lacking.

Slip and catch bonds refer to two opposite effects of physical force on biomolecular

interactions: increasing or decreasing their off-rate of dissociation, respectively[16,17]. Because force

tends to be disruptive and destabilizing, slip bonds are intuitive, whereas catch bonds are counter-

intuitive. Since excessive force can rupture even covalent bonds[18], continued force increase will

eventually overpower any catch bond, turning it to a slip bond after an "optimal" force where the

off-rate is minimal[4-9,17]. Slip bond behavior is commonly modeled by the Bell equation, which

assumes the dissociation rate k of a molecular bond along a single pathway in a one-dimensional

(1D), single-well energy landscape to be an exponential function of force, $k(F) = k_0 e^{\frac{\delta_0^* F}{k_B T}}$. Here,

k_0 is the transition rate at zero force, F is tensile force, δ_0^* is the force-free distance from the bound

state at the bottom of the energy well to the top of the energy barrier known as the "transition state",

k_B is the Boltzmann constant, and T absolute temperature[19]. Several models have been developed

to account for catch-slip bond behavior. Most introduced two dissociation pathways and/or two

bound states in a two-dimensional (2D) energy landscape that is tilted by force[20] (Fig. 1a). One

noticeable exception is that of Guo et al. where dissociation is modeled to start from a single bound

state along a single pathway through a 1D energy landscape based on the physical process of peeling

a polymer strand with force until the transition state is reached[21]. A distinct advantage of this model

is its ability to relate the force-induced tilting of the energy landscape to the force-induced

conformational change of the molecular complex (which all other models lack), thereby connecting

parameters of the abstract energy landscape to the structural-elastic properties of the interacting

molecules.

Besides binding properties of the TCR–pMHC complex, its structural features and

conformational changes have been suggested to be important for TCR triggering. For example,

TCR–pMHC docking orientation has been correlated to its ability to trigger T cell signaling[15,22,23].

Partial unfolding or allosteric regulation of either the TCR and/or MHC molecules has been inferred

from mechanical experiments and steered molecular dynamics (SMD) simulations of pulling single

TCR–pMHC bonds[6,8,24,25]. Whereas the extent of these conformational changes has been correlated

to the strength of TCR–pMHC catch bonds[6,8,24], the two phenomena have not been integrated into

a mathematical model to explore their potential connection.

Here we developed two such models, one for each MHC class, to describe both TCR catch-

slip and slip-only bonds. The model development follows Kramers' kinetic rate theory and uses

polymer physics models to construct a 1D energy landscape for single-state, single-pathway

dissociation that incorporates the structures, elastic properties, and force-induced conformational

changes of the TCR–pMHC-I/II complexes at the sub-molecular level, which includes domain

stretching, hinge rotation, and molecular extension. Incorporating the force-induced conformational

changes into the energy landscape formulation allows force to shift the energy barrier up in low

forces and down in high forces, thereby giving rise to catch-slop bonds along a single dissociation

pathway in a 1D energy landscape (Fig. 1b). Applying the models to analyze all TCR–pMHC bond

lifetime *vs* force datasets published to date in 10 papers by four laboratories measured using two

different techniques[4-8,13,15,26-29], including 50 datasets of 11 TCRs and their mutants interacting with

corresponding panels of both classes of pMHCs without coreceptor engagement, revealed the

ability of our models' structural and physical parameters to quantitatively integrate and classify a

broad range of catch-slip or slip-only bond behaviors and biological activities. The extensive

comparisons between theory and experiment allowed us to rigorously validate the models by

systematically testing their assumptions, to identify specific conformational changes that control

the bond profiles, to provide mechanistic insights into the inner workings of the TCR–pMHC

mechanosensory machinery by constructing the energy landscape and investigating its properties,

and to understand how force-elicited catch bond amplifies TCR signaling and antigen

123 discrimination by examining the correlation of the model parameters with the biological activities

124 of a large number of TCR–pMHC-I/II systems.

126 **Results**

127 **Model development**

128  *Model goal*: Kramers' kinetic rate theory treats bond dissociation as state transition in a 1D

129 energy landscape $\Delta G^*(\delta_1)$ from a free-energy well (bound state) over a barrier (transition state)

130 along the dissociation coordinate δ_1[30]. Following Guo et al.[21] to adapt the linear-cubic model of

131 Dudko et al.[31] but allow force F to tilt the original energy landscape by an amount of $-\delta_1\gamma(F)$, the

132 energy landscape takes the form of

$$\Delta G(\delta_1, F) = \frac{3\Delta G_0^*}{2}\left(\frac{\delta_1}{\delta_0^*} - \frac{1}{2}\right) - 2\Delta G_0^*\left(\frac{\delta_1}{\delta_0^*} - \frac{1}{2}\right)^3 - \delta_1\gamma(F) \ . \tag{1}$$

133 where δ_0^* and ΔG_0^* are the differences in dissociation coordinates and free-energy levels,

134 respectively, between the transition state and bound state of the original force-free energy

135 landscape. The corresponding force-dependent kinetic rate is

$$k(F) = k_0\sqrt{1 - \frac{2\delta_0^*\gamma(F)}{3\Delta G_0^*}}\exp\left(\frac{\Delta G_0^*}{k_\mathrm{B}T}\left(1 - \left(1 - \frac{2\delta_0^*\gamma(F)}{3\Delta G_0^*}\right)^{3/2}\right)\right) \tag{2}$$

136 where k_0 is the dissociation rate at zero force[21]. Letting $\gamma \sim F$ recovers from Eq. **2** the Dudko-

137 Hummer-Szabo (DHS) model[31], and further assuming $|2\delta_0^*\gamma(F)/(3\Delta G_0^*)| \ll 1$ reduces it to the

138 Bell model[19]. The condition for k to be able to model catch-slip bond is the derivative $k'(F_0) = 0$

139 where $F_0 > 0$. This translates to two conditions: the barrier height $\Delta G^* = \Delta G_0^*\left(1 - \frac{2\delta_0^*\gamma(F_0)}{3\Delta G_0^*}\right)^{3/2} =$

140 $k_BT/3$ or $\gamma'(F_0) = 0$. The first condition requires the energy change $-\delta_1\gamma$ induced by F_0 to lower

141 the energy barrier height to $k_\mathrm{B}T/3$ located at $\delta_0^*\left(\frac{k_BT}{2\Delta G_0^*}\right)^{1/3}$. The second condition requires γ to be a

142 biphasic function of F. This excludes the Bell model and the DHS model because both of their γ

143 functions depend on F monotonically. Our goal is to construct a biphasic $\gamma(F)$ with appropriate

144 structural-elastic dependency to account for TCR–pMHC conformational changes from the bound

state to the transition state, as analyzed by the biomembrane force probe (BFP) experiments where a tensile force is applied to its two ends to modulate bond dissociation[4-9].

Key model assumptions: Following the reasoning of Guo et al.[21], $\delta_l\gamma(F) = \int_0^F \delta_z(f)df$ where the integrand $\delta_z(f) = z(f) - z_0(f)$ is the projection on the force direction of the change induced by force f of the TCR–pMHC extension at the transition state relative to its extension at the bound state. For γ to depend on F biphasically as required for describing catch-slip bonds, δ_z should be a biphasic function of f as discussed later. Therefore, dissociation occurs because the system moves in the energy landscape along the dissociation coordinate δ_l from the bound state to the transition state by a distance $\delta^* = \delta_0^*\left(1 - \frac{2\delta_0^*\gamma(F)}{3\Delta G_0^*}\right)^{1/2}$ [21]. We assume that the differential contour length along the force transmission path across the TCR–pMHC structure (*i.e.*, summing up all contour lengths of various domains connected at nodes of force action, as depicted in red lines in Fig. 2a, for MHC-I) at the transition state l and bound state l_0 can serve as a dissociation coordinate, i.e., $\delta_l = l - l_0$. δ_z is the projection of δ_l on the z axis – the direction of the pulling force (Fig. 2a). When only contour lengths are considered, $\delta_l = \delta_0^*$, which serves as a criterion for finding the best-fit parameters (Fig. 2a, *left* and Supplementary Model Derivations, Eq. S7).

Suggested by single-molecule optical tweezers[6], BFP[8] and magnetic tweezers[8] experiments as well as steered MD (SMD) simulations[8], we assume that force-induced TCR–pMHC dissociation is accompanied by conformational changes in both the TCR and MHC molecules. Specifically, we assume that at the bound state, force induces elastic extension of the TCR–pMHC structure as a whole (Fig. 2a, *left*); but as the system moves toward the transition state for dissociation, conformational changes may occur, which may include disruption of intramolecular interfaces, hinge rotation, and partial unfolding of interdomain joints (Fig. 2a, *right*). To include appropriate details of these proposed conformational changes at the sub-molecular level into the expression of δ_z, we model the TCR–pMHC structure as a system of semi-rigid bodies representing the whole complex as well as various globular domains connected by semi-flexible polymers that allow

extension and hinge rotation under mechanical loads (Fig. 2a, *right*). Specifically, we assume that,

as the system moves along the dissociation pathway from the bound state toward the transition state,

force may induce disruption of the MHC $\alpha_1\alpha_2$–β_{2m} interdomain bond, thereby shifting the

mechanical load originally borne by this bond to the $\alpha_1\alpha_2$–α_3 joint to induce its partial unfolding,

as observed in SMD simulations[8]. As such, the MHC α_3 domain would change its length and rotate

about its C-terminus (Fig. 2b). Since the TCRβ subunit has also been proposed to undergo FG-

loop-regulated conformational change[6,24,25], we assume that disruption of the $\alpha_1\alpha_2$–β_{2m} joint would

also result in tilting of the TCR–pMHC bonding interface and shifting of the mechanical load from

the TCR Vβ–Cβ joint to the Vα–Cα joint, leading to partial unfolding of the Vα–Cα joint (Fig.

2b). This increased stretching of the Vα–Cα joint relative to the Vβ–Cβ joint is assumed to result

from strengthening of the Vβ–Cβ joint by the FG-loop[11]. At the transition state, therefore, we treat

the MHC α_3 domain ($d_{\alpha3}$), the MHC $\alpha_1\alpha_2$ domains bound to the TCR V$\alpha\beta$ domains ($d_{B.I}$), and the

TCR C$\alpha\beta$ domains ($d_{C\alpha\beta}$) as three semi-rigid bodies connected by two unfolded peptide chains of

the MHC $\alpha_1\alpha_2$–α_3 joint ($d_{p,MHC}$) and the TCR Vα–Cα joint ($d_{p,TCR}$) (Fig. 2b, *right*). At the bound

state, neither disruption of intramolecular bonds nor partial unfolding of interdomain joints occurs,

as mentioned earlier, allowing the whole TCR–pMHC ectodomain complex to be modeled as one

semi-rigid body (d_N).

Force-induced energy change: To derive an expression for the last term on the right-hand

side of Eq. 1, we model the semi-rigid bodies d_i (i = N, α_3, B.I., and C$\alpha\beta$ to denote the whole TCR–

pMHC ectodomain structure as well as its indicated domains) as three-dimensional freely-jointed

chains (FJC) and employ polymer physics to obtain their force-extended length $d_i(f)$ from their

force-free length $d_{i,c}$[32] (Supplementary Model Derivations, Eq. S1).

The assumed partial unfolding of the $\alpha_1\alpha_2$–α_3 joint and the Vα–Cα joint is based on

suggestions from single-molecule optical tweezers[6], BFP[8] and magnetic tweezers[8] experiments as

well as SMD simulations[8]. We model these unstructured polypeptides as extensible worm-like chains (eWLC) and employ polymer physics to obtain their force-induced extension $d_{p,i}(f)$ ($i =$ MHC and TCR) from their force-free, folded state, which has zero length[33] (Supplementary Model Derivations, Eq. S3).

Upon projecting the various force-induced extensions described above onto the force axis, we obtain z components of five contributions to the TCR–pMHC length increase at the transition state: extension of the MHC α_3 domain ($z_{\alpha 3}$), unfolding of the MHC $\alpha_1\alpha_2$–α_3 interdomain joint ($z_{p,MHC}$), extension of bonding interface that includes the MHC $\alpha_1\alpha_2$ domains bound to the TCR V$\alpha\beta$ domains ($z_{B,I}$), unfolding of the Vα–Cα joint ($z_{p,TCR}$), and extension of the TCR C$\alpha\beta$ domains ($z_{C\alpha\beta}$) (Fig. 2b). Finally, we obtain:

$$\delta_1\gamma(F) = \int_0^F \left[z_{\alpha 3}(f) + z_{p,MHC}(f) + z_{B,I}(f) + z_{p,TCR}(f) + z_{C\alpha\beta}(f) - z_N(f)\right]df \qquad (3)$$

Model characterization

Model constants and parameters: The FJC model constants for the 1st, 3rd, 5th, and 6th terms in the integrand on the right-hand side of Eq. 3 include the force-free lengths $d_{i,c}$ and the elastic modulus of the folded globular domain E_c, all of which are available from the literature. The 2nd and 4th terms are proportional the respective numbers of amino acids in the polypeptides of the partially unfolded MHC $\alpha_1\alpha_2$–α_3 joint ($n_{p,MHC}$) and TCR Vα–Cα joint ($n_{p,TCR}$), which can be combined as the product of the total unfolded amino acid number $n^* = n_{p,MHC} + n_{p,TCR}$, the average contour length per unfolded amino acid l_c, and the extension per unit contour length $z_{u,p}(f)$. The eWLC model constants for $z_{u,p}(f)$ include the average persistence length per unfolded amino acid l_p and the elastic modulus of polypeptides E_p (Supplementary Table 1).

After applying model constraints and the approximating $\Delta G_0^* \sim \ln(k_w/k_0)$ where $k_w \sim 10^6$ s^{-1} is known as the prefactor (Supplementary Model Derivations), the model parameters are reduced to 5: 3 structural parameters ($d_{\alpha 3}, \theta, n^*$) and 2 biophysical parameters (k_0, δ_0^*), for describing

dissociation of TCR–pMHC-I bonds. We will determine these parameters by comparing the model

predictions with experimental measurements, and by doing so, illustrate the ability of our model to

use a relatively low number of parameters to capture the coarse-grained structure and

conformational changes at the sub-molecular level during TCR–pMHC-I dissociation.

Model features and properties: To explore the general features and properties of the model,

we plotted δ_z vs F for two n^* values and a range of θ values as well as two θ values and a range of

n^* values (Fig. 2c). Conceptually, force-heightened energy barrier generates catch bonds and force-

lowered energy barrier produces slip bonds (Fig. 2d, *top left* and *right* panels). Since $-\delta_l \gamma$

represents the energy input by force F to the original energy landscape, a biphasic $\delta_l \gamma$ is required

to create catch-slip bonds (Fig. 2d, *top middle* panel); correspondingly, δ_z is required to have a root

at positive F where catch bond transitions to slip bond (Fig. 2c). The parameter domains capable

of generating catch, catch-slip, and slip bonds are mapped on an n^*-θ phase diagram (Fig. 2d),

showing that our model can describe catch-slop bond if and only if $n^* > 0$, $\theta > 0$, and $\delta_l \gamma(\infty) >$

0 (Fig. 2d, *top middle* panel). In other words, catch-slip bonds require partial unfolding of the MHC

$\alpha_1\alpha_2$–α_3 and/or TCR Vα–Cα joints and tilting of the TCR–pMHC bonding interface, a prediction

consistent with previous results of SMD simulations and single-molecule experiments[8].

For single-bond dissociation from a single bound state along a single pathway, the reciprocal

dissociation rate should equal to the average bond lifetime. Regardless of the bond type, the

reciprocal zero-force off-rate controls the *y*-intercept of the bond lifetime *vs* force curves. We

plotted the theoretical bond lifetime (normalized by its zero-force value) k_0/k *vs* force F for a

range of n^*, θ, and δ_0^* to examine how the model parameters control the bond lifetime *vs* force

profile (Fig. 2e). Consistent with Fig. 2c-d, only if $n^* > 0$ and $\theta > 0$ can our model describe catch-

slip bond. Increasing the tilting angle θ results in more pronounced catch-slip bonds with longer

lifetimes that peak at higher forces (Fig. 2e, *upper panel*). By comparison, increasing δ_0^* changes

the level of slip-only bonds if $n^* = 0$ and $\theta = 0$, but prolongs lifetime of catch-slip bonds (until

cross-over at a higher force) without changing the force where lifetime peaks if $n^* > 0$ and $\theta > 0$

(Fig. 2e, *lower panel*).

To understand physically how our model describes catch-slip bonds, we plotted the energy

landscape using Eq. 1 (Fig. 2f). Setting $\theta = 0$ in the upper panel generates a family of ΔG *vs* δ_l

curves where the energy barrier is suppressed monotonically with increasing force, indicating a

slip-only bond (Fig. 2f, *upper panel*). By comparison, setting $\theta > 0$ results in a family of ΔG *vs* δ_l

curves where the energy barrier height is initially raised, then lowered, by increasing force,

indicating a catch-slip bond (Fig. 2f, *lower panel*, also see Fig. 1b). We also examine how the

transition state location (Fig. 2g, *upper row*) and energy barrier height (Fig. 2g, *lower row*) change

with force for a range of θ and δ_0^* values that give rise to slip-only bonds and catch-slip bonds.

Noticeably, at fixed θ values, both rates by which the transition state location and the energy barrier

height change with force are accelerated by increasing δ_0^* (Fig. 2g, *right column*), suggesting that

this parameter can be used as a measure for force sensitivity. Interestingly, increasing θ slows the

decrease in both the transition state location and energy barrier height with force at higher values,

suggesting that the tilting angle controls the range at which force sensitivity can last (Fig. 2g, *left*

column).

Model validation

Model's capability to fit data: To test our model's validity, we used it to reanalyze data of

various TCRs forming catch-slip bonds and slip-only bonds with their respective panels of pMHCs.

These include three datasets published by us: 1) murine OT1 TCR expressed on either primary

naïve CD8$^+$ T cells or CD4$^+$CD8$^+$ thymocytes, which interacted with peptides presented by a mutant

(MT) MHC (H2-K^{b}α3A2) that abolished CD8 co-engagement[4,13] (Fig. 3a and Supplementary Fig.

1a); 2) wild-type (WT) or MT murine 2C TCRs either expressed on primary naïve CD8$^+$ T cells or

re-expressed on CD8$^-$ hybridoma cells, which interacted with peptides presented by H2-K^{b}α3A2

(for CD8$^+$ primary T cells) or H2-K^b (for CD8$^-$ hybridoma cells) to ensure no CD8 contribution

either way, or by H2-L^d(m31), a different MHC allele from H2-K^b (Fig. 3b, Supplementary Fig.

1b-c)[8]; and 3) WT and three MT human 1G4 TCRs were re-expressed on hybridoma cells, which

interacted with the melanoma peptide NY-ESO-1 bound to HLA-A2 (Supplementary Fig. 1d)[8]. In

addition, we fitted our model to two datasets published by other labs: 1) soluble mouse N15 TCR$\alpha\beta$

interacted with VSV and two MT peptides bound to H2-K^b (Supplementary Fig. 1e)[6] and 2) four

mouse TCRs expressed on hybridomas interacted with NP$_{366}$ bound to the D227K mutant of H-2D^b

to prevent CD8 binding[15] (Supplementary Fig. 1f). Gratifyingly, the theoretical reciprocal force-

dependent off-rate $1/k(F)$ fits all 39 experimental bond lifetime vs force curves well (Fig. 3a-b,

Supplementary Fig. 1, and Supplementary Table 2 and 3), demonstrating our model's capability to

describe a wide range of data.

   Characterization of force-lifetime relationship: Previous work reported qualitative

correlations between the TCR bond type, *i.e.*, catch-slip bond vs slip-only bond, with the biological

activity of the peptide to induce T cell activation, *i.e.*, pMHC potency[4,7-9,13]. To reduce data

representation and extract more information quantitatively from the bond lifetime vs force data, we

defined several metrics from their model fit for each TCR–pMHC system and examined their

correlation with T cell activation induced by a given interaction, using the OT1 system as an

example because the quantitative ligand potency data are available[4,34]. We measured the peak bond

lifetime, t_{peak}, and the change, Δt, from t_{peak} to the force-free bond lifetime, $t_0 = 1/k_0$ (Fig. 3a,

left panel). We found the relative metric Δt more suitable for comparison across different TCR

systems, and better correlates with ligand potency, than the absolute counterpart t_{peak} (Fig. 3c).

Although the force where catch-slip bond lifetime peaks, F_{opt}, occurs in a narrow range (10-20 pN),

the force increment, ΔF, from F_{opt} to the level where bond lifetime returns to t_0 defines the force

span of a catch-slip bond (Fig. 3a, *left panel*). One scaled parameter, $L = \Delta t/t_{peak}$ (relative length

of lifetime), does (Fig. 3d, *left panel*), but the other, $B = \Delta F/F_{opt}$ (relative breadth of lifetime),

does not (Supplementary Fig. 2a), correlate with ligand potency well. Remarkably, the combined

scaled parameter termed the catch bond intensity or catchiness, $I = L/(1 + B)$, correlates best with

the ligand potency across different TCR systems (Fig. 3d, *right panel* and 3e), supporting its

usefulness and reasonableness as a metric of reduced data representation for catch and slip bonds.

Model parameters' correlation to ligand potency: It seems reasonable to test the validity of

our model by examining the possible correlation of (or the lack thereof) the model parameters with

features of the biological system, e.g., the ligand potency. The rationale is that if its parameters are

capable of capturing and predicting such biological features, then the model would be more

meaningful and useful than merely a curve fitting tool. Therefore, we plotted the tilted angle of the

bonding interface θ, the number of the unfolded amino acids n^*, and the width of the zero-force

free-energy well δ_0^* that best-fit the force-lifetime curves of OT1, 2C, N15, TRBV TCRs interacting

with their corresponding panels of pMHCs (Fig. 3f). Gratifyingly, we observed good correlation

between each model parameter and the peptide potency for all published data of TCR–pMHC-I

catch-slip bonds and slip-only bonds from four independent laboratories[4,6,8,13,15].

In a previous study, we mutated residues in the 2C or 1G4 TCR and/or their corresponding

pMHCs to alter bond profiles as predicted by SMD simulations, which was confirmed by

experiment[8]. We therefore fitted our model to the force-lifetime curves of these mutant TCR–

pMHC bonds, which evaluated the best-fit model parameters, δ_0^* , θ , and n^* (Fig. 4a,

Supplementary Fig. 2b-c). In the absence of other functional data, we took an indirect approach to

examine their correlations with the catch bond intensity I of these bond lifetime *vs* force curve (Fig.

4b) since I and all three model parameters correlate with the peptide potency (Fig. 3e and f). Results

are exemplified by the δ_0^*, θ, and n^* *vs* I plots, which are graphed together with the data without

TCR and MHC mutations that already showed functional correlates. For the WT OT1, 2C, N15,

and TRBV TCRs interacting with their corresponding panels of pMHCs, δ_0^* (Fig. 4c), and to a lesser

extent, θ (Supplementary Fig. 2d) and n^* (Supplementary Fig. 2e), correlate with I that predicts

peptide potency (blue-open symbols). Remarkably, for the 2C and 1G4 TCRs specifically mutated

to alter bond profiles with the corresponding WT or mutant MHCs presenting the same agonist peptide, their best-fit δ_0^*, and to a lesser extent, θ and n^*, also correlate well with I (Fig. 4c and Supplementary Fig. 2d-e, green-closed symbols). Interestingly, $1/k_0$ shows a negative correlation with I (Supplementary Fig. 2f), consistent with reports that zero-force bond lifetime does not correspond to ligand potency in these cases[4,8,34].

It is worth pointing out that the above results not only support our model's validity but they also suggest that our model is more than a mere analytical framework to organize experimental data. Rather, the model parameters may be used to discriminate antigen and predict the force-amplified discriminative power. For example, the correlations of θ and δ_0^* with peptide potency (Figs. 3f and 4c) indicate that the more potent the peptide, the higher the force sensitivity of its TCR–pMHC interaction, and the narrower the force range over which the TCR–pMHC interaction is sensitive to force (Fig. 2g).

Comparison between coarse-grained and all-atom models: Bonding interface tilting has been observed to be associated with changes in the number of hydrogen bonds bridging the TCR and pMHC molecules as they were forced to unbind in SMD simulations[8]. Therefore, we investigated whether, and if so, how well the tilting angle would correlate with the change of hydrogen bonds between TCR and MHC. Remarkably, θ was found to be proportional to the net change in the total number of hydrogen bonds at the bonding interface (Fig. 4d and Supplementary Fig. 3). This finding is intuitive and supports the validity of our coarse-grained model because it is able to recapitulate the results of all-atom SMD simulations[8].

Classification of bond types by clustering analysis on phase diagrams: In Fig. 2e-g we have explored the model parameter space to identify regions that correspond to slip-only bonds and catch-slip bonds. Here we examined whether, and if so, how parameters that best-fit different experimental bond types map onto different regions of the parameter space. Since the model has four parameters, θ, δ_0^*, $d_{\alpha 3}$, and n^* (k_0 is not considered because if its lack of correlation with catch

bond intensity), we analyzed their clustering and projected their values in the 4D parameter space onto three phase diagrams spanning the θ-δ_0^* (Fig. 4e), θ-$d_{\alpha3}$ (Figs. 4f and Supplementary Fig. 4), and δ_0^*-n^* (Fig. 4g) 2D space. Clustering analysis of the model parameters that best-fit 36 published TCR–pMHC bond lifetime *vs* force curves (Supplementary Fig. 5) shows three distinct clusters in the δ_0^* *vs* θ and θ *vs* $d_{\alpha3}$ plots as well as δ_0^* and θ *vs* n^* plots (Fig. 4e-g), which classify the TCR–pMHC interactions into slip-only (SO), weak catch-slip (WC) and strong catch-slip (SC) bonds, which correspond to weak, intermediate, and strong potencies for pathogenic peptides and their variants. Whereas transition in bond type from SO to WC and SC requires monotonical increase in θ and n^* (Fig. 4f, g), the corresponding change in δ_0^* is non-monotonic (Fig. 4e-g). SO bonds show small n^*, δ_0^*, and θ values. WC and SC bonds observed from published experiments are best-fitted by similar n^* (5 for WC and 7 for SC) but oppositely ranked δ_0^* and θ values. To change from WC to SC bonds requires decreasing δ_0^* (from 2.7 to 1.9 nm) but increasing θ (from 14 to 32 °) (Fig. 4g). We also performed principal component analysis and calculated the Mahalanobis distances of the principal axes for the three bond types[35], which are statistically separated in the catch bond intensity *vs* Mahalanobis distance plot (Fig. 4h). Interestingly, WC and SC bonds show distinct conformational changes despite their similar *I* values measured from the force-lifetime curves. The corresponding structural features of these three types of bonds are depicted in Fig 4i, which have been observed in our previous SMD studies[8]. Of note, model parameters visualized by SMD simulations are usually larger than their best-fit values, which may have two explanations: First, to enable dissociation to be observed in the affordable computational time, much higher forces were used in simulations than experiments to accelerate the biophysical process, which likely induced much larger conformational changes. Secondly, our model describes the average conformational

change during the entire dissociation process, which is smaller than the maximum conformational

changes likely to occur right before unbinding and to be captured by SMD.

Our structure-based model is superior to the generic two-pathway model: It seems that

other published catch bond models should also be able to fit the experimental force-lifetime profiles

analyzed here, given their relative simple shapes. As an example, we examined the two-pathway

model below[36]:

$$k(F) = k_{0,p1}e^{\delta_{0,p1}^* F/k_B T} + k_{0,p2}e^{\delta_{0,p2}^* F/k_B T} \tag{4}$$

where $k_{0,p1}$ and $k_{0,p2}$ are the respective zero-force off-rates of the first and second pathway, $\delta_{0,p1}^*$

and $\delta_{0,p2}^*$ are the respective distances from the bound state to the transition states along the first and

second pathway (Fig. 1a). Thus, the off-rate for each pathway takes the form of the Bell model, but

the catch pathway parameter $\delta_{0,p1}^*$ has a negative value[36] (Supplementary Table 4). This model is

generic as it has previously been applied to TCR–pMHC catch-slip bonds without considering the

specific conformational changes[6].

As expected, Eq. 4 also fitted our experimental data with goodness-of-fit measures

statistically indistinguishable to Eq. 2 (Supplementary Figs. 1 and 10a-b). However, the fitting

parameters correlate with neither the TCR/pMHC-I potency to induce T cell function nor the

intensity of catch bond (Supplementary Fig. 6, see the negative or zero correlation and the poor R^2

values), hence have no biological relevance. This comparison indicates that the model developed

herein is superior to the previous two-pathway model.

Model for TCR catch bonds with class II pMHC

MHC class II differs from class I in three main aspects (comparing Fig. 2a and Fig. 5a): (1)

MHC-I has three α domains and a β$_{2m}$ domain whereas MHC-II has two α and two β domains. (2)

MHC-1 anchors to the T cell membrane through a single linker to the α$_3$ domain. The β$_{2m}$ domain

attaches to the α$_3$ domain instead of anchoring to the T cell membrane directly. By comparison,

MHC-II anchors to the membrane through two linkers, one to the α$_2$ domain and the other to the β$_2$

domain. (3) The peptide is presented by the α_1-α_2 domains of MHC-I but the α_1-β_1 domains of

MHC-II. These structural differences alter how forces are supported by and transmitted through,

and induce conformational changes in, the TCR complexes with pMHC-I vs pMHC-II. Thus, it is

necessary to modify the previous model in order for it to describe TCR catch and slip bonds with

MHC-II, which is done by using a different $\delta_1\gamma(F)$ expression than Eq. 3 (Supplementary Model

Derivations, Section B). This modification assumes force-induced partial unfolding and stretching

of the TCR Vα-Cα joint and the MHC α_1-α_2 and β_1-β_2 joints during dissociation, which results in

tilting of the bonding interface (Fig. 5a, b).

In the class II model, the same parameters δ_0^*, n^*, and θ are used but the MHC contribution

to n^*, $i.e.$, $n_{\mathrm{p,MHC}}$, represents the average number of amino acids in the polypeptides of the partially

unfolded MHC-II α_1-α_2 and β_1-β_2 joints instead of the MHC-I $\alpha_1\alpha_2$–α_3 joint, and the relationships

between θ to other structural parameters are also different from the class I model (Fig. 5b and

Supplementary Model Derivation, Section B). Like the class I model, the k_0/k vs F plots for a

range of n^*, θ, and δ_0^* in Fig. 5c show similar features to Fig. 2e and meet our objective of being

capable of describing catch-slip bonds if and only if $n^* > 0$ and $\theta \geq 0$. Unlike the class I model, a

much smaller θ value (< 10°) is seen in the class II model (compared Fig. 5c and Supplementary

Fig. 7 with Fig. 2e-g), indicating the main conformational change responsible for TCR–pMHC-II

catch-slip bond is unfolding rather than tilting. The validity of this model is supported by its

excellent fitting to our published data of mouse WT or MT TCRs on naïve T cells (3.L2) or

hybridomas (2B4) interacting with their respective p:I-E^k ligands[5,28] (Fig. 5d, Supplementary Table

5 and 6). The four metrics of the TCR–pMHC-II bond lifetime vs force curves Δt, L, I, and t_{peak}

correlate well with the published peptide potency[37,38] (Fig. 5e and Supplementary Fig. 8). Since

mutations on the 2B4 TCR were designed to perturb its C$\alpha\beta$ interaction with the CD3 subunits,

their effect on the bond profile and T cell function might be weaker than the effects of the altered

peptide ligands for the 3.L2 TCR[5,28]. Nevertheless, the correlations between ligand potency and the

three model parameters θ, n^*, and δ_0^* for the 2B4 system are comparable to those for the 3.L2

system (Fig. 5f), supporting the ability of the metrics of the bond profile and the model parameters

to recognize the change in the TCR-CD3 ectodomain interaction in addition to the ability to

discriminate antigen. These properties are desirable, intuitive, and are consistent with the parallel

properties found in the class I model. Similar to the class I model parameters, δ_0^* correlates well

with the catch bond intensity for the pooled results from all published class II data (Fig. 5g, *left*),

but θ and n^* correlate less well with I (Fig. 5g, *middle and right*). Thus, the validity of the class II

model is further supported by the faithful mapping of the relationship between biophysical

measurements of catch and slip bonds and biological activities of the TCR–pMHC-II interactions

onto a relationship between model parameters and biological function.

As expected, Eq. 4 also fitted our experimental data with goodness-of-fit measures

statistically indistinguishable to Eq. 2 (Fig. 5d and Supplementary Fig. 10c-d). Similar to the case

of class I system, the fitting parameters correlate with neither the TCR/pMHC-II potency to induce

T cell function nor the intensity of catch bond (Supplementary Fig. 6, see the negative or zero

correlation and the poor R^2 values), hence have no biological relevance. This comparison again

indicates that the model developed herein is superior to the previous two-pathway model.

Cross-examination of class I model against class II data and *vice versa*

Upon examining the catch-slip and slip-only bond lifetime *vs* force curves in Figs. 3a, 3b,

5d and Supplementary Fig. 1, it became apparent that the data seem very similar regardless of

whether they are for class I or class II pMHC. Indeed, applying the class I model to the class II data

and *vice versa* reveals that both models are capable of fitting both datasets well (Supplementary

Fig. 9) and produce statistically indistinguishable goodness-of-fit measures (Supplementary Fig.

10). This is not surprising because both models have five fitting parameters and the bond lifetime

vs force curves have relatively simple shapes. Nevertheless, fitting the same data by different

models returns different parameter values depending on the model used, because the two models

are constructed based on different structures and force-induced conformational changes of the

TCR–pMHC complexes. Therefore, we asked whether the best-fit model parameters were capable

of distinguishing data from the two classes of pMHCs and of telling whether a correct model was

used to analyze data of matched MHC class. To answer these questions, we plotted δ_0^* vs I (Fig. 6a,

b) and δ_0^* vs n^* (Fig. 6d, e) using values of the two models that best-fit the data of OT1, 2C, 1G4,

N15, and TRBV TCRs interacting with their respective panels pMHC-I ligands (Fig. 6a, d) as well

as 3.L2, 2B4, and E8 TCRs interacting with their respective panels of pMHC-II ligands (Fig. 6b,

e). Surprisingly, the dependency of δ_0^* on I is 2~3-fold stronger (i.e., steeper slope) (Fig. 6c),

indicating a greater discriminative power of ligand potency, for the matched than the mismatched

cases. Furthermore, it is well-known that the average contour length per a single amino acid l_c is ~

0.4 nm[21,39,40], which sets the biophysical limit for the slope of δ_0^* vs n^* plots. Indeed, we found that

the slopes of the δ_0^* vs n^* plots are within this limit for both model fits of the class I data (Fig. 6f).

For the class II data, the slope of the δ_0^* vs n^* plot reaches this limit for the matched case but exceeds

this limit for the mismatched case (Fig. 6f). Moreover, the goodness-of-fit (R^2) values of the linear

fit to the δ_0^* vs I (Fig. 6c) and δ_0^* vs n^* (Fig. 6f) data are much greater for the matched than the

mismatched cases, indicating more appropriate models for the data in the matched than the

mismatched cases. Indeed, the R^2 value for fitting the class I data by the class II model is too small

to be statistically reasonable, therefore telling the mismatch between the model and the data. These

results indicate that the model parameters are capable of distinguishing data from the two classes

of pMHCs.

Discussion

With the exception of a recent paper that observed T cell pulling on TCR by ~2 pN forces

using spider silk peptide-based force probe[41], five publications from two laboratories demonstrated

that TCR experienced endogenous forces of 12-19 pN using DNA-based force probes[13,14,42-44].

Also, except for a single paper that failed to observe catch bonds in the 1G4 TCRαβ ectodomain using a flow chamber[45], extensive data in 10 papers from four laboratories have demonstrated catch bonds in 11 membrane TCRs (including 1G4) using biomembrane force probe and optical tweezers[4-8,13,15,26-28]. These experiments prompted us to develop two mathematical models for TCR catch bonds, one with class I and the other with class II pMHC, based on Kramer's kinetic theory and accounted for the 3D coarse-grained structures, molecular elasticity, and conformational changes of the TCR–pMHC-I/II complexes. Previously, several models have been developed to describe catch-slip bonds of selectins–ligands[36,46-48], integrins–ligands[49,50], platelet glycoprotein Ibα–von Willebrand factor[51] cadherin–catenin[52], vinculin–actin[53], talin–actin[54] interactions. Except for the sliding-rebinding model, which is based on force-induced conformational changes in P-selectin–ligand observed from SMD simulations[48], none of these models have included any specific structural considerations of the interacting molecules. Instead, these models are based on a generic physical picture of dissociation along two pathways from either one or two bound states in a 2D energy landscape that is tilted by force[20,55]. Except for the two-pathway model tested here, which has 4 parameters[36], all other models have 5-10 parameters; therefore, over-fitting is a concern for applying them to some of the datasets analyzed here. Although the two-pathway model is capable of fitting the TCR–pMHC catch-slip bond data well, as shown previously[6] and tested more extensively by much larger datasets here (Supplementary Fig. 1 and Fig. 5d), the four best-fit model parameters correlate with neither T cell effector function nor the catch bond intensity for either class I or class II system (Supplementary Fig. 6), hence informing no biological insights. Consequently, it cannot distinguish the class I and class II systems because the parameters of the generic two-pathway model have nothing to do with the differential structures of the class I and II systems. This comparison highlights the utility and usefulness of our models.

Force-induced conformational changes of TCR–pMHC-I complexes have been observed or suggested by single-molecule experiments and SMD simulations[6,8]. Parameterizing these

conformational changes by the number of unfolded amino acids n^* and the bonding interface titling

angle θ in the class I model allows us to explain mechanistically and quantitatively the TCR–

pMHC-I catch-slip and slip-only bonds. Indeed, the criteria for catch-slip bond are $n^* > 0$ and $\theta >$

0; the greater their values the more pronounced the catch bond. Importantly, the validity of the class

I model has been supported by its capability to fit all force-lifetime data published to date and by

the correlation between the best-fit model parameters and the available biological activity data

induced by the TCR–pMHC-I interactions.

By comparison, the respective ranges of n^* and θ for the class II model are smaller,

consistent with the sturdier structure of the pMHC-II molecule[23]. Neither experimental nor

simulation data have been reported to suggest force-induced conformational changes in the TCR–

pMHC-II structure. Our class II model has also been tested by all published data and their best-fit

parameters also correlate well with the biological activities induced by the TCR–pMHC-II

interactions. Furthermore, the validity of models of both classes has been supported by the findings

that the best-fit model parameter δ_0^* correlates with the catch bond intensity I and that the δ_0^* *vs* I

and δ_0^* *vs* n^* plots have more appropriate slopes and R^2 values when the model matches than

mismatches the data.

We should note that since TCR–pMHC unbinding is assumed to be a spatially continuous

and temporally instantaneous process, all structural parameters determined here represent mean

field values, and they were evaluated by fitting the mean bond lifetime *vs* force data. However,

individual bond dissociation events are inherently stochastic; and as such, need not be

deterministically mapped onto any specific conformational changes in a one-on-one fashion.

Instead, any particular bond type and parameter sets are related on the average sense. Future studies

are required to extend the current framework to relate more detailed structural changes and bond

lifetime distributions, *e.g.*, to account for more sequential partial unfolding events prior to transition

state as suggested by experiments[6,11,29].

A strength of our agent-based models lies in their ability to incorporate many different ideas

and knowledge into a simple 1D formulation. This simplicity facilitates model application to both

class I and II experimental systems, enables quantitative interpretation of TCR–pMHC bond

lifetime *vs* force profiles, expresses biological functions by biophysical measurements, and suggests

structural mechanisms of how the TCR mechanotransduction machinery might work. However, the

1D simplification is also a weakness because theoretically these models can only describe single-

step dissociation by entropic conformational fluctuation in the low-force regime from a single-state

along a single-dissociation path, implicitly assuming that there is only a single energy barrier.

Although some catch-slip and slip-only bonds can be described by such simple models[5], more

complicated TCR–pMHC bonds has been reported. These are evidenced by the multi-exponential

bond lifetime distributions at constant forces, which have been fitted by data-driven multi-state,

multi-pathway models[27,28]. To address this weakness, future studies may extend the present 1D

model to 2D, e.g., by combining Eqs. 2 and 4, to enable proper description of multi-exponential

bond survival probabilities.

We introduced the catch-bond intensity I as a dimensionless scaled metric for the bond

lifetime *vs* force curve and generated four model parameters that describe the curve's geometric

features. Upon analyzing all catch-slip and slip-only bond profiles (a total of 50 curves) published

to date by four independent laboratories[4-8,13,15,26-28], we found that these quantities do a better job to

predict TCR function than any other quantities. This finding may explain how force amplifies TCR

signaling and antigen discrimination, because I is defined by a force curve and n^* and θ only

predict signaling when they assume none-zero values at $F > 0$.

A recent study showed surprising features of reversed-polarity of TRBV TCRs such that

interactions of NP_{366}:$H-2D^{bD227K}$ to TCRs B13.C1 and B17.C1 induced T cell signaling, whereas

interactions of the same pMHC to B17.R1 and B17.R2 TCRs did not[15]. Despite that the former two

TCRs formed catch-slip bonds with NP_{366}:$H-2D^{bD227K}$ and the latter two TCRs formed slip-only

bonds, the authors suggested that the signaling capability of the B13.C1 and B17.C1 TCRs could not be attributed to their force-prolonged bond lifetimes because the B17.C1 TCR–H-2D^{bD227K} bond was shorter-lived than the B17.R2 TCR–NP$_{366}$:H-2D^{bD227K} bond across the entire force range tested. Even at 9.4 pN, which was F_{opt} for the former with a $t_{peak} = 0.61$ s, the latter lived 2.48 s on average, and the longest lifetime of the latter was $t_0 = 2.83$ s occurred at zero force[15]. The authors hypothesized that the TCR–pMHC docking orientation, which was "canonical' for the B13.C1 and B17.C1 TCRs but "reversed" for the B17.R1 and B17.R2 TCRs, underlain the signaling outcomes by directing the position of Lck relative to the CD3. However, we suggest that even without knowing the docking orientation, our model parameters are capable of determining the signaling outcomes. Indeed, our analysis correctly maps the data of the B13.C1 and B17.C1 TCRs onto the high peptide potency region and the data of the B17.R1 and B17.R2 TCRs onto the low peptide potency region of the δ_0^* vs I (Fig. 4c), δ_0^* vs θ (Fig. 4e), and θ vs $d_{\alpha3}$ (Fig. 4f) phase diagrams. Thus, by mechanistically modeling the effect of force on bond dissociation, TCR signaling and antigen discrimination can be predicted by the model parameters.

The success in our model applications indicate that the conformational changes assumed in the models may be important to the TCR triggering, thereby suggesting testable hypotheses for future studies designed to investigate the inner workings of the TCR mechanotransduction machinery, *e.g.*, to extend and/or revise models regarding how TCR signaling is triggered. One of the TCR triggering models involves conformational changes and catch bond formation[1,2,56]. Our structure-based models relate catch and slip bonds to TCR–pMHC conformational changes. For the class I model, the parameterized structural changes include force-induced disruption of the MHC $\alpha_1\alpha_2$–β_{2m} interdomain bond, partial unfolding of the $\alpha_1\alpha_2$–α_3 joint, tilting of the TCR–pMHC bonding interface, and partial unfolding of the Vα-Cα and Vβ-Cβ joints. For the class II model, these are primarily limited to the force-induced partial unfolding of the MHC-II α_1-α_2 and β_1-β_2 joints as well as the Vα-Cα and Vβ-Cβ joints. Besides these, one additional conformational change

observed in the SMD simulations of TCRαβ–pMHC dissociation is unfolding of the connecting

peptides between the TCRαβ ectodomain and transmembrane domain[8]. We chose not to include

this conformational change in our models because such unfolding would likely be prevented by the

interaction of the Cαβ with the CD3 subunits[28]. Consistent with this assumption, the experimental

data used for model fitting to evaluate conformational change parameters (n^* and θ) are those of

pMHC bonds with TCR-CD3 complexes on the cell membrane that includes the TCRαβ ligand

binding subunits and the CD3 signaling subunits (except for the N15 TCRαβ case which is soluble

ectodomain only). Indeed, our previous work found that catch bonds of purified TCRαβ were

altered from those of cell surface TCR interacting with the same pMHCs[26,27], which is reflected by

their changed model parameters (Supplementary Fig. 11). As such, the TCRαβ conformational

changes predicted by our models provide a constraint for possible CD3 conformational changes in

the TCR-CD3 complex to be considered in future TCR triggering models.

Another constraint to be considered by future studies is that imposed by the coreceptor CD4

and CD8, as the co-ligation of the coreceptor prolongs bond lifetimes, amplifies catch bonds, and

may even changes slip-only bonds to catch-slip bonds[7,13,27]. Future studies should also consider

how to extend the current models to pre-TCR catch and slip bonds with a broad range of

ligands[29,57]. Instead of the TCRα, the pre-TCR uses the TCRβ chain to dimerize with a common

pre-Tα chain, which lacks the variable domain (hence no Vα-Cα hinge). Without extension, even

if our models are still able to fit the data of pre-TCR–ligand bonds or data of TCR–pMHC bonds

where the TCRβ F-G loop was deleted or bound by an anti-TCR antibody[6], the best-fit model

parameters may not correspond to the conformational changes of these bonds which are likely

different from the conformational changes in the TCR–pMHC bonds with intact F-G loop.

An objective of the present work is to explore the extent to which 1D models can describe

experimental data with a minimal set of meaningful parameters. Our parameters consider coarse-

grained structural features and relate catch and slip bonds to specific force-induced conformational

changes of the TCR–pMHC complex. This approach should be extendable to the modeling of other

receptor–ligand systems of different structural features yet also form catch and slip bonds, such as

selectins[17,58,59], integrins[49,50,60-63], cadherin[64], Fcγ receptor[65], notch receptor[66], platelet glycoprotein

Ibα[51,67], FimH[68], actin with myosin[69], actin with actin[70,71], cadherin-catenin complex with

actin[52], vinculin with F-actin[53], and microtubule with kinetochore particle[72].

Our models allow us to develop working hypotheses regarding how T cell function is

regulated through structural modulations of catch and slip bonds. For example, our class I model

predicts that strengthening of the $\alpha_1\alpha_2$–β_{2m} interdomain bond would weaken the TCR–pMHC catch

bond, which would in turn reduce T cell activation. This prediction has been supported by our

published data that somatic mutations in HLA-A2 found in some cancer patients impair TCR–

pHLA-A2 catch bonds, which may explain the suppressed anti-tumor T cell immunity[8]. More

interestingly, our models pave the way for engineering of TCR function for tumor immunotherapy

by modulating the TCR catch and slip bonds through alteration of its structures. For example,

mutations that weaken the Vα-Cα joint are predicted to strengthen TCR catch bonds and enhance

T cell effector function, which suggests a strategy that may be more advantageous compared to

mutations at the pMHC docking interface because mutations at the Vα-Cα joint would alter

function for any TCR specificity so it may be effective to many tumor antigens. By comparison,

mutations at the TCR binding interface may be applicable to a specific pMHC only, and may be

riskier in terms of cross-reactivity to self-pMHCs. Thus, rational design guided by catch bond

models may provide new TCR engineering strategies that warrant future studies.

Methods

The model developments, characterization, and validation are described in the main text with more details in the Supplementary Model Derivations. Model fitting to experimental data was done by nonlinear curve-fitting in the least-squares sense using the Levenberg-Marquardt algorithm, as detailed in Supplementary Model Derivation, A.4. Model applications, curve-fitting strategies, and biological relevance. Statistical and clustering analyses were done using Bayesian statistics and Lloyd's algorithm, respectively. All analyses were done using MATLAB. All experimental data of force *vs* bond lifetime were measured at room temperature as reported in Refs.[4-8,13,15,26-28] and the TCR–pMHC constructs used were described in the footnotes of Supplementary Tables 3, 4 and 6.

Data availability

Source data are provided with this paper through main text and Supplementary Information. Previously published bond lifetime data[4-8,13,15,26-28] re-analyzed for model fitting are summarized and deposited in Github (https://github.com/Chengzhulab/NCOMMS-22-20167).

Code availability

All code for three models used in this study are summarized and deposited in Github (https://github.com/Chengzhulab/NCOMMS-22-20167).

References

1   Feng, Y., Reinherz, E. L. & Lang, M. J. alphabeta T Cell Receptor Mechanosensing Forces out Serial Engagement. *Trends Immunol* **39**, 596-609, (2018).

2   Zhu, C., Chen, W., Lou, J., Rittase, W. & Li, K. Mechanosensing through immunoreceptors. *Nature Immunology* **20**, 1269-1278, (2019).

3   Liu, B., Kolawole, E. M. & Evavold, B. D. Mechanobiology of T Cell Activation: To Catch a Bond. *Annu Rev Cell Dev Biol* **37**, 65-87, (2021).

4   Liu, B., Chen, W., Evavold, B. D. & Zhu, C. Accumulation of dynamic catch bonds between TCR and agonist peptide-MHC triggers T cell signaling. *Cell* **157**, 357-368, (2014).

5 Hong, J., Persaud, S. P., Horvath, S., Allen, P. M., Evavold, B. D. & Zhu, C. Force-regulated
In situ TCR-peptide-bound MHC class II kinetics determine functions of CD4+ T cells. *Journal*
of immunology **195**, 3557-3564, (2015).
6 Das, D. K., Feng, Y., Mallis, R. J., Li, X., Keskin, D. B., Hussey, R. E., Brady, S. K., Wang, J.
H., Wagner, G., Reinherz, E. L. & Lang, M. J. Force-dependent transition in the T-cell receptor
beta-subunit allosterically regulates peptide discrimination and pMHC bond lifetime.
Proceedings of the National Academy of Sciences of the United States of America **112**, 1517-
1522, (2015).
7 Sibener, L. V., Fernandes, R. A., Kolawole, E. M., Carbone, C. B., Liu, F., McAffee, D.,
Birnbaum, M. E., Yang, X., Su, L. F., Yu, W., Dong, S., Gee, M. H., Jude, K. M., Davis, M.
M., Groves, J. T., Goddard, W. A., 3rd, Heath, J. R., Evavold, B. D., Vale, R. D. & Garcia, K.
C. Isolation of a Structural Mechanism for Uncoupling T Cell Receptor Signaling from Peptide-
MHC Binding. *Cell* **174**, 672-687 e627, (2018).
8 Wu, P., Zhang, T., Liu, B., Fei, P., Cui, L., Qin, R., Zhu, H., Yao, D., Martinez, R. J., Hu, W.,
An, C., Zhang, Y., Liu, J., Shi, J., Fan, J., Yin, W., Sun, J., Zhou, C., Zeng, X., Xu, C., Wang,
J., Evavold, B. D., Zhu, C., Chen, W. & Lou, J. Mechano-regulation of Peptide-MHC Class I
Conformations Determines TCR Antigen Recognition. *Mol Cell* **73**, 1015-1027 e1017, (2019).
9 Zhao, X., Kolawole, E. M., Chan, W., Feng, Y., Yang, X., Gee, M. H., Jude, K. M., Sibener, L.
V., Fordyce, P. M., Germain, R. N., Evavold, B. D. & Garcia, K. C. Tuning T cell receptor
sensitivity through catch bond engineering. *Science* **376**, eabl5282, (2022).
10 Courtney, A. H., Lo, W. L. & Weiss, A. TCR Signaling: Mechanisms of Initiation and
Propagation. *Trends Biochem Sci* **43**, 108-123, (2018).
11 Feng, Y., Brazin, K. N., Kobayashi, E., Mallis, R. J., Reinherz, E. L. & Lang, M. J.
Mechanosensing drives acuity of alphabeta T-cell recognition. *Proceedings of the National*
Academy of Sciences of the United States of America **114**, E8204-E8213, (2017).
12 McKeithan, T. W. Kinetic proofreading in T-cell receptor signal transduction. *Proceedings of*
the National Academy of Sciences of the United States of America **92**, 5042-5046, (1995).
13 Hong, J., Ge, C., Jothikumar, P., Yuan, Z., Liu, B., Bai, K., Li, K., Rittase, W., Shinzawa, M.,
Zhang, Y., Palin, A., Love, P., Yu, X., Salaita, K., Evavold, B. D., Singer, A. & Zhu, C. A TCR
mechanotransduction signaling loop induces negative selection in the thymus. *Nat Immunol* **19**,
1379-1390, (2018).
14 Liu, Y., Blanchfield, L., Ma, V. P., Andargachew, R., Galior, K., Liu, Z., Evavold, B. & Salaita,
K. DNA-based nanoparticle tension sensors reveal that T-cell receptors transmit defined pN
forces to their antigens for enhanced fidelity. *Proceedings of the National Academy of Sciences*
of the United States of America **113**, 5610-5615, (2016).
15 Zareie, P., Szeto, C., Farenc, C., Gunasinghe, S. D., Kolawole, E. M., Nguyen, A., Blyth, C.,
Sng, X. Y. X., Li, J., Jones, C. M., Fulcher, A. J., Jacobs, J. R., Wei, Q., Wojciech, L., Petersen,
J., Gascoigne, N. R. J., Evavold, B. D., Gaus, K., Gras, S., Rossjohn, J. & La Gruta, N. L.
Canonical T cell receptor docking on peptide-MHC is essential for T cell signaling. *Science*
372, (2021).
16 Dembo, M., Torney, D. C., Saxman, K. & Hammer, D. The reaction-limited kinetics of
membrane-to-surface adhesion and detachment. *Proc. R. Soc. Lond. B Biol. Sci.* **234**, 55-83,
(1988).
17 Marshall, B. T., Long, M., Piper, J. W., Yago, T., McEver, R. P. & Zhu, C. Direct observation
of catch bonds involving cell-adhesion molecules. *Nature* **423**, 190-193, (2003).
18 Grandbois, M., Beyer, M., Rief, M., Clausen-Schaumann, H. & Gaub, H. E. How strong is a
covalent bond? *Science* **283**, 1727-1730, (1999).
19 Bell, G. Models for the specific adhesion of cells to cells. *Science* **200**, 618-627, (1978).
20 Liu, B., Chen, W. & Zhu, C. Molecular force spectroscopy on cells. *Annu. Rev. Phys. Chem.*
66, 427-451, (2014).

Guo, S., Efremov, A. K. & Yan, J. Understanding the catch-bond kinetics of biomolecules on a one-dimensional energy landscape. *Communications Chemistry* **2**, 30, (2019).

Adams, Jarrett J., Narayanan, S., Liu, B., Birnbaum, Michael E., Kruse, Andrew C., Bowerman, Natalie A., Chen, W., Levin, Aron M., Connolly, Janet M., Zhu, C., Kranz, David M. & Garcia, K. C. T cell receptor signaling is limited by docking geometry to peptide-major histocompatibility complex. *Immunity* **35**, 681-693, (2011).

Rossjohn, J., Gras, S., Miles, J. J., Turner, S. J., Godfrey, D. I. & McCluskey, J. T cell antigen receptor recognition of antigen-presenting molecules. *Annual review of immunology* **33**, 169-200, (2015).

Banik, D., Hamidinia, M., Brzostek, J., Wu, L., Stephens, H. M., MacAry, P. A., Reinherz, E. L., Gascoigne, N. R. J. & Lang, M. J. Single Molecule Force Spectroscopy Reveals Distinctions in Key Biophysical Parameters of alphabeta T-Cell Receptors Compared with Chimeric Antigen Receptors Directed at the Same Ligand. *J Phys Chem Lett* **12**, 7566-7573, (2021).

Hwang, W., Mallis, R. J., Lang, M. J. & Reinherz, E. L. The alphabetaTCR mechanosensor exploits dynamic ectodomain allostery to optimize its ligand recognition site. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 21336-21345, (2020).

Liu, B., Chen, W., Natarajan, K., Li, Z., Margulies, D. H. & Zhu, C. The cellular environment regulates in situ kinetics of T-cell receptor interaction with peptide major histocompatibility complex. *European journal of immunology* **45**, 2099-2110, (2015).

Rushdi, M. N., Pan, V., Li, K., Travaglino, S., Choi, H.-K., Hong, J., Griffitts, F., Agnihotri, P., Mariuzza, R. A., Ke, Y. & Zhu, C. Cooperative binding of TCR and CD4 to pMHC enhances TCR sensitivity. *Nature Communications*, in press.

Yuan, Z., Cong, P., Natarajan, Ge, C., Travaglino, S., Beesam, S., Grazette, D., Krogsgaard, M. & Zhu, C. Cooperative ectodomain interaction among TCRαβ, CD3γε, and CD3δε enhances TCR mechanotransduction. *bioRxiv*, https://www.biorxiv.org/content/10.1101/2022.1104.1114.488403v488401, (2022).

Das, D. K., Mallis, R. J., Duke-Cohan, J. S., Hussey, R. E., Tetteh, P. W., Hilton, M., Wagner, G., Lang, M. J. & Reinherz, E. L. Pre-T Cell Receptors (Pre-TCRs) Leverage Vbeta Complementarity Determining Regions (CDRs) and Hydrophobic Patch in Mechanosensing Thymic Self-ligands. *The Journal of biological chemistry* **291**, 25292-25305, (2016).

Kramers, H. A. Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica* **7**, 284–304, (1940).

Dudko, O. K., Hummer, G. & Szabo, A. Intrinsic rates and activation free energies from single-molecule pulling experiments. *Physical review letters* **96**, 108101, (2006).

Smith, S. B., Finzi, L. & Bustamante, C. Direct mechanical measurements of the elasticity of single DNA molecules by using magnetic beads. *Science* **258**, 1122-1126, (1992).

Bouchiat, C., Wang, M. D., Allemand, J., Strick, T., Block, S. M. & Croquette, V. Estimating the persistence length of a worm-like chain molecule from force-extension measurements. *Biophysical journal* **76**, 409-413, (1999).

Huang, J., Zarnitsyna, V. I., Liu, B., Edwards, L. J., Jiang, N., Evavold, B. D. & Zhu, C. The kinetics of two-dimensional TCR and pMHC interactions determine T-cell responsiveness. *Nature* **464**, 932-936, (2010).

Xiang, S., Nie, F. & Zhang, C. Learning a Mahalanobis distance metric for data clustering and classification. *Pattern Recognition* **41**, 3600-3612, (2008).

Pereverzev, Y. V., Prezhdo, O. V., Forero, M., Sokurenko, E. V. & Thomas, W. E. The two-pathway model for the catch-slip transition in biological adhesion. *Biophysical journal* **89**, 1446-1454, (2005).

Kersh, G. J. & Allen, P. M. Structural basis for T cell recognition of altered peptide ligands: a single T cell receptor can productively recognize a large continuum of related ligands. *J Exp Med* **184**, 1259-1268, (1996).

Natarajan, A., Nadarajah, V., Felsovalyi, K., Wang, W., Jeyachandran, V. R., Wasson, R. A., Cardozo, T., Bracken, C. & Krogsgaard, M. Structural Model of the Extracellular Assembly of the TCR-CD3 Complex. *Cell Rep* **14**, 2833-2845, (2016).

Lapidus, L. J., Steinbach, P. J., Eaton, W. A., Szabo, A. & Hofrichter, J. Effects of Chain Stiffness on the Dynamics of Loop Formation in Polypeptides. Appendix: Testing a 1-Dimensional Diffusion Model for Peptide Dynamics. *J. Phys. Chem. B* **106**, 11628-11640, (2002).

Gebhardt, J. C., Bornschlogl, T. & Rief, M. Full distance-resolved folding energy landscape of one single protein molecule. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 2013-2018, (2010).

Gohring, J., Kellner, F., Schrangl, L., Platzer, R., Klotzsch, E., Stockinger, H., Huppa, J. B. & Schutz, G. J. Temporal analysis of T-cell receptor-imposed forces via quantitative single molecule FRET measurements. *Nature communications* **12**, 2502, (2021).

Ma, R., Kellner, A. V., Ma, V. P., Su, H., Deal, B. R., Brockman, J. M. & Salaita, K. DNA probes that store mechanical information reveal transient piconewton forces applied by T cells. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 16949-16954, (2019).

Ma, V. P., Liu, Y., Blanchfield, L., Su, H., Evavold, B. D. & Salaita, K. Ratiometric Tension Probes for Mapping Receptor Forces and Clustering at Intermembrane Junctions. *Nano letters* **16**, 4552-4559, (2016).

Wang, M. S., Hu, Y., Sanchez, E. E., Xie, X., Roy, N. H., de Jesus, M., Winer, B. Y., Zale, E. A., Jin, W., Sachar, C., Lee, J. H., Hong, Y., Kim, M., Kam, L. C., Salaita, K. & Huse, M. Mechanically active integrins target lytic secretion at the immune synapse to facilitate cellular cytotoxicity. *Nature communications* **13**, 3222, (2022).

Limozin, L., Bridge, M., Bongrand, P., Dushek, O., van der Merwe, P. A. & Robert, P. TCR-pMHC kinetics under force in a cell-free system show no intrinsic catch bond, but a minimal encounter duration before binding. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 16943-16948, (2019).

Barsegov, V. & Thirumalai, D. Dynamics of unbinding of cell adhesion molecules: transition from catch to slip bonds. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 1835-1839, (2005).

Evans, E., Leung, A., Heinrich, V. & Zhu, C. Mechanical switching and coupling between two dissociation pathways in a P-selectin adhesion bond. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 11281-11286, (2004).

Lou, J. & Zhu, C. A structure-based sliding-rebinding mechanism for catch bonds. *Biophys. J.* **92**, 1471-1485, (2007).

Chen, W., Lou, J. & Zhu, C. Forcing switch from short- to intermediate- and long-lived states of the alphaA domain generates LFA-1/ICAM-1 catch bonds. *J. Biol. Chem.* **285**, 35967-35978, (2010).

Kong, F., Garcia, A. J., Mould, A. P., Humphries, M. J. & Zhu, C. Demonstration of catch bonds between an integrin and its ligand. *J Cell Biol* **185**, 1275-1284, (2009).

Ju, L., Dong, J. F., Cruz, M. A. & Zhu, C. The N-terminal flanking region of the A1 domain regulates the force-dependent binding of von Willebrand factor to platelet glycoprotein Ibalpha. *The Journal of biological chemistry* **288**, 32289-32301, (2013).

Buckley, C. D., Tan, J., Anderson, K. L., Hanein, D., Volkmann, N., Weis, W. I., Nelson, W. J. & Dunn, A. R. Cell adhesion. The minimal cadherin-catenin complex binds to actin filaments under force. *Science* **346**, 1254211, (2014).

Huang, D. L., Bax, N. A., Buckley, C. D., Weis, W. I. & Dunn, A. R. Vinculin forms a directionally asymmetric catch bond with F-actin. *Science* **357**, 703-706, (2017).

Owen, L. M., Bax, N. A., Weis, W. I. & Dunn, A. R. The C-terminal actin-binding domain of talin forms an asymmetric catch bond with F-actin. *Proceedings of the National Academy of Sciences of the United States of America* **119**, e2109329119, (2022).

Thomas, W. Catch bonds in adhesion. *Annu Rev Biomed Eng* **10**, 39-57, (2008).

Mariuzza, R. A., Agnihotri, P. & Orban, J. The structural basis of T-cell receptor (TCR) activation: An enduring enigma. *The Journal of biological chemistry* **295**, 914-925, (2020).

Mallis, R. J., Bai, K., Arthanari, H., Hussey, R. E., Handley, M., Li, Z., Chingozha, L., Duke-Cohan, J. S., Lu, H., Wang, J. H., Zhu, C., Wagner, G. & Reinherz, E. L. Pre-TCR ligand binding impacts thymocyte development before alphabetaTCR expression. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 8373-8378, (2015).

Sarangapani, K. K., Yago, T., Klopocki, A. G., Lawrence, M. B., Fieger, C. B., Rosen, S. D., McEver, R. P. & Zhu, C. Low force decelerates L-selectin dissociation from P-selectin glycoprotein ligand-1 and endoglycan. *J. Biol. Chem.* **279**, 2291-2298, (2004).

Wayman, A. M., Chen, W., McEver, R. P. & Zhu, C. Triphasic force dependence of E-selectin/ligand dissociation governs cell rolling under flow. *Biophys. J.* **99**, 1166-1174, (2010).

Choi, Y. I., Duke-Cohan, J. S., Chen, W., Liu, B., Rossy, J., Tabarin, T., Ju, L., Gui, J., Gaus, K., Zhu, C. & Reinherz, E. L. Dynamic control of beta1 integrin adhesion by the plexinD1-sema3E axis. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 379-384, (2014).

Rosetti, F., Chen, Y., Sen, M., Thayer, E., Azcutia, V., Herter, J. M., Luscinskas, F. W., Cullere, X., Zhu, C. & Mayadas, T. N. A Lupus-Associated Mac-1 Variant Has Defects in Integrin Allostery and Interaction with Ligands under Force. *Cell Rep* **10**, 1655-1664, (2015).

Chen, Y., Lee, H., Tong, H., Schwartz, M. & Zhu, C. Force regulated conformational change of integrin alphaVbeta3. *Matrix Biol* **60-61**, 70-85, (2017).

Chen, Y., Ju, L. A., Zhou, F., Liao, J., Xue, L., Su, Q. P., Jin, D., Yuan, Y., Lu, H., Jackson, S. P. & Zhu, C. An integrin alphaIIbbeta3 intermediate affinity state mediates biomechanical platelet aggregation. *Nat Mater* **18**, 760-769, (2019).

Manibog, K., Li, H., Rakshit, S. & Sivasankar, S. Resolving the molecular mechanism of cadherin catch bond formation. *Nature communications* **5**, 3941, (2014).

Nishi, H., Furuhashi, K., Cullere, X., Saggu, G., Miller, M. J., Chen, Y., Rosetti, F., Hamilton, S. L., Yang, L., Pittman, S. P., Liao, J., Herter, J. M., Berry, J. C., DeAngelo, D. J., Zhu, C., Tsokos, G. C. & Mayadas, T. N. Neutrophil FcgammaRIIA promotes IgG-mediated glomerular neutrophil capture via Abl/Src kinases. *The Journal of clinical investigation* **127**, 3810-3826, (2017).

Luca, V. C., Choul Kim, B., Ge, C., Kakuda, S., Roein-Peikar, M., Haltiwanger, R. S., Zhu, C., Ha, T. & Garcia, K. C. Notch-Jagged complex structure implicates mechanical force in ligand discrimination. *Science* **355**, 1320-1324, (2017).

Yago, T., Lou, J., Wu, T., Yang, J., Miner, J. J., Coburn, L., Lopez, J. A., Cruz, M. A., Dong, J. F., McIntire, L. V., McEver, R. P. & Zhu, C. Platelet glycoprotein Ibalpha forms catch bonds with human WT vWF but not with type 2B von Willebrand disease vWF. *J. Clin. Invest.* **118**, 3195-3207, (2008).

Yakovenko, O., Sharma, S., Forero, M., Tchesnokova, V., Aprikian, P., Kidd, B., Mach, A., Vogel, V., Sokurenko, E. & Thomas, W. E. FimH forms catch bonds that are enhanced by mechanical force due to allosteric regulation. *The Journal of biological chemistry* **283**, 11596-11605, (2008).

Guo, B. & Guilford, W. H. Mechanics of actomyosin bonds in different nucleotide states are tuned to muscle contraction. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 9844-9849, (2006).

Lee, C. Y., Lou, J. Z., Wen, K. K., McKane, M., Eskin, S. G., Ono, S., Chien, S., Rubenstein, P. A., Zhu, C. & McIntire, L. V. Actin depolymerization under force is governed by lysine

113:glutamic acid 195-mediated catch-slip bonds. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 5022-
5027, (2013).
71 Lee, C. Y., Lou, J., Wen, K. K., McKane, M., Eskin, S. G., Rubenstein, P. A., Chien, S., Ono,
S., Zhu, C. & McIntire, L. V. Regulation of actin catch-slip bonds with a RhoA-formin module.
Sci Rep **6**, 35058, (2016).
72 Akiyoshi, B., Sarangapani, K. K., Powers, A. F., Nelson, C. R., Reichow, S. L., Arellano-
Santoyo, H., Gonen, T., Ranish, J. A., Asbury, C. L. & Biggins, S. Tension directly stabilizes
reconstituted kinetochore-microtubule attachments. *Nature* **468**, 576-579, (2010).
842

Acknowledgments

We thank Baoyu Liu, Jinsung Hong, Peng Wu, Tongtong Zhang, Aswin Natarajan, and Michelle

Krogsgaard for sharing their published BFP experimental data for re-analysis and model fitting.

This work was supported by a Postdoctoral Fellowship from the National Research Foundation of

South Korea (2021R1A6A3A03038382 to H.-K.C.), by a National Natural Science Foundation of

China grant (31971237 to W.C.), and by a National Natural Science Foundation of China grant

(32090044 to J.L.). This work was also supported by National Institutes of Health grants

(U01CA214354, U01CA250040, R01CA243486, and R01GM061126 to C.Z.).

Author contributions

H.-K.C. and C.Z. conceived the project and developed the models. H.-K.C., K.L., P.C., M.N.R.,

and C.G. provided data prior to their publication. H.-K.C. performed model analysis and data

fitting. H.-K.C., K.L. and C.Z. made key observations and conclusions from analyses. H.-K.C.,

K.L., and C.Z. prepared the manuscript with input from other authors.

Competing interests

No authors have competing interests to declare.

Figures

Figure 1 | Comparison between the two-pathway model and the TCR–pMHC-I model for catch bond. **a** *Upper*: The 2D energy landscape of the two-pathway model where the bond is trapped in the bound-state energy well by two energy barriers that resist dissociation along two pathways, p1 and p2, with two distinct transition states, p1‡ and p2‡. The application of force projects the energy landscape towards a dissociation pathway along force (δ_l). *Lower*: Force also tilts the energy-landscape, raising the energy barrier of the first pathway and lowering the energy barrier of the second pathway on the energy landscape projection (*red* and *blue*) relative to their positions in the absence of force (*black*). **b** *Lower*: The proposed 1D energy landscape of TCR–pMHC-I model with a single transition state δ^*. Below an optimal value (F_{opt}), force raises the energy barrier (*red*) relative to the zero-force conformation (*black*) by contraction of flexible regions due to entropic fluctuation. Above F_{opt}, force lowers the energy barrier (*blue*) by stretching the molecular complex. Together, these two mechanisms give rise to a catch-slip bond. *Upper*: Schematics of the TCR–pMHC-I structure (*left*) and its conformational changes that correspond to low (*middle*) and high (*right*) forces.

Figure 2 | Structure, mechanics, and characteristics of the TCR–pMHC-I catch bond model.

880 **a** Force-induced conformational changes of a TCR–pMHC-I complex as it traverses from the bound

881 state (*left*) to the transition state (*right*). The diagrams of the 2C TCR α (*yellow*) β (*green*) subunits

and the DEVA peptide (*red*) bound to the H2-K^b (various domains indicated) are based on snapshots

from SMD simulations performed on the complex structure (2CKB) at the initial time (bound state)

and a later time (transition state)[8]. The force-transmission path is shown as red lines connecting the

force-acting nodes. **b** Various contributions to the total extension projected on the force axis:

rotation of the α_3- β_{2m} domains about the MHC C-terminus upon dissociation of the β_{2m}–$\alpha_1\alpha_2$

interdomain bond ($z_{\alpha3}$), relative rotation between α_3 and $\alpha_1\alpha_2$ about their stretched interdomain

hinge ($z_{p,MHC}$), tilting of the MHC $\alpha_1\alpha_2$ complexed with TCR V$\alpha\beta$ ($z_{B,I}$), rotation about and

extension of the Vα–Cα interdomain hinge ($z_{p,TCR}$), and extension of the C$\alpha\beta$ and rotation about

their C-termini ($z_{C\alpha\beta}$). Two α_3-β_{2m} structures are shown: before (light colors) and after (dark colors)

β_{2m} dissociation from $\alpha_1\alpha_2$., with two parameters describing their contributions to the total

extension: $d_{\alpha3}$ = the distance between the $\alpha3$ C- and N-termini excluding the $\alpha_1\alpha_2$–$\alpha3$ hinge and θ

= the angle between the normal direction of the TCR–pMHC bonding interface at the bound state

(cf. (**a**), *left*) and the tilted direction at the transition state (cf. (**a**), *right*). **c** Extension change *vs*

force curves (*lower*) for the color-matched n^* and θ values (*upper*). The left panel (set 1 in *upper*

table) shows the effect of changing θ with ($n^* = 7$) and without ($n^* = 0$) partial unfolding. The right

panel (set 2) shows the effect of changing n^* with ($\theta = 30°$) and without ($\theta = 0°$) tilting. **d** n^*-θ

phase diagram showing three parameter domains: slip-only, catch-slip, and catch-only respectively

colored by red-purple, purple-blue, and blue-black. Upper insets indicate corresponding energy

change $\delta_l\gamma$ *vs* force curves for each bond type. **e** Theoretical normalized bond lifetime *vs* force

curves for indicated parameters. The upper and lower panels show the respective effects of changing

θ and δ_0^* from the set 1 parameters defined in (**c**). **f** Energy landscapes expressed as families ΔG *vs*

δ_l curves for a range of forces for slip-only (*upper*) and catch-slip (*lower*) bonds. The bound state

is located at the origin $\Delta G = 0$ and $\delta_l = 0$. The transition state has an energy of ΔG^* located at δ^*

when $F > 0$ and ΔG_0^* located at δ_0^* when $F = 0$. **g** Plots of transition state location δ^* (*upper*) and

906   height of energy barrier ΔG^* (*lower*) *vs* force F for changing θ while keeping δ_0^* constant (*left*) or

907   changing δ_0^* while keeping θ constant (*right*) for the indicated values from parameter table in (**c**).

Figure 3 | TCR bond type characterization and correlation with pMHC-I biological activity.

a-b Fitting of theoretical $1/k(F)$ curves to experimental bond lifetime *vs* force data (*points*, mean ± sem, re-analyzed from Refs. [4,8]) of OT1 (**a**) or 2C (**b**) TCR expressed on CD8$^+$ naïve T cells interacting with indicated p:H2-K^{b}α3A2 ((**a**) and (**b**) *left*) or on CD8$^-$ 2C hybridomas interacting with indicated p:H2-K^b ((**b**) *right*). Several metrics are defined to characterize the force-lifetime curve as indicated in the left panel of (**a**): F_{opt} is the "optimal force" where lifetime peaks (t_{peak}), Δt is the lifetime increase from the zero-force value t_0 to t_{peak}, and ΔF is the force increase from F_{opt} to the force where lifetime returns to t_0. **c-d** Two dimensional metrics, t_{peak} and Δt (**c**), and

two dimensionless metrics, $L = \Delta t / t_{\mathrm{peak}}$ and $I = L/(1+B)$ where $B = \Delta F / F_{\mathrm{opt}}$ (**d**), are plotted

vs the logarithm of the reciprocal peptide concentration required to stimulate half-maximal T cell

proliferation ($1/EC_{50}$) and fitted by a linear function. **e** Intensity of catch bond, I, calculated using

the fitted force-lifetime curves of OT1, 2C TCR on primary T cells, 2C TCR on hybridomas,

purified N15 TCRαβ, or TRBV TCRs (B13.C1/B17.C1 and B17.R1/B17.R2) expressed on

hybridomas interacting with their corresponding pMHCs are plotted according to the ranked-order

of peptide potencies (*bottom*). **f** Best-fit model parameters θ (the tilted angle of the bonding

interface, *upper*), n^* (the number of unfolded amino acids, *middle*), and δ_0^* (the width of zero-force

free-energy well, *lower*) are plotted according to the ranked order of the peptide potencies (*bottom*).

All error bars in 1st and 2nd rows are fitting error (Supplementary Table 3).

Figure 4 | Properties and biological relevance of class I model parameters. **a-b** The width of zero-force energy well δ_0^* (**a**) and the intensity of catch bond I (**b**) calculated from WT or mutant 2C TCRs (*left*) and WT 1G4 TCR (*right*) interacting with their corresponding WT or MT pMHCs. The MT 2C TCRs and H2-K^{b}s were designed to destabilize the TCR–pMHC interaction. The MT p:HLA-A2s were designed to either destabilize the TCR–pMHC interaction (R75A) or stabilize the MHC intramolecular interaction (A236T and F8V). **c** Data from **Fig. 3e** and **f** (3rd row) are re-graphed as δ_0^* *vs* I plot to show their correlation (*blue*). Additional δ_0^* *vs* I data from MT TCRs and/or MT pMHCs without functional data also show strong correlation (*green*). Different TCR systems are indicated by different symbols. The two datasets were separately fitted by two straight

lines with the goodness-of-fit indicated by R^2. **d** Tilting angle of the bonding interface (θ) *vs* normalized net gain of hydrogen bonds at the interface between 2C TCR and the indicated pMHCs is plotted (*points*) and fitted (*line*) with 95% confident interval shown as shaded area (errors in x- and y-axis represent propagation uncertainty from Supplementary Fig. 3 and fitting error, respectively). **e** Clustering analysis reveals three clusters in the δ_0^*-θ phase diagram: slip-only (SO, *black*), weak catch-slip (WC, *blue*), and strong catch-slip (SC, *red*) bonds. Closed-symbols represent bonds of the WT TCR with WT agonist peptides. **f** Tilting angle (θ) *vs* end-to-end distance of dissociated α3 domain ($d_{\alpha3}$). The three types of bonds, SO, WC, and SC, are also clustered in this phase diagram, which are separated by the dotted lines that predicted from the pulling constraints of the model. The two pairs of TRBV TCRs are indicated in e and f by green dots. **g** The average molecular extensions at zero force ($\langle\delta_0^*\rangle$, *left ordinate*) and the average rotation angle ($\langle\theta\rangle$, *right ordinate*) (mean ± sem) are plotted *vs* the total number of unfolded amino acids (n^*, *abscissa*) to show three clusters. Each bond type is indicated by a dotted line (n = 14, 9, and 16; number of data related to SO, WC, and SC, respectively, individual data of each cluster are shown in (**e-f**)). **h** Catch bond intensity *vs* Mahalanobis distance plot, again showing three clusters. Principal component analysis was used to find principal axes. Mahalanobis distances for each cluster were calculated using common principal axes from total dataset (number of data is same as (**g**)). **i** Structural models illustrating the conformations of three bond types based on their model parameters based on the previous SMD simulation of the 2C TCR system[8]. Two structural parameters (θ, *red*; $d_{\alpha3}$, *blue*) are indicated to show the differences between bond types. Unless otherwise described, all errors shown in (**a-f**) are fitting errors (Supplementary Table 3).

Figure 5 | The TCR–pMHC-II model. a Force-induced conformational changes of a TCR–pMHC-II complex as it traverses from the bound state (*left*) to the transition state (*right*). The diagrams are based on the published co-crystal structure (2IAM) of the E8 TCR α (*yellow*) β (*green*) subunits and the TPI peptide (*red*) bound to the HLA-DR1 α (*blue*) β (*pink*) subunits with various domains indicated. The force-transmission paths are shown as red lines connecting the force-acting nodes. **b** Various contributions to the total extension projected on the force axis: stretching of the TCR Cα and Cβ domains ($z_{C\alpha\beta}$), asymmetric partial unfolding of the TCR Vα-Cα and Vβ-Cβ

interdomain hinges ($z_{p,TCR}$), asymmetric partial unfolding of the MHC $\alpha1$-$\alpha2$ and $\beta1$-$\beta2$ interdomain hinges ($z_{p,MHC}$), and rotation between the $\alpha1$-$\beta1$ and $\alpha2$-$\beta2$ domain hinges and tilting of the bonding interface between the MHC $\alpha1$-$\beta1$ and the TCR $V\alpha$-$V\beta$ by an angle θ ($z_{B,I}$). **c** Theoretical normalized bond lifetime *vs* force curves. The effects of changing θ and δ_0^* are shown in the upper and lower panels, respectively, for the indicated parameter values. **d** Fitting of predicted $1/k(F)$ curves (*dashed lines* by two-pathway model and *solid lines* by TCR–pMHC-ΙΙ model) to experimental bond lifetime *vs* force data (*points*, mean ± sem) of 3.L2 TCR on CD4⁻CD8⁺ naive T cells interacting with indicated p:I-E^k's [5] (*upper*) or WT and indicated mutant 2B4 TCRs on hybridomas interacting with K5:I-E^k [28]. **e** Dimensional metrics, Δt (*left*), scaled relative length of bond lifetime L (*middle*), and intensity of catch bond I (*right*) *vs* reciprocal % change (relative to WT) of effector function, i.e., the peptide dose required for 3.L2 T cells to generate 40% B cell apoptosis $(1/EC_{40})$[37] (*red*) or the area under the dose response curve (AUC) of the 2B4 hybridoma IL-2 production[38] (*blue*) plots. **f** Best-fit model parameters δ_0^* (*left*), θ (*middle*), and n^* (*right*) are plotted *vs* reciprocal relative % change of effector function. **g** The three model parameters in f for both the 3.L2 and 2B4 TCR systems are plotted *vs* the catch-bond intensity I and fitted by a straight line. We also added to each panel an additional point obtained from data and model fit of E8 TCR–TPI:HLA-DR1 interactions[27]. All error bars in **f-g** are fitting error (Supplementary Table 6).

Figure 6 | Cross-examination of class I and II models against class I and II data. a-b δ_0^* *vs I*

plots obtained using class I (*black*) or class II (*red*) model to fit force-lifetime data of TCR

interacting with pMHC-I (**a**) or pMHC-II (**b**) molecules. In each panel, two sets of parameter values

were returned from fitting depending on whether class I (*black*) or class II (*red*) model was used

because they are based on different structures and conformational changes of the TCR–pMHC

complexes. **c** The slopes (*gray*, the level of correlation between δ_0^* and *I*) and goodness-of-fit (R^2)

(*blue*, the degree of appropriateness of the model for the data) of the linear fit in (**a**) and (**b**) are

shown in the matched (1st and 4th groups) and mismatched (2nd and 3rd groups) cases. **d-e** δ_0^* *vs* n^*

plots obtained using class I (*black*) or class II (*red*) model to fit force-lifetime data of TCR

interacting with pMHC-I (**d**) or pMHC-II (**e**) molecules. **f** The slopes and goodness-of-fit of the

linear fit in (**d**) and (**e**) are shown in the matched (1st and 4th groups) and mismatched (2nd and 3rd

groups) cases. The slopes indicate the average unfolding extension per amino acid (nm/a.a.) from

each model, which are compared to the maximum average contour length per amino acid of ~0.4

nm/a.a. (biophysical limit, black dashed line with considerable deviation)[21,39,40]. All error bars are

fitting error (Supplementary Table 6).

Supplementary Information

Catch bond models may explain how force amplifies TCR signaling and antigen discrimination

Hyun-Kyu Choi et al.

*Corresponding author. Email: cheng.zhu@bme.gatech.edu

This PDF file includes:

Supplementary Model Derivations
Supplementary Figures. 1 to 11
Supplementary Tables 1 to 6
References (73 to 79)

Supplementary Methods

Supplementary Model Derivations

A. Kinetic model for TCR–pMHC-I bonds
    A.1. Derivation of the force-induced energy change $\delta_l \gamma(F)$
    A.2. Simplifying assumptions and reducing parameters
    A.3. Defining the dissociation coordinate
    A.4. Model applications, curve-fitting strategies, and biological relevance
    A.5. Class I model constraints
B. Kinetic model for TCR–pMHC-II bonds
    B.1. Development, validation, and characterization
    B.2. Class II model constraints
C. A general biophysical limit of model parameters

Supplementary Figures

Supplementary Tables

Supplementary References

Supplementary Model Derivations

A. Kinetic model for TCR–pMHC-I bonds

A.1. Derivation of the force-induced energy change $\delta_1\gamma(F)$

The purpose of this subsection is to provide detailed derivation of the force-induced change in the energy landscape, or the work done by force as the TCR–pMHC complex is stretched at the two ends from the bound state to the transition state until dissociation (Fig. 2a, b). Mathematically, $\delta_1\gamma(F)$ is an integral over a force range $0 \leq f \leq F$, where F is the level of force under which the kinetic rate is being evaluated (Eq. 2). The integrand $\delta_z(f)$ is the projection on the force direction of the length change of the TCR–pMHC structure relative to its bound state during dissociation. Subtraction of this work from the interaction free energy tilts the energy landscape that governs the off-rate (Eq. 1, Fig. 1b). To calculate molecular stretch, we assume the TCR–pMHC complex to behave as a system of semi-rigid bodies of globular domains connected by semi-flexible polymers (Fig. 2b). As such, the total length change includes three components: 1) extension of individual globular domains, 2) various domain rotations about hinges, and 3) unfolding of secondary structures at specific regions.

For a globular domain without unfolding, the force-extension relationship is described by the three-dimensional freely-jointed chain model[1]:

$$d_i(f) = d_{i,c}\left\{\coth\left[fd_{i,c}/k_BT\right] - \frac{k_BT}{fd_{i,c}}\right\}\left(1 + \frac{f}{E_d}\right), \qquad \textbf{[S1]}$$

where d_i is end-to-end distance of the ith-domain, k_B is the Boltzmann constant, T is absolute temperature, f is force, and $E_d \sim 100$ pN is the elastic modulus of the folded globular domain[2]. The present work considers domain extension of the whole TCR–pMHC-I complex (d_N) and three of its parts: the MHC α_3 domain ($d_{\alpha3}$), bonding interface that includes the MHC $\alpha_1\alpha_2$ domains bound to the TCR V$\alpha\beta$ domains ($d_{B,I}$), and the TCR C$\alpha\beta$ domains ($d_{C\alpha\beta}$). Their contour lengths, $d_{i,c}$, have well-defined values depending on the TCR–pMHC complex in question, which are summarized in Supplementary Table 1. For example, $d_{N,c}$ = 12.7 and 10.9 nm for 2C TCR

complexed with H2-K^b (PDB codes 2CKB, 1MWA, and 1G6R) and H2-L^{d}m31 (2E7L), respectively, and 12.3 nm for 1G4 TCR complexed with HLA-A2 (2BNR and 2BNQ). 12.5 nm for NP1-B17 TCR complexed with H2-D^b. We also choose 11.6 nm as a reasonable guess value for the N15 TCR/OT1 TCR –p:H2-K^b complex (because average $d_{N,C}$ across 33 structures of TCR-pMHC class I complexes is 11.6 ± 1 nm (mean $\pm$ sd)[3]).

To calculate the work $\delta_l \gamma(F)$, we project the above domain extensions onto the direction of force, which is taken as the z direction (Fig. 2b) using two angles between the z axis and: 1) the normal direction of the bonding interface (θ) and 2) the line connecting the C- and N-termini of the MHC-I α_3 domain excluding any unfolded residues (φ):

$$z_N(f) = d_N(f) \qquad \qquad \text{[S2a]}$$

$$z_{\alpha 3}(f) = \sqrt{d_{\alpha 3}^2(f) - \left\{\left[d_N(f) - d_{C\alpha\beta}(f)\right]\sin\theta - d_{\alpha 3}(f)\sin(\varphi + \theta)\right\}^2}, \qquad \text{[S2b]}$$

$$z_{B.I}(f) = d_{B.I}(f) \equiv \left[d_N(f) - d_{C\alpha\beta}(f)\right]\cos\theta - d_{\alpha 3}(f)\cos(\varphi + \theta), \qquad \text{[S2c]}$$

$$z_{C\alpha\beta}(f) = d_{C\alpha\beta}(f). \qquad \qquad \text{[S2d]}$$

Note that θ is a model parameter as it describes the tilting of the bonding interface as part of the force-induced conformational change, whereas φ is a model constant is measured from the crystal structure (Supplementary Table 1, see A.2 below).

We assume that partial unfolding in the molecular complex may occur at connecting regions of globular domains, in particular, the $\alpha_1\alpha_2$–α_3 joint of the MHC-I and the Vα-Cα interdomain joint of the TCR (Fig. 2a). The former may be caused by dissociation of the noncovalent $\alpha_1\alpha_2$–β_{2m} interdomain bond, which shifts the mechanical load originally borne by this bond to the $\alpha_1\alpha_2$–α_3 hinge, resulting in its partial unfolding, as observed in SMD simulations[4] (Fig. 2a). Similarly, $\alpha_1\alpha_2$–β_{2m} dissociation results in tilting of the bonding interface and load shifting from the Vβ-Cβ joint to the Vα–Cα joint, leading to partial unfolding of the latter joint (Fig. 2a). The unfolded polypeptides

are flexible and can bear only tension but not moment, ensuring that their extension is along the direction of force, i.e., the z axis.

The force-extension relationship of the unfolded polypeptides can be described by an extensible worm-like chain (eWLC) model[5]:

$$\frac{f l_p}{k_B T} = \frac{1}{4(1 - z_p/(n l_c))} - \frac{1}{4} + \frac{z_p}{n l_c} - \frac{f}{K_p} + \sum_{j=1}^{j \leq 7} a_j \left(\frac{z_p}{n l_c} - \frac{f}{E_p} \right)^j, \qquad \text{[S3]}$$

where z_p is the extension of the unfolded coil under force with the subscript p indicating unstructured polypeptide, $l_c = 0.36$ nm and $l_p = 0.39$ nm are the average contour length and persistence length per unfolded amino acid, respectively[6,7,8], $E_p \sim 50$ μN is the elastic modulus of polypeptides[8], a_j are polynomial coefficients for the improved approximation, and n is the number of constituent amino acids in the unfolded polypeptide. In particular, we denote the respective numbers of amino acids in the unfolded MHC-I $\alpha_1\alpha_2$–α_3 and TCR Vα-Cα joints to be $n_{p,MHC}$ and $n_{p,TCR}$. Eq. S3 defines z_p as a function of f, which can be solved numerically to express in an explicit form: $z_p/n l_c = z_{u,p}(f)$ = the extension per unit contour length for the polypeptide under force f.

Thus, the length of the TCR–pMHC complex at the transition state is (Fig. 2b):

$$z(f) = z_{\alpha 3}(f) + n_{p,MHC} l_c z_{u,p}(f) + z_{B.I}(f) + n_{p,TCR} l_c z_{u,p}(f) + z_{C\alpha\beta}(f) \qquad \text{[S4]}$$

Since we do not have prior knowledge about either number of unfolded amino acids, we will evaluate their sum $n^* = n_{p,TCR} + n_{p,MHC}$ from curve-fitting of our model to the experimental data (see below). Since $d_N(f)$ is the length of the TCR–pMHC complex at the bound state (Fig. 2b), we have $z(f; 0) = d_N(f)$. Finally, the integrand on the right-hand side of Eq. 3 can be written as

$$\delta_z(f) = z_{\alpha 3}(f) + z_{B.I}(f) + z_{C\alpha\beta}(f) + n^* l_c z_{u,p}(f) - d_N(f). \qquad \text{[S5]}$$

A.2. Simplifying assumptions and reducing parameters

The purpose of this subsection is to provide the details omitted in the main text of how the model parameters are reduced to the smallest possible set and the underlying simplifying

assumptions. To begin, we take the φ values from PDB for the specific TCR–pMHC interactions in question (Supplementary Table 1). Due to their small range ($5° \leq \varphi \leq 23.5°$), we assume that the φ value remains constant during forced dissociation (*e.g.*, $\varphi = 12.7°$ and $5°$ for 2C TCR complexed with H2-K^b (PDB codes 2CKB, 1MWA, and 1G6R) and H2-L^{d}m31 (2E7L), respectively, and $13.3°$ for 1G4 TCR complexed with HLA-A2 (2BNR and 2BNQ). $15°$ for NP1-B17 TCR complexed with H2-D^b. We also choose $23.5°$ as a reasonable guess value for the N15 TCR/OT1 TCR –p:H2-K^b complex). Additionally, due to the semi-rigid approximation, the thickness of the constant domain of TCR α- and β-subunit ($d_{C\alpha\beta}$) is also treated as constant for each construct (Supplementary Table 1). This leaves only two structural parameters ($d_{\alpha3}$, θ) in our model to be evaluated by curve-fitting to data. Noting that the structure-based force function $\gamma(F; \delta_l)$ scales with the characteristic extension change per unit change of molecular length such that $\delta_l \gamma(F) = \int_0^F \delta_z(f) df$, the dissociation rate of TCR–pMHC-I bond can be written as:

$$k(F) = k_0 \sqrt{1 - \frac{2\int_0^F \delta_z(f)df}{3\Delta G_0^*}} \, e^{\frac{\Delta G_0^*}{k_B T}\left(1 - \left(1 - \frac{2\int_0^F \delta_z(f)df}{3\Delta G_0^*}\right)^{3/2}\right)}, \qquad \text{[S6]}$$

where $\delta_z(f)$ is given by Eq. S5.

To further reduce model parameters, we make additional assumptions as discussed in the main text. It is well-known that the fractions of free-energy change in biological interactions in liquid, such as unfolding and refolding of proteins, unbinding and rebinding of receptor–ligand bonds, and unzipping and rezipping of RNA or DNA, are small because of their limited dynamic transition time[9-12]. Such a rate limit results from the nature of biological interactions, *e.g.*, polar/non-polar interactions, hydrophobic interactions, and charged interactions, which typically yield finite range of transition kinetics. This enables to roughly estimate the free-energy barrier as $\Delta G_0^* \sim \ln(k_w/k_0)$ where $k_w \sim 10^6 \, s^{-1}$ known as the prefactor[9-12].

A.3. Defining the dissociation coordinate

The purpose of this subsection is to derive an operational way to determine the dissociation

coordinate variable δ_l used in the main text. We note that the total number of unfolded amino acids

n^* is zero at the bound state before unfolding occurs, increases monotonically during progressive

unfolding along the dissociation path, and reaches maximum at the dissociation point. Because n^*

is not known *a priori*, it must be treated as a fitting parameter similar to $d_{\alpha 3}$, θ, and δ_0^*. Since δ_l is

the contour length change along the dissociation path, we wish that δ_l approaches its upper bound

δ_0^* and depends on force implicitly through the model parameters n^*, $d_{\alpha 3}$, and θ, as given below:

$$\delta_l = n^* l_c + \Delta(d_{\alpha 3}, \theta) \rightarrow \delta_0^*, \qquad\qquad \text{[S7]}$$

where $\Delta(d_{\alpha 3}, \theta)$ is the difference of the contour lengths except for the partially unfolded regions.

Thus, Eq. S7 provides a constraint for δ_l instead of introducing another model parameter. $d_{\alpha 3}$ and

θ are determined for each n^* during the model fitting that searches for parameters to enable $\delta_l \rightarrow$

δ_0^*. Small errors, which may occur for the various contributions to Δ (red lines in Fig. 2a labeled as

force transmission lines), can be identified using a pair of $(d_{\alpha 3}, \theta)$ values and the crystal structure

for each complex. Specifically, average differences $\langle \Delta \rangle$ are - 0.9 nm for the strong catch bond (SC)

group ($d_{\alpha 3} > 3$ nm, $\theta > 20°$), 0.4 nm for the weak catch bond (WC) group (1 nm $< d_{\alpha 3} < 3$ nm,

$5° < \theta < 20°$), and 0 nm for slip-only bond (SO) group ($d_{\alpha 3} < 1$ nm, $\theta < 5°$), respectively (see Fig.

4e-i and associated text for the definitions of SC, WC, and SO groups). It has been well established

that the contour length of a single amino acid is $\sim 0.4 \pm 0.02$ nm/a.a[6,13,14], implying that the model

has a resolution of 3 amino acids. We further note that, even without conformational change, it is

possible for the slip-only bond group to have 3 unfolded amino acids due to the limited resolution

of the model. Finally, the best-fit model parameters can be determined by finding the subset of best-

fit parameters among possible n^* values that match the contour length change δ_l to the free-energy

well width at zero-force δ_0^*, *i.e.*, finding n^* such that $\delta_l(n^*, d_{\alpha 3}, \theta) \rightarrow \delta_0^*$ (see Supplementary Table

2). Under this condition, $\int_0^F \delta_z(f) df = \delta_l \gamma(F) \rightarrow \delta_0^* \gamma(F)$ and Eq. S6 becomes identical to Eq. 2.

A.4. Model applications, curve-fitting strategies, and biological relevance

The purpose of this subsection is to outline the procedures of applying our model to

experiments. Our procedures include four steps: 1) examine how the model parameters control the

model behaviors, 2) fit the model-predicted reciprocal off-rate (Eq. S6) to the experimental bond

lifetime *vs* force data, 3) construct the energy landscape and investigate its properties based on the

parameters evaluated in part 2, 4) elucidate the biological relevance of the model parameters.

To examine how the model parameters control the model behaviors, we varied one

parameter while keeping others constant. For example, to investigate the effect of varying the titling

angle θ, we kept the other parameters constants (*i.e.*, $n^* = 7$, and $\delta_0^* = 2$ nm, $k_0 = 3$ s^{-1} and $\Delta G_0^* =$

12.7 $k_\mathrm{B}T$), Also kept constant were several structural constants ($\varphi = 15°$, $d_N = 12$ nm and $d_{C\alpha\beta} =$

3.5 nm) introduced in section A.1. Of note, $d_{\alpha 3}$ varies as θ changes because of a pulling constraint

(see section B.1). With a fixed θ (*i.e.*, 30°) the effect of molecular extension at zero force was

investigated by varying extension from 0.5-3.5 nm. These were selected by the average values of

actual fitting results, which had been tested and confirmed as reasonable.

The free-energy landscape can be constructed by substituting the best-fit model parameters

into the following equations[13]:

$$\Delta G^*(F) = \Delta G_0^* \left(1 - \frac{2\delta_0^* \gamma(F)}{3\Delta G_0^*} \right)^{3/2},$$

$$\delta^*(F) = \delta_0^* \left(1 - \frac{2\delta_0^* \gamma(F)}{3\Delta G_0^*} \right)^{1/2},$$

[S8a,b]

Thus, by using model parameters, the dissociation state coordinates relative to the bound state

coordinates in the free-energy *vs* dissociation coordinate space can be defined as functions of force.

Note that this force-induced change of the barrier height should be under the condition of small

perturbation such that $|-\delta_0^* \gamma(F)| < \Delta G_0^*$. Since our fitting results show that the average free-energy

barrier height at zero force is ~12 $k_\mathrm{B}T$ ($\langle \Delta G_0^* \rangle$), the force range corresponding to a change of the

barrier height of <10 $k_\mathrm{B}T$ is reasonable for each dataset, *i.e.,* force range corresponding to energy

barrier heights in the range of 2 $k_\mathrm{B}T < \Delta G_0^* < 22$ $k_\mathrm{B}T$.

Firstly, to fit the model-predicted reciprocal off-rate (Eq. S6) to the experimental bond lifetime *vs* force data, all four parameters were changed simultaneously to search for the minimum of the chi squared error. Second, every fitting curve was then determined by the parameter set that has the closest δ_l value to δ_0^* (described as Eq. S7 in section A.3 and Supplementary Table 2). Fitting error of the best fit parameters could be calculated using standard error of mean of bond lifetime. Of the 50 datasets analyzed, only one has 4 data points and this dataset shows slip bond; as such, is governed by two fitting parameters because the other two parameters are nearly zero. All other datasets have 6-10 data points; therefore, over-fitting is not a problem.

To elucidate the biological relevance of the model parameters θ, n^*, $d_{C\alpha\beta}$, δ_0^*, and ΔG_0^*, we examine their changes with varying bond lifetime *vs* force data obtained from different TCR and pMHC interactions that induce a wide range of biological responses. Finding correspondence between a group of model parameters individually and/or collectively with the biological response would be considered as support for the biological relevance of the model, because such correspondence suggests that the model can discriminate different TCR–pMHC interactions that result in differential T cell functions.

A.5. Class I model constraints

The purpose of this subsection is to check whether the model parameters obtained from data fitting are consistent with the constraints to which the model is subjected. Our experiments applied tensile force through the two ends of the TCR–pMHC complex, such that the force direction would always align with the line connecting the C-termini of the respective TCR and MHC molecules during dissociation, giving rise to the so-called pulling constraint. To formulate this pulling constraint in our model, we note that the pulling line is maintained so that the coordinate perpendicular to the force direction is invariant. As depicted in Supplementary Fig. 4a, several angle and length variables can be related using model parameters and structural constants by:

$$\left(d_N - d_{C\alpha\beta} - d_{\alpha3}\frac{\cos(\varphi+\theta)}{\cos\theta}\right)\sin\theta = d_{\alpha3}\left(\frac{\sin\varphi}{\cos\theta} + \cos\left(2\psi + \varphi + \theta - \frac{\pi}{2}\right)\right), \qquad \text{[S9a]}$$

1224   where ψ is the angle in an isosceles triangle constructed by rotating the $\alpha3$ domain. By assuming

1225   that the $\alpha3$ domain would be aligned with force, we estimate that the angle in the last term is near

1226   90°, $i.e.$, $2\psi + \varphi + \theta - \frac{\pi}{2} \approx \frac{\pi}{2} - \delta$ where $\delta \ll \frac{\pi}{2}$. Under this small angle assumption, Eq. S9a can

1227   be approximated by:

$$d_{\alpha3} \approx \frac{d_N - d_{C\alpha\beta}}{\cot\theta\sin\varphi + \cos\varphi + \sin\delta/\sin\theta}. \qquad \text{[S9b]}$$

Upon inversing Eq. S9b, we found that the tilting angle θ is a function of the end-to-end distance

of the $\alpha3$ domain, $i.e.$, $\theta = f(d_{\alpha3})$. Setting $\delta = 25°$, which seems reasonable as it approximates the

maximum value of φ, the structural parameters obtained by fitting are scattered in-between two

black curves on the $d_{\alpha3} - \theta$ plane marked as pulling constraint: [$d_N = 12.5$ nm, $d_{C\alpha\beta} = 3.62$ nm,

$\varphi = 0°$] and [$d_N = 11.5$ nm, $d_{C\alpha\beta} = 3.42$ nm, $\varphi = 25°$]) (Fig. 4f).

Another constraint is the tilting constraint resulted from the asymmetric unfolding and

stretching of the interdomain links between the TCR constant and variable domains. This constraint

is introduced in the model to account for the potential regulatory effect of the FG-loop on catch

bond. Notwithstanding the total number of unfolded amino acids n^* can be determined by the

validation procedure demonstrated in A.2, its breakdown into the number of unfolded amino acids

for MHC ($n_{p,MHC}$) and TCR ($n_{p,TCR}$) remains undetermined. To do this, known structures from

PDB were used to determine $n_{p,MHC}$ first. Briefly, by matching $d_{\alpha3}$ (Supplementary Fig. 4b, c) with

the end-to-end distance between C-terminal end of $\alpha3$ domain and certain point following known

PDB structure, the exact starting position of partial unfolding in MHC can be found following

additional assumption that unfolding starts from C-terminal end of $\alpha2$ domain towards $\alpha3$ domain

(Supplementary Fig. 4b). Thus, $n_{p,TCR}$ can be simply calculated as $n_{p,TCR} = n^* - n_{p,MHC}$. Upon

combining all information, the tilting angle (θ_{TCR}) of variable domains of TCR can be described by

simple trigonometrical function:

$$\theta_{\text{TCR}} \approx \tan^{-1}\left(\frac{n_{\text{p,TCR}} l_{\text{c}}}{W_{\text{TCR}}}\right), \qquad\qquad \textbf{[S10]}$$

where W_{TCR} is width of two interdomain hinges of the TCR α- and β-subunits measured from the crystal structures (Supplementary Fig. 4a, tilting constraint). In this work we use $W_{\text{TCR}} = 3.7 \pm 0.3$ nm as a representative width due to structure-to-structure variations. Thus, by comparing the tilting angle of the bonding interface θ (model parameter) to the tilting angle of the TCR θ_{TCR} (derived from another model parameter $n_{\text{p,TCR}}$ and structural constants W_{TCR} and l_{c}), we can check the validity of tilting constraint using linear regression in the θ vs θ_{TCR} plot (Supplementary Fig. 4d).

B. Kinetic model for TCR–pMHC-II bonds

B.1. Development, validation, and characterization

The purpose of this subsection is to present details of the development, validation, and characterization of the TCR–pMHC-II catch bond model omitted in the main text for simplicity, in a similar fashion as the TCR–pMHC-I catch bond model described in Section A. The two models share exact the same framework but have different detailed form of the characteristic extension change $\delta_{\text{z}}(f)$ (Eq. S5). Comparing to the TCR–pMHC-I complex, the TCR–pMHC-II complex has different docking domains and pulling geometries (one vs two transmembrane domains on both TCR and MHC). For this reason, we assume that the force-induced bonding interface tilting angle (θ) would be much smaller in the TCR–pMHC-II than TCR–pMHC-I complex. The extension at the bound state can be defined as the end-to-end distance between both end-points identified by crystal structures (E8: 2IAM, 2IAN and 2B4 (as the substitution of 3.L2): 6BGA, 3QIB):

$$z(f; 0) = d_{\text{N}}(f) + d_{\text{linker}}(f), \qquad\qquad \textbf{[S11]}$$

where d_{N} is set to be 12.3 nm based on the crystal structures and $d_{\text{linker}} \approx 9.4$ nm represents the linker (e.g., a leucine zipper) engineered at the C-termini of soluble pMHC-II constructs to stabilize both the MHC α- and β-subunits, which is often used in experiments for measuring TCR–pMHC-II catch bonds. To account for domain rotation resulted from partial unfolding inside the TCR–

1270 pMHC-II complex, we introduce one more variable, φ, as the tilting angle of the TCR constant

1271 domains. Using the tilting constraint similarly to that used in the class I model, this angle can be

1272 approximately described by structural parameters $(d_{B.I}, \theta)$ and model constants (see subsection B.2).

1273 In short, each component in the right-hand side of Eq. S11 can be expressed by the model

1274 parameters $(d_{B.I}, \theta, n^*)$, and model constants as follows.

$$z_{B.I}(f) = \left(d_{B.I}(f) + z_{p,MHC}(f)\right) \cos \theta , \qquad \text{[S12a]}$$

$$z_{C\alpha\beta}(f) = \left(d_{C\alpha\beta}(f) + z_{p,TCR}(f)\right) \cos \varphi , \qquad \text{[S12b]}$$

$$z_{linker}(f) = d_{linker}(f) \cos \theta , \qquad \text{[S12c]}$$

where $z_{p,MHC}(f)$ and $z_{p,TCR}(f)$ are respective extensions of unfolded polypeptides given by

$n_{p,MHC} l_c z_{u,p}(f)$ and $n_{p,TCR} l_c z_{u,p}(f)$, respectively, $d_{B.I}$ is the length of structure consisting of the

MHC and the TCR variable domains, and other parameters defined previously. Finally, the rate

coefficient of the TCR–pMHC-II dissociation can be developed by employing the same framework

(see Eq. S6). However, when applying the model to experimental data, we can use the constraints

of the TCR–pMHC-II interaction to make the model much simpler (see section B.2).

Validation of the class II model follows exactly the same procedure as that used in the

validation of the pMHC I model, which is done by checking self-consistent through the definition

of the reaction coordinate. By varying the $n_{p,MHC}$ from 0 to 10 (see details explained in section B.2),

the molecular extension at zero force (δ_0^*) can be estimated by using the contour length-change (δ_l)

along the dissociation coordinate:

$$\delta_l = n_{p,MHC} l_c + \Delta \rightarrow \delta_0^* \qquad \text{[S13]}$$

where $\Delta(d_{B.I}, \theta) < 0.2$ nm because $\theta < 10°$. The best-fit model parameters can be determined by

finding a subset of best-fit parameters among possible $n_{p,MHC}$ values that match the contour length-

change along the dissociation coordinate (δ_l) to the width of the free-energy well at zero force (δ_0^*),

i.e., finding $n_{p,MHC}$ such that $\delta_l(n_{p,MHC}) \rightarrow \delta_0^*$ (see Supplementary Table 5).

To characterize the class II model, we examine the model predictions by varying the model

parameters one by one while fixing the others as constants. For example, to investigate force-

induced bonding interface titling, we fixed the other model parameters (*i.e.*, $n^* = 4$, $\delta_0^* = 1.2$ nm,

$k_0 = 10$ s^{-1}, and $\Delta G_0^* = 11.5$ $k_B T$) and structural constants ($d_N = 12$ nm, $d_{B.I} = 12.3$ nm, and

$d_{linker} = 9.4$ nm for pMHC-II constructs that have linkers). As another example, we fixed $\theta = 0°$

or 3° and examined the effect of the molecular extension at zero force by varying δ_0^* from 0.3-1.8

nm. The constants used for model characterization were selected by their averages from the

corresponding values used to fit actual experiments. The parameters for free-energy landscape

construction, the energy barrier height (ΔG^*) and energy well width (δ^*) as functions of force, are

given by Eq. S8, exactly the same as the pMHC I model.

Fitting of class II model to experimental data and examination of the biological relevance

of the best-fit parameters were done the same way as the class I model.

B.2. Class II model constraints

The purpose of this subsection is to describe the constraints of the class II model, which

share similar ideas to those of the class I model (*e.g.*, pulling and tilting constraints) but differ in

their specific expressions. To formulate the pulling constraint, we again used the fact that the pulling

force direction must aligns with the line connecting to the C-termini of the TCR and pMHC

molecules so that the coordinate perpendicular to the force direction is invariant. Using model

parameters and structural constants, this pulling constraint can be written as:

$$(d_{B.I} + z_{p,MHC} + d_{linker})\sin\theta = (d_{C\alpha\beta} + z_{p,TCR})\sin\varphi \,, \qquad \textbf{[S14a]}$$

1309   which can be solved for $z_{p,TCR}$ explicitly:

$$z_{p,TCR} = \left(d_{B.I} + z_{p,MHC} + d_{linker}\right)\frac{\sin\theta}{\sin\varphi} - d_{C\alpha\beta} .$$ [S14b]

By combining Eq. S12 and S14, the total extension (z) can be calculated as the sum of all component extensions ($\sum_i z_i$) at dissociation ($\delta_l > 0$):

$$z(f;\delta_l) = \left(d_{B.I}(f) + z_{p,MHC}(f) + d_{linker}(f)\right)(\cos\theta + \sin\theta\cot\varphi) .$$ [S15]

This equation states that only the number of unfolded amino acids in MHC ($n_{p,MHC}$) affects extension change during transition. The total number of unfolded amino acids from both TCR and MHC can be estimated by Eq. S14b:

$$n^* \approx \left| \frac{(d_{B.I}+d_{linker})\frac{\sin\theta}{\sin\varphi}-d_{C\alpha\beta}}{l_c} + \left(1+\frac{\sin\theta}{\sin\varphi}\right)n_{p,MHC} \right| .$$ [S16]

which can be approximately calculated using contour lengths of length components at force-free state.

By assuming small angle perturbation, which seems reasonable, we further reduce the number of model parameters after relating the tilting angle of the TCR constant domain (φ) and the titling angle of the bonding interface by the following equation:

$$\cot\varphi \approx \frac{d_N-d_{B.I}}{d_{B.I}+d_{linker}}\cot\theta .$$ [S17]

Thus, all terms including φ can be re-expressed by using Eq. S17.

Additionally, the tilting constraint can be expressed as follows:

$$\frac{\tan^{-1}\left(\frac{n_{p,TCR}l_c}{w_{TCR}}\right)}{\theta+\varphi} \approx 1 .$$ [S18]

Thus, by using Eq. S15 and S17, only 4 fitting parameters, two structural parameters ($d_{B.I}$, θ) and two biophysical parameters (k_0, δ_0^*), were used to fit the class II model to data.

C. A general biophysical limit of model parameters

The purpose of this section is to describe a general biophysical limit that constrains the fitted model parameters, which is used in Fig. 6 to accept the correct model application to data of matched MHC class and reject incorrect model application to data of mismatched MHC class. The idea is

that, even if the model is capable of fitting experimental data and the parameters are self-consistent

with one another within the model, their values should be within known limits. A prototypical

example of such a biophysical limit involves the molecular extension per unfolded amino acids. It

follows from Eqs. S7 and S13 that the average molecular extension at zero force over all data ($\langle \delta_0 \rangle$)

should be a linear function of n^* such that $\langle \delta_0 \rangle = an^* + b$. The y-axis intercept b can be

determined from the slip bond data because slip bonds are not expected to have unfolded amino

acids (i.e., $n^* = 0$) but still have a nonzero extension. The slope a is constrained by the fact that the

contour length of a single amino acid has a small range ($\sim 0.4 \pm 0.02$ nm/a.a.)[6,13,14]; thus, the average

molecular extension ($\langle \delta_0 \rangle$) per unfolded amino acid should be bounded by:

$$0 < a < 0.4 \,. \hspace{3cm} \textbf{[S19]}$$

Imposing this range limit of $\langle \delta_0 \rangle$/a.a. would enable us to rule out inappropriate application

of the model even if such application could achieve reasonable level of goodness-of-fit. An

exceedingly large value of a ($> 0.4 \pm 0.02$ nm/a.a.) estimated by the model would mean that the

model is inappropriate for the data.

Conversely, a nearly zero estimate of a would indicate that the model is inappropriate for

catch-slip bond data because, for the model to fit catch-slip bonds, it requires $n^* > 0$ (see Figs. 2c-

g and 3f, *middle row*). A parameter estimation of $a \approx 0$ indicates the lack of dependence of the

model behavior on n^*, which abolishes the model's ability to predict TCR signaling and antigen

discrimination, making the model irrelevant to biology.

The sturdier class II than class I pMHC structure also precludes large rotation during

conformational change at transition state, leaving only unfolding and stretching along the force.

Thus, the average molecular extension per amino acid should be close to 0.4 nm/a.a. On the other

hand, $a \approx 0$ is expected from fitting the slip-only data because such data correspond to the $n^* = 0$

case, which makes it difficult to robustly estimate the correct a value. Thus, the average molecular

extension per amino acid should be well-correlated with each other (high level of goodness-of-fit

as measured by R^2) and in the range between 0 to 0.4 nm/a.a. In Fig. 6, we used these criteria to

test the appropriateness of cross-applying the class I model to class II data and *vice versa*, showing

that it is appropriate to use either model to fit matched data but inappropriate to use either model to

fit mismatch data.

1357

Supplementary Figures

Supplementary Figure 1 | Fitting class I model and two-pathway model to data of TCR–

pMHC-I bond lifetime *vs* force. a-f Fitting of theoretical $1/k(F)$ curves of the class I model

(solid curves) and the two-pathway model (dashed curves) to experimental bond lifetime *vs* force

data (*points*, mean ± sem) of 5 TCRs and their mutants interacting with different pMHCs as

described below (re-analyzed from Refs.[4,15,16,17,18]). (**a**) OT1 TCR expressed on CD8$^+$ naïve T

cells (first three panels) or CD4$^+$CD8$^+$ thymocytes (last two panels) interacting with indicated

p:H2-K^{b}α3A2. (**b**) 2C TCR expressed on CD8$^+$ naïve T cells interacting with indicated p:H2-

K^{b}α3A2 (*top*, 1st panel) or on CD8$^-$ hybridomas interacting with indicated peptides presented by

WT (*top*, 2nd panel) or disulfate-locked MT (*top*, 3rd panel) H2-K^b, 2C TCR with indicated point

mutations expressed on CD8$^-$ hybridomas interacting with R4:H2-K^b (*top*, 4th and 5th panels), 2C

TCR expressed on CD8$^-$ hybridomas interacting with R4 peptide presented by H2-K^b with

indicated point mutations (*bottom*, 1st-3rd panels), or 2C TCR with indicated point mutations

expressed on CD8$^-$ hybridomas interacting with R4 peptide presented by H2-K^b with indicated

point mutations (*bottom*, 4th and 5th panels). (**c**) 2C TCR expressed on CD8$^+$ naïve T cells

interacting with indicated p:H2-L^d(m31) with truncated α3 domain. (**d**) 1G4 TCR expressed on

CD8$^-$ hybridomas interacting with indicated p:HLA-A2. The first panel of a and the first two top

panels of b are replotted from **Fig. 3a** and **b** for completeness. (**e**) Purified N15 TCRαβ

interacting with indicated p:H2-K^b. f, TRBV TCRs of canonical (B13.C1 and B17.C1) and

reverse (B17.R1 and B17.R2) pMHC-I docking orientation expressed on hybridomas interacting

with NP$_{366}$:H2-D^{bD227K}. See Supplementary Table 3 for a list of the interacting molecules.

1381

Supplementary Figure 2 | Correlation of model parameters with pMHC-I biological activity and TCR bond type. a Relative breadth *vs* the reciprocal peptide dose required to stimulate half maximal of OT1 T cell proliferation. **b-c** Model parameters θ (the tilted angle of the bonding interface, *upper*) and n^* (the number of unfolded amino acids, *lower*) that best-fit the data in Supplementary **Fig. 1b** and **d** are plotted *vs* the indicated WT and mutant 2C (**b**) and 1G4 (**c**) TCRs with their indicated pMHCs. **d-f** Scattergrams of θ (**d**), n^* (**e**), and $1/k_0$ (**f**) *vs* $I = L/(1 + B)$ (catch bond intensity) are plotted using the data from **Figs. 3e** & **3f**, **4b**, and **Supplementary Figs. 2b** & **2c** to examine correlation. Blue-open symbols indicate data of known T cell biological activities (ligand potencies) induced by the corresponding TCR–pMHC-I interactions that correlate to catch bond intensity. Green-closed symbols indicate data of known effects on catch bond metrics by targeted mutations on the TCR, MHC, or both. See Supplementary Table 3 for a list of the interacting molecules.

1397

Supplementary Figure 3 | Correlation of force-induced TCR–pMHC-I hydrogen (H) bonds

with bonding interface tilting angle. a H-bond distributions at bonding interface between 2C

TCR and the indicated pMHCs (R4, dEV8, EVSV, or L4) in the presence (*solid curves*) or

absence (*dotted curves*) of the force (obtained by re-analysis of SMD simulation results from Ref.

[4] and fitted by Gaussian functions). **b** Plots of average number of H-bonds (mean ± sd) with

respect to fitted tilting angle (with fitting error) of bonding interface in the absence (*red*) and

presence (*blue*) of force as well as their average (*gray*). Average numbers of H-bonds were

determined as mean value for each Gaussian distribution at each titling angle. All colored dashed

lines are linear fits.

1407

1408

1409

Supplementary Figure 4 | Pulling and tilting constraints of class I model. a An illustration of

both pulling and tilting constraints. Length of yellow line was used to set up the equation of the

pulling constraint between $d_{\alpha3}$ and θ. The tilted angle from the blue lines was used to check

asymmetric stretching of TCR. **b** A pMHC-I structure based on PDB code 2CKB. Partial

unfolding of MHC is assumed to start from the end of the α1-α2 domain towards the α3 domain

as suggested by SMD simulation[4]. **c** Representative end-to-end distance of $d_{\alpha3}$ *vs* the number of

unfolded amino acids. The distance is calculated from C-terminal end of the **α3** domain based on

the structure shown in (**b**). Distance is calculated using the PBD structure for each construct. $d_{\alpha3}$

distance obtained from fitting the model to data is converted to $n_{p,MHC}$ (*inset*). **d** Tilted angle of

the TCR variable domains (θ_{TCR}) *vs* the titled angle of bonding interface (θ) as one of the fitting

1420   parameters. The validity of asymmetric stretching of TCR was checked by the linear relationship

1421   between the two angles. All errors in (**d**) represent fitting errors.

1422

1423

1424

Supplementary Figure 5 | Determining the number of clusters in parameter space. a
Definitions of the *i*-th point-to-centroid Euclidean distance (blue line) and of the within-cluster
sums of square point-to-centroid distance when the number of clusters is 1. Scatters were
replotted from **Fig. 4e**. **b** Sum of all squared Euclidean distances *vs* number of possible clusters.
Point of abrupt reduction of steepness of slope leading to a straight line (*blue solid-line*) was
identified a number of clusters (*blue dotted-line*).

Supplementary Figure 6 | Lack of correlations between best-fit parameters of the two-pathway model and T cell function. a-d Correlation plots of $1/k_{0,p1}$ (**a**), $1/k_{0,p2}$ (**b**), $\delta^*_{0,p1}$ (**c**), and $\delta^*_{0,p2}$ (**d**) *vs* % changes in effector function. T cell effector function was defined by the peptide dose required to achieve half-maximal proliferation ($1/EC_{50}$) of OT1 T cells (*blue*) or to generate 40% B cell apoptosis ($1/EC_{40}$)[19] for 3.L2 T cells (*red*), or the area under the dose response curve (AUC) of 2B4 hybridoma IL-2 production[20] (*orange*). Color-matched solid lines are linear fits for the indicated TCR–pMHC systems. Also indicated are the slopes and R^2 values. **e-l** Correlation plots of $1/k_{0,p1}$ (**e, i**) $1/k_{0,p2}$ (**f, j**), $\delta^*_{0,p1}$ (**g, k**), and $\delta^*_{0,p2}$ (**h, l**) *vs* the intensity of catch bond evaluated from the indicated class I (**e-h**) or class II (**i-l**) datasets. Solid lines are linear fits, which are color-matched to indicate the 'potency-' or 'mutation-' related datasets for the class I system. All error bars are fitting error (Supplementary Table 4).

1445

Supplementary Figure 7 | Characterization of the energy landscape of TCR–pMHC-II

dissociation. a-d Plots of transition state location δ^* (**a**, **c**) and height of energy barrier ΔG^* (**b**, **d**)

vs force F for changing θ and n^* (**a**, **b**) or δ_0^* and n^* (**c**, **d**) while keeping $\delta_0^* = 1.2$ nm (**a**, **b**) or θ

$= 0, 3°$ (**c**, **d**). Inset color bars indicate parameter values used to plot the color-match theoretical

curves.

1451

1452

1453

Supplementary Figure 8 | Correlation of metrics of TCR–pMHC-II bond lifetime *vs* force curves with T cell biological activity. a-c Dimensional metrices, t_{peak} (**a**), ΔF (**b**) and relative breadth of catch-slip bond B (**c**) *vs* reciprocal % change (relative to WT) in T cell effector function, i.e., the peptide dose required for 3.L2 T cells to generate 40% B cell apoptosis $(1/EC_{40})$[19] (*red*) or the area under the dose response curve (AUC) of 2B4 hybridoma IL-2 production[20] (*blue*) plots.

1454

1455

1456

1457

1458

1459

1460

1461

1462

Supplementary Figure 9 | Comparing fitting curves of class I and class II models to the same data. a-i Fitting of theoretical $1/k(F)$ curves predicted by the class I (blue) or class II (*red*) model to experimental lifetime (*points*, mean ± sem) *vs* force data of TCR bonds with pMHC-I (**a-f**) or pMHC-II (**g-i**) ligands, as described below (partly presented in Supplementary Fig. 1, re-analyzed from Refs. [4,15,16,17,18,21,22,23]): OT1 (**a**) or 2C (**b**) TCR expressed on CD8[+] naïve T cells interacting with indicated p:H2-K[b]α3A2; WT or mutant 2C (**c**) or 1G4 (**d**) TCR expressed on hybridomas interacting with indicated peptides presented by WT or MT H2-K[b] or HLA-A2. B13.C1 and B17.C1 TCR expressed on hybridomas interacted with NP_{366} bound to the D227K MT of H-2D[b] to prevent CD8 binding (**e**). Soluble N15 TCRαβ interacting with indicated p:H2-K[b] (**f**). 3.L2 TCR expressed on CD4[-]CD8[+] T cells interacting with indicated p:I-E[k] (**g**). WT or MT 2B4 TCRs expressed on CD4[-] hybridomas interacting with K5:I-E[k] (**h**). E8 TCR expressed on CD4[-] Jurkat cells interacting with TPI:HLA-DR1 (**i**). See Supplementary Table 3 and 6 for lists of the interacting molecules.

1478

Supplementary Figure 10 | Comparison of goodness-of-fit measures among the class I and II models and the two-pathway model to class I and class II data. a-b Residual sum of squares (RSS, **a**) and Chi-square values (χ^2, **b**) obtained using three models – two-pathway model (*black*), TCR–pMHC-I model (red), and TCR–pMHC-II model (*blue*) – to fit pMHC-I data. **c-d** Residual sum of squares (RSS, **c**) and Chi-square values (χ^2, **d**) obtained using three models – two-pathway model (*black*), TCR-pMHC-I model (red), and TCR-pMHC-II model (*blue*) – to fit pMHC-II data. Individual scattered points, median, 25%, 75% and 5%, 95% whiskers are shown. Statistical tests were done using paired *t* test with two-tail.

1479

1480

1481

1482

1483

1484

1485

1486

1487

1488

Supplementary Figure 11 | Comparison of pMHC interactions with cell surface and soluble

TCRs. a Schematics of three biomembrane force probe (BFP) experiments. The BFP were set up

in three configurations (*upper row*) depending on the molecular systems tested (*lower row*). In the

first configuration (*left column*), the pMHC was coated on a BFP bead (*left*) and the TCR-CD3

complex was expressed on a T cell (*right*). In the second configuration (*middle column*), the

purified TCRαβ was coated on a BFP bead (*left*) and the pMHC was expressed on an APC

(*right*). In the third configuration, the pMHC was coated on a BFP bead (*left*) and the purified

TCRαβ was coated on a glass bead (*right*). The TCR–pMHC complexes were drawn based on the

cryoEM structure with CD3 (6JXR) or crystal structure without CD3 (2CKB). **b-c** Fitting of

theoretical $1/k(F)$ (*curves*) to experimental bond lifetime *vs* force data (*points*, mean ± sem) of the

following interactions: OT1 or 2C TCR respectively expressed on CD8[+] naïve T cells (*red*) or

CD8[-] hybridomas (*green*) or coated on beads (*pink* for OT1 and *yellow-green* for 2C) respectively

interacting with OVA:H2-K[b]α3A2 or QL9:H2-L[d](m3), both MHC class I molecules[24] (**b**) or of

E8 TCR expressed on CD8⁻ Jurkat (*sky-blue*) or coated on beads interacting with TPI:HLA-DR1, a MHC class II, coated on beads (*blue*) or expressed on THP-1 cells (*purple*)[22] (**c**). **d** Comparison of the catch bond intensity (*top*) and best-fit parameters δ_0^* (*middle*) and θ (*bottom*) evaluated from data in (**b**) and (**c**) of TCR–pMHC interactions measured using the three configurations depicted in (**a**). All error bars indicate fitting error. See Supplementary Table 3 and 6 for lists of the interacting molecules.

1510 **Supplementary Table 1 | Summary of model constants**

Parameters	Symbol	Value	Reference
Elastic modulus of the folded globular domain	E_d	100 pN	[2]
Average contour length	l_c	0.36 nm	[6,7,8]
Persistence length	l_p	0.39 nm	[6,7,8]
Elastic modulus of polypeptides	E_p	50 μN	[8]
Force-free extension of bound state	$d_{N,c}$	10.9 – 12.7 nm	[‡]
Force-free length of TCRαβ constant domain	$d_{C\alpha\beta,c}$	3.3 – 3.8 nm	[‡]
Angle shown in Fig. 2b and Supplementary Figure 4	φ	5 – 23.5 °	[‡]

1511 ‡Detailed values for each of the OT1, 2C, 1G4, N15, B13.C1, B17.C1, B17.R1, B17.R2, 3.L2, E8, and 2B4 TCR
1512 structures are described in Supplementary Model Derivations.
1513
1514

 Supplementary Table 2 | An example of finding the best-fit parameters for OT1 TCR–

1516 **OVA:H2-K^{b}a3A2 bond**

n^*	$d_{\alpha 3}$ (nm)[‡]	θ (°)[‡]	$n^* l_c$ (nm)	δ_l (nm)	δ_0 (nm)[‡]	k_0 (s^{-1})[‡]
1	5.2	61.9	0.36	-0.8	0.7	4.84
2	5.1	59.7	0.72	-0.5	1.1	4.66
3	4.9	57.3	1.08	0.0	1.4	4.54
4	4.7	55.4	1.44	0.2	1.7	4.44
5	4.5	54.1	1.80	0.6	1.9	4.34
6	4.4	53.5	2.16	1.0	2.1	4.26
7	4.3	53.4	2.52	1.3	2.2	4.20
8	4.2	53.6	2.88	1.7	2.4	4.14
9	4.1	54.1	3.24	2.0	2.5	4.09
10	4.1	54.8	3.60	2.4	2.6	4.04
11	**4.0**	**55.5**	**3.96**	**2.7**	**2.7**	**4.00**
12	4.0	56.4	4.32	3.1	2.8	3.73
13	1.8	25.8	4.68	3.5	3.9	3.76
14	2.4	70.5	5.04	3.9	3.3	3.87

1517 [‡]The fitting parameters derived from model (see Supplementary Model Derivation, Section A.3).

Supplementary Table 3 | TCR–pMHC-I bond summary and their best-fitting parameters

TCR	peptide	MHC class I	n^*	$d_{\alpha 3}$ ± error (nm)	θ ± error (°)	δ_0 ± error (nm)	k_0 ± error (s^{-1})	$\Delta G_0^{\ddagger}$ ± error ($k_B T$)
OT1[†]	OVA	H2-K^b (α3A2)	11	4.0 ± 0.2	55.5 ± 1.5	2.7 ± 0.3	4.0 ± 2.3	12.4 ± 0.5
OT1[ℙ]	OVA	H2-K^b (α3A2)	6	4.0 ± 0.2	34.9 ± 4.3	1.1 ± 0.1	9.6 ± 1.6	11.6 ± 0.1
OT1[†]	A2	H2-K^b (α3A2)	7	3.9 ± 0.2	44.0 ± 3.1	1.1 ± 0.1	4.1± 1.0	12.4 ± 0.2
OT1[†]	G4	H2-K^b (α3A2)	9	5.0 ± 0.4	35.2 ± 2.2	2.7 ± 0.4	4.2 ± 1.1	12.4 ± 0.2
OT1[†]	E1	H2-K^b (α3A2)	0	0.0 ± 0.0	1.0 ± 1.2	0.9 ± 0.1	2.4 ± 0.2	13.0 ± 0.1
OT1[†]	R4	H2-K^b (α3A2)	0	0.0 ± 0.0	0.0 ± 0.9	1.0 ± 0.3	2.0 ± 0.3	13.1 ± 0.1
OT1[*]	OVA	H2-K^b (α3A2)	6	1.6 ± 0.2	17.5 ± 2.9	1.6 ± 0.2	3.3 ± 0.6	12.6 ± 0.2
OT1[*]	OVA	H2-K^b	11	1.9 ± 0.3	26.6 ± 4.9	3.1 ± 0.3	3.6 ± 0.3	12.5 ± 0.1
OT1[*]	Q4H7	H2-K^b (α3A2)	1	0.0 ± 0.0	0.0 ± 0.2	0.4 ± 0.2	4.3 ± 0.8	12.4 ± 0.2
OT1[*]	Q4R7	H2-K^b (α3A2)	1	0.0 ± 0.0	0.0 ± 0.2	0.4 ± 0.1	3.8 ± 0.8	12.4 ± 0.2
2C[†]	R4	H2-K^b (α3A2)	7	2.5 ± 0.5	19.3 ± 1.3	2.3 ± 0.2	7.4 ± 4.8	11.8 ± 0.5
2C[†]	dEV8	H2-K^b (α3A2)	3	2.6 ± 0.5	13.3 ± 1.7	2.0 ± 0.2	75.6 ± 18.7	9.5 ± 0.2
2C[†]	L4	H2-K^b (α3A2)	0	0.0 ± 0.0	0.0 ± 0.2	0.4 ± 0.1	3.8 ± 1.5	12.5 ± 0.3
2C[*]	R4	H2-K^b (α3A2)	6	0.9 ± 0.1	7.3 ± 1.6	1.3 ± 0.2	3.7 ± 0.3	12.5 ± 0.1
2C[*]	R4	H2-K^b	12	1.7 ± 0.3	21.8 ± 2.5	2.8 ± 0.3	5.5 ± 0.6	12.1 ± 0.1
2C	R4	H2-K^b	7	4.0 ± 0.3	28.2 ± 2.0	1.5 ± 0.4	8.9 ± 4.7	11.6 ± 0.4
2C	R4	H2-K^b C-C locked[∗]	7	4.1 ± 0.2	24.9 ± 2.0	1.6 ± 0.3	14.8 ± 4.4	11.1 ± 0.3
2C	EVSV	H2-K^b	2	0.4 ± 0.2	6.4 ± 1.8	0.9 ± 0.3	9.0 ± 2.8	11.6 ± 0.3
2C	L4	H2-K^b	1	0.1 ± 0.1	2.1 ± 1.5	0.6 ± 0.1	6.8 ± 1.3	11.9 ± 0.1
2C E56A	R4	H2-K^b	9	4.1 ± 0.5	20.7 ± 3.3	1.2 ± 0.2	7.4 ± 3.5	11.8 ± 0.4
2C N30A	R4	H2-K^b	7	4.3 ± 0.1	23.6 ± 2.8	1.4 ± 0.4	9.5 ± 3.6	11.6 ± 0.3
2C	R4	H2-K^b Q72A	7	4.2 ± 0.4	25.3 ± 2.3	1.3 ± 0.3	8.0 ± 4.5	11.7 ± 0.4
2C	R4	H2-K^b K146A	4	0.4 ± 0.2	4.7 ± 1.0	1.1 ± 0.2	6.7 ± 1.3	11.9 ± 0.1
2C	R4	H2-K^b R79A	4	0.6 ± 0.3	9.1 ± 0.5	2.3 ± 0.6	13.2 ± 6.3	11.2 ± 0.4
2C E56A	R4	H2-K^b Q72A	4	0.4 ± 0.3	5.6 ± 1.2	1.1 ± 0.3	6.9 ± 2.7	11.9 ± 0.3
2C N30A	R4	H2-K^b K146A	2	0.4 ± 0.2	6.8 ± 0.9	1.5 ± 0.4	7.2 ± 3.8	11.8 ± 0.4
2C[†]	QL9	H2-L^d (m31)	6	2.5 ± 0.7	18.4 ± 4.5	3.2 ± 0.4	9.5 ± 7.9	11.6 ± 0.6
2C[ℙ]	QL9	H2-L^d (m31)	3	2.0 ± 0.6	2.0 ± 0.9	1.0 ± 0.1	26.9 ± 7.0	10.5 ± 0.2
1G4	9C	HLA-A2	4	1.8 ± 0.7	10.1 ± 0.5	2.7 ± 0.4	13.7 ± 3.5	11.2 ± 0.2

1G4	9C	HLA-A2 R75A	6	1.4 ± 0.2	9.4 ± 0.4	3.7 ± 0.3	8.1 ± 8.0	11.0 ± 0.7
1G4	9C	HLA-A2 A236T	5	4.2 ± 0.2	29.4 ± 2.6	1.8 ± 0.3	8.6 ± 2.0	11.7 ± 0.2
1G4	9C	HLA-A2 F8V	4	4.1 ± 0.4	26.8 ± 1.4	2.0 ± 0.4	19.7 ± 4.5	10.8 ± 0.2
N15[¶]	VSV8	H2-K^b	9	3.2 ± 0.3	32.0 ± 4.6	4.5 ± 0.5	32.9 ± 10.0	10.3 ± 0.2
N15[¶]	L4	H2-K^b	5	1.8 ± 0.3	13.7 ± 1.5	4.3 ± 0.5	48.1 ± 21.1	9.9 ± 0.4
N15[¶]	SEV9	H2-K^b	0	0.0 ± 0.0	0.0 ± 0.0	0.6 ± 0.1	0.6 ± 0.1	14.3 ± 0.2
B13.C1[#]	NP$_{366}$	H-2D^b (D227K)	8	4.2 ± 0.4	40.0 ± 6.2	1.4 ± 0.3	1.6 ± 0.8	13.3 ± 0.4
B17.C1[#]	NP$_{366}$	H-2D^b (D227K)	6	4.2 ± 00.5	33.2 ± 4.1	2.2 ± 0.5	11.3 ± 6.6	11.4 ± 0.2
B17.R1[#]	NP$_{366}$	H-2D^b (D227K)	0	0.0 ± 0.0	0.0 ± 0.0	0.6 ± 0.2	0.3 ± 0.1	14.9 ± 0.3
B17.R2[#]	NP$_{366}$	H-2D^b (D227K)	3	0.4 ± 0.3	0.4 ± 0.2	1.2 ± 0.2	0.4 ± 0.1	14.9 ± 0.2

[ᴾ]Purified recombinant TCRs coated on beads were used in BFP measurement.

[*]OT1 or 2C TCR expressed on CD4$^+$CD8$^+$ thymocytes were used.

[†]OT1 or 2C TCR expressed on naïve T cells from transgenic mice were used. In other cases, 2C or 1G4 TCR expressed on 58α^{-1}β^{-1} hybridomas and J76 Jurkat cells were used.

[¶]Soluble mouse N15 TCRαβ was used to measure bond lifetime in optical tweezers[16].

[#]Mouse TRBV TCRs (B13.C1 and B17.C1 with canonical docking orientation and B17.R1 and B17.R2 with reverse docking orientation) expressed on 5KC hybridomas interacted with NP$_{366}$ bound to the D227K MT of H-2D^b to prevent CD8 binding[18]

[ᵝ]I1C mutation in β2m domain, G120C and C121S mutations in α domain (H2-K^b) were introduced.

Note that the best-fit parameters were chosen by the procedure shown in Supplementary Table 2 or determined as the parameter set with the closest δ_1 value to δ_0^* as well as the smallest RSS. All errors are standard error of mean after optimizing the best-fit parameters. All values were rounded to one decimal place.

Supplementary Table 4 | Two-pathway model summary and their best-fitting parameters

TCR-pMHC class I

TCR	peptide	MHC class I	$k_{0,p1}$ (s^{-1})	$k_{0,p1}$ error	$\delta^*_{0,p1}$ (nm)	$\delta^*_{0,p1}$ error	$k_{0,p2}$ (s^{-1})	$k_{0,p2}$ error	$\delta^*_{0,p2}$ (nm)	$\delta^*_{0,p2}$ error
OT1[†]	OVA	H2-Kb(a3A2)	3.99	2.7	-1.36	1.13	0.41	0.49	0.43	0.31
OT1[‖]	OVA	H2-Kb(a3A2)	7.77	4.54	-1.81	2.16	3.29	2.38	0.44	0.28
OT1[†]	A2	H2-Kb(a3A2)	3.36	0.57	-1.01	0.41	0.86	0.36	0.35	0.1
OT1[†]	G4	H2-Kb(a3A2)	1.99	3.05	-3.13	37.34	1.47	3.02	1.04	1.17
OT1[†]	E1	H2-Kb(a3A2)	0.39	0.41	-2.32	20.95	2.01	17.59	0.95	5.24
OT1[†]	R4	H2-Kb(a3A2)	N.A	N.A	N.A	N.A	2	0.39	0.91	0.36
OT1*	OVA	H2-Kb(a3A2)	3.02	1.56	-0.84	0.71	0.35	0.44	0.33	0.24
OT1*	OVA	H2-Kb	2.47	0.88	-0.52	0.3	0.05	0.09	0.67	0.35
OT1*	Q4H7	H2-Kb(a3A2)	N.A	N.A	N.A	N.A	4.09	0.53	0.13	0.07
OT1*	Q4R7	H2-Kb(a3A2)	N.A	N.A	N.A	N.A	3.44	0.25	0.16	0.03
2C[†]	R4	H2-Kb(a3A2)	6.09	3.45	-1.13	0.71	0.28	0.39	0.56	0.35
2C[†]	dEV8	H2-Kb(a3A2)	15.44	16.59	-0.81	0.73	0.06	0.25	0.99	0.93
2C[†]	L4	H2-Kb(a3A2)	N.A	N.A	N.A	N.A	3.85	0.45	0.36	0.08
2C*	R4	H2-Kb(a3A2)	3.37	0.65	-0.26	0.31	0.13	0.39	0.75	0.58
2C*	R4	H2-Kb	5.44	5.35	-1.07	1.19	0.67	0.94	0.25	0.27
2C	R4	H2-Kb	8.04	3.69	-1.43	0.94	1.46	1.04	0.4	0.2
2C	R4	H2-Kb(C-C locked ⚹)	12.63	3.35	-1.43	0.64	2.48	1.36	0.49	0.18
2C	EVSV	H2-Kb	7.13	7.6	-0.08	2.73	0.28	6.44	1.02	5.08
2C	L4	H2-Kb	5.62	6.83	0	4.68	0.49	16.21	0.9	7.09
2C E56A	R4	H2-Kb	5.27	2.9e4	-12.7	23.38	2.79	0.9	0.41	0.13
2C N30A	R4	H2-Kb	7.91	4.32	-1.21	1.42	1.63	2.4	0.56	0.46
2C	R4	H2-Kb Q72A	7.52	1e4	-14.21	59.17	2.83	1.49	0.29	0.2
2C	R4	H2-Kb K146A	3.02	1.63	-1.21	1.35	3.83	1.57	0.2	0.11
2C	R4	H2-Kb R79A	12.37	2.24	-0.48	0.26	0.3	0.52	1.12	0.47
2C E56A	R4	H2-Kb Q72A	3.2	5e3	-15	69.15	4.26	1.13	0.29	0.1
2C N30A	R4	H2-Kb K146A	6.76	2.61	-0.39	1.06	0.26	1.57	1.37	2.12
2C[†]	QL9	H2-Ld (m31)	14.16	22	-2.24	1.3	0.19	0.17	0.53	0.25
2C[‖]	QL9	H2-Ld (m31)	15.91	5.12	-0.82	0.53	2.02	1.73	0.47	0.18
1G4	9C	HLA-A2	6.25	8.43	-0.98	1.23	0.05	0.27	0.97	1.4
1G4	9C	HLA-A2 R75A	7.3	7.73	-1.36	1.3	0.19	0.48	0.93	0.84
1G4	9C	HLA-A2 A236T	8.6	16.26	-2.01	3.81	0.82	1.29	0.4	0.46
1G4	9C	HLA-A2 F8V	7.72	33.1	-0.92	4.07	0.18	2.08	0.8	3.02
N15[¶]	VSV8	H2-Kb	33.35	222.82	-1.8	2.86	0.05	0.1	0.43	0.34
N15[¶]	L4	H2-Kb	7.96	20.51	-1.12	1.22	0.01	0.04	0.98	0.78
N15[¶]	SEV9	H2-Kb	N.A	N.A	N.A	N.A	0.64	0.47	0.58	0.38
B13.C1[#]	NP366	H-2Db (D227K)	4.29	6.09	-2.54	1.62	0.32	0.09	0.14	0.06
B17.C1[#]	NP366	H-2Db (D227K)	14.64	10.72	-1.93	0.66	0.54	0.18	0.41	0.08

B17.R1#	NP366	H-2Db (D227K)	N.A	N.A	N.A	N.A	0.35	0.14	0.61	0.34
B17.R2#	NP366	H-2Db (D227K)	0.29	0.1	0	0.75	0.02	0.14	0.78	1.16

TCR-pMHC class II

TCR	peptide	MHC class II	$k_{0,p1}$ (s^{-1})	$k_{0,p1}$ error	$\delta^*_{0,p1}$ (nm)	$\delta^*_{0,p1}$ error	$k_{0,p2}$ (s^{-1})	$k_{0,p2}$ error	$\delta^*_{0,p2}$ (nm)	$\delta^*_{0,p2}$ error
3.L2	Hb	I-E^k	1.54	1.03	-1.57	0.99	0.16	0.11	0.39	0.17
3.L2	T72	I-E^k	2.11	2.22	-1.01	2	0.24	0.8	0.49	0.8
3.L2	A72	I-E^k	4.25	7.93	-0.08	1.73	0.32	8.55	0.58	4.46
3.L2	I72	I-E^k	2.77	4.91	-0.24	1.15	1.13	5.13	0.23	0.54
E8	TPI	HLA-DR1	17.54	12.47	-0.39	0.45	0.29	1.06	0.58	0.53
E8ⁱ	TPI	HLA-DR1	10.13	11.77	-0.42	0.73	0.44	1.53	0.38	0.44
E8	TPI	HLA-DR1	53.09	70.85	-1.5	0.78	0.5	0.44	0.44	0.18
2B4 NP	K5	I-Ek	4.26	8.53	-2.12	2.11	0.14	0.13	0.45	0.26
2B4 WT	K5	I-Ek	5.99	10.17	-1.64	1.27	0.06	0.07	0.81	0.33
2B4 GK	K5	I-Ek	9.53	22.44	-1.88	1.89	0.18	0.16	0.51	0.23
2B4 HN	K5	I-Ek	2.53	1.57	-0.29	1.28	0.35	1.67	0.51	0.8

N.A indicates not applicable.

ⁱPurified recombinant TCRs coated on beads were used in BFP measurement.

*OT1 or 2C TCR expressed on CD4$^+$CD8$^+$ thymocytes were used.

†OT1 or 2C TCR expressed on naïve T cells from transgenic mice were used. In other cases, 2C or 1G4 TCR expressed on 58α^{-1}β^{-1} hybridomas and J76 Jurkat cells were used.

¶Soluble mouse N15 TCRαβ was used to measure bond lifetime in optical tweezers[16].

#Mouse TRBV TCRs (B13.C1 and B17.C1 with canonical docking orientation and B17.R1 and B17.R2 with reverse docking orientation) expressed on 5KC hybridomas interacted with NP$_{366}$ bound to the D227K MT of H-2D^b to prevent CD8 binding[18]

‡I1C mutation in β2m domain, G120C and C121S mutations in α domain (H2-K^b) were introduced.

Note that the best-fit parameters were extracted from equation (4) (two-pathway model) in the main text. Fitting errors are indicated.

 Supplementary Table 5 | An example of finding the best-fit parameters for the 3.L2 TCR–

1547   **Hb:I-E^k bond**

n_{MHC}^{*}†	$d_{\text{B.I}}$ (nm)‡	θ (°)‡	$n^{*}l_{\text{c}}$ (nm)	δ_{I} (nm)	δ_{0} (nm)‡	k_{0} (s^{-1})‡
1	8.9	0.0	0.4	0.6	0.7	2.0
2	8.8	0.0	0.7	0.9	1.2	1.9
3	8.6	0.5	1.1	1.3	1.5	1.8
4	**7.9**	**3.1**	**1.4**	**1.6**	**1.6**	**1.8**
5	7.9	2.6	1.8	2.0	1.7	1.7
6	7.8	2.1	2.2	2.4	1.9	1.7
7	8.0	1.5	2.5	2.7	2.1	1.6
8	7.8	0.6	2.9	3.1	2.2	1.6

1548   †n^{*} can be estimated from n_{pMHC}^{*}.
1549   ‡ See Supplementary Model Derivations, Section B.
1550

Supplementary Table 6 | TCR–pMHC-II bond summary and their best-fit parameters

TCR	peptide	MHC class II	n^*	$d_{B.I} \pm$ error (nm)	$\theta \pm$ error (°)	$\delta_0 \pm$ error (nm)	$k_0 \pm$ error (s^{-1})	$\Delta G_0^{\ddagger} \pm$ error (k_BT)
3.L2	Hb	I-E^k	4	8.0 ± 0.6	3.2 ± 0.6	1.6 ± 0.5	1.8 ± 0.3	13.2 ± 0.2
3.L2	T72	I-E^k	3	8.7 ± 0.1	0.1 ± 0.1	1.2 ± 0.3	2.6 ± 0.6	12.9 ± 0.3
3.L2	A72	I-E^k	2	9.3 ± 0.2	0.0 ± 0.3	0.2 ± 0.0	4.7 ± 0.5	12.3 ± 0.1
3.L2	I72	I-E^k	1	9.1 ± 0.1	0.0 ± 0.3	0.1 ± 0.0	3.9 ± 0.6	12.4 ± 0.2
E8	TPI	HLA-DR1	2	8.6 ± 0.1	4.1 ± 0.4	0.7 ± 0.3	78.3 ± 2.7	9.5 ± 0.1
E8▯	TPI	HLA-DR1	3	8.5 ± 0.1	4.0 ± 0.5	1.1 ± 0.3	41.4 ± 9.4	10.1 ± 0.2
E8*	TPI	HLA-DR1	4	8.8 ± 0.2	1.0 ± 0.3	1.2 ± 0.4	131.7 ± 40.5	8.9 ± 0.3
2B4 (NP)	K5	I-E^k	6	7.9 ± 0.3	7.3 ± 1.3	2.0 ± 0.2	4.5 ± 2.4	12.3 ± 0.4
2B4 (WT)	K5	I-E^k	4	8.8 ± 0.3	2.3 ± 0.7	1.8 ± 0.4	64.0 ± 9.1	9.7 ± 0.1
2B4 (GK)	K5	I-E^k	3	8.9 ± 0.2	1.9 ± 1.2	1.4 ± 0.5	40.1 ± 9.8	10.1 ± 0.2
2B4 (HN)	K5	I-E^k	2	8.9 ± 0.2	0.3 ± 0.1	0.5 ± 0.2	6.1 ± 0.8	12.0 ± 0.1
2B4 (NH)	K5	I-E^k	0	0.0 ± 0.0	0.0 ± 0.0	0.3 ± 0.1	2.6 ± 1.1	12.9 ± 0.1

▯The recombinant TCR coated to the bead was tested against pMHC on the BFP bead.

*The recombinant TCR coated to the BFP bead was tested against pMHC expressed on a THP-1 cell.

Note that the best-fit parameters were chosen using the procedure shown in Supplementary Table 5 or determined as the parameter set with the $\delta_z(n^*)$ value closest to δ_0^* as well as the smallest RSS. All errors are standard error of mean after optimizing the best-fit parameters. All values were rounded to one decimal place.

Supplementary References

1. Smith, S. B., Finzi, L. & Bustamante, C. Direct mechanical measurements of the elasticity of single DNA molecules by using magnetic beads. *Science* **258**, 1122-1126, (1992).

2. Guo, S., Tang, Q., Yao, M., You, H., Le, S., Chen, H. & Yan, J. Structural-elastic determination of the force-dependent transition rate of biomolecules. *Chem Sci* **9**, 5871-5882, (2018).

3. Rossjohn, J., Gras, S., Miles, J. J., Turner, S. J., Godfrey, D. I. & McCluskey, J. T cell antigen receptor recognition of antigen-presenting molecules. *Annual review of immunology* **33**, 169-200, (2015).

4. Wu, P., Zhang, T., Liu, B., Fei, P., Cui, L., Qin, R., Zhu, H., Yao, D., Martinez, R. J., Hu, W., An, C., Zhang, Y., Liu, J., Shi, J., Fan, J., Yin, W., Sun, J., Zhou, C., Zeng, X., Xu, C., Wang, J., Evavold, B. D., Zhu, C., Chen, W. & Lou, J. Mechano-regulation of Peptide-MHC Class I Conformations Determines TCR Antigen Recognition. *Mol Cell* **73**, 1015-1027 e1017, (2019).

5. Bouchiat, C., Wang, M. D., Allemand, J., Strick, T., Block, S. M. & Croquette, V. Estimating the persistence length of a worm-like chain molecule from force-extension measurements. *Biophysical journal* **76**, 409-413, (1999).

6. Gebhardt, J. C., Bornschlogl, T. & Rief, M. Full distance-resolved folding energy landscape of one single protein molecule. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 2013-2018, (2010).

7. Seol, Y., Li, J., Nelson, P. C., Perkins, T. T. & Betterton, M. D. Elasticity of short DNA molecules: theory and experiment for contour lengths of 0.6-7 microm. *Biophysical journal* **93**, 4360-4373, (2007).

8. Choi, H. K., Min, D., Kang, H., Shon, M. J., Rah, S. H., Kim, H. C., Jeong, H., Choi, H. J., Bowie, J. U. & Yoon, T. Y. Watching helical membrane proteins fold reveals a common N-to-C-terminal folding pathway. *Science* **366**, 1150-1156, (2019).

9. Schuler, B., Lipman, E. A. & Eaton, W. A. Probing the free-energy surface for protein folding with single-molecule fluorescence spectroscopy. *Nature* **419**, 743-747, (2002).

10. Yang, W. Y. & Gruebele, M. Folding at the speed limit. *Nature* **423**, 193-197, (2003).

11. Kubelka, J., Hofrichter, J. & Eaton, W. A. The protein folding 'speed limit'. *Curr Opin Struct Biol* **14**, 76-88, (2004).

12. Chung, H. S., Louis, J. M. & Eaton, W. A. Experimental determination of upper bound for transition path times in protein folding from single-molecule photon-by-photon trajectories. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 11837-11844, (2009).

13. Guo, S., Efremov, A. K. & Yan, J. Understanding the catch-bond kinetics of biomolecules on a one-dimensional energy landscape. *Communications Chemistry* **2**, 30, (2019).

14. Lapidus, L. J., Steinbach, P. J., Eaton, W. A., Szabo, A. & Hofrichter, J. Effects of Chain Stiffness on the Dynamics of Loop Formation in Polypeptides. Appendix: Testing a 1-Dimensional Diffusion Model for Peptide Dynamics. *J. Phys. Chem. B* **106**, 11628-11640, (2002).

15. Liu, B., Chen, W., Evavold, B. D. & Zhu, C. Accumulation of dynamic catch bonds between TCR and agonist peptide-MHC triggers T cell signaling. *Cell* **157**, 357-368, (2014).

16. Das, D. K., Feng, Y., Mallis, R. J., Li, X., Keskin, D. B., Hussey, R. E., Brady, S. K., Wang, J. H., Wagner, G., Reinherz, E. L. & Lang, M. J. Force-dependent transition in the T-cell receptor beta-subunit allosterically regulates peptide discrimination and pMHC bond lifetime. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 1517-1522, (2015).

17. Hong, J., Ge, C., Jothikumar, P., Yuan, Z., Liu, B., Bai, K., Li, K., Rittase, W., Shinzawa, M., Zhang, Y., Palin, A., Love, P., Yu, X., Salaita, K., Evavold, B. D., Singer, A. & Zhu, C. A TCR

mechanotransduction signaling loop induces negative selection in the thymus. *Nat Immunol* **19**, 1379-1390, (2018).

Zareie, P., Szeto, C., Farenc, C., Gunasinghe, S. D., Kolawole, E. M., Nguyen, A., Blyth, C., Sng, X. Y. X., Li, J., Jones, C. M., Fulcher, A. J., Jacobs, J. R., Wei, Q., Wojciech, L., Petersen, J., Gascoigne, N. R. J., Evavold, B. D., Gaus, K., Gras, S., Rossjohn, J. & La Gruta, N. L. Canonical T cell receptor docking on peptide-MHC is essential for T cell signaling. *Science* **372**, (2021).

Kersh, G. J. & Allen, P. M. Structural basis for T cell recognition of altered peptide ligands: a single T cell receptor can productively recognize a large continuum of related ligands. *J Exp Med* **184**, 1259-1268, (1996).

Natarajan, A., Nadarajah, V., Felsovalyi, K., Wang, W., Jeyachandran, V. R., Wasson, R. A., Cardozo, T., Bracken, C. & Krogsgaard, M. Structural Model of the Extracellular Assembly of the TCR-CD3 Complex. *Cell Rep* **14**, 2833-2845, (2016).

Hong, J., Persaud, S. P., Horvath, S., Allen, P. M., Evavold, B. D. & Zhu, C. Force-regulated In situ TCR-peptide-bound MHC class II kinetics determine functions of CD4+ T cells. *Journal of immunology* **195**, 3557-3564, (2015).

Rushdi, M. N., Pan, V., Li, K., Travaglino, S., Choi, H.-K., Hong, J., Griffitts, F., Agnihotri, P., Mariuzza, R. A., Ke, Y. & Zhu, C. Cooperative binding of TCR and CD4 to pMHC enhances TCR sensitivity. *Nature Communications*, in press.

Yuan, Z., Cong, P., Natarajan, Ge, C., Travaglino, S., Beesam, S., Grazette, D., Krogsgaard, M. & Zhu, C. Cooperative ectodomain interaction among TCRαβ, CD3γε, and CD3δε enhances TCR mechanotransduction. *bioRxiv*, https://www.biorxiv.org/content/10.1101/2022.1104.1114.488403v488401, (2022).

Liu, B., Chen, W., Natarajan, K., Li, Z., Margulies, D. H. & Zhu, C. The cellular environment regulates in situ kinetics of T-cell receptor interaction with peptide major histocompatibility complex. *European journal of immunology* **45**, 2099-2110, (2015).