Increased early sodium current causes familial atrial fibrillation and dampens effect of flecainide

O'Reilly M[1*], Sommerfeld LC[1,2,3*], O'Shea C[1], Broadway-Stringer S[1], Andaleeb S[1], Reyat JS[1], Kabir SN[1], Malinova A[1], Delbue D[2,3], Fortmueller L[2,3], Gehmlich K[1,5], Pavlovic D[1], Skryabin BV[6], Holmes AP[1,7], Kirchhof P[1,3,4], Fabritz L[1,2,3,4]

[1]*Institute of Cardiovascular Sciences, University of Birmingham, UK*

[2]*University Center of Cardiovascular Science, University Heart and Vascular Center Hamburg, UKE Hamburg, Germany*

[3]*DZHK Standort Hamburg/Kiel/Luebeck, Germany*

[4]*Department of Cardiology, University Heart and Vascular Center UKE Hamburg, Germany*

[5]*Division of Cardiovascular Medicine, Radcliffe Department of Medicine and British Heart Foundation Centre of Research Excellence Oxford, University of Oxford, UK*

[6]*Medical Faculty, Core Facility Transgenic animal and genetic engineering Models (TRAM), University of Muenster, Germany*

[7]*Institute of Clinical Sciences, University of Birmingham, UK*

*equal contribution

Corresponding author:

Professor Larissa Fabritz

University Center of Cardiovascular Science

University Heart and Vascular Center, UKE Hamburg

Hamburg, Germany

E-Mail: l.fabritz@uke.de

Abstract

Atrial fibrillation (AF) is the most common cardiac arrhythmia. Pathogenic variants in genes encoding ion channels are associated with familial AF. The point mutation M1875T in the *SCN5A* gene, which encodes the α-subunit of the cardiac sodium channel $Na_v1.5$, has been associated with increased atrial excitability and familial AF. Designing a new murine model carrying the *Scn5a*-M1875T mutation enabled us to study the effects of the $Na_v1.5$ mutation in detail.

Atrial cardiomyocytes from newly generated *Scn5a*-M1875T$^{+/-}$ mice showed a selective increase in the early (peak) cardiac sodium current, measured by patch clamp. Atrial microelectrode recordings revealed larger action potential amplitudes and a faster peak upstroke velocity. Conduction was studied using optical mapping. Conduction slowing by the sodium channel blocker flecainide was less pronounced in *Scn5a*-M1875T$^{+/-}$ compared to wildtype atria. *In vivo* analysis using electrocardiograms and echocardiography, as well as cardiac histology, excluded overt hypertrophy or heart failure in young adult *Scn5a*-M1875T$^{+/-}$ mice.

The *Scn5a*-M1875T point mutation causes a cardiac sodium channel gain-of-function. Our results suggest increased atrial peak sodium current as a potential trigger for increased atrial excitability and atrial fibrillation.

Statement of Significance

The observed changes induced by this point mutation suggest a selective increase in peak sodium current as a cause of familial AF. Our findings provide a possible explanation for the variable effectiveness of sodium channel blockers in patients with AF. Carriers of such sodium channel gain-of-function mutations may benefit more from tailored treatments.

1 Introduction

Atrial fibrillation (AF), the most common cardiac arrhythmia, is characterised by episodes of irregular and uncoordinated atrial electrical activity. It is associated with ischaemic stroke, cardiovascular death and frequent hospitalisations [1]. A range of different common factors, such as heart failure, diabetes and increased formation of fibrosis, can damage the atria, contributing to AF [2, 3]. These factors interact with a pre-existent, potentially inherited substrate to result in AF. Inherited forms are characterised by early onset of the condition. Pathogenic variants in several genes associated with cardiomyopathies have been identified in familial AF, including variants in sarcomeric and cell-cell contact genes, and others [4-6]. Within the group of ion channel genes, variants leading to dysfunction of the cardiac sodium channel are associated with familial AF [4, 7, 8], both via variants in the genes coding for the channel or through upstream mechanisms altering the expression of sodium channel genes [9, 10].

The cardiac voltage-gated sodium channel (Na$_v$1.5) facilitates movement of sodium (Na$^+$) ions across the cardiomyocyte membrane and hence elicits the cardiac Na$^+$ current (I$_{Na}$). I$_{Na}$ is vital for electrical excitation preceding mechanical contraction of the myocardium, as transient influx of Na$^+$ triggers the fast upstroke phase of the cardiac action potential and thus depolarisation of the cardiomyocyte membrane. The Na$_v$1.5 channel is composed of both α- and ß- subunits. The pore-forming α-subunit is encoded by the *SCN5A* gene. Variants in *SCN5A* have been linked to cardiac conditions, including AF [11-13], Brugada syndrome [14], conduction disease [15-17] and dilated cardiomyopathy [18, 19]. *SCN5A* genetic variants reported [20] show various underlying mechanisms mainly linked to channel dysfunction, defective channel trafficking or protein complex formation [17, 21, 22].

A missense *SCN5A* point mutation, Met1875Thr (M1875T), located in the C-terminus of the channel protein, was linked to autosomal dominant familial AF that spanned three generations of a family [23]. Atrial ectopy was evident in mutation carriers in adolescence, and persistent AF occurred as early as 27 years of age. Analyses in the human cell line HEK293 heterologous expression system [23] suggested an enhanced function of the mutated Na$_v$1.5 channel.

To investigate the impact of the M1875T mutation in a more physiological setting, we generated and characterised a novel knock-in murine model (*Scn5a*-M1875T$^{+/-}$). Mice that were heterozygous for the M1875T mutation were viable and studied herein. We investigated these mice from the whole organ *in vivo* to the level of the single cell *ex vivo*.

2 Materials and Methods

2.1   Generation and sequencing of the *Scn5a*-M1875T murine model

Mice heterozygous for the knock-in mutation M1875T in the *Scn5a* gene (*Scn5a*-M1875T[+/-]) were generated by T-C point-mutating exon 28 of the cardiac sodium channel *SCN5A* gene using pSCN5a_targ3 targeting vector and CRISPR/Cas9 system in murine embryonic stem (ES) cells (Figure 1a). Mutation-harboring ES cells were characterised using Southern blot analysis and sequenced in order to exclude genomic rearrangements (Figure 1b) [24, 25], and injected into B6D2F1 mouse blastocysts. Sequencing analysis of the mutation-containing region using DNA from adult wildtype (WT) and heterozygous *Scn5a*-M1875T[+/-] mice on C57Bl/6J x 129sv hybrid genetic background are shown in Figure 1c. Detailed steps of the generation are further explained in the supplement.

Methionine at position 1875 of the human $Na_v1.5$ protein sequence corresponds to position 1877 of the murine sequence. The latter is therefore point-mutated in this model. However, to underpin the bedside-to-bench nature of this investigation, we use the human annotation and refer to it as "Scn5a-M187**5**T" throughout this manuscript.

Mice were bred on an FVB or 129/sv genetic background and housed in individually ventilated cages with sex-matched littermates (2-5 mice/cage), under standard conditions: 12 hours light/dark circle, 22°C and 55% humidity. Food and water were available *ad libitum*. The health status of mice used in the study was monitored daily and prior to experiments.

Functional experiments were conducted on hearts of male and female young adult mice of pure background (8-20 weeks), heterozygous for the knock-in mutation M1875T in the *Scn5a* gene *(Scn5a*-M1875T[+/-]) and their WT littermates. Mice of both sexes and background were used evenly and analysed jointly, to make the data more widely applicable. Some of our investigations focus on the left atrium. This is mostly due to practical reasons, e.g. less spontaneous activity in the left compared to the right atrium and therefore the possibility to pace at a wider range of cycle lengths or better reproducibility of the LA in the parasternal view.

Figure 1 Targeting of exon 28 of the mouse *Scn5a* gene in order to introduce the M1875T coding mutation, and sequencing confirmation of its presence

a) The intronic and intergenic regions are shown as lines, exons are shown as filled boxes. The empty box corresponds to the neomycin resistance cassette (neo) flanked by the LoxP sites (vertical arrows). Exon numeration is shown above. The arrows below are corresponding to BamHI restriction endonuclease sites (B). The black box corresponds to Southern probe sequences (So3). The expected sizes of restriction DNA fragments are indicated below in kb. **ai)** Wildtype (WT) locus. **aii)** Targeted vector structure (without negative selection marker and plasmid backbone). **aiii)** Genomic locus after the homologous recombination. The neomycin cassette is present in intron 27 and flanked by two LoxP sites. **aiv)** Genomic locus after the CRE-mediated neo cassette deletion. **b)** Southern blot analysis of DNA isolated from mouse tail biopsy of the F1 offspring (1-13) and hybridized with the So3 probe. With help of the BamHI enzymatic digestion, we detect the WT allele 7.6 kb and targeted allele 5.8 kb. DNA samples 1, 2, 4, 5, 7, 8, 10, 11, and 13 contain correctly targeted *Scn5a* gene (Scn5a-M1875T$^{+/-}$). Positions of the size marker (in bp) are shown on the right. The WT control animal is labeled "wt". **c)** Sequencing analysis of the *Scn5a* gene region containing the mutation site in back-crossed adult mice (on pure genetic background). The T-C mutation on one allele causing the methionine-threonine exchange at position 1875 (1877) is indicated by an arrow.

 2.2   Study approval

 All procedures were performed in compliance with the guidelines from Directive 2010/63/EU of the

 European Parliament on the protection of animals used for scientific purposes and conducted in

 accordance with rules and regulations for experiments with animals and approved by the UK Home

 Office (PPL number 30/2967) and by the institutional review board of University of Birmingham.

2.3 Murine ECG recordings *in vivo*

Non-invasive electrocardiograms (ECG) were recorded in conscious young adult mice (8-19 weeks) using a tunnel system for gentle restraint (ecgTunnel, EMKA Technologies, Paris, France) [26]. ECG recordings were analysed using ECGauto software (EMKA Technologies, Paris, France). ECGs were also recorded in isoflurane-sedated mice during echocardiography as below.

2.4 Murine echocardiography *in vivo*

Echocardiography was performed in sedated mice (0.5-2% isoflurane, supplemented with 100% O_2) using Vevo® 2100 system (VisualSonics, Amsterdam, Netherlands) as reported previously [27]. Heart rate was maintained at 450 ± 70 bpm. Left atria (LA) were visualised in the parasternal long axis view in the plane of the aortic root. LA area and diameter were measured during pre-atrial contraction, using the P-wave of the limb ECG trace as a guide. Left ventricular parameters were also measured in the parasternal long axis view.

2.5 Atrial murine cardiomyocyte isolation

Murine hearts were excised under deep terminal anaesthesia (4% isoflurane inhalation in O_2, 1.5 L/min) and perfused at 4 mL.min^{-1} at 37°C on a vertical Langendorff apparatus with the following solutions, equilibrated with 100% O_2: (i) HEPES-buffered, Ca^{2+}-free, modified Tyrode's solution containing in mM: NaCl 145, KCl 5.4, $MgSO_4$ 0.83, Na_2HPO_4 0.33, HEPES 5, and glucose 11 (pH 7.4, NaOH) x 5 min; (ii) Tyrode's enzyme solution containing 640 µg/mL collagenase type II (270 U/mg), 600 µg/ml collagenase type IV (270 U/mg) and 50 µg/ml protease (Worthington, Lakewood, NJ), 20 mM taurine and 3 µM $CaCl_2$ × 8-12 min. The heart was removed from the Langendorff setup and perfused with 5 mL of modified Kraftbruhe (KB) solution containing in mM: DL-potassium aspartate 10, L-potassium glutamate 100, KCl 25, KH_2PO_4 10, $MgSO_4$ 2, taurine 20, creatine 5, EGTA 0.5, HEPES 5, 0.1% BSA, and glucose 20 (pH 7.2, KOH).

The LA was dissected free and cardiomyocytes were dissociated gently with fire-polished glass pipettes (2 to 1 mm diameter in sequence). Cells were re-suspended in 2 mL KB buffer and Ca^{2+} was gradually reintroduced to the cell suspension incrementally over a period of 2 hours to reach a final concentration of 1 mM. All experiments were performed within 8 hours of isolation.

2.6 Whole-cell patch clamp electrophysiology of isolated atrial cardiomyocytes

Dissociated murine LA cardiomyocytes were plated on, and allowed to adhere to, laminin-coated coverslips (10 mm diameter) for at least 20 minutes. Coverslips were transferred to a recording chamber and were continually superfused at 3 mL.min^{-1}, with a low Na$^+$ external solution containing

in mM; NaCl 10, KCl 4.5, $C_5H_{14}ClNO$ 130, $CaCl_2$ 1, $MgCl_2$ 1.2, HEPES 10 and glucose 10 (pH 7.4 with CsOH). To block L-type Ca^{2+} currents, 2 mM $NiCl_2$ was added to the superfusate. Experiments were performed at 22 ± 0.5°C. Whole-cell patch clamp recordings were obtained in voltage-clamp mode using borosilicate glass pipettes (tip resistances 1.5-3 MΩ) [28].

For Na^+ current recordings, the pipette solution contained in mM: CsCl 115, NaCl 5, EGTA 10, HEPES 10, MgATP 5, TEACl 20 and $MgCl_2$ 0.5 (pH 7.2, KOH). Voltage-dependent Na^+ currents were evoked by 5 mV step depolarisations (100 ms) from a holding potential of -100 mV to test potentials ranging from -95 mV to +40 mV. Cells were excluded from analysis if there was no reversal of the sodium current by +40mV. To investigate $Na_v1.5$ voltage-dependent inactivation kinetics, cells were subject to 500 ms pre-pulses ranging from -120 mV to -40 mV, followed by a 100 ms step to -30 mV. For $Na_v1.5$ time-dependent recovery kinetics, a standard two pulse protocol was used (-120 mV to -30 mV, 20 ms), with the time between the two pulses incrementally varying between 5 and 950 ms.

All recordings and analysis protocols were performed using an Axopatch 200B amplifier (Molecular Devices, USA) and digitized at 50 kHz using a CED micro1401 driven by Signal v6 software (Cambridge Electronic Design, Cambridge, UK). Series resistance was compensated, ranging between 60-100% for all cells. Experiments were terminated if series resistance abruptly changed or was above 10 MΩ.

2.7 Atrial murine microelectrode recordings

As previously described [28-30], following isolation the LA was immediately transferred into a dissecting chamber and continuously superfused at 10 mL.min^{-1} with a bicarbonate buffered Krebs-Henseleit (KH) solution containing in mM: NaCl 118; $NaHCO_3$ 24.88; KH_2PO_4 1.18; Glucose 11; $MgSO_4$ 0.83; $CaCl_2$ 1.8; KCl 3.52, equilibrated with 95% O_2/5% CO_2, 36-37°C, pH 7.4. Micro-dissection and pinning out of the LA was performed using a dissection microscope (Stemi SV 11, Zeiss, Germany). The LA was paced at 1–10 Hz via bipolar platinum electrodes. Action potentials (APs) were recorded from freely contracting LA using custom made glass floating microelectrodes containing 3 M KCl, (resistance 15-30 MΩ). Voltage signals were amplified and digitised at 20 kHz and were unfiltered (Axoclamp 2B; Molecular Devices, California, USA; Spike2 software Cambridge Electronic Design, Cambridge, UK). Measured parameters included the resting membrane potential (RMP), action potential amplitude (APA), peak depolarisation rate (dV/dt) and action potential duration (APD) at 30-90% repolarisation. APs were only analysed following sufficient rate adaptation achieved after at least 50 stimulated APs at each frequency.

2.8   Optical mapping of murine atria

Optical mapping of the LA was conducted as previously described [31, 32]. Isolated whole hearts were

loaded on to a vertical Langendorff apparatus and perfused with a standard KH solution. Hearts were

perfused at 4 mL.min^{-1} (equilibrated with 95%O_2/5%CO_2 and heated to 36-37°C, pH 7.4). Hearts were

loaded with 25 µL of voltage sensitive dye Di-4-ANEPPS at a concentration of 5 mg/mL, diluted in 1 mL

of KH solution and delivered via bolus port injection over 3-5 minutes. The LA was then isolated and

pinned in a superfusion chamber containing 37°C KH solution for transfer to the optical mapping

setup, anterior surface facing up.

In the optical mapping system, atria were superfused with KH solution (95%O_2/5%CO_2, 36-37°C)

containing contraction uncoupler Blebbistatin (35 µM). For imaging, atria were illuminated by two

dual LEDs at 530 nm. A 630 nm long-pass filter was used to sperate emitted fluorescence, imaged

using an ORCA flash 4.0 CMOS camera (Hamamatsu, Japan). Images were acquired at a framerate

0.987 kHz and pixel size of 71 µm/pixel^2. Atria were paced using bipolar platinum electrodes delivering

2 ms pulses at twice diastolic threshold (minimum voltage required to elicit APs).

One-minute baseline recording was taken following a 10 minutes equilibration period to ensure

contraction uncoupling and temperature re-stabilisation. During imaging, atria were initially paced at

330 ms pacing cycle length (PCL). A 'ramp' pacing protocol was then initiated, in which the atria were

paced at 120 ms PCL for 100 stimuli and then PCL was reduced from 120 ms to 80 ms in 10 ms intervals

every 20 stimuli. After taking baseline recordings, LED illumination was switched off and the

superfusion solution replaced with an identical solution containing flecainide at a concentration of

1 µM and then 5 µM (or control solution without flecainide for time control experiments). Subsequent

recordings were then made as described above after 20 minutes superfusion with 1 µM flecainide

solution, and then further 15 minutes with 5 µM flecainide solution. Atria were paced at 330 ms PCL

continuously in dark conditions between recordings.

From these recordings, APD and conduction velocity (CV) were mapped across the LA using

ElectroMap software [32]. Atria were removed from analysis at a given PCL if loss of 1:1 capture ratio

with pacing stimuli (i.e. missed beats) was observed.

2.9   Statistics

For all murine experiments presented herein, experimenters were blinded to the genotype of the

littermate pairs during data collection and analysis. Student t-tests were used for singular comparisons

for normally distributed data. A hierarchical nested t-test (Figure 2) and Mann-Whitney test (Table S3)

were used where appropriate. Multiple comparisons were made using 2-way ANOVA with

Bonferroni's post-hoc tests. For current-voltage graphs (Figure 2b), a Boltzmann curve was fit to the

data, using the modified Boltzmann equation: $I_{Na} = G_{max}(V_m - V_{rev})/(1+Exp[(V_{0.5}-V_m)/k])$, where I_{Na} is the

current density at an equivalent test potential (V_m), G_{max} is the peak conductance (nS), V_{rev} is the reverse

potential, $V_{0.5}$ is the membrane potential at 50% current activation, and k is the slope constant [33]. All

graphical representations display individual measurements. Means are quoted and shown in Figures

± SEM. Level of statistical significance is shown in Figures as follows: *p<0.05; **p<0.01; ***p<0.001;

****p<0.0001. Statistics and Figures were created using Prism 8 (GraphPad Software, San Diego,

California).

3 Results

3.1 Viable heterozygous *Scn5a*-M1875T[+/-] mice show normal cardiac size and basic function

The point mutation previously identified in patients with early familial AF was successfully introduced to exon 28 of the mouse *Scn5a* gene via homologous recombination of a targeting vector. The vector contained the T-C point mutation (CRISPR/cas9-mediated) resulting in methionine-threonine exchange in the $Na_v1.5$ protein (Figure 1). Offspring from both WT x heterozygote (=*Scn5a*-M1875T[+/-]) and heterozygote x heterozygote pairings were viable. No homozygous *Scn5a*-M1875T[-/-] offspring was born (Figure S1b), suggesting embryonic lethality, as previously reported for other *Scn5a* mutations [34, 35]. Accordingly, the ratio of WT and heterozygous animals shifted from 1:2 (expected) to approximately 1:3 when heterozygous animals were crossed (Figure S1b). The ratio of male:female sex in offspring approximated 1:1 as expected (Figure S1c).

Age-matched young adult WT and *Scn5a*-M1875T[+/-] mice displayed similar heart rate, PR-, QRS- and QT- interval regardless of genotype in electrocardiograms recorded awake (Table S1) and during sedation (Table S2). Echocardiography and histological examination excluded overt differences in atrial or ventricular structure (Table S3, Figure S2a). Accordingly, pro atrial natriuretic peptide protein expression (proANP) was detected in right atria as expected but was not elevated in ventricles (Figure S2b).

3.2 Atrial *Scn5a*-M1875T[+/-] cardiomyocytes have an augmented peak sodium current density

To determine the impact of the M1875T point mutation on peak I_{Na} amplitude and $Na_v1.5$ channel gating properties, left atrial cardiomyocytes from WT and *Scn5a*-M1875T[+/-] mice (8-13 weeks) were isolated and whole-cell patch clamp recordings were performed.

The M1875T variant increased I_{Na} over test potentials ranging from –100 to +40 mV (Figure 2a and b). At a peak test potential of -30 mV, mean left atrial cardiomyocyte I_{Na} density was higher in *Scn5a*-M1875T[+/-] (-28.0 ± 1.1 pA/pF, n=107 cells) than in WT littermates (-22.8 ± 0.7 pA/pF, n=109 cells, p<0.0001, Figure 2ci). The elevation in I_{Na} was also apparent when recordings were grouped by animal (*Scn5a*-M1875T[+/-] -29.1 ± 2.2 pA/pF, N=14 vs WT -22.3 ± 0.9 pA/pF, N=15, p=0.0075, Figure 2cii) and after applying hierarchical analysis (*Scn5a*-M1875T[+/-] -29.1 ± 1.1 pA/pF, n=107 cells, N=14 mice vs WT -22.3 ± 0.9 pA/pF, n=109 cells, N=15 mice, p=0.0093).

Capacitance measurements were not different between genotypes, indicative of similar cell size (Figure S3a).

219   Whole-cell I_{Na} voltage-dependent inactivation and time-dependent recovery kinetics were not altered

220   in *Scn5a*-M1875T$^{+/-}$ cardiomyocytes compared to WT (Figure S3b and c). Na_v1.5 expression in hearts

221   of *Scn5a*-M1875T$^{+/-}$ mice at mRNA and the protein level revealed no difference at either the whole cell

222   or isolated membrane fraction level (Figure S4).

Figure 2 Isolated atrial cardiomyocytes with the *Scn5a*-M1875T$^{+/-}$ mutation have a larger sodium current than wildtypes when measured with whole-cell patch clamp electrophysiology
a) Representative sodium current (I_{Na}) traces from whole-cell voltage clamp recording of isolated left atrial (LA) cardiomyocytes from wildtype (WT) (black) and *Scn5a*-M1875T$^{+/-}$ (blue) hearts. **b)** Normalised grouped data revealed that the *Scn5a*-M1875T$^{+/-}$ mutation increases peak I_{Na} in LA cardiomyocytes over test potentials ranging from −100 to +40 mV. **c)** At the peak I_{Na} test potential of -30 mV, *Scn5a*-M1875T$^{+/-}$ cardiomyocytes had a significantly larger I_{Na} than WTs, both when comparing individual cells (n=109 WT, n=107 *Scn5a*-M1875T$^{+/-}$) **(ci)** and hearts (N=15 WT, N=14 *Scn5a*-M1875T$^{+/-}$) **(cii)**. Data are presented as the mean ± SEM.

3.3   Action potentials from *Scn5a*-M1875T$^{+/-}$ atria have a larger amplitude and faster peak

upstroke velocity

Action potentials (APs) were measured in whole left atrial tissue isolated from WT and *Scn5a*-

M1875T$^{+/-}$ mice (9-13 weeks) using sharp microelectrodes. Representative AP traces are shown in

Figure 3 during stimulation at 100 ms (Figure 3c) and 1000 ms pacing cycle length (PCL) (Figure 3f).

Action potential amplitude was significantly larger in *Scn5a*-M1875T$^{+/-}$ murine left atria at all PCLs

tested and this effect was more pronounced at shorter cycle lengths (N=8, n=24-25, Figure 3a). The

variant resulted in a faster peak upstroke velocity (dV/dt), especially at the shorter cycle lengths

(100 ms PCL: WT 128.0 ± 3.3, n=24; *Scn5a*-M1875T$^{+/-}$ 142.8 ± 4.0 mV/ms, n=25, p=0.0282, Figure 3b).

Figure 3 Action potentials from atria with the *Scn5a*-M1875T$^{+/-}$ mutation have a larger action potential amplitude and a faster peak upstroke velocity when measured with the sharp microelectrode technique

a) *Scn5a*-M1875T$^{+/-}$ left atria (LA) had a significantly larger action potential (AP) amplitude at shorter pacing cycle lengths (PCLs) of 80-120 ms (p<0.05) and **b)** a significantly faster peak upstroke velocity (dV/dt). **d-f)** When paced at 1000 ms PCL, *Scn5a*-M1875T$^{+/-}$ LA had a significantly shorter AP duration (APD) when measured at 50% (APD50) **(d)** and 90% (APD90) **(e)** repolarisation. *p<0.05, **p<0.01, wildtype (WT) vs *Scn5a*-M1875T$^{+/-}$, N=8 per group, n=24 WT, n=25 *Scn5a*-M1875T$^{+/-}$; ANOVA statistics. Also shown are representative AP traces from wildtype (black) and *Scn5a*-M1875T$^{+/-}$ (blue) LA when stimulated at 100 ms **(c)** and 1000 ms PCL **(f)**. Data are presented as the mean ± SEM.

The resting membrane potential [36] was not different between genotypes (100 ms PCL: WT -72.4 ± 0.6;

Scn5a-M1875T$^{+/-}$ -73.1 ± 0.6 mV). Atrial activation times were also similar (100 ms PCL: WT 4.9 ± 0.2;

Scn5a-M1875T+/- 4.9 ± 0.2 ms) (Table S4). Only at the long PCL of 1000 ms, the AP duration (APD) at

50 and 90% repolarisation was shorter in *Scn5a*-M1875T$^{+/-}$ left atria than in WT (Figure 3d and e), while

the APD at 30 and 70% repolarisation was not significantly different (Table S4). Similarly, optical

mapping data showed no APD differences at PCLs tested.

3.4 Flecainide-induced atrial conduction slowing and post-repolarisation refractoriness is less pronounced in *Scn5a*-M1875T$^{+/-}$ atria

Optical mapping of WT and *Scn5a*-M1875T$^{+/-}$ whole left atrial tissue was performed to test effects of

the heterozygous *Scn5a*-M1875T mutation on atrial conduction.

Left atria were superfused with the open channel sodium channel blocker flecainide (1 µM, clinically

used concentration) to determine the response of *Scn5a*-M1875T$^{+/-}$ and WT left atria. While

conduction velocity was unchanged at baseline, flecainide slowed conduction less in *Scn5a*-M1875T$^{+/-}$

left atria (conduction velocity difference at 100 ms PCL -6 ± 1 cm/s, n=12, p=0.0357) than in WTs
(conduction velocity difference at 100 ms PCL -10 ± 1 cm/s, n=12, p=0.0357, Figure 4a and b). We also
investigated flecainide-induced changes in left atrial refractoriness as it is known that flecainide
induces post-repolarisation refractoriness using rapid atrial pacing. Representative optical AP
recordings show 1:1 capture in the *Scn5a*-M1875T[+/-] left atria with flecainide, while several stimuli in
the WT left atria did not elicit APs (Figure 4c). All time-controlled left atria (no flecainide, same
experimental duration) were successfully paced with 1:1 capture down to 80 ms PCL. Loss of 1:1
capture began at longer PCLs in WT (7/12 atria lost 1:1 capture at PCLs ≤ 120 ms) compared to *Scn5a*-
M1875T[+/-] (1/12 atria lost 1:1 capture at PCLs ≤ 120 ms, p=0.0185) left atria. The diastolic pacing
threshold remained consistent throughout experiments. Thus, flecainide induced less pronounced
post-repolarisation refractoriness in the *Scn5a*-M1875T[+/-] left atria.

Figure 4 The response to the anti-arrhythmic agent flecainide is reduced in *Scn5a*-M1875T+/- atria in optical mapping

a) Example left atrial (LA) activation maps from wildtype (WT, top panels) and *Scn5a*-M1875T+/- (bottom panels) mice. Left panels show activation maps at baseline, right panels show activation of the same atria following exposure to 1 µM flecainide for 20 minutes. **b)** Grouped data showing change in conduction velocity (ΔCV) following exposure to 1 µM flecainide in WT (black) and *Scn5a*-M1875T+/- (blue) LA for 20 mins. **c)** Example traces of optical action potentials (APs) recorded following further treatment of WT (top, black) and *Scn5a*-M1875T+/- (bottom, blue) LA with 5 µM flecainide for 15 mins. **d)** Grouped data showing minimum pacing cycle length (PCL) at which 1:1 stimulus capture was maintained in WT (black) and *Scn5a*-M1875T+/- (blue) LA following exposure to 5 µM flecainide. Time control data (grey) shows minimum PCL at which 1:1 stimulus capture was maintained in atria that were not exposed to flecainide but had been under experiment conditions for the same time period (35 mins from baseline recording). N=12 per group at 330 ms-100 ms PCL. Atria were excluded from further analysis at shorter PCLs if 1:1 capture was lost, only data in steady state was used. N=5 for time control experiments. *p<0.05, **p<0.01, WT vs *Scn5a*-M1875T+/-; ANOVA statistics. Data are represented as the mean ± SEM.

4 Discussion

4.1 Main findings

Our study describes the effects of the familial atrial fibrillation (AF) mutation *Scn5a*-M1875T[+/-] in a newly generated murine model. Key effects are that the M1875T *Scn5a* mutation leads to an increased atrial action potential upstroke velocity and amplitude and a selective increase in the early cardiac sodium current (I_{Na}) while atrial cardiomyocyte capacitance, and cardiac size and function are preserved. *Scn5a*-M1875T[+/-] atria exhibit a dampened effect to the sodium channel blocker flecainide. Our data in this new murine model suggest that a selective increase of I_{Na} can cause familial AF, and that commonly used concentrations of sodium channel blockers may be less effective in familial forms of AF with a selective increase in I_{Na} than in other types.

4.2 Gain-of-function properties of *Scn5a*-M1875T[+/-] sodium channels in the murine atrium

Our data show a gain-of-function variant, namely an increased early sodium current as evidenced by an augmented action potential amplitude and larger I_{Na} in the *Scn5a*-M1875T[+/-] variant.

In contrast to findings in HEK293 cells [23], there was no depolarising shift of $Na_v1.5$ channel inactivation in murine left atrial cardiomyocytes. Instead, we show a similarly augmented I_{Na} without alterations in channel gating properties. Generation of the murine mutant model allowed us to study mutated sodium channels in cardiac tissue within the presence of the greater protein complex including α- and β- subunits and other membrane proteins, a complete cardiomyocyte contractile apparatus and all other cardiac cell types. In contrast, HEK293 cells would not contract, exclusively contained the mutated version of $Na_v1.5$ in their membranes and are by necessity studied in isolation. Others have reported similar differences between HEK293 and cardiac model systems to the differences reported here [37].

Cardiac sodium channels have been observed to form dimers [38] and different regions of the channel protein have been implicated to play a part in dimerization. Structural analysis following the hypothesis of $Na_v1.5$ dimerization via C-terminal interaction reveals that the surface of residue Met1875 of one $Na_v1.5$ cytosolic C-terminus will interact with Ala1924 of a second $Na_v1.5$ [39]. This prediction suggests that the location site investigated in our model could provide a structural basis for altered $Na_v1.5$-$Na_v1.5$ channel interaction further to be investigated.

It is unlikely that the M1875T mutation increases the *late* sodium current ($I_{Na,l}$), as APs we measured were not prolonged by the mutation, in line with the initial report of the mutation in HEK cells [23]. A

lack of AP prolongation clearly differs to findings in the gain-of-function mutation *ΔKPQ-Scn5a*$^{+/-}$
mutant murine model which shows an increase in $I_{Na,I}$ [27, 28, 34] with prolonged atrial and ventricular AP
duration, especially at longer pacing cycle lengths. This suggests that the M1875T mutation acts
differently to, and is distinct from, *SCN5A* gain-of-function mutations leading to prolonged
repolarisation and long QT syndrome, in concordance with the broad spectrum of phenotypic
outcomes resulting from mutations in the same ion channel gene [40].
295
An enhanced sodium influx into cardiomyocytes should alter the sodium-calcium homeostasis [41, 42],
modifying intracellular sodium and calcium concentrations and signalling in cardiomyocyte
microdomains [43, 44]. This study found normal heart size and cardiomyocyte size. Nonetheless, chronic
sodium overload can result in structural changes in the heart with ageing. In the future, the M1875T
mouse model may be used to study the long-term effects of a selective increase in the early I_{Na}.

4.3   Effect of *Scn5a*-M1875T$^{+/-}$ on conduction and refractoriness
While structural alterations of the atria, contribute to atrial conduction disturbances [45, 46,47], defects in
the cardiac sodium channel can also cause conduction defects leading to AF [17-22]. Unlike other *SCN5A*
mutations [17-22], this variant shows a dampened response to flecainide both on conduction and post-
repolarisation refractoriness [48]. Flecainide is a clinically used sodium channel blocker that inhibits
cardiac I_{Na} via blocking the pores of open $Na_v1.5$ channels [49, 50]. The differential response to flecainide
is likely due to the increased early sodium influx through the mutated $Na_v1.5$ channels, enabling
preservation of conduction and activation properties in atrial cardiomyocytes when sodium channels
are inhibited, leading to an enhanced activation reserve.
310
311 Murine models have limitations due to differences between mice and men, but they have been useful
312 in characterising sodium channel mutations [27, 34, 35, 51]. Identified mutation-induced changes at the
313 cellular and organ level in this model appear sufficient to explain AF in the original family. Studies in
314 human cardiomyocytes [42] and in atrial engineered heart tissue [52, 53], would be desirable to assess the
315 effect of the M1875T mutation in human models in the future.
316

Conclusion

The *Scn5a*-M1875T$^{+/-}$ variant causes a selective increase in the early cardiac sodium current, potentially leading to an increased activation reserve and reduced refractoriness, whilst structure and contractile function are preserved. These findings can explain the familial occurrence of atrial ectopy and atrial fibrillation in the absence of reported severe heart disease. More widely they suggest a mechanism by which altered cardiomyocyte sodium currents can predispose to atrial fibrillation. Our data also show that the M1875T gain-of-function mutation decreases the effectiveness of sodium channel blockers such as flecainide, which may have implications for patient treatment.

Author Contribution Statement

LF and PK designed the research; BVS generated the mouse model in TRAM upon request by PK and LF;

MO, LCS, CO, SBS, SA, JSR, SNK, AM, DD, LFo, LF and APH conducted experiments, analysed data and performed statistical analysis;

KG, DP, APH, PK, LF supervised experiments and analysis.

MO, LCS, CO, SBS and LF wrote the manuscript together with all co-authors. All co-authors critically reviewed the manuscript.

Declaration of Interests

The authors have declared no direct conflict of interest in regards to the manuscript.

L.F. has received institutional research grants from governmental and charity funding agencies and several biomedical companies.

P.K. has received research support from several drug and device companies active in atrial fibrillation and has received honoraria from several such companies in the past.

L.F. and P.K. are listed as inventors on two patents held by University of Birmingham (Atrial Fibrillation Therapy WO 015140571, Markers for Atrial Fibrillation WO 2016012783).

Acknowledgments

We thank Hartwig Wieboldt, Daniel Stastny, Clara Apicella and Olivia Grech for technical assistance.

Supporting Citations

References (54-56) appear in the Supporting material.

References

1. Fabritz L, Crijns H, Guasch E, Goette A, Hausler KG, Kotecha D, Lewalter T, Meyer C, Potpara TS, Rienstra M, Schnabel RB, Willems S, Breithardt G, Camm AJ, Chan A, Chua W, de Melis M, Dimopoulou C, Dobrev D, Easter C, Eckardt L, Haase D, Hatem S, Healey JS, Heijman J, Hohnloser SH, Huebner T, Ilyas BS, Isaacs A, Kutschka I, Leclercq C, Lip GYH, Marinelli EA, Merino JL, Mont L, Nabauer M, Oldgren J, Purerfellner H, Ravens U, Savelieva I, Sinner MF, Sitch A, Smolnik R, Steffel J, Stein K, Stoll M, Svennberg E, Thomas D, Van Gelder IC, Vardar B, Wakili R, Wieloch M, Zeemering S, Ziegler PD, Heidbuchel H, Hindricks G, Schotten U, Kirchhof P. Dynamic risk assessment to improve quality of care in patients with atrial fibrillation: the 7th AFNET/EHRA Consensus Conference. *Europace* 2021;**23**:329-344.

2. Fabritz L, Guasch E, Antoniades C, Bardinet I, Benninger G, Betts TR, Brand E, Breithardt G, Bucklar-Suchankova G, Camm AJ, Cartlidge D, Casadei B, Chua WW, Crijns HJ, Deeks J, Hatem S, Hidden-Lucet F, Kaab S, Maniadakis N, Martin S, Mont L, Reinecke H, Sinner MF, Schotten U, Southwood T, Stoll M, Vardas P, Wakili R, West A, Ziegler A, Kirchhof P. Expert consensus document: Defining the major health modifiers causing atrial fibrillation: a roadmap to underpin personalized prevention and treatment. *Nat Rev Cardiol* 2016;**13**:230-237.

3. Benjamin EJ, Al-Khatib SM, Desvigne-Nickens P, Alonso A, Djousse L, Forman DE, Gillis AM, Hendriks JML, Hills MT, Kirchhof P, Link MS, Marcus GM, Mehra R, Murray KT, Parkash R, Pina IL, Redline S, Rienstra M, Sanders P, Somers VK, Van Wagoner DR, Wang PJ, Cooper LS, Go AS. Research Priorities in the Secondary Prevention of Atrial Fibrillation: A National Heart, Lung, and Blood Institute Virtual Workshop Report. *J Am Heart Assoc* 2021:e021566.

4. Chalazan B, Mol D, Darbar FA, Ornelas-Loredo A, Al-Azzam B, Chen Y, Tofovic D, Sridhar A, Alzahrani Z, Ellinor P, Darbar D. Association of Rare Genetic Variants and Early-Onset Atrial Fibrillation in Ethnic Minority Individuals. *JAMA Cardiol* 2021.

5. Gudbjartsson DF, Holm H, Sulem P, Masson G, Oddsson A, Magnusson OT, Saemundsdottir J, Helgadottir HT, Helgason H, Johannsdottir H, Gretarsdottir S, Gudjonsson SA, Njolstad I, Lochen ML, Baum L, Ma RC, Sigfusson G, Kong A, Thorgeirsson G, Sverrisson JT, Thorsteinsdottir U, Stefansson K, Arnar DO. A frameshift deletion in the sarcomere gene MYL4 causes early-onset familial atrial fibrillation. *Eur Heart J* 2017;**38**:27-34.

6. Hodgson-Zingman DM, Karst ML, Zingman LV, Heublein DM, Darbar D, Herron KJ, Ballew JD, de Andrade M, Burnett JC, Jr., Olson TM. Atrial natriuretic peptide frameshift mutation in familial atrial fibrillation. *N Engl J Med* 2008;**359**:158-165.

7. Ellinor PT, Nam EG, Shea MA, Milan DJ, Ruskin JN, MacRae CA. Cardiac sodium channel mutation in atrial fibrillation. *Heart Rhythm* 2008;**5**:99-105.

8. Johnson JN, Tester DJ, Perry J, Salisbury BA, Reed CR, Ackerman MJ. Prevalence of early-onset atrial fibrillation in congenital long QT syndrome. *Heart Rhythm* 2008;**5**:704-709.

9.   Rivaud MR, Delmar M, Remme CA. Heritable arrhythmia syndromes associated with
abnormal cardiac sodium channel function: ionic and non-ionic mechanisms.
Cardiovasc Res 2020;**116**:1557-1570.
10.   van Ouwerkerk AF, Bosada FM, Liu J, Zhang J, van Duijvenboden K, Chaffin M, Tucker
NR, Pijnappels D, Ellinor PT, Barnett P, de Vries AAF, Christoffels VM. Identification of
Functional Variant Enhancers Associated With Atrial Fibrillation. *Circ Res*
2020;**127**:229-243.
11.   Chen LY, Ballew JD, Herron KJ, Rodeheffer RJ, Olson TM. A common polymorphism in
SCN5A is associated with lone atrial fibrillation. *Clin Pharmacol Ther* 2007;**81**:35-41.
12.   Benito B, Brugada R, Perich RM, Lizotte E, Cinca J, Mont L, Berruezo A, Tolosana JM,
Freixa X, Brugada P, Brugada J. A mutation in the sodium channel is responsible for
the association of long QT syndrome and familial atrial fibrillation. *Heart Rhythm*
2008;**5**:1434-1440.
13.   Darbar D, Kannankeril PJ, Donahue BS, Kucera G, Stubblefield T, Haines JL, George AL,
Jr., Roden DM. Cardiac sodium channel (SCN5A) variants associated with atrial
fibrillation. *Circulation* 2008;**117**:1927-1935.
14.   Alings M, Wilde A. "Brugada" syndrome: clinical data and suggested
pathophysiological mechanism. *Circulation* 1999;**99**:666-673.
15.   Schott J-J, Alshinawi C, Kyndt F, Probst V, Hoorntje TM, Hulsbeek M, Wilde AAM,
Escande D, Mannens MMAM, Le Marec H. Cardiac conduction defects associate with
mutations in SCN5A. *Nat Genet* 1999;**23**:20-21.
16.   Tan HL, Bink-Boelkens MT, Bezzina CR, Viswanathan PC, Beaufort-Krol GC, van Tintelen
PJ, van den Berg MP, Wilde AA, Balser JR. A sodium-channel mutation causes isolated
cardiac conduction disease. *Nature* 2001;**409**:1043-1047.
17.   Zumhagen S, Veldkamp MW, Stallmeyer B, Baartscheer A, Eckardt L, Paul M, Remme
CA, Bhuiyan ZA, Bezzina CR, Schulze-Bahr E. A heterozygous deletion mutation in the
cardiac sodium channel gene SCN5A with loss- and gain-of-function characteristics
manifests as isolated conduction disease, without signs of Brugada or long QT
syndrome. *PLoS One* 2013;**8**:e67963.
18.   Hesse M, Kondo CS, Clark RB, Su L, Allen FL, Geary-Joo CT, Kunnathu S, Severson DL,
Nygren A, Giles WR, Cross JC. Dilated cardiomyopathy is associated with reduced
expression of the cardiac sodium channel Scn5a. *Cardiovasc Res* 2007;**75**:498-509.
19.   Remme CA, Bezzina CR. Sodium channel (dys)function and cardiac arrhythmias.
Cardiovasc Ther 2010;**28**:287-294.
20.   Wang T, Wehrens XH. Enhanced impact of SCN5A mutation associated with long QT
syndrome in fetal splice isoform. *Heart Rhythm* 2012;**9**:598-599.
21.   Rivolta I, Abriel H, Tateyama M, Liu H, Memmi M, Vardas P, Napolitano C, Priori SG,
Kass RS. Inherited Brugada and long QT-3 syndrome mutations of a single residue of
the cardiac sodium channel confer distinct channel and clinical phenotypes. *J Biol*
Chem 2001;**276**:30623-30630.
22.   Clerx M, Heijman J, Collins P, Volders PGA. Predicting changes to INa from missense
mutations in human SCN5A. *Sci Rep* 2018;**8**:12797.
23.   Makiyama T, Akao M, Shizuta S, Doi T, Nishiyama K, Oka Y, Ohno S, Nishio Y, Tsuji K,
Itoh H, Kimura T, Kita T, Horie M. A novel SCN5A gain-of-function mutation M1875T
associated with familial atrial fibrillation. *J Am Coll Cardiol* 2008;**52**:1326-1334.
24.   Skryabin BV, Kummerfeld DM, Gubar L, Seeger B, Kaiser H, Stegemann A, Roth J,
Meuth SG, Pavenstädt H, Sherwood J, Pap T, Wedlich-Söldner R, Sunderkötter C,

437   Schwartz YB, Brosius J, Rozhdestvensky TS. Pervasive head-to-tail insertions of DNA
438   templates mask desired CRISPR-Cas9-mediated genome editing events. *Sci Adv*
439   2020;**6**:eaax2941.

440 25.   Smirnov A, Fishman V, Yunusova A, Korablev A, Serova I, Skryabin BV, Rozhdestvensky
441   TS, Battulin N. DNA barcoding reveals that injected transgenes are predominantly
442   processed by homologous recombination in mouse zygote. *Nucleic Acids Res*
443   2020;**48**:719-735.

444 26.   Silbernagel N, Walecki M, Schafer MK, Kessler M, Zobeiri M, Rinne S, Kiper AK,
445   Komadowski MA, Vowinkel KS, Wemhoner K, Fortmuller L, Schewe M, Dolga AM,
446   Scekic-Zahirovic J, Matschke LA, Culmsee C, Baukrowitz T, Monassier L, Ullrich ND,
447   Dupuis L, Just S, Budde T, Fabritz L, Decher N. The VAMP-associated protein VAPB is
448   required for cardiac and neuronal pacemaker channel function. *FASEB J* 2018;**32**:6159-
449   6173.

450 27.   Blana A, Kaese S, Fortmüller L, Laakmann S, Damke D, van Bragt K, Eckstein J, Piccini I,
451   Kirchhefer U, Nattel S, Breithardt G, Carmeliet P, Carmeliet E, Schotten U, Verheule S,
452   Kirchhof P, Fabritz L. Knock-in gain-of-function sodium channel mutation prolongs
453   atrial action potentials and alters atrial vulnerability. *Heart Rhythm* 2010;**7**:1862-1869.

454 28.   Lemoine MD, Duverger JE, Naud P, Chartier D, Qi XY, Comtois P, Fabritz L, Kirchhof P,
455   Nattel S. Arrhythmogenic left atrial cellular electrophysiology in a murine genetic long
456   QT syndrome model. *Cardiovasc Res* 2011;**92**:67-74.

457 29.   Holmes AP, Yu TY, Tull S, Syeda F, Kuhlmann SM, O'Brien SM, Patel P, Brain KL, Pavlovic
458   D, Brown NA, Fabritz L, Kirchhof P. A Regional Reduction in Ito and IKACh in the Murine
459   Posterior Left Atrial Myocardium Is Associated with Action Potential Prolongation and
460   Increased Ectopic Activity. *PLoS One* 2016;**11**:e0154077.

461 30.   Syeda F, Holmes AP, Yu TY, Tull S, Kuhlmann SM, Pavlovic D, Betney D, Riley G, Kucera
462   JP, Jousset F, de Groot JR, Rohr S, Brown NA, Fabritz L, Kirchhof P. PITX2 Modulates
463   Atrial Membrane Potential and the Antiarrhythmic Effects of Sodium-Channel
464   Blockers. *J Am Coll Cardiol* 2016;**68**:1881-1894.

465 31.   Yu TY, Syeda F, Holmes AP, Osborne B, Dehghani H, Brain KL, Kirchhof P, Fabritz L. An
466   automated system using spatial oversampling for optical mapping in murine atria.
467   Development and validation with monophasic and transmembrane action potentials.
468   *Prog Biophys Mol Biol* 2014;**115**:340-348.

469 32.   O'Shea C, Holmes AP, Yu TY, Winter J, Wells SP, Correia J, Boukens BJ, De Groot JR,
470   Chu GS, Li X, Ng GA, Kirchhof P, Fabritz L, Rajpoot K, Pavlovic D. ElectroMap: High-
471   throughput open-source software for analysis and mapping of cardiac
472   electrophysiology. *Sci Rep* 2019;**9**:1389.

473 33.   Ackers-Johnson M, Li PY, Holmes AP, O'Brien SM, Pavlovic D, Foo RS. A Simplified,
474   Langendorff-Free Method for Concomitant Isolation of Viable Cardiac Myocytes and
475   Nonmyocytes From the Adult Mouse Heart. *Circ Res* 2016;**119**:909-920.

476 34.   Nuyens D, Stengl M, Dugarmaa S, Rossenbacker T, Compernolle V, Rudy Y, Smits JF,
477   Flameng W, Clancy CE, Moons L, Vos MA, Dewerchin M, Benndorf K, Collen D,
478   Carmeliet E, Carmeliet P. Abrupt rate accelerations or premature beats cause life-
479   threatening arrhythmias in mice with long-QT3 syndrome. *Nat Med* 2001;**7**:1021-
480   1027.

481 35.   Remme CA, Verkerk AO, Nuyens D, van Ginneken AC, van Brunschot S, Belterman CN,
482   Wilders R, van Roon MA, Tan HL, Wilde AA, Carmeliet P, de Bakker JM, Veldkamp MW,

Bezzina CR. Overlap syndrome of cardiac sodium channel disease in mice carrying the
equivalent mutation of human SCN5A-1795insD. *Circulation* 2006;**114**:2584-2594.

36. Holmes AP, Saxena P, Kabir SN, O'Shea C, Kuhlmann SM, Gupta S, Fobian D, Apicella
C, O'Reilly M, Syeda F, Reyat JS, Smith GL, Workman AJ, Pavlovic D, Fabritz L, Kirchhof
P. Atrial resting membrane potential confers sodium current sensitivity to
propafenone, flecainide, and dronedarone. *Heart Rhythm* 2021.

37. Casini S, Albesa M, Wang Z, Portero V, Ross-Kaschitza D, Rougier JS, Marchal GA,
Chung WK, Bezzina CR, Abriel H, Remme CA. Functional Consequences of the SCN5A-
p.Y1977N Mutation within the PY Ubiquitylation Motif: Discrepancy between HEK293
Cells and Transgenic Mice. *Int J Mol Sci* 2019;**20**.

38. Clatot J, Hoshi M, Wan X, Liu H, Jain A, Shinlapawittayatorn K, Marionneau C, Ficker E,
Ha T, Deschênes I. Voltage-gated sodium channels assemble and gate as dimers.
Nature Communications 2017;**8**:2077.

39. Gabelli SB, Boto A, Kuhns VH, Bianchet MA, Farinelli F, Aripirala S, Yoder J, Jakoncic J,
Tomaselli GF, Amzel LM. Regulation of the NaV1.5 cytoplasmic domain by calmodulin.
Nat Commun 2014;**5**:5126.

40. Cerrone M, Remme CA, Tadros R, Bezzina CR, Delmar M. Beyond the One Gene-One
Disease Paradigm: Complex Genetics and Pleiotropy in Inheritable Cardiac Disorders.
Circulation 2019;**140**:595-610.

41. Bogeholz N, Pauls P, Kaese S, Schulte JS, Lemoine MD, Dechering DG, Frommeyer G,
Goldhaber JI, Seidl MD, Kirchhefer U, Eckardt L, Muller FU, Pott C. Triggered activity in
atrial myocytes is influenced by Na(+)/Ca(2+) exchanger activity in genetically altered
mice. *J Mol Cell Cardiol* 2016;**101**:106-115.

42. Voigt N, Li N, Wang Q, Wang W, Trafford AW, Abu-Taha I, Sun Q, Wieland T, Ravens U,
Nattel S, Wehrens XH, Dobrev D. Enhanced sarcoplasmic reticulum Ca2+ leak and
increased Na+-Ca2+ exchanger function underlie delayed afterdepolarizations in
patients with chronic atrial fibrillation. *Circulation* 2012;**125**:2059-2070.

43. Heijman J, Muna AP, Veleva T, Molina CE, Sutanto H, Tekook M, Wang Q, Abu-Taha
IH, Gorka M, Kunzel S, El-Armouche A, Reichenspurner H, Kamler M, Nikolaev V,
Ravens U, Li N, Nattel S, Wehrens XHT, Dobrev D. Atrial Myocyte NLRP3/CaMKII Nexus
Forms a Substrate for Postoperative Atrial Fibrillation. *Circ Res* 2020;**127**:1036-1055.

44. Molina CE, Abu-Taha IH, Wang Q, Rosello-Diez E, Kamler M, Nattel S, Ravens U,
Wehrens XHT, Hove-Madsen L, Heijman J, Dobrev D. Profibrotic, Electrical, and
Calcium-Handling Remodeling of the Atria in Heart Failure Patients With and Without
Atrial Fibrillation. *Front Physiol* 2018;**9**:1383.

45. Schotten U, Verheule S, Kirchhof P, Goette A. Pathophysiological mechanisms of atrial
fibrillation: a translational appraisal. *Physiol Rev* 2011;**91**:265-325.

46. Nattel S, Heijman J, Zhou L, Dobrev D. Molecular Basis of Atrial Fibrillation
Pathophysiology and Therapy: A Translational Perspective. *Circ Res* 2020;**127**:51-72.

47. Goette A, Kalman JM, Aguinaga L, Akar J, Cabrera JA, Chen SA, Chugh SS, Corradi D,
D'Avila A, Dobrev D, Fenelon G, Gonzalez M, Hatem SN, Helm R, Hindricks G, Ho SY,
Hoit B, Jalife J, Kim YH, Lip GY, Ma CS, Marcus GM, Murray K, Nogami A, Sanders P,
Uribe W, Van Wagoner DR, Nattel S, Document R. EHRA/HRS/APHRS/SOLAECE expert
consensus on atrial cardiomyopathies: definition, characterization, and clinical
implication. *Europace* 2016;**18**:1455-1490.

48. Kirchhof PF, Fabritz CL, Franz MR. Postrepolarization refractoriness versus conduction
slowing caused by class I antiarrhythmic drugs: antiarrhythmic and proarrhythmic
effects. *Circulation* 1998;**97**:2567-2574.

49. Liu H, Atkins J, Kass RS. Common molecular determinants of flecainide and lidocaine
block of heart Na+ channels: evidence from experiments with neutral and quaternary
flecainide analogues. *J Gen Physiol* 2003;**121**:199-214.

50. Ramos E, O'Leary M E. State-dependent trapping of flecainide in the cardiac sodium
channel. *J Physiol* 2004;**560**:37-49.

51. Fabritz L, Damke D, Emmerich M, Kaufmann SG, Theis K, Blana A, Fortmuller L,
Laakmann S, Hermann S, Aleynichenko E, Steinfurt J, Volkery D, Riemann B, Kirchhefer
U, Franz MR, Breithardt G, Carmeliet E, Schafers M, Maier SK, Carmeliet P, Kirchhof P.
Autonomic modulation and antiarrhythmic therapy in a model of long QT syndrome
type 3. *Cardiovasc Res* 2010;**87**:60-72.

52. Uzun AU, Mannhardt I, Breckwoldt K, Horvath A, Johannsen SS, Hansen A,
Eschenhagen T, Christ T. Ca(2+)-Currents in Human Induced Pluripotent Stem Cell-
Derived Cardiomyocytes Effects of Two Different Culture Conditions. *Front Pharmacol*
2016;**7**:300.

53. Lemoine MD, Lemme M, Ulmer BM, Braren I, Krasemann S, Hansen A, Kirchhof P,
Meyer C, Eschenhagen T, Christ T. Intermittent Optogenetic Tachypacing of Atrial
Engineered Heart Tissue Induces Only Limited Electrical Remodelling. *J Cardiovasc*
Pharmacol 2020;**77**:291-299.

54. Mali P, Yang L, Esvelt KM, Aach J, Guell M, DiCarlo JE, Norville JE, Church GM. RNA-
guided human genome engineering via Cas9. Science 2013;339:823-826.

55. Thomas KR, Capecchi MR. Site-directed mutagenesis by gene targeting in mouse
embryo-derived stem cells. Cell 1987;51:503-512.

56. Reyat JS, Chua W, Cardoso VR, Witten A, Kastner PM, Kabir SN, Sinner MF, Wesselink
R, Holmes AP, Pavlovic D, Stoll M, Kääb S, Gkoutos GV, de Groot JR, Kirchhof P, Fabritz
L. Reduced left atrial cardiomyocyte PITX2 and elevated circulating BMP10 predict
atrial fibrillation after ablation. JCI Insight 2020;5.