**Infection strategy and biogeography distinguish cosmopolitan groups of marine jumbo bacteriophages**

Alaina R. Weinheimer[1], Frank O. Aylward[1,2]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA. [2] Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0913

Correspondence to email: faylward@vt.edu; address: 5092 Derring Hall, 926 W Campus Drive, Blacksburg, VA 24061; telephone: (540) 231-8657

Competing Interests: The authors declare no competing interests.
Funding Support: This work was supported by grants from the Institute for Critical Technology and Applied Science, the NSF (IIBR-1918271) and a Simons Early Career Award in Marine Microbial Ecology and Evolution to F.O.A. ARW was supported by an ICTAS Doctoral Scholars Fellowship.

**Infection strategy and biogeography distinguish cosmopolitan groups of marine jumbo bacteriophages**

Alaina R. Weinheimer[1], Frank O. Aylward[1,2]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA. [2] Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0913

Correspondence to faylward@vt.edu

**Keywords**: Jumbo bacteriophage, viral diversity, viral gigantism, marine microbiology, microbial oceanography

## ABSTRACT

Recent research has underscored the immense diversity and key biogeochemical roles of large DNA viruses in the ocean. Although they are important constituents of marine ecosystems, it is often difficult to detect these viruses due to their large size and complex genomes. This is true for "jumbo" bacteriophages, which have genome sizes >200 kbp and large capsids reaching up to 0.45 µm in diameter. In this study, we sought to assess the genomic diversity and distribution of these bacteriophages in the ocean by developing a bioinformatic pipeline to generate and validate jumbo phage genomes from metagenomes. We recover 85 marine jumbo phages that ranged in size from 201-498 kilobases, and we examine their genetic similarities and biogeography together with a reference database of marine jumbo phage genomes. By analyzing Tara Oceans metagenomic data we show that jumbo phages are less common in <0.22 µm size fractions but are widespread in larger fractions, consistent with their large size. Our network-based analysis of gene sharing patterns reveals that jumbo bacteriophage belong to five genome clusters that are typified by diverse replication strategies, genomic repertoires, and potential host ranges. Our analysis of jumbo phage distributions in the ocean reveals that depth is a major factor shaping their biogeography, with some phage genome clusters exhibiting higher relative abundance in either surface or mesopelagic waters, respectively. Taken

together, our findings demonstrate that jumbo phages are widespread community members in

the ocean with complex genomic repertoires and ecological impacts that warrant further

targeted investigation.

**Introduction**

Although historically noted for their small virion sizes and simple genomes [1], viruses with large

particles and elaborate genomes have been discovered in recent decades throughout the

biosphere [2–4]. These complex viruses not only invite intriguing evolutionary questions [5–7],

but also expand the potential roles viruses have in shaping microbial community structure and

biogeochemical cycling [4, 8–10]. One group of these larger viruses are jumbo bacteriophages

(jumbo phages), which have traditionally been defined as *Caudovirales* with genomes over 200

kilobases in length [11]. While a recent survey of cultured jumbo phages by Iyer and colleagues

[6] showed jumbo phages share some universal features and genes, such as encoding DNA

polymerases and the terminase large subunit (TerL), these features do not distinguish them

from smaller phages, and several lines of evidence suggest that jumbo phages emerged from

smaller phages multiple times independently. For example, a recent phylogenetic study of

cultured *Caudovirales* used concatenated protein alignments to generate phylogenies and found

that the most supported clades within the *Caudovirales* family do not consistently correspond to

genome length [12]. Furthermore, Iyer et al. found that jumbo phages cluster with smaller

phages based on gene content and are best grouped by replication machinery, among other

infection apparati [6]. Taken together, jumbo phages likely emerged from diverse smaller

phages multiple times independently and form distinct clades within the *Caudovirales*.

Although the first jumbo phages were isolated as early as the 1970s [13], these viruses have

remained relatively sparse in culture, representing less than 3% (n=93) of complete

*Caudovirales* genomes on NCBI's RefSeq Viral Genome Portal (downloaded July 5, 2020). All

cultured jumbo phage capsids have morphologies belonging to the *Myoviridae* and *Siphoviridae*

families and undergo infection cycles that reflect temporal patterns of lytic *Caudovirales* [14].

Some jumbo phages are known to stall infections resulting in "pseudolysogeny", however, which

has been proposed as a competitive strategy against other phages to prevent superinfection

[15]. Jumbo phages that have been studied extensively are primarily investigated for their

unusually complex functional capabilities, such as encoding entire transcriptional apparatti [16]

or sophisticated anti-CRISPR defense mechanisms [17, 18]. Although most cultivated jumbo

phages have been isolated from a variety of Gram positive and Gram negative model and

pathogenic bacteria [14], a recent metagenomic survey uncovered these viruses in diverse

ecosystems around the world [4].

Despite this apparent broad environmental distribution, commonly-used methods for viral

isolation and diversity surveys bias against the inclusion of jumbo phages. Because viruses

have historically been considered to be smaller than cells, many viral diversity surveys

specifically examine only small particle sizes. For example, in plaque assays, agar

concentrations are often too high for larger phage particles to diffuse through compared to

smaller particles [19]. Moreover, filters are often used to remove cells when preparing viral

enrichments for metagenomic sequencing [20], which excludes larger viruses [9, 21].

Particularly in marine studies, the <0.22 µm fraction, sometimes even referred to as the "viral

fraction" [22], is most commonly examined for viruses[23–25]. Jumbo phages can have particles

over 0.45 µm in length (i.e. *Pseudomonas aeruginosa* phage PhiKZ)[13], however, and will

therefore be excluded from <0.22 µm size fractions. Lastly, in bioinformatic pipelines, phage

sequences are typically only assembled to the contig or scaffold level, which is sometimes

sufficient for the assembly of most known smaller phage genomes [26], but often leaves larger

phage genomes fragmented into multiple contigs and may require additional joining of contigs

into bins [27]. Overall, considering these biases and the recently-discovered broad distribution

of these viruses [4], jumbo phages may represent underappreciated components of marine

microbial communities and food webs that warrant further examination.

In this study, we examine the diversity and prevalence of jumbo phages in the global ocean. We

developed a workflow for generating and validating high-quality jumbo bacteriophage bins from

metagenomic data, with which we identified 85 bins of jumbo phages. We then compared the

genetic content of these jumbo phages with other cultured phages of all sizes and metagenomic

jumbo phages from other studies. We found that the jumbo phages of this study grouped into

five distinct clusters that are distinguished by diverse replication machinery and infection

strategies, implicating a broad range of potential jumbo-phage host interactions in the ocean.

We then assessed the distribution of jumbo phages belonging to these genome clusters in the

ocean by using metagenomic data from Tara Oceans [28,29]. Mapping the Tara Oceans

metagenomic data onto the jumbo phage sequences revealed that these jumbos phages are

collectively widely distributed in the ocean, but vary in biogeography based on cluster, with

some more enriched in surface waters relative to deeper waters and vice versa. Upon

examining the collective presence of jumbo phages in different filter fractions, we also found that

commonly used size fractions in viral diversity surveys reduce the ability to recover and detect

jumbo phage sequences. As these complications apply to many viral diversity studies in other

environments, our results indicate that jumbo phages may play underappreciated roles in

ecosystems around the globe.

**RESULTS AND DISCUSSION**

**Detection and validation of high-quality jumbo phage bins**

Due to the large size of jumbo bacteriophage genomes, it is likely that they are present in

multiple distinct contigs in metagenomic datasets and therefore require binning to recover high-

quality metagenome-assembled genomes (MAGs) [27]. This has been shown for large DNA

viruses that infect eukaryotes, where several recent studies have successfully employed binning

approaches to recover viral MAGs [2, 3]. Here, we used the same 1,545 high-quality
metagenomic assemblies [30] used in a recent study to recover giant viruses of eukaryotes [3],
but we modified the bioinformatic pipeline to identify bins of jumbo bacteriophages. We first
binned the contigs from these assemblies with MetaBat2 [31], which groups contigs together
based on similar nucleotide composition and coverage profiles, and focused on bins of at least
200 kilobases in total length. We subsequently identified bins composed of bacteriophage
contigs through analysis with VirSorter2 [32], VIBRANT [33], and CheckV [34] (see Methods for
details).

The occurrence of multiple copies of highly-conserved marker genes is typically used to assess
the level of contamination present in metagenome-derived genomes of bacteria and archaea
[35]. Because bacteriophage lack these marker genes, we developed alternative strategies to
assess possible contamination in our jumbo phage bins. Firstly, we refined the set of bins by
retaining only those with no more than 5 contigs that were each at least 5 kilobases in length to
reduce the possibility that spurious contigs were put together. Secondly, we assessed the
possibility that two strains of smaller phages with similar nucleotide composition may be binned
together by aligning the contigs in a bin to each other. Bins that had contigs with high sequence
similarity across the majority of their lengths were discarded (Supplementary Figure 1). Thirdly,
we discarded bins if their contigs exhibited aberrant co-abundance profiles compared to other
contigs in the same bin. We identified these bins through read mapping and co-abundance
analysis of 225 marine metagenomes from Tara Oceans. Coverage variation was benchmarked
based on read-mapping results from artificially-fragmented reference genomes present in the
samples (See Methods for details). Only bins with coverage variation below our empirically-
derived threshold were retained. Using this stringent filtering, we identified 85 bins belonging to
jumbo bacteriophages that are present in Tara Oceans metagenomes. These bins ranged in
length from 202 bp to 498 kbp, and 31 consisted of a single contig, while 54 consisted of 2-5
contigs (Supplemental Figure 2).

To assess global diversity patterns of jumbo bacteriophage we combined these jumbo phage

bins together with a compiled database of previously-identified jumbo phages that included all

complete jumbo *Caudovirales* genomes on RefSeq (downloaded July 5th, 2020), the

INPHARED database [36], the Iyer et al 2021 study [6], the Al-Shayeb et al. 2020 study [4], and

marine jumbo phage contigs from metagenomic surveys of GOV 2.0 [25] (60 jumbo phages),

ALOHA 2.0 [37] (8 jumbo phages), and one megaphage MAG recovered from datasets of the

English Channel [38]. Ultimately, we arrived at a set of 223 jumbo phages present in at least

one Tara Oceans sample that we analyzed further in this study and refer to as marine jumbo

phages. Statistics on genomic features can be found in Supplemental Dataset 1.

**Marine jumbo phages belong to distinct groups with diverse infection strategies**

Because bacteriophages lack high-resolution, universal marker genes for classification, such as

16S rRNA in bacteria, phages are often grouped by gene content [39, 40]. Here, we generated

a bipartite network that included the 85 bins of jumbo phages with a dataset of available

*Caudovirales* complete genomes in RefSeq (3,012 genomes; downloaded July 5th, 2020) and

the set of jumbo phages described above. To construct the bipartite network, we compared

proteins encoded in all the phage genomes to the VOG database (vogdb.org), and each

genome was linked to VOG hits that were present (Figure 1, Supplemental Dataset 2, see

Methods for details). To identify groups of phage genomes with similar VOG profiles, we

employed a spinglass community detection algorithm [41] to generate genome clusters. The

marine jumbo phages of this study clustered into five groups that all included both jumbo and

non-jumbo phage genomes (Figure 2a). We refer to these five clusters as Phage Genome

Clusters (PGCs): PGC_A, PGC_B, PGC_C, PGC_D, and PGC_E. These PGCs included

cultured phages and metagenome-derived jumbo phages found in various environments (i.e.

aquatic, engineered) and isolated on a diversity of hosts (i.e. Firmicutes, Proteobacteria,

Bacteroidetes) (Figure 2b,c). Of the marine jumbo phages, 134 belonged to PGC_A, 11 to

PGC_B, 90 to PGC_C 27, 7 to PGC_D, and 1 to PGC_E (Figure 1b). In addition to this network-

based analysis, we also examined phylogenies of the major capsid protein (MCP) and the

terminase large subunit (TerL) encoded by the marine jumbo phages and the same reference

phage set examined in the network (Figure 1c, 1d). With the exception of PGC_A, the marine

jumbo phages that belong to the same PGC appeared more closely related to each other than

those belonging to different clusters. The evolutionary history of these marker genes therefore

supports the results of the bipartite network genome clustering and is consistent with the view

that genome gigantism evolved multiple times independently within the *Caudovirales* [6].

We then compared functional content encoded by the marine jumbo phages in the PGCs to

identify functional differences that distinguish these groups. PGC_E was excluded from this

analysis because this genome cluster contained only a single jumbo phage. Collectively, most

genes of the marine jumbo phages could not be assigned a function (mean: 86.60%, std dev:

7.01%; Supplemental Dataset 3), which is common with environmental viruses [42, 43]. Genes

with known functions primarily belonged to functional categories related to viral replication

machinery, such as information processing and virion structure (Figure 3a), and these genes

drove the variation between the genome clusters of marine jumbo phages (Figure 3b). Iyer et al

provided a comparative genomic analysis that was able to identify three types of jumbo phages

that are defined by different infection strategies and host interactions (referred to as Groups 1-3)

[6]. We cross-referenced our PGCs and found that PGCs B, C, and D of this study

corresponded to Iyer Groups 1, 2, and 3, respectively, suggesting that these genome clusters

contain phages with distinct infection and replication strategies. PGC_A corresponded to

multiple phage groups in the Iyer study, indicating that this genome cluster contains a

particularly broad diversity of phages.

PGC_B consists of 238 phages (11 (4.6%) marine jumbo phages, including 10 bins generated

here), and included cultured phage of Group 1 of Iyer et al, which is typified by *Pseudomonas*

*aeruginosa* phage PhiKZ. Supporting this correspondence with Iyer Group 1, all marine jumbo

phages of PGC_B encoded the same distinct replication and transcription machinery, including

a divergent family B DNA polymerase and a multi-subunit RNA polymerase (Figure 3b,

Supplemental Dataset 3). These marine jumbo phages also encoded a PhiKZ internal head

protein, and they uniquely encoded shell and tubulin homologs which has recently been found in

PhiKZ phages to assist in the formation of a nucleus-like compartment during infection that

protects the replicating phage from host defenses [17,18]. Although we could not confidently

predict hosts for the 11 metagenomic marine jumbo phages in this PGC_B (Supplemental

Dataset 1), the cultured phages of this genome cluster infect pathogenic bacteria belonging to

the phyla Proteobacteria (178 phages) and Firmicutes (6 phages) (Figure 2c), implicating a

potential host range for marine jumbo phages in PGC_B.

PGC_C is comprised of 156 phages total (90 marine jumbo phages (57.7%); 4 bins generated

from this study) and included reference jumbo phages in Iyer Group 2 (31, 19.9%) which are

typified by Alphaproteobacteria and Cyanobacteria phages. Likewise, the host range of other

cultured phages in PGC_C support the Group 2 correspondence, either infecting Cyanobacteria

(139 phages) or Proteobacteria (4 phages) (Figure 2c). Furthermore, all 3 marine metagenomic

phages in PGC_C for which hosts could be predicted were matched to Cyanobacteria hosts

(Supplemental Dataset 1). Functional annotations of PGC_C marine jumbo phages revealed

nearly all encode a family B DNA polymerase (97.8% phages) and the photosystem II D2

protein (PF00124, VOG04549) characteristic of cyanophages (90% phages) (Figure 3b). This

PGC included the reference *Prochlorococcus* phage P-TIM68 (NC_028955.1), which encodes

components of both photosystem I and II as a mechanism to enhance cyclic electron flow during

infection [44]. This suggests that an enhanced complement of genes used to manipulate host

physiology during infection may be a driver of large genome sizes in this group. Additionally,

most of the PGC_C marine jumbo phages encoded Lipopolysaccharide (LPS) biosynthesis

proteins (76%), which have been found in cyanophage genomes that may induce a

"pseudolysogeny" state, when infected host cells are dormant, by changing the surface of the

host cell and preventing additional phage infections [15] (Supplemental Dataset 3). Taken

together, most marine jumbo phages of PGC_C likely follow host interactions of jumbo

cyanophages, such as potentially manipulating host metabolism by encoding their own

photosynthetic genes and potentially inducing a pseudolysogenic state.

PGC_D comprised 47 phages, of which 7 were marine jumbo phages generated in this study

(14.9%). This group included Iyer Group 3 jumbo phages (15, 31.9%), which is primarily

distinguished by encoding a T7-type DNA polymerase but is not typified by a particular phage

type or host range. Supporting this grouping, all marine jumbo phages in this study encoded this

T7 DNA polymerase which belongs to family A DNA polymerases (Figure 3b, Supplemental

Dataset 3). Most of the other genes distinctively encoded by the marine jumbo phages in this

group included structural genes related to T7 (T7 baseplate, T7 capsid proteins), a eukaryotic

DNA topoisomerase I catalytic core (PF01028), and a DNA structural modification genes

(MmcB-like DNA repair protein, DNA gyrase B). Hosts of metagenomic marine jumbo phages in

PGC_D could not be predicted (Supplemental Dataset 1); however, cultured Group 3 jumbo

phages in PGC_D all infect Proteobacteria, primarily Enterobacteria and other pathogens.

PGC_A was the largest of the PGCs with 469 phages in total, including 134 marine jumbo

phages (63 bins from this study). This genome cluster contained the largest jumbo phages,

such as *Bacillus* phage G (498 kb) and the marine megaphage Mar_Mega_1 (656 kb) recently

recovered from the English Channel [38]. Unlike other PGCs, PGC_A contained mostly

metagenomic phages (401, 85%, Figure 2b,c). Considering PGC_A contains the largest jumbo

phages (Figure 1b, 2a), the vast genetic diversity in this PGC might explain why few genes were

found to distinguish this group. Of the genes unique to PGC_A, only one was present in at least

half of the phages (51.9%), which was a Bacterial DNA polymerase III alpha NTPase domain

(PF07733). The host ranges of cultured phages from this PGC further reflect the phage diversity

and included a variety of phyla and genera that can perform complex metabolisms or lifestyles,

such as the nitrogen-fixing Cyanobacteria of the *Nodularia* genus isolated from the Baltic Sea

(corresponding to phage sequences NC_048756.1 and NC_048757.1) and the Bacteroidetes

bacteria *Rhodothermus* isolated from a hot spring in Iceland [45].

**Relative abundance of jumbo bacteriophages across size fractions**

To explore the distribution of the marine jumbo phages in the ocean, we first examined the size

fractions in which the jumbo phages were most prevalent. To remove redundancy, we clustered

the 223 jumbo phages into populations if they were highly similar at the nucleotide level ( >80%

genes with >95% average nucleotide identity) [23], which resulted in 120 jumbo phage

populations. Out of the 225 Tara Oceans metagenomes examined, 202 (89.8%) contained at

least one jumbo phage population (median: 24, Supplemental Dataset 4). Jumbo phages were

less frequently detected in samples below 0.22 μm (<-0.22 μm, 0.1-0.22 μm) than those above

0.22 μm (0.45-0.8 μm, 0.22-0.45 μm, 0.22-1.6 μm, 0.22-3 μm). Only 54.5% of samples in the <-

0.22 μm fraction and 82.4% of samples in the 0.1-0.22 μm fraction had at least one jumbo

phage present. Conversely, all samples in all larger size fractions above 0.22 μm had at least

one jumbo phage present (Figure 4a). Furthermore, we detected 21 populations (14.9%)

exclusively in samples above 0.22 um, compared to only one population (0.71%) exclusive to

samples below 0.22 μm. A similar disparity in virus detection between size fractions has been

reported for large eukaryotic viruses, where roughly 41% of phylotypes were present in samples

of the picoplankton fraction of 0.22-3 μm but absent in size fractions below 0.22 μm [9]. In

contrast to this study, where certain viral groups were more prevalent in larger size fractions

than smaller, a jumbo phage's PGC membership or genome size did not affect its probability of

detection at different size fractions (Supplemental Figure 3, Supplemental Figure 4).

We also compared jumbo phage diversity (defined as population richness), relative abundance,

and community composition between the size fractions. Samples of the size fraction <-0.22 μm

were significantly less diverse (P-value 0.0003, Wilcox test) and had significantly lower relative

abundances of jumbo phages relative to the large size fractions of 0.1-0.22 and 0.22-1.6 (P-

values 0.003, 0.031 Wilcox test, respectively). Samples of the 0.1-0.22 μm size fraction had the

highest diversity and relative abundances compared to the other fractions (Figure 4b,c). The

enhanced signal of jumbo phages relative to other entities in the 0.1-0.22 µm size fraction may

result from the reduction of cells below the 0.22 µm threshold and the reduction of most other

viruses above the 0.1 µm limit. Despite these differences in diversity and relative abundances,

jumbo phage community composition did not significantly differ between the >0.22 and <0.22

µm size fractions (P-value 0.8109, ANOSIM, presence/absence Bray-Curtis distance matrix,

Figure 4d,e).

To directly test the effect of the 0.22 µm size fraction cut-off on jumbo phage recovery, we

examined a subset of the samples that were co-collected at the same station and depth for the

fractions below 0.22 µm (<-0.22 or 0.1-022) and above 0.22 µm (0.1-0.22 µm or 0.22-3 µm).

The number of detected jumbo phage populations was significantly higher in samples above

0.22 µm than below 0.22 µm (P-value = 0.0004, Wilcox test, Supplemental Figure 5a). The

relative abundance of jumbo phages was not significantly different between this size distinction

(P-value = 0.1309, Wilcox test, Supplemental Figure 5b), however, nor was community

composition (P-value = 0.1229, ANOSIM, presence/absence Bray-Curtis distance matrix,

Supplemental Figure 5c,d). Taken together, these findings suggest that using size fractions

below 0.22 µm to analyze phages limits the recovery of jumbo phage populations detected,

likely due the large size of these viruses. The enhanced relative abundance of jumbo phages in

the 0.1-0.22 µm relative to the other size fractions suggests that this may be the optimal size

fraction to examine for studies that seek to target jumbo phages specifically.

**Biogeography of jumbo bacteriophages in the global ocean**

*Jumbo phage populations varied in depth distribution by PGC.* Jumbo populations in this study

were found in all three Longhurst biomes (Trades, Westerlies, Coastal) and depths (Surface,

Deep Chlorophyll Maximum, Mesopelagic) examined (Figure 5, Supplemental Figure 6;

Supplemental Dataset 4). We focused these biogeographic analyses on the 0.22-1.6 or 0.22-3

µm size fractions because the most sites were available for these samples, and because we

found that jumbo phages were recovered at a higher frequency in larger size fractions. Jumbo

phage communities differed significantly between depths (P-value = 0.0001, ANOSIM based on

presence/absence Bray-Curtis distance matrix, Supplemental Figure 7c,d), consistent with the

dramatic transition in community composition that occurs from surface waters to below the deep

chlorophyll maximum [37, 46]. Specifically, the diversity of jumbo phages across depths varied

by genome cluster, with PGC_A and PGC_C exhibiting higher abundance and diversity in the

epipelagic (SRF and DCM), and PGC_B and PGC_D exhibiting higher abundance in the

mesopelagic (Figure 5d, Supplemental Figure 7a,b). PGC_C is typified by cyanophages,

providing a clear reason why this phage group is enriched in surface waters. Conversely,

PGC_B is typified by *Pseudomonas aeruginosa* PhiKZ phages, suggesting these PGC_B

marine jumbo phages may be infecting heterotrophic bacteria that potentially thrive in deeper

waters. Overall relative abundance and diversity of jumbo phages in this study were significantly

higher in the epipelagic zone, partly because most of these viruses are in PGC_A. This pattern

held when examining only those samples that were co-collected at all three depths

(Supplemental Figure 8,9). This general pattern therefore reflects what has been found in

previous studies on the depth distribution and viral protein clusters, where more were unique in

the euphotic (i.e. epipelagic) than aphotic depths [9, 25, 47, 48], although this contrasts what

has recently been found in the Pacific Ocean, where overall viral diversity increased in the

mesopelagic [37].

*Jumbo phage biogeography across biomes.* Jumbo phage communities in this study

significantly differed in composition in surface samples between the Trades, Coastal, and

Westerlies biomes, albeit weakly (P-value 0.0021, ANOSIM, presence/absence Bray-Curtis

distance matrix, Supplemental Figure 10c,d). However, no biome appeared to be a particular

hotspot for jumbo phages, as they were not significantly more diverse in any particular biome

(Supplemental Figure 10a). Although the relative abundances of jumbo phages were higher in

Trades samples relative to Coastal samples (Supplemental Figure 10b), this pattern did not

always hold when comparing biomes at different depths (Supplemental Figure 11). Similarly,

jumbo phages from all PGCs could be detected in all biomes, but no PGC was consistently

enriched in diversity and relative abundance in a particular biome (Supplemental Figure 12). A

recent global study on marine viruses has found that viral diversity was better explained by

ecological zones defined by physicochemical factors like temperature, rather than by Longhurst

biomes defined by patterns of chlorophyll *a* concentrations [25], suggesting that Longhurst

biomes may not be good predictors of viral diversity in general.

Jumbo phage populations ranged in endemicity with seven populations (5.8%) detected in only

one Tara station and eleven (9.2%) present in 50 or more stations (Figure 5c). More endemic

populations, present in only one Tara station, belonged to PGCs A, B, and C, while the most

prevalent jumbo phage, those present in 50 or more stations, belonged to PGC_A and PGC_C.

PGC_A and PGC_C contained the most populations, which likely explains the wide range of

endemicity of phages in these clusters. Moreover, the cyanobacterial hosts that are known for

many of the jumbo phages in PGC_C are widespread in the ocean, which may also explain the

prevalence of this group of phages. In general, the heterogeneous distribution and abundance

of these jumbo phages is consistent with the seed bank hypothesis, which postulates that

viruses are passively dispersed throughout the ocean and viral community structure is shaped

by local selective forces [23, 49]. This framework has previously been used to explain why

phage distributions range from extremely cosmopolitan to extremely rare, which is a pattern that

also appears to hold for jumbo bacteriophages.

**CONCLUSION**

Large DNA viruses are becoming increasingly recognized as critical components of the

virosphere, notable for their intriguing evolutionary histories [50, 51], vast functional capacities

[3, 4], and global distribution [2, 4]. Here, we uncover the diversity and ecology of marine jumbo

bacteriophages, which have historically been difficult to study due to biases in filtration and

isolation strategies commonly used in virology. We employed a binning strategy to generate and

quality-check genomes of jumbo phages and used it to identify 85 high quality bins. We

combined these bins together with reference jumbo bacteriophage genomes, and ultimately

identified 120 populations that are present in Tara Oceans metagenomes. When compared with

other metagenomic jumbo phages and cultured phages of all sizes, we found that marine jumbo

phages primarily belong to five phage genome clusters (PGCs) that encode distinct replication

machinery, biogeography, and potential hosts. For example, marine jumbo phages in PGC_C

follow cyanophage infection strategies and ecology, as this cluster included cultured marine

cyanophages and encoded classic family B DNA polymerases and photosynthesis enzymes

characteristic of cyanophages. Furthermore, we found they are enriched in surface waters

relative to the mesopelagic, consistent with the geographic range of their hosts. In contrast,

marine jumbo phages of PGC_B included cultured PhiKZ phages of *Pseudomonas aeruginosa*

and uniquely encoded multi-subunit RNA polymerases and tubulin, which are thought to play a

role in the remarkable nucleus-like structures that these viruses employ as an anti-CRIPSR

defense [17, 18]. PGC_B was particularly enriched in mesopelagic waters, suggesting that this

complex infection strategy may be more common in the deep ocean. These results demonstrate

the diverse biology and ecology of jumbo phages, consistent with the hypothesis that jumbo

phages evolved from smaller phages on independent occasions [6].

We show that jumbo phages are widespread throughout the ocean and are typically more

diverse and abundant in epipelagic waters, which reflect previous findings that surface waters

usually harbor a higher per-sample alpha diversity of viral groups compared to deeper waters [9,

25]. Larger phages therefore appear to coexist in patterns broadly similar to smaller viruses

despite the disadvantages of their size, such as smaller burst sizes and lower host contact rates

[52]. In eukaryotic giant viruses it has been hypothesized that these disadvantages are offset by

higher infection efficiency, broader host ranges, potentially decreased decay rates, and higher

rates of successful attachments compared to smaller viruses [52]. Although some of these

advantages to viral gigantism may also apply to jumbo bacteriophages, it is unlikely that they

are all applicable. For example, given that they are tailed *Caudovirales* [6], they likely possess

higher host specificity in part due to their non-phagocytotic mode of infection. Nonetheless, the

large genomes of jumbo bacteriophage often encode an expanded complement of genes used

to manipulate host physiology during infection, and these may play critical roles in promoting

infection efficiency or offsetting host defense mechanisms. The impressive complement of

photosynthesis genes in PGC_C is at least partially responsible for the large genomes in this

lineage, while the genes involved in anti-CRISPR defense found in PGC_B indicate that a host-

virus arms race may be responsible for genome gigantism in this group. Interestingly, the

largest number of jumbo phage genomes we identified belong to PGC_A, which is largely

uncharacterized and composed of primarily metagenome-derived genomes, suggesting that

these viruses have as-yet unidentified infection strategies. Overall, it is likely that the factors

leading to genome gigantism in each of these genome clusters are different, given that they

represent distinct lineages of *Caudovirales* that have independently evolved large genomes.

Future work further characterizing the hosts of these jumbo phages and the details of their

infection program, particularly in PGC_A, will therefore be critical to understanding mechanisms

that underlie complexity in the virosphere and maintain diversity.

**Methods Summary**

**Jumbo phage bin detection.** Metagenomic scaffolds were downloaded from 1,545 assembled

by Parks et al. 2017 [30] and binned with MetaBAT2 [31] (-s 200000 --unbinned -t 32 -m 5000 --

minS 75 --maxEdges 75). Bins were retained if they summed to at least 200,000 base-pairs and

comprised <= 5 contigs (min. contig size 5kb). Proteins were predicted with prodigal [53] using

default settings on each bin individually. Bins were retained if they lacked more than one

ribosomal protein, lacked overlapping regions (via promer and gnuplot [54] with MUMmer 3.0

[55]), had fewer hits to NCLDV than phage (via LASTp [56] against RefSeq r99), lacked more

than one NCLDV marker gene (via HMM search (hmmer.org) against NCLDV marker gene

HMM profiles [57]) and had even read coverage of Tara Ocean metagenomes via coverM [58]

(https://github.com/wwood/CoverM). Jumbo phages were then detected by running the bins

through VirSorter2 [32], VIBRANT [33], and CheckV [34]. Bins were considered putative phages

if they had at least an average dsDNAphage score of >0.9 from VirSorter2 or VirSorter2

average score >0.5 and either i) CheckV quality of medium or higher (via pseudocontigs) or ii)

VIBRANT consensus classification as viral. 85 bins were retained for downstream analyses.

Bins were clustered into populations with a compiled set of jumbo bacteriophages (RefSeq

phages over 200 kilobases, phage sequences over 200 kilobases from the INPHARED

database [36], Iyer et al 2021 [6], Al-Shayeb et al. 2020 [4] jumbo phage genomes, GOV 2.0

[25], ALOHA 2.0 [37], and a megaphage assembled from the English Channel [38]) based on

nucleotide sequences of genes (predicted with prodigal -d flag) aligned with BLASTn [59] (>95%

average nucleotide identity, >80% genes) [23]. See Supplemental Methods for details.

**Evolutionary analyses.** Reference phage sequences were compiled from RefSeq's

*Caudovirales* complete genomes (downloaded July 2020 from NCBI's Virus genome portal;

3,012 genomes) along with the curated jumbo phage set used in the population analysis.

Proteins of jumbo bins and this reference set were predicted with prodigal and searched against

the Virus Orthologous Groups (VOGs, vogdb.org) via HMM searches (E-value 0.001). Bipartite

network was made based on shared VOGs using igraph (graph.incidence) (1.2.5) [60] in R

(version 3.5.1) [61] with RStudio (version 1.1.456) [62]. Clusters were detected by the spinglass

community detection algorithm [41] via igraph (50 spins for 100 iterations). Final clusters

determined by iteration with highest modularity. Phages were further classified with vContact2

[39]; however, none of the jumbo phage bins belonged to clusters that could be assigned an

ICTV taxa. Plots of the cluster composition from the bipartite network analysis were made with

ggplot2 (3.1.1) [63] in R with Rstudio. TerL (terminase large subunit) and MCP (major capsid

protein) trees were made with hits to TerL VOG families and MCP VOG families encoded by the

jumbo phages and reference hits (HMM searches, E-value < 0.001; Supplemental Dataset 2).

Reference hits were de-replicated with CD-HIT (-c 0.9) [64] and filtered for size (See

Supplementary Methods). Proteins were then aligned with Clustal Omega [65], trimmed with

trimAl (-gt 0.1) [66] and constructed with IQ-TREE [67] (TEST model selection with ModelFinder

[68]).

**Size fraction and ecological analyses.** Metagenomic reads from Tara Oceans were mapped

onto the population representatives of the jumbo phage set (535 populations) with coverM

(coverm genome; >10% genome covered, min ID 95%), and reads per kilobase per million

(RPKM) was then calculated by dividing the number of reads aligned to a phage by the length of

the phage in kilobases and then dividing that by the number of reads in the sample in millions,

which accounts for differences in phage sequence length and differences in sample sequencing

depth. Statistical analyses and plots were carried out in R with vegan (2.5-5) [69], ggplot2, maps

(3.3.0) [70], and ggpubr (0.2.4) [71] packages. Community composition was compared between

variables using ANOSIMs based on Bray-Curtis distances using both presence/absence and

RPKM matrices with a significance p-value cutoff of 0.05. Statistical tests were carried out with

the ggplot2 function stat_compare_means(label="p.signif").

**Annotation.** Amino acid sequences of genes were annotated with HMM searches (Evalue <

0.001) against the Pfam [72] (version 32), eggNOG [73], and VOG databases. Virion structural

protein families were identified based on VOG hit descriptions (Supplemental Dataset 3).

Consensus annotation was based on Pfam annotations and then the highest bit score between

eggNOG and VOG hits. Functions were grouped into larger categories (Supplemental Dataset

3). Clusters from the network analyses were compared for functional composition by averaging

the proportion of genes in a functional category using python (version 3) statistics and plotted

with ggplot2 in R. Genes with the highest variance between clusters were identified based on

the variance in proportion of genomes in a cluster with that gene. Those with over 0.2 variance

and a known function were visualized with pheatmap [74] in R.

**Host prediction.** Hosts of the jumbo phage bins were predicted based on matching CRISPR

spacers, tRNAs, and gene content. CRISPR spacers were predicted on the Genome Taxonomy

Database (release 95) [75], metagenome assembled genomes (MAGs) of bacteria and archaea

from the metagenomes that the jumbo phage bins derived (provided by Parks et al. 2017 [30]),

and on the jumbo phage bins with minCED (derived from reference [76]). Spacers were aligned

with BLASTn and matches were >24 bp with <= 1 mismatches [4]. tRNA sequences were

predicted with tRNAscan-SE (-bacteria option) [77] on the MAGs and jumbo phage bins.

Promiscuous tRNAs [78] were removed (BLASTn hits 100% ID, <= 1 mismatches). Jumbo

phage tRNAs were aligned against the MAGs tRNAs and NCBI nr database (BLASTn 100% ID,

<= 1 mismatches) [4]. Lastly, hosts were assigned based on the taxonomy of coding sequence

matches to the MAGs (BLASTn). Hits to phyla were summed and a putative host phylum had

three times the number of hits as the phylum with the next most hits as used in a previous study

[4].

**Acknowledgements**

We thank members of the Aylward Lab for helpful feedback. This work was performed using

compute nodes available at the Virginia Tech Advanced Research and Computing Center.

**Competing Interests**

The authors declare no competing financial interests.

**References**

1.   Brüssow H, Hendrix RW. Phage Genomics: Small is Beautiful. *Cell* 2002; **108**: 13–16.

2.   Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh DA, Denef VJ, et al. Giant virus

diversity and host interactions through global metagenomics. *Nature* 2020; **578**: 432–436.

3.   Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO. Dynamic genome

evolution and complex virocell metabolism of globally-distributed giant viruses. *Nat*

*Commun* 2020; **11**: 1710.

4.    Al-Shayeb B, Sachdeva R, Chen L-X, Ward F, Munk P, Devoto A, et al. Clades of huge

phages from across Earth's ecosystems. *Nature* 2020; **578**: 425–431.

5.    Koonin EV, Yutin N. Origin and Evolution of Eukaryotic Large Nucleo-Cytoplasmic DNA

Viruses. *Intervirology* . 2010; **53**: 284–292.

6.    M Iyer L, Anantharaman V, Krishnan A, Burroughs AM, Aravind L. Jumbo Phages: A

Comparative Genomic Overview of Core Functions and Adaptions for Biological Conflicts.

*Viruses* 2021; **13**.

7.    Raoult D, Forterre P. Redefining viruses: lessons from Mimivirus. *Nat Rev Microbiol* 2008;

**6**: 315–319.

8.    Ha AD, Moniruzzaman M, Aylward FO. High Transcriptional Activity and Diverse Functional

Repertoires of Hundreds of Giant Viruses in a Coastal Marine System. *mSystems* 2021; **6:**

e00293-21.

9.    Endo H, Blanc-Mathieu R, Li Y, Salazar G, Henry N, Labadie K, et al. Biogeography of

marine giant viruses reveals their interplay with eukaryotes and ecological functions. *Nat*

*Ecol Evol* 2020; **4**: 1639–1649.

10.   Kaneko H, Blanc-Mathieu R, Endo H, Chaffron S, Delmont TO, Gaia M, et al. Eukaryotic

virus composition can predict the efficiency of carbon export in the global ocean. *iScience*

2021; **24**: 102002.

11.   Hendrix RW. Jumbo bacteriophages. *Curr Top Microbiol Immunol* 2009; **328**: 229–240.

12.   Low SJ, Džunková M, Chaumeil P-A, Parks DH, Hugenholtz P. Evaluation of a

concatenated protein phylogeny for classification of tailed double-stranded DNA viruses

belonging to the order Caudovirales. *Nat Microbiol* 2019; **4**: 1306–1315.

13.   Krylov VN, Zhazykov IZ. Pseudomonas bacteriophage phiKZ--possible model for studying

the genetic control of morphogenesis. *Genetika* 1978; **14**: 678–685.

14.   Yuan Y, Gao M. Jumbo Bacteriophages: An Overview. *Frontiers in Microbiology* 2017; **8**:

403.

15.  Bertani B, Ruiz N. Function and Biogenesis of Lipopolysaccharides. *EcoSal Plus* 2018; **8**:

https://doi.org/10.1128/ecosalplus.ESP-0001-2018.

16.  Ceyssens P-J, Minakhin L, Van den Bossche A, Yakunina M, Klimuk E, Blasdel B, et al.

Development of Giant Bacteriophage KZ Is Independent of the Host Transcription

Apparatus. *Journal of Virology.* 2014; **88**: 10501–10510.

17.  Mendoza SD, Nieweglowska ES, Govindarajan S, Leon LM, Berry JD, Tiwari A, et al. A

bacteriophage nucleus-like compartment shields DNA from CRISPR nucleases. *Nature*

2020; **577**: 244–248.

18.  Malone LM, Warring SL, Jackson SA, Warnecke C, Gardner PP, Gumy LF, et al. A jumbo

phage that forms a nucleus-like structure evades CRISPR–Cas DNA targeting but is

vulnerable to type III RNA-based immunity. *Nature Microbiology* . 2020; **5**: 48–55.

19.  Serwer P, Hayes SJ, Thomas JA, Hardies SC. Propagating the missing bacteriophages: a

large bacteriophage in a new class. *Virol J* 2007; **4**: 21.

20.  Thurber RV, Haynes M, Breitbart M, Wegley L, Rohwer F. Laboratory procedures to

generate viral metagenomes. *Nat Protoc* 2009; **4**: 470–483.

21.  Palermo CN, Shea DW, Short SM. Analysis of Different Size Fractions Provides a More

Complete Perspective of Viral Diversity in a Freshwater Embayment. *Appl Environ*

*Microbiol* 2021; **87**: e00197-21.

22.  Sieburth JM, Smetacek V, Lenz J. Pelagic ecosystem structure: Heterotrophic

compartments of the plankton and their relationship to plankton size fractions 1. *Limnology*

*and Oceanography* 1978; **23**: 1256–1263.

23.  Brum JR, Ignacio-Espinoza JC, Roux S, Doulcier G, Acinas SG, Alberti A, et al. Ocean

plankton. Patterns and ecological drivers of ocean viral communities. *Science* 2015; **348**:

1261498.

24.  Roux S, Brum JR, Dutilh BE, Sunagawa S, Duhaime MB, Loy A, et al. Ecogenomics and

potential biogeochemical impacts of globally abundant ocean viruses. *Nature* 2016; **537**:

689–693.

25. Gregory AC, Zayed AA, Conceição-Neto N, Temperton B, Bolduc B, Alberti A, et al. Marine

DNA Viral Macro- and Microdiversity from Pole to Pole. *Cell* 2019; **177**: 1109–1123.e14.

26. Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. Benchmarking viromics: an

evaluation of metagenome-enabled estimates of viral community composition and diversity.

*PeerJ* 2017; **5**: e3817.

27. García-López R, Vázquez-Castellanos JF, Moya A. Fragmentation and Coverage Variation

in Viral Metagenome Assemblies, and Their Effect in Diversity Calculations. *Front Bioeng*

*Biotechnol* 2015; **3**: 141.

28. Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, et al. Ocean

plankton. Structure and function of the global ocean microbiome. *Science* 2015; **348**:

1261359.

29. Sunagawa S, Coordinators TO, Acinas SG, Bork P, Bowler C, Eveillard D, et al. Tara

Oceans: towards global ocean ecosystems biology. *Nature Reviews Microbiology* 2020; **18**:

428–445.

30. Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, et al.

Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree

of life. *Nat Microbiol* 2017; **2**: 1533–1542.

31. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies.

*PeerJ* 2019; **7**: e7359.

32. Guo J, Bolduc B, Zayed AA, Varsani A, Dominguez-Huerta G, Delmont TO, et al.

VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA and RNA

viruses. *Microbiome* 2021; **9**: 37.

33. Kieft K, Zhou Z, Anantharaman K. VIBRANT: automated recovery, annotation and curation

of microbial viruses, and evaluation of viral community function from genomic sequences.

*Microbiome* 2020; **8**: 90.

34.  Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. CheckV

assesses the quality and completeness of metagenome-assembled viral genomes. *Nat*

*Biotechnol* 2021; **39**: 578–585.

35.  Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

*Genome research* 2015; **7**: 1043-1055.

36.  Cook R, Brown N, Redgwell T, Rihtman B, Barnes M, Clokie M, et al. INfrastructure for a

PHAge REference Database: Identification of Large-Scale Biases in the Current Collection

of Cultured Phage Genomes. *PHAGE: Therapy, Applications, and Research* 2021; e-pub

ahead of print 5 October 2021; http://doi.org/10.1089/phage.2021.0007

37.  Luo E, Eppley JM, Romano AE, Mende DR, DeLong EF. Double-stranded DNA

virioplankton dynamics and reproductive strategies in the oligotrophic open ocean water

column. *ISME J* 2020; **14**: 1304–1315.

38.  Michniewski S, Rihtman B, Cook R, Jones MA, Wilson WH, Scanlan DJ, et al. Identification

of a new family of 'megaphages' that are abundant in the marine environment.

39.  Jang HB, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al. Gene sharing

networks to automate genome-based prokaryotic viral taxonomy. *Nat Biotechnology* 2019;

**37**: doi:10.1101/533240.

40.  Iranzo J, Krupovic M, Koonin EV. The Double-Stranded DNA Virosphere as a Modular

Hierarchical Network of Gene Sharing. *MBio* 2016; **7**: e00978-16.

41.  Reichardt J, Bornholdt S. Statistical mechanics of community detection. *Phys Rev E Stat*

*Nonlin Soft Matter Phys* 2006; **74**: 016110.

42.  Hurwitz BL, Sullivan MB. The Pacific Ocean virome (POV): a marine viral metagenomic

dataset and associated protein clusters for quantitative viral ecology. *PLoS One* 2013; **8**:

e57355.

43. Pappas N, Dutilh BE. Finding functional associations between prokaryotic virus orthologous

groups: a proof of concept. *BMC Bioinformatics* 2021; **22**: 1–11.

44. Fridman S, Flores-Uribe J, Larom S, Alalouf O, Liran O, Yacoby I, et al. A myovirus

encoding both photosystem I and II proteins enhances cyclic electron flow in infected

*Prochlorococcus* cells. *Nat Microbiol* 2017; **2**: 1350–1357.

45. Hjorleifsdottir S, Aevarsson A, Hreggvidsson GO, Fridjonsson OH, Kristjansson JK.

Isolation, growth and genome of the *Rhodothermus* RM378 thermophilic bacteriophage.

*Extremophiles* 2014; **18**: 261–270.

46. Mende DR, Bryant JA, Aylward FO, Eppley JM, Nielsen T, Karl DM, et al. Environmental

drivers of a microbial genomic transition zone in the ocean's interior. *Nat Microbiol* 2017; **2**:

1367–1373.

47. Goldsmith DB, Brum JR, Hopkins M, Carlson CA, Breitbart M. Water column stratification

structures viral community composition in the Sargasso Sea. *Aquatic Microbial Ecology* .

2015; **76**: 85–94.

48. Hurwitz BL, Brum JR, Sullivan MB. Depth-stratified functional and taxonomic niche

specialization in the 'core' and 'flexible' Pacific Ocean Virome. *The ISME Journal* 2015; **9**:

472–484.

49. Breitbart M, Rohwer F. Here a virus, there a virus, everywhere the same virus? *Trends in*

*Microbiology* 2005; **13**: 278–284.

50. Brüssow H. Huge bacteriophages: bridging the gap? *Environmental Microbiology* 2020; **22**:

1965–1970.

51. Koonin EV, Yutin N. Evolution of the Large Nucleocytoplasmic DNA Viruses of Eukaryotes

and Convergent Origins of Viral Gigantism. *Adv Virus Res* 2019; **103**: 167–202.

52. Edwards KF, Steward GF, Schvarcz CR. Making sense of virus size and the tradeoffs

shaping viral fitness. *Ecol Lett* 2021; **24**: 363–373.

53. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic

gene recognition and translation initiation site identification. *BMC Bioinformatics* 2010; **11**:

119.

54. Williams T, Kelley C. Gnuplot 5.2 Manual: An Interactive Plotting Program. 2017.

55. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile and

open software for comparing large genomes. *Genome Biol* 2004; **5**: 1-9.

56. Kiełbasa SM, Wan R, Sato K, Horton P, Frith MC. Adaptive seeds tame genomic sequence

comparison. *Genome Res* 2011; **21**: 487–493.

57. Aylward FO, Moniruzzaman M. ViralRecall-A Flexible Command-Line Tool for the Detection

of Giant Virus Signatures in 'Omic Data. *Viruses* 2021; **13**.

58. wwood. GitHub - wwood/CoverM: Read coverage calculator for metagenomics.

https://github.com/wwood/CoverM. Accessed 23 Jul 2021.

59. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J*

*Mol Biol* 1990; **215**: 403–410.

60. igraph – Network analysis software. http://igraph.org. Accessed 23 Jul 2021.

61. R Core Team. R: A Language and Environment for Statistical Computing. 2019. R

Foundation for Statistical Computing, Vienna, Austria.

62. RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA

URL http://www.rstudio.com/.

63. Wickham H. ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics* 2011; **3**:

180–185.

64. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation

sequencing data. *Bioinformatics* 2012; **28**: 3150–3152.

65. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of

high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol*

2011; **7**: 539.

66. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment

trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009; **25**: 1972–1973.

67. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015;

**32**: 268–274.

68. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast

model selection for accurate phylogenetic estimates. *Nat Methods* 2017; **14**: 587–589.

69. Dixon P. VEGAN, a package of R functions for community ecology. *Journal of Vegetation

Science* 2003; **14**: 927–930.

70. maps: Draw Geographical Maps. https://CRAN.R-project.org/package=maps. Accessed 22

Jul 2021.

71. Kassambara A. 'ggplot2' Based Publication Ready Plots [R package ggpubr version 0.4.0].

2020.

72. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al.

Pfam: The protein families database in 2021. *Nucleic Acids Res* 2020; **49**: D412–D419.

73. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, et al.

eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource

based on 5090 organisms and 2502 viruses. *Nucleic Acids Research* 2019; **47**: D309–

D314.

74. pheatmap: Pretty Heatmaps. https://CRAN.R-project.org/package=pheatmap. Accessed 22

Sep 2021.

75. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, et al. A

standardized bacterial taxonomy based on genome phylogeny substantially revises the tree

of life. *Nat Biotechnol* 2018; **36**: 996–1004.

76. Bland C, Ramsey TL, Sabree F, Lowe M, Brown K, Kyrpides NC, et al. CRISPR

Recognition Tool (CRT): a tool for automatic detection of clustered regularly interspaced

palindromic repeats. *BMC Bioinformatics* 2007; **8**: 1-8.

77. Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of

transfer RNA genes. *Nucleic Acids Res* 2016; **44**: W54–7.

78. Paez-Espino D, Eloe-Fadrosh EA, Pavlopoulos GA, Thomas AD, Huntemann M, Mikhailova

N, et al. Uncovering earth's virome. *Nature* 2016; **536**: 425–430.

**Figure Legends**

**Figure 1. Comparison of bins to reference gene content and evolutionary histories. (a)**

Network with marine jumbos and references as nodes and edges based on shared VOGs.

Marine jumbo phage nodes are colored by PGC as detected with spinglass community

detection analysis, other nodes are in gray. (**b**) Barchart of the number of members in each

PGC. PGCs with marine jumbo phages are denoted with a star and the number of marine jumbo

phages in that PGC. Proportion of marine jumbo phages in that PGC is colored. (**c,d**)

Phylogenies of TerL (**c**) and MCP (**d**) proteins with references and bins. Inner ring and branches

are colored by the 5 PGCs that marine jumbo phages belong to. Navy blue circles in the outer

ring denote marine jumbo phages.

**Figure 2. (a)** Boxplot of genome length in each network cluster (x-axis is PGC number). Star

denotes PGC with a marine jumbo phage and the color matches the PGC letters of Figure 1. **(b)**

Stacked barplot of the metagenome environment from which each phage derives from in each

PGC (x-axis). Reference (yellow) are cultured phages, black is the bins of jumbo phages from

this study **(c)** Stacked barplot of the host phylum of the RefSeq cultured phages in each cluster;

metagenomic phages are in gray.

**Figure 3** (**a**) Functional categories for genes encoded by jumbo phages. (**b**) Heatmap of

proportion of genomes in each cluster that contain the listed genes. Listed genes were selected

based on containing a known function and variance between the clusters above 0.2.

Dendrogram was generated based on hierarchical clustering in pheatmap.

**Figure 4. Comparison of jumbo abundance and presence in samples of different filter**

**size fractions**. Dark teal are fractions with minimum sizes of 0.22 μm or higher. Light teal are

fractions with a maximum size of 0.22 μm or lower. **a)** Barchart of the proportion of samples with

at least one marine jumbo phage (x-axis) by size fraction (y-axis) sorted from highest to lowest.

**b)** Boxplot with x-axis as the number of marine jumbo phages found in a sample with size

fraction on the y-axis sorted by median. **c)** Boxplot with x-axis as the relative abundance of

marine jumbo phages found in a sample (RPKM) with size fraction on the y-axis sorted by

median. Significance bars in c,d correspond to Wilcox test, with stars corresponding to p-values

< 0.05 and those with p-values > 0.05 as not significant "ns" (stat_compare_means

function).  (**d,e**) NMDS plots (Bray Curtis dissimilarity distances) of jumbo phage composition in

each sample using presence absence data (**d**) and relative abundance data (**e**). Samples are

colored by size fraction distinction above 0.22 (dark teal) and below 0.22 (light teal). Ellipses

calculated based on multivariate normal distribution. Samples above and below 0.22 were not

significantly different from each other using ANOSIM (p-values > 0.05).

715
**Figure 5. a,b**) Maps of the relative abundance (**a**) of jumbo phages (in RPKM) and (**b**) number

of jumbo populations present in each surface (SRF) sample of the picoplankton size fraction

(either 0.22-3μm or 0.22-1.6μm depending on availability). Dots sizes are proportional to the

number of populations or RPKM and colored by biome (Coastal - pink, Westerlies - purple,

Trades - blue). (**c**) Scatterplot of the mean RPKM of a jumbo population in SRF picoplankton

samples versus the number of SRF picoplankton stations it was present. Populations are

colored by PGC and size corresponds to putative genome length in 100 kilobases. (**d**) Boxplot

of the number of jumbo phage populations in a sample separated by depth sorted by median for

each PGC. Significance bars correspond to Wilcox test, with stars corresponding to p-values <

0.05 and those with pvalues > 0.05 as not significant (stat_compare_means function)