Infection strategy and biogeography distinguish cosmopolitan groups of marine jumbo bacteriophages

Alaina R. Weinheimer[1], Frank O. Aylward[1,2]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA. [2] Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0913

Correspondence to email: alainarw@vt.edu; address: 5092 Derring Hall, 926 W Campus Drive, Blacksburg, VA 24061; telephone: (540) 231-8657

Competing Interests: The authors declare no competing interests.
Funding Support: This work was supported by grants from the Institute for Critical Technology and Applied Science, the NSF (IIBR-1918271) and a Simons Early Career Award in Marine Microbial Ecology and Evolution to F.O.A. ARW was supported by an ICTAS Doctoral Scholars Fellowship.

Infection strategy and biogeography distinguish cosmopolitan groups of marine jumbo bacteriophages

Alaina R. Weinheimer[1], Frank O. Aylward[1,2]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA. [2] Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0913

Correspondence to alainarw@vt.edu

Keywords: Jumbo bacteriophage, viral diversity, viral gigantism, marine microbiology, microbial oceanography

ABSTRACT

Recent research has underscored the immense diversity and key biogeochemical roles of large DNA viruses in the ocean. Although they are important constituents of marine ecosystems, it is sometimes difficult to detect these viruses due to their large size and complex genomes. This is true for "jumbo" bacteriophages, which have genome sizes >200 kbp and large capsids reaching up to 0.45 µm in diameter. In this study, we sought to assess the genomic diversity and distribution of these bacteriophages in the ocean by generating and analyzing jumbo phage genomes from metagenomes. We recover 85 marine jumbo phages that ranged in size from 201-498 kilobases, and we examine their genetic similarities and biogeography together with a reference database of marine jumbo phage genomes. By analyzing Tara Oceans metagenomic data, we show that although most jumbo phages can be detected in a range of different size fractions, 17 of our bins tend to be found in those greater than 0.22 um, potentially due to their large size. Our network-based analysis of gene sharing patterns reveals that jumbo bacteriophages belong to five genome clusters that are typified by diverse replication strategies, genomic repertoires, and potential host ranges. Our analysis of jumbo phage distributions in the ocean reveals that depth is a major factor shaping their biogeography, with some phage genome clusters occurring preferentially in either surface or mesopelagic waters, respectively. Taken together, our findings indicate that jumbo

phages are widespread community members in the ocean with complex genomic repertoires and

ecological impacts that warrant further targeted investigation.

Introduction

Although historically noted for their small virion sizes and simple genomes [1], viruses with large

particles and elaborate genomes have been discovered in recent decades throughout the

biosphere [2–4]. These complex viruses not only invite intriguing evolutionary questions [5–7], but

also expand the potential roles viruses have in shaping microbial community structure and

biogeochemical cycling [4, 8–10]. One group of these larger viruses are jumbo bacteriophages

(jumbo phages), which have traditionally been defined as *Caudovirales* with genomes over 200

kilobases in length [11]. While a recent survey of cultured jumbo phages showed jumbo phages

share some universal features and genes, such as encoding DNA polymerases and the terminase

large subunit (TerL), these features do not distinguish them from smaller phages, and several

lines of evidence suggest that jumbo phages emerged from smaller phages multiple times

independently [6]. For example, a recent phylogenetic study of cultured *Caudovirales* used

concatenated protein alignments to generate phylogenies and found that the most supported

clades within the *Caudovirales* family do not consistently correspond to genome length [12].

Furthermore, a previous study found that jumbo phages cluster with smaller phages based on

gene content and are best grouped by replication machinery, among other infection apparati [6].

Taken together, jumbo phages likely emerged from diverse, smaller phages multiple times

independently and form distinct clades within the *Caudovirales*.

Although the first jumbo phages were isolated as early as the 1970s [13], these viruses have

remained relatively sparse in culture, representing less than 3% (n=93) of complete *Caudovirales*

genomes on NCBI's RefSeq Viral Genome Portal (downloaded July 5, 2020) and 2.2% of the

INPHARED database [14]. All cultured jumbo phage capsids have morphologies of myoviruses

or siphoviruses, and they undergo infection cycles that reflect temporal patterns of lytic

Caudovirales [15,16]. Some jumbo phages are known to stall infections resulting in

"pseudolysogeny", however, which has been proposed as a competitive strategy against other

phages to prevent superinfection [6]. Jumbo phages that have been studied extensively are

primarily investigated for their unusually complex functional capabilities, such as encoding entire

transcriptional apparatti [17] or sophisticated anti-CRISPR defense mechanisms [18, 19].

Regarding their ecological range, cultivated jumbo phages have been isolated on both Gram

negative and Gram positive bacteria [15], and a recent metagenomic survey uncovered these

viruses in diverse, global ecosystems [4].

Despite this apparent broad environmental distribution, common methods for viral isolation and

diversity surveys often bias against the inclusion of jumbo phages. Because viruses have

historically been considered smaller than cells, many viral diversity surveys specifically examine

only small particle sizes. For example, in plaque assays, agar concentrations are often too high

for larger phage particles to diffuse through compared to smaller particles [20]. Moreover, filters

are often used to remove cells when preparing viral enrichments for metagenomic sequencing

[21], which excludes larger viruses [9, 22]. Particularly in marine studies, the <0.22 μm fraction,

sometimes even referred to as the "viral fraction" [23], is most commonly examined for viruses

[24–26]. Jumbo phages can have particles over 0.45 μm in length (i.e. *Pseudomonas aeruginosa*

phage PhiKZ) [13], however, and will therefore be excluded from <0.22 μm size fractions. Lastly,

in bioinformatic pipelines, phage sequences are often only assembled to the contig or scaffold

level, which is sometimes sufficient for the assembly of most known smaller phage genomes [27],

but often leaves larger phage genomes fragmented into multiple contigs and may require

additional joining of contigs into bins [28]. Overall, considering these biases and the recently-

discovered broad distribution of these viruses [4], jumbo phages may represent underappreciated

components of marine microbial communities and food webs that warrant further examination.

In this study, we examine the diversity and prevalence of jumbo phages in the global ocean. We develop a workflow for generating and validating high-quality jumbo bacteriophage bins from metagenomic data with which we identify 85 bins of jumbo phages. We then compare the genetic content of these jumbo phages with other cultured phages of all sizes and metagenomic jumbo phages from other studies. We find that the jumbo phages of this study group into five distinct clusters that are distinguished by diverse replication machinery and infection strategies, implicating a broad range of potential jumbo-phage host interactions in the ocean. We then assess the distribution of jumbo phages belonging to these genome clusters in the ocean by using metagenomic data from Tara Oceans [29]. Mapping the Tara Oceans metagenomic data onto the jumbo phage sequences reveals that these jumbos phages are collectively widely distributed in the ocean, but vary in biogeography both within and between clusters, with some more enriched in surface waters relative to deeper waters and vice versa. Upon examining the collective presence of jumbo phages in different filter fractions, we also find that most could be detected in a range of different size fractions, although 34 were recovered from only >0.22 um fractions. Our results support the view that jumbo phages are widespread in the biosphere and may play underappreciated roles in ecosystems around the globe.

RESULTS AND DISCUSSION

Detection and validation of high-quality jumbo phage bins

Due to the large size of jumbo bacteriophage genomes, it is likely that they are present in multiple distinct contigs in metagenomic datasets and therefore require binning to recover high-quality metagenome-assembled genomes (MAGs) [28]. This has been shown for large DNA viruses that infect eukaryotes, where several recent studies have successfully employed binning approaches to recover viral MAGs [2, 3]. Here, we used the same 1,545 high-quality metagenomic assemblies [30] used in a recent study to recover giant viruses of eukaryotes [3], but we modified the bioinformatic pipeline to identify bins of jumbo bacteriophages. These metagenomes were

compiled by Parks et al. 2017 [30] and included all available metagenomes on the NCBI's Short

Read Archive by December 31, 2015 (see Parks et al. 2017 [30]). This dataset includes a wide

variety of marine metagenomes (n=469) with many non-Tara metagenomes (n=165). We focused

our benchmarking and distribution analyses on Tara data [29] because of the well-curated

metadata and size fractions in this dataset. We first binned the contigs from these assemblies

with MetaBat2 [31], which groups contigs together based on similar nucleotide composition and

coverage profiles and focused on bins of at least 200 kilobases in total length. We subsequently

identified bins composed of bacteriophage contigs through analysis with VirSorter2 [32],

VIBRANT [33], and CheckV [34] (see Methods for details).

The occurrence of multiple copies of highly conserved marker genes is typically used to assess

the level of contamination present in metagenome-derived genomes of bacteria and archaea [35].

Because bacteriophage lack these marker genes, we developed alternative strategies to assess

possible contamination in our jumbo phage bins. Firstly, we refined the set of bins by retaining

those with no more than 5 contigs that were each at least 5 kilobases in length to reduce the

possibility that spurious contigs were put together. Secondly, we assessed the possibility that two

strains of smaller phages with similar nucleotide composition may be binned together by aligning

the contigs in a bin to each other. Bins that had contigs with high sequence similarity across the

majority of their lengths were discarded (Supplemental Figure 1). Thirdly, we discarded bins if

their contigs exhibited aberrant co-abundance profiles in different metagenomes (see

Supplemental Methods). To generate these co-abundance profiles, we mapped reads from 225

marine metagenomes provided by Tara Oceans [29] onto the bins. Coverage variation between

contigs was benchmarked based on read-mapping results from artificially-fragmented reference

genomes present in the samples (See Methods for details). Only bins with coverage variation

below our empirically-derived threshold were retained. Using this stringent filtering, we identified

85 bins belonging to jumbo bacteriophages. These bins ranged in length from 202 bp to 498 kbp,

and 31 consisted of a single contig, while 54 consisted of 2-5 contigs (Supplemental Figure 2).

To assess global diversity patterns of jumbo bacteriophages, we combined these jumbo phage

bins together with a compiled database of previously-identified jumbo phages that included all

complete jumbo *Caudovirales* genomes on RefSeq (downloaded July 5th, 2020), the INPHARED

database [36], the cultured jumbo phage survey [6], the Al-Shayeb et al. 2020 study [4], and

marine jumbo phage contigs from metagenomic surveys of GOV 2.0 [26] (60 jumbo phages),

ALOHA 2.0 [37] (8 jumbo phages), and one megaphage MAG recovered from datasets of the

English Channel [38]. Ultimately, we arrived at a set of 244 jumbo phages, including the 85 bins,

that were present in at least one Tara Oceans sample (min. 20% genome covered, see Methods)

or deriving from a marine dataset (i.e. ALOHA, GOV 2.0) which we analyzed further in this study

and refer to as marine jumbo phages. Statistics on genomic features can be found in

Supplemental Dataset 1.

Marine jumbo phages belong to distinct groups with diverse infection strategies

Because bacteriophages lack high-resolution, universal marker genes for classification, such as

16S rRNA in bacteria, phages are often grouped by gene content [39, 40]. Here, we generated a

bipartite network that included the 85 bins of jumbo phages with a dataset of available

Caudovirales complete genomes in RefSeq (3,012 genomes; downloaded July 5th, 2020) and

the full set of reference jumbo phages described above. To construct the bipartite network, we

compared proteins encoded in all the phage genomes to the VOG database (vogdb.org), and

each genome was linked to VOG hits that were present (Figure 1, Supplemental Dataset 2, see

Methods for details). To identify groups of phage genomes with similar VOG profiles, we

employed a spinglass community detection algorithm [41] to generate genome clusters. Similar

methods have recently been used to analyze evolutionary relationships in other dsDNA viruses

[40]. The marine jumbo phages of this study clustered into five groups that included both jumbo

and non-jumbo phage genomes (Figure 2a). We refer to these five clusters as Phage Genome

Clusters (PGCs): PGC_A, PGC_B, PGC_C, PGC_D, and PGC_E. These PGCs included cultured

phages and metagenome-derived jumbo phages found in various environments (i.e. aquatic,

engineered) and isolated on a diversity of hosts (i.e. Firmicutes, Proteobacteria, Bacteroidetes)

(Figure 2b,c). Of the marine jumbo phages, 135 belonged to PGC_A, 11 to PGC_B, 90 to PGC_C,

7 to PGC_D, and 1 to PGC_E (Figure 1b). In addition to this network-based analysis, we also

examined phylogenies of the major capsid protein (MCP) and the terminase large subunit (TerL)

encoded by the marine jumbo phages and the same reference phage set examined in the network

(Figure 1c, 1d). With the exception of PGC_A, the marine jumbo phages that belong to the same

PGC appeared more closely related to each other than those belonging to different clusters. The

polyphyletic placement of jumbo phages in these marker gene phylogenies is consistent with the

view that genome gigantism evolved multiple times, independently within the *Caudovirales* [6].

We then compared functional content encoded by the marine jumbo phages in the PGCs to

identify functional differences that distinguish these groups. PGC_E was excluded from this

analysis because this genome cluster contained only a single jumbo phage. Collectively, most

genes of the marine jumbo phages could not be assigned a function (mean: 86.60%, std dev:

7.01%; Supplemental Dataset 3), which is common with environmental viruses [42, 43]. Genes

with known functions primarily belonged to functional categories related to viral replication

machinery, such as information processing and virion structure (Figure 3a), and these genes

drove the variation between the genome clusters of marine jumbo phages (Figure 3b). A recent

comparative genomic analysis of cultivated jumbo phages was able to identify three types of

jumbo phages that are defined by different infection strategies and host interactions (referred to

as Groups 1-3) [6]. We cross-referenced our PGCs and found that PGCs B, C, and D of this study

corresponded to Groups 1, 2, and 3, respectively, suggesting that these genome clusters contain

phages with distinct infection and replication strategies. PGC_A corresponded to multiple groups,

indicating that this genome cluster contains a particularly broad diversity of phages.

The second largest phage cluster with marine jumbo phages, PGC_B, consists of 238 phages

(11 (4.6%) marine jumbo phages, including 10 bins generated here), and included cultured

phages of Group 1, which is typified by *Pseudomonas aeruginosa* phage PhiKZ. Supporting this

correspondence with Group 1, all marine jumbo phages of PGC_B encoded the same distinct

replication and transcription machinery, including a divergent family B DNA polymerase and a

multi-subunit RNA polymerase (Figure 3b, Supplemental Dataset 3). These marine jumbo phages

also encoded a PhiKZ internal head protein, and they uniquely encoded shell and tubulin

homologs which has recently been found in PhiKZ phages to assist in the formation of a nucleus-

like compartment during infection that protects the replicating phage from host defenses [18, 19].

Although we could not confidently predict hosts for the 11 metagenomic marine jumbo phages in

this PGC_B (Supplemental Dataset 1), the cultured phages of this genome cluster infect

pathogenic bacteria belonging to the phyla Proteobacteria (178 phages) and Firmicutes (6

phages) (Figure 2c), implicating a potential host range for marine jumbo phages in PGC_B.

The next largest phage genome cluster, PGC_C, comprised of 156 phages total (90 marine jumbo

phages (57.7%); 4 bins generated from this study) and included reference jumbo phages in Group

2 (31, 19.9%) which are typified by Alphaproteobacteria and Cyanobacteria phages. Likewise,

the host range of other cultured phages in PGC_C support the Group 2 correspondence, either

infecting Cyanobacteria (139 phages) or Proteobacteria (4 phages) (Figure 2c). Furthermore, all

3 marine metagenomic phages in PGC_C for which hosts could be predicted were matched to

Cyanobacteria hosts (Supplemental Dataset 1). Functional annotations of PGC_C marine jumbo

phages revealed nearly all encode a family B DNA polymerase (97.8% phages) and the

photosystem II D2 protein (PF00124, VOG04549) characteristic of cyanophages (90% phages)

(Figure 3b). This PGC included the reference *Prochlorococcus* phage P-TIM68 (NC_028955.1),

which encodes components of both photosystem I and II as a mechanism to enhance cyclic

electron flow during infection [44]. This suggests that an enhanced complement of genes used to

manipulate host physiology during infection may be a driver of large genome sizes in this group.

Additionally, most of the PGC_C marine jumbo phages encoded lipopolysaccharide (LPS)

biosynthesis proteins (76%), which have been found in cyanophage genomes that may induce a

"pseudolysogeny" state, when infected host cells are dormant, by changing the surface of the

host cell and preventing additional phage infections [6] (Supplemental Dataset 3). Taken together,

most marine jumbo phages of PGC_C likely follow host interactions of jumbo cyanophages, such

as potentially manipulating host metabolism by encoding their own photosynthetic genes and

potentially inducing a pseudolysogenic state.

Finally, phages of PGC_D totaled at 47 phages, of which 7 were marine jumbo phages generated

in this study (14.9%). This group included Group 3 jumbo phages (15, 31.9%), which is primarily

distinguished by encoding a T7-type DNA polymerase but is not typified by a particular phage

type or host range. Supporting this grouping, all marine jumbo phages in this study encoded a T7

DNA polymerase which belongs to family A DNA polymerases (Figure 3b, Supplemental Dataset

3). Most of the other genes distinctively encoded by the marine jumbo phages in this group

included structural genes related to T7 (T7 baseplate, T7 capsid proteins), a eukaryotic DNA

topoisomerase I catalytic core (PF01028), and DNA structural modification genes (MmcB-like

DNA repair protein, DNA gyrase B). Hosts of metagenomic marine jumbo phages in PGC_D could

not be predicted (Supplemental Dataset 1); however, cultured Group 3 jumbo phages in PGC_D

all infect Proteobacteria, primarily Enterobacteria and other pathogens.

The largest of the phage genome clusters, PGC_A, contained 469 phages, including 135 marine

jumbo phages (63 bins from this study). This genome cluster contained the largest jumbo phages,

such as *Bacillus* phage G (498 kb) and the marine megaphage Mar_Mega_1 (656 kb) recently

recovered from the English Channel [38]. Unlike other PGCs, PGC_A contained mostly

metagenomic phages (401, 85%, Figure 2b,c). Considering PGC_A contains the largest jumbo

phages (Figure 1b, 2a), the vast genetic diversity in this PGC might explain why few genes were

found to distinguish this group. Of the genes unique to PGC_A, only one was present in at least

half of the phages (51.9%), which was a Bacterial DNA polymerase III alpha NTPase domain

(PF07733). The host ranges of cultured phages from this PGC further reflect the large diversity

of this group and included a variety of phyla and genera that can perform complex metabolisms

or lifestyles, such as the nitrogen-fixing Cyanobacteria of the *Nodularia* genus isolated from the

Baltic Sea (accessions NC_048756.1 and NC_048757.1) and the Bacteroidetes bacteria

Rhodothermus isolated from a hot spring in Iceland (NC_004735.1) [45].

Relative abundance of jumbo bacteriophages across size fractions

To explore the distribution of the marine jumbo phages in the ocean, we first examined the size

fractions in which the jumbo phages were most prevalent. To remove redundancy, we examined

the 244 jumbo phage the population-level (>80% genes with >95% average nucleotide identity

[24]), corresponding to 142 populations (11 unique to this study, corresponding to 47 bins). We

then mapped Tara Oceans metagenomes onto the 142 jumbo phage populations, and 102 of

these populations could be detected [min. 20% of genome covered], resulting in 74 populations

in PGC_A, 2 in PGC_B, 22 in PGC_C, 3 in PGC_D, and 1 in PGC_E. Out of the 225 Tara Oceans

metagenomes examined, 213 (94.6%) contained at least one jumbo phage population (median:

7, Supplemental Dataset 4). Jumbo phages were more frequently detected in samples below 0.22

µm (<-0.22 µm, 0.1-0.22 µm) than those above 0.22 µm (0.45-0.8 µm, 0.22-0.45 µm, 0.22-1.6

µm, 0.22-3 µm) (Figure 4A). All samples in the <-0.22 µm fraction and the 0.1-0.22 µm fraction

had at least one jumbo phage present, while the larger fractions ranged from 89% to 97%.

Interestingly, we detected 34 populations (33.3%) exclusive to samples above 0.22 um, compared

to only one population (0.98%) exclusive to samples below 0.22 µm. A similar disparity in virus

detection between size fractions has been reported for large eukaryotic viruses, where roughly

41% of phylotypes were present in the 0.22-3 µm size fraction but absent in fractions below 0.22

µm [9]. In contrast to this study, where certain viral groups were more prevalent in larger size

fractions than smaller, a jumbo phage's PGC membership or genome size generally did not affect

its probability of detection at different size fractions (Supplemental Figure 3, Supplemental Figure

4).

We also compared jumbo phage diversity (defined as population richness), relative abundance

(calculated in reads per kilobase per million (RPKM)), and community composition between the

size fractions (based on Bray-Curtis distance matrices). Collectively, samples of the size fractions

below 0.22 µm were significantly more diverse (*p value* <0.0001, Wilcox test) and had significantly

higher relative abundances (*p value* < 0.0001, Wilcox test) of jumbo phages relative to the size

fractions above 0.22 µm. Despite these differences in diversity and relative abundances, jumbo

phage community composition did not significantly differ between the >0.22 and <0.22 µm size

fractions when comparing samples based on presence/absence data (*p value* = 0.1082, ANOSIM,

presence/absence Bray-Curtis distance matrix, Figure 4d), but did differ when using relative

abundance data (*p value* = 0.0001, ANOSIM, RPKM Bray-Curtis distance matrix, Figure 4e).

To directly test the effect of the 0.22 µm size fraction cut-off on jumbo phage recovery, we

examined a subset of the samples that were co-collected at the same station and depth for the

fractions below 0.22 µm (<-0.22 or 0.1-022) and above 0.22 µm (0.22-1.6 µm or 0.22-3 µm). The

number of detected jumbo phage populations was significantly higher in samples below 0.22 µm

than above 0.22 µm (*p value* = 0.000138, Wilcox test, Supplemental Figure 5a). The relative

abundance of jumbo phages was also significantly higher in size fractions below 0.22 µm than

above 0.22 µm (*p value* = 0.00001, Wilcox test, Supplemental Figure 5b). Likewise, community

composition significantly differed between samples above and below 0.22 µm (*p value* = 0.0001,

ANOSIM, presence/absence Bray-Curtis distance matrix, Supplemental Figure 5c,d). Taken

together, these findings suggest that using size fractions below 0.22 µm to analyze phages

enhances the signal of jumbo phage sequences, relative to samples of larger size fractions, likely

due to cellular sequences present in the larger sizes. Notwithstanding, roughly 33% of jumbo

phages in this study were exclusive to size fractions above 0.22 µm, indicating that analyzing a

range of size fractions is necessary for a more synoptic view of jumbo phages in the environment.

Biogeography of jumbo bacteriophages in the global ocean

Jumbo phage populations varied by depth in different PGCs. Jumbo populations in this study

varied in their distribution (Supplemental Figure 6) but were collectively found in all three depths

(Surface (SRF), Deep Chlorophyll Maximum (DCM), Mesopelagic (MES)) examined (Figure 5,

Supplemental Figure 7; Supplemental Dataset 4). Although jumbo phages were more prevalent

in samples of the viral size fractions (<-0.22 or 0.1-0.22 µm), we focused biogeographic analyses

on the 0.22-1.6 or 0.22-3 µm size fractions because the most sites were available for these

samples enabling comparisons between depths and biomes. For instance, only four MES

samples and no Westerlies samples were available in the viral fractions. When applicable,

analyses were also completed with viral fraction samples, and results are deposited at the end of

the Supplement (Supplemental Figures 16 - 24). Jumbo phage communities differed significantly

between depths (*p value* = 0.0001, ANOSIM based on presence/absence and RPKM Bray-Curtis

distance matrices, Supplemental Figure 8), consistent with the dramatic transition in community

composition that occurs from surface waters to below the deep chlorophyll maximum [37, 46].

Specifically, the diversity of jumbo phages across depths varied by genome cluster (Figure 5d,

Supplemental Figure 10), with PGC_A and PGC_C exhibiting higher prevalence in the epipelagic

(SRF and DCM). Although PGC_B and PGC_D had too few populations detected to generalize

for these clusters (2 and 3, respectively), our results for these phages showed that PGC_B and

PGC_D were typically less prevalent in SRF samples compared to DCM and MES samples.

PGC_C is typified by cyanophages, providing a clear reason why this phage group is enriched in

surface waters. Conversely, PGC_B is typified by *Pseudomonas aeruginosa* PhiKZ phages,

suggesting these PGC_B marine jumbo phages may be infecting heterotrophic bacteria that are

potentially less prevalent in surface waters. Overall relative abundance and diversity of jumbo

phages in this study were significantly higher in the epipelagic zone (Supplemental Figure 8),

partly because most of these phages are in PGC_A. These collective and PGC-specific patterns

held when examining only those samples that were co-collected at all three depths (Figure 5c,

Supplemental Figures 8, 9, 10). This general pattern therefore reflects what has been found in

previous studies on the depth distribution and viral protein clusters, where more were unique in

the euphotic (i.e. epipelagic) than aphotic depths [9, 26, 47, 48], although this contrasts what has

recently been found in the Pacific Ocean, where overall viral diversity increased in the

mesopelagic [37].

Jumbo phage biogeography across biomes. Collectively, jumbo phages could be found in all three

Longhurst biomes (Coastal, Westerlies, Trades) (Supplemental Figure 11), and jumbo phage

communities in this study significantly differed in composition between the biomes (*p value* < 0.05,

ANOSIM, presence/absence Bray-Curtis distance matrix, Supplemental Figure 11c,). However,

no biome appeared to be a particular hotspot for jumbo phages, as they were not significantly

more diverse in any biome (Supplemental Figure 11a). When looking at depths separately, the

relative abundance of jumbo phages in SRF samples was significantly higher in Coastal samples

(Supplemental Figure 12b), but no clear biome was a hotspot for phages in DCM and MES

samples (Supplemental Figure 11). Likewise, jumbo phage community composition only differed

between samples in the SRF and DCM, but not in MES samples (*p value* cutoff < 0.05, ANOSIM,

presence/absence Bray-Curtis distance matrices). Upon examining jumbo phages by group,

jumbo phages from PGCs A, C, and D could be detected in all biomes, while PGB_B phages

could not be detected in Westerlies samples (Supplemental Figure 13). Similar to their collective

results, no PGC was enriched in a single biome (Supplemental Figure 13). A recent global study

on marine viruses has found that viral diversity is better explained by ecological zones defined by

physicochemical factors like temperature, rather than by Longhurst biomes defined by patterns of

chlorophyll *a* concentrations [26], suggesting that Longhurst biomes may not be good predictors

of viral diversity in general.

Jumbo phage populations ranged in endemicity. Fifteen populations (14.7%) were detected in

only one Tara station, and six (5.6%) were present in over half of the stations (>=34) (Figure 5c).

Both the more endemic populations and the more prevalent jumbo phages belonged to PGCs A

and C. PGC_A and PGC_C contained the most populations, which likely explains the wide range

of endemicity of phages in these clusters (Supplemental Figures 14, 15). Moreover, the

cyanobacterial hosts that are known for many of the jumbo phages in PGC_C are widespread in

the ocean, which may also explain the prevalence of this group of phages. In general, the

heterogeneous distribution and abundance of these jumbo phages is consistent with the seed

bank hypothesis, which postulates that viruses are passively dispersed throughout the ocean and

viral community structure is shaped by local selective forces [24, 49]. This framework has

previously been used to explain why phage distributions range from extremely cosmopolitan to

extremely rare, which is a pattern that also appears to hold for jumbo bacteriophages.

CONCLUSION

Large DNA viruses are becoming increasingly recognized as critical components of the

virosphere, notable for their intriguing evolutionary histories [50, 51], vast functional capacities [3,

4], and global distribution [2, 4]. Here, we uncover the diversity and ecology of marine jumbo

bacteriophages, which have historically been difficult to study due to biases in filtration and

isolation strategies commonly used in virology. We employed a binning strategy to generate and

quality-check genomes of jumbo phages and used it to identify 85 high quality bins. We employed

a conservative approach to genome binning because binning has traditionally not been used for

bacteriophage, and as a result these bins likely represent a small fraction of total jumbo phages

in these marine samples. We combined these bins together with reference jumbo bacteriophage

genomes, and ultimately identified 102 populations that are present in Tara Oceans

metagenomes. When compared with other metagenomic jumbo phages and cultured phages of

all sizes, we found that marine jumbo phages primarily belong to four phage genome clusters

(PGCs) that encode distinct replication machinery, biogeography, and potential hosts. For

example, marine jumbo phages in PGC_C follow cyanophage infection strategies and ecology,

as this cluster included cultured marine cyanophages and encoded classic family B DNA

polymerases and photosynthesis enzymes characteristic of cyanophages. Furthermore, we found

they are enriched in surface waters relative to the mesopelagic, consistent with the geographic

range of their hosts. In contrast, marine jumbo phages of PGC_B included cultured PhiKZ phages

of *Pseudomonas aeruginosa* and uniquely encoded multi-subunit RNA polymerases and tubulin,

which are thought to play a role in the remarkable nucleus-like structures that these viruses

employ as an anti-CRIPSR defense [18, 19]. PGC_B was particularly enriched in mesopelagic

waters, suggesting that this complex infection strategy may be more common in the deep ocean.

These results demonstrate the diverse biology and ecology of jumbo phages, consistent with the

hypothesis that jumbo phages are an incredibly diverse set of phages with unique evolutionary

histories [6].

The jumbo phages we analyze are collectively widespread throughout the ocean and are typically

more diverse and abundant in epipelagic waters, which reflect previous findings that surface

waters usually harbor a higher per-sample alpha diversity of viral groups compared to deeper

waters [9, 26]. Larger phages therefore appear to coexist in patterns broadly similar to smaller

viruses despite the disadvantages of their size, such as smaller burst sizes and lower host contact

rates [52]. In eukaryotic giant viruses, it has been hypothesized that these disadvantages are

potentially offset by higher infection efficiency, broader host ranges, decreased decay rates, and

higher rates of successful attachments compared to smaller viruses [52]. Although some of these

advantages to viral gigantism may also apply to jumbo bacteriophages, it is unlikely that they are

all applicable. For example, given that they are tailed *Caudovirales* [6], jumbo phages likely

possess higher host specificity in part due to their non-phagocytotic mode of infection.

Nonetheless, the large genomes of jumbo bacteriophage often encode an expanded complement

of genes used to manipulate host physiology during infection, and these may play critical roles in

promoting infection efficiency or offsetting host defense mechanisms. The impressive

complement of photosynthesis genes in PGC_C is at least partially responsible for the large

genomes in this lineage, while the genes involved in anti-CRISPR defense found in PGC_B

indicate that a host-virus arms race may be responsible for genome gigantism in this group.

Interestingly, the largest number of jumbo phage genomes we identified belong to PGC_A, which

is largely uncharacterized and composed of primarily metagenome-derived genomes, suggesting

that these viruses have as-yet unidentified infection strategies. Overall, it is likely that the factors

leading to genome gigantism in each of these genome clusters are distinct. Future work further

characterizing the hosts of these jumbo phages and the details of their infection program,

particularly in PGC_A, will therefore be critical to understanding mechanisms that underlie

complexity in the virosphere and maintain diversity. Moreover, future in-depth examination of the

genomics and evolutionary histories of jumbo phages will be an important step to integrating these

viruses into a meaningful taxonomy and clarifying their evolutionary relationships to other

Caudovirales.

Methods Summary

Jumbo phage binning and detection. Overview of pipeline can be found in Supplemental Figure

25. Metagenomic scaffolds were downloaded from 1,545 assemblies by Parks et al. 2017 [30]

and binned with MetaBAT2 [31] (-s 200000 --unbinned -t 32 -m 5000 --minS 75 --maxEdges 75)

using the coverage files provided by Parks et al. 2017. Bins were retained if they summed to at

least 200,000 base-pairs and comprised <= 5 contigs (min. contig size 5kb). Proteins were

predicted with Prodigal [53] using default settings on each bin individually. Bins were retained if

they lacked more than one ribosomal protein, lacked overlapping regions (via promer and gnuplot

[54] with MUMmer 3.0 [55]), had fewer hits to NCLDV than phage (via LASTp [56] against RefSeq

r99), lacked more than one NCLDV marker gene (via hmmsearch (hmmer.org) against NCLDV

marker gene HMM profiles [57]) and had even read coverage of Tara Ocean metagenomes via

coverM [58] (https://github.com/wwood/CoverM). Briefly, "even read coverage" means that the

read coverage of each contig in the bin varied between one another below a variation threshold

determined by reference mapping results (See Supplemental Methods for details). Jumbo phages

were then detected by running the bins through VirSorter2 [32], VIBRANT [33], and CheckV [34].

Bins were considered putative phages if they had at least an average dsDNAphage score of >0.9

from VirSorter2 or a VirSorter2 average score >0.5 and either i) CheckV quality of medium or

higher or ii) VIBRANT consensus classification as viral. Ultimately, 85 bins were retained for

downstream analyses. Prior to further gene-based analyses, we checked if the jumbo bins used

alternative genetic codes with Codetta [59], and all were found to use the standard bacteria code

11; we therefore proceeded with the initial Prodigal predictions. Bins were grouped into

populations based on single-linkages with a compiled set of 898 jumbo bacteriophages (RefSeq

phages over 200 kilobases (93), phage sequences over 200 kilobases from the INPHARED

database (354) [36], Iyer et al 2021 (46, non-overlapping with INPHARED) [6], Al-Shayeb et al.

2020 [4] jumbo phage genomes (336), GOV 2.0 (60) [26], ALOHA 2.0 (8) [37], and a megaphage

assembled from the English Channel [38]) based on nucleotide sequences of genes (predicted

with prodigal -d flag) aligned with BLASTn [60] (>95% average nucleotide identity, >80% genes)

[24]. See Supplemental Methods for details.

Bipartite network and phylogenetic analyses. Because phages lack universal, high resolution

phylogenetic marker genes, gene-sharing networks have typically been used to classify phages

[39, 40]. Bipartite networks are commonly used to examine evolutionary relationships in divergent

viral lineages [3, 40, 61]. To classify the jumbo bins of this study with a bipartite network, reference

phage sequences were compiled from RefSeq's *Caudovirales* complete genomes (downloaded

July 2020 from NCBI's Virus genome portal; 3,012 genomes) along with the curated jumbo phage

set used in the population analysis. Proteins of jumbo bins and this reference set were predicted

with Prodigal and searched against the Virus Orthologous Groups (VOGs, vogdb.org) via HMM

searches (E-value 0.001). A bipartite network was made based on shared VOGs using igraph

(graph.incidence) (1.2.5) [62] in R (version 3.5.1) [63] with RStudio (version 1.1.456) [64]. A

previous study classified divergent viral lineages via the spinglass community detection algorithm

[41], which we used on the bipartite network generated here via igraph (50 spins for 100

iterations). Final clusters were determined by using those of the iteration with the highest

modularity. Plots of the cluster composition from the bipartite network analysis were made with

ggplot2 (3.1.1) [65] in R with Rstudio. TerL (terminase large subunit) and MCP (major capsid

protein) trees were made with hits to TerL VOG families and MCP VOG families encoded by the

jumbo phages and reference hits (hmmsearch, E-value < 0.001; See Data Availability). Reference

hits were de-replicated with CD-HIT (version 4.8.1 -c 0.9) [66] and filtered for size (See

Supplemental Methods). Proteins were then aligned with Clustal Omega [67], trimmed with trimAl

(-gt 0.1) [68] and constructed with IQ-TREE [69] (TEST model selection with ModelFinder [70]).

Size fraction and ecological analyses. Metagenomic reads from Tara Oceans were trimmed

with trim_galore (--paired --length 50 -e 0.1 -q 50) and subsampled to an even depth of 20 million

reads per sample with seqkit sample (-s 1000 -n 20000000 -2). These reads were then mapped

onto the population representatives of the jumbo phage set (535 populations). For the mapping,

the reference database of the representative jumbo phage sequences was created with minimap2

(minimap2 -x sr -d)[71], and the mapping was carried out with coverM (coverm genome --min-

read-percent-identity 95 -m covered_fraction rpkm count variance length -t 32 --minimap2-

reference-is-index --coupled)).   Mapping results were retained if at least 20% of the phage

genome was covered (see Supplemental Methods for benchmarking, Supplemental Figure 26).

Relative abundance of a phage in each sample was calculated in reads per kilobase per million

(RPKM). Statistical analyses and plots were carried out in R with vegan (2.5-5) [72], ggplot2,

maps (3.3.0) [73], and ggpubr (0.2.4) [74] packages. Community composition was compared

between variables using ANOSIMs based on Bray-Curtis distances using both presence/absence

and RPKM matrices with a significance *p values* < 0.05. Statistical tests were carried out with the

ggplot2 function stat_compare_means(label="p.signif").

Annotation. Amino acid sequences of genes were annotated with HMM searches (E-value <

0.001) against the Pfam [75] (version 32), eggNOG (5.0) [76], and VOG (release 98) databases.

Virion structural protein families were identified based on VOG hit descriptions (Supplemental

Dataset 3). Consensus annotation was based on Pfam annotations and then the highest bit score

between eggNOG and VOG hits. Functions were grouped into larger categories (Supplemental

Dataset 3). Clusters from the network analyses were compared for functional composition

between marine jumbo phages by averaging the proportion of genes in a functional category.

Genes with the highest variance between PGCs were identified based on the variance in the

proportion of genomes in a PGC with that gene. Those with a variance >0.2 and a known function

were visualized with pheatmap [77] in R.

Host prediction. Hosts of the jumbo phage bins were predicted based on matching CRISPR

spacers, tRNAs, and gene content. CRISPR spacers were predicted on the Genome Taxonomy

Database (release 95) [78], metagenome assembled genomes (MAGs) of bacteria and archaea

from the metagenomes that the jumbo phage bins derived (provided by Parks et al. 2017 [30]),

and on the jumbo phage bins with minCED (derived from reference [79]). Spacers were aligned

with BLASTn (-task blastn-short) and matches were >24 bp with <= 1 mismatches [4]. Only one

jumbo phage contained a CRISPR array, but the spacers did not match any other jumbo phages

or MAGs. tRNA sequences were predicted with tRNAscan-SE (-bacteria option) [80] on the MAGs

and jumbo phage bins. Promiscuous tRNAs [81] were removed (BLASTn hits 100% ID, <= 1

mismatches). Jumbo phage tRNAs were aligned against the MAGs tRNAs and NCBI nr database

(BLASTn 100% ID, <= 1 mismatches) [4]. Lastly, hosts were assigned based on the taxonomy of

coding sequence matches to the MAGs (BLASTn). Hits to phyla were summed and a putative

host phylum had three times the number of hits as the phylum with the next most hits as used in

a previous study [4]. If a putative host could be predicted by multiple methods, a consensus host

was assigned if all approaches agreed on a phylum. If the methods disagreed at the phylum-level,

no putative host was assigned.

Data Availability

Nucleic acid sequences and protein predictions for the 85 bins analyzed in this study and the

proteins and files for the phylogenetic analyses (proteins, HMM profiles, treefiles) can be found

on FigShare (https://figshare.com/projects/Marine_jumbo_phages/127391).

Acknowledgements

We thank members of the Aylward Lab for helpful feedback. We thank Anh Ha for assistance with

read mapping. This work was performed using compute nodes available at the Virginia Tech

Advanced Research and Computing Center.

Competing Interests

The authors declare no competing financial interests.

References

1. Brüssow H, Hendrix RW. Phage Genomics. Cell. 2002;108:13–16.

2. Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh DA, Denef VJ, et al. Giant virus

diversity and host interactions through global metagenomics. Nature. 2020;578:432–436.

3. Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO. Dynamic genome

evolution and complex virocell metabolism of globally-distributed giant viruses. Nat

Commun. 2020;11:1710.

4. Al-Shayeb B, Sachdeva R, Chen L-X, Ward F, Munk P, Devoto A, et al. Clades of huge

phages from across Earth's ecosystems. Nature. 2020;578:425–431.

5. Koonin EV, Yutin N. Origin and evolution of eukaryotic large Nucleo-Cytoplasmic DNA

viruses. Intervirology. 2010;53:284–292

6. M Iyer L, Anantharaman V, Krishnan A, Burroughs AM, Aravind L. Jumbo phages: a

comparative genomic overview of core functions and adaptions for biological conflicts.

Viruses. 2021;13.1:63.

7. Raoult D, Forterre P. Redefining viruses: lessons from Mimivirus. Nat Rev Microbiol.

2008;6:315–319.

8. Ha AD, Moniruzzaman M, Aylward FO. High transcriptional activity and diverse functional

repertoires of hundreds of giant viruses in a coastal marine system. mSystems.

2021;6:e00293-21.

9.   Endo H, Blanc-Mathieu R, Li Y, Salazar G, Henry N, Labadie K, et al. Biogeography of

marine giant viruses reveals their interplay with eukaryotes and ecological functions. Nat

Ecol Evol 2020; 4: 1639–1649.

10.   Kaneko H, Blanc-Mathieu R, Endo H, Chaffron S, Delmont TO, Gaia M, et al. Eukaryotic

virus composition can predict the efficiency of carbon export in the global ocean. iScience.

2021;24:102002.

11.   Hendrix RW. Jumbo bacteriophages. Curr Top Microbiol Immunol. 2009;328:229–240.

12.   Low SJ, Džunková M, Chaumeil P-A, Parks DH, Hugenholtz P. Evaluation of a

concatenated protein phylogeny for classification of tailed double-stranded DNA viruses

belonging to the order Caudovirales. Nat Microbiol. 2019;4:1306–1315.

13.   Krylov VN, Zhazykov IZ. Pseudomonas bacteriophage phiKZ--possible model for studying

the genetic control of morphogenesis. Genetika. 1978;14:678–685.

14.   Cook R, Brown N, Redgwell T, Rihtman B, Barnes M, Stekel DJ, et al. INfrastructure for a

PHAge REference Database: Identification of large-scale biases in the current collection of

phage genomes. PHAGE. 2021;2.4:214-223.

15.   Yuan Y, Gao M. Jumbo bacteriophages: an overview. Frontiers in Microbiology.

2017;8:403.

16.   Nazir, A., Ali, A., Qing, H., & Tong, Y. Emerging aspects of jumbo bacteriophages. Infect.

Drug Resist. 2021;14:5041.

17.   Ceyssens P-J, Minakhin L, Van den Bossche A, Yakunina M, Klimuk E, Blasdel B, et al.

Development of giant bacteriophage KZ Is independent of the host transcription apparatus.

J Virol. 2014;88:10501–10510.

18.   Mendoza SD, Nieweglowska ES, Govindarajan S, Leon LM, Berry JD, Tiwari A, et al. A

bacteriophage nucleus-like compartment shields DNA from CRISPR nucleases. Nature.

2020;577:244–248.

19.   Malone LM, Warring SL, Jackson SA, Warnecke C, Gardner PP, Gumy LF, et al. A jumbo

phage that forms a nucleus-like structure evades CRISPR–Cas DNA targeting but is

vulnerable to type III RNA-based immunity. Nat. Microbiol. 2020;5:48–55.

20. Serwer P, Hayes SJ, Thomas JA, Hardies SC. Propagating the missing bacteriophages: a

large bacteriophage in a new class. Virol J. 2007;4:21.

21. Thurber RV, Haynes M, Breitbart M, Wegley L, Rohwer F. Laboratory procedures to

generate viral metagenomes. Nat Protoc. 2009;4:470–483.

22. Palermo CN, Shea DW, Short SM. Analysis of different size fractions provides a more

complete perspective of viral diversity in a freshwater embayment. Appl Environ Microbiol.

2021;87.11:e00197-21.

23. Sieburth JM, Smetacek V, Lenz J. Pelagic ecosystem structure: Heterotrophic

compartments of the plankton and their relationship to plankton size fractions 1. Limnol and

Oceanogr. 1978;23:1256–1263.

24. Brum JR, Ignacio-Espinoza JC, Roux S, Doulcier G, Acinas SG, Alberti A, et al. Ocean

plankton. Patterns and ecological drivers of ocean viral communities. Science.

2015;348:1261498.

25. Roux S, Brum JR, Dutilh BE, Sunagawa S, Duhaime MB, Loy A, et al. Ecogenomics and

potential biogeochemical impacts of globally abundant ocean viruses. Nature.

2016;537:689–693.

26. Gregory AC, Zayed AA, Conceição-Neto N, Temperton B, Bolduc B, Alberti A, et al. Marine

DNA Viral Macro- and Microdiversity from Pole to Pole. Cell. 2019;177:1109–1123.e14.

27. Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. Benchmarking viromics: an

evaluation of metagenome-enabled estimates of viral community composition and diversity.

PeerJ. 2017;5:e3817.

28. García-López R, Vázquez-Castellanos JF, Moya A. Fragmentation and coverage variation

in viral metagenome assemblies, and their effect in diversity calculations. Front Bioeng

Biotechnol. 2015;3:141.

29. Sunagawa S, Pedro Coelho L, Chaffron S, Roat Kultima J, Labadie K, Salazar G, et al.

Structure and function of the global ocean microbiome. Science. 2015;348.6237:1261359.

30. Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, et al.

Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree

of life. Nat Microbiol. 2017;2:1533–1542.

31. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies.

PeerJ. 2019;7:e7359.

32. Guo J, Bolduc B, Zayed AA, Varsani A, Dominguez-Huerta G, Delmont TO, et al.

VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA and RNA

viruses. Microbiome. 2021;9.1:1-13.

33. Kieft K, Zhou Z, Anantharaman K. VIBRANT: automated recovery, annotation and curation

of microbial viruses, and evaluation of viral community function from genomic sequences.

Microbiome. 2020;8.1:1-23.

34. Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. CheckV

assesses the quality and completeness of metagenome-assembled viral genomes. Nat

Biotechnol. 2021;39:578–585.

35. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome Res. 2015;7:1043-1055.

36. Cook R, Brown N, Redgwell T, Rihtman B, Barnes M, Clokie M, et al. INfrastructure for a

PHAge REference Database: Identification of large-scale biases in the current collection of

cultured phage genomes. PHAGE. 2021;2.4:214-223.

37. Luo E, Eppley JM, Romano AE, Mende DR, DeLong EF. Double-stranded DNA

virioplankton dynamics and reproductive strategies in the oligotrophic open ocean water

column. ISME J. 2020;14:1304–1315.

38. Michniewski S, Rihtman B, Cook R, Jones MA, Wilson WH, Scanlan DJ, et al. Identification

of a new family of 'megaphages' that are abundant in the marine environment. ISME

Comm. 2021;1.1:1-4.

39. Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al. Taxonomic

assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing

networks. Nat Biotechnol. 2019;37:632–639.

40. Iranzo J, Krupovic M, Koonin EV. The double-stranded DNA virosphere as a modular

hierarchical network of gene sharing. MBio 2016;7.4:e00978-16.

41. Reichardt J, Bornholdt S. Statistical mechanics of community detection. Phys Rev E Stat

Nonlin Soft Matter Phys. 2006;74.1:016110.

42. Hurwitz BL, Sullivan MB. The Pacific Ocean virome (POV): a marine viral metagenomic

dataset and associated protein clusters for quantitative viral ecology. PLoS One.

2013;8:e57355.

43. Pappas N, Dutilh BE. Finding functional associations between prokaryotic virus orthologous

groups: a proof of concept. BMC Bioinform. 2021;22:1–11.

44. Fridman S, Flores-Uribe J, Larom S, Alalouf O, Liran O, Yacoby I, et al. A myovirus

encoding both photosystem I and II proteins enhances cyclic electron flow in infected

Prochlorococcus cells. Nat Microbiol. 2017;2:1350–1357.

45. Hjorleifsdottir S, Aevarsson A, Hreggvidsson GO, Fridjonsson OH, Kristjansson JK.

Isolation, growth and genome of the Rhodothermus RM378 thermophilic bacteriophage.

Extremophiles. 2014;18:261–270.

46. Mende DR, Bryant JA, Aylward FO, Eppley JM, Nielsen T, Karl DM, et al. Environmental

drivers of a microbial genomic transition zone in the ocean's interior. Nat Microbiol.

2017;2:1367–1373.

47. Goldsmith DB, Brum JR, Hopkins M, Carlson CA, Breitbart M. Water column stratification

structures viral community composition in the Sargasso Sea. Aquat. Microb. Ecol.

2015;76:85–94.

48. Hurwitz BL, Brum JR, Sullivan MB. Depth-stratified functional and taxonomic niche

specialization in the 'core' and 'flexible' Pacific Ocean Virome. ISME J. 2015;9:472–484.

49. Breitbart M, Rohwer F. Here a virus, there a virus, everywhere the same virus? Trends

Microbiol. 2005;13:278–284.

50. Brüssow H. Huge bacteriophages: bridging the gap? Environ Microbiol. 2020;22:1965–

1970.

51. Koonin EV, Yutin N. Evolution of the large nucleocytoplasmic DNA viruses of eukaryotes

and convergent origins of viral gigantism. Adv Virus Res. 2019;103:167–202.

52. Edwards KF, Steward GF, Schvarcz CR. Making sense of virus size and the tradeoffs

shaping viral fitness. Ecol Lett. 2021;24:363–373.

53. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic

gene recognition and translation initiation site identification. BMC Bioinform. 2010;11.1:1-

11.

54. Williams T, Kelley C. Gnuplot 5.2 Manual: An interactive plotting program. 2017.

55. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile and

open software for comparing large genomes. Genome Biol. 2004;5.2:1-9.

56. Kiełbasa SM, Wan R, Sato K, Horton P, Frith MC. Adaptive seeds tame genomic sequence

comparison. Genome Res. 2011;21:487–493.

57. Aylward FO, Moniruzzaman M. ViralRecall - a flexible command-line tool for the detection

of giant virus signatures in 'omic data. Viruses. 2021;13.2:150.

58. wwood. GitHub - wwood/CoverM: Read coverage calculator for metagenomics.

https://github.com/wwood/CoverM. Accessed 23 Jul 2021.

59. Shulgina Y, Eddy SR. A computational screen for alternative genetic codes in over 250,000

genomes. eLife 2021;10:e71402.

60. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J

672  Mol Biol. 1990;215:403–410.

673 61. Iranzo J, Koonin EV, Prangishvili D, Krupovic M. Bipartite network analysis of the archaeal

674   virosphere: evolutionary connections between viruses and capsidless mobile elements. J

675   Virol. 2016;90.24:11043–11055.

676 62. igraph – Network analysis software. http://igraph.org. Accessed 23 Jul 2021.

677 63. R Core Team. R: A Language and Environment for Statistical Computing. 2019. R

678   Foundation for Statistical Computing, Vienna, Austria.

679 64. RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA

680   URL http://www.rstudio.com/.

681 65. Wickham H. ggplot2. Wiley Interdisciplinary Reviews: Comput Stat. 2011;3.2:180–185.

682 66. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation

683   sequencing data. Bioinformatics. 2012;28:3150–3152.

684 67. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of

685   high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol.

686   2011;7:539.

687 68. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment

688   trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25:1972–1973.

689 69. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

690   stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.

691   2015;32:268–274.

692 70. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast

693   model selection for accurate phylogenetic estimates. Nat Methods. 2017;14:587–589.

694 71. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34:

695   3094–3100.

696 72. Dixon P. VEGAN, a package of R functions for community ecology. Journal of Vegetation

697   Science. 2003;14:927–930.

73. maps: Draw Geographical Maps. https://CRAN.R-project.org/package=maps. Accessed 22

Jul 2021.

74. Kassambara A. 'ggplot2' Based Publication Ready Plots [R package ggpubr version 0.4.0].

2020.

75. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al.

Pfam: The protein families database in 2021. Nucleic Acids Res. 2020;49.D1:D412–D419.

76. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, et al.

eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource

based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 2019;47:D309–D314.

77. pheatmap: Pretty Heatmaps. https://CRAN.R-project.org/package=pheatmap. Accessed 22

Sep 2021.

78. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, et al. A

standardized bacterial taxonomy based on genome phylogeny substantially revises the tree

of life. Nat Biotechnol. 2018;36:996–1004.

79. Bland C, Ramsey TL, Sabree F, Lowe M, Brown K, Kyrpides NC, et al. CRISPR

Recognition Tool (CRT): a tool for automatic detection of clustered regularly interspaced

palindromic repeats. BMC Bioinform. 2007;8.1:1-8

80. Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of

transfer RNA genes. Nucleic Acids Res. 2016;44.W1:W54–W57.

81. Paez-Espino D, Eloe-Fadrosh EA, Pavlopoulos GA, Thomas AD, Huntemann M, Mikhailova

N, et al. Uncovering earth's virome. Nature 2016;536:425–430.

Figure Legends

Figure 1. Comparison of bins to reference gene content and evolutionary histories. (a)

Network with marine jumbos and references as nodes and edges based on shared VOGs.

Marine jumbo phage nodes are colored by PGC as detected with spinglass community

detection analysis, other nodes are in gray. Node size corresponds to the natural log of genome

length in kilobases. **(b)** Barchart of the number of members in each PGC. PGCs with marine

jumbo phages are denoted with a star and the number of marine jumbo phages in that PGC.

Proportion of marine jumbo phages in that PGC is colored. c,d) Phylogenies of TerL **(c)** and

MCP **(d)** proteins with references and bins. Inner ring and branches are colored by the 5 PGCs

that marine jumbo phages belong to. Navy blue circles in the outer ring denote marine jumbo

phages.

Figure 2. (a) Boxplot of genome length in each network cluster (x-axis is PGC number). Star

denotes PGC with a marine jumbo phage and the color matches the PGC letters of Figure 1. **(b)**

Stacked barplot of the metagenome environment from which each phage derives from in each

PGC (x-axis). Reference (yellow) are cultured phages, black is the bins of jumbo phages from

this study **(c)** Stacked barplot of the host phylum of the RefSeq cultured phages in each cluster;

metagenomic phages are in gray.

Figure 3 (a) Functional categories for genes encoded by jumbo phages. **(b)** Heatmap of

proportion of genomes in each PGC that contain the listed genes. Listed genes were selected

based on containing a known function and variance between the PGCs above 0.2. Dendrogram

was generated based on hierarchical clustering in pheatmap.

Figure 4. Comparison of jumbo abundance and presence in samples of different filter

size fractions. Dark teal are fractions with minimum sizes of 0.22 μm or higher. Light teal are

fractions with a maximum size of 0.22 μm or lower. **(a)** Barchart of the proportion of samples

with at least one marine jumbo phage (x-axis) by size fraction (y-axis) sorted from highest to

lowest. **(b)** Boxplot with x-axis as the number of marine jumbo phages found in a sample with

size fraction on the y-axis sorted by median. **(c)** Boxplot with x-axis as the relative abundance of

marine jumbo phages found in a sample (RPKM) with size fraction on the y-axis sorted by

median. Significance bars in c,d correspond to Wilcox test, with stars corresponding to p-values

< 0.05 and those with p-values > 0.05 as not significant "ns" (stat_compare_means function).

(d,e) NMDS plots (Bray Curtis dissimilarity distances) of jumbo phage composition in each

sample using presence absence data **(d)** and relative abundance data **(e)**. Samples are colored

by size fraction distinction above 0.22 (dark teal) and below 0.22 (light teal). Ellipses calculated

based on multivariate normal distribution. Samples above and below 0.22 were not significantly

different from each other using ANOSIM (p-values > 0.05).

Figure 5. (a,b) Maps of the relative abundance **(a)** of total jumbo phages (in RPKM) and **(b)**

total number of jumbo populations present regardless of phage cluster membership in each

surface (SRF) sample of the picoplankton size fraction (either 0.22-3µm or 0.22-1.6µm

depending on availability). Dots sizes are proportional to the number of populations or RPKM.

(c) Scatterplot of the mean RPKM of a jumbo population in SRF picoplankton samples versus

the number of SRF picoplankton stations it was present. Populations are colored by PGC and

size corresponds to putative genome length in 100 kilobases. **(d)** Boxplot of the number of

jumbo phage populations in samples co-collected at each depth sorted by mean for each PGC.

Significance bars correspond to Wilcox test, with stars corresponding to *p values* < 0.05

(stat_compare_means function).