An early decline in ETPs reflects reduced pre-thymic progenitors and altered signals from the thymus microenvironment

Short title: Early ETP decline reflects BM and thymus changes

Jayashree Srinivasan[1*], Anusha Vasudev[2*], Hilary J. Selden[1], Encarnacion Perez Jr.[2], Bonnie LaFleur[3], Shripad A. Sinari[3], Andreas Krueger[4], Ellen R. Richie[2§], Lauren I. R. Ehrlich[1§]

[1]Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX; [2]Department of Epigenetics and Molecular Carcinogenesis, The University of Texas MD Anderson Cancer Center, Houston, TX; [3]Center for Biomedical Informatics and Statistics, The University of Arizona, Tucson, AZ; [4]Institute for Molecular Medicine, Goethe University Frankfurt am Main, 60590 Frankfurt am Main, Germany

* These authors contributed equally to the work.
§ Co-senior authors

Corresponding authors:

Name: Lauren I. R. Ehrlich
Address: 100 East 24th St., STOP A5000, Austin, TX 78712-1191
Email: lehrlich@austin.utexas.edu
Phone: 512-475-7125

Name: Ellen R. Richie
Address: 1515 Holcombe Blvd, Unit 1951, Houston, TX 77030-4009
Email: erichie@mdanderson.org
Phone: 832-750-7181

Abstract

Age-associated thymus involution results in decreased generation and export of T cells, with an associated increase in susceptibility to pathogens. Elucidating mechanisms underlying diminished T-cell production during involution will inform strategies to restore thymopoiesis with age. T-cell development depends on periodic thymic colonization by circulating bone marrow (BM)-derived thymus seeding progenitors (TSPs). Upon thymic entry, TSPs differentiate into early T-cell progenitors (ETPs), which give rise to all downstream thymocyte subsets. Here, we find a significant decline in ETP cellularity between 1 and 3 months (3MO) of age, mirrored by reduced total thymocyte cellularity. This early thymic involution could be driven by changes in pre-thymic progenitors and/or thymic stromal cells. Using a multicongenic progenitor transfer approach, we determined that the number of TSP-supportive niches does not diminish through 12MO. However, the number of lymphoid progenitors in BM and blood declines substantially by 3MO of age, although their cell-intrinsic capacity to seed and differentiate in the thymus is unimpaired, indicating that pre-thymic hematopoietic changes contribute to the early decline in ETPs. Moreover, sustaining thymic epithelial cells maintains ETP cellularity with age, despite the reduction in circulating TSPs, supporting the idea that age-associated defects in ETP niche quality also contribute to early involution. Consistent with this possibility, Notch signaling, which is essential for T-lineage differentiation, diminishes in 3MO BM progenitors and in ETPs. Together, our findings suggest that a reduction in both BM lymphopoiesis and thymic stromal support contribute to the initial age-associated decline in ETPs, setting the stage for progressive thymic involution.

Introduction

T cells develop in the thymus which provides a unique and complex stromal microenvironment essential for T-cell lineage commitment, differentiation and selection[1,2]. Since thymocytes do not have the ability to self-renew, the thymus periodically recruits bone marrow (BM)-derived thymus seeding progenitors (TSPs) which are essential to maintain T cell production throughout life[3]. TSP entry is a well-orchestrated multistep process mediated by adhesion molecules P-selectin, ICAM-1, and VCAM-1, chemokines CCL25 and CCL21, and other niche factors, like KITL, supplied by thymic endothelial cells[4-8]. After entering the thymus via vasculature at the cortico-medullary junction, TSPs differentiate into highly proliferative $CD4^-$ $CD8^-$ $CD44^{hi}$ $c-kit^{hi}$ early T-cell progenitors (ETPs) in the thymic cortex. Commitment of developing thymocytes to the T-cell lineage depends on inductive signals provided by cortical thymic epithelial cells (cTECs)[1,2,9,10]. In particular, activation of Notch signaling in ETPs via Delta-like Notch ligand 4 (DLL4) expressed by cortical thymic epithelial cells (cTECs) is indispensable for T-cell lineage commitment and differentiation[11-13]. Since ETPs give rise to all downstream T cell subsets and differentiation stages, thymopoiesis and T cell development depend on both continued influx of TSPs into the thymus and the availability of functional stromal niches that support TSP entry and downstream ETP differentiation.

During age-associated thymus involution, thymus size and cellularity undergo a progressive decline accompanied by reduced output of naïve T cells. Thymus involution is initiated early in life, by ~7 weeks of age in mice[14,15]. Thus, we first asked if ETP cellularity is adversely affected during the earliest stages of thymus involution. Here, we find that the decline in ETP cellularity is initiated early, between 1MO and 3MO of age in mice, and further declines with advancing

age. Thymus involution involves not only a reduction in thymocyte cellularity, but also a decline in the number, composition, proliferative capacity, and organization of the TEC compartment[16-20]. TEC degeneration is a key driver of age-associated thymic involution, as sustaining or restoring TECs prevents or reverses involution, respectively[14,21-24]. Given that thymic stromal niche factors play a key role in recruitment of TSPs and that thymic stromal signals are required for proliferation and T-lineage commitment of ETPs, we anticipated the early age-associated decline in ETPs would be driven by changes in the thymic microenvironment.

Here, we used several approaches to determine whether the early loss of ETPs is due to age-related changes in availability of functional TSP/ETP niches and/or in pre-thymic lymphoid progenitors. First, we used a multicongenic progenitor transfer approach to quantify the number of functional TSP/ETP niches from the onset of involution through middle-age[25]. These studies revealed that the number of TSP/ETP niches is sustained, despite substantial involution, through 12MO of age. We thus considered whether the reduction in ETP cellularity could be due to a decline in the hematopoietic progenitors that seed the thymus. Surprisingly, we observe a significant decline in the number of lymphoid progenitors with T-lineage potential in the 3MO BM and blood, although their cell-intrinsic ability to seed and differentiate in the thymus remains intact. Because Notch signaling plays a key role in promoting T-lineage specification and progenitor differentiation, we also interrogated niche function by evaluating *Dll4* expression by cTECs and assessing Notch signaling activity in ETPs and BM progenitors. Notably, we find reduced Notch signaling in T-lineage precursors in both the BM and thymus by 3MO of age. Collectively, these findings demonstrate a surprisingly early decline in BM lymphoid progenitors and circulating TSPs, along with diminished Notch signaling in the BM and

thymus microenvironments, implicating both pre-thymic and thymic changes in the early decline of ETP cellularity at the outset of thymus involution.

Methods

Mice

C57BL/6J (CD45.2, Thy1.2), B6.PL-Thy1a/CyJ 9 (CD45.2, Thy1.1), B6.SJL-*Ptprc[a] Pepc[b]*/BoyJ (CD45.1, Thy1.2), C57BL/6-Tg(CAG-EGFP)1Osb/J (GFP)[26], Tg(Cp-HIST1H2BB/Venus)47Hadj/J (Notch-Venus reporter)[27], and K5.cyclin D1[28] (Conti lab, MD Anderson Cancer Center, Smithville, TX). Mice were obtained from Jackson Laboratories, except as specified, and bred in-house. F1 mice for multicongenic barcoding experiments were bred in-house from CD45 congenic, Thy1 congenic, and EGFP strains (supplemental Figure 2B). C57BL/6J recipient mice for multicongenic barcoding experiments were obtained from the National Institute of Aging. All mice were maintained under specific pathogen-free conditions at the animal facilities at the University of Texas at Austin and the University of Texas MD Anderson Cancer Center, Smithville. All experimental procedures were performed in accordance with the Institutional Animal Care and Use Committees.

Multicongenic barcoding progenitor transfer assay

BM cells were obtained from long bones of 4-6-week-old mice of 8 different congenic backgrounds (supplemental Figure 2B). Bones were crushed and single cell suspensions were prepared in FACS wash buffer (PBS+ 2% FBS), followed by RBC lysis (RBC lysis buffer, BioLegend). Cells were stained with lineage (Lin)-specific antibodies against CD11b, Gr-1, Ter-119, B220, CD19, CD3 and CD8 (all from BioXCell), and Lin[+] cells were depleted with sheep anti-rat IgG immunomagnetic beads (Dynabeads, Invitrogen). Cells were then immunostained and

Lin⁻Flk2⁺CD27⁺ BM progenitors from F1 congenic strains were FACS purified (supplemental

Figure 2A). Progenitors from each of the 8 congenic strains were mixed at equal ratios, and a

total of 100,000 cells were retro-orbitally injected into non-irradiated 1MO, 3MO, 6MO and

12MO C57BL/6J recipient mice. Thymocytes were analyzed after 21 days by flow cytometry to

distinguish the contributions of distinct congenic donor strains. 2.5 million events were

recorded per recipient thymus, and successful thymic colonization was scored if a minimum of

40 cells of a given donor strain was detected. Quantification of niches was performed using a

multinomial sampling algorithm (see Statistical Analysis).

Competitive heterochronic progenitor seeding assay

Lin⁻Flk2⁺CD27⁺ progenitors were isolated from the long bones of 1MO and 3MO congenic

donor mice as described above. 1MO and 3MO progenitors were mixed at 1:1 ratio and

100,000 cells were retro-orbitally injected into 1MO and 3MO non-irradiated C57BL/6J

recipient mice. After 21 days, donor chimerism was analyzed in the recipient thymi by flow

cytometry.

Statistical analysis

All statistical analyses were performed using Prism (V9.2.0, GraphPad Software). Statistical

significance was determined using an unpaired Student t-test, 1-way analysis of variance

(ANOVA) with Kruskal-Wallis multiple comparisons test or 2-way ANOVA with Sidak's multiple

comparisons test for normally distributed data.

For quantification of thymic niches, a Bayesian technique was implemented to estimate the

number of niches present in recipient mice from the missing number of donors detected.

Generalized linear models were used to test if the distributions of the estimated number of

niches present in each age category declines with increasing age. General linear modeling was performed with Poisson response and a log link function with age as the only categorical covariate. Test statistic (χ^2) was used to test the trend of number of niches over age groups. For estimation, the donor probabilities from the proportion of donor cells injected in each mouse was calculated. A posterior likelihood was computed using sampling from a multinomial distribution where L is the number of iterations, N is the number of niches and k is the number of distinct donors. This likelihood was then normalized to give a probability distribution for each likely value of missing donors. The maximum estimate of N that are drawn from the posterior distributions (MAP) were used as the most likely estimate of available niches. Batch to batch variation (i.e. replicate experiments) was used to test for potential bias due to batch effects. Regression analysis was used to assess if there are differences in available niches by age.

The supplemental methods section contains additional experimental details.

Results

ETP and total thymocyte cellularity decline by 3MO of age

We first quantified age-associated changes in ETPs and downstream thymocyte subsets between 1 and 12MO of age (Figure 1; supplemental Figure 1A). Total thymus cellularity declines with age, dropping significantly by 2MO, consistent with previous reports that thymus involution begins early in life (Figure 1A)[14,15,20]. Notably, ETP cellularity also declines progressively, starting at 1.5MO, which precedes, but otherwise mirrors the reduction in total thymocytes (Figure 1A). There is a strong correlation between ETP and total thymocyte numbers through 12MO of age (Figure 1B), and this correlation extends to subsequent

thymocyte subsets (supplemental Figure 1B), showing that increased numbers of downstream thymocytes do not compensate for the loss of ETPs with age. Moreover, the consistent DN2 to ETP ratio over the first year of life further suggests that ETP differentiation is not impaired with age (Figure 1C).

The number of available thymic niches for TSPs does not decrease with age

The age-associated decline in ETP cellularity could be a function of changes in the thymic stromal microenvironment and/or defects in pre-thymic lymphoid progenitors. Given the important role of thymic microenvironment in supporting all stages of developing T cells and its known impact on involution, we first tested whether the reduction in ETPs is due to an age-related decline in the number of functional progenitor niches that support early stages of thymopoiesis. We used a modification of a previously reported multicongenic progenitor transfer approach to quantify available, functional TSP niches from 1MO-12MO of age. This approach previously revealed that a young thymus contains ~10 available niches that can be colonized by thymocyte progenitors at any one time[25]. We FACS sorted $Lin^-Flk2^+CD27^+$ BM progenitors, which encompass all thymus seeding and T cell progenitor activity in the BM[29], from 8 different congenic strains, mixed them at equal ratios, and transplanted them into non-irradiated recipient mice of different ages (Figure 2A; supplemental Figure 2A-B). Recipient thymuses were analyzed after 21 days to allow sufficient time for donor progenitors to complete one wave of T-cell differentiation[29]. To determine the number of missing congenic tags, we quantified the number of different donor strains present in each recipient thymus by flow cytometry (supplemental Figure 2C). If the number of available niches that support T cell

development declines with age, then fewer donor strains will be detected in thymuses of older recipients. However, we found that the number of available, functional niches does not decline with age; to the contrary there is a trend toward an increased number of niches in 6 and 12MO recipients (Figure 2B-C). Therefore, the age-associated decline in ETP cellularity is not due to a reduction in the number of functional niches available to thymus seeding progenitors.

1MO and 3MO T-cell progenitors seed and thrive comparably in the thymus environment

We next asked whether the decline in ETPs reflects intrinsic defects in 3MO TSPs, compromising their ability to seed and differentiate in available thymic niches. Thus, we performed competitive heterochronic progenitor transfer assays in which isolated Lin$^-$ Flk2$^+$CD27$^+$BM progenitors obtained from 1 and 3MO congenic donor strains were mixed at an equal ratio and injected i.v. into 1 versus 3MO non-irradiated recipients. If BM progenitors from 3MO donors have intrinsic defects, then we would expect to detect fewer 3MO compared to 1MO donor-derived cells in recipient thymuses. However, comparable percentages and numbers of 1MO and 3MO donor cells were recovered from recipient thymuses of both ages (Figure 3A-B), indicating that TSPs from 3MO donors are fully competent to seed and thrive in the thymus. Moreover, there is no apparent defect in the differentiation capacity of 3MO TSPs, as a similar percentage of DP thymocytes are generated from 1MO and 3MO donor cells (Figure 3C).

We next considered the possibility that changes in proliferation or survival could account for the decline in ETPs at 3MO. However, a comparable percentage of 1MO and 3MO ETPs incorporated BrdU (Figure 3D-E), and analysis of intracellular cleaved caspase-3 showed that

apoptosis is negligible in ETPs at both 1MO and 3MO of age (supplemental Figure 3). Taken

together, the data suggest that the decline in ETP numbers at 3MO of age is not a consequence

of defects in the ability of 3MO thymus seeding progenitors to thrive in the thymus.

Lymphoid progenitors with T-lineage potential decline in the BM and blood of 3MO mice

As we found no apparent defects in either the availability of functional TSP niches or the

ability of TSPs to enter and thrive in those niches at 3MO of age, we asked whether the

numbers of BM or circulating lymphoid progenitors are decreased by 3MO of age. A significant

reduction in the number of TSPs could play a key role in the age-associated decline in ETP

cellularity. To assess this possibility, we used flow cytometry to quantify circulating Lin$^-$

Flk2$^+$CD27$^+$ cells, which comprise all thymic seeding activity in the blood[29] (Figure 4A). We also

quantified upstream Lin$^-$Flk2$^+$CD27$^+$ lymphoid progenitors in the BM (Figure 4C). Interestingly,

the frequency and number of Flk2$^+$ CD27$^+$ lymphoid progenitors in both blood and BM

undergoes a significant decline between 1MO and 3MO of age (Figure 4B, D). Flk2$^+$ MPPs and

Ly6D$^-$ CLPs are the lymphoid progenitors within the Lin$^-$Flk2$^+$CD27$^+$ compartment that have T-

lineage potential, and both subsets also decline significantly by 3MO of age (Figure 4E-F). These

results demonstrate a surprisingly early reduction in the number of BM lymphoid progenitors

and circulating TSPs. These data suggest that the decrease in ETP cellularity by 3MO is likely

due, at least in part, to a reduction in the number of circulating TSPs that are available to enter

the thymus and differentiate.

Sustaining TEC proliferation prevents the age-associated decline in ETP cellularity

cTECs provide a critical supportive niche for ETP development, suggesting the age-associated deterioration of the TEC compartment could contribute to the reduction in ETPs by 3MO of age. This scenario is supported by analysis of thymocyte and ETP cellularity in K5.D1 transgenic mice in which a keratin 5 promoter drives Cyclin D1 transgene expression in TECs, but not in hematopoietic or non-TEC stromal cells. The K5.D1 transgene promotes cell cycle progression and sustained growth of the TEC compartment, which in turn, prevents age-associated thymus involution[21]. Sustaining the TEC compartment in K5.D1 mice prevents the decline in ETP and total thymocyte cellularity with age (Figure 5A-B). However, the number of circulating Flk2$^+$CD27$^+$ progenitors is comparable in K5.D1 versus non-transgenic wild-type (WT) control mice at 1MO of age and undergoes a similar decline by 3MO of age in both genotypes (Figure 5C). Moreover, the number of Ly6d$^-$ CLPs declines comparably in K5.D1 and WT mice by 3MO, with a similar trend for BM Flk2$^+$ MPPs (Figure 5D-E). These results indicate that despite a decrease in BM lymphoid progenitors and circulating TSPs, sustaining the TEC compartment is sufficient to prevent the decline in ETPs. Thus, both a decline in lymphoid progenitors and reduced TEC function could contribute to the age-associated reduction in ETPs in WT mice.

Notch signaling is downregulated by 3MO in ETPs and BM lymphoid progenitors

Activation of the Notch signaling pathway in ETPs is required for their development and T-cell lineage commitment. Our previously reported microarray data show that expression of the Notch ligand *Dll4* is downregulated in cTECs between 1 and 3MO of age[30]. Quantitative RT-PCR analysis of isolated 1MO and 3MO cTECs confirms reduced *Dll4* expression by 3MO of age

(Figure 6A). In addition, the level of Notch1 expression on ETPs declines significantly by 3MO

(Figure 6B). To determine if the decreased levels of Notch1 on ETPs and reduced *Dll4*

expression by cTECs at 3MO of age are associated with diminished Notch signaling activity in

ETPs, we used the CBF-H2B-Venus strain in which expression of an H2B-Venus fusion protein is

driven by Notch signaling[27]. Due to perdurance of the Venus reporter fluorescent signal, this

reporter identifies cells that are currently undergoing or have recently undergone Notch

signaling. Flow cytometry analysis revealed that the frequency of ETPs undergoing Notch

signaling declines between 1MO and 3MO of age (Figure 6C-D). Also, the lower relative mean

fluorescence intensity (RFI) of the Venus reporter in 3MO ETPs indicates a reduced level of

Notch signaling.

NOTCH signaling in lymphoid progenitors in the bone marrow also plays a critical role in

generation of T-lineage progenitors[31,32]; thus, we considered the possibility that changes in

Notch signaling could contribute to the reduction in BM lymphoid progenitors at 3MO of age.

Although the frequency of NOTCH-signaled Flk2$^+$ MPPs and Ly6d$^-$ CLPs was comparable at 1MO

and 3MO (Figure 6E-H), there was a notable enrichment in Ly6d- CLPs expressing low levels of

the Venus reporter in 3MO relative to 1MO mice (Figure 6I-J). This suggests that by 3MO of age,

NOTCH signaling is waning in Ly6d-CLPs in the BM. Taken together, these data indicate that the

early decline in ETP cellularity may be due, at least in part, to diminished Notch signaling

activity in ETPs as well as in BM lymphoid progenitors.

Discussion

It is well established that thymus cellularity and T-cell output decline during age-associated thymus involution[17]. While prior studies focused on comparative analysis of thymuses from young mice (1-4MO of age) versus severely involuted thymuses from middle aged and old mice (9-28MO of age), the present investigation focuses on changes that occur during the initial stages of thymus involution between 1 and 3MO of age. In agreement with other reports[14,15,20], we find a significant decline in thymus cellularity by 3MO of age. Furthermore, the progressive reduction in total thymocyte numbers with age is highly correlated with a reduction in the number of ETPs, which contain progenitors that give rise to all subsequent stages of developing T cells. These results indicate that the early decline in ETPs plays a key role in initiating age-associated thymus atrophy. Given that TSPs, ETPs and all subsequent stages of developing T cells rely on inductive and supportive signals from the thymic microenvironment, which undergoes profound age-associated changes that have been shown to be critical in driving thymus involution, we first tested whether a reduction in the thymic niches that support TSPs and ETPs are diminished by 3MO of age. Unexpectedly, we did not find an age-related decline in the number of functional, available TSP niches between 1 and 12MO of age. In fact, there was a slight increase in these niches by 12MO of age. Thus, we subsequently investigated whether the early decline in ETP numbers reflects a decline in BM and/or circulating T-cell progenitors. Our studies reveal a dramatic decline in circulating Flk2$^+$CD27$^+$ progenitors, which contain TSPs, as well as a reduction in BM lymphoid progenitors by 3MO of age. Furthermore, expression of the Notch ligand DLL4 by cTECs is reduced by 3MO, as is Notch signaling activity in ETPs and BM-resident CLPs. Overall, our results suggest that a decline in pre-thymic lymphoid

progenitors in the BM and blood, along with changes in the BM and thymic stromal

microenvironments, contribute to the early onset of thymus involution by 3MO of age.

It has previously been established that hematopoietic progenitors in the BM become

myeloid-biased with age. For example, transplantation experiments revealed that HSCs from

aged mice are biased towards a myeloid fate[7,33-36]. Moreover, fate mapping studies have shown

that MPPs from old mice transit more efficiently into common myeloid progenitors (CMPs) than

into CLPs[36]. Nevertheless, the differentiation potential of lymphoid-biased HSCs on a per cell

basis does not decline with age[37,38]. These prior studies assessed the decline in BM lymphoid

potential between young mice (1-4MO) and much older mice (10-28MO). Since we find that

ETP numbers decline by 3MO of age, we explored the possibility that a reduction in BM

lymphoid progenitors as early as 3MO could contribute to the early reduction in ETPs.

Importantly, we found a striking decrease in the number of circulating TSPs at 3MO, as well as

the number and percentage of the pre-TSP BM progenitors Ly6D$^-$ CLPs and Flk2$^+$ MPPs. These

findings indicate that diminished production and/or egress of T-lineage progenitors from the

BM likely contribute to the early reduction in ETPs. Competitive heterochronic progenitor

transfer assays, in which equivalent numbers of congenic 1 and 3MO Flk2$^+$ CD27$^+$ BM

progenitors were injected into non-irradiated recipients of either age, showed that there is no

cell-intrinsic defect in the ability of 3MO lymphoid progenitors to seed the thymus or

differentiate therein. To our knowledge, the present study is the first to show a substantial

reduction in lymphoid-primed BM progenitors and circulating TSPs by 3MO of age.

As lymphoid progenitor activity is regulated by niche factors in the thymus and/or BM

microenvironments, age-associated changes in stromal cells could also play a role in the early

decline of ETPs and BM lymphoid progenitors. In this regard, heterochronic BM transplantation

experiments showed that successful T-cell reconstitution depends on the age of the recipient

thymus rather than on the age of donor hematopoietic progenitors[39,40], consistent with the

notion that thymic stromal support of thymopoiesis wanes with age. Alterations in the

composition, organization and/or function of the thymus microenvironment could impair

seeding and/or differentiation of TSPs, which enter the thymus on a periodic basis to occupy a

limited number of stromal cell niches[25,41,42]. Therefore, we asked whether the availability of

functional TSP niches diminishes with age. Using multicongenic barcoding progenitor transfers,

combined with mathematical modeling, thymuses from 1MO and 3MO old mice were found to

contain a comparable number of available TSP niches. Thus, the early decline in ETP numbers is

not a function of reduced TSP niche availability. To the contrary, our results showed an upward

trend in the number of TSP niches with age. Previous investigations have reported evidence for

a feedback loop in which key TSP niche factors and T-cell progenitor entry are restricted by the

number of intrathymic T-cell precursors[7,25,43]. Since we find a substantial reduction in the

number of circulating TSPs and intrathymic ETPs as early as 3MO of age, it is possible that the

observed increase in available TSP niches with advancing age is due, at least in part, to a decline

in feedback from TSPs and/or ETPs.

   After TSPs transition to the ETP stage, cTECs play a critical role in supporting their survival,

proliferation and T-lineage commitment by providing niche factors, like DLL4, KITL and IL-7[2,3,44].

Given that the TEC compartment deteriorates during thymus involution, we considered the

possibility that ETP niche factors are diminished with age. The essential role of TECs in

sustaining thymopoiesis was demonstrated by reversing the age-dependent loss of FOXN1, a

key transcription factor required for TEC differentiation and proliferation. Upregulating *Foxn1* in the TEC compartment of old, severely involuted thymuses partially reversed thymus involution by restoring TEC cellularity, which resulted in increased numbers of both ETPs and total thymocytes[23]. Consistent with these findings, we show that the number of ETPs and downstream thymocyte subsets do not decline with age when TEC deterioration and thymus involution are prevented by expression of a Cyclin D1 transgene in TECs of K5.D1 mice. Together, these data support the concept that TEC-derived signals play an essential role in ETP homeostasis and suggest that age-related changes in the TEC compartment contribute to the decline in ETPs.

It is well established that Notch signaling plays a key role in T-lymphopoiesis[2,11,13,31,32,45]. NOTCH1 is expressed on lymphoid progenitors in the BM and on ETPs in the thymus, while the critical Notch ligand DLL4 is expressed by vascular endothelial cells in both BM and thymus as well as on cTECs[2,31,32]. Several observations in the present report suggest that diminished Notch signaling contributes to the early decline in BM lymphoid progenitors and ETPs between 1MO and 3MO of age. We found that lower levels of Notch1 are expressed on ETPs at 3MO of age, when DLL4 expression is reduced on cTECs. Subsequent analysis of NOTCH signaling using the CBF-H2B-Venus NOTCH reporter strain[27] demonstrated an age-related decline in Notch signaling in ETPs, as well diminished Notch signaling in BM-resident CLPs. This decline may be due to deterioration of functional vascular and/or TEC niches. Collectively, these results implicate an early age-associated decline in Notch signaling in T-cell progenitors in the early decline in BM lymphoid progenitors, circulating TSPs and thymic ETPs by 3MO of age.

In summary, our collective results suggest that an early age-associated decline in BM lymphoid potential and alterations in thymic stromal cells during the first 3MO of life synergize to initiate the earliest stages of thymus involution.

Acknowledgements

We thank Janko Nikolich-Zugich, Nancy Manley, Marcel van den Brink, Jarrod Dudakov, Gregory Sempowski, Laura Hale, and Jen Uhrlaub for constructive feedback on experiments and manuscript preparation. This research was supported by a grant from the National Institutes of Health, P01AG052359, to L.I.R.E. and E.R.R.

Authorship contributions

J.S., A.V., E.R. and L. E. designed the experiments and wrote the manuscript; J.S., A.V., H.S., E.P., performed experiments and analyzed data; B.L. and S.S. performed statistical analyses; A.K., E.R. and L.E. edited the manuscript.

Disclosure of conflict of interest

The authors declare no conflicts of interest.

References

1. Yui, M. A. & Rothenberg, E. V. Developmental gene networks: a triathlon on the course to T cell identity. *Nat Rev Immunol* **14**, 529-545, doi:10.1038/nri3702 (2014).
2. Han, J. & Zuniga-Pflucker, J. C. A 2020 View of Thymus Stromal Cells in T Cell Development. *J Immunol* **206**, 249-256, doi:10.4049/jimmunol.2000889 (2021).
3. Krueger, A., Zietara, N. & Lyszkiewicz, M. T Cell Development by the Numbers. *Trends Immunol* **38**, 128-139, doi:10.1016/j.it.2016.10.007 (2017).
4. Krueger, A., Willenzon, S., Lyszkiewicz, M., Kremmer, E. & Forster, R. CC chemokine receptor 7 and 9 double-deficient hematopoietic progenitors are severely impaired in seeding the adult thymus. *Blood* **115**, 1906-1912, doi:10.1182/blood-2009-07-235721 (2010).
5. Zlotoff, D. A. *et al.* CCR7 and CCR9 together recruit hematopoietic progenitors to the adult thymus. *Blood* **115**, 1897-1905, doi:10.1182/blood-2009-08-237784 (2010).
6. Scimone, M. L., Aifantis, I., Apostolou, I., von Boehmer, H. & von Andrian, U. H. A multistep adhesion cascade for lymphoid progenitor cell homing to the thymus. *Proc Natl Acad Sci U S A* **103**, 7006-7011, doi:10.1073/pnas.0602024103 (2006).
7. Rossi, F. M. *et al.* Recruitment of adult thymic progenitors is regulated by P-selectin and its ligand PSGL-1. *Nat Immunol* **6**, 626-634, doi:10.1038/ni1203 (2005).
8. Buono, M. *et al.* A dynamic niche provides Kit ligand in a stage-specific manner to the earliest thymocyte progenitors. *Nat Cell Biol* **18**, 157-167, doi:10.1038/ncb3299 (2016).
9. Petrie, H. T. & Zuniga-Pflucker, J. C. Zoned out: functional mapping of stromal signaling microenvironments in the thymus. *Annu Rev Immunol* **25**, 649-679, doi:10.1146/annurev.immunol.23.021704.115715 (2007).
10. Thompson, P. K. & Zuniga-Pflucker, J. C. On becoming a T cell, a convergence of factors kick it up a Notch along the way. *Semin Immunol* **23**, 350-359, doi:10.1016/j.smim.2011.08.007 (2011).
11. Hozumi, K. *et al.* Delta-like 4 is indispensable in thymic environment specific for T cell development. *J Exp Med* **205**, 2507-2513, doi:10.1084/jem.20080134 (2008).
12. Hozumi, K. *et al.* Notch signaling is necessary for GATA3 function in the initiation of T cell development. *Eur J Immunol* **38**, 977-985, doi:10.1002/eji.200737688 (2008).
13. Koch, U. *et al.* Delta-like 4 is the essential, nonredundant ligand for Notch1 during thymic T cell lineage commitment. *J Exp Med* **205**, 2515-2523, doi:10.1084/jem.20080829 (2008).
14. Chen, L., Xiao, S. & Manley, N. R. Foxn1 is required to maintain the postnatal thymic microenvironment in a dosage-sensitive manner. *Blood* **113**, 567-574, doi:10.1182/blood-2008-05-156265 (2009).
15. Hale, J. S., Boursalian, T. E., Turk, G. L. & Fink, P. J. Thymic output in aged mice. *Proc Natl Acad Sci U S A* **103**, 8447-8452, doi:10.1073/pnas.0601040103 (2006).
16. Gray, D. H. *et al.* Developmental kinetics, turnover, and stimulatory capacity of thymic epithelial cells. *Blood* **108**, 3777-3785, doi:10.1182/blood-2006-02-004531 (2006).
17. Chinn, I. K., Blackburn, C. C., Manley, N. R. & Sempowski, G. D. Changes in primary lymphoid organs with aging. *Semin Immunol* **24**, 309-320, doi:10.1016/j.smim.2012.04.005 (2012).

Baran-Gale, J. *et al.* Ageing compromises mouse thymus function and remodels epithelial cell differentiation. *Elife* **9**, doi:10.7554/eLife.56221 (2020).

Griffith, A. V., Fallahi, M., Venables, T. & Petrie, H. T. Persistent degenerative changes in thymic organ function revealed by an inducible model of organ regrowth. *Aging Cell* **11**, 169-177, doi:10.1111/j.1474-9726.2011.00773.x (2012).

Lepletier, A. *et al.* Interplay between Follistatin, Activin A, and BMP4 Signaling Regulates Postnatal Thymic Epithelial Progenitor Cell Differentiation during Aging. *Cell Rep* **27**, 3887-3901 e3884, doi:10.1016/j.celrep.2019.05.045 (2019).

Klug, D. B. *et al.* Transgenic expression of cyclin D1 in thymic epithelial precursors promotes epithelial and T cell development. *J Immunol* **164**, 1881-1888, doi:10.4049/jimmunol.164.4.1881 (2000).

Garfin, P. M. *et al.* Inactivation of the RB family prevents thymus involution and promotes thymic function by direct control of Foxn1 expression. *J Exp Med* **210**, 1087-1097, doi:10.1084/jem.20121716 (2013).

Bredenkamp, N., Nowell, C. S. & Blackburn, C. C. Regeneration of the aged thymus by a single transcription factor. *Development* **141**, 1627-1637, doi:10.1242/dev.103614 (2014).

Cheng, L. *et al.* Postnatal tissue-specific disruption of transcription factor FoxN1 triggers acute thymic atrophy. *J Biol Chem* **285**, 5836-5847, doi:10.1074/jbc.M109.072124 (2010).

Zietara, N. *et al.* Multicongenic fate mapping quantification of dynamics of thymus colonization. *J Exp Med* **212**, 1589-1601, doi:10.1084/jem.20142143 (2015).

Okabe, M., Ikawa, M., Kominami, K., Nakanishi, T. & Nishimune, Y. 'Green mice' as a source of ubiquitous green cells. *FEBS Lett* **407**, 313-319, doi:10.1016/s0014-5793(97)00313-x (1997).

Nowotschin, S., Xenopoulos, P., Schrode, N. & Hadjantonakis, A. K. A bright single-cell resolution live imaging reporter of Notch signaling in the mouse. *BMC Dev Biol* **13**, 15, doi:10.1186/1471-213X-13-15 (2013).

Robles, A. I. *et al.* Expression of cyclin D1 in epithelial tissues of transgenic mice results in epidermal hyperproliferation and severe thymic hyperplasia. *Proc Natl Acad Sci U S A* **93**, 7634-7638, doi:10.1073/pnas.93.15.7634 (1996).

Serwold, T., Ehrlich, L. I. & Weissman, I. L. Reductive isolation from bone marrow and blood implicates common lymphoid progenitors as the major source of Thymopoiesis. *Blood* **113**, 807-815, doi:10.1182/blood-2008-08-173682 (2009).

Ki, S. *et al.* Global transcriptional profiling reveals distinct functions of thymic stromal subsets and age-related changes during thymic involution. *Cell Rep* **9**, 402-415, doi:10.1016/j.celrep.2014.08.070 (2014).

Chen, E. L. Y., Thompson, P. K. & Zuniga-Pflucker, J. C. RBPJ-dependent Notch signaling initiates the T cell program in a subset of thymus-seeding progenitors. *Nat Immunol* **20**, 1456-1468, doi:10.1038/s41590-019-0518-7 (2019).

Tikhonova, A. N. *et al.* The bone marrow microenvironment at single-cell resolution. *Nature* **569**, 222-228, doi:10.1038/s41586-019-1104-8 (2019).

Young, K. *et al.* Progressive alterations in multipotent hematopoietic progenitors underlie lymphoid cell loss in aging. *J Exp Med* **213**, 2259-2267, doi:10.1084/jem.20160168 (2016).

Min, H., Montecino-Rodriguez, E. & Dorshkind, K. Reduction in the developmental potential of intrathymic T cell progenitors with age. *J Immunol* **173**, 245-250, doi:10.4049/jimmunol.173.1.245 (2004).

Beerman, I. *et al.* Functionally distinct hematopoietic stem cells modulate hematopoietic lineage potential during aging by a mechanism of clonal expansion. *Proc Natl Acad Sci U S A* **107**, 5465-5470, doi:10.1073/pnas.1000834107 (2010).

Busch, K. *et al.* Fundamental properties of unperturbed haematopoiesis from stem cells in vivo. *Nature* **518**, 542-546, doi:10.1038/nature14242 (2015).

Dorshkind, K., Hofer, T., Montecino-Rodriguez, E., Pioli, P. D. & Rodewald, H. R. Do haematopoietic stem cells age? *Nat Rev Immunol* **20**, 196-202, doi:10.1038/s41577-019-0236-2 (2020).

Montecino-Rodriguez, E. *et al.* Lymphoid-Biased Hematopoietic Stem Cells Are Maintained with Age and Efficiently Generate Lymphoid Progeny. *Stem Cell Reports* **12**, 584-596, doi:10.1016/j.stemcr.2019.01.016 (2019).

Zhu, X. *et al.* Lymphohematopoietic progenitors do not have a synchronized defect with age-related thymic involution. *Aging Cell* **6**, 663-672, doi:10.1111/j.1474-9726.2007.00325.x (2007).

Mackall, C. L., Punt, J. A., Morgan, P., Farr, A. G. & Gress, R. E. Thymic function in young/old chimeras: substantial thymic T cell regenerative capacity despite irreversible age-associated thymic involution. *Eur J Immunol* **28**, 1886-1893, doi:10.1002/(SICI)1521-4141(199806)28:06<1886::AID-IMMU1886>3.0.CO;2-M (1998).

Donskoy, E. & Goldschneider, I. Thymocytopoiesis is maintained by blood-borne precursors throughout postnatal life. A study in parabiotic mice. *J Immunol* **148**, 1604-1612 (1992).

Foss, D. L., Donskoy, E. & Goldschneider, I. The importation of hematogenous precursors by the thymus is a gated phenomenon in normal adult mice. *J Exp Med* **193**, 365-374, doi:10.1084/jem.193.3.365 (2001).

Prockop, S. E. & Petrie, H. T. Regulation of thymus size by competition for stromal niches among early T cell progenitors. *J Immunol* **173**, 1604-1611, doi:10.4049/jimmunol.173.3.1604 (2004).

Lancaster, J. N., Li, Y. & Ehrlich, L. I. R. Chemokine-Mediated Choreography of Thymocyte Development and Selection. *Trends Immunol* **39**, 86-98, doi:10.1016/j.it.2017.10.007 (2018).

Wilson, A., MacDonald, H. R. & Radtke, F. Notch 1-deficient common lymphoid precursors adopt a B cell fate in the thymus. *J Exp Med* **194**, 1003-1012, doi:10.1084/jem.194.7.1003 (2001).

Figure 1. ETP and total thymocyte cellularity declines by 3MO of age. (A) Quantification of total thymocytes and ETPs in C57BL/6J mice at the indicated ages. (B) Linear regression analysis of total thymic cellularity and ETP cellularity at each age. R^2 = coefficient of correlation (C) Ratio of DN2 to ETP cellularity at each age. Symbols represent A) and C) data from individual mice and B) an average of 7-14 mice at each age. Bars represent means +/- SEM from 3-4 independent experiments for each age group. Statistical analysis in (A) was performed using one-way ANOVA with Dunnett's multiple comparisons test where *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 2. The number of available, functional TSP/ETP niches does not decline with age. (A) Schematic of the multicongenic barcoding experiment to quantify available niches in mice of different ages. Lin⁻Flk2⁺CD27⁺ BM progenitors were isolated from 8 different congenic strains (supplemental Figure 2B), mixed at equal ratios and i.v. injected into non-irradiated 1MO, 3MO, 6MO and 12MO mice. After 21 days, the thymi from recipient mice were analyzed by flow cytometry to quantify the number of detectable donor strains that underwent T-cell differentiation. (B) Pie charts illustrate the number and relative frequency of distinct color-coded donor strains in individual recipients of the indicated ages. (C) The estimated number of available functional niches in each age recipient as determined by multinomial sampling. Symbols indicate data from individual mice; bars represent means +/- SEM compiled from 4 independent experiments (n=10-12 mice per recipient age).

Figure 3. 3MO T-cell progenitors show comparable seeding and differentiation capacity relative to 1MO progenitors. (A-C) Donor chimerism in 1MO and 3MO recipient thymuses 21d after transplantation of an equal number of 1MO and 3MO Lin⁻Flk2⁺CD27⁺ BM progenitors into 1MO or 3MO non-irradiated recipients. Quantification of the (A) frequency and (B) number of 1MO and 3MO donor-derived thymocytes in recipients of the indicated ages. (C) Quantification of the percentage of DP thymocytes derived from 1MO and 3MO donors detected in recipient mice of the indicated ages. (D) Representative flow cytometry plots and (E) quantification of the percentage of ETPs that incorporated bromodeoxyuridine (BrdU) 8 hours after i.p. injection in mice of the indicated ages. (A-C,E) Symbols represent data from individual mice. Bars represent mean +/- SEM from 4 independent experiments. p value for (A) was obtained by two-way ANOVA with Sidak's multiple comparison test.

Figure 4. The number of circulating TSPs and BM lymphoid progenitors declines between 1MO and 3MO of age. (A) Representative flow cytometry plots showing the percentage of Flk2⁺CD27⁺ progenitors within the Lin- compartment of the blood in 1MO and 3MO mice. (B) Frequency and absolute cell numbers of circulating lymphoid progenitors in the blood of 1MO and 3MO mice. (C) Representative gating strategy for quantification of Lin⁻Flk2⁺CD27⁺ progenitors, Ly6D⁻ common lymphoid progenitors (CLPs), and Flk2⁺ multipotent progenitors (MPPs) in the BM. (D-F) Frequency and absolute numbers of (D) Flk2⁺CD27⁺ progenitors, (E)

Flk2$^+$ MPPs and (F) Ly6D$^-$ CLPs in BM of 1MO and 3MO mice. Symbols represent data from individual mice at each age. Bars represent mean +/- SEM of data compiled from 3 independent experiments. Statistical analysis was performed using Student *t-test* *p < 0.05, **p <0.01, ***p<0.001, ****p<0.0001.

Figure 5. Sustaining TEC proliferation prevents the age-associated decline in total thymocyte and ETP cellularity, despite a reduction in circulating progenitors by 3MO. Quantification of (A) total thymocyte and (B) ETP cell numbers in WT versus K5.D1 transgenic littermates at the indicated ages. (C) Absolute numbers of circulating lymphoid progenitor cells in the blood of 1MO and 3MO WT and K5.D1 littermates. Quantification of (D) Flk2$^+$ MPP and (E) Ly6D$^-$ CLP cells in the BM of mice at 1MO and 3MO of age. (A-B) symbols indicate the average cell numbers of 7-14 mice pooled from 3-4 independent experiments per age. Error bars show Mean +/- SD. (C-E) Symbols represent data from individual mice. Error bars show Mean +/- SEM for data pooled from 3 independent experiments. Statistical analysis was performed using Student *t-test* **p <0.01, ***p<0.001.

Figure 6. Age-associated changes in Notch signaling correlate with the decline in ETPs at 3MO.

(A) qPCR analysis showing relative *Dll4* expression by EpCAM⁺ Ly51⁺ cTECs from 1MO and 3MO

C57BL/6J mice. Data are pooled from 2 biological experiments with 3-4 technical replicates per experiment, and expression levels were normalized to those at 1MO of age. (B) Representative histogram (left) and quantification of relative fluorescence intensity (RFI) (right) of Notch1 cell surface expression on ETPs. Data are normalized to the average Notch1 MFI of 1MO mice in each individual experiment. Data are pooled from 3 independent experiments. (C-H) Representative flow cytometry plots and the frequencies and RFIs of Notch-Venus reporter expression in (C, D) thymic ETPs, (E,F) BM Flk2$^+$ MPPs and (G,H) BM Ly6D$^-$ CLPs. Symbols represent data from individual mice at each age. Bars represent mean +/- SEM of data pooled from 3 independent experiments. Statistical analysis was performed using Student *t-test* *p < 0.05, ***p<0.001.