Hummingbird blood traits track oxygen availability across space and time

Jessie L. Williamson[1,2,3*], Ethan B. Linck[1], Emil Bautista[4], Ashley Smiley[1,5], Jimmy A.

McGuire[5], Robert Dudley[5], and Christopher C. Witt[1*]

[1]Museum of Southwestern Biology and Department of Biology, University of New Mexico,

Albuquerque, NM, USA

[2]Cornell Lab of Ornithology, Cornell University, Ithaca, NY, USA

[3]Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY, USA

[4]Centro de Ornitología y Biodiversidad (CORBIDI), Lima, Perú

[5]Museum of Vertebrate Zoology and Department of Integrative Biology, University of

California, Berkeley, Berkeley, CA, USA

*Corresponding authors: Jessie L. Williamson (jlw432@cornell.edu) and Christopher C. Witt

(cwitt@unm.edu)

Abstract

Predictable trait variation across environmental gradients suggests that adaptive evolution
repeatedly finds the same solution to a challenge. Trait-environment associations can reflect
long-term, genetic evolution across phylogenies, or short-term, plastic responses of individuals.
When phylogenetic and population-level patterns match, it implies consistency between the long
timescale of adaptation and the short timescale of acclimatization. Alternatively, genetic
adaptation can find solutions that 'break the rules' of trait-environment covariation. For example,
blood-hemoglobin concentration ([Hb]) increases at high elevations in animals, but genetic
adaptations in some populations have been shown to augment tissue oxygenation while curtailing
hemoglobin production, altering this predictable [Hb]-elevation association. Here, we tested
whether species adaptation to elevation generally alters trait-environment relationships for blood.
To do this, we measured blood traits of 1,217 individuals representing 77 species of Andean
hummingbirds, across a 4,600 m elevational gradient. We used hierarchical Bayesian modeling
to estimate blood trait responses to elevation, environmental temperature, precipitation,
individual and species characteristics, and phylogeny. Strikingly, the effects of elevation on [Hb]
and hematocrit (Hct) were nearly identical for individuals and species, implying that rules of
elevational blood variation are set by physics of gas exchange in the hummingbird respiratory
system and are unchanged by species adaptation. However, when we looked at mechanisms of
[Hb] adjustment—by changes in red blood cell size or number—we did find a signal of species
adaptation: To adjust [Hb], species at low and high elevations, respectively, tended to adjust cell
size, whereas species at mid-elevations tended to adjust cell number. Despite scale-independent
elevational variation in [Hb] and Hct, the species-specific balance of red blood cell size versus

38   number appears to have been affected by adaptations that distinguish hummingbird species living

39   at moderate versus high elevations.

41   **Keywords:** Andes, blood-oxygen, cell size, ecophysiology, elevation, hummingbird

Introduction

Predictable variation of traits across environmental gradients can reveal both the power and limitations of adaptive evolution. Trait-environment relationships among species tend to reflect long-term, genetically fixed trait differences across phylogenies; in contrast, trait-environment relationships within species tend to reflect short-term, plastic responses of individuals with shared genetic backgrounds. When within-species and among-species patterns for a specific trait match, it implies a single, shared solution to an environmental challenge. For example, cooler waters at high latitudes are consistently associated with higher numbers of vertebrae in fish, whether comparing species or conspecific individuals (Jordan's Rule) [1,2]. When within-species and among-species patterns conflict, it implies that genetic adaptation has found a qualitatively different solution to the same environmental challenge, altering the trait-environment relationship. The tendency for larger body sizes at colder ambient temperatures among species (Bergmann's Rule) and within species (James's Rule; [3]) is frequently subject to conflicting patterns at these two scales of comparison; it is detectable in only a subset of animal clades, and is reversed in others [4]. These conflicting trait patterns have been attributed to multifarious selection pressures impacting species differences in body size and thermoregulatory physiology [3], and they reflect the expected erosion of trait-environment (or trait-trait) associations over macroevolutionary time [5]. Trait-trait relationships are similarly scale-dependent [6]; for example, individual common milkweed (*Asclepia syriaca*) plants that produce higher concentrations of cardenolide toxins have reduced growth rates [7], while other species of milkweed that produce up to seven times the concentrations of cardenolide toxins as *A. syriaca* show no reduction in growth rate [8]. In this way, genetic adaptations can 'break the rules' that govern trait relationships. Explaining such 'trait gradients' across biological scales will be

essential not only for understanding the functional basis of adaptation, but for predicting the

functional consequence of global change for species and communities [9].

In animals, blood traits associated with oxygen transport change predictably with

declining oxygen (O_2) availability at elevation [10–13]. Most O_2 in the blood is carried by

hemoglobin (Hb) [14], the concentration of which ([Hb]) is a key determinant of blood-O_2

carrying capacity. When lowland-adapted birds ascend to altitude, they experience reduced

arterial O_2 saturation; to compensate, they undergo erythropoiesis, which can increase [Hb] and

therefore blood-O_2 carrying capacity, enhancing O_2 delivery to the tissues and compensating for

reduced arterial O_2 saturation [13]. In the absence of compensatory increases in plasma volume

[15], increased Hb mass leads to elevated hematocrit (Hct) [16]. A suite of interrelated physical

characteristics of blood—total red blood cell count (TRBC), mean cell volume (MCV), mean

cell hemoglobin (MCH), and mean cell hemoglobin concentration (MCHC; Table S1)—can be

modulated to optimize blood-O_2 transport, or to compensate for other cardiac, vascular, or

hematological responses to elevation. For example, MCV typically increases in acclimatized,

lowland-adapted animals undergoing accelerated erythropoiesis [17].

However, genetic adaptation to high elevation can break the 'rules' of trait-environment

covariation in blood traits. For example, evidence from high-altitude-adapted human populations

and theoretical work suggests that optimal [Hb] and Hct levels may be similar to those of sea-

level populations [13,18–20]. Adaptive genetic changes to hemoglobin proteins have been

demonstrated in many high-altitude-native birds, including hummingbirds, House Wrens,

Andean waterfowl, Bar-headed Geese, and a few dozen other bird taxa [21–25]. In

hummingbirds, there is a highly predictable association between species-typical Hb β^A13–β^A83

genotype and elevation, with derived increases in Hb-O_2 affinity in highland lineages and

reductions in Hb-O_2 affinity in lowland lineages [22]. Hummingbirds occurring at high altitudes

exhibit two amino-acid substitutions on the β^A-globin subunit of Hb (Hb β^A-$13_{GLY \rightarrow SER}$ and Hb

β^A-$83_{GLY \rightarrow SER}$) that are known to affect O_2 binding [22]. In particular, β^A-13_{SER} is known from

only four hummingbird genera (*Patagona*, *Chalcostigma*, *Pterophanes*, and *Oreotrochilus*), all

of which occur at extreme high altitudes [21,22,26]. O_2-binding affinity of hemoglobin has

critical effects on tissue oxygenation, and, in birds, binding affinity increases predictably with

elevations [21,27]. It is not yet clear whether the genetic adaptations that change Hb-O_2 binding

affinity predictably affect other blood traits or their covariation with elevation. It is plausible that

species-typical Hb β^A genotypes associated with different O_2-binding affinities induce

predictable differences in blood traits and their reaction norms to elevation; such an association

could reflect a direct functional tradeoff or an indirect effect of accrued, multi-locus adaptation

to elevation at multiple loci.

The macro-physiological 'rules' governing blood trait variation over phylogenetic and

population timescales have remained elusive because adequate comparative data are scarce.

Blood traits are known to vary among avian clades (e.g., [28,29]) and exhibit elevational

variation both within species (e.g., [30–32]) and among species (e.g., [30,31]). At least two

additional recent studies have conducted broad-scale interspecific comparative analyses of avian

[Hb] and Hct data, but both datasets were compiled from heterogeneous published data and

lacked elevational sampling sufficient to estimate elevational effects [33,34]. Furthermore, no

previous comparative analysis of hematological variation across elevation has included the six

key parameters that affect blood-O_2 carrying capacity per unit volume (Table S1). Thus, we

continue to lack a general understanding of within- and among-species blood trait-environment

relationships.

Examining the underlying mechanisms of adjustment for blood-oxygen carrying capacity

can offer insight on trait-environment relationships. [Hb] is equivalent to O_2-carrying capacity

per unit volume of blood, and it can be adjusted via adjustments to cell number (TRBC), cell size

(MCV), within-cell [Hb] (MCHC), or plasma volume. Cell number and cell size are both

measurable components that vary among individuals and species, are subject to plastic and

evolutionary adjustments, and have strong direct effects on [Hb]. Cell number and cell size are

reciprocally constrained by physical space available in the circulatory system, such that

individual birds can have a few large erythrocytes, many small erythrocytes, or an intermediate

state for both traits [35,36]. MCHC tends to be relatively invariant across the vertebrate tree of

life [35], with Hb typically comprising ~1/3 of erythrocytes. Plasma volume adjustments can

decouple Hb mass from [Hb] and, although plasma volume appears variable among human

populations [15], we currently have insufficient data on elevational variation in avian plasma

volume to assess its importance. Quantifying the relative contributions of TRBC and MCV in

driving [Hb] variation within and among species may show that species use different

mechanisms of adjusting blood-O_2 carrying capacity, revealing signatures of adaptation that

distinguish hummingbirds living at moderate versus high elevations.

In this study, we sought to test whether phylogenetic and population patterns match for

blood-trait variation with environment. Specifically, we asked: Does species adaptation change

trait-environment relationships? To address this question, we spent 2006–2020 collecting

specimen-vouchered hematological data from 1,217 wild hummingbirds of 77 species,

representing all 9 clades, and spanning ~4,600 meters in elevation. Using these data, we then

constructed hierarchical Bayesian models to estimate the responses of six blood traits to

elevation, climate, individual and species characteristics, and phylogeny. The hummingbirds are

a potential model clade in which to study rules of hematological variation due to their

documented evolutionary genetic responses to shifts in PO_2, high species richness in mountains,

phylogenetically conserved elevational ranges, and relative uniformity of body size, clutch size,

diet, mating system, and flight mode [21,22,37–39]. Notably, matching O_2 supply and demand is

an acute challenge for all hummingbirds because they exhibit the highest mass-specific

metabolic rates among endotherms, and because they employ sustained hovering, one of the

most energetically demanding forms of animal locomotion [38,40]. This study posits new rules

of elevational blood trait variation, at least for hummingbirds, providing new insights into the

origins and maintenance of trait-environment relationships.

Results and Discussion

We quantified three primary ([Hb], Hct, and TRBC) and three secondary hematological indices

(MCV, MCH, and MCHC; Table S1), and separately analyzed two datasets: an individual-level

dataset, wherein each row contained data from one individual hummingbird (n=1,217), and a

species-level dataset, wherein each row contained mean blood values calculated for each species,

with a minimum of 2 individuals sampled (n=58–65 species, depending on blood trait modeled).

In the individual-level data, mean [Hb] was 18.84 g/dl (±1.65), mean Hct was 58.54% (±5.15),

mean TRBC was 6.37 RBC x 10^6/mm^3 or 6,370,000 cells per µl (±1.16), mean MCV was 93.36

fl (±17.1), mean MCH was 30.4 pg (±5.64), and mean MCHC was 32.44 g/dl (± 2.12). Complete

reference values—the first such values ever published for >90% of our study species—are

reported in detail in Table S2.

Within-species blood trait-environment variation

We used the individual-level dataset to model within species (i.e., interspecific) variation

using multilevel Bayesian models. For each of six blood parameters, we estimated the effects of

elevation, body mass, precipitation, air temperature, species-typical Hb β^A13–β^A83 genotype, as

well as within-species variation in elevational position (an individual's sampled elevation

relative to its species elevational range), body mass, air temperature, and precipitation. We

accounted for between-species differences by incorporating a species grouping variable.

Separately, we excluded sex and wing-loading as potentially important predictors (see Methods).

In total, we compared six models per set using LOOIC and WAIC (Tables S3-S4; Methods). In

addition to fixed effects, models included species as a random effect, and a phylogenetically

correlated random intercept (i.e., covariance matrix of phylogenetic distances among taxa). Thus,

our full model in each set reflects the hypothesis that all predictors, phylogeny, and species

identity influence blood characteristics.

Based on the top-performing model in each set, we found that for every 1,000-m increase

in elevation, [Hb] is expected to increase by 0.39 g/dl, Hct is expected to increase by 1.42

percentage points, TRBC is expected to increase by 310,000 cells per µl, and MCH is expected

to decrease by 0.48 pg. Elevation strongly predicted [Hb], Hct, TRBC, and MCH (Figs 2 A–F

and 3A–F). Elevational position was also a strong predictor of Hct, indicating that individuals

located higher in their species' elevational range tended to have higher Hct (Fig 3B). [Hb],

TRBC, and MCH also increased with increasing elevational position, although these

relationships were subtle (Fig 3A, C, E).

The finding that [Hb], Hct, and TRBC increase with elevation was consistent with

theoretical expectations for O_2-carrying capacity of blood and with prior empirical findings in

birds [26,30,31,33,34], humans [41,42], mammals [11,43], and other taxa [44,45]. In prior work,

substantial variation in slopes of these parameters across elevation may have been attributable to

variation in temporal, taxonomic, and spatial scales of comparison, as well as constraints on the

time-course of evolutionary adaptation [22,42,46]. We expect increases in Hct at higher

elevations because an increase in the total mass of Hb enhances oxygenation under lower

ambient PO_2 [18]; unless there is a proportional increase in plasma volume or MCHC, an

increase in Hb mass is expected to be accompanied by an increase in the proportion of blood

comprised of red blood cells. TRBC also increased with elevation and was, along with MCV,

one of two key drivers of [Hb] adjustment (Fig 2, Fig 4; [38]. MCV did not vary predictably in

our linear models (Fig 2D). Although increasing within-cell [Hb], or MCHC, would seem to be

an alternative mechanism of increasing O_2-carrying capacity, MCHC was less variable than other

blood parameters (Table S2), likely due to physical constraints [47,48]; accordingly, MCH and

MCV were strongly correlated (Table S5; Fig 4C). Contrary to recent empirical findings [30,31],

adjustments to MCHC appeared to be neither substantial nor predictable with respect to elevation

within or among species (Fig 2F).

We found that MCV and MCH decreased in individuals occurring in warmer

environments (Figs 3D, E), and that MCHC decreased in individuals occurring at warmer

localities within a species' range (Fig 3F). The directions of these relationships were consistent

with previous experimental work: Comparisons of juvenile broiler chickens reared from ~1-7

weeks of age in cool, ambient, and hot environments showed decreases in [Hb], Hct, MCV, and

TRBC as temperature increased [49,50]. These findings add to a limited but intriguing pool of

evidence that warmer ambient temperatures may lead to developmental reductions in O_2-carrying

capacity of blood.

Models that included the species grouping variable fit substantially better than those

without; this effect explained 13–21% of the variance in the top model, depending on the model

set (Tables S3-S4; Methods). Moderate effects of the species grouping variable suggest that

variation in blood traits is partly explained by aspects of species biology that evolve too fast to

have resulted in phylogenetic autocorrelation. Species-typical Hb β^A13–β^A83 genotype did not

predict [Hb] or any other blood characteristics (Fig 3A–F), providing no evidence that

evolutionary adjustments to O_2-binding affinity affect the physical composition of blood or its

variation with elevation.

Among-species blood trait-environment variation

Using the species-level dataset, we evaluated the effects of elevation, body mass, species-typical

Hb β^A13–β^A83 genotype, and ambient temperature and precipitation on each of our 6 blood

response variables. We compared sets of models with LOOIC and WAIC (Table S3),

incorporating a quadratic elevation effect for TRBC, MCV, MCH, and MCHC models based on

curvature evident in the data (Fig 2G–L; Methods).

Within- and among-species findings for elevational effects were similar (Fig 3). Based on

the top model in each set, for every 1,000-m increase in elevation, [Hb] is expected to increase

by 0.29 g/dl, and Hct is expected to increase by 1.55 percentage points. Elevation was a

consistently strong predictor of [Hb], Hct, and TRBC, all of which increased with increasing

elevation (Figs 2G–I and 3A–C). The importance of the quadratic effect of elevation for MCV

and MCH, and overall shape of the species-level data, suggest that O_2 carrying capacity

adjustment differs for middle-elevation species versus those found at low and high elevations

(Fig 2J–K; Fig 3D–E). Neither species-typical Hb β^A13–β^A83 genotype, body mass,

environmental temperature, nor precipitation explained variation in any species-mean blood

characteristic (Fig 3).

Mechanisms of blood-O_2 carrying capacity adjustment

The 'banana' shape of the relationship between cell number versus size indicates that the highest

levels of blood-O_2 carrying capacity per unit blood are achieved at high values of TRBC or

MCV, respectively. Our data support that the axis of variation in cell number versus size space is

tightly constrained (Fig 4A–B), mirroring patterns evident from comparisons across multiple

vertebrate classes [35,36]. Individual hummingbirds can possess few large erythrocytes or many

small erythrocytes, but cell number and size must be balanced [36]. Because of the

proportionately greater surface area for gas exchange, having many small cells is predicted to be

advantageous for fast metabolism or when the partial pressure of O_2 in inspired air is low [36].

Similarly, having large cells may be advantageous for modulating the time-course of gas

exchange [51]. One way to shed light on this tradeoff is to test whether individuals and species,

respectively, adjust O_2 carrying capacity per unit blood primarily through increases in cell

number or size.

We designed models to understand the relative contributions of cell number versus size to

blood O_2-carrying capacity per unit volume ([Hb]), within and among species over ~22 Mya of

hummingbird evolution. These models demonstrated that hummingbirds co-equally modulated

cell number and cell size to adjust [Hb] (Table S6). The fact that this was true both within and

among species suggests that developmental and evolutionary fine-tuning of blood-O_2 carrying

capacity occur similarly.

Using data for 32 well-sampled species, we estimated species-specific standardized

TRBC and MCV coefficients (β) in models with [Hb] as the response variable (Fig S1), and we

used these to calculate a 0–1 index that we refer to as the Cell Number-Size Index (hereafter

CeNS). *CeNS* represents the proportional contribution of TRBC versus MCV to increasing [Hb],

calculated as:

252
$$\frac{(\beta_{TRBC})}{(\beta_{TRBC} + \beta_{MCV})}$$

Species specific *CeNS* values ranged from 0.36-0.69 ($\bar{x} = 0.52$), with higher values indicating a

proportionally greater contribution of TRBC than MCV in adjusting [Hb]; conversely, lower

values indicate a proportionally greater contribution of MCV than TRBC.

Extending our analysis, we designed a set of models to test potential predictors of *CeNS*:

elevation, a quadratic elevation effect (*elevation*^{2}; to account for notable curvature in the data),

body mass, wing loading, species-typical Hb β^A13–β^A83 genotype, and phylogeny (see

Methods). Elevation, *elevation*^{2}, and body mass strongly predicted *CeNS* (Fig 4D-F; Table S7).

We found a trend of decreasing *CeNS* with higher wing loading, but the 95% credible intervals

overlapped zero (Fig S2A–B). Species-typical Hb β^A13–β^A83 genotype did not predict *CeNS*

(Fig S3), nor did we find phylogenetic signal in *CeNS* (Fig S4).

Examining the tradeoff between cell number and volume provides a potential explanation

for the striking pattern that middle elevation species tend to adjust [Hb] by different mechanisms

than low and high elevation species (Fig 4E). *CeNS* was uncorrelated with TRBC coefficients

($R^2 = 0.07$, $r = -0.27$) but strongly correlated with MCV coefficients ($R^2 = 0.51$, $r = -0.71$),

indicating that variation in *CeNS* was primarily driven by the degree to which species modulate

cell size (Fig S2, Fig S5. We suggest that because low and high elevation species tend to possess

relatively small cells, they may be more able to rely on shifts in cell size, rather than cell number,

to increase [Hb] (Fig 2J; Fig 4). However, as cell size increased, species were more likely to

approach maximum size for red blood cells, beyond which they could incur a functional

detriment. As a result, species with larger red blood cells were more reliant on cell number to

adjust [Hb]. For middle elevation hummingbird species that have evolved under moderate (~15–

30%) reductions in O_2 availability, TRBC appears to be the primary axis for adjustment of

blood-O_2 carrying capacity per unit volume of blood. Species that have evolved at the highest

elevations are expected to be the best adapted to hypobaric hypoxia, and it is therefore

remarkable that their mode of [Hb] adjustment (primarily by MCV) resembles that of

hummingbird species in the lowlands, the putative ancestral elevation for hummingbirds [37].

Hummingbirds at extreme elevations

Do hummingbirds at extreme high elevations follow the same rules as hummingbirds

distributed at low and moderate elevations with respect to elevational blood variation? Our

models encompassed the entire elevational gradient and therefore could have missed important

phenomena such as threshold effects. To address this, we compared individual hummingbirds

sampled above and below 4,200 m elevation, a threshold that marks the upper ~10% of the

gradient. Hummingbirds sampled from these extreme high elevations had significantly smaller

cells and higher erythrocyte counts than others (Fig S6). This finding suggests that high-altitude

specialist hummingbirds tend to have smaller cells, consistent with findings in other high altitude

vertebrates [11,29,44,52,53]; however, this pattern was only detectable under the extreme

conditions at the uppermost limits of the Andean elevational gradient. We proceeded to formally

test for elevational breakpoints using segmented regression (see Methods). Evidence for

objective blood trait thresholds along the elevational gradient was mixed, with breakpoint

estimates for individual traits ranging from ~1,400 to ~4,000 m (Fig S7). Lack of consilience in

these estimates suggests that trait-specific thresholds are challenging to identify and may not be

generalizable across species and clades.

Phylogenetic signal in blood traits

Phylogenetic signal was low ($\lambda = 0.0$; K range = 0.14–0.21) across all blood traits for both

within- and among-species models. Blomberg's K and Pagel's λ results were consistent with

visual assessment of continuous trait maps, even after removing species with <3 individuals

sampled (Table S8; Fig S8). We additionally found no phylogenetic signal in *CeNS* (Fig S4).

Non-existent to low phylogenetic signal in blood traits within and among hummingbird

species was surprising given demonstrated, phylogenetically conserved Hb genetic adaptations to

elevation [21,22] and in light of strong phylogenetic signal reported from recent comparative

analyses of blood traits that had broader taxonomic sampling [33,34]. However, it is likely that

diverse avian clades possess different blood trait optima; in previous comparative analyses that

included multiple families or orders, phylogenetic signal may have been driven primarily by

differences among clades. For example, Minias et al. [33] reported that small passerines had

consistently high [Hb] and Hct estimates, and that including the shorebird order,

Charadriiformes, which was characterized by high [Hb] but medium Hct values, strongly

influenced patterns of phylogenetic conservatism [33]. We tested for and found no interaction of

clade and elevation on the six studied blood traits, with the exception of the interaction of

elevation and the Hermit clade in the [Hb] model set (Fig S9). Interestingly, hermits are a large,

early diverging hummingbird clade that contains diminishing species diversity with altitude (and

no species that exceeds 3,000 m), and its apparent increased sensitivity of [Hb] to elevation hints

at a functional explanation for this biogeographic tendency. The existence of clade-specific trait

optima [33] and strong species effects, demonstrated in this study, may at first seem difficult to

reconcile with our findings that there was no within-family phylogenetic signal in

hummingbirds; however, we view these findings as being consistent with steady evolution of

trait optima by a mean-reverting, Ornstein-Uhlenbeck process, as described by Minias et al. [33],

and marked by occasional major jumps that coincide with changes in body plan or respiratory

demands.

A strength of our study was that hummingbirds vary only subtly in aspects of their

biology that could plausibly confound our analyses, such as metabolic rate, energy budget, torpor

use, mode of locomotion, foraging strategy, or clutch size. Phylogenetic conservatism in these

traits facilitates explanation of blood trait variation across the entire clade. A strong effect of

species identity in spite of these intrinsic controls hints that species optimize O_2-carrying

functions of blood for subtle changes in lifestyle and environment, and on rapid timescales. This

finding is consistent with the rapid diversification of human hematological responses to

elevation, as illustrated by the copious variation among populations in patterns of elevational

variation [19,41,42,54–56].

Conclusions

Across ~22 Mya of the hummingbird evolution, ambient O_2-availability and species

idiosyncrasies—not phylogenetic affinities—were the primary drivers of variation in blood traits

affecting O_2-carrying functions. In response to elevational changes in O_2 partial pressure,

hummingbirds make predictable adjustments to the physical composition of blood. Remarkably,

after controlling for many potentially confounding variables, decreasing PO_2 drove increases in

[Hb] and Hct that were nearly identical in slope within and among hummingbird species,

respectively. This implies that developmental and evolutionary optimization of blood traits to

ambient PO_2 occurs in fundamentally similar ways, responding to the same set of physical rules

that govern gas exchange at the blood-gas barrier. In addition to O_2 availability, species identity

itself explained 13–21% of blood trait variation, suggesting that fast-evolving, unmodeled

aspects of species biology underpin substantial hematological variation. Our data revealed a

striking constraint space that defines the cell number-size tradeoff for hummingbird erythrocytes;

within this space, individuals and species co-equally modulated cell number and size to adjust

[Hb]. However, species varied in their proportional reliance on adjustments to cell number

versus size (*CeNS*) to increase [Hb], and both elevation and body mass were strongly predictive

of species' tendencies. Middle-elevation species that have been evolving under moderate

hypoxia adjusted [Hb] primarily by adjusting cell number; in contrast, low and high elevation

species tended to adjust [Hb] by adjusting cell size, a pattern partly underpinned by

hummingbirds at extreme elevations exhibiting smaller cells. Together, these results suggest that

evolutionary solutions to reduced O_2 availability can differ qualitatively in conjunction with

species-specific histories of exposure to hypoxia challenges of varying severity.

Materials and Methods

Field Sampling

We collected data from 1,217 individuals of 77 hummingbird species from Peru and Chile from

2006–2020 (Fig 1). After filtering, our final within-species dataset included 1,151 individuals of

77 species from 75 localities. The best-sampled clades were the Coquettes (387 individuals, 20

species) and Brilliants (269 individuals, 19 species). The species with greatest sampling depth

were *Metallura tyrianthina* (95 individuals), *Colibri coruscans* (73 individuals), and *Amazilia*

amazilia and *Patagona gigas* (64 individuals each; Fig 1). Sampling across sexes was balanced

(553 females, 592 males, and 6 unknown). Most birds sampled were adults (936 adults, 179

juveniles, and 36 unknown). Detailed data on specimens, collection localities, and blood traits

are available in Witt et al. (*In prep*) and in the ARCTOS database (arctosdb.org) for

reproducibility and extension of our work.

Blood trait measurements

To calculate primary indices (hemoglobin concentration ([Hb]), hematocrit (Hct), and total red

blood cell count (TRBC)), we obtained whole blood samples within ~5 mins to ~3 hours of

capture by venipuncture on the underside of the wing and collection with heparinized

microcapillary tubes and Hemocue HB201+ cuvettes. Hct (%) was measured with digital calipers

after centrifuging the sealed microcapillary tube for 5 min at 13,000 r.p.m. When >1 Hct sample

was taken, weighted average values were used in analyses. [Hb] (g/dL blood) was measured on

~5 µL of blood using a HemoCue HB201+ hemoglobin photometer. The HemoCue proprietary

photometric method produces values for avian blood that are ~1.0 g/dL greater than those

generated using cyanomethemoglobin spectrophotometry [57]. Accordingly, we corrected [Hb]

values by subtracting 1.0 g/dL. When possible, at the time of blood collection, 10 µl whole blood

was preserved in a 1:200 NaCl dilution to obtain TRBC estimates. Within 12-hours of blood

collection, we pipetted a 10 µl sub-sample of blood-NaCl dilution into a hemocytometer and

allowed cells to settle for 1 min. After focusing the microscope, one or more photos were

obtained of cells in the hemocytometer grid. Later, cells were manually counted in the ImageJ

software [58] and TRBC was estimated following standard protocols [28,59].

After drawing blood, birds were humanely killed by rapid cardiac compression. We

recorded collection elevation, latitude, longitude, sex, body mass, and date. Body mass was

typically recorded prior to blood sampling; however, body mass was corrected for estimated

volume of blood drawn for any individuals massed after blood sampling. Each collected bird was

prepared as a study skin. Specimens and tissues are housed at the Museum of Southwestern

Biology (MSB) at the University of New Mexico, USA, the Centro de Ornitología y

Biodiversidad (CORBIDI) in Peru, and the Universidad Pontificia Católica de Chile in Chile.

Secondary blood indices (mean cell volume (MCV), mean cell hemoglobin (MCH), and

mean cell hemoglobin concentration (MCHC)) were calculated from primary indices (Table S1)

and following standard protocols [28]. We examined distributions of the data using histograms

and qqPlots, and we eliminated outliers that fell outside of the expected distribution and that

differed from known 'typical' values [28,31,60]. To reduce age-related outliers, we examined

values at the tails of each blood trait distribution that were associated with presence of a bursa

measuring ~2 x 2 mm. All juvenile values fell within the expected distribution of blood values.

We did not age individuals by plumage, as juvenile plumage lasts much longer than juvenile

blood values, which stabilize one to several months after fledging.

Environmental and life history traits

We compiled elevational range limits for each species using a combination of published data,

field guides, and specimen records from the MSB [26,61–63]. We calculated each individual's

elevational range position relative to its species elevational range breadth (hereafter 'elevational

position') as *1 – ((maximum species elevation – individual sampling elevation)/species*

elevational range).

We incorporated species-typical Hb β^A13–β^A83 genotypes from published data [22,37].

For 7 species, we identified these genotypes from Hb β^A sequences aligned to a GenBank

reference from Projecto-Garcia et al. [22]. For 12 species (*Amazilia franciae, Campylopterus*

villaviscensio, Eutoxeres aquila, Klais guimeti, Leucippus taczanowskii, Myrmia micrura,

Phaethornis atrimentalis, Sephanoides sephaniodes, Chaetocercus mulsant, Eriocnemis vestita,

Myrtis fanny, and *Oreotrochilus leucopleurus*) species-typical Hb β^A13–β^A83 genotypes were

inferred from sister taxa or nearest relatives.

We characterized climatic variation among latitudinal and elevational gradients using 19

bioclimatic variables from the WorldClim database [64]. We downloaded rasters of each

BioClim variable using the 'raster' package in R (version 4.0.0), cropped them to our study areas

of Peru and Chile, and extracted temperature and precipitation values for each sampling locality.

We used principal components analysis (PCA) for temperature (Bio1–11) and precipitation

(Bio12–19) variables to create a composite measure of temperature and precipitation across the

gradient. PC1 of the temperature variables (hereafter "temperature") explained 73.4% of the

variation in composite temperature, and loadings suggested that this axis corresponded to

increasing temperature across sites (see GitHub). PC1 of the precipitation variables (hereafter

"precipitation") explained 79.8% of the variation in composite precipitation, and loadings

suggested that this axis corresponded to increasing precipitation across sites (see GitHub).

To account for multiple measurement effects (i.e., traits measured from multiple

individuals within species), we calculated between- and within-species differences following

Villemereuil et al. [65]. This approach uses within-group centering to separate each predictor

into two components: We first calculated species mean values (i.e., between-species variability)

and then subtracted the mean value of individual observations (i.e., within-species differences).

These measures of within-species variation for temperature, precipitation, and body mass were

included in models. Because hummingbird body masses may differ by sex, within-species

differences were calculated using mean body mass of males for males, mean body mass of

females for females, and species-level means for individuals of unknown sex (n=6).

We used wing area data from Skandalis et al. [66] to calculate wing loading in g/cm^2,

calculated as species mean body mass (g) divided by total wing area (cm^2). We analyzed wing

loading due its relevance to aerodynamic theory and hummingbird flight at high altitudes [67–

70].

Hummingbird evolutionary history

We used the rate-calibrated molecular hummingbird phylogeny from McGuire et al. [37] as a

backbone to generate a phylogeny pruned to the 77 species in our dataset using the R package

'phytools' [71]. Two species, *Oreotrochilus leucopleurus* and *Phaethornis stuarti*, were missing

from the McGuire et al. [37] tree and were added to our phylogeny using the bind.tip() function.

Branch lengths and position of *O. leucopleurus* were calculated from genetic distances and

recently described sister relationships [72]. Approximate branch length and position of *P. stuarti*

were inferred from BirdTree.org [73,74] with divergence rate estimates from a multi-locus, time-

calibrated phylogeny of Hawaiian honeycreepers [75]. We generated model-specific tree subsets

by including only those tips that corresponded to the species in each modeling subset.

Phylogenetic signal

We initially constructed phylogenetic models (results not shown), and calculated phylogenetic

signal using the full and reduced models for each blood parameter, following the vignette and

recommendations of P. Bürkner (https://cran.r-project.org/web/packages/brms/

vignettes/brms_phylogenetics.html), using the 'hypothesis' method. Estimates of phylogenetic

signal for all model sets (full and reduced in both within- and among species analyses) were low

($\lambda < 0.004$); these estimates are supported by visualizations of continuous trait maps of mean

blood parameters (SI Appendix; Fig S8). Accordingly, we excluded phylogeny as an effect in

models and evaluated phylogenetic signal using Blomberg's K and Pagel's λ. Results from the

two tests were concordant and consistent across model subsets (Table S8).

Within species trait-environment models

We analyzed within species variation in blood characteristics using generalized linear multilevel

models in the R package brms [80]. Brms uses Markov chain Monte Carlo (MCMC) sampling to

obtain draws from posterior distributions. We designed hierarchical models for each of six

continuous responses variables: [Hb], Hct, TRBC, MCV, MCH, and MCHC. For each model,

predictor variables included one species-level categorical variable (species-typical Hb $\beta^A 13$–

$\beta^A 83$ genotype) and eight individual-level continuous variables (elevation, elevational position,

body mass (log-transformed), precipitation, temperature, and within species variation in body

mass, temperature, and precipitation). All variables were standardized to a mean of zero and

standard deviation of one.

We compared 6 models in each model set: 1) an intercept-only null model, 2) an

intercept-only model with a species grouping variable (1|species), 3) a model with all predictors

and no grouping variable, 4) a model with all predictors and the species grouping variable, 5) a

reduced-predictor model with no grouping variable (reduced model 3), 6) a reduced-predictor

model with the species grouping variable (reduced model 4). Reduced models included only

predictors whose 95% credible intervals (CIs) did not overlap zero in full models. Distribution

family differed by model set: Gaussian for [Hb], Hct, and TRBC models; skew-normal for MCV

and MCH models; and student's t-distribution for MCHC models. We ran four Markov chains

for 10,000 iterations with a burn-in period of 5,000, thinned every 10 steps.

We tested for interactions between clade and elevation by comparing three models: 1) the

top within-species model for each blood parameter; 2) the top model + (1|Clade); and 3) the top

model with an elev:Clade interaction. We found no meaningful interaction of elevation and clade

except for elev:CladeHermit in the [Hb] model, whose 95% credible intervals did not overlap

zero (Fig S9). In accordance with lack of phylogenetic signal in our data, the effect of clade

explained minimal variation in model sets for each parameter (Hb=13%; Hct=8%; TRBC=4%;

MCV=2%; MCH=2%; MCHC=4%).

Among species trait-environment models

Final among-species datasets varied in the number of species they contained: ([Hb]=60 species;

Hct=65 species; TRBC=58 species, MCV=60 species; MCH=59 species; MCHC=64 species).

Species mean values were calculated for all species with n=2 or more individuals sampled. We

split *Patagona gigas* into "north" and "south" haplotypes in light of evidence that the complex

may be comprised of two functionally and genetically distinct populations separated by strong

migratory divides [77], and we analyzed data only from individuals sampled during the breeding

months to ensure correct population assignment. We designed model sets for each of our six

blood traits. For each response, we analyzed distributions (qqPlots, Cook's Distance, leverage)

and removed outliers that strongly influenced fit. For each model, predictor variables included

species-typical Hb β^A13–β^A83 genotype, mean elevation, mean body mass (log-transformed; one

mean for all individuals due to balanced sampling between sexes), mean temperature, and mean

precipitation. All variables were standardized to a mean of zero and standard deviation of one.

For each response, we compared: 1) an intercept-only null model, 2) a model with all

predictors, and 3) a reduced-predictor model with only those predictors whose 95% CIs did not

overlap zero in full models (reduced model 2). Given the notable quadratic shape of species

mean data for TRBC, MCV, MCH, and MCHC (Fig 2G–L), for those model sets we additionally

compared 4) a model with all predictors plus a quadratic component of elevation and 5) a

reduced-predictor model with quadratic component including only those predictors whose 95%

CIs did not overlap zero in the full quadratic model (reduced model 4). Distribution family

differed by model set: Gaussian for [Hb; skew-normal for Hct; and student's *t*-distribution for

TRBC, MCV, MCH, and MCHC models. We ran four Markov chains for 20,000 iterations with

a burn-in period of 10,000, thinned every 10 steps.

Cell number versus size analyses

First, to assess within species variation in cell number versus size, we compared a null model

(intercept only) to a single full model *[Hb] ~ MCV + TRBC + (1|species)* for all individuals of

59 species that had [Hb], MCV, and TRBC measurements, distinguishing between *Patagona*

gigas "north" and "south". The species grouping variable allowed us to account for repeated

sampling of individuals within a species and to hold species identity constant. Second, to

evaluate among-species variation in cell number versus size, we generated a mean dataset for

each species with ≥3 individuals sampled (48 species total). We again compared a null (intercept

only) model to a single full model *[Hb] ~ MCV + TRBC*. Third, to assess how individual species

differ in their relative contributions of MCV and TRBC to increasing [Hb], we repeated our

approach for individual species: For each species with ≥8 sample individuals (32 species total),

we generated a species-specific data subset, carefully and conservatively evaluated influential

outlier points, and ran a single model: *[Hb] ~ MCV + TRBC* (Fig S1). For all three model sets

(within species, among species, and single species), both predictors were standardized to a mean

of zero and standard deviation of one.

Using standardized, model-estimated coefficients from each single-species model, for

each species we calculated the Cell Number-Size Index (hereafter *CeNS*) as described in the

main text. *CeNS* ranged from 0.0–1.0 and represented the proportional contribution of TRBC

relative to MCV in driving [Hb], with higher values corresponding to greater relative

contributions of cell number versus cell size.

We then analyzed the factors that contribute to variation in *CeNS*. Our model response

variable was *CeNS* (0–1 index). Predictors, all species mean values, included elevation, a

quadratic elevation term "elevation^2" to account for notable curvature in the data, body mass

(log-transformed), wing loading, and species-typical Hb $\beta^A 13$–$\beta^A 83$ genotype. We compared six

models: 1) an intercept-only null model, 2) a model with core predictors (elevation, body mass,

and wing loading), 3) a model with core predictors plus elevation^2, 4) a model with core

predictors plus elevation^2 and species-typical Hb $\beta^A 13$–$\beta^A 83$ genotype, 5) a reduced version of

model 3, and 6) a reduced version of model 4. Reduced models included only those predictors

whose 95% CIs did not overlap zero in full models. In addition to model fit diagnostics, we

assessed correlations between variables by analyzing correlation matrices and by regressing

CeNS on each R^2 values from standardized species-specific *CeNS* models, unstandardized TRBC

and MCV beta estimates, and raw TRBC and MCV values (Fig S5). Distribution family differed

by model set: Gaussian for individual-level, species mean, and *CeNS* prediction models, and

student's *t*-distribution for species-specific models. For all *CeNS* models we ran four Markov

chains for 20,000 iterations with a burn-in period of 10,000, thinned every 10 steps. We assessed

phylogenetic signal with Blomberg's K, Pagel's λ, and visually with contmaps (Fig S4).

Model comparison and diagnostics

All models and sets were run using the brm() function with default priors (weakly informative

priors for the intercept and sigma; improper flat priors for predictors). To assess convergence we

visually examined trace plots, checked diagnostics, and confirmed that Rhat was <1.01 [76]. We

used a posterior predictive check to assess whether model assumptions yielded good

approximations of the data generating process. There were no problems in any of our models

resulting from Bayesian fraction of missing information or divergent transitions. We compared

models and verified fits using the widely applicable information criterion (WAIC) [78], Bayesian

expected log pointwise predictive densities (ELPD) [79], and approximate leave-one-out cross-

validation information criterion based on the posterior likelihoods (LOOIC; Tables S3, S6). We

estimated fixed effects (means and 95% CIs) from the posterior distributions for each predictor.

For relevant models, we estimated the proportion of the total variance attributed to the species

grouping variable, which accounts for unique aspects of blood parameters that are not captured

by the individual, life history, or environmental predictors.

Breakpoint analysis

We tested for objective thresholds in each blood trait using piecewise linear regression of

elevation (standardized) regressed on each blood trait. In this procedure, we first fit a simple

linear regression model, iteratively searched for objective "breakpoints" along our sampled

elevational gradient, selected the breakpoint with the lowest residual error, and refit the linear

regression model with separate estimates for slopes and intercepts below and above the

breakpoint, respectively. We constrained the model to estimate breakpoints that were each 400 m

from the lowest and highest points in our data (i.e., from 400 m to 4,178 m in elevation), to avoid

biasing breakpoints towards small sample sizes and each low and high elevations, and anomalous

species at both ends of the distribution. We evaluated goodness of fit using fit metrics and by

visually assessing QQplots, Cook's D, leverage, and residual distributions. We compared models

using AIC values. All breakpoint models fit significantly better than simple linear regression

models except for the TRBC breakpoint model; this model did fit better in all respects than the

simple linear regression model, but the fit was not significant at the p=0.05 level. We used a one-

way ANOVA to compare whether individuals below and above model-estimated breakpoints

differed in their blood characteristics (Fig S7).

Data and code availability

All data and code for analyses are available in an open GitHub repository:

https://github.com/jlwilliamson/ComparativeHummingbirdBlood. Specimen data are available

from the Arctos database (https://www.arctosdb.org). Additional detailed data on specimens,

collection localities, and blood values are reported in a forthcoming data paper, Witt et al. (*In*

Prep).

Acknowledgements

We thank Luis Alza, Christopher Barger, Selina Bauernfeind, Matthew Baumann, Elizabeth Beckman, Phred Benham, José Miguel Bogdanovich, Francisco Bozinovic, Homan Castillo Benitez, Jessica Castillo, Marlon Chagua, Jennifer Clark, Mariela Combe, Lida Crooks, Robert Driver, Shane DuBay, Margarita Espinoza, L. Monica Flores, Chauncey Gadek, Ariel Gaffney, Spencer Galen, Avia González-Méndez, Paige Handley, Zachary Hanna, Mike Hartshorne, José Ernesto Huaroto Tornero, Andrew Johnson, Matthew Jones, Daniel Lane, LyAndra Lujan, Mattías Marzfeld, Sabrina McNew, Kyana Montoya, Jano Nuñez-Zapata, Iris Olivas, Paloma Ordoñez, Alessandra Quiñonez, Javier Reinoso, Natalia Ricote, C. Gregory Schmitt, Donna Schmitt, C. Jonathan Schmitt, Frank Solano Bravo, Dora Susanibar, William Talbot, Jorge Tiravanti C., Abraham Urbay T., Thomas Valqui, Walter Vargas Campos, Karen Verde-Guerra, Natalie Wright, and Alfredo Zelada. We are grateful to the following communities in Peru for site access: San Pedro de Casta, Comunidad Campesina Santiago de Carampoma, Ocros, Macate, Oncoy, Yanahara, Huacarpay, Sianbal, Plataforma, Agua Azul, Incahuasi, and Anchicha. This research was funded by the National Science Foundation (DEB-0543556, DEB-1146491, and DBI-1907353), and dissertation research grants to JLW from the Nuttall Ornithological Club Blake-Nuttall Fund Grants (2016 & 2018), the American Philosophical Society Lewis & Clark Grant, the Explorers Club Exploration Fund Grant, American Museum of Natural History Chapman grants (2017 & 2019), the UNM Biology Graduate Student Association Graduate Research Allocations Committee grant, UNM Graduate and Professional Student Association grants (2017–2021), the UNM Latin American & Iberian Institute Field Research Grant, UNM Department of Biology Grove Research Scholarships (2018, 2019, and 2021), and the UNM Department of Biology Melinda Bealmear Memorial Scholarship and Dr. Jones & Dr. Wong

Scholarship. Permits in Chile were granted by Servicio Agrícola y Ganadero (SAG; RE Noºs:

7593/2016, 6817/2017, 7903/2018, 6691/2018, 6692/2018, 6693/2018, 6694/2018). Permits in

Peru were granted by Servicio Nacional Forestal y de Fauna Silvestre (SERFOR; RDG Noºs:

004-2007-INRENA-IFFS-DCB, 135-2009-AG-DGFFS-DGEFFS, 0377-2010-AG-DGFFS-

DGEFFS, 0199-2012-AG-DGFFS-DGEFFS, and 006-2013-MINAGRI-DGFFS/DGEFFS, 244-

2020-MINAGRI-SERFOR/DGGSPFFS-DGSPFS). UNM IACUC approval was granted under

protocols 16-200596-MC and 19-200804-MC.

Author Contributions

622   JLW, EBL, EB, JAM, RD, and CCW designed the research and methods; JLW, EB, AS, JAM,

623   RD, and CCW collected the data; JLW analyzed the data; JLW, JAM, RD, and CCW acquired

624   funds and resources; and JLW and CCW wrote the paper with input from all authors.

Competing interests

627   The authors declare no competing interests.

References

1. McDowall RM. Variation in vertebral number in galaxiid fishes (Teleostei: Galaxiidae): A legacy of life history, latitude and length. Environ Biol Fishes. 2003;66: 361–381. doi:10.1023/A:1023902922843

2. Jordan DS. Relations of temperature to vertebrae among fishes. Proc United States Natl Museum. 1891;14: 107–120. doi:10.5479/si.00963801.14-845.107

3. Blackburn TM, Gaston KJ, Loder N. Geographic gradients in body size: A clarification of Bergmann's rule. Divers Distrib. 1999;5: 165–174. doi:10.1046/j.1472-4642.1999.00046.x

4. De Queiroz A, Ashton KG. The phylogeny of a species-level tendency: Species heritability and possible deep origins of Bergmann's rule in tetrapods. Evolution (N Y). 2004;58: 1674–1684. doi:10.1111/j.0014-3820.2004.tb00453.x

5. McGlothlin JW, Kobiela ME, Wright H V., Mahler DL, Kolbe JJ, Losos JB, et al. Adaptive radiation along a deeply conserved genetic line of least resistance in Anolis lizards. Evol Lett. 2018;2: 310–322. doi:10.1002/evl3.72

6. Agrawal AA. A scale-dependent framework for trade-offs, syndromes, and specialization in organismal biology. Ecology. 2020;101: 1–24. doi:10.1002/ecy.2924

7. Züst T, Rasmann S, Agrawal AA. Growth-defense tradeoffs for two major anti-herbivore traits of the common milkweed Asclepias syriaca. Oikos. 2015;124: 1404–1415. doi:10.1111/oik.02075

8. Agrawal AA, Fishbein M. Phylogenetic escalation and decline of plant defense strategies. Proc Natl Acad Sci U S A. 2008;105: 10057–10060. doi:10.1073/pnas.0802368105

9. Enquist BJ, Norberg J, Bonser SP, Violle C, Webb CT, Henderson A, et al. Scaling from

Traits to Ecosystems: Developing a General Trait Driver Theory via Integrating Trait-

Based and Metabolic Scaling Theories. In: Samraat Pawar, Guy Woodward AID, editor.

Advances in Ecological Research. Elsevier Ltd.; 2015. pp. 249–318.

doi:10.1016/bs.aecr.2015.02.001

10.   Borras A, Cabrera J, Senar JC. Hematocrit Variation in Response to Altitude Changes in

Wild Birds: A Repeated-Measures Design. Condor. 2010;112: 622–626.

doi:10.1525/cond.2010.090113

11.   Tufts DM, Revsbech IG, Cheviron ZA, Weber RE, Fago A, Storz JF. Phenotypic

plasticity in blood-oxygen transport in highland and lowland deer mice. J Exp Biol.

2013;216: 1167–1173. doi:10.1242/jeb.079848

12.   Levett D, Radford E, Menassa D, Graber E, Morash A, Hoppeler H, et al. Acclimatization

of skeletal muscle mitochondria to high-altitude hypoxia during an ascent of Everest.

FASEB J. 2012;26: 1431–1441. doi:10.1096/fj.11-197772

13.   Storz JF, Scott GR, Cheviron ZA. Phenotypic plasticity and genetic adaptation to high-

altitude hypoxia in vertebrates. J Exp Biol. 2010;213: 4125–36. doi:10.1242/jeb.048181

14.   Pittman RN. Chapter 4: Oxygen Transport. Regulation of Tissue Oxygentation. Morgan &

Claypool Life Sciences; 2011. Available:

https://www.ncbi.nlm.nih.gov/books/NBK54103/

15.   Stembridge M, Williams AM, Gasho C, Dawkins TG, Drane A, Villafuerte FC, et al. The

overlooked significance of plasma volume for successful adaptation to high altitude in

Sherpa and Andean natives. Proc Natl Acad Sci U S A. 2019;116: 16177–16179.

doi:10.1073/pnas.1909002116

16.   Storz JF. Genes for High Altitudes. Science (80-). 2010;329: 40–41.

674  doi:10.1126/science.1192481

675 17. Heinicke K, Prommer N, Cajigal J, Viola T, Behn C, Schmidt W. Long-term exposure to

676  intermittent hypoxia results in increased hemoglobin mass, reduced plasma volume, and

677  elevated erythropoietin plasma levels in man. Eur J Appl Physiol. 2003;88: 535–543.

678  doi:10.1007/s00421-002-0732-z

679 18. Scott GR, Milsom WK. Flying high: A theoretical analysis of the factors limiting exercise

680  performance in birds at altitude. Respir Physiol Neurobiol. 2006;154: 284–301.

681  doi:10.1016/j.resp.2006.02.012

682 19. Beall CM. Two routes to functional adaptation: Tibetan and Andean high-altitude natives.

683  Proc Natl Acad Sci. 2007;104: 8655–8660. doi:10.1073/pnas.0701985104

684 20. Simonson TS, Yang Y, Huff CD, Yun H, Qin G, Witherspoon DJ, et al. Genetic Evidence

685  for High-Altitude Adaptation in Tibet. Science (80-). 2010;329: 72–75.

686  doi:10.1126/science.1189406

687 21. Natarajan C, Hoffmann FG, Weber RE, Fago A, Witt CC, Storz JF. Predictable

688  convergence in hemoglobin function has unpredictable molecular underpinnings. Science

689  (80-). 2016;354: 336–339.

690 22. Projecto-Garcia J, Natarajan C, Moriyama H, Weber RE, Fago A, Cheviron ZA, et al.

691  Repeated elevational transitions in hemoglobin function during the evolution of Andean

692  hummingbirds. Proc Natl Acad Sci. 2013;110: 20669–20674.

693 23. Natarajan C, Projecto-Garcia J, Moriyama H, Weber RE, Muñoz-Fuentes V, Green AJ, et

694  al. Convergent Evolution of Hemoglobin Function in High-Altitude Andean Waterfowl

695  Involves Limited Parallelism at the Molecular Sequence Level. PLoS Genet. 2015;11: 1–

696  25. doi:10.1371/journal.pgen.1005681

697   24.   Storz JF, Moriyama H. Mechanisms of hemoglobin adaptation to high altitude hypoxia.

698   High Alt Med Biol. 2008;9: 148–157.

699   25.   Zhu X, Guan Y, Qu Y, David G, Song G, Lei F. Elevational divergence in the great tit

700   complex revealed by major hemoglobin genes. Curr Zool. 2017;64: 455–464.

701   doi:10.1093/cz/zox042

702   26.   Williamson JL, Witt CC. Elevational niche-shift migration: Why the degree of elevational

703   change matters for the ecology, evolution, and physiology of migratory birds.

704   Ornithology. 2021;138: 1–26. doi:10.1093/ornithology/ukaa087

705   27.   Storz JF. Hemoglobin–oxygen affinity in high-altitude vertebrates: is there evidence for an

706   adaptive trend? J Exp Biol. 2016;219: 3190–3203. doi:10.1242/jeb.127134

707   28.   Campbell TW, Ellis CK. Avian and Exotic Animal Hematology and Cytology. Blackwell

708   Publishing Inc.; 2007. doi:10.1017/CBO9781107415324.004

709   29.   Zhang H, Wu CX, Chamba Y, Ling Y. Blood characteristics for high altitude adaptation in

710   Tibetan chickens. Poult Sci. 2007;86: 1384–1389. doi:10.1093/ps/86.7.1384

711   30.   Barve S, Dhondt AA, Mathur VB, Cheviron ZA. Life-history characteristics influence

712   physiological strategies to cope with hypoxia in Himalayan birds. Proc R Soc B.

713   2016;283: 1–8. doi:dx.doi.org/10.1098/rspb.2016.2201

714   31.   Dubay SG, Witt CC. Differential high-altitude adaptation and restricted gene flow across a

715   mid-elevation hybrid zone in Andean tit-tyrant flycatchers. Mol Ecol. 2014;23: 3551–

716   3565. doi:10.1111/mec.12836

717   32.   Linck EB, Williamson JL, Bautista E, Beckman EJ, Benham PM, DuBay SG, et al. Blood

718   variation implicates respiratory limits on elevational ranges of Andean birds. bioRxiv.

719   2021; 1–28. doi:https://doi.org/10.1101/2021.09.30.462673

33.   Minias P. Ecology and evolution of blood oxygen-carrying capacity in birds. Am Nat.

2020;195: 788–801. doi:10.1086/707720

34.   Yap KN, Tsai OH-I, Williams TD. Haematological traits co-vary with migratory status,

altitude and energy expenditure: a phylogenetic, comparative analysis. Sci Rep. 2019;9:

6351. doi:10.1038/s41598-019-42921-4

35.   Wintrobe MM. Variation in the size and hemoglobin content of erythrocytes in the blood

of various vertebrates. Folia Haematol. 1933;51: 32–59.

36.   Hawkey CM, Bennett PM, Gascoyne SC, Hart MG, Kirkwood JK. Erythrocyte size,

number and haemoglobin content in vertebrates. Br J Haematol. 1991;77: 392–397.

doi:10.1111/j.1365-2141.1991.tb08590.x

37.   McGuire JA, Witt CC, Remsen J V, Corl A, Rabosky DL, Altshuler DL, et al. Molecular

phylogenetics and the diversification of hummingbirds. Curr Biol. 2014;24: 1–7.

doi:10.1016/j.cub.2014.03.016

38.   Suarez RK. Hummingbird flight: Sustaining the highest mass-specific metabolic rates

among vertebrates. Experientia. 1992;48: 565–570. doi:10.1007/BF01920240

39.   Graham CH, Parra JL, Rahbek C, McGuire JA. Phylogenetic structure in tropical

hummingbird communities. Proc Natl Acad Sci. 2009;106: 19673–19678.

doi:10.1073/pnas.0912879107

40.   Buermann W, Chaves JA, Dudley R, Mcguire JA, Smith TB, Altshuler DL. Projected

changes in elevational distribution and flight performance of montane Neotropical

hummingbirds in response to climate change. Glob Chang Biol. 2011;17: 1671–1680.

doi:10.1111/j.1365-2486.2010.02330.x

41.   Beall CM, Brittenham GM, Strohl KP, Blangero J, Williams-Blangero S, Goldstein MC,

et al. Hemoglobin concentration of high-altitude Tibetans and Bolivian Aymara. Am J

Phys Anthropol. 1998;106: 385–400.

42. Beall CM. Andean, Tibetan, and Ethiopian patterns of adaptation to high-altitude hypoxia.

Integr Comp Biol. 2006;46: 18–24. doi:10.1093/icb/icj004

43. Hammond KA, Szewczak JOE, Król E. Effects of altitude and temperature on organ

phenotypic plasticity along an altitudinal gradient. J Exp Biol. 2001;204: 1991–2000.

44. Lu S, Xin Y, Tang X, Yue F, Wang H, Bai Y, et al. Differences in hematological traits

between high- and low-altitude lizards (Genus Phrynocephalus). PLoS One. 2015;10: 1–

16. doi:10.1371/journal.pone.0125751

45. Hadley NF, Burns TA. Intraspecific Comparison of the Blood Properties of the Side-

Blotched Lizard, Uta stansburiana. Copeia. 1968;1968: 737–740.

46. Dawson NJ, Alza L, Nandal G, Scott GR, McCracken KG. Convergent changes in muscle

metabolism depend on duration of high-altitude ancestry across andean waterfowl. Elife.

2020;9: 1–35. doi:10.7554/eLife.56259

47. Fischer SL, Fischer SP. Mean Corpuscular Volume. JAMA Intern Med. 1983;143: 282–

283. doi:10.1001/archpedi.1981.02130320078028

48. Rose MS. Epitaph for the M.C.H.C. Br Med J. 1971;4: 169.

49. Donkoh A. Ambient temperature: a factor affecting performance and physiological

response of broiler chickens. Int J Biometeorol. 1989;33: 259–265.

doi:10.1007/BF01051087

50. Moye RJ, Washburn KW, Huston TM. Effects of environmental temperature on

erythrocyte numbers and size. Poult Sci. 1969;48: 1683–1686. doi:10.3382/ps.0481683

51. Glomski CA, Pica A. The Avian Erythrocyte: Its Phylogenetic Odyssey. 1st ed. Enfield,

New Hampshire, USA: Science Publishers; 2011.

52.   Ruiz G, Rosenmann M, Veloso A. Altitudinal distribution and blood values in the toad,

Bufo spinulosus Wiegmann. Comp Biochem Physiol -- Part A Physiol. 1989;94: 643–646.

doi:10.1016/0300-9629(89)90609-9

53.   Yamaguchi K, Jürgens KD, Bartels H, Piiper J. Oxygen transfer properties and

dimensions of red blood cells in high-altitude camelids, dromedary camel and goat. J

Comp Physiol B Biochem Syst Environ Physiol. 1987;157: 1–9. doi:10.1007/BF00702722

54.   Brutsaert TD, Araoz M, Soria R, Spielvogel H, Haas JD. Higher arterial oxygen saturation

during submaximal exercise in Bolivian Aymara compared to European sojourners and

Europeans born and raised at high altitude. Am J Phys Anthropol. 2000;113: 169–181.

doi:10.1002/1096-8644(200010)113:2<169::AID-AJPA3>3.0.CO;2-9

55.   Grimminger J, Richter M, Tello K, Sommer N, Gall H, Ghofrani HA. Thin Air Resulting

in High Pressure: Mountain Sickness and Hypoxia-Induced Pulmonary Hypertension. Can

Respir J. 2017; 1–17. doi:10.1155/2017/8381653

56.   Guo Y-B, He Y-X, Cui C-Y, Ouzhuluobu, Baimakangzhuo, Duojizhuoma, et al. GCH1

plays a role in the high-altitude adaptation of Tibetans. Zool Res. 2017;38: 155–162.

doi:10.24272/j.issn.2095-8137.2017.037

57.   Simmons P, Lill A. Development of parameters influencing blood oxygen carrying

capacity in the welcome swallow and fairy martin. Comp Biochem Physiol - A Mol Integr

Physiol. 2006;143: 459–468. doi:10.1016/j.cbpa.2005.12.018

58.   Rasband WS. ImageJ. Maryland, USA: U.S. National Institutes of Health; Available:

https://imagej.nih.gov/ij/

59.   Samour J. Diagnostic Value of Hematology. In: Harrison G, Lightfoot T, editors. Clinical

Avian Medicine, Volume II. 2005. pp. 587–610. Available:

http://avianmedicine.net/publication_cat/clinical-avian-medicine/

60.   Elarabany N. A comparative study of some haematological and biochemical parameters

between two species from the Anatidae family within migration season. J Basic Appl

Zool. 2018;79. doi:10.1186/s41936-018-0044-4

61.   Parker TAI, Stotz DF, Fitzpatrick JW. Ecological and Distributional Databases. In: D.F.

Stotz, J.W. Fitzpatrick, T.A. Parker, III DKM, editor. Neotropical Birds: Ecology and

Conservation. Chicago, IL, USA: University of Chicago Press; 1996.

62.   Jaramillo A. Birds of Chile. Princeton, New Jersey, USA: Princeton University Press;

2003.

63.   Schulenberg T, Thomas T, Stotz D. Birds of Peru: revised and updated edition. Princeton

Univeristy Press; 2010.

64.   Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution interpolated

climate surfaces for global land areas. Int J Climatol. 2005;25: 1965–1978.

doi:10.1002/joc.1276

65.   Villemereuil P de, Nakagawa S. Modern phylogenetic comparative methods and their

application in evolutionary biology. Mod Phylogenetic Comp Methods their Appl Evol

Biol. 2014; 1–552. doi:10.1007/978-3-662-43550-2

66.   Skandalis DA, Segre PS, Bahlman JW, Groom DJE, Welch KC, Witt CC, et al. The

biomechanical origin of extreme wing allometry in hummingbirds. Nat Commun. 2017;8:

1–9. doi:10.1038/s41467-017-01223-x

67.   Dudley R. Mechanisms and Implications of Animal Flight Maneuverability. Integr Comp

Biol. 2002;42: 135–140. doi:10.1093/icb/42.1.135

68.   Roslyn Dakin, Paolo S. Segre, Andrew D. Straw DLA. Morphology, muscle capacity,

813   skill, and maneuvering ability in hummingbirds. Science (80-). 2018;657: 653–657.

814   69.   Lee SY, Scott GR, Milsom WK. Have wing morphology or flight kinematics evolved for

815   extreme high altitude migration in the bar-headed goose? Comp Biochem Physiol Part C.

816   2008;148: 324–331. doi:10.1016/j.cbpc.2008.05.009

817   70.   Altshuler DL, Dudley R, McGuire JA. Resolution of a paradox: Hummingbird flight at

818   high elevation does not come without a cost. Proc Natl Acad Sci U S A. 2004;101: 17731–

819   17736. doi:10.1073/pnas.0405260101

820   71.   Revell LJ. phytools: an R package for phylogenetic comparative biology (and other

821   things). Methods Ecol Evol. 2012;3: 217–223. doi:10.1111/j.2041-210X.2011.00169.x

822   72.   Sornoza-Molina F, Freile JF, Nilsson J, Krabbe N, Bonaccorso E. A striking, critically

823   endangered, new species of hillstar (Trochilidae: Oreotrochilus) from the southwestern

824   Andes of Ecuador. Auk. 2018;135: 1146–1171. doi:10.1642/AUK-18-58.1

825   73.   Rubolini D, Liker A, Garamszegi LZ, Moller AP, Saino N. Using the BirdTree. org

826   website to obtain robust phylogenies for avian comparative studies: A primer. Curr Zool.

827   2015;61: 959–965.

828   74.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds in

829   space and time. Nature. 2012;491: 444–498. doi:10.1038/nature11631

830   75.   Lerner HRL, Meyer M, James HF, Hofreiter M, Fleischer RC. Multilocus resolution of

831   phylogeny and timescale in the extant adaptive radiation of Hawaiian honeycreepers. Curr

832   Biol. 2011;21: 1838–1844. doi:10.1016/j.cub.2011.09.039

833   76.   Bürkner PC. brms: An R package for Bayesian multilevel models using Stan. J Stat Softw.

834   2017;80. doi:10.18637/jss.v080.i01

835   77.   Williamson JL, Witt CC. A lightweight backpack harness for tracking hummingbirds. J

836   Avian Biol. 2021; 1–9. doi:10.1111/jav.02802

837   78.   Watanabe S. Asymptotic equivalence of Bayes cross validation and widely applicable

838   information criterion in singular learning theory. J Mach Learn Res. 2010;11: 3571–3594.

839   79.   Vehtari A, Gelman A, Gabry J. Practical Bayesian model evaluation using leave-one-out

840   cross-validation and WAIC. Stat Comput. 2017; 1413–1432.

Fig 1. Elevational, latitudinal, and phylogenetic sampling from 2006–2020. **(A)** Sampling localities spanned 28° latitude and 4,578 meters in elevation in Peru and Chile. **(B)** We studied 77 species ('n' column indicates sampling depth) from all nine major hummingbird clades. Sampling (gray opaque points) is shown across each species' elevational range (colored horizontal bars). Time-calibrated phylogeny is modified from McGuire et al. [37]. Illustrations depicting a representative from each clade are from *Birds of the World* and are reproduced courtesy of the Cornell Lab of Ornithology (©).

Fig 2. Relationship between elevation and the six studied blood parameters (hemoglobin concentration ([Hb]), hematocrit (Hct), total red blood cell count (TRBC), mean cell volume (MCV), mean cell hemoglobin content (MCH), and mean cell hemoglobin concentration (MCHC). **(A–F)** Within species comparisons, wherein each point represents a value from a single individual. **(G–L)** Among species comparisons, wherein each point represents a mean value from a single species. In all panels, solid trend lines correspond to relationships where 95% posterior credible intervals in models did not overlap zero, while dashed trend lines correspond to relationships where 89% credible intervals in models did not overlap zero (see Fig 3). Credible intervals for both elevation (at the 95% level) and *elevation²* (at the 89% level) did not overlap zero in the TRBC model (Panel I). Colors correspond to clades in Fig 1. R^2 values are reported from Bayesian models reported in Fig 3.

Fig 3. Posterior probabilities and credible intervals from full within-species (main plots) and among-species (insets) models for each of six blood traits: **(A)** hemoglobin concentration ([Hb]), **(B)** hematocrit (Hct), **(C)** total red blood cell (TRBC), **(D)** mean cell volume (MCV), **(E)** mean cell hemoglobin (MCH), **(F)** mean cell hemoglobin concentration (MCHC). Circles indicate beta estimates, thick and thin bars illustrate 89% and 95% credible intervals, respectively. Red circles denote parameters that did not overlap zero in top-ranked models. For Hb, Hct, and TRBC within-species analyses, the full model was the top model. See Tables S3-S4 for model comparison and parameter estimates from top models in other model sets.

Fig 4. Theoretical and empirical relationships among hummingbird blood parameters underlying

hemoglobin concentration ([Hb]) adjustment and predictors of the Cell Number-Size Index

(*CeNS*). **(A–B)** Hummingbirds have evolved optimized blood-O_2 carrying capacity, achieved by

balancing total red blood cell count (TRBC) with mean cell volume (MCV). **(A)** Predictions for

O_2 carrying-capacity. Background is shaded by O_2 carrying-capacity, calculated as *MCV x TRBC*

x 1/3 scaling factor to represent constant MCHC. **(B)** Empirical findings. Points are colored by

[Hb], with lighter colors representing higher concentrations. **(C)** Mean cell hemoglobin (MCH)

and MCV are tightly linked, suggesting little variance in mean cell hemoglobin concentration

(MCHC). Points are colored by TRBC (units presented on the y-axis), with lighter colors

representing higher erythrocyte counts. In panels A–C, each point represents a value from a

single individual. **(C)** Posterior probabilities and credible intervals from the best fitting reduced

CeNS model. **(D)** *CeNS* shows a strong quadratic relationship with elevation, revealing that

middle elevation species adjust blood-O_2 carrying capacity differently than low- and high-

elevation species. **(E)** *CeNS* increases with increasing body mass. In panels E–F, R^2 values are

reported from generalized linear models of each x regressed on y, and thus do not account for

other modeled variables.