Main Manuscript for

Oxygen availability drives blood traits and the cell number-size tradeoff across Andean hummingbirds

Jessie L. Williamson[1*], Ethan B. Linck[1], Emil Bautista[2], Ashley Smiley[1,3], Jimmy A. McGuire[3], Robert Dudley[3], and Christopher C. Witt[1*]

[1]Museum of Southwestern Biology and Department of Biology, University of New Mexico, 302 Yale Blvd NE, Albuquerque, NM 87131, USA
[2]Centro de Ornitología y Biodiversidad (CORBIDI), Lima, Perú
[3]Museum of Vertebrate Zoology and Department of Integrative Biology, University of California, Berkeley, Berkeley, CA 94720, USA

*Corresponding authors: Jessie L. Williamson and Christopher C. Witt

Email: williamson@unm.edu and cwitt@unm.edu

Author Contributions: JLW, EBL, EB, JAM, RD, CCW designed the research and methods; JLW conducted formal analysis; JLW, JAM, RD, CCW acquired funds and resources; JLW, EB, AS, JAM, RD, CCW collected the data; and JLW and CCW wrote the paper with input from all authors.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences/Ecology.

Keywords: Andes, blood-oxygen, cell, elevation, hummingbird

This PDF file includes:
>   Main Text
>   Figures 1 to 4

ABSTRACT

For endotherms with sustained high metabolism, like hummingbirds, blood-oxygen (O_2) carrying

capacity should be finely tuned to supply O_2 to respiring tissues. Hematological adjustments are

expected in response to changes in O_2 availability that occur along elevational gradients;

however, eco-physiological rules by which this variation occurs have remained elusive. If

evolutionary divergence in elevational niches is accompanied by predictable shifts in blood trait

reaction norms, we would expect different patterns of elevational variation for within-species

versus among-species comparisons; but this has not been systematically tested. Here, we sought

to elucidate these rules using hummingbirds as a model clade. We measured blood traits that

affect O_2-carrying functions: total hemoglobin concentration ([Hb]), hematocrit (Hct),

erythrocyte count (TRBC), mean cell volume (MCV), mean cellular hemoglobin content (MCH),

and mean cellular hemoglobin concentration (MCHC). For 1,217 Andean hummingbirds

representing 77 species, spanning sea level to 4,578 meters in elevation, we modeled blood traits

as responses to environmental conditions and individual and species traits. [Hb] and Hct

increased with elevation similarly within species and across the phylogeny—for every 1,000-m

ascent, [Hb] increased by one third of a gram per deciliter and Hct increased by one and a half

percentage points, corresponding to about three-hundred thousand additional cells per microliter.

Hummingbirds adjusted blood cell number and cell size by similar proportions to modulate

blood-O_2 carrying capacity; however, species at the lowest and highest elevations relied

relatively more on adjustments to cell volume, revealing qualitatively different evolutionary

responses elicited by moderate versus extreme reductions in O_2 availability.

SIGNIFICANCE STATEMENT

Trait variation along environmental gradients reveals how individuals adjust or populations

evolve to improve performance, foreshadowing climate change effects. For blood traits such as

hemoglobin concentration ([Hb]), rules of elevational variation have remained elusive; it is

unknown whether species adaptation affects reaction of blood phenotypes to local oxygen

availability. We tested this in a model clade, hummingbirds, that have undergone many

elevational shifts. Elevational variation was predictable, but similar within and among species,

indicating no predictable evolution of reaction norms. However, for the cell number-size tradeoff

that underpins [Hb], species varied by elevational range. Cell number explained [Hb] at

intermediate elevations, while cell size was more important at extremes, indicating qualitative

differences in respiratory adaptation to moderate versus high elevations.

INTRODUCTION

Organisms that sustain high levels of aerobic activity under hypoxia, such as birds flying at high

altitude, must supply oxygen (O_2) to respiring tissues despite reduced O_2 availability in 'thin' air.

To achieve this, the O_2 transport functions of blood must be finely tuned to match O_2 supply and

demand under a variety of ambient conditions (1–6). Most O_2 in the blood is carried by

hemoglobin (Hb; (7), the concentration of which ([Hb]) is a key determinant of blood-O_2

carrying capacity. Under hypobaric hypoxia, animals experience reduced arterial O_2 saturation;

to compensate, they undergo erythropoiesis, which can increase [Hb] and therefore blood-O_2

carrying capacity, enhancing O_2 delivery to the tissues and compensating for reduced arterial O_2

saturation (8). In the absence of compensatory increases in plasma volume (9), increased Hb

mass leads to elevated hematocrit (Hct) and, beyond a certain threshold, corresponding increases

in blood viscosity will begin to compromise O_2 transport (10). A suite of interrelated physical

characteristics of blood—total red blood cell count (TRBC), mean cell volume (MCV), mean

cell hemoglobin (MCH), and mean cell hemoglobin concentration (MCHC; Table S1)—can be

modulated to optimize blood-O_2 transport, or to compensate for other cardiac, vascular, or

hematological responses to elevation.

Birds are well-known for their abilities to maintain tissue oxygenation in hypoxic

environments, which can be attributed to both plastic and constitutive aspects of respiratory

physiology that enhance O_2 uptake, transport, and delivery (3, 11–19). Along with many factors

that have been shown to affect the physical characteristics of avian blood, including age, sex,

body condition, hormones, environmental factors, and disease (20, 21), high-altitude hypoxia has

predictable short-term effects on blood composition that are mediated by the hypoxia-inducible

factor (HIF) pathway (8, 22–24). Accordingly, when lowland-adapted birds ascend to altitude,

HIF-induced production of erythropoietin is expected to increase [Hb] and Hct. Birds that are

chronically exposed to high elevations have been found to display either normal or elevated

[Hb], Hct, and MCHC (25, 26). Evidence from high-altitude-adapted human populations and

theoretical work suggests that optimal [Hb] and Hct levels may be similar to those of sea-level

populations (8, 13, 27). MCV typically increases in acclimatized, lowland-adapted animals

undergoing accelerated erythropoiesis (28); however, bird and other vertebrate lineages that have

evolved for extended periods at altitude have often been found to possess smaller cells, likely as

part of a suite of adaptive traits for efficient tissue oxygenation (23, 29, 30). Sensitivity of blood

traits to low partial pressure of O_2 on the short timescale of acclimatization is well-documented

(8, 23). Over longer timescales, natural selection on genes that affect hypoxia performance leads

predictably to genetic adaptation (18, 19, 31, 32), but it is not known how this evolutionary

process may affect the reaction norms that govern elevational variation in blood traits.

Systematic trait variation across environmental gradients reflects the outcome of

evolutionary and developmental processes, and explaining such 'trait gradients' will be essential

to predicting the functional consequence of climate warming for species and communities (33).

Despite its critical importance for tissue oxygenation, many questions remain unanswered

regarding elevational variation in blood traits that underpin blood-O_2 carrying capacity. Blood

traits vary among avian clades (e.g., 20, 33) and exhibit elevational variation both within species

(e.g., 4, 25, 34) and among species (e.g., 4, 25, 33, 35). Trait shifts with elevation are expected to

differ systematically for conspecific versus heterospecific comparisons; whereas within-species

variation should be driven primarily by developmental plasticity, among-species variation should

be influenced by genetic changes that have accrued during differences in elevational exposure

lasting thousands to millions of generations (37). Linck et al. (35) estimated elevational effects

on [Hb] and Hct within each of 136 species of Andean birds and showed that, within the median

species, [Hb] increased by 0.7 g/dL per 1,000 m in elevation, while Hct increased by 2.0

percentage points per 1,000 m. At least two additional recent studies have conducted broad-scale

interspecific comparative analyses of avian [Hb] and Hct data, but both datasets were compiled

from heterogeneous published data and lacked elevational sampling sufficient to estimate

elevational effects (34, 38). Furthermore, no previous comparative analysis of hematological

variation across elevation has included the six key parameters that affect blood-O_2 carrying

capacity per unit volume (Table S1). We continue to lack a general understanding of within- and

among-species effects of elevation on blood traits, and specifically, an understanding of whether

different rules govern elevational variation at these different levels of biological organization.

89   Another outstanding question is the degree to which known genetic adaptations to

hematological function may predict blood-O_2 carrying capacity. Adaptive genetic changes to

hemoglobin proteins have been demonstrated in many high-altitude-native birds, including

hummingbirds, House Wrens, Andean waterfowl, Bar-headed Geese, and a few dozen other bird

taxa (18, 19, 39–41). In hummingbirds, there is a highly predictable association between species-

typical Hb $\beta^A 13$–$\beta^A 83$ genotype and elevation, with derived increases in Hb-O_2 affinity in

highland lineages and reductions in Hb-O_2 affinity in lowland lineages (19). Hummingbirds

occurring at high altitudes exhibit two amino-acid substitutions on the β^A-globin subunit of Hb

(Hb β^A-$13_{GLY \to SER}$ and Hb β^A-$83_{GLY \to SER}$) that are known to affect O_2 binding (19). In particular,

β^A-13_{SER} is known from only four hummingbird genera (*Patagona*, *Chalcostigma*, *Pterophanes*,

and *Oreotrochilus*), all of which occur at extreme high altitudes (3, 18, 19). Both O_2-binding

affinity and blood-O_2 carrying capacity have critical effects on tissue oxygenation, but are part of

a complex O_2 uptake, transport, and delivery system that contains multiple convective and

diffusive steps (42). It is plausible that species-typical Hb β^A genotypes associated with different

O_2-binding properties induce predictable differences in blood traits and their reaction norms to

elevation; such an association could reflect a direct functional tradeoff or an indirect effect of

accrued genetic adaptation to elevation at multiple loci.

106   The constraints on blood trait values that are imposed by physical tradeoffs and the

relative contributions of each blood trait to overall blood-O_2 carrying capacity remain poorly

understood. [Hb] defines O_2-carrying capacity per unit volume of blood, and it can be adjusted

via adjustments to cell number (TRBC), cell size (MCV), within-cell [Hb] (MCHC), or plasma

volume. MCHC tends to be relatively invariant across the vertebrate tree of life, as demonstrated

nearly a century ago by Wintrobe (43), with Hb typically comprising ~1/3 of erythrocytes.

Plasma volume adjustments can decouple Hb mass from [Hb] and, although plasma volume

appears to vary among human populations (9), we currently have insufficient data on elevational

variation in avian plasma volume to assess its importance. Cell number and cell size are both

measurable components that are variable among individuals and species, subject to plastic and

evolutionary adjustments, and have strong direct effects on [Hb]. Cell number and cell size are

reciprocally constrained by physical space available in the circulatory system, such that

individual birds can have a few large erythrocytes, many small erythrocytes, or an intermediate

state for both traits (43, 44). Quantifying the relative contributions of TRBC and MCV in driving

[Hb] variation within and among species may reveal that species utilize different mechanisms of

adjusting blood-O_2 carrying capacity; if such mechanisms are predictably associated with

elevational environments or evolutionary histories of altitude exposure, they could lead to new

biogeographic insights.

In this study, we sought to elucidate the rules governing variation in blood traits that

affect O_2-carrying functions, including the effects of environment, individual characteristics, and

evolved species differences. Specifically, we asked: 1) To what extent do blood traits shift

predictably with elevational changes in O_2 availability, while controlling for other sources of

variation?; 2) Do blood traits vary with O_2 availability differently across developmental (within

species) and evolutionary (among species) time scales?; 3) How are blood traits affected by

phylogenetic history and evolved species differences, such as body mass, metabolic intensity,

and known genetic variants that affect Hb-O_2 binding affinity?; 4) To what extent are individual

and species variation in O_2-carrying capacity per unit volume of blood attributable to changes in

cell number versus cell size? And 5) does manifestation of the cell number-size tradeoff vary

predictably among species?

We collected specimen-vouchered hematological data from hummingbirds between

2006–2020 across a ~4,600-meter elevational gradient in the Andes. Data for six blood traits

from 1,217 individuals of 77 species were used to model within- and among-species variation,

respectively (Figure 1). Hummingbirds could be considered a model clade in which to study

hematological variation due to their documented evolutionary genetic responses to shifts in $P$$O_2$,

high species richness in mountains, phylogenetically conserved elevational ranges, and relative

uniformity of body size, clutch size, diet, mating system, and flight mode (18, 19, 45–47).

Notably, matching O_2 supply and demand is an acute challenge for all hummingbirds because

they exhibit the highest mass-specific metabolic rates among endotherms, and because they

employ sustained hovering, one of the most energetically demanding forms of animal

locomotion (46, 48).

RESULTS AND DISCUSSION

We quantified three primary ([Hb], Hct, and TRBC) and three secondary hematological indices
(MCV, MCH, and MCHC; Table S1), and separately analyzed two datasets: an individual-level
dataset, wherein each row contained data from one individual hummingbird (*n*=1,217), and a
species-level dataset, wherein each row contained mean blood values calculated for each species,
with a minimum of 2 individuals sampled (*n*=58–65 species, depending on blood trait modeled).
In the individual-level data, mean [Hb] was 18.84 g/dl (±1.65), mean Hct was 58.54% (±5.15),
mean TRBC was 6.37 RBC x 10^6/mm^3 or 6,370,000 cells per µl (±1.16), mean MCV was 93.36
fl (±17.1), mean MCH was 30.4 pg (±5.64), and mean MCHC was 32.44 g/dl (± 2.12). Complete
reference values—the first such values ever published for >90% of our study species—are
reported in detail in Table S2.

Within-species blood comparison

We used the individual-level dataset to model within species variation and accounted for
between-species differences by incorporating a species grouping variable. Using multilevel
Bayesian models for each of the six blood parameters, we estimated the effects of elevation,
body mass, precipitation, air temperature, species-typical Hb $\beta^A 13$–$\beta^A 83$ genotype, as well as
within-species variation in elevational position (an individual's sampled elevation relative to its
species elevational range), body mass, air temperature, and precipitation. Separately, we
excluded sex and wing-loading as potentially important predictors (see Methods). In total, we
compared six models per set using LOOIC and WAIC (Tables S3-S4; Methods). In addition to
fixed effects, models included species as a random effect, and a phylogenetically correlated
random intercept (i.e., covariance matrix of phylogenetic distances among taxa). Thus, our full

model in each set reflects the hypothesis that all predictors, phylogeny, and species identity

influence blood characteristics.

Based on the top-performing model in each set, we found that for every 1,000-m increase

in elevation, [Hb] is expected to increase by 0.39 g/dl, Hct is expected to increase by 1.42

percentage points, TRBC is expected to increase by 310,000 cells per µl, and MCH is expected

to decrease by 0.48 pg. Elevation strongly predicted [Hb], Hct, TRBC, and MCH (Figures 2 A–F

and 3A–F). Elevational position was also a strong predictor of Hct, indicating that individuals

located higher in their species' elevational range tended to have higher Hct (Figure 3B). [Hb],

TRBC, and MCH also increased with increasing elevational position, although these

relationships were subtle (Figure 3A, C, E).

The finding that [Hb], Hct, and TRBC increase with elevation was consistent with

theoretical expectations for O_2-carrying capacity of blood and with prior empirical findings in

birds (3, 4, 25, 34, 38), humans (49, 50), mammals (23, 51), and other taxa (52, 53). In prior

work, substantial variation in slopes of these parameters across elevation may have been

attributable to variation in temporal, taxonomic, and spatial scales of comparison, as well as

constraints on the time-course of evolutionary adaptation (19, 32, 50). We expect increases in

Hct at higher elevations because an increase in the total mass of Hb enhances oxygenation under

lower ambient PO_2 (13); unless there is a proportional increase in plasma volume or MCHC, an

increase in Hb mass is expected to be accompanied by an increase in the proportion of blood

comprised of red blood cells. TRBC also increased with elevation and was, along with MCV,

one of two key drivers of [Hb] adjustment (Figure 2, Figure 4; 45). MCV did not vary

predictably in our linear models (Figure 2D). Although increasing within-cell [Hb], or MCHC,

would seem to be an alternative mechanism of increasing O_2-carrying capacity, MCHC was less

variable than other blood parameters (Table S2), likely due to physical constraints (54, 55);

accordingly, MCH and MCV were strongly correlated (Table S5; Figure 4C). Contrary to recent

empirical findings (4, 25), adjustments to MCHC appeared to be neither substantial nor

predictable with respect to elevation within or among species (Figure 2F).

We found that MCV and MCH decreased in individuals occurring in warmer

environments (Figures 3D, E), and that MCHC decreased in individuals occurring at warmer

localities within a species' range (Figure 3F). The directions of these relationships were

consistent with previous experimental work: Comparisons of juvenile broiler chickens reared

from ~1-7 weeks of age in cool, ambient, and hot environments showed decreases in [Hb], Hct,

MCV, and TRBC as temperature increased (56, 57). Our findings add to a limited but intriguing

pool of evidence that warmer ambient temperatures may lead to developmental reductions in O_2-

carrying capacity of blood.

Models that included the species grouping variable fit substantially better than those

without; this effect explained 13–21% of the variance in the top model, depending on the model

set (Tables S3-S4; Methods). Moderate effects of the species grouping variable suggest that

variation in blood traits is partly explained by aspects of species biology that evolve too fast to

have resulted in phylogenetic autocorrelation. Species-typical Hb $\beta^A 13$–$\beta^A 83$ genotype did not

predict [Hb] or any other blood characteristics (Figure 3A–F), providing no evidence that

evolutionary adjustments to O_2-binding affinity affect the physical composition of blood or its

variation with elevation.

Among-species blood comparison

Using the species-level dataset, we evaluated the effects of elevation, body mass, species-typical

Hb $\beta^A13–\beta^A83$ genotype, and ambient temperature and precipitation on each of our 6 blood

response variables. We compared sets of models with LOOIC and WAIC (Table S3),

incorporating a quadratic elevation effect for TRBC, MCV, MCH, and MCHC models based on

curvature evident in the data (Figure 2G–L; Methods).

Within- and among-species findings for elevational effects were similar (Figure 3). Based

on the top model in each set, for every 1,000-m increase in elevation, [Hb] is expected to

increase by 0.29 g/dl, and Hct is expected to increase by 1.55 percentage points. Elevation was a

consistently strong predictor of [Hb], Hct, and TRBC, all of which increased with increasing

elevation (Figures 2G–I and 3A–C). The importance of the quadratic effect of elevation for

MCV and MCH, and overall shape of the species-level data, suggest that O_2 carrying capacity

adjustment differs for middle-elevation species versus those found at low and high elevations

(Figure 2J–K; Figure 3D–E). Neither species-typical Hb $\beta^A13–\beta^A83$ genotype, body mass,

environmental temperature, nor precipitation explained variation in any species-mean blood

characteristic (Figure 3).

Influence of cell number versus size on blood-O_2 carrying capacity

The axis of variation in cell number versus size space was tightly constrained (Figure 4A–B); the

'banana' shape of Figure 4A–B mirrors patterns from Wintrobe (43) and Hawkey et al. (44),

which present similar plots based on species representing multiple vertebrate classes. The shape

of this relationship indicates that the highest levels of blood-O_2 carrying capacity per unit blood

are achieved at high values of TRBC or MCV, respectively. Individual hummingbirds can

possess few large erythrocytes or many small erythrocytes, but cell number and size must be

balanced (44). Because of the proportionately greater surface area for gas exchange, having

many small cells is predicted to be advantageous for fast metabolism or when the partial pressure

of O_2 in inspired air is low (44). Similarly, having large cells may be advantageous for

modulating the time-course of gas exchange (58). One way to shed light on this tradeoff is to test

whether individuals and species, respectively, adjust O_2 carrying capacity per unit blood

primarily through increases in cell number or size.

We designed models to understand the relative contributions of cell number versus size to

blood O_2-carrying capacity per unit volume ([Hb]), within and among species over ~22 Mya of

hummingbird evolution. These models demonstrated that hummingbirds co-equally modulated

cell number and cell size to adjust [Hb] (Table S6). The fact that this was true both within and

among species suggests that developmental and evolutionary fine-tuning of blood-O_2 carrying

capacity occur similarly.

Using data for 32 well-sampled species, we estimated species-specific standardized

TRBC and MCV coefficients (β) in models with [Hb] as the response variable (Figure S1), and

we used these to calculate a 0–1 index that we refer to as the Cell Number-Size Index (hereafter

CeNS). *CeNS* represents the proportional contribution of TRBC versus MCV to increasing [Hb],

calculated as:

255
$$\frac{(\beta_{TRBC})}{(\beta_{TRBC} + \beta_{MCV})}$$

Species specific *CeNS* values ranged from 0.36-0.69 ($\bar{x} = 0.52$), with higher values indicating a

proportionally greater contribution of TRBC than MCV in adjusting [Hb]; conversely, lower

values indicate a proportionally greater contribution of MCV than TRBC.

Extending our analysis, we designed a set of models to test potential predictors of *CeNS*:

elevation, a quadratic elevation effect (elevation^2; to account for notable curvature in the data),

body mass, wing loading, species-typical Hb β^A13–β^A83 genotype, and phylogeny (see

Methods). Elevation, elevation^2, and body mass strongly predicted *CeNS* (Figure 4D-F; Table

S7). We found a trend of decreasing *CeNS* with higher wing loading, but the 95% credible

intervals overlapped zero (Figure S2A–B). Species-typical Hb β^A13–β^A83 genotype did not

predict *CeNS* (Figure S3), nor did we find phylogenetic signal in *CeNS* (Figure S4).

Examining the tradeoff between cell number and volume provides a potential explanation

for the striking pattern that middle elevation species tend to adjust [Hb] by different mechanisms

than low and high elevation species (Figure 4E). *CeNS* was uncorrelated with TRBC coefficients

($R^2 = 0.07$, $r = -0.27$) but strongly correlated with MCV coefficients ($R^2 = 0.51$, $r = -0.71$),

indicating that variation in *CeNS* was primarily driven by the degree to which species modulate

cell size (Figure S2, Figure S5. We suggest that because low and high elevation species tend to

possess relatively small cells, they may be more able to rely on shifts in cell size, rather than cell

number, to increase [Hb] (Figure 2J; Figure 4). However, as cell size increased, species were

more likely to approach maximum size for red blood cells, beyond which they could incur a

functional detriment. As a result, species with larger red blood cells were more reliant on cell

number to adjust [Hb]. For middle elevation hummingbird species that have evolved under

moderate (~15–30%) reductions in O_2 availability, TRBC appears to be the primary axis for

adjustment of blood-O_2 carrying capacity per unit volume of blood. Species that have evolved at

the highest elevations are expected to be the best adapted to hypobaric hypoxia, and it is

therefore remarkable that their mode of [Hb] adjustment (primarily by MCV) resembles that of

hummingbird species in the lowlands, the putative ancestral elevation for hummingbirds (45).

Hummingbirds at extreme elevations

Do hummingbirds at extreme high elevations follow the same rules as hummingbirds

distributed at low and moderate elevations with respect to elevational blood variation? Our

models encompassed the entire elevational gradient and therefore could have missed important

phenomena such as threshold effects. To address this, we compared individual hummingbirds

sampled above and below 4,200 m elevation, a threshold that marks the upper ~10% of the

gradient. Hummingbirds sampled from these extreme high elevations had significantly smaller

cells and higher erythrocyte counts than others (Figure S6). This finding suggests that high-

altitude specialist hummingbirds tend to have smaller cells, consistent with findings in other high

altitude vertebrates (23, 29, 30, 52, 59); however, this pattern was only detectable under the

extreme conditions at the uppermost limits of the Andean elevational gradient. We proceeded to

formally test for elevational breakpoints using segmented regression (see Methods). Evidence for

objective blood trait thresholds along the elevational gradient was mixed, with breakpoint

estimates for individual traits ranging from ~1,400 to ~4,000 m (Figure S7). Lack of consilience

in these estimates suggests that trait-specific thresholds are challenging to identify and may not

be generalizable across species and clades.

Phylogenetic signal in blood characteristics

Phylogenetic signal was low ($\lambda = 0.0$; *K* range = 0.14–0.21) across all blood traits for both

within- and among-species models. Blomberg's K and Pagel's λ results were consistent with

visual assessment of continuous trait maps, even after removing species with <3 individuals

sampled (Table S8; Figure S8). We additionally found no phylogenetic signal in *CeNS* (Figure

S4).

Non-existent to low phylogenetic signal in blood traits within and among hummingbird

species was surprising given demonstrated, phylogenetically conserved Hb genetic adaptations to

elevation (18, 19) and in light of strong phylogenetic signal reported from recent comparative

analyses of blood traits that had broader taxonomic sampling (34, 38). However, it is likely that

diverse avian clades possess different blood trait optima; in previous comparative analyses that

included multiple families or orders, phylogenetic signal may have been driven primarily by

differences among clades. For example, Minias et al. (34) reported that small passerines had

consistently high [Hb] and Hct estimates, and that including the shorebird order,

Charadriiformes, which was characterized by high [Hb] but medium Hct values, strongly

influenced patterns of phylogenetic conservatism (34). We tested for and found no interaction of

clade and elevation on the six studied blood traits, with the exception of the interaction of

elevation and the Hermit clade in the [Hb] model set (Figure S9). Interestingly, hermits are a

large, early diverging hummingbird clade that contains diminishing species diversity with

altitude (and no species that exceeds 3,000 m), and its apparent increased sensitivity of [Hb] to

elevation hints at a functional explanation for this biogeographic tendency. The existence of

clade-specific trait optima (34) and strong species effects, demonstrated in this study, may at first

seem difficult to reconcile with our findings that there was no within-family phylogenetic signal

in hummingbirds; however, we view these findings as being consistent with steady evolution of

trait optima by a mean-reverting, Ornstein-Uhlenbeck process, as described by Minias et al. (34),

and marked by occasional major jumps that coincide with changes in body plan or respiratory

demands.

A strength of our study was that hummingbirds vary only subtly in aspects of their

biology that could plausibly confound our analyses, such as metabolic rate, energy budget, torpor

use, mode of locomotion, foraging strategy, or clutch size. Phylogenetic conservatism in these

traits facilitates explanation of blood trait variation across the entire clade. A strong effect of

species identity in spite of these intrinsic controls hints that species optimize O_2-carrying

functions of blood for subtle changes in lifestyle and environment, and on rapid timescales. This

finding is consistent with the rapid diversification of human hematological responses to

elevation, as illustrated by the copious variation among populations in patterns of elevational

variation (27, 49, 50, 60–62).

Conclusions

Across ~22 Mya of the hummingbird evolution, ambient O_2-availability and species

idiosyncrasies—not phylogenetic affinities—were the primary drivers of variation in blood traits

affecting O_2-carrying functions. In response to elevational changes in O_2 partial pressure,

hummingbirds make predictable adjustments to the physical composition of blood. Remarkably,

after controlling for many potentially confounding variables, decreasing PO_2 drove increases in

[Hb] and Hct that were nearly identical in magnitude within and among hummingbird species,

respectively. This implies that developmental and evolutionary optimization of blood traits to

ambient PO_2 occurs in fundamentally similar ways, responding to the same set of physical rules

that govern gas exchange at the blood-gas barrier. In addition to O_2 availability, species identity

itself explained 13–21% of blood trait variation, suggesting that fast-evolving, unmodeled

aspects of species biology underpin substantial hematological variation. Our data revealed a

striking constraint space that defines the cell number-size tradeoff for hummingbird erythrocytes;

within this space, individuals and species co-equally modulated cell number and size to adjust

[Hb]. However, species varied in their proportional reliance on adjustments to cell number

versus size (*CeNS*) to increase [Hb], and both elevation and body mass were strongly predictive

of species' tendencies. Middle-elevation species that have been evolving under moderate

hypoxia adjusted [Hb] primarily by adjusting cell number; in contrast, low and high elevation

species tended to adjust [Hb] by adjusting cell size, a pattern partly underpinned by

hummingbirds at extreme elevations exhibiting smaller cells. Together, these results suggest that

evolutionary solutions to reduced O_2 availability can differ qualitatively in conjunction with

species-specific histories of exposure to hypoxia challenges of varying severity.

MATERIALS AND METHODS

Field Sampling

We collected data from 1,217 individuals of 77 hummingbird species from Peru and Chile from

2006–2020 (Figure 1). After filtering, our final within-species dataset included 1,151 individuals

of 77 species from 75 localities. The best-sampled clades were the Coquettes (387 individuals,

20 species) and Brilliants (269 individuals, 19 species). The species with greatest sampling depth

were *Metallura tyrianthina* (95 individuals), *Colibri coruscans* (73 individuals), and *Amazilia*

amazilia and *Patagona gigas* (64 individuals each; Figure 1). Sampling across sexes was

balanced (553 females, 592 males, and 6 unknown). Most birds sampled were adults (936 adults,

179 juveniles, and 36 unknown). Detailed data on specimens, collection localities, and blood

traits are available in Witt et al. (*In prep*) and in the ARCTOS database (arctosdb.org) for

reproducibility and extension of our work.

Hematological measurements

To calculate primary indices (hemoglobin concentration ([Hb]), hematocrit (Hct), and total red

blood cell count (TRBC)), we obtained whole blood samples within ~5 mins to ~3 hours of

capture by venipuncture on the underside of the wing and collection with heparinized

microcapillary tubes and Hemocue HB201+ cuvettes. Hct (%) was measured with digital calipers

after centrifuging the sealed microcapillary tube for 5 min at 13,000 r.p.m. When >1 Hct sample

was taken, weighted average values were used in analyses. [Hb] (g/dL blood) was measured on

~5 µL of blood using a HemoCue HB201+ hemoglobin photometer. The HemoCue proprietary

photometric method produces values for avian blood that are ~1.0 g/dL greater than those

generated using cyanomethemoglobin spectrophotometry (63). Accordingly, we corrected [Hb]

values by subtracting 1.0 g/dL. When possible, at the time of blood collection, 10 µl whole blood

was preserved in a 1:200 NaCl dilution to obtain TRBC estimates. Within 12-hours of blood

collection, we pipetted a 10 µl sub-sample of blood-NaCl dilution into a hemocytometer and

allowed cells to settle for 1 min. After focusing the microscope, one or more photos were

obtained of cells in the hemocytometer grid. Later, cells were manually counted in the ImageJ

software (64) and TRBC was estimated following standard protocols (20, 21).

After drawing blood, birds were humanely killed by rapid cardiac compression. We

recorded collection elevation, latitude, longitude, sex, body mass, and date. Body mass was

typically recorded prior to blood sampling; however, body mass was corrected for estimated

volume of blood drawn for any individuals massed after blood sampling. Each collected bird was

prepared as a study skin. Specimens and tissues are housed at the Museum of Southwestern

Biology (MSB) at the University of New Mexico, USA, the Centro de Ornitología y

Biodiversidad (CORBIDI) in Peru, and the Universidad Pontificia Católica de Chile in Chile.

Secondary blood indices (mean cell volume (MCV), mean cell hemoglobin (MCH), and

mean cell hemoglobin concentration (MCHC)) were calculated from primary indices (Table S1)

and following standard protocols (20). We examined distributions of the data using histograms

and qqPlots, and we eliminated outliers that fell outside of the expected distribution and that

differed from known 'typical' values (20, 25, 65). To reduce age-related outliers, we examined

values at the tails of each blood trait distribution that were associated with presence of a bursa

measuring ~2 x 2 mm. All juvenile values fell within the expected distribution of blood values.

We did not age individuals by plumage, as juvenile plumage lasts much longer than juvenile

blood values, which stabilize one to several months after fledging.

Environmental and life history variables

We compiled elevational range limits for each species using a combination of published data,

field guides, and specimen records from the MSB (3, 66–68). We calculated each individual's

elevational range position relative to its species elevational range breadth (hereafter 'elevational

position') as *1 – ((maximum species elevation – individual sampling elevation)/species*

elevational range).

We incorporated species-typical Hb $\beta^A 13$–$\beta^A 83$ genotypes from published data (19, 45).

For 7 species, we identified these genotypes from Hb β^A sequences aligned to a GenBank

reference from Projecto-Garcia et al. (19). For 12 species (*Amazilia franciae, Campylopterus*

villaviscensio, Eutoxeres aquila, Klais guimeti, Leucippus taczanowskii, Myrmia micrura,

Phaethornis atrimentalis, Sephanoides sephaniodes, Chaetocercus mulsant, Eriocnemis vestita,

Myrtis fanny, and *Oreotrochilus leucopleurus*) species-typical Hb $\beta^A 13$–$\beta^A 83$ genotypes were

inferred from sister taxa or nearest relatives.

We characterized climatic variation among latitudinal and elevational gradients using 19

bioclimatic variables from the WorldClim database (69). We downloaded rasters of each

BioClim variable using the 'raster' package in R (version 4.0.0), cropped them to our study areas

of Peru and Chile, and extracted temperature and precipitation values for each sampling locality.

We used principal components analysis (PCA) for temperature (Bio1–11) and precipitation

(Bio12–19) variables to create a composite measure of temperature and precipitation across the

gradient. PC1 of the temperature variables (hereafter "temperature") explained 73.4% of the

variation in composite temperature, and loadings suggested that this axis corresponded to

increasing temperature across sites (see GitHub). PC1 of the precipitation variables (hereafter

"precipitation") explained 79.8% of the variation in composite precipitation, and loadings

suggested that this axis corresponded to increasing precipitation across sites (see GitHub).

To account for multiple measurement effects (i.e., traits measured from multiple

individuals within species), we calculated between- and within-species differences following

Villemereuil et al. (70). This approach uses within-group centering to separate each predictor

into two components: We first calculated species mean values (i.e., between-species variability)

and then subtracted the mean value of individual observations (i.e., within-species differences).

These measures of within-species variation for temperature, precipitation, and body mass were

included in models. Because hummingbird body masses may differ by sex, within-species

differences were calculated using mean body mass of males for males, mean body mass of

females for females, and species-level means for individuals of unknown sex (n=6).

We used wing area data from the 59 MSB species studied in Skandalis et al. (71) to

calculate wing loading in g/cm^2, calculated as species mean body mass (g) divided by total wing

area (cm^2). We analyzed wing loading due its relevance to aerodynamic theory and hummingbird

flight at high altitudes (72–75).

Hummingbird phylogeny

We used the rate-calibrated molecular hummingbird phylogeny from McGuire et al. (45) as a

backbone to generate a phylogeny pruned to the 77 species in our dataset using the R package

'phytools' (76). Two species, *Oreotrochilus leucopleurus* and *Phaethornis stuarti*, were missing

from the McGuire et al. (45) tree and were added to our phylogeny using the bind.tip() function.

Branch lengths and position of *O. leucopleurus* were calculated from genetic distances and

recently described sister relationships (77). Approximate branch length and position of *P. stuarti*

were inferred from BirdTree.org (78, 79) with divergence rate estimates from a multi-locus,

time-calibrated phylogeny of Hawaiian honeycreepers (80). We generated model-specific tree

subsets by including only those tips that corresponded to the species in each modeling subset.

Phylogenetic signal

We initially constructed phylogenetic models (results not shown), and calculated phylogenetic

signal using the full and reduced models for each blood parameter, following the vignette and

recommendations of P. Bürkner (https://cran.r-project.org/web/packages/brms/

vignettes/brms_phylogenetics.html), using the 'hypothesis' method. Estimates of phylogenetic

signal for all model sets (full and reduced in both within- and among species analyses) were low

($\lambda < 0.004$); these estimates are supported by visualizations of continuous trait maps of mean

blood parameters (SI Appendix; Figure S8). Accordingly, we excluded phylogeny as an effect in

models and evaluated phylogenetic signal using Blomberg's K and Pagel's λ. Results from the

two tests were concordant and consistent across model subsets (Table S8).

Within species models

We analyzed within species variation in blood characteristics using generalized linear multilevel

models in the R package brms (81). Brms uses Markov chain Monte Carlo (MCMC) sampling to

obtain draws from posterior distributions. We designed hierarchical models for each of six

continuous responses variables: [Hb], Hct, TRBC, MCV, MCH, and MCHC. For each model,

predictor variables included one species-level categorical variable (species-typical Hb β^A13–

β^A83 genotype) and eight individual-level continuous variables (elevation, elevational position,

body mass (log-transformed), precipitation, temperature, and within species variation in body

mass, temperature, and precipitation). All variables were standardized to a mean of zero and

standard deviation of one.

We compared 6 models in each model set: 1) an intercept-only null model, 2) an

intercept-only model with a species grouping variable (1|species), 3) a model with all predictors

and no grouping variable, 4) a model with all predictors and the species grouping variable, 5) a

reduced-predictor model with no grouping variable (reduced model 3), 6) a reduced-predictor

model with the species grouping variable (reduced model 4). Reduced models included only

predictors whose 95% credible intervals (CIs) did not overlap zero in full models. Distribution

family differed by model set: Gaussian for [Hb], Hct, and TRBC models; skew-normal for MCV

and MCH models; and student's *t*-distribution for MCHC models. We ran four Markov chains

for 10,000 iterations with a burn-in period of 5,000, thinned every 10 steps.

We tested for interactions between clade and elevation by comparing three models: 1) the

top within-species model for each blood parameter; 2) the top model + (1|Clade); and 3) the top

model with an elev:Clade interaction. We found no meaningful interaction of elevation and clade

except for elev:CladeHermit in the [Hb] model, whose 95% credible intervals did not overlap

zero (Figure S9). In accordance with lack of phylogenetic signal in our data, the effect of clade

explained minimal variation in model sets for each parameter (Hb=13%; Hct=8%; TRBC=4%;

MCV=2%; MCH=2%; MCHC=4%).

Among species models

Final among-species datasets varied in the number of species they contained: ([Hb]=60 species;

Hct=65 species; TRBC=58 species, MCV=60 species; MCH=59 species; MCHC=64 species).

Species mean values were calculated for all species with n=2 or more individuals sampled. We

split *Patagona gigas* into "north" and "south" haplotypes in light of evidence that the complex

may be comprised of two functionally and genetically distinct populations separated by strong

migratory divides (82), and we analyzed data only from individuals sampled during the breeding

months to ensure correct population assignment. We designed model sets for each of our six

blood traits. For each response, we analyzed distributions (qqPlots, Cook's Distance, leverage)

and removed outliers that strongly influenced fit. For each model, predictor variables included

species-typical Hb β^A13–β^A83 genotype, mean elevation, mean body mass (log-transformed; one

mean for all individuals due to balanced sampling between sexes), mean temperature, and mean

precipitation. All variables were standardized to a mean of zero and standard deviation of one.

For each response, we compared: 1) an intercept-only null model, 2) a model with all

predictors, and 3) a reduced-predictor model with only those predictors whose 95% CIs did not

overlap zero in full models (reduced model 2). Given the notable quadratic shape of species

mean data for TRBC, MCV, MCH, and MCHC (Figure 2G–L), for those model sets we

additionally compared 4) a model with all predictors plus a quadratic component of elevation and

5) a reduced-predictor model with quadratic component including only those predictors whose

95% CIs did not overlap zero in the full quadratic model (reduced model 4). Distribution family

differed by model set: Gaussian for [Hb; skew-normal for Hct; and student's *t*-distribution for

TRBC, MCV, MCH, and MCHC models. We ran four Markov chains for 20,000 iterations with

a burn-in period of 10,000, thinned every 10 steps.

Cell number versus size analyses

First, to assess within species variation in cell number versus size, we compared a null model

(intercept only) to a single full model *[Hb] ~ MCV + TRBC + (1|species)* for all individuals of

59 species that had [Hb], MCV, and TRBC measurements, distinguishing between *Patagona*

gigas "north" and "south". The species grouping variable allowed us to account for repeated

sampling of individuals within a species and to hold species identity constant. Second, to

evaluate among-species variation in cell number versus size, we generated a mean dataset for

each species with ≥3 individuals sampled (48 species total). We again compared a null (intercept

only) model to a single full model *[Hb] ~ MCV + TRBC.* Third, to assess how individual species

differ in their relative contributions of MCV and TRBC to increasing [Hb], we repeated our

approach for individual species: For each species with ≥8 sample individuals (32 species total),

we generated a species-specific data subset, carefully and conservatively evaluated influential

outlier points, and ran a single model: *[Hb] ~ MCV + TRBC* (Figure S1). For all three model sets

(within species, among species, and single species), both predictors were standardized to a mean

of zero and standard deviation of one.

Using standardized, model-estimated coefficients from each single-species model, for

each species we calculated the Cell Number-Size Index (hereafter *CeNS*) as described in the

main text. *CeNS* ranged from 0.0–1.0 and represented the proportional contribution of TRBC

relative to MCV in driving [Hb], with higher values corresponding to greater relative

contributions of cell number versus cell size.

We then analyzed the factors that contribute to variation in *CeNS*. Our model response

variable was *CeNS* (0–1 index). Predictors, all species mean values, included elevation, a

quadratic elevation term "elevation^2" to account for notable curvature in the data, body mass

(log-transformed), wing loading, and species-typical Hb β^A13–β^A83 genotype. We compared six

models: 1) an intercept-only null model, 2) a model with core predictors (elevation, body mass,

and wing loading), 3) a model with core predictors plus elevation^2, 4) a model with core

predictors plus elevation^2 and species-typical Hb β^A13–β^A83 genotype, 5) a reduced version of

model 3, and 6) a reduced version of model 4. Reduced models included only those predictors

whose 95% CIs did not overlap zero in full models. In addition to model fit diagnostics, we

assessed correlations between variables by analyzing correlation matrices and by regressing

CeNS on each R^2 values from standardized species-specific *CeNS* models, unstandardized TRBC

and MCV beta estimates, and raw TRBC and MCV values (Figure S5). Distribution family

differed by model set: Gaussian for individual-level, species mean, and *CeNS* prediction models,

and student's *t*-distribution for species-specific models. For all *CeNS* models we ran four Markov

chains for 20,000 iterations with a burn-in period of 10,000, thinned every 10 steps. We assessed

phylogenetic signal with Blomberg's K, Pagel's λ, and visually with contmaps (Figure S4).

Model comparison and diagnostics

All models and sets were run using the brm() function with default priors (weakly informative

priors for the intercept and sigma; improper flat priors for predictors). To assess convergence we

visually examined trace plots, checked diagnostics, and confirmed that Rhat was <1.01 (81). We

used a posterior predictive check to assess whether model assumptions yielded good

approximations of the data generating process. There were no problems in any of our models

resulting from Bayesian fraction of missing information or divergent transitions. We compared

models and verified fits using the widely applicable information criterion (WAIC; (83), Bayesian

expected log pointwise predictive densities (ELPD; (84), and approximate leave-one-out cross-

validation information criterion based on the posterior likelihoods (LOOIC; Tables S3, S6). We

estimated fixed effects (means and 95% CIs) from the posterior distributions for each predictor.

For relevant models, we estimated the proportion of the total variance attributed to the species

grouping variable, which accounts for unique aspects of blood parameters that are not captured

by the individual, life history, or environmental predictors.

Breakpoint analysis

We tested for objective thresholds in each blood trait using piecewise linear regression of

elevation (standardized) regressed on each blood trait. In this procedure, we first fit a simple

linear regression model, iteratively searched for objective "breakpoints" along our sampled

elevational gradient, selected the breakpoint with the lowest residual error, and refit the linear

regression model with separate estimates for slopes and intercepts below and above the

breakpoint, respectively. We constrained the model to estimate breakpoints that were each 400 m

from the lowest and highest points in our data (i.e., from 400 m to 4,178 m in elevation), to avoid

biasing breakpoints towards small sample sizes and each low and high elevations, and anomalous

species at both ends of the distribution. We evaluated goodness of fit using fit metrics and by

visually assessing QQplots, Cook's D, leverage, and residual distributions. We compared models

using AIC values. All breakpoint models fit significantly better than simple linear regression

models except for the TRBC breakpoint model; this model did fit better in all respects than the

simple linear regression model, but the fit was not significant at the p=0.05 level. We used a one-

way ANOVA to compare whether individuals below and above model-estimated breakpoints

differed in their blood characteristics (Figure S7).

Data and code availability

Specimen information is available from the Arctos database (https://www.arctosdb.org). Detailed

data on specimens, collection localities, and blood values are reported in a forthcoming data

paper, Witt et al. (*In Prep*). All data and code for analyses are available in an open GitHub

repository: https://github.com/jlwilliamson/ComparativeHummingbirdBlood.

ACKNOWLEDGEMENTS

We thank Luis Alza, Christopher Barger, Selina Bauernfeind, Matthew Baumann, Elizabeth

Beckman, Phred Benham, José Miguel Bogdanovich, Francisco Bozinovic, Homan Castillo

Benitez, Jessica Castillo, Marlon Chagua, Jennifer Clark, Mariela Combe, Lida Crooks, Robert

Driver, Shane DuBay, Margarita Espinoza, L. Monica Flores, Chauncey Gadek, Ariel Gaffney,

Spencer Galen, Avia González-Méndez, Paige Handley, Zachary Hanna, Mike Hartshorne, José

Ernesto Huaroto Tornero, Andrew Johnson, Matthew Jones, Daniel Lane, LyAndra Lujan,

Mattías Marzfeld, Sabrina McNew, Kyana Montoya, Jano Nuñez-Zapata, Iris Olivas, Paloma

Ordoñez, Alessandra Quiñonez, Javier Reinoso, Natalia Ricote, C. Gregory Schmitt, Donna

Schmitt, C. Jonathan Schmitt, Frank Solano Bravo, Dora Susanibar, William Talbot, Jorge

Tiravanti C., Abraham Urbay T., Thomas Valqui, Walter Vargas Campos, Karen Verde-Guerra,

Natalie Wright, and Alfredo Zelada. We are grateful to the following communities in Peru for

site access: Comunidades de San Pedro de Casta, Campesina Santiago de Carampoma, Ocros,

Macate, Oncoy, Yanahara, Huacarpay, Sianbal, Plataforma, Agua Azul, Incahuasi, and

Anchicha. This research was funded by the National Science Foundation (DEB-0543556, DEB-

1146491, and DBI-1907353), and dissertation research grants to JLW from the Nuttall

Ornithological Club Blake-Nuttall Fund Grants (2016 & 2018), the American Philosophical

Society Lewis & Clark Grant, the Explorers Club Exploration Fund Grant, American Museum of

Natural History Chapman grants (2017 & 2019), the UNM Biology Graduate Student

Association Graduate Research Allocations Committee grant, UNM Graduate and Professional

Student Association grants (2017–2021), the UNM Latin American & Iberian Institute Field

Research Grant, UNM Department of Biology Grove Research Scholarships (2018, 2019, and

2021), and the UNM Department of Biology Melinda Bealmear Memorial Scholarship and Dr.

Jones & Dr. Wong Scholarship. Permits in Chile were granted by Servicio Agrícola y Ganadero

(SAG; RE Noºs: 7593/2016, 6817/2017, 7903/2018, 6691/2018, 6692/2018, 6693/2018,

6694/2018). Permits in Peru were granted by Servicio Nacional Forestal y de Fauna Silvestre

(SERFOR; RDG Noºs: 004-2007-INRENA-IFFS-DCB, 135-2009-AG-DGFFS-DGEFFS, 0377-

2010-AG-DGFFS-DGEFFS, 0199-2012-AG-DGFFS-DGEFFS, and 006-2013-MINAGRI-

DGFFS/DGEFFS, 244-2020-MINAGRI-SERFOR/DGGSPFFS-DGSPFS). UNM IACUC

approval was granted under protocols 16-200596-MC and 19-200804-MC.

REFERENCES

1.   J. S. Krause, *et al.*, Annual Hematocrit Profiles in Two Subspecies of White-Crowned Sparrow: A Migrant and a Resident Comparison. *Physiol. Biochem. Zool.* **89**, 51–60 (2015).

2.   J. F. Storz, G. R. Scott, Life Ascending: Mechanism and Process in Physiological Adaptation to High-Altitude Hypoxia. *Annu. Rev. Ecol. Evol. Syst.* **50**, 1–24 (2019).

3.   J. L. Williamson, C. C. Witt, Elevational niche-shift migration: Why the degree of elevational change matters for the ecology, evolution, and physiology of migratory birds. *Ornithology* **138**, 1–26 (2021).

4.   S. Barve, A. A. Dhondt, V. B. Mathur, Z. A. Cheviron, Life-history characteristics influence physiological strategies to cope with hypoxia in Himalayan birds. *Proc. R. Soc. B* **283**, 1–8 (2016).

5.   S. L. Laguë, High-altitude champions: birds that live and migrate at altitude. *J. Appl. Physiol.* **123**, 942–950 (2017).

6.   J. F. Storz, Hemoglobin–oxygen affinity in high-altitude vertebrates: is there evidence for an adaptive trend? *J. Exp. Biol.* **219**, 3190–3203 (2016).

7.   R. N. Pittman, "Chapter 4: Oxygen Transport" in *Regulation of Tissue Oxygentation*, (Morgan & Claypool Life Sciences, 2011) (June 23, 2021).

8.   J. F. Storz, G. R. Scott, Z. A. Cheviron, Phenotypic plasticity and genetic adaptation to high-altitude hypoxia in vertebrates. *J. Exp. Biol.* **213**, 4125–36 (2010).

9.   M. Stembridge, *et al.*, The overlooked significance of plasma volume for successful adaptation to high altitude in Sherpa and Andean natives. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 16177–16179 (2019).

10.   J. F. Storz, Genes for High Altitudes. *Science (80-.).* **329**, 40–41 (2010).

11.   S. L. Laguë, *et al.*, Divergent respiratory and cardiovascular responses to hypoxia in bar-

headed geese and Andean birds. *J. Exp. Biol.* **220**, 4186–4194 (2017).

12.   F. M. Faraci, Adaptations to hypoxia in birds: how to fly high. *Annu. Rev. Physiol.* **53**, 59–

70 (1991).

13.   G. R. Scott, W. K. Milsom, Flying high: A theoretical analysis of the factors limiting

exercise performance in birds at altitude. *Respir. Physiol. Neurobiol.* **154**, 284–301

(2006).

14.   C. M. Bishop, Heart mass and the maximum cardiac output of birds and mammals:

Implications for estimating the maximum aerobic power input of flying animals. *Philos.*

Trans. R. Soc. B Biol. Sci. **352**, 447–456 (1997).

15.   B. R. Grubb, Allometric relations of cardiovascular function in birds. *Am. J. Physiol.* **245**,

H567–H572 (1983).

16.   G. R. Scott, *et al.*, How Bar-Headed Geese Fly Over the Himalayas. *Physiology* **30**, 107–

115 (2015).

17.   M. C. W. Lim, C. C. Witt, C. H. Graham, L. M. Dávalos, Parallel Molecular Evolution in

Pathways, Genes, and Sites in High-Elevation Hummingbirds Revealed by Comparative

Transcriptomics. *Genome Biol. Evol.* **11**, 1552–1572 (2019).

18.   C. Natarajan, *et al.*, Predictable convergence in hemoglobin function has unpredictable

molecular underpinnings. *Science (80-.).* **354**, 336–339 (2016).

19.   J. Projecto-Garcia, *et al.*, Repeated elevational transitions in hemoglobin function during

the evolution of Andean hummingbirds. *Proc. Natl. Acad. Sci.* **110**, 20669–20674 (2013).

20.   T. W. Campbell, C. K. Ellis, *Avian and Exotic Animal Hematology and Cytology*

(Blackwell Publishing Inc., 2007).

21.   J. Samour, "Diagnostic Value of Hematology" in *Clinical Avian Medicine, Volume II*, G.

Harrison, T. Lightfoot, Eds. (2005), pp. 587–610.

22.   A. Borras, J. Cabrera, J. C. Senar, Hematocrit Variation in Response to Altitude Changes

in Wild Birds: A Repeated-Measures Design. *Condor* **112**, 622–626 (2010).

23.   D. M. Tufts, *et al.*, Phenotypic plasticity in blood-oxygen transport in highland and

lowland deer mice. *J. Exp. Biol.* **216**, 1167–1173 (2013).

24.   D. Levett, *et al.*, Acclimatization of skeletal muscle mitochondria to high-altitude hypoxia

during an ascent of Everest. *FASEB J.* **26**, 1431–1441 (2012).

25.   S. G. Dubay, C. C. Witt, Differential high-altitude adaptation and restricted gene flow

across a mid-elevation hybrid zone in Andean tit-tyrant flycatchers. *Mol. Ecol.* **23**, 3551–

3565 (2014).

26.   D. T. Clemens, Interspecific variation and effects of altitude on blood properties of rosy

finches (Leucosticte arctoa) and House Finches (Carpodacus mexicanus). *Physiol. Zool.*

62, 288–307 (1990).

27.   C. M. Beall, Two routes to functional adaptation: Tibetan and Andean high-altitude

natives. *Proc. Natl. Acad. Sci.* **104**, 8655–8660 (2007).

28.   K. Heinicke, *et al.*, Long-term exposure to intermittent hypoxia results in increased

hemoglobin mass, reduced plasma volume, and elevated erythropoietin plasma levels in

man. *Eur. J. Appl. Physiol.* **88**, 535–543 (2003).

29.   G. Ruiz, M. Rosenmann, A. Veloso, Altitudinal distribution and blood values in the toad,

Bufo spinulosus Wiegmann. *Comp. Biochem. Physiol. -- Part A Physiol.* **94**, 643–646

(1989).

692 30. H. Zhang, C. X. Wu, Y. Chamba, Y. Ling, Blood characteristics for high altitude

693   adaptation in Tibetan chickens. *Poult. Sci.* **86**, 1384–1389 (2007).

694 31. Y. Hao, *et al.*, Comparative transcriptomics of 3 high-altitude passerine birds and their

695   low-altitude relatives. *Proc. Natl. Acad. Sci.* **116**, 11851–11856 (2019).

696 32. N. J. Dawson, L. Alza, G. Nandal, G. R. Scott, K. G. McCracken, Convergent changes in

697   muscle metabolism depend on duration of high-altitude ancestry across andean waterfowl.

698   *Elife* **9**, 1–35 (2020).

699 33. B. J. Enquist, *et al.*, "Scaling from Traits to Ecosystems: Developing a General Trait

700   Driver Theory via Integrating Trait-Based and Metabolic Scaling Theories" in *Advances*

701   *in Ecological Research*, A. I. D. Samraat Pawar, Guy Woodward, Ed. (Elsevier Ltd.,

702   2015), pp. 249–318.

703 34. P. Minias, Ecology and evolution of blood oxygen-carrying capacity in birds. *Am. Nat.*

704   **195**, 788–801 (2020).

705 35. E. B. Linck, *et al.*, Blood variation implicates respiratory limits on elevational ranges of

706   Andean birds. *bioRxiv*, 1–28 (2021).

707 36. C. Carey, M. L. Morton, Aspects of circulatory physiology of montane and lowland birds.

708   *Comp. Biochem. Physiol. Part A* **54**, 61–74 (1976).

709 37. C. J. Axelrod, F. Laberge, B. W. Robinson, Intraspecific brain size variation between

710   coexisting sunfish ecotypes. *Proc. R. Soc. B Biol. Sci.* **285** (2018).

711 38. K. N. Yap, O. H.-I. Tsai, T. D. Williams, Haematological traits co-vary with migratory

712   status, altitude and energy expenditure: a phylogenetic, comparative analysis. *Sci. Rep.* **9**,

713   6351 (2019).

714 39. C. Natarajan, *et al.*, Convergent Evolution of Hemoglobin Function in High-Altitude

Andean Waterfowl Involves Limited Parallelism at the Molecular Sequence Level. *PLoS*

Genet. **11**, 1–25 (2015).

40.   J. F. Storz, H. Moriyama, Mechanisms of hemoglobin adaptation to high altitude hypoxia.

High Alt. Med. Biol. **9**, 148–157 (2008).

41.   X. Zhu, *et al.*, Elevational divergence in the great tit complex revealed by major

hemoglobin genes. *Curr. Zool.* **64**, 455–464 (2017).

42.   J. F. Storz, High-Altitude Adaptation: Mechanistic Insights from Integrated Genomics and

Physiology. *Mol. Biol. Evol.* (2021) https:/doi.org/10.1093/molbev/msab064.

43.   M. M. Wintrobe, Variation in the size and hemoglobin content of erythrocytes in the

blood of various vertebrates. *Folia Haematol.* **51**, 32–59 (1933).

44.   C. M. Hawkey, P. M. Bennett, S. C. Gascoyne, M. G. Hart, J. K. Kirkwood, Erythrocyte

size, number and haemoglobin content in vertebrates. *Br. J. Haematol.* **77**, 392–397

(1991).

45.   J. A. McGuire, *et al.*, Molecular phylogenetics and the diversification of hummingbirds.

Curr. Biol. **24**, 1–7 (2014).

46.   R. K. Suarez, Hummingbird flight: Sustaining the highest mass-specific metabolic rates

among vertebrates. *Experientia* **48**, 565–570 (1992).

47.   C. H. Graham, J. L. Parra, C. Rahbek, J. A. McGuire, Phylogenetic structure in tropical

hummingbird communities. *Proc. Natl. Acad. Sci.* **106**, 19673–19678 (2009).

48.   W. Buermann, *et al.*, Projected changes in elevational distribution and flight performance

of montane Neotropical hummingbirds in response to climate change. *Glob. Chang. Biol.*

17, 1671–1680 (2011).

49.   C. M. Beall, *et al.*, Hemoglobin concentration of high-altitude Tibetans and Bolivian

Aymara. *Am. J. Phys. Anthropol.* **106**, 385–400 (1998).

50.   C. M. Beall, Andean, Tibetan, and Ethiopian patterns of adaptation to high-altitude

hypoxia. *Integr. Comp. Biol.* **46**, 18–24 (2006).

51.   K. A. Hammond, J. O. E. Szewczak, E. Król, Effects of altitude and temperature on organ

phenotypic plasticity along an altitudinal gradient. *J. Exp. Biol.* **204**, 1991–2000 (2001).

52.   S. Lu, *et al.*, Differences in hematological traits between high- and low-altitude lizards

(Genus Phrynocephalus). *PLoS One* **10**, 1–16 (2015).

53.   N. F. Hadley, T. A. Burns, Intraspecific Comparison of the Blood Properties of the Side-

Blotched Lizard , Uta stansburiana. *Copeia* **1968**, 737–740 (1968).

54.   S. L. Fischer, S. P. Fischer, Mean Corpuscular Volume. *JAMA Intern. Med.* **143**, 282–283

(1983).

55.   M. S. Rose, Epitaph for the M.C.H.C. *Br. Med. J.* **4**, 169 (1971).

56.   A. Donkoh, Ambient temperature: a factor affecting performance and physiological

response of broiler chickens. *Int. J. Biometeorol.* **33**, 259–265 (1989).

57.   R. J. Moye, K. W. Washburn, T. M. Huston, Effects of environmental temperature on

erythrocyte numbers and size. *Poult. Sci.* **48**, 1683–1686 (1969).

58.   C. A. Glomski, A. Pica, *The Avian Erythrocyte: Its Phylogenetic Odyssey*, 1st Ed.

(Science Publishers, 2011).

59.   K. Yamaguchi, K. D. Jürgens, H. Bartels, J. Piiper, Oxygen transfer properties and

dimensions of red blood cells in high-altitude camelids, dromedary camel and goat. *J.*

Comp. Physiol. B Biochem. Syst. Environ. Physiol. **157**, 1–9 (1987).

60.   T. D. Brutsaert, M. Araoz, R. Soria, H. Spielvogel, J. D. Haas, Higher arterial oxygen

saturation during submaximal exercise in Bolivian Aymara compared to European

sojourners and Europeans born and raised at high altitude. *Am. J. Phys. Anthropol.* **113**,

169–181 (2000).

61. J. Grimminger, *et al.*, Thin Air Resulting in High Pressure: Mountain Sickness and

Hypoxia-Induced Pulmonary Hypertension. *Can. Respir. J.*, 1–17 (2017).

62. Y.-B. Guo, *et al.*, GCH1 plays a role in the high-altitude adaptation of Tibetans. *Zool. Res.*

38, 155–162 (2017).

63. P. Simmons, A. Lill, Development of parameters influencing blood oxygen carrying

capacity in the welcome swallow and fairy martin. *Comp. Biochem. Physiol. - A Mol.*

Integr. Physiol. **143**, 459–468 (2006).

64. W. S. Rasband, ImageJ.

65. N. Elarabany, A comparative study of some haematological and biochemical parameters

between two species from the Anatidae family within migration season. *J. Basic Appl.*

Zool. **79** (2018).

66. T. A. I. Parker, D. F. Stotz, J. W. Fitzpatrick, "Ecological and Distributional Databases" in

Neotropical Birds: Ecology and Conservation, D. K. M. D.F. Stotz, J.W. Fitzpatrick, T.A.

Parker, III, Ed. (University of Chicago Press, 1996).

67. A. Jaramillo, *Birds of Chile* (Princeton University Press, 2003).

68. T. S. Schulenberg, D. F. Stotz, D. F. Lane, J. P. O'Neill, T. A. P. III, *Birds of Peru*

(Princeton University Press, 2010).

69. R. J. Hijmans, S. E. Cameron, J. L. Parra, P. G. Jones, A. Jarvis, Very high resolution

interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**, 1965–1978 (2005).

70. P. de Villemereuil, S. Nakagawa, Modern phylogenetic comparative methods and their

application in evolutionary biology. *Mod. Phylogenetic Comp. Methods their Appl. Evol.*

Biol., 1–552 (2014).

71.   D. A. Skandalis, *et al.*, The biomechanical origin of extreme wing allometry in

hummingbirds. *Nat. Commun.* **8**, 1–9 (2017).

72.   R. Dudley, Mechanisms and Implications of Animal Flight Maneuverability. *Integr.*

Comp. Biol. **42**, 135–140 (2002).

73.   D. L. A. Roslyn Dakin, Paolo S. Segre, Andrew D. Straw, Morphology, muscle capacity,

skill, and maneuvering ability in hummingbirds. *Science (80-.).* **657**, 653–657 (2018).

74.   S. Y. Lee, G. R. Scott, W. K. Milsom, Have wing morphology or flight kinematics

evolved for extreme high altitude migration in the bar-headed goose? *Comp. Biochem.*

Physiol. Part C **148**, 324–331 (2008).

75.   D. L. Altshuler, R. Dudley, J. A. McGuire, Resolution of a paradox: Hummingbird flight

at high elevation does not come without a cost. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 17731–

17736 (2004).

76.   L. J. Revell, phytools: an R package for phylogenetic comparative biology (and other

things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

77.   F. Sornoza-Molina, J. F. Freile, J. Nilsson, N. Krabbe, E. Bonaccorso, A striking,

critically endangered, new species of hillstar (Trochilidae: Oreotrochilus) from the

southwestern Andes of Ecuador. *Auk* **135**, 1146–1171 (2018).

78.   D. Rubolini, A. Liker, L. Z. Garamszegi, A. P. Moller, N. Saino, Using the BirdTree. org

website to obtain robust phylogenies for avian comparative studies: A primer. *Curr. Zool.*

61, 959–965 (2015).

79.   W. Jetz, G. H. Thomas, J. B. Joy, K. Hartmann, A. O. Mooers, The global diversity of

birds in space and time. *Nature* **491**, 444–498 (2012).

80.   H. R. L. Lerner, M. Meyer, H. F. James, M. Hofreiter, R. C. Fleischer, Multilocus

808   resolution of phylogeny and timescale in the extant adaptive radiation of Hawaiian

809   honeycreepers. *Curr. Biol.* **21**, 1838–1844 (2011).

810   81.   P. C. Bürkner, brms: An R package for Bayesian multilevel models using Stan. *J. Stat.*

811   *Softw.* **80** (2017).

812   82.   J. L. Williamson, C. C. Witt, A lightweight backpack harness for tracking hummingbirds.

813   *J. Avian Biol.*, e02802 (2021).

814   83.   S. Watanabe, Asymptotic equivalence of Bayes cross validation and widely applicable

815   information criterion in singular learning theory. *J. Mach. Learn. Res.* **11**, 3571–3594

816   (2010).

817   84.   A. Vehtari, A. Gelman, J. Gabry, Practical Bayesian model evaluation using leave-one-out

818   cross-validation and WAIC. *Stat. Comput.*, 1413–1432 (2017).

Figure 1. Elevational, latitudinal, and phylogenetic sampling from 2006–2020. **(A)** Sampling localities spanned 28° latitude and 4,578 meters in elevation in Peru and Chile. **(B)** We studied 77 species ('n' column indicates sampling depth) from all nine major hummingbird clades. Sampling (gray opaque points) is shown across each species' elevational range (colored horizontal bars). Time-calibrated phylogeny is modified from McGuire et al. (45). Illustrations depicting a representative from each clade are from *Birds of the World* and are reproduced courtesy of the Cornell Lab of Ornithology (©).

Figure 2. Relationship between elevation and the six studied blood parameters (hemoglobin concentration ([Hb]), hematocrit (Hct), total red blood cell count (TRBC), mean cell volume (MCV), mean cell hemoglobin content (MCH), and mean cell hemoglobin concentration (MCHC). **(A–F)** Within species comparisons, wherein each point represents a value from a single individual. **(G–L)** Among species comparisons, wherein each point represents a mean value from a single species. In all panels, solid trend lines correspond to relationships where 95% posterior credible intervals in models did not overlap zero, while dashed trend lines correspond to relationships where 89% credible intervals in models did not overlap zero (see Figure 3). Credible intervals for both elevation (at the 95% level) and elevation^2 (at the 89% level) did not overlap zero in the TRBC model (Panel I). Colors correspond to clades in Figure 1. R^2 values are reported from Bayesian models reported in Figure 3.

Figure 3. Posterior probabilities and credible intervals from full within-species (main plots) and among-species (insets) models for each of six blood traits: **(A)** hemoglobin concentration ([Hb]), **(B)** hematocrit (Hct), **(C)** total red blood cell (TRBC), **(D)** mean cell volume (MCV), **(E)** mean cell hemoglobin (MCH), **(F)** mean cell hemoglobin concentration (MCHC). Circles indicate beta estimates, thick and thin bars illustrate 89% and 95% credible intervals, respectively. Red circles denote parameters that did not overlap zero in top-ranked models. For Hb, Hct, and TRBC within-species analyses, the full model was the top model. See Tables S3-S4 for model comparison and parameter estimates from top models in other model sets.

Figure 4. Theoretical and empirical relationships among hummingbird blood parameters underlying hemoglobin concentration ([Hb]) adjustment and predictors of the Cell Number-Size Index (*CeNS*). **(A–B)** Hummingbirds have evolved optimized blood-O_2 carrying capacity, achieved by balancing total red blood cell count (TRBC) with mean cell volume (MCV). **(A)** Predictions for O_2 carrying-capacity. Background is shaded by O_2 carrying-capacity, calculated as *MCV x TRBC x 1/3 scaling factor* to represent constant MCHC. **(B)** Empirical findings. Points are colored by [Hb], with lighter colors representing higher concentrations. **(C)** Mean cell hemoglobin (MCH) and MCV are tightly linked, suggesting little variance in mean cell

hemoglobin concentration (MCHC). Points are colored by TRBC (units presented on the y-axis),

with lighter colors representing higher erythrocyte counts. In panels A–C, each point represents a

value from a single individual. **(C)** Posterior probabilities and credible intervals from the best

fitting reduced *CeNS* model. **(D)** *CeNS* shows a strong quadratic relationship with elevation,

revealing that middle elevation species adjust blood-O_2 carrying capacity differently than low-

and high-elevation species. **(E)** *CeNS* increases with increasing body mass. In panels D–E, R^2

values are reported from generalized linear models of each *x* regressed on *y*, and thus do not

account for other modeled variables.