Oxygen availability drives blood traits and the cell number-size tradeoff across Andean

hummingbirds

Jessie L. Williamson[1,4*], Ethan B. Linck[1], Emil Bautista[2], Ashley Smiley[1,3], Jimmy A. McGuire[3],

Robert Dudley[3], and Christopher C. Witt[1*]

[1]Museum of Southwestern Biology and Department of Biology, University of New Mexico, 302

Yale Blvd NE, Albuquerque, NM 87131, USA

[2]Centro de Ornitología y Biodiversidad (CORBIDI), Lima, Perú

[3]Museum of Vertebrate Zoology and Department of Integrative Biology, University of

California, Berkeley, Berkeley, CA 94720, USA

[4]Lead contact

*Corresponding authors: Jessie L. Williamson, williamson@unm.edu, @jl_williamson;

Christopher C. Witt, cwitt@unm.edu, @MSBbirds). Address: Department of Biology, University

of New Mexico, Castetter Hall 1480, MSC03-2020, 219 Yale Blvd. NE, Albuquerque, NM

87131, USA

SUMMARY

For endotherms with sustained high metabolism, like hummingbirds, blood-oxygen (O_2) carrying

capacity should be finely tuned to supply O_2 to respiring tissues. Hematological adjustments are

expected in response to changes in O_2 availability that occur along elevational gradients;

however, the rules by which this variation occurs have remained elusive. For example,

elevational variation among conspecific individuals may not be comparable to variation among

species that have evolved differences in traits, genes, and environmental niches. We sought to

elucidate these eco-physiological rules in the hummingbird radiation by analyzing blood traits

that affect O_2-carrying functions: total hemoglobin concentration ([Hb]), hematocrit (Hct),

erythrocyte count (TRBC), mean cell volume (MCV), mean cellular hemoglobin content (MCH),

and mean cellular hemoglobin concentration (MCHC). For 1,217 Andean hummingbirds

representing 77 species and spanning sea level to 4,578 meters in elevation, we modeled blood

traits as responses to environmental conditions and individual and species traits. [Hb] and Hct

increased with elevation similarly within species and across the phylogeny—for every 1,000-m

ascent, [Hb] increased by one third of a gram per deciliter and Hct increased by one and a half

percentage points, corresponding to about three-hundred thousand additional cells per microliter.

Hummingbirds adjusted blood cell number and cell size by similar proportions to modulate

blood-O_2 carrying capacity; however, species at the lowest and highest elevations relied

relatively more on adjustments to cell volume, revealing qualitatively different evolutionary

responses elicited by moderate versus extreme reductions in O_2 availability.

Key words: Andes, blood, blood-oxygen carrying capacity, cell, elevation, hummingbird,

physiology, elevational gradient

INTRODUCTION

Organisms that sustain high levels of aerobic activity under hypoxia, such as birds flying at high altitude, must supply oxygen (O_2) to respiring tissues despite reduced O_2 availability in 'thin' air. To achieve this, the O_2 transport functions of blood must be finely tuned to match O_2 supply and demand under a variety of ambient conditions[1–6]. Most O_2 in the blood is carried by hemoglobin (Hb)[7], the concentration of which ([Hb]) is a key determinant of blood-O_2 carrying capacity. Under hypobaric hypoxia, animals experience reduced arterial O_2 saturation; to compensate, they undergo erythropoiesis, which can increase [Hb] and therefore blood-O_2 carrying capacity, enhancing O_2 delivery to the tissues and compensating for reduced arterial O_2 saturation[8]. In the absence of compensatory increases in plasma volume[9], increased Hb mass leads to elevated hematocrit (Hct) and, beyond a certain threshold, corresponding increases in blood viscosity will begin to compromise O_2 transport[10]. A suite of interrelated physical characteristics of blood— total red blood cell count (TRBC), mean cell volume (MCV), mean cell hemoglobin (MCH), and mean cell hemoglobin concentration (MCHC; Table S1)—can be modulated to optimize blood-O_2 transport, or to compensate for other cardiac, vascular, or hematological responses to elevation.

Birds are well-known for their abilities to maintain tissue oxygenation in hypoxic environments, which can be attributed to both plastic and constitutive aspects of respiratory physiology that enhance O_2 uptake, transport, and delivery[3,11–19]. Along with many factors that have been shown to affect the physical characteristics of avian blood, including age, sex, body condition, hormones, environmental factors, and disease[20,21], high-altitude hypoxia has predictable short-term effects on blood composition that are mediated by the hypoxia-inducible factor (HIF) pathway[8,22–24]. Accordingly, when lowland-adapted birds ascend to altitude, HIF-

induced production of erythropoietin is expected to increase [Hb] and Hct. Birds that are

chronically exposed to high elevations have been found to display either normal or elevated

[Hb], Hct, and MCHC[25,26]. Evidence from high-altitude-adapted human populations and

theoretical work suggests that optimal [Hb] and Hct levels may be similar to those of sea-level

populations[8,13,27]. MCV typically increases in acclimatized, lowland-adapted animals undergoing

accelerated erythropoiesis[28]; however, bird and other vertebrate lineages that have evolved for

extended periods at altitude have often been found to have smaller cells, likely as part of a suite

of adaptive traits for efficient tissue oxygenation[23,29,30]. Plasticity in blood characteristics to low

partial pressure of O_2 over short-term timescales of acclimatization is well-documented[8,23]. Over

longer timescales, natural selection on genes that affect hypoxia performance leads predictably to

genetic adaptation[17–19,31], but the effects of this evolutionary process on the physical composition

of blood are unclear.

Systematic trait variation across environmental gradients reflects the outcome of

evolutionary and developmental processes, and explaining such 'trait gradients' will be essential

to predicting the functional consequence of climate warming for species and communities[32].

Despite its critical importance for tissue oxygenation, many questions remain unanswered

regarding elevational variation in blood traits that underpin blood-O_2 carrying capacity. Blood

traits vary among avian clades (e.g.,[20,33]) and exhibit elevational variation both within species

(e.g.,[4,25,34]) and among species (e.g.,[4,25,33,35]). Trait shifts with elevation are expected to differ

systematically for conspecific versus heterospecific comparisons; whereas within-species

variation should be driven primarily by developmental plasticity, among-species variation should

be influenced by genetic changes that have accrued during differences in elevational exposure

lasting thousands to millions of generations[36]. Linck et al.[34] estimated elevational effects on [Hb]

and Hct within each of 136 species of Andean birds and showed that, within the median species,

[Hb] increased by 0.7 g/dL per 1,000 m in elevation, while Hct increased by 2.0% per 1,000 m.

At least two additional recent studies have conducted broad-scale interspecific comparative

analyses of avian [Hb] and Hct data, but both datasets were compiled from heterogeneous

published data and lacked elevational sampling sufficient to estimate elevational effects[33,37].

Furthermore, no previous comparative analysis of hematological variation across elevation has

included the six key parameters that affect blood-O_2 carrying capacity per unit volume (Table

S1). We continue to lack a general understanding of within- and among-species effects of

elevation on blood traits, and specifically, an understanding of whether different rules govern

elevational variation at these different levels of biological organization.

97   Another outstanding question is the degree to which known genetic adaptations to

hematological function may predict blood-O_2 carrying capacity. Adaptive genetic changes to

hemoglobin proteins have been demonstrated in many high-altitude-native birds, including

hummingbirds, House Wrens, Andean waterfowl, Bar-headed Geese, and a few dozen other bird

taxa[17,19,38–41]. In hummingbirds, there is a highly predictable association between species-typical

Hb β^A_{13}–β^A_{83} genotype and elevation, with derived increases in Hb-O_2 affinity in highland

lineages and reductions in Hb-O_2 affinity in lowland lineages[17]. Hummingbirds occurring at high

altitudes exhibit two amino-acid substitutions on the β^A-globin subunit of Hb (Hb β^A-13$_{GLY \to SER}$

and Hb β^A-83 $_{GLY \to SER}$) that are known to confer increased O_2-binding affinity[17]. In particular,

β^A-13$_{SER}$ is known from only four hummingbird genera (*Patagona*, *Chalcostigma*, *Pterophanes*,

and *Oreotrochilus*), all of which occur at extreme high altitudes[3,19,42]. Both O_2-binding affinity

and blood-O_2 carrying capacity have critical effects on tissue oxygenation, but are part of a

complex O_2 uptake, transport, and delivery system that contains multiple convective and

diffusive steps[43]. It is plausible that species-typical Hb β^A genotypes associated with different

O_2-binding affinities predict hematological phenotypes that relate to blood-O_2 carrying functions;

such an association could reflect a direct functional tradeoff or an indirect effect of accrued

genetic adaptation to elevation at multiple loci.

The constraints on blood trait values that are imposed by physical tradeoffs and the

relative contributions of each blood trait to overall blood-O_2 carrying capacity remain poorly

understood. [Hb] defines O_2-carrying capacity per unit volume of blood, and it can be adjusted

via adjustments to cell number (TRBC), cell size (MCV), within-cell [Hb] (MCHC), or plasma

volume. MCHC tends to be relatively invariant across the vertebrate tree of life, as demonstrated

nearly a century ago by Wintrobe[44], with Hb typically comprising ~1/3 of erythrocytes. Plasma

volume adjustments can decouple Hb mass from [Hb] and, although plasma volume appears to

vary among human populations[9], we currently have insufficient data on elevational variation in

avian plasma volume to assess its importance. Cell number and cell size are both measurable

components that are variable among individuals and species, subject to plastic and evolutionary

adjustments, and have strong direct effects on [Hb]. Cell number and cell size are reciprocally

constrained by physical space available in the circulatory system, such that individual birds can

have a few large erythrocytes, many small erythrocytes, or an intermediate state for both

traits[44,45]. Quantifying the relative contributions of TRBC and MCV in driving [Hb] variation

within and among species may reveal that species utilize different mechanisms of adjusting

blood-O_2 carrying capacity; if such mechanisms are predictably associated with elevational

environments or evolutionary histories of altitude exposure, they could lead to new

biogeographic insights.

In this study, we sought to elucidate the rules governing variation in blood traits that

affect O_2-carrying functions, including the effects of environment, individual characteristics, and

evolved species differences. Specifically, we asked: (1) To what extent do blood traits shift

predictably with elevational changes in O_2 availability, while controlling for other sources of

variation?; (2) Do blood traits vary with O_2 availability differently over developmental (within

species) and evolutionary (among species) time scales?; (3) How are blood traits affected by

phylogenetic history and evolved species differences, such as body mass, metabolic intensity,

and known genetic variants that affect Hb-O_2 binding affinity?; (4) To what extent are individual

and species variation in O_2-carrying capacity per unit volume of blood attributable to changes in

cell number versus cell size? And (5) does manifestation of the cell number-size tradeoff vary

predictably among species?

We collected specimen-vouchered hematological data from hummingbirds between

2006–2020 across a ~4,600-meter elevational gradient in the Andes. Data for six blood traits

from 1,217 individuals of 77 species were used to model within- and among-species variation,

respectively (Figure 1). Hummingbirds are a useful system in which to study hematological

variation due to their documented evolutionary genetic responses to shifts in PO$_2$, high species

richness in mountains, phylogenetically conserved elevational ranges, and relative uniformity of

body size, clutch size, diet, mating system, and flight mode[19,42,46–48]. Notably, matching O_2

supply and demand is an acute challenge for all hummingbirds because they exhibit the highest

mass-specific metabolic rates among endotherms, and because they employ sustained hovering,

one of the most energetically demanding forms of animal locomotion[47,49].

RESULTS AND DISCUSSION

We quantified three primary ([Hb], Hct, and TRBC) and three secondary hematological indices

(MCV, MCH, and MCHC; Table S1), and separately analyzed two datasets: an individual-level

dataset, wherein each row contained data from one individual hummingbird (n=1,217), and a

species-level dataset, wherein each row contained mean blood values calculated for each species,

with a minimum of 2 individuals sampled (n=58–65 species, depending on blood trait modeled).

In the individual-level data, mean [Hb] was 18.84 g/dl (±1.65), mean Hct was 58.54% (±5.15),

mean TRBC was 6.37 RBC x 10^6/mm^3 or 6,370,000 cells per µl (±1.16), mean MCV was 93.36

fl (±17.1), mean MCH was 30.4 pg (±5.64), and mean MCHC was 32.44 g/dl (± 2.12). Complete

reference values—the first such values ever published for >90% of our study species—are

reported in detail in Table S2.

Within-species blood comparison

We used the individual-level dataset to model within species variation and accounted for

between-species differences by incorporating a species grouping variable. Using multilevel

Bayesian models for each of the six blood parameters, we estimated the effects of elevation,

body mass, precipitation, air temperature, species-typical Hb β^A_{13}–β^A_{83} genotype, as well as

within-species variation in elevational position (an individual's sampled elevation relative to its

species elevational range), body mass, air temperature, and precipitation. Separately, we

excluded sex and wing-loading as potentially important predictors (see Methods). In total, we

compared six models per set using LOOIC and WAIC (Tables 1 and S5; Methods). In addition to

fixed effects, models included species as a random effect, and a phylogenetically correlated

random intercept (i.e., covariance matrix of phylogenetic distances among taxa). Thus, our full

model in each set reflects the hypothesis that all predictors, phylogeny, and species identity

influence blood characteristics.

Based on the top-performing model in each set, we found that for every 1,000-m increase

in elevation, [Hb] is expected to increase by 0.39 g/dl, Hct is expected to increase by 1.42%,

TRBC is expected to increase by 310,000 cells per μl, and MCH is expected to decrease by 0.48

pg. Elevation strongly predicted [Hb], Hct, TRBC, and MCH (Figures 2 A–F and 4A–F).

Elevational position was also a strong predictor of Hct, indicating that individuals located higher

in their species' elevational range tended to have higher Hct (Figure 4B). [Hb], TRBC, and MCH

also increased with increasing elevational position, although these relationships were subtle

(Figure 4A, C, E).

The finding that [Hb], Hct, and TRBC increase with elevation was consistent with

theoretical expectations for O_2-carrying capacity of blood and with prior empirical findings in

birds[3,4,25,33,37], humans[50,51], mammals[23,52], and other taxa[53,54]. In prior work, substantial variation

in slopes of these parameters across elevation may have been attributable to variation in

temporal, taxonomic, and spatial scales of comparison, as well as constraints on the time-course

of evolutionary adaptation[17,31,51]. We expect increases in Hct at higher elevations because an

increase in the total mass of Hb enhances oxygenation under lower ambient PO_2[13]; unless there

is a proportional increase in plasma volume or MCHC, an increase in Hb mass is expected to be

accompanied by an increase in the proportion of blood comprised of red blood cells. TRBC also

increased with elevation and was, along with MCV, one of two key drivers of [Hb] adjustment

(Figure 2, Figure 3, Figure 5C–D)[45]. MCV did not vary predictably in our linear models (Figure

2D). Although increasing within-cell [Hb], or MCHC, would seem to be an alternative

mechanism of increasing O_2-carrying capacity, MCHC was less variable than other blood

parameters (Table S2), likely due to physical constraints[55,56]; accordingly, MCH and MCV were

strongly correlated (Table S3; Figure 3C). Contrary to recent empirical findings[4,25], adjustments

to MCHC appeared to be neither substantial nor predictable with respect to elevation within or

among species (Figure 2F).

We found that MCV and MCH decreased in individuals occurring in warmer

environments (Figures 4D, E), and that MCHC decreased in individuals occurring at warmer

localities within a species' range (Figure 4F). The directions of these relationships were

consistent with previous experimental work: Comparisons of juvenile broiler chickens reared

from ~1-7 weeks of age in cool, ambient, and hot environments showed decreases in [Hb], Hct,

MCV, and TRBC as temperature increased[57,58]. Our findings add to a limited but intriguing pool

of evidence that warmer ambient temperatures may lead to developmental reductions in O_2-

carrying capacity of blood.

Models that included the species grouping variable fit substantially better than those

without; this effect explained 13–21% of the variance in the top model, depending on the model

set (Tables 1 and S5; Methods). Moderate effects of the species grouping variable suggest that

variation in blood traits is partly explained by aspects of species biology that evolve too fast to

have resulted in phylogenetic autocorrelation. Species-typical Hb β^A_{13}–β^A_{83} genotype did not

predict [Hb] or any other blood characteristics (Figure 4A–F), providing no evidence that

evolutionary adjustments to O_2-binding affinity affect the physical composition of blood or its

variation with elevation.

Among-species blood comparison

Using the species-level dataset, we evaluated the effects of elevation, body mass, species-typical

Hb β^A_{13}–β^A_{83} genotype, and ambient temperature and precipitation on each of our 6 blood

response variables. We compared sets of models with LOOIC and WAIC (Table 1),

incorporating a quadratic elevation effect for TRBC, MCV, MCH, and MCHC models based on

curvature evident in the data (Figure 2G–L; Methods).

Within- and among-species findings for elevational effects were similar (Figure 4). Based

on the top model in each set, for every 1,000-m increase in elevation, [Hb] is expected to

increase by 0.29 g/dl, and Hct is expected to increase by 1.55%. Elevation was a consistently

strong predictor of [Hb], Hct, and TRBC, all of which increased with increasing elevation

(Figures 2G–I and 4A–C). The importance of the quadratic effect of elevation for MCV and

MCH, and overall shape of the species-level data, suggest that O_2 carrying capacity adjustment

differs for middle-elevation species versus those found at low and high elevations (Figure 2J–K;

Figure 4D–E). Neither species-typical Hb $\beta^A 13$–$\beta^A 83$ genotype, body mass, environmental

temperature, nor precipitation explained variation in any species-mean blood characteristic

(Figure 4).

Influence of cell number versus size on blood-O_2 carrying capacity

The axis of variation in cell number versus size space was tightly constrained (Figure 3A–B); the

'banana' shape of Figure 3 mirrors patterns from Wintrobe[44] and Hawkey et al.[45], which present

similar plots based on species representing multiple vertebrate classes. The shape of this

relationship indicates that the highest levels of blood-O_2 carrying capacity per unit blood are

achieved at high values of TRBC or MCV, respectively. Individual hummingbirds can possess

few large erythrocytes or many small erythrocytes, but cell number and size must be balanced[45].

Because of the proportionately greater surface area for gas exchange, having many small cells is

predicted to be advantageous for fast metabolism or when the partial pressure of O_2 in inspired

air is low[45]. Similarly, having large cells may be advantageous for modulating the time-course of

gas exchange[59]. One way to shed light on this tradeoff is to test whether individuals and species,

respectively, adjust O_2 carrying capacity per unit blood primarily through increases in cell

number or size.

We designed models to understand the relative contributions of cell number versus size to

blood O_2-carrying capacity per unit volume ([Hb]), within and among species over ~22 Mya of

hummingbird evolution. These models demonstrated that hummingbirds co-equally modulated

cell number and cell size to adjust [Hb] (Table S6). The fact that this was true both within and

among species suggests that developmental and evolutionary fine-tuning of blood-O_2 carrying

capacity occur similarly.

Using data for 32 well-sampled species, we estimated species-specific standardized

TRBC and MCV coefficients (β) in models with [Hb] as the response variable (Figure S2), and

we used these to calculate a 0–1 index that we refer to as the Cell Number-Size Index (hereafter

CeNS). *CeNS* represents the proportional contribution of TRBC versus MCV to increasing [Hb],

calculated as:

262
$$\frac{(\beta_{TRBC})}{(\beta_{TRBC} + \beta_{MCV})}$$

Species specific *CeNS* values ranged from 0.36-0.69 ($\bar{x} = 0.52$), with higher values indicating a

proportionally greater contribution of TRBC than MCV in adjusting [Hb]; conversely, lower

values indicate a proportionally greater contribution of MCV than TRBC.

Extending our analysis, we designed a set of models to test potential predictors of *CeNS*:

elevation, a quadratic elevation effect (elevation^2; to account for notable curvature in the data),

body mass, wing loading, species-typical Hb β^A_{13}–β^A_{83} genotype, and phylogeny (see Methods).

Elevation, elevation^2, and body mass strongly predicted *CeNS* (Figure 5C-D; Tables S9–S10).

We found a trend of decreasing *CeNS* with higher wing loading (Figure 5F), but the 95%

credible intervals overlapped zero (Figure 5C). Species-typical Hb β^A_{13}–β^A_{83} genotype did not

predict *CeNS* (Figure S5), nor did we find phylogenetic signal in *CeNS* (Figure S4).

Examining the tradeoff between cell number and volume provides a potential explanation

for the striking pattern that middle elevation species tend to adjust [Hb] by different mechanisms

than low and high elevation species (Figure 5A–B). *CeNS* was uncorrelated with TRBC

coefficients ($R^2 = 0.07$, $r = -0.27$) but strongly correlated with MCV coefficients ($R^2 = 0.51$, $r = -$

0.71), indicating that variation in *CeNS* was primarily driven by the degree to which species

modulate cell size (Figure 5, Figure S3). We suggest that because low and high elevation species

tend to possess relatively small cells, they may be more able to rely on shifts in cell size, rather

than cell number, to increase [Hb] (Figure 2J; Figure 5). However, as cell size increased, species

were more likely to approach maximum size for red blood cells, beyond which they could incur a

functional detriment. As a result, species with larger red blood cells were more reliant on cell

number to adjust [Hb]. For middle elevation hummingbird species that have evolved under

moderate (~15–30%) reductions in O_2 availability, TRBC appears to be the primary axis for

adjustment of blood-O_2 carrying capacity per unit volume of blood. Species that have evolved at

the highest elevations are expected to be the best adapted to hypobaric hypoxia, and it is

therefore remarkable that their mode of [Hb] adjustment (primarily by MCV) resembles that of

hummingbird species in the lowlands, the putative ancestral elevation for hummingbirds[46].

Hummingbirds at extreme elevations

Do hummingbirds at extreme high elevations follow the same rules as hummingbirds

distributed at low and moderate elevations with respect to elevational blood variation? Our

models encompassed the entire elevational gradient and therefore could have missed important

phenomena such as threshold effects. To address this, we compared individual hummingbirds

sampled above and below 4,200 m elevation, a threshold that marks the upper ~10% of the

gradient. Hummingbirds sampled from these extreme high elevations had significantly smaller

cells and higher erythrocyte counts than others (Figure 6). This finding suggests that high-

altitude specialist hummingbirds tend to have smaller cells, consistent with findings in other high

altitude vertebrates[23,29,30,53,60]; however, this pattern was only detectable under the extreme

conditions at the uppermost limits of the Andean elevational gradient. We proceeded to formally

test for elevational breakpoints using segmented regression (see Methods). Evidence for

objective blood trait thresholds along the elevational gradient was mixed, with breakpoint

estimates for individual traits ranging from ~1,400 to ~4,000 m (Figure S6). Lack of consilience

in these estimates suggests that trait-specific thresholds are challenging to identify and may not

be generalizable across species and clades.

Phylogenetic signal in blood characteristics

Phylogenetic signal was low ($\lambda = 0.0$; K range = 0.14–0.21) across all blood traits for both

within- and among-species models. Blomberg's K and Pagel's λ results were consistent with

visual assessment of continuous trait maps, even after removing species with <3 individuals

312   sampled (Table S4; Figure S1). We additionally found no phylogenetic signal in *CeNS* (Figure

313   S4).

314   Non-existent to low phylogenetic signal in blood traits within and among hummingbird

species was surprising given demonstrated, phylogenetically conserved Hb genetic adaptations to

elevation[17,19] and in light of strong phylogenetic signal reported from recent comparative

analyses of blood traits that had broader taxonomic sampling[33,37]. However, it is likely that

diverse avian clades possess different blood trait optima; in previous comparative analyses that

included multiple families or orders, phylogenetic signal may have been driven primarily by

differences among clades. For example, Minias et al.[33] reported that small passerines had

consistently high [Hb] and Hct estimates, and that including the shorebird order,

Charadriiformes, which was characterized by high [Hb] but medium Hct values, strongly

influenced patterns of phylogenetic conservatism[33]. We tested for and found no interaction of

clade and elevation on the six studied blood traits, with the exception of the interaction of

elevation and the Hermit clade in the [Hb] model set (Figure S7). Interestingly, hermits are a

large, early diverging hummingbird clade that contains diminishing species diversity with

altitude (and no species that exceeds 3,000 m), and its apparent increased sensitivity of [Hb] to

elevation may hint at a functional explanation for this biogeographic tendency. The existence of

clade-specific trait optima[33] and strong species effects, demonstrated in this study, may at first

seem difficult to reconcile with our findings that there was no within-family phylogenetic signal

in hummingbirds; however, we view these findings as being consistent with steady evolution of

trait optima by a mean-reverting, Ornstein-Uhlenbeck process, as described by Minias et al.[33],

and marked by occasional major jumps that coincide with changes in body plan or respiratory

demands.

A strength of our study was that hummingbirds vary only subtly in aspects of their

biology that could plausibly confound our analyses, such as metabolic rate, energy budget, torpor

use, mode of locomotion, foraging strategy, or clutch size. Phylogenetic conservatism in these

traits facilitates comparisons of blood characteristics across the entire clade to an extent that

would be difficult in other taxa. A strong species effect in spite of these intrinsic controls hints

that species optimize O_2-carrying functions of blood for subtle changes in lifestyle and

environment, on rapid timescales. This finding is consistent with the rapid diversification of

human hematological responses to elevation, as illustrated by the copious variation among

populations in patterns of elevational variation[27,50,51,61–63].

Conclusions

Across ~22 Mya of the hummingbird evolution, ambient O_2-availability and species

idiosyncrasies—not phylogenetic affinities—were the primary drivers of variation in blood traits

affecting O_2-carrying functions. In response to elevational changes in O_2 partial pressure,

hummingbirds make predictable adjustments to the physical composition of blood. Remarkably,

after controlling for many potentially confounding variables, decreasing PO_2 drove increases in

[Hb] and Hct that were nearly identical in magnitude within and among hummingbird species,

respectively. This implies that developmental and evolutionary optimization of blood traits to

ambient PO_2 occurs in fundamentally similar ways, responding to the same set of physical rules

that govern gas exchange at the blood-gas barrier. In addition to O_2 availability, species identity

itself explained 13–21% of blood trait variation, suggesting that fast-evolving, unmodeled

aspects of species biology underpin substantial hematological variation. Our data revealed a

striking constraint space that defines the cell number-size tradeoff for hummingbird erythrocytes;

within this space, individuals and species co-equally modulated cell number and size to adjust

[Hb]. However, species varied in their proportional reliance on adjustments to cell number

versus size (*CeNS*) to increase [Hb], and both elevation and body mass were strongly predictive

of species' tendencies. Middle-elevation species that have been evolving under moderate

hypoxia adjusted [Hb] primarily by adjusting cell number; in contrast, low and high elevation

species tended to adjust [Hb] by adjusting cell size, a pattern partly underpinned by

hummingbirds at extreme elevations exhibiting smaller cells. Together, these results suggest that

evolutionary solutions to reduced O_2 availability can differ qualitatively depending on species-

specific histories of exposure to hypoxia challenges of varying severity.

ACKNOWLEDGEMENTS

We thank Luis Alza, Christopher Barger, Selina Bauernfeind, Matthew Baumann, Elizabeth

Beckman, Phred Benham, José Miguel Bogdanovich, Francisco Bozinovic, Homan Castillo

Benitez, Jessica Castillo, Marlon Chagua, Jennifer Clark, Mariela Combe, Lida Crooks, Robert

Driver, Shane DuBay, Margarita Espinoza, L. Monica Flores, Chauncey Gadek, Ariel Gaffney,

Spencer Galen, Avia González-Méndez, Paige Handley, Zachary Hanna, Mike Hartshorne, José

Ernesto Huaroto Tornero, Andrew Johnson, Matthew Jones, Daniel Lane, LyAndra Lujan,

Mattías Marzfeld, Sabrina McNew, Kyana Montoya, Jano Nuñez-Zapata, Iris Olivas, Paloma

Ordoñez, Alessandra Quiñonez, Javier Reinoso, Natalia Ricote, C. Gregory Schmitt, Donna

Schmitt, C. Jonathan Schmitt, Frank Solano Bravo, Dora Susanibar, William Talbot, Jorge

Tiravanti C., Abraham Urbay T., Thomas Valqui, Walter Vargas Campos, Karen Verde-Guerra,

Natalie Wright, and Alfredo Zelada. We are grateful to the following communities in Peru for

site access: Comunidades de San Pedro de Casta, Campesina Santiago de Carampoma, Ocros,

Macate, Oncoy, Yanahara, Huacarpay, Sianbal, Plataforma, Agua Azul, Incahuasi, and

Anchicha. This research was funded by the National Science Foundation (DEB-0543556, DEB-

1146491, and DBI-1907353), and dissertation research grants to JLW from the Nuttall

Ornithological Club Blake-Nuttall Fund Grants (2016 & 2018), the American Philosophical

Society Lewis & Clark Grant, the Explorers Club Exploration Fund Grant, American Museum of

Natural History Chapman grants (2017 & 2019), the UNM Biology Graduate Student

Association Graduate Research Allocations Committee grant, UNM Graduate and Professional

Student Association grants (2017–2021), the UNM Latin American & Iberian Institute Field

Research Grant, UNM Department of Biology Grove Research Scholarships (2018, 2019, and

2021), and the UNM Department of Biology Melinda Bealmear Memorial Scholarship and Dr.

AUTHOR CONTRIBUTIONS

Conceptualization: JLW, EBL, JAM, RD, CCW; Formal analysis: JLW; Funding and resource

acquisition: JLW, JAM, RD, CCW; Investigation: JLW, EB, AS, JAM, RD, CCW;

Methodology: JLW, EBL, EB, CCW; Writing – original draft: JLW, CCW. All authors reviewed

and edited the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1. Krause, J.S., Németh, Z., Pérez, J.H., Chmura, H.E., Ramenofsky, M., and Wingfield, J.C. (2015). Annual Hematocrit Profiles in Two Subspecies of White-Crowned Sparrow: A Migrant and a Resident Comparison. Physiol. Biochem. Zool. *89*, 51–60.

2. Storz, J.F., and Scott, G.R. (2019). Life Ascending: Mechanism and Process in Physiological Adaptation to High-Altitude Hypoxia. Annu. Rev. Ecol. Evol. Syst. *50*, 1–24.

3. Williamson, J.L., and Witt, C.C. (2021). Elevational niche-shift migration: Why the degree of elevational change matters for the ecology, evolution, and physiology of migratory birds. Ornithology *138*, 1–26.

4. Barve, S., Dhondt, A.A., Mathur, V.B., and Cheviron, Z.A. (2016). Life-history characteristics influence physiological strategies to cope with hypoxia in Himalayan birds. Proc. R. Soc. B *283*, 1–8.

5. Laguë, S.L. (2017). High-altitude champions: birds that live and migrate at altitude. J. Appl. Physiol. *123*, 942–950.

6. Storz, J.F. (2016). Hemoglobin–oxygen affinity in high-altitude vertebrates: is there evidence for an adaptive trend? J. Exp. Biol. *219*, 3190–3203.

7. Pittman, R.N. (2011). Chapter 4: Oxygen Transport. In Regulation of Tissue Oxygentation (Morgan & Claypool Life Sciences).

8. Storz, J.F., Scott, G.R., and Cheviron, Z.A. (2010). Phenotypic plasticity and genetic adaptation to high-altitude hypoxia in vertebrates. J. Exp. Biol. *213*, 4125–36.

9. Stembridge, M., Williams, A.M., Gasho, C., Dawkins, T.G., Drane, A., Villafuerte, F.C., Levine, B.D., Shave, R., and Ainslie, P.N. (2019). The overlooked significance of plasma

volume for successful adaptation to high altitude in Sherpa and Andean natives. Proc.

Natl. Acad. Sci. U. S. A. *116*, 16177–16179.

10.   Storz, J.F. (2010). Genes for High Altitudes. Science (80-.). *329*, 40–41.

11.   Laguë, S.L., Chua, B., Alza, L., Scott, G.R., Frappell, P.B., Zhong, Y., Farrell, A.P.,

McCracken, K.G., Wang, Y., and Milsom, W.K. (2017). Divergent respiratory and

cardiovascular responses to hypoxia in bar-headed geese and Andean birds. J. Exp. Biol.

220, 4186–4194.

12.   Faraci, F.M. (1991). Adaptations to hypoxia in birds: how to fly high. Annu. Rev. Physiol.

53, 59–70.

13.   Scott, G.R., and Milsom, W.K. (2006). Flying high: A theoretical analysis of the factors

limiting exercise performance in birds at altitude. Respir. Physiol. Neurobiol. *154*, 284–

301.

14.   Bishop, C.M. (1997). Heart mass and the maximum cardiac output of birds and mammals:

Implications for estimating the maximum aerobic power input of flying animals. Philos.

Trans. R. Soc. B Biol. Sci. *352*, 447–456.

15.   Grubb, B.R. (1983). Allometric relations of cardiovascular function in birds. Am. J.

Physiol. *245*, H567–H572.

16.   Scott, G.R., Hawkes, L.A., Frappell, P.B., Butler, P.J., Bishop, C.M., and Milsom, W.K.

(2015). How Bar-Headed Geese Fly Over the Himalayas. Physiology *30*, 107–115.

17.   Projecto-Garcia, J., Natarajan, C., Moriyama, H., Weber, R.E., Fago, A., Cheviron, Z.A.,

Dudley, R., McGuire, J.A., Witt, C.C., and Storz, J.F. (2013). Repeated elevational

transitions in hemoglobin function during the evolution of Andean hummingbirds. Proc.

Natl. Acad. Sci. *110*, 1–6.

18. Lim, M.C.W., Witt, C.C., Graham, C.H., and Dávalos, L.M. (2019). Parallel Molecular

Evolution in Pathways, Genes, and Sites in High-Elevation Hummingbirds Revealed by

Comparative Transcriptomics. Genome Biol. Evol. *11*, 1552–1572.

19. Natarajan, C., Hoffmann, F.G., Weber, R.E., Fago, A., Witt, C.C., and Storz, J.F. (2016).

Predictable convergence in hemoglobin function has unpredictable molecular

underpinnings. Science (80-.). *354*, 336–339.

20. Campbell, T.W., and Ellis, C.K. (2007). Avian and Exotic Animal Hematology and

Cytology (Blackwell Publishing Inc.).

21. Samour, J. (2005). Diagnostic Value of Hematology. In Clinical Avian Medicine, Volume

II, G. Harrison and T. Lightfoot, eds., pp. 587–610.

22. Borras, A., Cabrera, J., and Senar, J.C. (2010). Hematocrit Variation in Response to

Altitude Changes in Wild Birds: A Repeated-Measures Design. Condor *112*, 622–626.

23. Tufts, D.M., Revsbech, I.G., Cheviron, Z.A., Weber, R.E., Fago, A., and Storz, J.F.

(2013). Phenotypic plasticity in blood-oxygen transport in highland and lowland deer

mice. J. Exp. Biol. *216*, 1167–1173.

24. Levett, D., Radford, E., Menassa, D., Graber, E., Morash, A., Hoppeler, H., Clarke, K.,

Martin, D., Ferguson-Smith, A., Montgomery, H., et al. (2012). Acclimatization of

skeletal muscle mitochondria to high-altitude hypoxia during an ascent of Everest. FASEB

J. *26*, 1431–1441.

25. Dubay, S.G., and Witt, C.C. (2014). Differential high-altitude adaptation and restricted

gene flow across a mid-elevation hybrid zone in Andean tit-tyrant flycatchers. Mol. Ecol.

23, 3551–3565.

26. Clemens, D.T. (1990). Interspecific variation and effects of altitude on blood properties of

rosy finches (Leucosticte arctoa) and House Finches (Carpodacus mexicanus). Physiol.

Zool. *62*, 288–307.

27. Beall, C.M. (2007). Two routes to functional adaptation: Tibetan and Andean high-

altitude natives. Proc. Natl. Acad. Sci. *104*, 8655–8660.

28. Heinicke, K., Prommer, N., Cajigal, J., Viola, T., Behn, C., and Schmidt, W. (2003).

Long-term exposure to intermittent hypoxia results in increased hemoglobin mass,

reduced plasma volume, and elevated erythropoietin plasma levels in man. Eur. J. Appl.

Physiol. *88*, 535–543.

29. Ruiz, G., Rosenmann, M., and Veloso, A. (1989). Altitudinal distribution and blood

values in the toad, Bufo spinulosus Wiegmann. Comp. Biochem. Physiol. -- Part A

Physiol. *94*, 643–646.

30. Zhang, H., Wu, C.X., Chamba, Y., and Ling, Y. (2007). Blood characteristics for high

altitude adaptation in Tibetan chickens. Poult. Sci. *86*, 1384–1389.

31. Dawson, N.J., Alza, L., Nandal, G., Scott, G.R., and McCracken, K.G. (2020). Convergent

changes in muscle metabolism depend on duration of high-altitude ancestry across andean

waterfowl. Elife *9*, 1–35.

32. Enquist, B.J., Norberg, J., Bonser, S.P., Violle, C., Webb, C.T., Henderson, A., Sloat,

L.L., and Savage, V.M. (2015). Scaling from Traits to Ecosystems: Developing a General

Trait Driver Theory via Integrating Trait-Based and Metabolic Scaling Theories. In

Advances in Ecological Research, A. I. D. Samraat Pawar, Guy Woodward, ed. (Elsevier

Ltd.), pp. 249–318.

33. Minias, P. (2020). Ecology and evolution of blood oxygen-carrying capacity in birds. Am.

Nat. *195*, 788–801.

34.   Linck, E.B., Williamson, J.L., Bautista, E., Beckman, E.J., Benham, P.M., DuBay, S.G.,

499   Flores, L.M., Gadek, C.R., Johnson, A.B., Jones, M.R., et al. (2021). Blood variation

500   implicates respiratory limits on elevational ranges of Andean birds. bioRxiv, 1–28.

501   35.   Carey, C., and Morton, M.L. (1976). Aspects of circulatory physiology of montane and

502   lowland birds. Comp. Biochem. Physiol. Part A *54*, 61–74.

503   36.   Axelrod, C.J., Laberge, F., and Robinson, B.W. (2018). Intraspecific brain size variation

504   between coexisting sunfish ecotypes. Proc. R. Soc. B Biol. Sci. *285*.

505   37.   Yap, K.N., Tsai, O.H.-I., and Williams, T.D. (2019). Haematological traits co-vary with

506   migratory status, altitude and energy expenditure: a phylogenetic, comparative analysis.

507   Sci. Rep. *9*, 6351.

508   38.   Natarajan, C., Projecto-Garcia, J., Moriyama, H., Weber, R.E., Muñoz-Fuentes, V., Green,

509   A.J., Kopuchian, C., Tubaro, P.L., Alza, L., Bulgarella, M., et al. (2015). Convergent

510   Evolution of Hemoglobin Function in High-Altitude Andean Waterfowl Involves Limited

511   Parallelism at the Molecular Sequence Level. PLoS Genet. *11*, 1–25.

512   39.   Galen, S.C., Natarajan, C., Moriyama, H., Weber, R.E., Fago, A., Benham, P.M., Chavez,

513   A.N., Cheviron, Z.A., Storz, J.F., and Witt, C.C. (2015). Contribution of a mutational hot

514   spot to hemoglobin adaptation in high-altitude Andean house wrens. Proc. Natl. Acad. Sci.

515   *112*, 13958–13963.

516   40.   Storz, J.F., and Moriyama, H. (2008). Mechanisms of hemoglobin adaptation to high

517   altitude hypoxia. High Alt. Med. Biol. *9*, 148–157.

518   41.   Zhu, X., Guan, Y., Qu, Y., David, G., Song, G., and Lei, F. (2017). Elevational divergence

519   in the great tit complex revealed by major hemoglobin genes. Curr. Zool. *64*, 455–464.

42.   Projecto-Garcia, J., Natarajan, C., Moriyama, H., Weber, R.E., Fago, A., Cheviron, Z.A.,

Dudley, R., McGuire, J.A., Witt, C.C., and Storz, J.F. (2013). Repeated elevational

transitions in hemoglobin function during the evolution of Andean hummingbirds. Proc.

Natl. Acad. Sci. *110*, 20669–20674.

43.   Storz, J.F. (2021). High-Altitude Adaptation: Mechanistic Insights from Integrated

Genomics and Physiology. Mol. Biol. Evol.

44.   Wintrobe, M.M. (1933). Variation in the size and hemoglobin content of erythrocytes in

the blood of various vertebrates. Folia Haematol. *51*, 32–59.

45.   Hawkey, C.M., Bennett, P.M., Gascoyne, S.C., Hart, M.G., and Kirkwood, J.K. (1991).

Erythrocyte size, number and haemoglobin content in vertebrates. Br. J. Haematol. *77*,

392–397.

46.   McGuire, J.A., Witt, C.C., Remsen, J. V, Corl, A., Rabosky, D.L., Altshuler, D.L., and

Dudley, R. (2014). Molecular phylogenetics and the diversification of hummingbirds.

Curr. Biol. *24*, 1–7.

47.   Suarez, R.K. (1992). Hummingbird flight: Sustaining the highest mass-specific metabolic

rates among vertebrates. Experientia *48*, 565–570.

48.   Graham, C.H., Parra, J.L., Rahbek, C., and McGuire, J.A. (2009). Phylogenetic structure

in tropical hummingbird communities. Proc. Natl. Acad. Sci. *106*, 19673–19678.

49.   Buermann, W., Chaves, J.A., Dudley, R., Mcguire, J.A., Smith, T.B., and Altshuler, D.L.

(2011). Projected changes in elevational distribution and flight performance of montane

Neotropical hummingbirds in response to climate change. Glob. Chang. Biol. *17*, 1671–

1680.

50.   Beall, C.M., Brittenham, G.M., Strohl, K.P., Blangero, J., Williams-Blangero, S.,

Goldstein, M.C., Decker, M.J., Vargas, E., Villena, M., Soria, R., et al. (1998).

544  Hemoglobin concentration of high-altitude Tibetans and Bolivian Aymara. Am. J. Phys.

545  Anthropol. *106*, 385–400.

546 51. Beall, C.M. (2006). Andean, Tibetan, and Ethiopian patterns of adaptation to high-altitude

547  hypoxia. Integr. Comp. Biol. *46*, 18–24.

548 52. Hammond, K.A., Szewczak, J.O.E., and Król, E. (2001). Effects of altitude and

549  temperature on organ phenotypic plasticity along an altitudinal gradient. J. Exp. Biol. *204*,

550  1991–2000.

551 53. Lu, S., Xin, Y., Tang, X., Yue, F., Wang, H., Bai, Y., Niu, Y., and Chen, Q. (2015).

552  Differences in hematological traits between high- and low-altitude lizards (Genus

553  Phrynocephalus). PLoS One *10*, 1–16.

554 54. Hadley, N.F., and Burns, T.A. (1968). Intraspecific Comparison of the Blood Properties of

555  the Side-Blotched Lizard , Uta stansburiana. Copeia *1968*, 737–740.

556 55. Fischer, S.L., and Fischer, S.P. (1983). Mean Corpuscular Volume. JAMA Intern. Med.

557  *143*, 282–283.

558 56. Rose, M.S. (1971). Epitaph for the M.C.H.C. Br. Med. J. *4*, 169.

559 57. Donkoh, A. (1989). Ambient temperature: a factor affecting performance and

560  physiological response of broiler chickens. Int. J. Biometeorol. *33*, 259–265.

561 58. Moye, R.J., Washburn, K.W., and Huston, T.M. (1969). Effects of environmental

562  temperature on erythrocyte numbers and size. Poult. Sci. *48*, 1683–1686.

563 59. Glomski, C.A., and Pica, A. (2011). The Avian Erythrocyte: Its Phylogenetic Odyssey 1st

564  ed. (Science Publishers).

565 60. Yamaguchi, K., Jürgens, K.D., Bartels, H., and Piiper, J. (1987). Oxygen transfer

566  properties and dimensions of red blood cells in high-altitude camelids, dromedary camel

and goat. J. Comp. Physiol. B Biochem. Syst. Environ. Physiol. *157*, 1–9.

61. Brutsaert, T.D., Araoz, M., Soria, R., Spielvogel, H., and Haas, J.D. (2000). Higher

arterial oxygen saturation during submaximal exercise in Bolivian Aymara compared to

European sojourners and Europeans born and raised at high altitude. Am. J. Phys.

Anthropol. *113*, 169–181.

62. Grimminger, J., Richter, M., Tello, K., Sommer, N., Gall, H., and Ghofrani, H.A. (2017).

Thin Air Resulting in High Pressure: Mountain Sickness and Hypoxia-Induced Pulmonary

Hypertension. Can. Respir. J., 1–17.

63. Guo, Y.-B., He, Y.-X., Cui, C.-Y., Ouzhuluobu, Baimakangzhuo, Duojizhuoma,

Dejiquzong, Bianba, Peng, Y., Bai, C., et al. (2017). GCH1 plays a role in the high-

altitude adaptation of Tibetans. Zool. Res. *38*, 155–162.

64. Simmons, P., and Lill, A. (2006). Development of parameters influencing blood oxygen

carrying capacity in the welcome swallow and fairy martin. Comp. Biochem. Physiol. - A

Mol. Integr. Physiol. *143*, 459–468.

65. Rasband, W.S. ImageJ.

66. Elarabany, N. (2018). A comparative study of some haematological and biochemical

parameters between two species from the Anatidae family within migration season. J.

Basic Appl. Zool. *79*.

67. Parker, T.A.I., Stotz, D.F., and Fitzpatrick, J.W. (1996). Ecological and Distributional

Databases. In Neotropical Birds: Ecology and Conservation, D. K. M. D.F. Stotz, J.W.

Fitzpatrick, T.A. Parker, III, ed. (University of Chicago Press).

68. Jaramillo, A. (2003). Birds of Chile (Princeton University Press).

69. Schulenberg, T.S., Stotz, D.F., Lane, D.F., O'Neill, J.P., and III, T.A.P. (2010). Birds of

Peru (Princeton University Press).

70.   Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G., and Jarvis, A. (2005). Very high

resolution interpolated climate surfaces for global land areas. Int. J. Climatol. *25*, 1965–

1978.

71.   Villemereuil, P. de, and Nakagawa, S. (2014). Modern phylogenetic comparative methods

and their application in evolutionary biology. Mod. Phylogenetic Comp. Methods their

Appl. Evol. Biol., 1–552.

72.   Skandalis, D.A., Segre, P.S., Bahlman, J.W., Groom, D.J.E., Welch, K.C., Witt, C.C.,

McGuire, J.A., Dudley, R., Lentink, D., and Altshuler, D.L. (2017). The biomechanical

origin of extreme wing allometry in hummingbirds. Nat. Commun. *8*, 1–9.

73.   Dudley, R. (2002). Mechanisms and Implications of Animal Flight Maneuverability.

Integr. Comp. Biol. *42*, 135–140.

74.   Roslyn Dakin, Paolo S. Segre, Andrew D. Straw, D.L.A. (2018). Morphology, muscle

capacity, skill, and maneuvering ability in hummingbirds. Science (80-.). *657*, 653–657.

75.   Lee, S.Y., Scott, G.R., and Milsom, W.K. (2008). Have wing morphology or flight

kinematics evolved for extreme high altitude migration in the bar-headed goose? Comp.

Biochem. Physiol. Part C *148*, 324–331.

76.   Altshuler, D.L., Dudley, R., and McGuire, J.A. (2004). Resolution of a paradox:

Hummingbird flight at high elevation does not come without a cost. Proc. Natl. Acad. Sci.

U. S. A. *101*, 17731–17736.

77.   Revell, L.J. (2012). phytools: an R package for phylogenetic comparative biology (and

other things). Methods Ecol. Evol. *3*, 217–223.

78.   Sornoza-Molina, F., Freile, J.F., Nilsson, J., Krabbe, N., and Bonaccorso, E. (2018). A

striking, critically endangered, new species of hillstar (Trochilidae: Oreotrochilus) from

the southwestern Andes of Ecuador. Auk *135*, 1146–1171.

79.   Rubolini, D., Liker, A., Garamszegi, L.Z., Moller, A.P., and Saino, N. (2015). Using the

BirdTree. org website to obtain robust phylogenies for avian comparative studies: A

primer. Curr. Zool. *61*, 959–965.

80.   Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K., and Mooers, A.O. (2012). The global

diversity of birds in space and time. Nature *491*, 444–498.

81.   Lerner, H.R.L., Meyer, M., James, H.F., Hofreiter, M., and Fleischer, R.C. (2011).

Multilocus resolution of phylogeny and timescale in the extant adaptive radiation of

Hawaiian honeycreepers. Curr. Biol. *21*, 1838–1844.

82.   Bürkner, P.C. (2017). brms: An R package for Bayesian multilevel models using Stan. J.

Stat. Softw. *80*.

83.   Vehtari, A., Gelman, A., and Gabry, J. (2017). Practical Bayesian model evaluation using

leave-one-out cross-validation and WAIC. Stat. Comput., 1413–1432.

84.   Williamson, J.L., and Witt, C.C. (2021). A lightweight backpack harness for tracking

hummingbirds. J. Avian Biol., e02802.

85.   Watanabe, S. (2010). Asymptotic equivalence of Bayes cross validation and widely

applicable information criterion in singular learning theory. J. Mach. Learn. Res. *11*,

3571–3594.

Table 1. Comparison of Bayesian models to evaluate within- and among-species variation in the 6 studied blood traits: hemoglobin

concentration ([Hb]), hematocrit (Hct), total red blood cell count (TRBC), mean cell volume (MCV), mean cell hemoglobin (MCH),

and mean cell hemoglobin concentration (MCHC). Only within species models included the species grouping variable and only

among species TRBC, MCV, MCH, and MCHC models included a quadratic component (elev^2). Reduced models included the set of

predictors from full models whose 95% credible intervals did not overlap zero. Dashes indicate that a reduced model was not present

(i.e., all credible intervals from full models overlapped zero). Model fit was assessed using the widely applicable information criterion

(WAIC) and leave-one-out cross-validation (LOO), which were consistent. The difference between the best fit model and each other

model in the set is shown as ΔELPD with difference in standard error (ΔSE) in parentheses (LOOIC values are calculated as –

2*ELPD, which converts ELPD to a deviance scale). ΔELPD < 4 indicates that differences between two models are small.

	Model description						Response					
	Predictors					**Grouping Variable**	**[Hb] ΔELPD (ΔSE)**	**Hct ΔELPD (ΔSE)**	**TRBC ΔELPD (ΔSE)**	**MCV ΔELPD (ΔSE)**	**MCH ΔELPD (ΔSE)**	**MCHC ΔELPD (ΔSE)**
	None (null)	**All (full)**	**Reduced**	**All + elev^2**	**Reduced + elev^2**	**Species**						
Within species	X			NA	NA	X	-29 (8.4)	-22.6 (8.2)	-3.9 (4.7)	-0.5 (1.0)	**0.0 (1.5)**	-11.5 (5.0)
		X		NA	NA	X	**0.0**	**0.0**	**0.0**	-0.9 (3.6)	-1.2 (3.1)	-0.1 (2.6)
			X	NA	NA	X	-3.3 (4.2)	-0.3 (2.8)	-0.1 (3.8)	**0.0**	**0.0**	**0.0**
	X			NA	NA		-85.5 (13.1)	-70.5 (13.6)	-34.6 (9.3)	-24.9 (6.4)	-19.6 (6.5)	-39.4 (8.7)
		X		NA	NA		-29.9 (7.6)	-29.9 (7.7)	-29.8 (7.7)	-26.5 (7.3)	-22.8 (6.7)	-29.3 (7.1)
			X	NA	NA		-29.7 (8.8)	-26.9 (7.9)	-25.0 (8.2)	-23.3 (7.1)	-20.1 (6.5)	-29.9 7.1
Among species	X					NA	-4.5 (3.9)	-2.3 (4.1)	**0.0**	**0.0**	-1.3 (2.2)	**0.0**
		X				NA	-0.1 (3.9)	**0.0**	-2.7 (2.6)	-3.6 (2.5)	-4.5 (3.5)	-5.4 (1.8)
			X			NA	**0.0**	0.1 (2.0)	-0.5 (0.8)	–	–	–
				X		NA	NA	NA	-2.0 (2.5)	-2.4 (2.7)	-2.8 (2.1)	-4.5 (2.1)
					X	NA	NA	NA	–	–	**0.0**	–

Figure 1. Elevational, latitudinal, and phylogenetic sampling from 2006–2020. **(A)** Sampling localities spanned 28° latitude and 4,578 meters in elevation in Peru and Chile. **(B)** We studied 77 species ('n' column indicates sampling depth) from all nine major hummingbird clades. Sampling (gray opaque points) is shown across each species' elevational range (colored horizontal bars). Time-calibrated phylogeny is modified from McGuire et al.[46]. Illustrations depicting a representative from each clade are from *Birds of the World* and are reproduced courtesy of the Cornell Lab of Ornithology (©).

Figure 2. Relationship between elevation and the six studied blood parameters (hemoglobin concentration ([Hb]), hematocrit (Hct), total red blood cell count (TRBC), mean cell volume (MCV), mean cell hemoglobin content (MCH), and mean cell hemoglobin concentration (MCHC). **(A–F)** Within species comparisons, wherein each point represents a value from a single individual. **(G–L)** Among species comparisons, wherein each point represents a mean value from a single species. In all panels, solid trend lines correspond to relationships where 95% posterior credible intervals in models did not overlap zero, while dashed trend lines correspond to relationships where 89% credible intervals in models did not overlap zero (see Figure 4). Credible intervals for both elevation (at the 95% level) and elevation^2 (at the 89% level) did not overlap zero in the TRBC model (Panel I). Colors correspond to clades in Figure 1. R^2 values are reported from Bayesian models reported in Figure 4.

Figure 3. Theoretical and empirical relationships among hummingbird blood parameters

underlying hemoglobin concentration ([Hb]) adjustment. **(A–B)** Hummingbirds have evolved

optimized blood-O_2 carrying capacity, achieved by balancing total red blood cell count (TRBC)

with mean cell volume (MCV). **(A)** Predictions for O_2 carrying-capacity. Background is shaded

by O_2 carrying-capacity, calculated as *MCV x TRBC x 1/3 scaling factor* to represent constant

MCHC. **(B)** Empirical findings. Points are colored by [Hb], with lighter colors representing

higher concentrations. **(C)** Mean cell hemoglobin (MCH) and MCV are tightly linked,

suggesting little variance in mean cell hemoglobin concentration (MCHC). Points are colored by

TRBC (units presented on the y-axis), with lighter colors representing higher erythrocyte counts.

In all panels, each point represents a value from a single individual.

Figure 4. Posterior probabilities and credible intervals from full within-species (main plots) and among-species (insets) models for each of six blood traits: (A) hemoglobin concentration ([Hb]), (B) hematocrit (Hct), (C) total red blood cell (TRBC), (D) mean cell volume (MCV), (E) mean cell hemoglobin (MCH), (F) mean cell hemoglobin concentration (MCHC). Circles indicate beta estimates, thick and thin bars illustrate 89% and 95% credible intervals, respectively. Red circles denote parameters that did not overlap zero in top-ranked models. For Hb, Hct, and TRBC within-species analyses, the full model was the top model. See Tables 1 and S5 for model comparison and parameter estimates from top models in other model sets.

Figure 5. (A–B) Effects of mean cell volume (MCV) and total red blood cell count (TRBC) on *CeNS*, elucidating how, once cells reach a certain size, TRBC becomes the primary mechanism through which birds adjust [Hb]. **(C–F)** Results from *CeNS* models. **(C)** Posterior probabilities and credible intervals from the best fitting full model (main plot) and reduced model (inset) for *CeNS*. **(D)** *CeNS* shows a strong quadratic relationship with elevation, revealing that middle elevation species adjust blood-O_2 carrying capacity differently than low- and high-elevation species. **(E)** *CeNS* increases with increasing body mass. **(F)** *CeNS* shows a marginal decrease with increased wing loading. R^2 values are reported from generalized linear models of each *x* regressed on *y*, and thus do not account for other modeled variables.

Figure 6. Blood values differ significantly at 'extreme' high elevations. **(A)** Total red blood cell count (TRBC) is significantly higher ≥4,200 m ($p < 0.001$). TRBC increases by 530,000 cells per µl with each 100-meter increase in elevation ≥4,200 m. **(B)** Mean cell volume (MCV) is significantly lower ≥4,200 m (p=0.010). MCV decreases by 5.52 fl with each 100-meter increase in elevation ≥4,200 m. **(C)** Mean cell hemoglobin content (MCH) is significantly lower ≥4,200 m (p=0.020). MCH decreases by 2.37 pg with each 100-meter increase in elevation ≥4,200 m. Data were analysed by subsetting the individual-level dataset by sampling elevation ("<4,200 m" or "≥4,200 m"), consistent with notable trends in the data (Figure 2). Significance and direction of relationships were assessed with one-way ANOVAs. Rates of increase and decrease were calculated using regression coefficients generalized linear models.

METHODS

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and code should be directed to and will be fulfilled by the Lead Contacts, Jessie Williamson (williamson@unm.edu) and Christopher Witt (cwitt@unm.edu).

Materials availability
This study did not generate new unique reagents.

Data and code availability
This paper analyzes original data collected from hummingbirds in Peru and Chile. All data are linked to vouchered specimens at the Museum of Southwestern Biology (MSB), Centro de Ornitología y Biodiversidad (CORBIDI), and the Universidad Pontificia Católica de Chile, accessible in the Arctos database (www.arctosdb.org). Detailed data on specimens, collection localities, and blood values are reported in a forthcoming data paper, Witt et al. (*In Prep*). Additional information about key resources is provided in the Key Resources Table. All code and extensive annotations for data processing, analyses, and figures are available on GitHub: https://github.com/jlwilliamson/ComparativeHummingbirdBlood. Any additional information required to reanalyze data reported in this paper is available from the lead contacts upon request.

METHOD DETAILS

Field Sampling
We collected data from 1,217 individuals of 77 hummingbird species from Peru and Chile during 2006–2020 (Figure 1). After filtering (see below), our final within-species dataset included 1,151 individuals of 77 species collected from 75 localities (Figure 1). The best-sampled clades were the Coquettes (387 individuals, 20 species) and Brilliants (269 individuals, 19 species). The species with greatest sampling depth were *Metallura tyrianthina* (95 individuals), *Colibri coruscans* (73 individuals), and *Amazilia amazilia* and *Patagona gigas* (64 individuals each; Figure 1). Sampling across sexes was balanced (553 females, 592 males, and 6 unknown). The majority of birds sampled were adults (936 adults, 179 juveniles, and 36 unknown). Detailed data on specimens, collection localities, and blood values are described in Witt et al. (*In prep*) and are available in the ARCTOS database (arctosdb.org) for reproducibility and extension of our work.

Hematological measurements
To calculate primary indices (hemoglobin concentration ([Hb]), hematocrit (Hct), and total red blood cell count (TRBC)), we obtained whole blood samples within ~5 mins to ~3 hours of capture by venipuncture on the underside of the wing and collection with heparinized microcapillary tubes (Hct) and Hemocue HB201+ cuvettes ([Hb]). Hct (%) was measured with digital calipers after centrifuging the sealed microcapillary tube for 5 min at 13,000 r.p.m. to separate the red blood cells and plasma. When >1 Hct sample was taken, values were averaged,

and weighted Hct values were used in analyses. [Hb] (g/dL blood) was measured on ~5 µL of
blood using a HemoCue HB201+ hemoglobin photometer. The HemoCue proprietary
photometric method produces values for avian blood that are ~1.0 g/dL greater than those
generated using cyanomethemoglobin spectrophotometry[64]. To account for this, we corrected
[Hb] values by subtracting 1.0 g/dL. When possible, at time of blood collection, 10 µl whole
blood was preserved in a 1:200 NaCl dilution to obtain TRBC estimates. Within 12-hours of
blood collection, we pipetted a 10 µl sub-sample of blood-NaCl dilution into a hemocytometer
and allowed cells to settle for 1 min. After focusing the microscope, one or more photos were
obtained of cells in the hemocytometer grid. Later, cells were manually counted in the ImageJ
software[65] and TRBC was estimated following standard protocols[20,21].
After drawing blood, birds were humanely killed by rapid cardiac compression. We
recorded collection elevation, latitude, longitude, sex, body mass, and date. Body mass was
typically taken prior to blood sampling; however, when individuals were massed after blood
sampling, body mass was corrected for estimated volume of blood drawn. Each collected bird
was prepared as a study skin. Specimens and tissues are housed at the Museum of Southwestern
Biology (MSB) at the University of New Mexico, USA, the Centro de Ornitología y
Biodiversidad (CORBIDI) in Peru, and the Universidad Pontificia Católica de Chile in Chile.
Secondary blood indices (mean cell volume (MCV), mean cell hemoglobin (MCH), and
mean cell hemoglobin concentration (MCHC)) were calculated from primary indices (Table S1)
following standard protocols[20]. We examined blood data distributions using histograms and
qqPlots and eliminated outliers based on distribution, in combination with known 'typical'
values[20,25,66]. To reduce the possibility of incorporating anomalous blood values resulting from
juvenile birds (i.e., age-related outliers), we examined values on the tail ends of distributions
associated with presence of a bursa measuring ~2 x 2 mm. All values fell within the expected
distribution of blood values. We did not age individuals by plumage, as juvenile plumage lasts
much longer than juvenile blood values, which stabilize one to several months after fledging.
802
Environmental and life history variables
We compiled elevational range limits for each species in our dataset using a combination of
published data, field guides, and specimen records from the Museum of Southwestern Biology
(MSB)[3,67–69]. We calculated each individual's elevational range position, hereafter 'elevational
position', as *1 – ((maximum species elevation – individual sampling elevation) / species*
elevational range). This 0.0–1.0 index allowed us to quantify each individual's position relative
to its species elevational range breadth. Generally, we expect that individuals sampled near the
higher end of their species' elevational range (i.e., that have higher elevational position values)
will have higher blood trait values for most parameters.
We incorporated species-typical Hb β^A_{13}–β^A_{83} genotypes using published sequence
values[42,46]. For *n*=7 species, we identified species-typical Hb β^A_{13}–β^A_{83} genotype from Hb β^A
sequences aligned to a GenBank reference from Projecto-Garcia et al.[42]. For *n*=12 species
(*Amazilia franciae, Campylopterus villaviscensio, Eutoxeres aquila, Klais guimeti, Leucippus*
taczanowskii, Myrmia micrura, Phaethornis atrimentalis, Sephanoides sephaniodes,
Chaetocercus mulsant, Eriocnemis vestita, Myrtis fanny, and *Oreotrochilus leucopleurus*)
species-typical Hb β^A_{13}–β^A_{83} genotypes were inferred from sister taxa or nearest relatives.
We characterized climatic variation among latitudinal and elevational gradients using 19
bioclimatic variables from the WorldClim database[70]. We downloaded rasters of each BioClim
variable using the 'raster' package in R (version 4.0.0), cropped them to our study areas of Peru

and Chile, and extracted temperature and precipitation values for each sampling locality. We then used principal components analysis (PCA) for temperature (Bio1–11) and precipitation variables (Bio12–19) to create a composite measure of temperature and precipitation across the gradient. PC1 of the temperature variables (hereafter "temperature") explained 73.4% of the variation in composite temperature, and loadings suggested that this axis corresponded to increasing temperature across sites (see GitHub). PC1 of the precipitation variables (hereafter "precipitation") explained 79.8% of the variation in composite precipitation, and loadings suggested that this axis corresponded to increasing precipitation across sites (see GitHub).

        To account for multiple measurement effects (i.e., traits measured from multiple individuals within species), we calculated between- and within-species differences following Villemereuil et al.[71]. This approach uses within-group centering to separate each predictor into two components: We first calculated species mean values (between-species variability) and then subtracted the mean value of individual observations (within-species differences). The latter measures of within-species variation for temperature, precipitation, and body mass were included in models. Because hummingbird body masses may differ by sex, within-species differences were calculated using mean body mass of males for males, mean body mass of females for females, and species-level means for birds of unknown sex (n=6).

        We used wing area data from the 59 species from the Museum of Southwestern Biology studied in Skandalis et al.[72] to calculate wing loading in g/cm^2, calculated as species mean body mass (g) divided by total wing area (cm^2). We analyzed wing loading due its relevance to aerodynamic theory and hummingbird flight at high altitudes[73–76].

Hummingbird phylogeny

We used the substitution rate-calibrated molecular phylogeny of hummingbirds from McGuire et al.[46] as a backbone to generate a phylogeny pruned to the 77 species in our dataset using the R package 'phytools'[77]. Two species, *Oreotrochilus leucopleurus* and *Phaethornis stuarti*, are missing from the McGuire et al.[46] tree and were added to our phylogeny using the bind.tip() function. Branch lengths and position of *O. leucopleurus* were calculated from genetic distances and recent described sister relationships[78]. Approximate branch length and position of *P. stuarti* was inferred from BirdTree.org[79,80] with divergence rate estimates from a multi-locus, time-calibrated phylogeny of Hawaiian honeyceepers[81]. We generated other model-specific tree subsets by including only those tips that corresponded to the species in each modeling subset (see annotated GitHub analysis scripts).

Phylogenetic signal

Phylogenetic signal indicates the proportion of the total variance attributed to phylogenetic variance. We originally ran sets of phylogenetic models (results not shown) and calculated phylogenetic signal, or lambda (λ) from the models as the phylogenetic heritability described by Lynch (1991). We estimated phylogenetic signal using the full and reduced models for each blood parameter following the vignette and recommendations of P. Bürkner (https://cran.r-project.org/web/packages/brms/ vignettes/brms_phylogenetics.html), using the 'hypothesis' method. However, all estimates of phylogenetic signal for all model sets (full and reduced for within- and among species analyses) were low ($\lambda < 0.004$); these estimates are supported by visualizations of continuous trait maps of mean blood parameters (Figure S1). Because estimates from preliminary models and continuous trait maps of mean blood values—across all datasets— indicated low phylogenetic signal (Supplement; Figure S1), we excluded phylogeny as an effect

868 in models and evaluated phylogenetic signal prior to modeling using Blomberg's K and Pagel's
869 λ. Results among the two tests were similar and consistent across model subsets (e.g., Table S4).
870

Within species analysis
We built generalized linear multilevel models using the R package brms[82] to analyze within
species variation in blood characteristics. Brms uses Markov chain Monte Carlo (MCMC)
sampling to obtain draws from posterior distributions and it easily interfaces with the R package
loo[83] to compute different information criteria. We designed hierarchical models for each of six
continuous responses variables: [Hb], Hct, TRBC, MCV, MCH, and MCHC. For each model,
predictor variables included one species-level categorical variable (species-typical Hb β^A_{13}–β^A_{83}
genotype) and eight individual-level continuous variables (elevation, elevational position, body
mass (log-transformed), precipitation, temperature, and within species variation in body mass,
temperature, and precipitation). All variables were standardized to a mean of zero and standard
deviation of one.
For each response, we compared 6 models: (1) an intercept-only null model, (2) an
intercept-only model with a species grouping variable (1|species), (3) a model with all predictors
and no grouping variable, (4) a model with all predictors and the species grouping variable, (5) a
reduced-predictor model with no grouping variable (reduced model 3), (6) a reduced-predictor
model with the species grouping variable (reduced model 4). Reduced models included only
predictors whose 95% credible intervals (CIs) did not overlap zero in full models, suggesting
high importance. Distribution family differed by model set: Gaussian for [Hb], Hct, and TRBC
models; skew-normal for MCV and MCH models; and student's *t*-distribution for MCHC
models. We ran four Markov chains for 10,000 iterations with a burn-in period of 5,000, thinned
every 10 steps.
The full predictor models with species grouping variable had the lowest (best) WAIC and
ELPD values for [Hb] (R^2 = 0.21), Hct (R^2 = 0.17), and TRBC (R^2 = 0.17) models. For MCV,
the reduced model with temperature and the species grouping variable had the lowest scores (R^2
= 0.094) and for MCHC, the reduced model that included within species variation in both body
mass and temperature plus the species grouping variable ranked the highest (R^2 = 0.12). Both top
MCV and MCHC models ranked similarly to the full model with species grouping variable, as
well as the null model. For MCH, the reduced model with elevational position, elevation, and the
species grouping variable ranked the highest and was comparable to the null model (R^2 = 0.092;
Table 1).
We tested for interactions between clade and elevation by comparing three models: (1) The
top within-species model for each blood parameter; (2) The top model + (1|Clade); and (3) The
top model with an elev:Clade interaction. We evaluated models and conducted model
comparison as described for within and among species comparisons. We found no meaningful
interaction of elevation and clade except for elev:CladeHermit in the [Hb] model, whose 95%
credible intervals did not overlap zero (Figure S7). In accordance with lack of phylogenetic
signal in our data, the effect of clade explained minimal variation in model sets for each
parameter (Hb=13%; Hct=8%; TRBC=4%; MCV=2%; MCH=2%; MCHC=4%).
909
910

Among-species analysis
Our final among-species dataset included 58–65 species, depending on the subset for each blood
characteristic ([Hb] = 60 species; Hct = 65 species; TRBC = 58 species, MCV = 60 species;

MCH = 59 species; MCHC = 64 species). Species mean values were calculated for all species
with $n=2$ or more individuals sampled. We split *Patagona gigas* into "north" and "south"
haplotypes in light of evidence that the complex may be comprised of two functionally and
genetically distinct populations separated by strong migratory divides[84], and we analyzed data
only from individuals sampled during the breeding months to ensure correct population
assignment. We designed model sets for each of our six continuous blood response variables. For
each response we analyzed distributions (qqPlots, Cook's Distance, leverage) and removed
outliers that strongly influenced fit. For each model, predictor variables included species-typical
Hb β^A_{13}–β^A_{83} genotype, mean elevation, mean body mass (log-transformed; one mean for all
individuals due to balanced sampling between sexes), mean air temperature, and mean
precipitation. All variables were standardized to a mean of zero and standard deviation of one.
For each response, we compared: (1) an intercept-only null model, (2) a model with all
predictors, and (3) a reduced-predictor model with only those predictors whose 95% CIs did not
overlap zero in full models (reduced model 2). Given the notable quadratic shape of species
mean data for TRBC, MCV, MCH, and MCHC (Figure 2G–L), for those model sets we
additionally compared (4) a model with all predictors plus a quadratic component of elevation
and (5) a reduced-predictor model with quadratic component including only those predictors
whose 95% CIs did not overlap zero in the full quadratic model (reduced model 4). Distribution
family differed by model set: Gaussian for [Hb; skew-normal for Hct; and student's *t*-distribution
for TRBC, MCV, MCH, and MCHC models. We ran four Markov chains for 20,000 iterations
with a burn-in period of 10,000, thinned every 10 steps.
We compared sets of models with LOOIC and WAIC (Table 1). The [Hb] model with the
best (lowest) WAIC and ELPD values was a reduced model with only elevation ($R^2 = 0.18$),
though this model was only marginally better than the full model with all predictors ($R^2 = 0.33$;
Table 1). The top Hct model was the full model with all predictors ($R^2 = 0.24$; Table 1). For
MCH, the top model was a reduced model containing elevation and a quadratic parameter for
elevation, both of which were important predictors ($R^2 = 0.12$; Tables 1 and S5; Figures 2K and
4E). For TRBC, MCV, and MCHC model sets, the intercept-only model received the strongest
support, though other models were closely ranked (Table 1).
943
Cell number versus size analyses
First, to assess within species variation in cell number versus size, we compared a null model
(intercept only) to a single full model *[Hb] ~ MCV + TRBC + (1|species)* for all individuals of
59 species that had [Hb], MCV, and TRBC measurements, distinguishing between *Patagona*
gigas "north" and "south". The species grouping variable allowed us to account for repeated
sampling of individuals within a species and to hold species identity constant. Second, to assess
among species cell number versus size variation, we generated a species mean dataset that
included data from all species with $n=3$ or more individuals sampled (48 species total). We again
compared a null (intercept only) model to a single full model *[Hb] ~ MCV + TRBC.* Third, to
assess how individual species differ in their relative contributions of MCV and TRBC to
increasing [Hb], we repeated our approach for individual species: For each species with $n=8$ or
more individuals sampled (32 species total), we generated a species-specific data subset,
carefully and conservatively evaluated influential outlier points, and ran a single model, *[Hb] ~*
MCV + TRBC (Figure S2). For all three model sets (within species, among species, and single
species), both predictors were standardized to a mean of zero and standard deviation of one.

Using standardized, model-estimated coefficients from each single-species model, for
each species we calculated the Cell Number-Size Index (hereafter *CeNS*) as *TRBC coefficient /*
(MCV coefficient + TRBC coefficient); see main text. *CeNS* ranged from 0.0–1.0 and represented
the proportional contribution of TRBC relative to MCV in driving [Hb], with higher values
corresponding to greater relative contributions of cell number versus cell size.
For both within- and among-species *CeNS* models, full models fit substantially better
than the intercept-only null model (full within species model R^2 = 0.56; full among species
model R^2 = 0.62; Table S6). Based on model estimates, for every increase in TRBC of one
million cells per µl, we expect that [Hb] will increase by 1.81 g/dl within species and 2.15 g/dl
among species, and for every increase in MCV of one fl, we expect that [Hb] will increase by
0.12 g/dl within species and 0.14 g/dl among species.
We then designed a set of models to analyze what contributes to variation in *CeNS*. We
pruned a phylogeny to the subset of 32 species with *CeNS* data for estimation of phylogenetic
signal. Our model response variable was *CeNS* (0–1 index). Predictors, all species mean values,
included elevation, a quadratic elevation term "elevation^2" to account for notable curvature in the
data, body mass (log-transformed), wing loading, and species-typical Hb β^A_{13}–β^A_{83} genotype. We
compared six models: (1) an intercept-only null model, (2) a model with core predictors
(elevation, body mass, and wing loading), (3) a model with core predictors plus elevation^2, (4) a
model with core predictors plus elevation^2 and species-typical Hb β^A_{13}–β^A_{83} genotype, (5) a
reduced version of model 3, and (6) a reduced version of model 4. Reduced models included
only those predictors whose 95% CIs did not overlap zero in full models. In addition to model fit
diagnostics, we assessed correlations between variables by analyzing correlation matrices and by
regressing *CeNS* on each R^2 values from standardized species-specific *CeNS* models,
unstandardized TRBC and MCV beta estimates, and raw TRBC and MCV values (Figure S3).
Distribution family differed by model set: Gaussian for individual-level, species mean, and *CeNS*
prediction models, and student's *t*-distribution for species-specific models. For all *CeNS* models
we ran four Markov chains for 20,000 iterations with a burn-in period of 10,000, thinned every
10 steps.
In our *CeNS* analysis, we compared six models using LOOIC and WAIC (Table S7). The
model with elevation, elevation^2, body mass, and wing loading received the strongest support (R^2
= 0.38) but was closely ranked with a model including only elevation, elevation^2, and body mass
(R^2 = 0.33; Figure 5C; Table S7). Models with elevation^2 fit substantially better than those
without. All but *Coeligena coeligena* showed a positive relationship between *CeNS* and
predictors (Figure S2).
993

Model comparison and diagnostics
All models were run using the brm() function with default priors (weakly informative priors for
the intercept and sigma; improper flat priors for predictors). For all models, to assess
convergence we visually examined trace plots, checked diagnostics, and confirmed that Rhat
values were less than 1.01[82]. We used a posterior predictive check to assess whether model
assumptions yielded good approximations of the data generating process. There were no
problems in any of our models resulting from Bayesian fraction of missing information or
divergent transitions. We compared models using the widely applicable information criterion
(WAIC)[85] values as well as approximate leave-one-out cross-validation information criterion
based on the posterior likelihoods (LOOIC) and Bayesian expected log pointwise predictive
densities (ELPD)[83] to verify full model fits (Table 1). We estimated fixed effects (means and

95% CI) from the posterior distributions for each predictor. For relevant model sets, we
estimated the proportion of the total variance attributed to the species grouping variable, which
accounts for unique aspects of blood parameters that are not captured by the individual, life
history, or environmental predictors.
1009
Breakpoint analysis
We tested for objective thresholds in each blood trait using piecewise linear regression of
elevation (standardized) regressed on each blood trait. In this procedure, we first fit a simple
linear regression model, iteratively searched for objective "breakpoints" along our sampled
elevational gradient, selected the breakpoint with the lowest residual error, and refit the linear
regression model with separate estimates for slopes and intercepts below and above the
breakpoint, respectively. We constrained the model to estimate breakpoints that were each 400 m
from the lowest and highest points in our data (i.e., from 400 m to 4,178 m in elevation), to avoid
biasing breakpoints towards small sample sizes and each low and high elevations, and anomalous
species at both ends of the distribution. We evaluated goodness of fit using fit metrics and by
visually assessing QQplots, Cook's D, leverage and residual distributions. We compared models
using AIC values. All breakpoint models fit significantly better than simple linear regression
models with the exception of the TRBC breakpoint model; this model did fit better in all respects
than the simple linear regression model, but the fit was not significant at the $p = 0.05$ level. We
used a one-way ANOVA to compare whether individuals below and above model-estimated
breakpoints differed in their blood characteristics. Results are presented in Figure S6.