Environmental and taxonomic drivers of bacterial extracellular vesicle production in

marine ecosystems

Steven J. Biller[1,2*], Allison Coe[1], Aldo A. Arellano[1], Keven Dooley[1], Jacqueline S. Gong[2], Emily

A. Yeager[2], Jamie W. Becker[1,3], and Sallie W. Chisholm[1,4]

[1]Department of Civil and Environmental Engineering, Massachusetts Institute of Technology,

Cambridge MA

[2]Department of Biological Sciences, Wellesley College, Wellesley MA

[3]Department of Natural and Biomedical Sciences, Alvernia University, Reading PA

[4]Department of Biology, Massachusetts Institute of Technology, Cambridge MA

*Correspondence: Steven Biller, sbiller@wellesley.edu

Abstract

Extracellular vesicles are small (~50-250 nm diameter) membrane-bound structures

released by cells into their surrounding environment. Vesicles are abundant in the global oceans

and likely play a number of ecological roles in these microbially dominated ecosystems, yet we

know nothing about what influences their production and distributions. Here we examine how

vesicle production varies among different strains of cultivated marine microbes and explore the

degree to which this is influenced by some key environmental variables. We show that vesicle

production rates – the number of vesicles produced per cell per generation – vary across an order

of magnitude in cultures of marine Proteobacteria, Cyanobacteria, and Bacteroidetes. Vesicle

production rates further differ among strains of the cyanobacterium *Prochlorococcus,* and vary

across temperature and light gradients. These data suggest that both community composition and

local environmental conditions modulate the production and standing stock of vesicles in the

oceans. Examining samples from the oligotrophic North Pacific Gyre, we show depth-dependent

changes in the abundance of vesicle-like particles in the upper water column in a manner broadly

consistent with culture observations: highest vesicle abundances are found near the surface,

where light irradiances and temperatures are greatest, and then decrease with depth. This work

represents the beginnings of a quantitative framework for describing extracellular vesicle

dynamics in the oceans – essential as we begin to incorporate vesicles into our ecological and

biogeochemical understanding of marine ecosystems.

Importance

Bacteria secrete extracellular vesicles containing a wide variety of cellular compounds, including lipids, proteins, nucleic acids, and small molecules, into their surrounding environment. These structures are found in diverse microbial habitats, including the oceans, where their distributions vary throughout the water column. Differences in vesicle abundances likely affect their functional impacts within microbial ecosystems, but the factors influencing vesicle distributions in the environment remain poorly understood. Using quantitative analysis of marine microbial cultures, we show that bacterial vesicle production in the oceans is shaped by a combination of biotic and abiotic factors. Our data indicate that different marine taxa release vesicles at rates varying across an order of magnitude, and that vesicle production can change dynamically as a function of environmental conditions. Taken together with direct measurements of vesicle concentrations in the oceans, these culture-based measurements further provide a window into estimating vesicle loss rates. These findings represent a step forward in our understanding of marine vesicle distributions and provide a basis for quantitatively exploring vesicle dynamics in natural ecosystems.

Keywords

Extracellular vesicles, *Prochlorococcus*, *Pelagibacter*, *Alteromonas*, *Thalassospira, Alcanivorax, Marinobacter, Dokdonia, Polaribacter,* Station ALOHA

Introduction

Cells from all domains of life produce membrane-bound structures known as extracellular vesicles. Most of what we know about vesicle release by bacteria is derived from studies of model microbes relevant to advances in molecular biology or biomedical research. Recently, however, the theater has been expanded to examine the microorganisms that serve as the biogeochemical engines of natural ecosystems. In previous work, we demonstrated that the globally abundant marine cyanobacterium *Prochlorococcus* releases vesicles containing a wide range of biomolecules including lipids, nucleic acids, proteins, and metabolites (1–3). Extracellular vesicles are abundant in both coastal and oligotrophic waters, and they are produced by a diverse suite of marine bacteria, archaea, and eukaryotes from both temperate and polar regions (1, 2, 4–7). Though the precise functional roles of vesicles in marine ecosystems remain unknown, lab and field studies suggest that these structures contribute to many biological processes, including horizontal gene transfer, host-phage interactions, and nutrient exchange (1, 8–10). Vesicles represent an essentially unknown feature of the dissolved organic matter in ocean ecosystems (1). Still, we know nothing about what factors influence their production and, in turn, distributions along ocean environmental gradients.

While the production of extracellular vesicles appears to be essentially ubiquitous among bacteria, the cellular mechanisms of vesicle production remain unclear. In Gram-negative bacteria, vesicles released by intact, viable cells are largely thought to arise via 'blebbing' of the outer membrane, where small regions of the lipid bilayer protrude away from the cell and eventually separate in the form of a spherical vesicle (11, 12). These membrane protrusions could originate from numerous sources, such as localized membrane changes, uneven membrane synthesis, membrane turgor pressure, or the influence of small molecules on local regions of the membrane (13–17). Extracellular vesicles can arise through other mechanisms as well, such as through the pinching off of both outer and inner membrane to yield dual-membrane vesicles (18, 19), or via processes associated with the rotation of sheathed flagella (20). Self-annealing of membrane fragments released by cellular lysis can also lead to vesicle formation (21, 22).

Given the underlying complexity and diversity of vesicle biogenesis and release, understanding how this might be regulated – and even the degree to which it is directly regulated in the first place – represents a challenge. In some Gram-negative pathogens and human commensals, vesicle formation appears to involve some degree of genetic control, though it is

not known whether these impacts arise through direct or indirect effects (23–28). Further, stress

induced by changes in temperature (6, 29), nutrient availability, reactive oxygen species, and UV

exposure has also been correlated with increased vesicle production (29–31). These examples

show the potential for vesicle production to vary in response to intracellular and extracellular

conditions, but the magnitude to which such variables affect vesicle release by different microbes

remains an open question.

92   Here we examine factors that may influence vesicle production within marine microbial

ecosystems, focusing on potential roles for community composition and environmental variables.

To this end, we measured vesicle production rates along a spectrum of environmentally relevant

conditions for a suite of taxonomically diverse, and numerically abundant, marine bacteria. We

further explored the degree to which vesicle production varies among closely related strains of

Prochlorococcus, and how vesicle production rates change across gradients of light and

temperature. Finally, we examined the depth-distribution of vesicles at a site in the oligotrophic

North Pacific Gyre to explore how natural vesicle abundances correlate with environmental

parameters.

Results and Discussion

Vesicle production rates vary among marine microbial taxa

We first wondered about the degree to which marine bacteria vary in their production of

extracellular vesicles. Focusing solely on vesicles released by intact, growing cells, we measured

production rates by several diverse marine microbes during exponential growth in terms of

vesicles produced per cell per generation, a standardized metric that enables comparisons across

differences in both cell abundances and growth rates (Methods and (1)). We surveyed

representatives from several of the most abundant bacterial phyla in marine systems –

Proteobacteria, Cyanobacteria, and Bacteroidetes – and found that their vesicle release rates,

when growing under similar conditions, spanned across an order of magnitude (Fig. 1a). Median

vesicle production rates during exponential growth ranged from ~4 vesicles cell^{-1} generation^{-1}

among strains of *Prochlorococcus* to ~58 vesicles cell^{-1} generation^{-1} in the heterotroph

Marinobacter, with other taxa producing vesicles at intermediate values.

Though the total amount of cellular surface area available to form membrane vesicles

might be expected to play some role in bounding the number of vesicles that can be formed at

any time, vesicle production was not significantly associated with cell surface area (Fig. S1a).

Cellular growth rates, on the other hand, were positively correlated with vesicle release rates

(Pearson's correlation $p < 0.05$), but growth rates only explained a portion of the variation in

vesicle production (see below). Together, our results indicate that vesicle production rates differ

markedly among marine taxa, and are due at least in part to physiological differences between

strains.

Growth conditions impact vesicle production by marine heterotrophs

To explore how environmental conditions might influence vesicle production, we first

measured vesicle production rates of selected strains of marine heterotrophic bacteria under

different culture conditions. Focusing first on media composition, we compared vesicle

production rates by *Pelagibacter* (SAR11) grown in either a defined artificial seawater medium

(AMS1) or a variant based on natural seawater but with otherwise equivalent nutrient

amendments (ProMS*; see Methods). *Pelagibacter* grew 3.2x faster in ProMS* than AMS1, and

vesicle release rates increased by ~15x (from ~0.4 vesicles cell^{-1} generation^{-1} in AMS1 to ~6

vesicles cell^{-1} generation^{-1} in ProMS*; Fig. 1b). While we do not know which component(s) of

the natural seawater background are responsible for the difference, this indicates that nutrient

availability can affect vesicle production. We next asked whether changes in temperature

influenced vesiculation by comparing production rates at two temperatures in *Marinobacter* and

Thalassospira. An increase in growth temperature from 21 °C to 24 °C increased the rate of

vesicle production by a factor of ~4-8x in both strains (Fig. 1c), broadly consistent with trends

observed in host-associated and terrestrial microbes such as *Pseudomonas aeruginosa* (29) and

Acinetobacter baylyi (32).

Looking across all heterotrophs and growth conditions examined, cells tended to produce

more vesicles per generation when they grew faster (Fig. S2). On average, each growth rate

increase of 1 day^{-1} resulted in ~2.6 additional vesicles released per cell per generation. However,

changes in growth rate only accounted for ~26% of the observed variance (linear regression $p <$

0.01; Fig. S2), indicating that other factors are at play. One possible explanation for the influence

of growth rate on vesicle production is that increasing growth rate requires increased cell

wall/membrane biosynthesis activity, which could in turn transiently weaken the connections

between the outer membrane and the rest of the cell to create more opportunities for vesicle

release. Environmental conditions might also modulate vesicle production via changes in

membrane composition and/or fluidity (17).

Combinatorial impacts of light and temperature on Prochlorococcus *vesicle production*

153   For abundant marine photosynthetic organisms such as *Prochlorococcus*, light and

temperature are key environmental variables impacting cellular growth and physiology (33–35),

and we suspected that they also influence vesicle production. To explore this, we measured

production rates under different combinations of light irradiance and temperature using five

different strains of *Prochlorococcus* that encompass phylogenetically and ecologically distinct

high-light and low-light adapted clades (Table 1). Median production rates varied significantly

among strains, and ranged from 1.2 – 7.7 vesicles cell^{-1} generation^{-1} across all conditions tested

(two-way ANOVA, $p < 0.01$; Table 2; Fig. 2a). Vesicle production by individual

Prochlorococcus strains was remarkably dynamic, varying by an average factor of ~27 between

the lowest and highest production rates observed (Table 2). On average, high light-adapted

Prochlorococcus produced fewer vesicles per cell per generation than low light-adapted strains;

the high-light strains also had a lower dynamic range of production rates (Table 2). What might

account for this? High light-adapted *Prochlorococcus* are smaller than low light-adapted cells

but, as with the heterotrophs, differences in cell size had little explanatory value for differences

in vesicle production rates (Fig. S1b). Unlike the heterotrophs, however, differences in

Prochlorococcus growth rate were not significantly correlated with variation in vesicle

production (linear model $R^2 = 0.065$; Figs. 2a and S4), suggesting that other physiological factors

must play key roles.

171   Looking at the impact of light and temperature, we found that individual strains

responded differently. Vesicle production by the high-light adapted strains did not significantly

change as a function of irradiance (Table 2; Fig. 2b), as consistent trends were only observed at

some, but not all, temperatures (Figs. 2a, S3). By contrast, the low-light adapted cells exhibited

significant variation in production rates with both light and temperature, again with higher light

or increased temperature leading to increased vesicle production rates (Figs. 2a, 2b, S3; Table 2).

Differences in these production patterns thus join the properties that contribute to the ecological

differentiation between high-light and low-light adapted *Prochlorococcus*.

Vesicle release as a response to light stress in Prochlorococcus

Our experiments indicate that light and temperature impact *Prochlorococcus* vesicle

production in complex ways. Given that most strains showed an increase in production at the

highest light intensities tested (Fig. 2b), we wondered whether differences in production might

be associated with light stress. Irradiance levels higher than that which can be utilized by

photosystems can disrupt intracellular redox balance, inhibit photosynthesis, or generate reactive

oxygen species in cyanobacteria. Cells must then activate one or more stress response pathways

to repair the damage and/or help to manage the excess energy input (36–38). Across all

Prochlorococcus strains examined, production rates were not influenced by irradiance when cells

were grown at or below the levels which maximized growth rates (Figs 2c, S4). Above this light

optimum, when cells are assumed to experience some degree of light stress, production was

significantly and positively correlated with relative increases in light (R^2=0.23, linear model *p* <

0.01; Fig. 2c). Changepoint threshold regression analysis corroborated this difference, indicating

a significant step increase in vesiculation when light levels were increased above the growth

optima at each particular temperature (threshold range: 1-2x optimal light level, *p* < 0.001). This

suggests that chronic light stress may directly or indirectly lead to increases in vesicle release.

Might vesicle production directly contribute to mitigating light stress, or does it instead

reflect a side effect of other physiological changes occurring within the cell? Though we can

only speculate at this point, cyanobacteria can utilize several strategies to dissipate excess light

energy during light stress, including modifying their light-harvesting apparatus,

nonphotochemical quenching, and increased production of enzymes with antioxidant functions

(37). In addition, *Prochlorococcus* can utilize excess light energy to power increased levels of

carbon fixation and then 'dump' the unneeded organic molecules into the extracellular space to

maintain redox balance (36). While perhaps an indirect metabolic route, increased synthesis of

lipids and secretion of fixed organic compounds via vesicles might conceivably provide yet

another route for releasing the products of overflow metabolism. Intracellular oxidative stress,

such as might arise from high light conditions, increases the number of vesicles produced by

Pseudomonas aeruginosa (29) and a similar mechanism could be at play here. *Prochlorococcus*

vesicles can contain oxidized metabolites, raising the possibility that they could aid the cell by

removing cellular material damaged by oxidative stress (3, 12). While there remains much to

uncover about this association, differences in light stress responses may explain, in part, the

different degree to which light impacts vesicle release by high-light vs low-light adapted

Prochlorococcus.

Vesicle size varies across microbial taxa and environmental conditions

Bacteria release extracellular vesicles of varying sizes, often ranging between ~50 - 250

nm diameter (12). Vesicle size distributions can differ among strains of the same organism (39),

as a function of growth phase (40), and in response to changes in membrane composition (41).

As vesicle size is associated with differences in both proteomic (42) and DNA content (43),

shifts in size distributions may have functional consequences. Among the marine microbes we

studied, vesicle sizes differed significantly by taxonomic group (one-way ANOVA $p < 0.001$;

Fig. 3), with members of the Bacteroidetes releasing the largest (~170 nm mode diameter) and

Prochlorococcus releasing the smallest (~70 nm). Vesicle size was not significantly correlated

with cellular growth rate or production rate across the heterotrophs or *Prochlorococcus* strains.

However, within *Prochlorococcus*, vesicle size did vary with light intensity and, to a lesser

extent, growth temperature (Fig. S5; two-way ANOVA $p < 1e-6$ and $p < 0.01$, respectively).

Strains grown at lower light intensities tended to release larger vesicles than cells experiencing

higher light fluxes, even at levels below the optimum (Fig. S5a). The largest vesicles also tended

to come from cultures grown in the middle of the temperature range tested (Fig. S5b).

Do taxa differ in the relative investment of resources they put into vesicles? As an initial

approach to addressing this question, we calculated how much membrane material is required to

produce the median number of vesicles released per generation compared to each cell's total

surface area. Despite differences in cell size, vesicle diameter, and vesicle production rates, we

estimate that most strains, including *Prochlorococcus, Pelagibacter*, and *Alteromonas*, release

vesicles equivalent to a median of ~3 – 6% of their cell surface lipid membranes per generation

(Fig. S6). *Alcanivorax* and *Marinobacter* were notably distinct from the others and released

vesicles equivalent to ~40% of their surface area per generation (Fig. S6). Forming a given size

vesicle requires proportionately less membrane material from larger cells than from smaller cells,

but it appears that increased production rates can balance these lower proportional investments in

individual vesicles. Given the dynamic range of vesicle production rates, and thus relative

cellular investment, cells must substantially increase membrane synthesis under certain growth

conditions, and this burden may not fall just on the smallest cells. The costs and benefits of

changing vesicle production rates under a given environmental context, especially when

considering other resources invested besides just the membranes, are clearly complex (44).

Extracellular vesicle abundances across environmental gradients in the North Pacific

We have shown that extracellular vesicle concentrations released by the entire microbial

community decreases with depth at an oligotrophic site in the Atlantic ocean (1). To explore

whether those results may be generalizable to other regions of the ocean, we examined vesicle

abundances at Station ALOHA in the North Pacific subtropical gyre (Fig. 4a). As at the Atlantic

site, vesicle-like particles were most abundant near the surface, with their concentration

decreasing by approximately an order of magnitude over the upper 500 m. Vesicle-like particle

concentrations in this set of Pacific ocean samples were of the same magnitude as those

previously seen in the Atlantic and declined with depth to a similar degree. This overall

congruence of results across different oceans, seasons, and years, provides an early indication

that this general pattern of vesicle distributions in the water column might be generally

representative for mid-latitude oligotrophic waters.

Assuming loss rates are constant with depth, the data from our culture studies would

broadly predict that, all else being equal, vesicle abundances should be greatest where light

irradiance and temperature are highest, and this is what we observed (Fig. 4b-c). Not only were

overall vesicle-like particle concentrations greatest at the surface, but there were proportionally

more vesicles per cell at the surface than in deeper samples, where environmental conditions

would be expected to yield lower vesicle production rates (Fig. 4). These data are consistent with

local environmental factors, such as those explored above, influencing *in situ* vesicle distribution

patterns.

Given production rates in pure cultures, measurements of microbial standing stocks in the

wild, and estimates of *in situ* cellular growth rates, we can also begin to estimate vesicle loss

rates – an aspect of vesicle dynamics in the environment that is completely unknown. Assuming

that marine phytoplankton grow at a rate of ~0.5 per day (45, 46), with heterotrophs growing at

~0.1 per day (45), the vesicle production rates measured here suggest that the surface community

at Station ALOHA might produce in excess of 1 vesicle cell^{-1} day^{-1}, or $>10^5$ vesicles mL^{-1} day^{-1}.

This estimated daily input vesicle flux is comparable to the measured standing stocks in the

oceans; if we further assume that standing vesicle concentrations are roughly constant, this

would suggest that vesicle loss rates must be similar to production rates. Vesicle production rates

in the ocean could certainly differ from those measured under laboratory conditions due to strain

differences, nutrient availability, the influence of biotic or abiotic interactions, consideration of

vesicle production through cell lysis mechanisms, or other factors. Regardless, loss rates must be

significant, and could arise from processes including natural decay, consumption or degradation,

fusion with other cells, or physical transport. More work is needed to arrive at a detailed estimate

of losses and examine the relative contribution of these and other mechanisms.

Conclusions

This study shows that extracellular vesicle production rates differ among marine

microbes and highlights the dynamic nature of vesicle production by individual bacterial cells.

The composition of extracellular vesicle populations in a given location will be influenced by a

complex set of factors involving differences in cellular community structure and the vesicle

production rate of each taxa under local environmental conditions. As differences in cell and

vesicle concentrations would be expected to impact their encounter rates in an aqueous,

planktonic environment, vesicle-mediated interaction dynamics may also changes as a result of

environmental shifts. Studies addressing the fate and turnover of vesicles will be essential for

developing detailed models of vesicle distributions and functions in the oceans.

Materials and Methods

Culturing conditions

Axenic *Prochlorococcus* cells were grown in 0.2 μm filtered and autoclaved Sargasso

Sea water amended with Pro99 nutrients (47). Triplicate cultures were grown on a 14:10 hr

light:dark cycle at temperatures ranging from 15 °C to 26 °C and at four light levels: very high

light (VHL), high light (HL), medium light (ML), and low light (LL), with specific values

determined by each strain's light tolerance (Table 1). Culture axenicity was regularly verified

using a suite of purity test broths (MPTB, ProAC, ProMM) (48–50) and by flow cytometry.

Cultures were maintained at the indicated temperature and light intensity for at least 12

generations (approximately four complete transfers), and were not sampled until the culture

reached balanced growth. Growth was routinely monitored by culture fluorescence with a 10AU

or TD700 Fluorometer (Turner Designs), and growth rates were calculated by exponential

regression from the log-linear portion of the growth curve.

Alcanivorax MIT1350 was isolated from a water sample collected at 150m depth from

Station ALOHA (22.75 °N 158 °W) in the North Pacific in 2013. This strain was flow-sorted

from a 'small cell, low nucleic acid' population derived from a *Prochlorococcus* enrichment

culture. Following isolation, *Alcanivorax* MIT1350, along with *Alteromonas macleodii* strain

MIT1002 (51), *Thalassospira* strain MIT1004 (2), and *Marinobacter* MIT1353 (52) were

maintained in ProMM medium. The genome sequence of *Alcanivorax* MIT1350 is available at

the IMG database (53) under Genome ID 2681813576.

Pelagibacterales sp. HTCC7211, a member of the abundant warm-water surface-

dwelling Ia.3 ecotype of the SAR11 bacterial clade (54), was cultured in either a defined

artificial medium (AMS1; (55)) or Sargasso seawater-based ProMS* (a modified version of

ProMS media (52) with pyruvate, glycine, methionine, and vitamin mixes added to

concentrations equal to those in AMS1). *Dokdonia* MED134 (56) and *Polaribacter* MED152

(57) were grown in in Pro99 media supplemented with 5 g L^{-1} peptone and 1 g L^{-1} yeast extract

(BD Difco). All heterotrophs were grown at 24 °C, except for HTCC7211, which was

maintained at 22 °C.

We manually examined the genome sequence of each strain listed above and found no

evidence for the presence of lysogenic phage and/or gene transfer agents in these organisms

(based on BLASTp searches using the *Rhodobacter capsulatus* gene transfer agent sequences

RCAP_rcc01682 - RCAP_rcc01698), and the cultures contained no evidence of lytic phage

activity. Given this and prior electron microscopy data (2), our data assume that all particles

between ~50-250 nm diameter in these culture supernatants are extracellular vesicles.

Cell and vesicle enumeration

At each time point sampled, cells were preserved for flow cytometry by fixation in

0.125% glutaraldehyde, flash frozen in liquid nitrogen, and stored at -80 ºC until analysis.

Vesicles were sampled by collecting 1 mL of culture and filtering it through a 0.22 µm Supor

syringe filter (Pall). The <0.22 µm fraction was stored at -20 ºC until ready for analysis.

Prochlorococcus cells were enumerated using a Guava easyCyte 12HT flow cytometer

(Luminex). Heterotrophs were stained with 1x SYBR Green I (Molecular Probes) for 1 hr and

then counted using either an easyCyte 12HT or a ZE5 flow cytometer (Bio-Rad).

Prochlorococcus cell sizes were normalized to 2 μm beads (Polysciences). Cells were excited

with a blue 488 nm laser and analyzed for red fluorescence (692/40 nm), green fluorescence

(530/40 nm), and size (forward scatter). Cell lengths were determined by imaging >500 cells on

a Zeiss Axioskop microscope equipped with a 100x objective lens and a Nikon D90 digital

camera. Cell lengths were determined using ImageJ, and calibrated using a 1mm / 0.01mm scale

stage micrometer (Meiji Techno).

Vesicles were measured from samples collected at two or more timepoints during

exponential growth phase. Vesicles were enumerated by nanoparticle tracking analysis using a

NanoSight LM10HS instrument (Malvern/NanoSight), equipped with a LM14 blue laser module

and NTA software V3.1. Three technical replicate videos (60s each) were collected from each

sample using a camera level setting of 11. Samples containing >80 particles per frame were

diluted with clean seawater media to a final concentration between 20-80 particles per frame.

The sample chamber was thoroughly flushed with 18.2 MΩ cm^{-1} water (Milli-Q; Millipore)

between samples, and visually examined to ensure that no particles were carried over between

samples. Videos were analyzed using NTA V3.1 software with a threshold of 1. Vesicle

concentrations in each sample are based only on particles between 50 - 250 nm diameter. Media

blank controls were routinely run for each batch of media used, and any background

concentration was subtracted. A small number of samples containing abnormally high particle

counts (defined as containing >10x higher concentrations than other replicates of that timepoint

and/or the surrounding timepoints) or in which large, highly refractive particles were present

(which mask the underlying particle counts) were removed from the final dataset. The source of

these contaminants is unknown and their occurrence in samples did not follow any obvious

systematic patterns; we hypothesize that they could have arisen from manufacturing variation or

defects in individual filters, plastic tubes used to store samples, or plastic syringes used in fluid

handling.

Vesicle production and surface area calculations

We measured vesicle production rates as the average number of vesicles produced on a

per cell, per generation basis based on averaged cell and vesicle measurements from all

biological replicates at each timepoint. These calculations are based on a model that incorporates

the relative rates of change in cell and vesicle concentrations in pure cultures over time as

detailed in (1). To better account for variation in slope across the time course, we computed the

average number of vesicles produced per cell per generation between each successive pair of

timepoints obtained during the exponential growth phase.

The surface area of vesicles from an individual strain was calculated assuming that

vesicles are spherical structures with a diameter equivalent to the mode value measured. Cellular

surface area is based on microscopy measurements and literature values (58, 59).

Prochlorococcus was estimated to be spherical, and the surface area of other cells were

estimated as cylinders with a hemisphere on each end. Relative cellular investment in vesicles

per generation was defined as (individual vesicle surface area * vesicle production rate per cell

per generation) / cell surface area.

Statistics

All statistical analyses were performed in R (V 4.0.2) and plots were produced using the

ggplot2 package (60). ANOVA analysis of vesicle production rate data was performed on log-

transformed values. Model selection for two-way ANOVAs was carried out using the Akaike

information criterion metric, as implemented in the 'AICcmodavg' R package (61). Changepoint

threshold regression analysis was carried out using the 'chngpt' R package (62). Contour plots of

growth rate data were generated using the 'metR' and 'interp' R packages (63, 64).

Field sampling

Seawater samples for vesicle analysis were collected from Station ALOHA on cruise

HOT263 (June 2014). At each depth, we collected 100 - 175 L using a Niskin bottle array and

concentrated the < 0.2 μm fraction by tangential flow filtration following previously described

protocols (1). Vesicles were enriched across an Optiprep density gradient (Iodixanol; Sigma-

Aldrich) as previously described (1). Enumeration of field concentrations were based on

Nanosight concentration measurements of gradient-purified fractions containing the highest

abundance of vesicles (between ~1.14 – 1.19 g/mL) and consideration of the original sampled

volume. We note that particle losses incurred during the sample processing and analysis pipeline,

thus these *in situ* measurements should be considered lower-bound estimates. Cellular

abundance, PAR and temperature values were provided by the Hawaii Ocean Time-series

program (https://hahana.soest.hawaii.edu/hot/hot-dogs/interface.html). As vesicle-like particles

could be produced by microbes from all domains of life, cell abundances represent the combined

counts for cyanobacteria, heterotrophic bacteria, and eukaryotes from the indicated depths.

Acknowledgments

We thank the captain and crew of HOT263 for their help with obtaining field samples.

Stephen Giovannoni (Oregon State U.) kindly supplied *Pelagibacteriales sp.* HTCC7211, and

Jarone Pinhassi (Linnaeus U.) shared the *Polaribacter* and *Dokdonia* cultures. We are also

grateful to Christina Bliem and Madeline Williams for providing laboratory assistance. HOT

data were obtained via the Hawaii Ocean Time-series HOT-DOGS application; University of

Hawai'i at Mānoa National Science Foundation Award # 1756517.

This work was supported by the National Science Foundation (OCE-2049004 and DBI-

2018337 to SJB, OCE-1356460 to SWC) and the Simons Foundation (SCOPE Award ID

329108, SWC). This is a contribution of the Simons Collaboration on Ocean Processes and

Ecology (SCOPE).

The authors declare no conflicts of interest.

References

1. Biller SJ, Schubotz F, Roggensack SE, Thompson AW, Summons RE, Chisholm SW. 2014. Bacterial vesicles in marine ecosystems. Science 343:183–186.

2. Biller SJ, McDaniel LD, Breitbart M, Rogers E, Paul JH, Chisholm SW. 2017. Membrane vesicles in sea water: heterogeneous DNA content and implications for viral abundance estimates. ISME J 11:394–404.

3. Biller SJ, Lundeen RA, Hmelo LR, Becker KW, Arellano AA, Dooley K, Heal KR, Carlson LT, Van Mooy BAS, Ingalls AE, Chisholm SW. 2021. *Prochlorococcus* extracellular vesicles: molecular composition and adsorption to diverse microbes. Environ Microbiol 1462-2920.15834.

4. Baeza N, Mercade E. 2021. Relationship between membrane vesicles, extracellular ATP and biofilm formation in Antarctic Gram-negative bacteria. Microb Ecol 81:645–656.

5. Erdmann S, Tschitschko B, Zhong L, Raftery MJ, Cavicchioli R. 2017. A plasmid from an Antarctic haloarchaeon uses specialized membrane vesicles to disseminate and infect plasmid-free cells. Nat Microbiol 2:1446–1455.

6. Frias A, Manresa A, de Oliveira E, López-Iglesias C, Mercade E. 2010. Membrane vesicles: A common feature in the extracellular matter of cold-adapted Antarctic bacteria. Microb Ecol 59:476–486.

7. Yokoyama F, Kawamoto J, Imai T, Kurihara T. 2017. Characterization of extracellular membrane vesicles of an Antarctic bacterium, *Shewanella livingstonensis* Ac10, and their enhanced production by alteration of phospholipid composition. Extremophiles 21:723–731.

8. Biller SJ. 2020. Extracellular Vesicles in the Environment, p. 75–99. *In* Kaparakis-Liaskos, M, Kufer, TA (eds.), Bacterial Membrane Vesicles. Springer International Publishing, Cham.

9. Lynch JB, Alegado RA. 2017. Spheres of hope, packets of doom: the good and bad of outer membrane vesicles in interspecies and ecological dynamics. J Bacteriol 199:e00012-17.

10. Schatz D, Vardi A. 2018. Extracellular vesicles - new players in cell-cell communication in aquatic environments. Curr Opin Microbiol 43:148–154.

11. Pathirana R, Kaparakis-Liaskos M. 2016. Bacterial membrane vesicles: biogenesis, immune regulation and pathogenesis. Cell Microbiol 18:1518–1524.

12. Schwechheimer C, Kuehn MJ. 2015. Outer-membrane vesicles from Gram-negative bacteria: biogenesis and functions. Nat Rev Microbiol 13:605–619.

13. Baumgarten T, Vazquez J, Bastisch C, Veron W, Feuilloley MGJ, Nietzsche S, Wick LY, Heipieper HJ. 2012. Alkanols and chlorophenols cause different physiological adaptive responses on the level of cell surface properties and membrane vesicle formation in *Pseudomonas putida* DOT-T1E. Appl Microbiol Biotechnol 93:837–845.

14. Kobayashi H, Uematsu K, Hirayama H, Horikoshi K. 2000. Novel toluene elimination system in a toluene-tolerant microorganism. J Bacteriol 182:6451–6455.

15. Mashburn LM, Whiteley M. 2005. Membrane vesicles traffic signals and facilitate group activities in a prokaryote. Nature 437:422–425.

16.   Schertzer JW, Whiteley M. 2012. A bilayer-couple model of bacterial outer membrane
vesicle biogenesis. mBio 3:e00297-11.

17.   Toyofuku M, Nomura N, Eberl L. 2019. Types and origins of bacterial membrane
vesicles. Nat Rev Microbiol 17:13–24.

18.   Perez-Cruz C, Carrion O, Delgado L, Martinez G, Lopez-Iglesias C, Mercade E. 2013.
New type of outer membrane vesicle produced by the Gram-negative bacterium *Shewanella*
vesiculosa M7T: Implications for DNA content. Appl Environ Microbiol 79:1874–1881.

19.   Pérez-Cruz C, Delgado L, López-Iglesias C, Mercade E. 2015. Outer-inner membrane
vesicles naturally secreted by Gram-negative pathogenic bacteria. PLoS ONE 10:e0116896.

20.   Aschtgen M-S, Lynch JB, Koch E, Schwartzman J, McFall-Ngai M, Ruby E. 2016.
Rotation of *Vibrio fischeri* flagella oroduces outer membrane vesicles that induce host
development. J Bacteriol 198:2156–2165.

21.   Mandal PK, Ballerin G, Nolan LM, Petty NK, Whitchurch CB. 2021. Bacteriophage
infection produces membrane vesicles in *Escherichia coli* via both explosive cell lysis and
membrane blebbing. Microbiology 167: 001021.

22.   Turnbull L, Toyofuku M, Hynen AL, Kurosawa M, Pessi G, Petty NK, Osvath SR,
Cárcamo-Oyarce G, Gloag ES, Shimoni R, Omasits U, Ito S, Yap X, Monahan LG, Cavaliere R,
Ahrens CH, Charles IG, Nomura N, Eberl L, Whitchurch CB. 2016. Explosive cell lysis as a
mechanism for the biogenesis of bacterial membrane vesicles and biofilms. Nat Commun
7:11220–13.

23.   Bernadac A, Gavioli M, Lazzaroni JC, Raina S, Lloubès R. 1998. *Escherichia coli tol-pal*
mutants form outer membrane vesicles. J Bacteriol 180:4872–4878.

24.   Kulp AJ, Sun B, Ai T, Manning AJ, Orench-Rivera N, Schmid AK, Kuehn MJ. 2015.
Genome-wide assessment of outer membrane vesicle production in *Escherichia coli*. PLoS ONE
10:e0139200-16.

25.   McBroom AJ, Johnson AP, Vemulapalli S, Kuehn MJ. 2006. Outer membrane vesicle
production by *Escherichia coli* is independent of membrane instability. J Bacteriol 188:5385–
5392.

26.   Nakayama-Imaohji H, Hirota K, Yamasaki H, Yoneda S, Nariya H, Suzuki M, Secher T,
Miyake Y, Oswald E, Hayashi T, Kuwahara T. 2016. DNA inversion regulates outer membrane
vesicle production in *Bacteroides fragilis*. PLoS ONE 11:e0148887-21.

27.   Rath P, Huang C, Wang T, Wang T, Li H, Prados-Rosales R, Elemento O, Casadevall A,
Nathan CF. 2013. Genetic regulation of vesiculogenesis and immunomodulation in
Mycobacterium tuberculosis. Proc Natl Acad Sci 110:E4790–E4797.

28.   Resch U, Tsatsaronis JA, Le Rhun A, Stübiger G, Rohde M, Kasvandik S, Holzmeister S,
Tinnefeld P, Wai SN, Charpentier E. 2016. A two-component regulatory system impacts
extracellular membrane-derived vesicle production in Group A *Streptococcus*. mBio 7:e00207-
16.

29.   MacDonald IA, Kuehn MJ. 2013. Stress-induced outer membrane vesicle production by
Pseudomonas aeruginosa. J Bacteriol 195:2971–2981.

30.   Gamalier JP, Silva TP, Zarantonello V, Dias FF, Melo RCN. 2017. Increased production
of outer membrane vesicles by cultured freshwater bacteria in response to ultraviolet radiation.
Microbiol Res 194:38–46.

31.   Zarantonello V, Silva TP, Noyma NP, Gamalier JP, Mello MM, Marinho MM, Melo
RCN. 2018. The cyanobacterium *Cylindrospermopsis raciborskii* (CYRF-01) responds to
environmental stresses with increased vesiculation detected at single-cell resolution. Front
Microbiol 9:272.

32.   Fulsundar S, Harms K, Flaten GE, Johnsen PJ, Chopade BA, Nielsen KM. 2014. Gene
transfer potential of outer membrane vesicles of *Acinetobacter baylyi* and effects of stress on
vesiculation. Appl Environ Microbiol 80:3469–3483.

33.   Moore L, Chisholm S. 1999. Photophysiology of the marine cyanobacterium
Prochlorococcus: Ecotypic differences among cultured isolates. Limnol Oceanogr 44:628–638.

34.   Moore L, Goericke R, Chisholm S. 1995. Comparative physiology of *Synechococcus* and
Prochlorococcus: influence of light and temperature on growth, pigments, fluorescence and
absorptive properties. Mar Ecol Prog Ser 116:259–275.

35.   Moore LR, Rocap G, Chisholm SW. 1998. Physiology and molecular phylogeny of
coexisting *Prochlorococcus* ecotypes. Nature 393:464–467.

36.   Braakman R, Follows MJ, Chisholm SW. 2017. Metabolic evolution and the self-
organization of ecosystems. Proc Natl Acad Sci 114:E3091–E3100.

37.   He Q, Dolganov N, Bjorkman O, Grossman AR. 2001. The high light-inducible
polypeptides in *Synechocystis* PCC6803. Expression and function in high light. J Biol Chem
276:306–314.

38.   Muramatsu M, Hihara Y. 2011. Acclimation to high-light conditions in cyanobacteria:
from gene expression to physiological responses. J Plant Res 125:11–39.

39.   Tartaglia NR, Nicolas A, Rodovalho V de R, Luz BSR da, Briard-Bion V, Krupova Z,
Thierry A, Coste F, Burel A, Martin P, Jardin J, Azevedo V, Loir YL, Guédon E. 2020.
Extracellular vesicles produced by human and animal *Staphylococcus aureus* strains share a
highly conserved core proteome. 1. Sci Rep 10:8467.

40.   Zavan L, Bitto NJ, Johnston EL, Greening DW, Liaskos MK. 2019. *Helicobacter pylori*
growth stage determines the size, protein composition, and preferential cargo packaging of outer
membrane vesicles. Proteomics 19:e1800209–e1800223.

41.   Murphy K, Park AJ, Hao Y, Brewer D, Lam JS, Khursigara CM. 2014. Influence of O
polysaccharides on biofilm development and outer membrane vesicle biogenesis in
Pseudomonas aeruginosa PAO1. J Bacteriol 196:1306–17.

42.   Turner L, Bitto NJ, Steer DL, Lo C, D'Costa K, Ramm G, Shambrook M, Hill AF,
Ferrero RL, Kaparakis-Liaskos M. 2018. *Helicobacter pylori* outer membrane vesicle size
determines their mechanisms of host cell entry and protein content. Front Immunol 9:1466.

43.   Bitto NJ, Chapman R, Pidot S, Costin A, Lo C, Choi J, Cruze TD x02019, Reynolds EC,
Dashper SG, Turnbull L, Whitchurch CB, Stinear TP, Stacey KJ, Ferrero RL. 2017. Bacterial
membrane vesicles transport their DNA cargo into host cells. Sci Rep 7:7072.

44.   McMillan HM, Kuehn MJ. 2021. The extracellular vesicle generation paradox: a bacterial
point of view. EMBO J 40:e108174.

45.   Kirchman DL. 2016. Growth rates of microbes in the oceans. Annu Rev Mar Sci 8:285–
309.

46.   Vaulot D, Marie D, Olson RJ, Chisholm SW. 1995. Growth of *Prochlorococcus*, a
photosynthetic prokaryote, in the equatorial Pacific Ocean. 268:1480–1482.

47.   Moore L, Coe A, Zinser ER, Saito M, Sullivan M, Lindell D, Frois-Moniz K, Waterbury
J, Chisholm S. 2007. Culturing the marine cyanobacterium *Prochlorococcus*. Limnol Ocean
Methods 5:353–362.

48.   Berube PM, Biller SJ, Kent AG, Berta-Thompson JW, Roggensack SE, Roache-Johnson
KH, Ackerman M, Moore LR, Meisel JD, Sher D, Thompson LR, Campbell L, Martiny AC,
Chisholm SW. 2015. Physiology and evolution of nitrate acquisition in *Prochlorococcus*. ISME
J 9:1195–1207.

49.   Morris JJ, Kirkegaard R, Szul MJ, Johnson ZI, Zinser ER. 2008. Facilitation of robust
growth of *Prochlorococcus* colonies and dilute liquid cultures by "helper" heterotrophic bacteria.
Appl Environ Microbiol 74:4530–4534.

50.   Saito M, Moffett J, Chisholm S, Waterbury J. 2002. Cobalt limitation and uptake in
Prochlorococcus. Limnol Oceanogr 47:1629–1636.

51.   Biller SJ, Coe A, Martin-Cuadrado A-B, Chisholm SW. 2015. Draft genome sequence of
Alteromonas macleodii strain MIT1002, isolated from an enrichment culture of the marine
cyanobacterium *Prochlorococcus*. Genome Announc 3:e00967-15.

52.   Becker JW, Hogle SL, Rosendo K, Chisholm SW. 2019. Co-culture and biogeography of
Prochlorococcus and SAR11. ISME J 13:1506–1519.

53.   Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A, Huang
J, Woyke T, Huntemann M, Anderson I, Billis K, Varghese N, Mavromatis K, Pati A, Ivanova
NN, Kyrpides NC. 2014. IMG 4 version of the integrated microbial genomes comparative
analysis system. Nucleic Acids Res 42:D560-7.

54.   Brown MV, Lauro FM, Demaere MZ, Muir L, Wilkins D, Thomas T, Riddle MJ,
Fuhrman JA, Pfannkoch CA, Hoffman JM, McQuaid JB, Allen A, Rintoul SR, Cavicchioli R.
2012. Global biogeography of SAR11 marine bacteria. Mol Syst Biol 8:595.

55.   Carini P, Steindler L, Beszteri S, Giovannoni SJ. 2013. Nutrient requirements for growth
of the extreme oligotroph "*Candidatus* Pelagibacter ubique" HTCC1062 on a defined medium.
ISME J 7:592–602.

56.   Gómez-Consarnau L, González JM, Coll-Lladó M, Gourdon P, Pascher T, Neutze R,
Pedros-Alio C, Pinhassi J. 2007. Light stimulates growth of proteorhodopsin-containing marine
Flavobacteria. Nature 445:210–213.

57.   González JM, Fernandez-Gomez B, Fernàndez-Guerra A, Gómez-Consarnau L, Sánchez
O, Coll-Lladó M, del Campo J, Escudero L, Rodríguez-Martínez R, Alonso-Sáez L, Latasa M,
Paulsen I, Nedashkovskaya O, Lekunberri I, Pinhassi J, Pedros-Alio C. 2008. Genome analysis
of the proteorhodopsin-containing marine bacterium *Polaribacter* sp. MED152 (Flavobacteria).
Proc Natl Acad Sci 105:8724–8729.

58.   Rappé MS, Connon SA, Vergin KL, Giovannoni SJ. 2002. Cultivation of the ubiquitous
SAR11 marine bacterioplankton clade. Nature 418:630–633.

59.   Ting CS, Hsieh C, Sundararaman S, Mannella C, Marko M. 2007. Cryo-electron
tomography reveals the comparative three-dimensional architecture of *Prochlorococcus*, a
globally important marine cyanobacterium. J Bacteriol 189:4485–4493.

60.   Wickham H. 2009. ggplot2: elegant graphics for data analysis. Springer New York.

61.   Mazerolli MJ. 2020. AICcmodavg: Model selection and multimodel inference based on
(Q)AIC(c). R package version 2.3-1. https://cran.r-project.org/package=AICcmodavg.

62.   Fong Y, Huang Y, Gilbert PB, Permar SR. 2017. chngpt: threshold regression model
estimation and inference. BMC Bioinformatics 18:454.

63.   Campitelli E. 2021. metR: Tools for Easier Analysis of Meteorological Fields. R package
version 0.10.0. https://github.com/eliocamp/metR.

64.   Gebhardt A, Bivand R, Sinclair L. 2020. interp: Interpolation Methods. R package
version 1.0-33. https://CRAN.R-project.org/package=interp.

FIG 1 Extracellular vesicle production rates among marine microbes and as a function of growth

environment. (A) Vesicle production rates for selected strains (indicated here by genus) of

marine microbes at 24 °C. *Prochlorococcus* data represent median values from 5 strains grown at

different light levels (see Fig. 2). (B) *Pelagibacter* (SAR11) vesicle release when grown at 22 °C

on either a defined minimal medium (AMS1) or a natural seawater variant (ProMS*). (C) Impact

of temperature on vesicle production by two heterotrophs. Colors represent taxonomic groupings

of microbes at the Class level: Cyanophyceae (green), Alphaproteobacteria (red),

Gammaproteobacteria (purple), Flavobacteriia (blue).

FIG 2 *Prochlorococcus* vesicle production rates as a function of light and temperature. (A)

Vesicle production rates (black circles) for four *Prochlorococcus* strains grown at different

combinations of light irradiance and temperature. Culture growth rate is indicated by background

color. (B) Production rate data summarized across all growth temperatures measured. Light

levels: LL= low light, ML= medium light, HL= high light, VHL=very high light (see Table 1).

(C) Relative vesicle production rates as a function of relative light level. Vesicle release is

normalized relative to the minimum measured value for each strain; light levels are compared to

the irradiance that resulted in maximal growth rate for each strain (optimal = 1). Dashed lines

represent the changepoint regression fit.

FIG 3 Extracellular vesicle size variation among marine microbes. Vesicle diameters indicate

the mode size value from replicate vesicle populations. Colors represent taxonomic groupings of

microbes at the Class level: Cyanophyceae (green), Alphaproteobacteria (red),

Gammaproteobacteria (purple), Flavobacteriia (blue).

FIG 4 Distribution of vesicle-like particles in the North Pacific Subtropical Gyre. (A)

Concentration of vesicle-like particles (open circles) and cells (filled squares) from Station

ALOHA in June, 2014. Contextual environmental data are shown for (B) photosynthetically

active radiation (PAR) levels and (C) temperature.

TABLE 1 Light conditions examined for each strain. Light treatments are defined as very high

light (VHL), high light (HL), medium light (ML), and low light (LL), corresponding to intensity

values relative to the specific optima for that strain.

Strain	Ecotype/Clade	Light treatment	Light intensity (μmol photons m^{-2} sec^{-1})
MED4	high-light adapted I	VHL	144±4
MED4	high-light adapted I	HL	76±1
MED4	high-light adapted I	ML	40±1
MED4	high-light adapted I	LL	11±1[a]
MIT9312	high-light adapted II	VHL	144±4
MIT9312	high-light adapted II	HL	76±1
MIT9312	high-light adapted II	ML	40±1
MIT9312	high-light adapted II	LL	11±1
NATL2A	low-light adapted I	VHL	96±1
NATL2A	low-light adapted I	HL	45±1
NATL2A	low-light adapted I	ML	20±1
NATL2A	low-light adapted I	LL	10±0.5
MIT9313	low-light adapted IV	HL	45±1
MIT9313	low-light adapted IV	ML	20±1
MIT9313	low-light adapted IV	LL	10±0.5
MIT1223	low-light adapted VIII group	HL	20±1

626
627 [a]This strain was unable to grow at the indicated light level at 24 ˚C, and was instead grown at
628 17±0.5 μmol photons m^{-2} s^{-1}.
629

630   **TABLE 2** *Prochlorococcus* vesicle production rates. All rate values are in units of vesicles cell^{-1}

631   generation^{-1}.

Strain	Ecotype/Clade	Median vesicle production rate across all conditions tested ($\pm$ SD)	Range of vesicle production rate (lowest-highest values in any combination of light/temp)	Significant variation with light (L) or temperature (T)[1]?
MED4	high-light adapted I	1.2 ($\pm$1.5)	0.2 – 3.5	T
MIT9312	high-light adapted II	2.2 ($\pm$9.1)	0.6 – 8.9	-
NATL2A	low-light adapted I	2.5 ($\pm$8.9)	0.4 – 19.0	T, L
MIT9313	low-light adapted IV	3.9 ($\pm$6.2)	0.3 – 15.4	T, L
MIT1223	low-light adapted VIII group	7.7 ($\pm$7.6)	7.7 – 11.1	*nd*[2]

634   [1]Two-way ANOVA, $p < 0.05$.

635   [2]Not determined due to limited number of conditions where this strain grew