Title: Unexpected growth of a classic yeast auxotroph

Authors: S. Branden Van Oss[1,2,7], Saurin Bipin Parikh[1,2,7], Nelson Castilho Coelho[1,2,7], Aaron Wacholder[1,2], Ivan Belashov[3], Manuel Michaca[1,2], Jiazhen Xu[1], Yun Pyo Kang [4,5], Katherine M. McCourt[1,2], Jake McKee[1], Trey Ideker[6], Andrew P. VanDemark[3], Gina M. DeNicola[4], Anne-Ruxandra Carvunis[1,2,*]

[1]Department of Computational and System Biology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, 15213

[2]Pittsburgh Center for Evolutionary Biology and Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, 15213

[3]Department of Biological Sciences, University of Pittsburgh, Dietrich School of Arts & Sciences, Pittsburgh, Pennsylvania, 15213

[4]Department of Cancer Physiology, H. Lee Moffitt Cancer Center, Tampa, Florida, 33612

[5]College of Pharmacy and Research Institute of Pharmaceutical Sciences, Seoul National University, Seoul, Republic of Korea (current address)

[6]Departments of Medicine, Bioengineering, Computer Science and Engineering, Institute for Genomic Medicine, University of California San Diego, La Jolla, California, 92093

[7]Co-first author: These authors contributed equally

*Correspondence: anc201@pitt.edu

Keywords: microbiology, genetics, auxotrophy, methionine, hydrogen sulfide, sulfur, metabolism, conditional essentiality, *MET17*, homocysteine, microbial physiology

Corresponding Author:

Dr. Anne-Ruxandra Carvunis

Department of Computational and Systems Biology

Pittsburgh Center for Evolutionary Biology and Medicine

University of Pittsburgh School of Medicine

3501 Fifth Avenue

Pittsburgh, PA 15260, U.S.A.

Phone: +1 (412) 648-3335

Email: anc201@pitt.edu

ABSTRACT

In budding yeast, the Met15 enzyme has long been assumed to be the sole homocysteine synthase facilitating *de novo* synthesis of sulfur-containing organic compounds (organosulfurs) from inorganic precursors. Here we show that an alternative homocysteine synthase encoded by the previously uncharacterized gene YLL058W supports growth of mutants lacking *MET15* in the absence of exogenous organosulfurs. This growth is observed specifically when cells are deposited in an automated fashion to seed colonies, but not with traditional cell propagation techniques such as thick patches of cells or liquid cultures. We show that the lack of growth in these contexts, which has historically justified the status of *MET15* as a classic auxotrophic marker, is largely due to toxic levels of hydrogen sulfide accumulation rather than an inability to perform *de novo* homocysteine biosynthesis. These data suggest that cell fate decisions in yeast are influenced by hydrogen sulfide levels in the microenvironment. Our results have broad implications for investigations of sulfur metabolism, including studies of methionine restriction and aging.

The metabolism of organic sulfur-containing compounds ("organosulfurs") is critical for

all domains of life. There exists substantial diversity among the components of the

cellular networks that govern this metabolism. This is particularly true of plants and

microbes which, unlike animals, can synthesize organosulfurs from inorganic sulfates

via the sulfate assimilation pathway (SAP) (Thomas and Surdin-Kerjan, 1997). Enzymes

in the SAP reduce inorganic sulfates into sulfides and, in the terminal step, generate

homocysteine from O-acetyl-L-homoserine (OAH) and hydrogen sulfide (H_2S). In

Saccharomyces cerevisiae, Met15 (also referred to as Met17 and Met25) catalyzes this

reaction both *in vivo* (Masselot and De Robichon-Szulmajster, 1975) and *in vitro* (Chen

et al., 2018; Yamagata, 1971). Homocysteine can then be converted either directly to

methionine or indirectly to cysteine. These amino acids may be incorporated into

nascent polypeptides; alternatively, they may be modified to produce other essential

organosulfurs, including the "universal methyl donor" S-adenosylmethionine (SAM) and

glutathione, an important buffer against oxidative stress. Reactions downstream of

homocysteine biosynthesis are reversible; thus all of these organosulfurs may be

recycled back to homocysteine (Thomas and Surdin-Kerjan, 1997).

The first precise and complete deletion of *MET15* (*met15Δ0*) in *S. cerevisiae* was made

in 1998, in the S288C strain background, as part of a collection of "designer deletion"

strains containing one or more deletions of common auxotrophic markers (Brachmann

et al., 1998). Many commonly used laboratory strains contain this deletion, including

BY4741, which serves as the genetic background of the haploid *MAT***a** version of the

yeast deletion collection that has been extensively utilized for functional genomics and

other purposes (Douglas et al., 2012; Giaever and Nislow, 2014). The use of this

deletion as an auxotrophic marker, and the corresponding assumption that *MET15* is

required for growth in organosulfur-free media, has been applied to the interpretation of

a wide range of studies (Johnson and Johnson, 2014; Plummer and Johnson, 2019;

Zou et al., 2017).

Some data, however, suggest that *MET15* may not be strictly required for growth in the

absence of exogenous organosulfurs. *MET15* was first identified in a screen for

methylmercury resistance (Singh and Sherman, 1974). In this study, *met15* mutants

were shown to be auxotrophic for methionine and resistant to methylmercury, but, at a

critical concentration of methylmercury, the methionine requirement appeared to be

alleviated. There are also reports of papillae seen when replica plating thick patches of

met15Δ0 cells to medium lacking organosulfurs (Boeke Lab, 2003).

Here, we demonstrate the existence of an alternative pathway for homocysteine

production in *S. cerevisiae*. We show that *met15Δ* cells grow robustly on media lacking

organosulfurs when propagated by automated colony transfer. This growth requires

exogenous inorganic sulfates and an intact SAP, as well as the uncharacterized gene

YLL058W. We demonstrate that Yll058w catalyzes homocysteine biosynthesis directly

from OAH and H_2S in a purified *in vitro* system. Hence, *met15Δ* cells can synthetize

organosulfurs from inorganic sulfates despite a long-held assumption to the contrary.

The ability of *met15Δ* cells to grow in the absence of exogenous organosulfurs is

context-dependent and highly sensitive to the way the cells are propagated. Specifically,

stable growth is observed with automated colony transfer, but very low or no growth is

observed when using conventional techniques such as thick patches or liquid cultures.

We show that this appearance of auxotrophy is at least partially explained by toxic

levels of accumulated H_2S. Our results have implications for a broad range of studies

that presume organosulfur auxotrophy in *met15Δ* cells, as well as for our understanding

of sulfur metabolism more generally.

RESULTS

met15Δ **cells show unexpected growth in the absence of exogenous**

organosulfurs

We serendipitously observed robust growth of BY4741 when cells were automatedly

transferred in a 384-colony array (**Figure 1A, Table S1**) from a rich medium (YPDA) to

a "synthetic defined" (SD) medium lacking methionine (Met) and cysteine (Cys) and

containing glucose as the carbon source (hereafter referred to as SD-Met+Glu). We first

observed this using a BY4741 control strain from the haploid deletion collection, which

has a *KanMX* cassette that confers resistance to the antibiotic Geneticin (G418). To

confirm that this surprising finding was not unique to our stock of this strain or related to

the presence of the *KanMX* cassette, we purchased BY4741 (lacking the cassette) and

BY4742 (a *MET15$^+$* control) from a commercial vendor (Dharmacon). These BY4741

cells again showed unexpected growth on SD-Met+Glu (**Figure 1B**). Both BY4741 and

BY4742 harbor complete deletions of *URA3* and *LEU2* (Brachmann *et al.*, 1998),

rendering them auxotrophic for uracil (Ura) and leucine (Leu). There was a large,

statistically significant difference in relative colony size between BY4741 colonies that

grew on SD-Met+Glu and the very small colonies detectable on SD-Ura+Glu or SD-

Leu+Glu (**Figure 1C**; effect size = 0.627, p-value = 0.021 [Kruskal-Wallis test] and effect

size = 0.757, p-value = 0.007 [Kruskal-Wallis test], respectively). These experiments

reveal that, in this automated colony transfer procedure, BY4741 does not behave as an

organosulfur auxotroph despite lacking *MET15*.

To confirm this unanticipated observation, we first ensured that our media were devoid

of organosulfur contamination. We analyzed both SD-Met and SD+Met media by mass

spectrometry, specifically looking for methionine, cysteine, homocysteine and SAM. No

contaminants were detected (**Table S2**). We next performed a cross between BY4741

and BY4742 and dissected eight four-spore tetrads. Monogenic phenotypes are

expected to exhibit 2:2 segregation in tetrads. When patched to SD-Met+Glu, we

observed such segregation in all eight tetrads, with two patches growing normally and

two patches displaying extremely weak growth, indicating that our observations were

not likely the result of an unknown secondary mutation (**Figure S1**). The extremely

weak growth of thick patches of cells lacking *MET15* was consistent with auxotrophy, in

contrast to the robust growth observed with the automated colony transfer. Thus, the

growth of cells lacking *MET15* on organosulfur-free media is dependent on how cells

are propagated.

To further investigate the impact of cell propagation technique on growth, we grew

BY4741 and BY4742 in liquid SD-Met+Glu medium (**Figure 1D**). In this context, very

little growth was seen in BY4741 cultures (**Figure 1E**). Drops taken from these cultures

were used to make spots and patches on SD-Met+Glu solid medium (**Figure 1D**). Here

again BY4741 cells displayed an auxotrophic-like phenotype, while BY4742 cells grew

normally (**Figure 1F**). Hence, BY4741 displays auxotroph-like growth when propagated

using traditional techniques.

A

Automated Colony Transfer

B

SD-Met+Glu

BY4742

BY4741

C

Glucose   Galactose   Ethanol

BY4742

BY4741

Relative Colony Size

```
+ Met   - Met   + Met   + Met        - Met   + Met        - Met   + Met
+ Ura   + Ura   - Ura   + Ura        + Ura   - Ura        + Ura   - Ura
+ Leu   + Leu   + Leu   - Leu        + Leu   + Leu        + Leu   + Leu
```

Presumed Auxotrophy   ■ Uracil-Leucine   ■ Methionine-Uracil-Leucine

D

SD-Met+Glu

SD-Met+Glu

E

p = 4.9e-06

OD_{600} at Saturation

BY4742   BY4741

F

SD-Met+Glu

BY4742

BY4741

G

FY4

FY4 − $met15\Delta$

SD-Met+Glu

Colony Size (pixels)

0.65   0.60   0.55   0.46   0.53   0.46   0.49

FY4

FY4 − $met15\Delta$

Time (hours)

Presumed Auxotroph   ■ Yes   ■ No

114
115

Figure 1. The unexpected growth of *met15Δ* cells on organosulfur-free media is both stable and
context-dependent. A) Automated colony transfer technique used in this study. See **Materials and**
Methods and **Table S1**. For some experiments, cells were transferred to the condition plate once; for
others, cells were transferred to the condition media multiple times (light gray plates). **B)** Unexpected
growth of a strain harboring the *met15Δ0* deletion (BY4741) on a medium lacking organosulfurs (SD-
Met+Glu). Cropped images of representative 384-density plates after 366 hours of growth are shown. See
also **Figure S3A**. **C)** Robust growth of *met15Δ0* cells on SD-Met+Glu and SD-Met+Gal, but not on SD-
Met+EtOH. Relative colony size distribution (y-axis) at saturation (~358 to 366 hours) of BY4742 and
BY4741 grown on SD media with glucose, galactose or ethanol, and containing or lacking the indicated
nutrients (x-axis). Relative colony size is normalized to the prototrophic FY4 grown in the same conditions
(**Figure S3B; Materials and Methods**). *: significant Kruskal-Wallis test (p-value < 0.05) comparing the
median relative colony sizes of biological replicates between two strains. **D)** Experimental setup for
traditional cell propagation techniques used in this study (see **Materials and Methods**). **E)** BY4741 cells
fail to grow in liquid SD-Met+Glu medium. Mean OD_{600} after ~51 hours in SD-Met+Glu liquid cultures (y-
axis) over three biological replicates for each strain (x-axis). Error bars: standard error. p: two-tailed
Student's T-test. **F)** BY4741 cells fail to grow on solid SD-Met+Glu medium in large spots or patches.
Cropped representative images of cells grown for 192 hours are shown. **G)** Growth of FY4-*met15Δ* on an
organosulfur-free medium is stable over six consecutive automated colony transfers. Top: experimental
design for comparing the growth of FY4-*met15Δ* (blue) and FY4 (yellow) over consecutive transfers on
SD-Met+Glu. Bottom: colony size (y-axis) over time (x-axis). Lines represent the mean colony size at
each time point. Shaded area: one standard deviation. Numbers next to the blue curves indicate the
fitness of FY4-*met15Δ* relative to FY4 as determined by dividing median colony sizes at the final time
point of each curve.
139

We next interrogated the extent to which the unexpected growth of BY4741 colonies

depends on the specific composition of the growth medium. We repeated the automated

colony transfer procedure using SD-Met and SD-Ura media with either galactose (SD-

Met+Gal and SD-Ura+Gal) or ethanol (SD-Met+EtOH and SD-Ura+EtOH) as the carbon

source (**Figure 1C**). Similar to our observations in SD-Met+Glu, when galactose was

used as a carbon source, BY4741 colonies grew to much larger sizes on SD-Met+Gal

than on SD-Ura+Gal (effect size = 0.561, p-value =0.021 [Kruskal-Wallis test]). In

contrast, the miniscule growth of BY4741 on SD-Met+EtOH was consistent with

auxotrophy. We also repeated the procedure using "synthetic complete" (SC) medium

containing all amino acids except Met and Cys (SC-Met-Cys). We observed robust

growth of BY4741 cells similar to that seen on SD-Met, whether glucose or galactose

was used as a carbon source (**Figure S2**). As was the case with SD media, the growth

of BY4741 colonies on SC-Leu+Glu was substantially weaker than was growth on SC-

Met-Cys+Glu (**Figure S2;** effect size = 0.914, p-value = 0.007 [Kruskal-Wallis test]).

To study the specific consequences of *MET15* loss independent of the other

auxotrophies present in BY4741, we used CRISPR to make a scarless deletion of

MET15 in the prototrophic FY4 background (FY4-*met15Δ*) (see **Materials and**

Methods) and used this strain for all subsequent experiments. Similarly to BY4741,

FY4-*met15Δ* cells grew robustly when automatically transferred on SD-Met with glucose

or galactose, but not ethanol, used as carbon sources (**Figure S3B**). FY4-*met15Δ* also

grew similar to BY4741 on SC-Met-Cys medium whether glucose or galactose was

used (**Figure S2**). Thus, the unexpected growth of cells lacking *MET15* is not

dependent on the genetic background particular to BY4741.

To determine whether the robust growth of *met15Δ* colonies may be explained by

met15Δ recycling and/or scavenging of preexisting organosulfurs, we transferred FY4-

met15Δ from and to SD-Met+Glu six consecutive times. FY4-*met15Δ* cells displayed

robust growth on SD-Met+Glu over repeated transfers (**Figure 1G**). Relative to FY4, the

fitness of these colonies was 0.65 on the initial SD-Met+Glu plates and stabilized at

~0.50. The amount of new biomass generated throughout this experiment demonstrates

that the growth of FY4-*met15Δ* cells cannot be explained by recycling or scavenging.

Despite a longstanding assumption of auxotrophy, *met15Δ* strains can in fact grow

stably in media lacking organosulfurs. This growth is context-dependent, manifesting

when cells are grown by automated colony transfer, but not in thick patches or liquid

cultures, and when cells utilize glucose or galactose, but not ethanol, as a carbon

source.

Stable growth of *met15Δ* cells on organosulfur-free media requires inorganic

sulfates and is not seen in other presumed auxotrophs

To investigate whether the growth of *met15Δ* colonies that we observed in our

automated colony transfer assay was specific to *met15Δ* mutants, we again used

CRISPR to make complete deletions in the FY4 background of nine additional genes

encoding enzymes involved in organosulfur biosynthesis (**Figure 2A)**. These mutants,

along with FY4 and the FY4-*met15Δ* strain, were transferred as described above

(**Figure 1A**) to either YPDA or SD-Met+Glu plates.

All mutant strains displayed growth (or lack thereof) on SD-Met+Glu that was consistent

with current knowledge of organosulfur biosynthesis (**Figure 2B**), with the exception of

FY4-*met3Δ.* In agreement with previous studies (Masselot and De Robichon-

Szulmajster, 1975; Ono et al., 1988), the FY4-*met5Δ,* FY4-*met10Δ,* FY4-*met2Δ*, FY4-

met6Δ, FY4-*met13Δ*, and FY4-*cys4Δ* strains all displayed negligible or no growth on

SD-Met+Glu (effect sizes = 0.887, 0.869, 0.930, 0.904, 0.903, and 0.981, respectively;

p-value < 0.05 [Kruskal-Wallis test] for all comparisons) (**Figure 2B**). The extent to

which *str3Δ* mutants can utilize sulfates has not, to our knowledge, been directly

examined, and loss of *MET12* has not been associated with any *in vivo* phenotypes

(Raymond et al., 1999). The FY4-*str3Δ* strain had a moderate growth defect in SD-

Met+Glu that was also observed in YPDA, while the FY4-*met12Δ* strain grew normally

(**Figure 2B**). Only the FY4-*met3Δ* mutant displayed an unexpected growth phenotype.

Met3 catalyzes the first step in the reduction of intracellular sulfate and is thought to be

essential for growth when unreduced sulfates are the sole sulfur source (Masselot and

De Robichon-Szulmajster, 1975); however, the FY4-*met3Δ* mutant displayed growth on

SD-Met+Glu that was roughly equivalent to that seen with the FY4-*met15Δ* strain

(relative to FY4, effect size = 0.651, p-value < 0.001 [Kruskal-Wallis test] and effect size

= 0.618, p-value < 0.001 [Kruskal-Wallis test], respectively; **Figure 2B**). The growth of

FY4-*met3Δ* was largely unaffected by carbon source, with only a slightly greater growth

defect in SD-Met+Gal and SD-Met+EtOH than in SD-Met+Glu (effect size = 0.289, p-

value = 0.021 [Kruskal-Wallis test] and effect size = 0.298, p-value = 0.021 [Kruskal-

Wallis test] respectively, **Figure S4**).

Figure 2. Growth of *met15Δ* colonies in organosulfur-free media is specific and dependent on the utilization of inorganic sulfates. A) The *S. cerevisiae* organosulfur biosynthesis pathway. Exogenous organosulfurs such as methionine, cysteine, and SAM can be directly imported (gray arrows); inorganic sulfates can also be imported and reduced to sulfides via the SAP. Sulfides are converted into organosulfurs by Met15, which catalyzes *de novo* homocysteine biosynthesis. Homocysteine is then converted to numerous critical downstream organosulfurs, a selection of which are indicated. The *in vivo* biosynthesis of methionine requires both the methionine synthase Met6 and the methylenetetrahydrofolate reductase Met13. The Met13 isozyme, Met12 is shown in gray. Enzymes are indicated within boxes, next to the reactions they catalyze, which are illustrated with arrows that indicate reaction directionality. Drawing inspired by (Walvekar and Laxman, 2019). **B)** *met3Δ* cells also show unexpected growth. Cells grown in each condition as described in **Figure 1A** and **Materials and Methods**. Box plots show the growth of FY4 and its derivative mutant deletion strains, on rich medium (YPDA, left) and on medium lacking organosulfurs (SD-Met+Glu, right). The box plots show relative colony size data (y-axis; see **Materials and Methods**) per strain (subpanels) at saturation (~83.5-89.0 hours in YPDA; 160.0-173.5 hours in SD-Met+Glu). **C-D)** Disruptions of the organosulfur biosynthesis pathway result in a range of H_2S accumulation in cells. Representative crops of 384-density plate images (Panel C) and qualitative assessment, by eye, of relative H_2S (Panel D) in the strains transferred to rich medium containing bismuth for ~160-173.5 hours (BiGGY; see **Table S9**). Error bars: standard error. **E)** Growth of FY4-*met15Δ* colonies requires inorganic sulfates. This experimental procedure is indicated in the schema at the top of the panel. The initial transfer steps were as described in **Figure 1A** The indicated strains were then grown either on SD-Met+Glu medium containing inorganic sulfates (black) or on an analogous medium completely devoid of inorganic sulfates (orange) (see **Materials and Methods** and **Supplementary Material**). After reaching saturation, all strains were transferred to the same condition. Beneath the schema are growth curves (colony size, y-axis) of the indicated strains (subpanels) through the entire experimental procedure (time, x-axis) with the mean colony size depicted. The orange (SD-Met+Glu) and black curves (SD-Met+Glu) in the lower subpanel (FY4-*met3Δ*) overlap.

This surprising growth of *met3Δ* cells led us to examine the level of free sulfides present in our mutant strains. To do so we utilized a colony color assay using bismuth. Bismuth reacts with free sulfides to form a dark precipitate on the surface of cells. (Nickerson, 1953). The darkness of the colonies is therefore a proxy readout for *de novo* homocysteine biosynthesis through the canonical pathway, with strains deficient in this activity expected to accumulate sulfides, as well as for the activity of the SAP, where a broken or inefficient pathway would be expected to result in little to no sulfide accumulation. The strains fell loosely into three groups. The FY4-*met15Δ* and FY4-*met2Δ* mutants both formed very dark colonies when grown on complete media containing bismuth (BiGGY), confirming that they are deficient in homocysteine biosynthesis. The FY4-*cys4Δ* mutant also formed very dark colonies, consistent with the upregulation of the SAP that occurs when cells cannot make cysteine (Hansen and

Johannesen, 2000). The FY4-*met6Δ,* FY4-*met13Δ*, and FY4-*str3Δ* mutants formed an

intermediate amount of precipitate, as did the FY4 control and the FY4-*met12Δ* mutant.

Finally, the FY4-*met3Δ*, FY4-*met5Δ*, and FY4-*met10Δ* mutants had no discernable

precipitate (**Figures 2C-D**), confirming that the SAP is indeed disrupted in these strains

as expected. We saw consistent results in a bismuth-containing medium lacking

organosulfurs (SD-Met+Glu+Bi; **Figure S5**). Altogether, these observations are

consistent with the known roles of the deleted genes in organosulfur biosynthesis

(**Figure 2A**). Therefore, the unexpected growth of FY4-*met15Δ* and FY4-*met3Δ* cells is

not explained by previous mischaracterizations of the positions of the Met15 and Met3

enzymes in the organosulfur biosynthesis pathway.

To determine whether the surprising growth of FY4-*met15Δ* and FY4-*met3Δ* cells in

media lacking organosulfurs depends on the utilization of inorganic sulfates, we

examined how these mutants grown when automatically transferred onto plates entirely

devoid of both organosulfurs and inorganic sulfates. We observed reduced, but not

abolished, growth in SD-Met+Glu without inorganic sulfates for FY4-*met15Δ* and FY4

control colonies. In contrast, the growth of FY4-*met3Δ* colonies was unaffected by the

presence or absence of inorganic sulfates (**Figure 2E**). We then transferred cells from

SD-Met+Glu with or without inorganic sulfates to the same medium a second time.

Upon so doing, the growth of the FY4-*met15Δ* strain on the no sulfate medium was

essentially abolished, indicating that said growth depends on the utilization of inorganic

sulfates. The FY4-*met3Δ* strain failed to show any substantial growth upon a second

transfer, irrespective of the presence or absence of inorganic sulfates. Therefore, unlike

the stable growth seen in FY4-*met15Δ* colonies, the growth of FY4-*met3Δ* colonies

observed after a single transfer to organosulfur-free media is not stable (**Figure 2E**).

Altogether, these results demonstrate that the unexpected stable growth of *met15Δ*

cells is not observed when other key enzymes in the pathway are missing, and requires

inorganic sulfates.

To determine whether the cells that stopped growing when deprived of inorganic

sulfates had arrested or died, we asked whether we could restore growth by transferring

them to SD medium containing both methionine and sulfates. Growth was restored for

both the FY4-*met15Δ* and FY4-*met3Δ* strains that had previously stopped growing

(**Figure 2E**). An increase in colony size was also observed for FY4, which had

surprisingly continued to grow on media lacking both organosulfurs and inorganic

sulfates even after five successive transfers albeit to a significantly reduced colony size

(**Figure 2E**). These data are consistent with studies showing that cells starved for

inorganic sulfates and other natural nutrients tend to arrest rather than die (Boer et al.,

2008).

YLL058W is critical for the growth of organosulfur-starved *met15Δ* cells

Our findings led us to hypothesize that the *S. cerevisiae* genome encoded at least one

enzyme other than Met15 capable of catalyzing *de novo* homocysteine biosynthesis.

We therefore examined the targets of the sole known transcriptional activator of the

sulfur metabolic network, Met4. Of the 45 genes that compose the Met4 "core regulon"

(Lee et al., 2010), six constituted possible candidates: three that are classified by the

Saccharomyces Genome Database (SGD) as "uncharacterized," and three that are

classified as verified genes but have functions described as "unknown" or "putative"

(Stanford University, 2020). Sequence analyses placed two of these candidates –

YHR112C and YLL058W - as a part of the PLP-dependent transferase superfamily,

together with *MET15* and nine other genes primarily involved in sulfur metabolism

(Wilson et al., 2009). Of these two genes, YLL058W was the more compelling candidate

due to the additional fact that its encoded protein is predicted to have the ability to

catalyze homocysteine biosynthesis by a yeast genome-scale metabolic model (Wang

et al., 2021).

YLL058W is located near the telomere of chromosome 12 within a cluster of genes that

are transcriptionally regulated by Met4, several of which have a direct role in sulfur

metabolism (**Figure 3A)** (Lee *et al.*, 2010). There has been speculation that it may have

arisen as a result of horizontal transfer (Lee *et al.*, 2010) - a relatively rare occurrence in

yeast. We thus pursued further investigation of the YLL058W ORF, beginning with its

evolutionary origins. YLL058W has two close paralogs in the *S. cerevisiae* genome:

STR2, which encodes an enzyme that converts cysteine into cystathionine (**Figure 2A**),

and the uncharacterized YML082W, which is a copy of *STR2* generated during the

whole-genome duplication event (Byrne and Wolfe, 2005). We identified bacterial or

fungal homologs of YLL058W, *STR2,* and YML082W using BLASTP and HMMER

(Eddy, 2011) and performed K-means clustering on these sequences. If YLL058W were

derived from a recent horizontal transfer event, we would expect it to be more similar to

some of its homologs outside *Ascomycota* than to its *Ascomycota* homologs. Instead,

all homologs outside *Ascomycota* grouped together along with all budding yeast

homologs more distantly related to *S. cerevisiae* than the *Hanseniaspora* genus. The

homologs found in all descendants of the common ancestor of *Saccharomyces* and

Hanseniaspora comprised two additional groups, one containing YLL058W and the

other containing both *STR2* and YML082W (**Figure S6**). The genes belonging to the

YLL058W class and the *STR2*/YML082W class appear at approximately the same point

in the phylogenetic tree (**Figure 3B**), prior to whole-genome duplication. Therefore, the

most plausible evolutionary scenario is duplication of an ancestral gene into YLL058W

and the *STR2/*YML082W ancestor, followed by substantial divergence of both copies

from the ancestral state, and later by duplication of the *STR2/*YML082W ancestor.

Interestingly, in all species that contain genes of the YLL058W class, the YLL058W

homolog is located near the telomeres among a collection of other sulfur metabolism

related genes (**Figure 3A and Table S3**). The preservation of this pattern in the

genomically-unstable near-telomere region is consistent with a strong selective

constraint keeping YLL058W in proximity to other genes involved in sulfur metabolism.

Figure 3. Yll058w functions as an inefficient homocysteine synthase. A) Genomic context of YLL058W (Stanford University, 2020). In *S. cerevisiae*, YLL058W (magenta) is located near a cluster of genes that are transcriptionally activated by Met4 ((Lee *et al.*, 2010); red boxes), several of which have a direct role in sulfur metabolism (red boxes with black stripes). Black boxes indicate ORFs that fall into neither category. **B)** Distribution of YLL058W homolog clusters across the budding yeast phylogenetic tree. The tree topology follows (Shen et al., 2018). Circles adjacent to each taxon indicate whether YLL058W homologs belonging to each of the three classes identified by K-means clustering in the MDS plot (**Figure S6**; see **Materials and Methods**) are found in the genome of that taxa, while black boxes indicate the presence of a cluster of sulfur-related genes, such as is depicted in Panel A for *S. cerevisiae*. **C)** Yll058w has structural similarity to a homocysteine synthase. Structural overlap of MetY from *Thermatoga maritima* (PDB ID: 7KB1) (gray) with the predicted structure of Yll058w (magenta) and Met15 (violet) calculated using AlphaFold. For clarity, only the active site residues are shown. An O-acetyl-homoserine derivative reaction intermediate captured within the MetY structure (Brewster et al., 2021) is shown in semi-transparent spheres. Residue labeling corresponds to the Yll058w sequence. See also **Figure S7**. **D)** Yll058w is critical for growth in the absence of *MET15*. The indicated strains (x-axis) bearing high-copy (2μ ORI) plasmids conferring resistance to G418, and, where indicated, overexpressing YLL058W, were grown on SD-Met+Gal+G418 medium. The box plots show relative colony size (y-axis) data at saturation (336 hours). Cropped, representative images of the plates corresponding to the strains tested are shown on top. See also **Figure S8**. **E)** Yll058w catalyzes homocysteine biosynthesis. An *in vitro* homocysteine biosynthesis assay was carried out using recombinantly expressed, purified *S. cerevisiae* Met15 (violet) or Yll058w (magenta), or a no-enzyme control (pink), as the catalyst (see **Materials and Methods**). Homocysteine levels (y-axis) were measured at various time points (x-axis). Solid lines: mean. Shaded area: one standard deviation. p: two-tailed, paired Student's T-test paired combining all replicates and time points. Homocysteine levels were significantly higher than the negative control for both Met15 (p=4.46 X10^{-5}) and Yll058w (p=0.00018). The y-axis is shown on a log2 scale.

The sequence similarity and evolutionary relationship of Yll058w to enzymes involved in

sulfur metabolism led us to examine its predicted structure. To do so, we utilized the

crystal structure of MetY (**Figure S7A**), a bacterial *O*-acetylhomoserine

aminocarboxypropyltransferase that catalyzes the homocysteine biosynthesis from *O*-

acetylhomoserine and bisulfide (Brewster *et al.*, 2021), as well as the predicted

structures of Met15 and Yll058w generated by AlphaFold2 (Jumper et al., 2021). The

MetY crystal structure captured a reaction intermediate, allowing for a precise

determination of active site residues. The predicted structures of both Met15 and

Yll058w are highly similar to the experimentally determined structure of MetY within the

active site, including the catalytic lysine (**Figures 3C and S7B**). One difference is

position 280 in the Yll058w sequence, which is a phenylalanine in Yll058w but a

tyrosine in both Met15 and MetY. Another difference in the overall fold is that MetY and

Met15 function as homotetramers (Brewster *et al.*, 2021; Yamagata, 1976), with

multimerization domains allowing the distinct monomers to come together to complete

the active site (**Figure S7C**). In contrast, Yll058w lacks a multimerization domain and is

predicted to be monomeric. However, it contains an approximately 180 amino acid

region before the catalytic domain that has no homology to MetY or Met15. This region

folds back onto the core of the protein, supplying the missing amino acids to the active

site (**Figure S7D**). Altogether, these comparative structural analyses suggest that a

Yll058w monomer may be capable of carrying out catalysis similar to that of the MetY

and Met15 tetramers.

We thus sought to determine whether the putative Yll058w enzyme is necessary and

sufficient to support the growth of *met15Δ* cells in the absence of exogenous

organosulfurs using genetic approaches. We used CRISPR to delete YLL058W in the

FY4 background, and made a *met15Δ yll058wΔ* double mutant in the same

background. We also constructed high copy-number plasmids overexpressing

YLL058W. The FY4-*yll058wΔ* single mutant resulted in only a very modest growth

defect on SD-Met+Glu (effect size = 0.054, p-value > 0.05 [Kruskal-Wallis test]) (**Figure**

S8A), and when grown on bismuth-containing medium displayed an intermediate level

of precipitate (**Figure S8B-C**); its relatively normal growth and intermediate H_2S

accumulation was comparable to FY4 and FY4-*met12Δ* (**Figures 2B-D and S5A-B**).

Strikingly however, the FY4-*met15Δ yll058wΔ* double mutant showed little to no growth

in the absence of exogenous organosulfurs (**Figure 3D**), indicating a synthetic lethal

interaction between *MET15* and YLL058W. To verify that this effect was due to the loss

of the YLL058W gene product, we compared the growth of FY4-*met15Δ* and FY4-

met15Δyll058wΔ strains carrying an empty vector plasmid to the same strains carrying

a plasmid overexpressing YLL058W. Overexpression of YLL058W in the FY4-

met15Δyll058wΔ background resulted in a large, statistically significant increase in

growth (**Figure 3D**; effect size= 9.478, p-value < 0.001 [Kruskal-Wallis test]).

Overexpression of YLL058W also significantly enhanced the growth of the FY4-*met15Δ*

single mutant (effect size= 1.703, p-value < 0.001 [Kruskal-Wallis test]) (**Figure 3D**).

This strongly suggests that the growth of *met15Δ* cells in the absence of exogenous

organosulfurs is mediated by the putative enzyme encoded by YLL058W.

To directly examine the ability of Yll058w to catalyze homocysteine biosynthesis, we

purified recombinant protein from *E. coli*, along with Met15 as a positive control, and

carried out *in vitro* reactions (see **Materials and Methods**). This assay uses the

canonical O-acetyl-homoserine and H_2S substrates and includes in the reaction mixture

pyridoxal 5'-phosphate (PLP), a cofactor critical for Met15 activity. Yll058w was able to

catalyze homocysteine production significantly above background, albeit much less

efficiently than Met15, with the Met15 reactions yielding ~9.9 times as much

homocysteine at the 15-minute time point (**Figure 3E**). Thus, YLL058W does encode an

alternative homocysteine synthase capable of catalyzing the same reaction as Met15.

We sought to investigate whether such inefficient *de novo* homocysteine biosynthesis

would be sufficient to support growth in the absence of Met15 using metabolic

modeling. To do so, we adapted an enzyme-constrained version of Yeast8, the current

consensus genome-scale metabolic model in *S. cerevisiae* (Lu et al., 2019) (see

Materials and Methods). We first removed all of the hypothetical Yll058w-"consuming"

reactions and replaced them with our observed *in vitro* reaction. This had little effect on

the flux of the biomass equation, a proxy for growth in this *in silico* system. We next

removed *MET15* from the model and evaluated the flux through the biomass equation at

a range of arbitrarily assigned catalytic efficiencies (Kcat) for Yll058w. We found that an

appreciable reduction in biomass occurred only when Yll058w was assigned a catalytic

efficiency ~100x lesser than that assigned to Met15 in the model (**Figure S9**). These

simulations, together with our structural, *in vivo*, and biochemical findings, demonstrate

that it is plausible that Yll058w activity enables cells to grow in the absence of *MET15*.

Removal of H_2S enhances the growth of *met15Δ* cells in organosulfur-free media

Our observations (**Figure 1C-F**) show that *met15Δ* cells do indeed display auxotrophic-

like phenotypes when propagated using the type of conventional methods that were

almost exclusively in use at the time of the gene's initial characterization, such as liquid

cultures in flasks or thick patches on Petri dishes, and that these same cells also fail to

grow when ethanol is used as a carbon source. It may be that *met15Δ* cells fail to grow

in these contexts not due to organosulfur starvation *per se*, but to some downstream

cause brought about by the loss of Met15 activity. One such cause may be the large

excess of H_2S present in *met15Δ* cells (**Figures 2C-D and S5A-B**). This accumulation

of H_2S is expected in *met15Δ* cells since the reaction catalyzed by Met15 is one of the

major sulfide-consuming reactions in the cell (Linderholm et al., 2008). Although some

amount of H_2S is required for cell growth, at high levels it is toxic (Blackstone et al.,

2005).

To investigate whether the auxotroph-like behavior of *met15Δ* cells propagated in liquid

cultures or thick patches might be partly due to toxic levels of H_2S, we examined the

growth of FY4 and FY4-*met15Δ* cells in liquid SD-Met+Glu cultures with or without the

H_2S chelator ferric (Fe)-EDTA, which bonds free sulfides that exit the cell by diffusion

(Gaensly et al., 2014; Philip and Brooks, 1974) (**Figure 4A**). In the presence of Fe-

EDTA, we observed a striking increase in the growth of FY4-*met15Δ* cultures (p-value =

0.01 [Student's t-test]; **Figure 4B**, top right panel). To rule out spontaneous mutations

as the reason for this growth increase, we repassaged the cultures in SD-Met+Glu

medium lacking chelator (**Figure 4A**). The repassaged FY4-*met15Δ* cultures showed

little to no growth (**Figure 4B**, bottom right panel). Interestingly, FY4 cultures that had

previously been propagated in the presence of chelator saturated at a higher OD_{600} than

those that had not (p-value = 0.0012 [Student's t-test]; **Figure 4B**, bottom left panel).

We then placed cells from both the original and the repassaged cultures on solid media

plates in large drops and patches (**Figures 4A and 4C**). The FY4-*met15Δ* cells failed to

grow on SD-Met+Glu and SC-Met-Cys+Glu in drops or patches, irrespective of their

previous exposure to the chelator (**Figure 4C**). These results demonstrate that the

growth of FY4-*met15Δ* cultures in the presence of chelator is not facilitated by heritable

mutations.

Drops and patches of FY4-*met15Δ* cells on BiGGY plates retained the expected dark

brown color (**Figure 4C**). This indicates that the cells are still producing an excess of

H_2S in the absence of the chelator, suggesting that the chelator also failed to induce

epigenetic or other heritable changes affecting H_2S levels. However, upon adding Fe-

EDTA to the solid medium, the lack of growth of drops and patches of FY4-*met15Δ* cells

in organosulfur-free media was partially rescued (**Figure 4C**). Thus, the growth defect

(in some settings) and auxotrophic-like behavior (in other settings) of *met15Δ* cells

grown without organosulfurs is at least partially attributable to H_2S toxicity.

Figure 4. Chelation of H₂S facilitates the growth of *met15Δ* cells in the absence of exogenous organosulfurs. A) Experimental setup. Strains were grown in Erlenmeyer flasks in either SD-Met+Glu or SD-Met+Glu+Fe-EDTA liquid media (top two flasks, original cultures). Once saturation was reached for all cultures, those cultures were then repassaged in SD-Met+Glu media (bottom two flasks, repassaged cultures) and again grown to saturation (Panel B). Ten μl from each culture were then spotted and struck to the indicated solid media (Panel C). See also **Figure S11**. **B)** The H₂S chelator Fe-EDTA partially rescues the growth of *met15Δ* cells grown in liquid media. Mean OD_{600} values of the cultures after ~51 hours of growth with or without chelator (top), and ~51 hours after repassage (bottom). Error bars: standard error; p: two-tailed, unpaired Student's t-test. Individual values for each replicate, from both independent experiments, are shown in **Table S4**; outlier FY4-*met15Δ* cultures that showed unexpected growth in SD-Met+Glu were excluded from the data presented here and indicated in green text in the table. **C)** Chelation of H₂S partially rescues the growth of FY4-*met15Δ* cells grown in spots or streaks. Cropped, representative images of cells grown for 192 hours are shown (see **Panel A** and **Materials and Methods**). **D)** Growth defect on media lacking organosulfurs of *met15Δ* petite (ρ^-) derivatives is rescued in presence of H₂S chelator. FY4 and the FY4-*met15Δ* petite (ρ^-) derivatives were grown in similar fashion as the original cultures from Panel A. Data is represented as in Panel B. **E)** A growth defect of *met15Δ* cells grown with ethanol is seen in the presence and absence of organosulfurs in the media. Growth curves of FY4 and the FY4-*met15Δ* mutant (solid line), along with their respective petite (ρ^-) derivatives (dashed line), grown on SD media containing or lacking methionine and utilizing either glucose or ethanol as the sole carbon source are shown. For each condition (subpanel) the cells were grown as described in **Figure 1A and Materials and Methods**. All colony sizes were normalized to the median colony size of FY4 at the final time point in the same condition. Line: mean relative colony size. Shaded regions: one standard deviation. **Table S5** contains the Kruskal-Wallis Test results comparing the growth of a particular strain as estimated by area under the curve in the presence or absence of methionine in media containing either glucose or ethanol as the carbon source.

Excess H₂S is often associated with mitochondrial dysfunction (Paul et al., 2021). Yeast strains with damaged or absent mitochondrial genomes display what is known as the petite phenotype (Ferguson and von Borstel, 1992). To determine whether the partial rescue of *met15Δ* cells by an H₂S chelator is dependent on the mitochondrial genome, we constructed petite (ρ-) derivatives of FY4 and FY4-*met15Δ* (see **Materials and Methods**) and grew them in SD-Met+Glu cultures with or without Fe-EDTA. We observed a strong rescue of FY4-*met15Δ* ρ- growth upon addition of the chelator (p-value = 1.1×10^{-5} [Student's t-test]; **Figure 4D**, bottom panel). We next transferred these strains and their respective non-petite parental strains to solid media containing or lacking methionine, and with either glucose or ethanol as the carbon source (**Figure 4E**). In SD medium containing methionine, non-petite FY4-*met15Δ* cells grew poorly relative to their $MET15^+$ counterparts when ethanol was used as the carbon source. Still,

FY4-*met15Δ* ρ- strains grew on SD-Met+Glu almost as well as their non-petite parental

strains. These results suggest that *met15Δ* cells experience some degree of

mitochondrial dysfunction that impairs their ability to metabolize ethanol, but that neither

their unexpected growth in automated colony transfer settings, nor the chelator-

mediated rescue of their lack of growth in traditional propagation techniques, require a

functional mitochondrial genome.

DISCUSSION

Here we show that, contrary to longstanding assumptions, the organosulfur auxotrophy-like phenotype in a *met15Δ* background is context-dependent. Our analyses demonstrate that the *S. cerevisiae* genome, and possibly those of several closely related fungal species, encode another enzyme, the previously uncharacterized Yll058w, that can catalyze the same reaction as Met15, albeit less efficiently. This activity compensates for the loss of *MET15* and enables the growth of *met15Δ* cells in the absence of exogenous organosulfurs when cells are propagated using automated colony transfer or grown in the presence of an H_2S chelator. Altogether, our observations support a model according to which *met15Δ* cells can synthesize enough homocysteine to support growth through the activity of Yll058w, but experience chronic H_2S toxicity that prevents growth in certain contexts (**Figure 5**). Given its evolutionary conservation, Yll058w may well have other biological activities that are not investigated here.

533
534
Figure 5. H₂S toxicity inhibits growth when Yll058w catalyzes *de novo* homocysteine biosynthesis.
Model representing the hypothesized response to organosulfur starvation in cells containing or lacking
MET15. Extracellular inorganic sulfates are imported through the cell membrane (yellow line) and
reduced to H₂S. When Met15 is present (left), cells efficiently convert H₂S and O-acetylhomoserine (OAH)
into homocysteine, leading to a moderate amount of H₂S (light gray) and an adequate amount of
homocysteine (HC; dark green) to support normal growth. When Met15 is absent, homocysteine
biosynthesis is catalyzed by the comparatively inefficient Yll058w enzyme (right). The resulting lower
amount of homocysteine (light green) is sufficient to support growth in some contexts, but the high
amount of H₂S that accumulates (dark gray), exacerbated by an upregulated intake of sulfates, can
become toxic and lead to cell death or arrest in other contexts.
545

H₂S-mediated context-dependence of a classic auxotrophy

H₂S levels are extremely high when *met15Δ* cells are growing without exogenous

organosulfurs (**Figure S5**), and our results indicate that H₂S toxicity is at least partially

responsible for the arrest and/or death that occurs in *met15Δ* cells grown using

conventional techniques (**Figure 5**). A primary mechanism for the toxic effects of

elevated H₂S, seen in many organisms, involves mitochondrial disfunction (Paul *et al.*,

2021). The failure of *met15Δ* cells to utilize ethanol (**Figures 1C, 4E, and S3B**) is

consistent with mitochondrial dysfunction. However, the ability of the H₂S chelator Fe-

EDTA to restore the growth of petite FY4-*met15Δ* cells grown in liquid cultures (**Figure**

4D), as well as the ability of these cells to grow on solid SD-Met+Glu medium (**Figure**

4E), strongly suggests that H₂S is limiting the growth of *met15Δ* cells in part through a

pathway that does not involve the mitochondrial genome. The precise molecular

mechanisms at play may be dissected in future genome-scale investigations of

suppressor and synthetic lethal mutations, similar to how a recent study dissected the

mechanisms underlying the growth defect associated with petites (Vowinckel et al.,

2021).

We show that *met15Δ* cells behave like organosulfur auxotrophs under conventional

growth techniques, but not when propagated by automated colony transfer. We

hypothesize that robust growth is possible in an automated colony transfer context

because excess sulfides can be eliminated through diffusion between the evenly

spaced small colonies whereas, in the context of thick patches of cells, excess sulfides

are secreted and affect neighboring cells, creating group-level toxicity and self-limiting

growth. Similarly, we hypothesize that H_2S toxicity is high in liquid cultures because the

semi-sealed environment limits gas exchange. While the conditional essentiality of

genes has been well studied in the context of differing genetic backgrounds or

environments (Hou et al., 2018; Zhang and Ren, 2015), to our knowledge, *MET15*

represents the first report of a case in which the essential nature of gene is dependent

on the way cells are propagated.

An expanded yeast sulfur metabolic network

Sulfur-containing amino acids act as cellular sensors for growth control (Laxman et al.,

2013; Sutter et al., 2013). There has therefore been renewed interest in sulfur

metabolism. In particular, methionine restriction is known to increase longevity in a

broad range of organisms (Green et al., 2021), and may serve as a therapeutic

treatment for certain cancers (Endicott et al., 2021). Mutants lacking *MET15* have been

widely used in the study of lifespan extension in yeast (Johnson and Johnson, 2014;

Plummer and Johnson, 2019; Ruckenstuhl et al., 2014; Zou *et al.*, 2017). Our data add

a potential layer of nuance to such studies and hint at the possibility of a role for H_2S in

lifespan extension. More generally, our discovery of the alternative homocysteine

synthase Yll058w may influence the interpretation of the plethora of published studies,

including genome-wide screens conducted in the BY4741 background, wherein the

results or methodologies assumed *met15Δ* cells to be auxotrophic for organosulfurs.

Our findings that *S. cerevisiae* can grow in the absence of exogenous organosulfurs

despite very low levels of *de novo* homocysteine biosynthesis, that the previously

observed auxotroph-like phenotypes are dependent on cell propagation techniques, and

that H_2S levels control group-level growth behaviors, may represent a novel avenue for

studying the interplay between sulfur metabolism and its many dependent phenotypes.

It is of note that even the prototrophic FY4 strain displayed enhanced growth in liquid

media when cells were previously exposed to an H_2S chelator (**Figure 4B**). This

suggests that 1) even wild-type cells suffer some degree of H_2S-induced stress when

grown in organosulfur-free media, and 2) removal of excess H_2S allows the cells to

adapt to this stress, likely at the level of gene expression or epigenetics. This may have

implications for anyone seeking to maximize cellular growth in media lacking, or

containing low amounts of, organosulfurs, whether in an industrial or a research setting.

Also surprising was the limited, but continued, growth of FY4 cells even after multiple

successive transfers to media lacking both organosulfurs and inorganic sulfates (**Figure

2E). This continued growth was not anticipated given that to our knowledge the SAP

relies on inorganic sulfates (**Figure 2A**). Perhaps Met15, but not Yll058w, can

metabolize derivatives of other sulfur-containing compounds present in our growth

media (*e.g.* biotin). Altogether, our work suggests there may yet be more unknown

factors that contribute to the yeast sulfur metabolic network.

MATERIALS AND METHODS

Yeast strains

The well-characterized S288C-derived *Saccharomyces cerevisiae* strains FY4, BY4741 and BY4742 were used throughout this work (Brachmann *et al.*, 1998; Winston et al., 1995).

<u>CRISPR/Cas9 deletion strains</u>

All other strains (**Table S6**) were derived by CRISPR/Cas9 based scarless whole-ORF deletions of genes from the prototrophic strain, FY4. Co-transformations were made with 5μg of repair fragment and 1μg of plasmid expressing cas9 (Addgene 60847, **Table S7**) (Ryan et al., 2014) with the respective sgRNA (DiCarlo et al., 2013). Identification of PAM sites and sgRNA selection were chosen using Benchling (https://www.benchling.com/) and the sgRNA and repair fragment sequences are given in **Table S8**. Transformed cells were plated on YPDA+G418 and colonies were PCR genotyped to select those with the correct gene deletion. The positive clones were then grown on YPDA to induce the loss of the cas9- and sgRNA-containing plasmids. Strains carrying plasmids for overexpression experiments were made with the LiAc/PEG/ssDNA protocol (Dunham, 2015) using 500ng of each plasmid and selected on YPDA+G418.

<u>Petite strains</u>

Petite derivatives of FY4 and FY4-*met15Δ* were made using a slightly modified version of the standard LiAc/PEG/ssDNA yeast transformation protocol (Dunham, 2015), with

shorter incubation times and without a DNA template, followed by replica platting on

YPDA and YP-Glycerol plates to identify the petites.

<u>Strain stock plates</u>

Stock plates were constructed for each strain by selecting four independent colonies

from streakouts, which were used to inoculate liquid YPDA medium and grown

overnight at 30°C. These cultures were considered biological replicates. The cultures

were then used to create 384-well glycerol stocks (glycerol 25% v/v) for each strain,

with each biological replicate occupying one-fourth of the plate, distributed uniformly.

<u>Plasmid construction</u>

The overexpression plasmids used in **Figure 3D** were made by replacing the Leu2 gene

by KanMx cassette into the destination plasmid pAG425GPD-ccdB (Alberti, 2007). This

plasmid was used as an empty vector plasmid (pARC0112, **Table S7**) while an LR

recombination (Gateway LR Clonase II Enzyme Mix, ThermoFisher) of this same

plasmid with an entry clone containing the ORF YLL058W was used to produce the

plasmid overexpressing YLL058W (pARC0172, **Table S7**).

Plasmids pARC0190 and pARC0191 (**Table S7**) were made by amplification of *MET15*

and YLL058W coding sequences, respectively, from FY4 genomic DNA, and cloned into

a pET28a-derived vector expressing an N-terminal, cleavable 10XHis-Ruby tag (Kredel

at.al., 2009).

Growth Media

The recipes for all the various growth media used in this study are given in **Table S9**.

When required, the analogous liquid medium was made with the same composition,

excluding the agar.

Strain Crossing

The cross depicted in **Figure S1** was carried out by standard techniques. Briefly, the

BY4741 and BY4742 deletion collection parent strains (**Table S6**) were mated on YPDA

solid medium and incubated at room temperature (RT) for ~6 hours, and then struck to

select for colonies formed from single diploids on solid SD+His+Leu+Ura medium

(Table S9). After incubation at 30°C for ~2 days on the diploid selection medium,

colonies were patched to "GNA" plates (**Table S9**) and incubated at 30°C for ~2 days.

These plates were then replica plated to "sporulation (SPO)" medium (**Table S9)**

containing histidine, lysine, methionine, uracil, and leucine, and incubated at RT for ~11

days. Four-spore tetrads were selected and dissected using standard techniques and

initially grown on YPDA. Individual spores were then patched to petri dishes as shown

in **Figure S1**.

Automated Colony Transfer

Using the benchtop RoToR HDA robotic plate handler (Singer Instruments Co Ltd,

Roadwater, UK) strains were transferred from glycerol stocks to solid rich medium

plates (YPDA, **Table S9**) as described on **Figure 1A** and incubated at 30°C for 72

hours (see **Table S1** for a description of the settings used). Each of these plates were

677 copied to two YPDA plates, which represented the two technical replicates of plates per

678 strain, and again incubated at 30°C for 48 hours. This step reduces the variability in

679 colony size that is introduced when cells are transferred from glycerol stocks. Finally,

680 these plates were 'one-to-many' copied onto various solid 'condition' media plates (**see**

681 **Table S9)** depending on the experiment. The plates were incubated at 30°C and

682 periodically imaged until the colonies reached saturation. This experimental protocol,

683 depicted in **Figure 1A**, was carried out at least twice per experiment to control for batch

684 effect. **Figures 1C, 1G, 2B, 2E, 3D and 4E** (indicated with a dull yellow background) as

685 well as **Figures S2, S3B, S4 and S8A** were carried out in this manner.

687 **Colony size estimation**

688 Serial imaging of the plates was done either using spImager Automated Imaging

689 System (S & P Robotics Inc., Ontario, Canada) or manually using a custom-made

690 lightbox with an overhead camera mount using a commercially available SLR camera

691 (18Mpixel Rebel T6, Canon USA Inc., Melville, NY, USA). The plates were imaged at

692 regular intervals beginning right after transfer until the colonies reached saturation. The

693 images were analyzed in bulk using a custom script made using functions from the

694 MATLAB Colony Analyzer Toolkit (Bean et al., 2014) to provide colony size estimations

695 (https://github.com/sauriiiin/lid_personal/blob/master/justanalyze.m). The output files

696 containing colony size information along with the images is available at

697 https://bit.ly/3pOe6aT.

Relative fitness measurement

A spatially cognizant colony size database was built using the colony size estimates per

the LI Detector framework for high-throughput colony-based screening analysis (Parikh

et al., 2021). For all experiments, first, the outermost border colonies were removed

from the dataset because these usually tend to grow more due to excess access to

nutrients, leaving a maximum of 77 colony size observations per strain per time point

per plate per biological replicate. Outlier colony size values per experiment per technical

replicate per time point per plate per biological replicate were removed using two

median adjusted deviations as a cutoff. These raw colony size values are reported in

Figures 1G and **2E**.

For **Figures 1C**, **2B**, **3D, 4E, S2, S3B, S4,** and **S8A,** the median colony size per time

point of the prototrophic strain, FY4, was used to calculate relative colony size of every

colony per strain per time point. At saturation, the relative colony size distribution per

strain is determined by the median relative colony sizes per plate per biological

replicate. This relative colony size distribution is compared amongst strains using the

non-parametric Kruskal-Wallis Test to determine significant changes in fitness. Effect

size values are calculated as the difference in median colony sizes divided by the

median colony size of the strain being used as reference in the comparison.

Additionally for **Figure 4E and Table S5**, growth curve analysis was done with the

growthcurver (Sprouffske and Wagner, 2016) package in R (Team, 2021), using the

time series colony size data per colony per strain per technical replicate per biological

replicate. This time series data was first converted into an equally spaced time series

using a second degree locally weighed scatterplot smoothing (loess) function to

generate 100 equally spaced data points. This was then fed into the

'*SummarizeGrowthByPlate()*' function of the gro*wthcurver* package (Sprouffske and

Wagner, 2016) to make estimates of the area under the curve (AUC) per colony per

strain in the experiment. AUC summarizes a growth curve by integrating the

contributions of the initial population size, growth rate, and carrying capacity into a

single value, making it a more robust estimation of growth-curve-based 'fitness.'

Relative area under the curve, effect sizes and significant changes were estimated the

same way as the colony size data described above. The AUCs from the data in **Figure**

4E were used to calculate the statistics in **Table S5**.

Relative H$_2$S measurement

Bismuth containing plates (BiGGY and SD-Met+Glu+Bi; **Table S9**) were used to assess

the level of excess sulfides in various strains. Cells were transferred from YPDA plates

onto bismuth containing plates, as described on **Figure 1A**. The qualitative assessment

involved two replicates scored by eye in duplicate by independent observers on a scale

of 1-7, with the darkest strain(s) assigned a value of seven and the lightest strains

assigned a value of one.

Mass spectrometry analysis of media by semi-targeted high-resolution LC-HRMS

<u>**Sample preparation**</u>

Metabolic quenching and polar metabolite pool extraction was performed by adding

400µL ice cold methanol to 100µL of sample. Deuterated (D$_3$)-creatinine and (D$_3$)-

alanine, (D_4)-taurine and (D_3)-lactate (Sigma-Aldrich) was added to the sample lysates

as an internal standard for a final concentration of 10µM. Samples were vortexed, and

then homogenized using a 25°C water bath sonicator for 5 minutes. The supernatant

was then cleared of protein by centrifugation at 16,000xg. Two µL of cleared

supernatant was subjected to online LC-MS analysis.

LC-HRMS Method

Analyses were performed by untargeted LC-HRMS. Briefly, Samples were injected via a

Thermo Vanquish UHPLC and separated over a reversed phase Thermo HyperCarb

porous graphite column (2.1×100mm, 3µm particle size) maintained at 55°C. For the 20

minute LC gradient, the mobile phase consisted of the following: solvent A (water /

0.1% formic acid) and solvent B (ACN / 0.1% formic acid). The gradient was the

following: 0-1min 1% B, increase to 15%B over 5 minutes, continue increasing to 98%B

over 5 minutes, hold at 98%B for five minutes, reequillibrate at 1%B for five minutes.

The Thermo IDX tribrid mass spectrometer was operated in positive ion mode, scanning

in ddMS^2 mode (2 µscans) from 70 to 800 m/z at 120,000 resolution with an AGC target

of 2e5 for full scan, 2e4 for ms^2 scans using HCD fragmentation at stepped 15,35,50

collision energies. Source ionization setting was 3.0kV spray voltage for positive mode.

Source gas parameters were 35 sheath gas, 12 auxiliary gas at 320°C, and 8 sweep

gas. Calibration was performed prior to analysis using the Pierce^{TM} FlexMix Ion

Calibration Solutions (Thermo Fisher Scientific). Integrated peak areas were then

extracted manually using Quan Browser (Thermo Fisher Xcalibur ver. 2.7)

(Metabolomics & Lipidomics Core, University of Pittsburgh Health Sciences Core

Research Facilities).

Evolutionary origins of YLL058W

Genome sequences of 332 budding yeasts were taken from a published dataset (Shen

et al., 2018). Each genome was scanned for ORFs (consisting of the sequence between

an ATG start codon and a stop codon in the same frame) larger than 300 bp. We then

used BLASTP on the predicted protein products of these ORFs to identify the strongest

match among *S. cerevisiae* annotated genes. BLASTP was run with default settings and

a 10^{-4} e-value cutoff. All ORFs for which the best match (i.e., the lowest BLASTP e-

value) was either the *S. cerevisiae* YLL058W sequence or the sequence of one of its

two *S. cerevisiae* homologs (*STR2* and YML082W) were retained. For each ortholog

identified, we also noted the best *S. cerevisiae* matches for the 5 nearest ORFs in each

direction that had a match. To supplement this list with orthologs from more distant

species, we submitted the *S. cerevisiae* YLL058W sequence to the HMMER homology

search tool (Eddy, 2011). From these results, we selected the strongest matching

sequences (lowest e-values) from a variety of bacterial and fungal taxa outside of the

Ascomycota phylum (four bacterial and nine fungal outgroups). The full list of YLL058W

homologs and their sequences is given in **Supplementary File 1**.

We next constructed a multiple alignment of all homologs using the MUSCLE aligner

(Madeira et al., 2019). We used the R package bio2mds to perform K-means clustering

on the sequence alignment and to visualize the results using multidimensional scaling.

To investigate the phylogenetic distribution of each cluster within budding yeasts, we

used the phylogeny generated by the same study of budding yeasts that formed the

foundation of our sequence dataset (Shen *et al.*, 2018).

Purification of recombinantly expressed proteins

Both Met15 or Yll058w were expressed as fusion proteins with an N-terminal His_{10}-

mRuby2 tag which can be removed by cleavage with TEV protease. Fusion proteins

were expressed in the BL21 (DE3) RIPL CodonPlus *E. coli* strain (Stratagene). Cells

were grown at room temperature in 2L of standard LB media containing kanamycin and

chloramphenicol to an OD600 of ~0.6, induced with 0.5mM IPTG, and grown for eight

additional hours at room temperature post-induction. Purification of both proteins was

performed essentially as described previously (Van Oss et al., 2016). Briefly, both

proteins were purified by two rounds (pre- and post-cleavage with TEV protease) of

nickel affinity chromatography followed by ion exchange chromatography. Met15 was

further purified by size exclusion chromatography. The final purified fractions that were

used in the *in vitro* homocysteine assay are shown in **Figure S10**.

in vitro **homocysteine assay**

in vitro homocysteine biosynthesis was carried out largely as described previously

(Chen *et al.*, 2018). Aliquots of Met15 and Yll058w, purified as described above, were

dialyzed into a final storage buffer of 20 mM Tris-Cl pH 8.0, 500 mM NaCl, 20 mM

Imidazole, 0.2 mM pyridoxal 5'-phosphate (PLP), and 10% glycerol, snap frozen in

liquid N_2, and stored at -80°C. Aliquots were thawed on ice, and the enzymes were

diluted in the storage buffer to a concentration of 60 nM and preincubated at room

temperature for 10 minutes. The enzyme solution was then mixed with diluted 10x,

giving a final enzyme concentration of 6nM, with a reaction mixture containing 50 mM

Potassium phosphate buffer (pH 7.8), 1.0 mM EDTA pH 8.0, 15.0 mM O-acetyl-L-

homoserine, 5.0 mM NaHS, and 0.6 mM PLP. The reaction components were then

mixed by gentle pipetting and incubated at 30°C. Each reaction was performed in

triplicate. Samples were removed at 6 timepoints (0, 1, 3, 5, 10, and 15 minutes), and

the reactions were stopped by adding 1.0 M HCl at a ratio of 10mL for every 100mL of

sample, then snap frozen with liquid N_2. In the same buffer as the reaction mixture but

without any enzyme or substrate (but with acid added), we also generated a

homocysteine standard curve with seven serial 1:5 dilutions ranging from 15 mM to

0.00096 mM, as well as a 0 mM "blank" sample.

Ten µL of each enzyme reaction were extracted and derivatized with N-ethylmaleamide

(NEM; Alfa Aesar, Cat# 40526-06) in 990 µL of ice-cold extraction solvent (80% MeOH:

20% H_2O containing 25 mM NEM and 10 mM ammonium formate, pH=7.0) containing

0.5 uM of [D_4]-Homocysteine (Cambridge Isotope labs, Cat# DLM-8259-PK) followed by

incubation at 4°C for 30 min. After centrifugation (17,000 g, 20 min, 4°C), the

supernatants were analyzed by LC-MS following previously established conditions

(Kang et al., 2019). For the chromatographic metabolite separation, a Vanquish UPLC

system was coupled to a Q Exactive HF (QE-HF) mass spectrometer equipped with

HESI (Thermo Fisher Scientific, Waltham, MA). The column was a SeQuant ZIC-pHILIC

LC column, 5 mm, 150 x 4.6mm (MilliporeSigma, Burlington, MA) with a SeQuant ZIC-

pHILIC guard column, 20 x 4.6 mm (MilliporeSigma, Burlington, MA). Mobile phase A

was 10 mM $(NH_4)_2CO_3$ and 0.05% NH_4OH in H_2O while mobile phase B was 100%

ACN. The column chamber temperature was set to 30°C. The mobile phase condition

was set according to the following gradient: 0-13min: 80% to 20% of mobile phase B,

13-15min: 20% of mobile phase B. The ESI ionization mode was positive. The MS scan

range (m/z) was set to 60-900. The mass resolution was 120,000 and the AGC target

was 3×10^6. Capillary voltage and capillary temperature were set to 3.5 KV and 320°C,

respectively. Five μL of sample was loaded. The homocysteine and $[D_4]$-Homocysteine

peaks were manually identified and integrated with EL-Maven (Version 0.11.0) by

matching with an in-house library. Homocysteine levels were calculated using the

standard curve and corrected using the $[D_4]$-Homocysteine internal standard.

Metabolic Modeling

Metabolic modeling simulations were performed with Python script making use of the

COBRApy toolkit (Ebrahim et al., 2013) and the ecYeastGEM_batch model (Lu *et al.*,

2019), an "enzyme-constrained" variant (Sanchez et al., 2017) of Yeast8. The enzyme-

constrained feature of the steady-state model is achieved by treating enzymes as

metabolites that are temporarily "consumed" (i.e. removed from the available pool) while

they are engaged in catalysis. The script first removes all preexisting reactions (which

have not been experimentally validated) contributing to the "consumption" of Yll058w

(r_0815No1, r_0815REVNo1, and arm_r_0815). Second, the script adds the

hypothetical homocysteine-generating reaction that we observe in our *in vitro* assay,

which uses H_2S and *O*-acetylhomoserine as substrates and yields homocysteine,

acetate, and hydrogen as products, with Yll058w as the catalyst. This reaction was

added with a range of Kcat values assigned to Yll058w in order to evaluate the

"biomass" generated over a broad range of enzyme efficiencies. The reaction catalyzed

by Yll058w was made to be bidirectional; however, flux was only observed in the

forward (homocysteine-producing) reaction. Thirdly, *MET15* was removed from the

model, eliminating all reactions that the encoded enzyme was associated with. Finally,

the model was "solved" to obtain the flux through the biomass equation, a proxy for

fitness. The default growth medium associated with the model, which contains sulfates

but lacks organosulfurs, was used for all simulations.

Chelation of H_2S with Ferric-EDTA sodium salt

For each strain (FY4, FY4-*met15Δ*, and their respective petite derivatives), five single

colonies were scraped from a YPDA agar plate and cells were resuspended in SD-

Met+Glu medium. The optical density at 600 nm (OD_{600}) of the suspension was

measured using a spectrophotometer, and 25mL cultures were inoculated at a starting

OD_{600} of 0.1 in SD-Met+Glu or SD-Met+Glu supplemented with 0.048M Ferric (Fe)-

EDTA (Ethylenediaminetetraacetic acid ferric sodium salt, E6067, Millipore Sigma) in

125mL Erlenmeyer flasks (optimal Fe-EDTA dosage was determined by a titration

experiment (**Figure S11**)). For the non-petite strains, the experiment was performed

twice; each time, cultures were prepared in triplicate. The petite strains were tested in a

single experiment using four biological replicates. Cultures were incubated at 30^{o}C with

150rpm shaking and OD_{600} was measured regularly until saturation was reached (~51 to

53 hours). For the repassaging of the non-petite strains (**Figure 4A-C**), cells were then

pelleted to facilitate the exchange to fresh SD-Met+Glu medium, again using a starting

OD_{600} of 0.1, and incubated as before for ~51 hours with regular OD_{600} measurements.

A portion of each culture was stored at 4°C. Ten μl drops of each culture (both the

original culture and after the repassaged growth) were then both spotted and struck on

agar plates (YPDA, SD-Met+Glu, SC-Met-Cys+Glu, SC-Met-Cys+Glu+(Fe-EDTA) and

BiGGY) that were incubated at 30°C (**Figure 4C**) and imaged regularly with spImager

Automated Imaging System (S & P Robotics Inc., Ontario, Canada). All experiments

carried out in liquid media are indicated with a gray background.

Data availability

All data generated/analyzed in this study are available in the main text, in the

Supplementary Figures and Tables, and as Supplementary Data files. Supplementary

material is available at https://bit.ly/3GIgpld.

Code availability

All images and statistical analyses in this study were generated using code available at

https://github.com/sauriiiin/methionine/tree/master/paper.

ACKNOWLEDGEMENTS

We thank all the members of the Carvunis Lab, and in particular Carly Houghton, for

helpful suggestions during lab meetings and presentations. We also thank Kate

Metedgul-Ernar, Cameron Hines and Brian Hsu for their precious help in the early

stages of the project and S.G. Wendell and Steven Mullett for their assistance with

mass spectrometry analysis.

FUNDING

This work was supported by the National Institutes of Health grant R00GM108865 to A.-

R.C, F32GM129929 to S.B.V.O, P01CA250984 to G.M.D, NIHS10OD023402 to S.G.W.,

and the AACR-Takeda Oncology Lung Cancer Research Fellowship (19-40-38-KANG)

to Y.P.K.

AUTHOR CONTRIBUTIONS

Conceptualization: S.B.V.O., S.B.P., N.C.C., G.M.D., A-R.C. Methodology: S.B.V.O.,

S.B.P., N.C.C., A.W., A.P.V., G.M.D., A-R.C. Investigation: S.B.V.O., S.B.P., N.C.C.,

A.W., I.B., M.M., J.X., Y.P.K., K.M.M., J.M., A.P.V. Writing: S.B.V.O., S.B.P., N.C.C.,

A.W., A-R.C. Review and Editing: S.B.V.O., S.B.P., N.C.C., T.I., A.P.V., G.M.D., A-R.C.

Supervision: A-R.C.

DECLARATION OF INTERESTS

A.-R. C. is a member of the scientific advisory board for Flagship Labs 69, Inc. T.I. is a

co-founder of Data4Cure and has an equity interest. T.I. is on the Scientific Advisory

925   Board of Ideaya BioSciences, Inc., has an equity interest, and receives income. The

926   terms of these arrangements have been reviewed and approved by the University of

927   California San Diego in accordance with its conflict of interest policies.

REFERENCES

Bean, G.J., Jaeger, P.A., Bahr, S., and Ideker, T. (2014). Development of ultra-high-density screening tools for microbial "omics". PLoS One *9*, e85177. 10.1371/journal.pone.0085177.

Blackstone, E., Morrison, M., and Roth, M.B. (2005). H2S induces a suspended animation-like state in mice. Science *308*, 518. 10.1126/science.1108581.

Boeke Lab (2003). The Met15 Resource: Update. http://www.bs.jhmi.edu/mbg/boekelab/Resources/Met15/Met15updt.html.

Boer, V.M., Amini, S., and Botstein, D. (2008). Influence of genotype and nutrition on survival and metabolism of starving yeast. Proc Natl Acad Sci U S A *105*, 6930-6935. 10.1073/pnas.0802601105.

Brachmann, C.B., Davies, A., Cost, G.J., Caputo, E., Li, J., Hieter, P., and Boeke, J.D. (1998). Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. Yeast *14*, 115-132. 10.1002/(SICI)1097-0061(19980130)14:2<115::AID-YEA204>3.0.CO;2-2.

Brewster, J.L., Pachl, P., McKellar, J.L.O., Selmer, M., Squire, C.J., and Patrick, W.M. (2021). Structures and kinetics of Thermotoga maritima MetY reveal new insights into the predominant sulfurylation enzyme of bacterial methionine biosynthesis. J Biol Chem *296*, 100797. 10.1016/j.jbc.2021.100797.

Byrne, K.P., and Wolfe, K.H. (2005). The Yeast Gene Order Browser: combining

curated homology and syntenic context reveals gene fate in polyploid species. Genome

Res *15*, 1456-1461. 10.1101/gr.3672305.

Chen, Z., Zhang, X., Li, H., Liu, H., Xia, Y., and Xun, L. (2018). The Complete Pathway

for Thiosulfate Utilization in Saccharomyces cerevisiae. Appl Environ Microbiol *84*.

10.1128/AEM.01241-18.

DiCarlo, J.E., Norville, J.E., Mali, P., Rios, X., Aach, J., and Church, G.M. (2013).

Genome engineering in Saccharomyces cerevisiae using CRISPR-Cas systems.

Nucleic Acids Res *41*, 4336-4343. 10.1093/nar/gkt135.

Douglas, A.C., Smith, A.M., Sharifpoor, S., Yan, Z., Durbic, T., Heisler, L.E., Lee, A.Y.,

Ryan, O., Gottert, H., Surendra, A., et al. (2012). Functional analysis with a barcoder

yeast gene overexpression system. G3 (Bethesda) *2*, 1279-1289.

10.1534/g3.112.003400.

Dunham, M.J., Gartenberg, M. R., Brown, G. W. (2015). Methods in Yeast Genetics and

Genomics, 2015 Edition: A CSHL Course Manual (Cold Spring Harbor Laboratory

Press).

Ebrahim, A., Lerman, J.A., Palsson, B.O., and Hyduke, D.R. (2013). COBRApy:

COnstraints-Based Reconstruction and Analysis for Python. BMC Syst Biol *7*, 74.

10.1186/1752-0509-7-74.

Eddy, S.R. (2011). Accelerated Profile HMM Searches. PLoS Comput Biol *7*, e1002195.

10.1371/journal.pcbi.1002195.

Endicott, M., Jones, M., and Hull, J. (2021). Amino acid metabolism as a therapeutic

target in cancer: a review. Amino Acids *53*, 1169-1179. 10.1007/s00726-021-03052-1.

Ferguson, L.R., and von Borstel, R.C. (1992). Induction of the cytoplasmic 'petite'

mutation by chemical and physical agents in Saccharomyces cerevisiae. Mutat Res

265, 103-148. 10.1016/0027-5107(92)90042-z.

Gaensly, F., Picheth, G., Brand, D., and Bonfim, T.M. (2014). The uptake of different

iron salts by the yeast Saccharomyces cerevisiae. Braz J Microbiol *45*, 491-494.

10.1590/s1517-83822014000200016.

Giaever, G., and Nislow, C. (2014). The yeast deletion collection: a decade of functional

genomics. Genetics *197*, 451-465. 10.1534/genetics.114.161620.

Green, C.L., Lamming, D.W., and Fontana, L. (2021). Molecular mechanisms of dietary

restriction promoting health and longevity. Nat Rev Mol Cell Biol. 10.1038/s41580-021-

00411-4.

Hansen, J., and Johannesen, P.F. (2000). Cysteine is essential for transcriptional

regulation of the sulfur assimilation genes in Saccharomyces cerevisiae. Mol Gen Genet

263, 535-542. 10.1007/s004380051199.

Hou, J., van Leeuwen, J., Andrews, B.J., and Boone, C. (2018). Genetic Network

Complexity Shapes Background-Dependent Phenotypic Expression. Trends Genet *34*,

578-586. 10.1016/j.tig.2018.05.006.

Johnson, J.E., and Johnson, F.B. (2014). Methionine restriction activates the retrograde

response and confers both stress tolerance and lifespan extension to yeast, mouse and

human cells. PLoS One *9*, e97729. 10.1371/journal.pone.0097729.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate

protein structure prediction with AlphaFold. Nature *596*, 583-589. 10.1038/s41586-021-

03819-2.

Kang, Y.P., Torrente, L., Falzone, A., Elkins, C.M., Liu, M., Asara, J.M., Dibble, C.C.,

and DeNicola, G.M. (2019). Cysteine dioxygenase 1 is a metabolic liability for non-small

cell lung cancer. Elife *8*, e45572.

Laxman, S., Sutter, B.M., Wu, X., Kumar, S., Guo, X., Trudgian, D.C., Mirzaei, H., and

Tu, B.P. (2013). Sulfur amino acids regulate translational capacity and metabolic

homeostasis through modulation of tRNA thiolation. Cell *154*, 416-429.

10.1016/j.cell.2013.06.043.

Lee, T.A., Jorgensen, P., Bognar, A.L., Peyraud, C., Thomas, D., and Tyers, M. (2010).

Dissection of combinatorial control by the Met4 transcriptional complex. Mol Biol Cell

21, 456-469. 10.1091/mbc.E09-05-0420.

Linderholm, A.L., Findleton, C.L., Kumar, G., Hong, Y., and Bisson, L.F. (2008).

Identification of genes affecting hydrogen sulfide formation in Saccharomyces

cerevisiae. Appl Environ Microbiol *74*, 1418-1427. 10.1128/AEM.01758-07.

Lu, H., Li, F., Sanchez, B.J., Zhu, Z., Li, G., Domenzain, I., Marcisauskas, S., Anton,

P.M., Lappa, D., Lieven, C., et al. (2019). A consensus S. cerevisiae metabolic model

Yeast8 and its ecosystem for comprehensively probing cellular metabolism. Nat

Commun *10*, 3586. 10.1038/s41467-019-11581-3.

Madeira, F., Park, Y.M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P.,

Tivey, A.R.N., Potter, S.C., Finn, R.D., and Lopez, R. (2019). The EMBL-EBI search

and sequence analysis tools APIs in 2019. Nucleic Acids Res *47*, W636-W641.

10.1093/nar/gkz268.

Masselot, M., and De Robichon-Szulmajster, H. (1975). Methionine biosynthesis in

Saccharomyces cerevisiae. I. Genetical analysis of auxotrophic mutants. Mol Gen

Genet *139*, 121-132. 10.1007/BF00264692.

Nickerson, W.J. (1953). Reduction of inorganic substances by yeasts. I. Extracellular

reduction of sulfite by species of Candida. J Infect Dis *93*, 43-56. 10.1093/infdis/93.1.43.

Ono, B., Shirahige, Y., Nanjoh, A., Andou, N., Ohue, H., and Ishino-Arao, Y. (1988).

Cysteine biosynthesis in Saccharomyces cerevisiae: mutation that confers cystathionine

beta-synthase deficiency. J Bacteriol *170*, 5883-5889. 10.1128/jb.170.12.5883-

5889.1988.

Parikh, S.B., Castilho Coelho, N., and Carvunis, A.R. (2021). LI Detector: a framework

for sensitive colony-based screens regardless of the distribution of fitness effects. G3

(Bethesda) *11*. 10.1093/g3journal/jkaa068.

Paul, B.D., Snyder, S.H., and Kashfi, K. (2021). Effects of hydrogen sulfide on

mitochondrial function and cellular bioenergetics. Redox Biol *38*, 101772.

10.1016/j.redox.2020.101772.

Philip, C.V., and Brooks, D.W. (1974). Iron(III) chelate complexes of hydrogen sulfide

and mercaptans in aqueous solution. Inorganic Chemistry *13*, 384-386.

10.1021/ic50132a030.

Plummer, J.D., and Johnson, J.E. (2019). Extension of Cellular Lifespan by Methionine

Restriction Involves Alterations in Central Carbon Metabolism and Is Mitophagy-

Dependent. Front Cell Dev Biol *7*, 301. 10.3389/fcell.2019.00301.

Raymond, R.K., Kastanos, E.K., and Appling, D.R. (1999). Saccharomyces cerevisiae

expresses two genes encoding isozymes of methylenetetrahydrofolate reductase. Arch

Biochem Biophys *372*, 300-308. 10.1006/abbi.1999.1498.

Ruckenstuhl, C., Netzberger, C., Entfellner, I., Carmona-Gutierrez, D., Kickenweiz, T.,

Stekovic, S., Gleixner, C., Schmid, C., Klug, L., Sorgo, A.G., et al. (2014). Lifespan

extension by methionine restriction requires autophagy-dependent vacuolar

acidification. PLoS Genet *10*, e1004347. 10.1371/journal.pgen.1004347.

Ryan, O.W., Skerker, J.M., Maurer, M.J., Li, X., Tsai, J.C., Poddar, S., Lee, M.E.,

DeLoache, W., Dueber, J.E., Arkin, A.P., and Cate, J.H. (2014). Selection of

chromosomal DNA libraries using a multiplex CRISPR system. Elife *3*.

10.7554/eLife.03703.

Sanchez, B.J., Zhang, C., Nilsson, A., Lahtvee, P.J., Kerkhoven, E.J., and Nielsen, J.

(2017). Improving the phenotype predictions of a yeast genome-scale metabolic model

by incorporating enzymatic constraints. Mol Syst Biol *13*, 935. 10.15252/msb.20167411.

Shen, X.X., Opulente, D.A., Kominek, J., Zhou, X., Steenwyk, J.L., Buh, K.V., Haase,

M.A.B., Wisecaver, J.H., Wang, M., Doering, D.T., et al. (2018). Tempo and Mode of

Genome Evolution in the Budding Yeast Subphylum. Cell *175*, 1533-1545 e1520.

10.1016/j.cell.2018.10.023.

Singh, A., and Sherman, F. (1974). Association of methionine requirement with methyl

mercury resistant mutants of yeast. Nature *247*, 227-229. 10.1038/247227a0.

Sprouffske, K., and Wagner, A. (2016). Growthcurver: an R package for obtaining

interpretable metrics from microbial growth curves. BMC Bioinformatics *17*, 172.

10.1186/s12859-016-1016-7.

Stanford University (2020). Saccharomyces Genome Database.

https://www.yeastgenome.org/.

Sutter, B.M., Wu, X., Laxman, S., and Tu, B.P. (2013). Methionine inhibits autophagy

and promotes growth by inducing the SAM-responsive methylation of PP2A. Cell *154*,

403-415. 10.1016/j.cell.2013.06.041.

Team, R.C. (2021). R: A language and environment for statistical computing. (R

Foundation for Statistical Computing, Vienna, Austria.).

Thomas, D., and Surdin-Kerjan, Y. (1997). Metabolism of sulfur amino acids in

Saccharomyces cerevisiae. Microbiol Mol Biol Rev *61*, 503-532.

Van Oss, S.B., Shirra, M.K., Bataille, A.R., Wier, A.D., Yen, K., Vinayachandran, V.,

Byeon, I.L., Cucinotta, C.E., Heroux, A., Jeon, J., et al. (2016). The Histone Modification

Domain of Paf1 Complex Subunit Rtf1 Directly Stimulates H2B Ubiquitylation through

an Interaction with Rad6. Mol Cell *64*, 815-825. 10.1016/j.molcel.2016.10.008.

Vowinckel, J., Hartl, J., Marx, H., Kerick, M., Runggatscher, K., Keller, M.A., Mulleder,

M., Day, J., Weber, M., Rinnerthaler, M., et al. (2021). The metabolic growth limitations

of petite cells lacking the mitochondrial genome. Nat Metab *3*, 1521-1535.

10.1038/s42255-021-00477-6.

Walvekar, A.S., and Laxman, S. (2019). Methionine at the Heart of Anabolism and

Signaling: Perspectives From Budding Yeast. Front Microbiol *10*, 2624.

10.3389/fmicb.2019.02624.

Wang, H., Robinson, J.L., Kocabas, P., Gustafsson, J., Anton, M., Cholley, P.E., Huang,

S., Gobom, J., Svensson, T., Uhlen, M., et al. (2021). Genome-scale metabolic network

reconstruction of model animals as a platform for translational research. Proc Natl Acad

Sci U S A *118*. 10.1073/pnas.2102344118.

Wilson, D., Pethica, R., Zhou, Y., Talbot, C., Vogel, C., Madera, M., Chothia, C., and

Gough, J. (2009). SUPERFAMILY--sophisticated comparative genomics, data mining,

visualization and phylogeny. Nucleic Acids Res *37*, D380-386. 10.1093/nar/gkn762.

Winston, F., Dollard, C., and Ricupero-Hovasse, S.L. (1995). Construction of a set of

convenient Saccharomyces cerevisiae strains that are isogenic to S288C. Yeast *11*, 53-

55. 10.1002/yea.320110107.

Yamagata, S. (1971). Homocysteine synthesis in yeast. Partial purification and

properties of O-acetylhomoserine sulfhydrylase. J Biochem *70*, 1035-1045.

10.1093/oxfordjournals.jbchem.a129712.

Yamagata, S. (1976). O-Acetylserine and O-acetylhomoserine sulfhydrylase of yeast.

Subunit structure. J Biochem *80*, 787-797. 10.1093/oxfordjournals.jbchem.a131339.

Zhang, Z., and Ren, Q. (2015). Why are essential genes essential? - The essentiality of

Saccharomyces genes. Microb Cell *2*, 280-287. 10.15698/mic2015.08.218.

Zou, K., Ouyang, Q., Li, H., and Zheng, J. (2017). A global characterization of the

translational and transcriptional programs induced by methionine restriction through

ribosome profiling and RNA-seq. BMC Genomics *18*, 189. 10.1186/s12864-017-3483-2.

1101

1 **SUPPLEMENTARY MATERIAL**
2
3
4 **SUPPLEMENTARY FIGURES**
5
6

Spore 1 **Spore 2** **Spore 3** **Spore 4**

Tetrads

BY4741 X BY4742
SD-Met+Glu

7
8
Figure S1. *met15Δ*-associated growth phenotype segregates 2:2 in a heterozygous cross. The
BY4741 and BY4742 strains were mated, sporulated, and dissected using standard techniques. Eight
four-spore tetrads were grown on YPDA, and then patched onto SD-Met+Glu medium. Individual plates
were used to prevent cross-feeding between *MET15*[+] and *met15Δ* cells. Each tetrad resulted in 2:2
segregation of the growth phenotype on organosulfur-deficient media. Notably, the growth of *met15Δ*
thick patches was substantially weaker than that observed in the automated colony transfer assays used
throughout the manuscript.

17
18
Figure S2. *met15Δ* **cells also grow robustly in "synthetic complete" (SC) media lacking**
organosulfurs. The same automated colony transfer and analysis pipeline shown in **Figure 1A-C** was
repeated, here on SC-Met-Cys media containing either glucose or galactose as the sole carbon source,
or on SC-Leu+Glu medium, where the negligible growth of the BY4741 and BY4742 strains is consistent
with auxotrophy. The prototrophic strain FY4 and a mutant derivative lacking *MET15* were also assayed.
Box plots represent relative colony sizes (y-axis) at saturation (~140 hours (SC-Leu+Glu) or 165 hours
(SC-Met+Glu/Gal)) of BY4742, BY4741, FY4 and FY4-*met15Δ* (x-axis). Relative colony size is normalized
to the mean colony size of FY4 grown in the same condition (see **Materials and Methods**).

A   Synthetic Defined Media

B

27
28
Figure S3. Robust growth of *met15Δ* cells in an organosulfur-deficient medium. **A)** In "synthetic
defined" medium lacking methionine and with glucose comprising the carbon source (SD-Met+Glu), cells
harboring the *met15Δ0* deletion (BY4741) or a "scarless deletion" of *MET15* made by CRISPR (FY4-
met15Δ) show surprisingly robust growth inconsistent with their presumed auxotrophy, while in
"conventional" SD medium (SD+Met+Glu), these mutant cells form slightly larger colonies than their
nearly isogenic *MET15*[+] counterparts, BY4741 and FY4, respectively. Colonies shown here have grown
for 366 hours. Related to **Figure 1B**. **B)** The "scarless" CRISPR *met15Δ* mutant shows similarly robust
growth, inconsistent with auxotrophy, when glucose or galactose comprise the carbon source, but growth
is much weaker when the same strain is grown with ethanol as the carbon source. Box plots represent
relative colony sizes (y-axis) at saturation (~358 to 366 hours). Relative colony size is normalized to the
mean colony size of FY4 grown in the same condition (see **Materials and Methods**).

40
41
42 **Figure S4. The growth of FY4-*met3Δ* cells on SD-Met media is generally unaffected by the carbon**
43 **source.** The FY4-*met3Δ* mutant was grown in the same manner as the strains presented in **Figures 1C**
44 and **S3A-B**. Box plots represent relative colony sizes (y-axis) at saturation (366 hours). Relative colony
45 size is normalized to the mean colony size of FY4 grown in the same condition (see **Materials and**
46 **Methods**).

SD-Met+Glu+Bi

47
48
49 **Figure S5. H₂S levels of mutant strains in an organosulfur-free medium.** Cropped images (**Panel A**)
50 and qualitative assessment of colony color (**Panel B**, performed in the same manner described in **Figure**
51 **2D**) of the mutant strains tested in **Figures 2C-D and S8B-C** pinned to SD-Met+Glu medium containing
52 bismuth (SD-Met+Glu+Bi). The FY4-*cys4Δ* mutant showed no observable growth hence has been
53 excluded from Panel B. The darkness of the colonies is a proxy for the level of excess H₂S in each strain.

54
55
56 **Figure S6. YLL058W homologs cluster into three distinct classes. Related to Figure 3B.** The
57 sequence distances between YLL058W homologs found in budding yeasts and outgroup taxa are
58 visualized using multidimensional scaling (MDS). Points are colored according to k-means clustering
59 based on these sequence distances using three clusters. The three genes contained in the *S. cerevisiae*
60 genome are labeled.

61
62
Figure S7. Yll058w functions as a monomer, in contrast to the tetrameric conformation seen in
canonical homocysteine synthases. Related to Figure 3C. A) A cartoon view of the MetY monomer,
from the MetY crystal structure, pdbID:7KB1 (Brewster et al., 2021). The N-terminal tetramerization
domain is shown in orange, the PLP-binding domain is shown in light purple, and the C-terminal substrate
binding domain is shown in pale green. **B)** The predicted structure of the monomeric Yll058w (white) is
highly similar structurally to MetY within the PLP-binding and C-terminal domains and aligns well with a
MetY dimer (reaction intermediate colored red for clarity). The Yll058w unique domain (Panel D), which
replaces the tetramerization domain found in MetY, mimics part of the neigboring subunit from the MetY
dimer, effectively preventing Yll058w from multimerizing in a manner similar to MetY. **C)** Cartoon view of
the MetY N-terminal tetramerization domains (Tet_A and Tet_B; orange) and the C-terminal substrate
binding domains (CBD_A, CBD_B; pale green) in contact with the reaction intermediate (red). The PLP-
binding domains are hidden. Met15 (not shown) is predicted to adopt a similar conformation. **D)** The
unique portion of the Yll058w monomer supplies the additional catalytic residues (pale green) provided by
the second subunit of the MetY dimer. The portion of the predicted structure of Yll058w in contact with the
reaction intermediate (red) is shown, with areas showing sequence and predicted structural similarity
shown in white, and the region lacking homology to MetY and Met15 (Yll058w_unique) shown in gray.
This unique region supports the possibility of Yll058w catalysis as a monomer, as opposed the confirmed
(MetY) and predicted (Met15) multimeric structures of other homocysteine synthases. Portions of the
PLP-binding domain that contribute putative catalytic residues are shown in black.

82
83
Figure S8. Deletion of YLL058W has negligible effect on fitness and H₂S accumulation in the FY4
background. Related to Figure 3D. A) Deletion of YLL058W in the prototrophic FY4 background has a
negligible effect on growth in SD-Met+Glu medium. The strains were examined in the same manner as
the mutants shown in **Figure 2B**, with colony size normalized to that of FY4**. B and C)** The relative H₂S
level of FY4-*yll058wΔ*, as determined by growth on BiGGY and SD-Met+Glu+Bi, is comparable to that of
wild-type FY4 cells. For Panel B, the darkness of the colonies was scored on the same qualitative scale
applied to the mutants in **Figures 2C-D and S5B**.

91
92
Figure S9. Simulated growth utilizing the consensus, genome-scale *S. cerevisiae* metabolic model
(Lu et al., 2019) shows that a low-efficiency homocysteine synthase can support growth. Growth in
an organosulfur-free medium is predicted *in silico* even with a homocysteine synthase with a dramatically
lower catalytic efficiency than Met15. Two modifications were made to the default model – the addition of
the *in vitro* reaction carried out in **Figure 3E** utilizing Yll058w as the catalytic enzyme (black), and the
subsequent removal of the *MET15* gene from the model (gray). The y-axis indicates the flux through the
"biomass equation", a proxy for growth/fitness, and the x-axis indicates different catalytic efficiency (Kcat)
values assigned to Yll058w, relative to the default Kcat assigned to Met15, with the efficiency of Yll058w
in the hypothetical reaction decreasing from left to right.

102
103
104 **Figure S10. Coomassie-stained SDS-PAGE gel showing the final fractions that contained the**
105 **recombinantly-expressed, purified protein used in the *in vitro* homocysteine biosynthesis assay.**
106 A lane containing molecular weight standards (kDa = kilodaltons) is shown on the left. Lanes containing
107 less pure fractions that were not used were cropped, but the image is otherwise unaltered.

108
109
110 **Figure S11. Titration of the H$_2$S chelator Fe-EDTA. Related to Figure 4.** To determine the optimal
111 dosage of Fe-EDTA for the experiments depicted in **Figures 4B-D**, FY4 and FY4-met15Δ cultures were
112 grown in SD+Met+Glu and SD-Met+Glu supplemented with Fe-EDTA, with concentrations raging from 0
113 to 0.136 M. The concentration of 0.048 M yielded the greatest restoration of growth in FY4-met15Δ
114 cultures. Bar plot: mean OD$_{600}$. Error bars: standard error.
115

SUPPLEMENTARY TABLES

Table S1. Settings for the RoToR HDA robotic plate handler used for all of the automated pinning experiments described in the manuscript. Screening of the deletion collection involved all five unique settings, while the remaining pinning experiments all followed the general schema described in **Figure 1B** and used only the "384 Glycerol to 384 Plate" and "384 Plate to 384 Plate" settings.

SETTINGS				384 Glycerol to 384 Plate	384 Plate to 384 Plate
Source				384 MTP	384 Agar
Target				384 Agar	384 Agar
Pads				384 Long Pin	384 Short
Program				Spot Many	Replicate
	General				
		Quantity		2	N/A
		Recycle		None	None
		Plate Protection		Uncheck	Uncheck
		Pairs			
		Start Pair			
	Source				
		Offset			Automatic
			Max Radius		0.4 mm
		Offsets…			
		Pinning			
			Pin Pressure		32%
			Speed	19 mm per sec	19 mm per sec
			Backoff	0.5 mm	2 mm
			Repeat Pin	1 times	1 times
		Dry Mix			
			Clearance		0.5 mm
			Diameter		1 mm
			Cycles		1 rotations
		Wet Mix			
			Diameter	1 mm	
			Speed	25 mm per sec	
			Cycles	3 rotations	
			Travel	3	
			2D/3D	3D	
		Permanent Offset			
	Target				
		Pinning			
			Pin Pressure	32%	32%
			Speed	19 mm per sec	19 mm per sec
			Backoff		
			Overshoot	2 mm	2 mm
			Repeat Pin	1 times	1 times
		Dry Mix			
			Clearance	0.5 mm	0.5 mm
			Diameter	1 mm	1 mm
			Cycles	1 rotations	1 rotations
		Wet Mix			
			Diameter		
			Speed		
			Cycles		
			Travel		
			2D/3D		
	Pads				
		Pressure			
		Ejject Time			
	Selected Options				
		Recycle		Off	Off
		Revist		Off	Off
		Protect Source		Off	Off
		Offset			Auto
		Source Mix		On	On
		Target Mix		On	Off

126 **Table S2.** Mass spectrometry analysis confirms that the growth medium is not contaminated with
127 organosulfurs. ND= Not Detected. See **Materials and Methods**.
128

	Cysteine	Methionine	Homocysteine	SAM
Yeast Nitrogen Base	ND	ND	ND	ND
2% Glucose	ND	ND	ND	ND
Leucine	ND	ND	ND	ND
Histidine	ND	ND	ND	ND
Uracil	ND	ND	ND	ND
Lysine	ND	ND	ND	ND
Glutamic Acid	ND	ND	ND	ND
SD-Met	ND	ND	ND	ND
SD+Met	ND	7.15 mM	ND	ND
ddH$_2$O	ND	ND	ND	ND
Pure H$_2$O	ND	ND	ND	ND

130 **Table S3.** *YLL058W* and its orthologs are found in a cluster of genes near the telomere. *S. cerevisiae*
131 gene names are used for all species shown. The ten nearest genes (5 upstream (5') genes, numbered -1
132 to -5, and 5 downstream (3') genes, numbered 1 to 5) to *YLL058W* orthologs are shown. Asterisks
133 indicate a known role in sulfur metabolism.
134

		Position Relative to YLL058W										
		-5	-4	-3	-2	-1	YLL058W	1	2	3	4	5
Species	Saccharomyces cerevisiae	PAU18	AYT1	MHT1*	MMP1*	GTT2*	YLL058W*	JLP1*	YLL056C	YCT1*	YLL054C	YLL053C
	Zygosaccharomyces bisporus	RRP42	TRM3	ATG20	YDL114W	ARG8	YLL058W*	YLL058W*	YCT1*	AYR1	DAL5	PUT3
	Torulaspora delbrueckii	RPT6	ALG23	DSE4	MHT1*	MMP1*	YLL058W*	YCT1*	DLD2	GRX1*	DIP5	YRF1-1
	Torulaspora pretoriensis	THI73	AMD2	SOA1*	JLP1*	PHO84	YLL058W*	MMP1*	MHT1*	DSE4	ALG23	RPT6
	Lachancea quebecensis	YCT1*	SOA1*	ZRT1	PHO11	ZPS1	YLL058W*	VEL1	FLO5	COS9	AIF1	YER152C

Table S4. OD$_{600}$ values at saturation (~51 hours) for cultures of the indicated strains grown in SD-
Met+Glu medium with or without the H$_2$S chelator Ferric(Fe)-EDTA (column 4), and OD$_{600}$ values for the
same cultures diluted and repassaged in SD-Met-Glu medium lacking chelator (column 5). Each strain
has 3 replicates (1-3) from two independent experiments (A&B). Although FY4-*met15Δ* cultures generally
failed to show any substantial growth in liquid medium without an H$_2$S chelator present (**Figure 4B**), we
occasionally observed "escape" in four FY4-*met15Δ* cultures (indicated in green text) grown in SD-
Met+Glu without an H$_2$S chelator present, independent of whether the cells had previously been in the
presence of chelator.
144

Strain	Replicate	Initial Medium	OD600	OD600 (Repassage)
FY4	1A	SD-Met+Glu	4.98	6.13
FY4	1B	SD-Met+Glu	5.45	4.75
FY4	2A	SD-Met+Glu	4.9	5.88
FY4	2B	SD-Met+Glu	5.71	4.64
FY4	3A	SD-Met+Glu	5.13	6.45
FY4	3B	SD-Met+Glu	5.76	5.8
FY4	1A	SD-Met+Glu+Fe-EDTA	5.86	7.71
FY4	1B	SD-Met+Glu+Fe-EDTA	6.76	7.03
FY4	2A	SD-Met+Glu+Fe-EDTA	5.5	7.82
FY4	2B	SD-Met+Glu+Fe-EDTA	6.24	6.78
FY4	3A	SD-Met+Glu+Fe-EDTA	5.26	7.87
FY4	3B	SD-Met+Glu+Fe-EDTA	6.15	6.61
FY4-*met15Δ*	1A	SD-Met+Glu	1.39	0.48
FY4-*met15Δ*	1B	SD-Met+Glu	0.21	0.09
FY4-*met15Δ*	2A	SD-Met+Glu	0.29	5.93
FY4-*met15Δ*	2B	SD-Met+Glu	0.22	0.07
FY4-*met15Δ*	3A	SD-Met+Glu	0.34	0.14
FY4-*met15Δ*	3B	SD-Met+Glu	0.21	0.07
FY4-*met15Δ*	1A	SD-Met+Glu+Fe-EDTA	2.55	0.39
FY4-*met15Δ*	1B	SD-Met+Glu+Fe-EDTA	1.02	3.29
FY4-*met15Δ*	2A	SD-Met+Glu+Fe-EDTA	2.23	0.26
FY4-*met15Δ*	2B	SD-Met+Glu+Fe-EDTA	0.56	0.08
FY4-*met15Δ*	3A	SD-Met+Glu+Fe-EDTA	2.52	4.22
FY4-*met15Δ*	3B	SD-Met+Glu+Fe-EDTA	1.12	3.32

146 **Table S5.** Kruskal-Wallis Test results comparing the growth of a given strain in condition_1 (+Met) vs.
147 condition_2 (-Met), with EtOH or Glucose. The statistic and p-value columns are the Kruskal-Wallis test
148 statistic and p-value output, and the effect size column is the difference between the median fitness value
149 in media lacking methionine and that in media containing methionine divided by the median fitness value
150 in media containing methionine.
151

Strain	condition_1 (+Met)	condition_2 (-Met)	statistic	p-value	effect size
FY4	SD+Met+EtOH	SD-Met+EtOH	0.0300	0.8625	0.0000
FY4-*met15Δ*	SD+Met+EtOH	SD-Met+EtOH	17.2800	0.0000	-0.8926
FY4-*met15Δ* (ρ−)	SD+Met+EtOH	SD-Met+EtOH	12.5952	0.0004	-0.4500
FY4 (ρ−)	SD+Met+EtOH	SD-Met+EtOH	0.4821	0.4875	0.1471
FY4	SD+Met+Glu	SD-Met+Glu	0.3333	0.5637	0.0000
FY4-*met15Δ*	SD+Met+Glu	SD-Met+Glu	17.2800	0.0000	-0.6168
FY4-*met15Δ* (ρ−)	SD+Met+Glu	SD-Met+Glu	17.2800	0.0000	-0.4874
FY4 (ρ−)	SD+Met+Glu	SD-Met+Glu	3.8533	0.0496	-0.0683

153 **Table S6.** *Saccharomyces cerevisiae* strains used in this study (related to **Materials and Methods**).
154

Strain #	Strain name	Genotype	Used in Figure(s)	Source
yARC0052	BY4741	*MAT**a** his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*	1B, 1C, 1E, 1F, S2, S3A, S4A	Winzeler et al., 1999 (acquired from Horizon Discovery)
yARC0053	BY4742	*MAT**α** his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	1B, 1C, 1E, 1F, S2, S3A, S4A	Winzeler et al., 1999 (acquired from Horizon Discovery)
yARC0055	BY4741 *HOΔ0 ::KanMX*	*MAT**a** his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 HOΔ0::KanMX*	S1	Giaever et al., 2002
yARC0056	BY4742 *HOΔ0 ::KanMX*	*MAT**α** his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 HO::KanMX*	S1	Giaever et al., 2002
yARC0083	FY4	*MAT**a***	1G, 2B, 2C, 2D, 2E, 4B, 4C, 4E, S2, S3A, S3B, S4, S5A, S5B, S8, S10	Winston et al., 1995
yARC0293	FY4-*cys4Δ0*	*MAT**a** cys4Δ0*	2B, 2C, 2D, S5A	This Study
yARC0224	FY4-*met2Δ0*	*MAT**a** met2Δ0*	2B, 2C, 2D, S5A, S5B	This Study
yARC0312	FY4-*met3Δ0*	*MAT**a** met3Δ0*	2B, 2C, 2D, 2E, S4, S5A, S5B	This Study
yARC0458	FY4-*met5Δ0*	*MAT**a** met5Δ0*	2B, 2C, 2D, S5A, S5B	This Study
yARC0300	FY4-*met6Δ0*	*MAT**a** met6Δ0*	2B, 2C, 2D, S5A, S5B	This Study
yARC0459	FY4-*met10Δ0*	*MAT**a** met10Δ0*	2B, 2C, 2D, S5A, S5B	This Study
yARC0222	FY4-*met12Δ0*	*MAT**a** met12Δ0*	2B, 2C, 2D, S5A, S5B	This Study
yARC0310	FY4-*met13Δ0*	*MAT**a** met13Δ0*	2B, 2C, 2D, S5A, S5B	This Study
yARC0178	FY4-*met15Δ0*	*MAT**a** met15Δ0*	1G, 2B, 2C, 2D, 2E, 4B, 4C, 4E, S2, S3A, S3B, S5A, S5B, S10	This Study
yARC0295	FY4-*str3Δ0*	*MAT**a** str3Δ0*	2B, 2C, 2D, S5A, S5B	This Study
yARC0220	FY4-*yll058wΔ0*	*MAT**a** yll058wΔ*	S8	This Study
yARC0460	FY4	*MAT**a** pGDP-ccdB::KanMx*	3D	This Study
yARC0461	FY4-*met15Δ0*	*MAT**a** met15Δ0 pGDP-ccdB::KanMx*	3D	This Study
yARC0462	FY4-*yll058wΔ0*	*MAT**a** yll058wΔ pGDP-ccdB::KanMx*	3D	This Study
yARC0463	FY4- *met15Δ0 yll058wΔ0*	*MAT**a** met15Δ0 yll058wΔ pGDP-ccdB::KanMx*	3D	This Study
yARC0464	FY4-*met15Δ0*	*MAT**a** met15Δ0 pGDP-yll058w::KanMx*	3D	This Study
yARC0465	FY4- *met15Δ0 yll058wΔ0*	*MAT**a** met15Δ0 yll058wΔ pGDP-yll058w::KanMx*	3D	This Study
yARC0207	FY4 (ρ-)	*MAT**a** (ρ-)*	4D, 4E	This Study
yARC0208	FY4-met15Δ (ρ-)	*MAT**a** met15Δ0 (ρ-)*	4D, 4E	This Study

155
156

157 **Table S7.** Plasmids used in this study (related to **Materials and Methods**).
158

Plasmid #	Construction	Yeast Origin	Promoter	Gene Product(s)	Markers	Used in Figure(s)
pARC0028	See (Ryan et al., 2014)	2μ		Cas9+sgRNA	KanR	
pARC0112	LEU2 was replaced by *KanMX* on plasmid pAG425GPD-ccdB from Yeast Gateway Kit (Alberti et al., 2007)	2μ	GPD	ccdB (empty plasmid)	G418R (*KanMX*)	3D
pARC0172	Made by LR recombination between destination plasmid pARC0112 and entry clone containing the Yll058w gene cds	2μ	GPD	Yll058w	G418R (KanMX)	3D
pARC0190	The coding sequence for *MET15* was amplified from *S. cerevisiae* genomic DNA and cloned into a pET28a-derived vector expressing an N-terminal, cleavable 10XHis-Ruby (Kredel et al., 2009) tag.		T7	10XHis-Ruby-TEV-Met15	KanR	S11
pARC0191	The coding sequence for YLL058W was amplified from *S. cerevisiae* genomic DNA and cloned into a pET28a-derived vector expressing an N-terminal, cleavable 10XHis-Ruby (Kredel et al., 2009) tag.		T7	10XHis-Ruby-TEV-Yll058w	KanR	S11

160   **Table S8.** sgRNA, repair fragment template (HR_template) and amplification primer (HR_Fw and HR_rv)
161   sequences used for CRISPR/Cas9-mediated ORF deletions (related to **Materials and Methods**).
162

ORF name	Primers	Sequence	Used to construct strain #
CYS4	sgRNA	GTTGTAGGCCACTTGCTCAA	yARC0293
	HR_template	CATCTAGATAAATACGACGTAAGAATAAAAATAAGAACCCACGCTTCAAATAAAAGCAAA	
	HR_Fw	AGTAAAAGGCAACACTTGAAGATTTCGTTGTAGGCCACTTGCTCAAACGACATCTAGATA	
	HR_Rv	AGAGAACGGTGCAATTGAATAGGAAAGGAATGACGGATTTTGCTTCTATGTTTGCTTTTA	
MET2	sgRNA	ATCGAAAACGCTCCAAGAGC	yARC0224
	HR_template	AAAGAAAGAAAAAAACGTAGTATAAAAGGAGGATAGATACCACACATACCTCAGGCATAA	
	HR_Fw	TTTTTAGTCACAGGGTCCGTCTAAAGTTTCTCTTTATTTGGAATAATAGAAAAGAAAGAA	
	HR_Rv	GAAAGGACCTCAGTGATGTCATCTACTCTGTCTCTTGCCAAATCGTGCGTATCCAGTTTA	
MET3	sgRNA	AACATGCCTGCTCCTCACGG	yARC0312
	HR_template	TAATAATAACAAGAATAAAGTATAATTAACTCGTCATAAAATGCTCCCATCTCAAAAGTA	
	HR_Fw	ACTTATTTGCAAATCTATTTATTTTGCGCGGTCGATCATGAATTTTGCCCTACTTTTGAG	
	HR_Rv	ACTTATTTGCAAATCTATTTATTTTGCGCGGTCGATCATGAATTTTGCCCTACTTTTGAG	
MET5	sgRNA	ACAATTGTGGGAGCGTCAAG	yARC0458
	HR_template	TAGTATGTCCTACTATGTCATATGCTATCATCGCCAATCTTTTGTTTTTCTCTGGTTCCC	
	HR_Fw	AGTTATATTCAATTGCGAAAAAAATAAATAAATGTGGGAAGAAAACCCAATAGTATGTCC	
	HR_Rv	TTGATATTTATTCAGATCCCTTATATACTAATAAAGTAACAGTAGGGAACGGGAACCAGA	
MET6	sgRNA	TTCAAAGTACATATCAAAAA	yARC0300
	HR_template	ACCAATATAATTTCAAAGTACATATCAAAAGGTTTTAAAAAGGAAGCAAAGTAATGATAT	
	HR_Fw	ATAAAAAAGCAAGCATCTAAGAGCATTGACAACACTCTAAGAAACAAAATACCAATATAA	
	HR_Rv	ATTAAATCACCGATGTTGAATCCCAGAATAAAAAACAAAAAGTTCAGAAAATATCATTAC	
MET10	sgRNA	GGATTGGTAGCAAACTCAAC	yARC0459
	HR_template	TAGATATTTAGTTTTTATTACTATATTAATCTCGTTGTATATTTGGGTGACCTCGAGGAA	
	HR_Fw	AAAACTAAAAGTAAAAAAAAAAATAGAAATGTTCAAGACAGGTTCAATAAATAGATATTTA	
	HR_Rv	TGGAAATAGTTTCGTTTAGATGGGGTGCACATAGTTTTTTTTCAACTGCTTTTCCTCGAGG	
MET12	sgRNA	TCAGAGATTTATATCATGCG	yARC0222
	HR_template	GACGGGACAGGTTGATTACATTTTTTAAACTATTTCACTGAATATAAATACAAAAATGCG	
	HR_Fw	CTCACATCATCTATTTTGTTTCAAGTTCGTACATTTTTTGAAGCGTGTTGGACGGGACAG	
	HR_Rv	TTAGAAGGGCGTTCTGCCCGTTATGTATACGTTAAATATTACATTATTTTCGCATTTTTG	
MET13	sgRNA	GAGCAACATAGACAGACCTC	yARC0310
	HR_template	CCACAGTTACTACTACAACCACATCGCAATTAACACAGACAACGATAACAGTTCTTTAAC	
	HR_Fw	CCTTGATTGAAATAGTCTCCCTAAACTAAAGTTATCAGCAAACAGAACCACCACAGTTAC	
	HR_Rv	AATGCAATGTAATGTTGAATGTAATGCAAAGAGGGGGGGGTTGACAATTCTGTTAAAGAAC	
MET15	sgRNA	GATACTGTTCAACTACACGC	yARC0178
	HR_template	AGATACAATTCTATTACCCCCATCCATACAGTGTGCGTAATGAGTTGTAAAATTATGTAT	
	HR_Fw	GTCTTTTCATCTACTATTTCCTTCGTGTAATACAGGGTCGTCAGATACATAGATACAATT	
	HR_Rv	AATAAAGTTCGTTTTATAAAAGTATAGTACTTGTGAGAGAAAGTAGGTTTATACATAATT	
STR3	sgRNA	CAAGAGATTAGATACAGTTG	yARC0295
	HR_template	agcaaacaacaaaagcatagaagcaaaaagGGGAAAATATGCATGTACCttttttttttt	
	HR_Fw	ACATCCCATAGCTCTGTGTGCTTACAGTTCATTaaaaacaaaaaatatcaagcaaacaac	
	HR_Rv	TATACTTGACTATTTAAAGTTACTATCTTTGGATTTGAACCTTATaaaaaaaaaaaaaaa	
YLL0558W	sgRNA	TGAACAACTATTAATCATCG	yARC0220
	HR_template	GAACAACTATTAATCATCGTGGAAATTATTAGCAGATCATGAGTAGAATATCAAAAATCA	
	HR_Fw	TCTCTGGTGCGGTAAAACATAAATTTTAAATCTAGTACTCAACCACCACTGAACAACTAT	
	HR_Rv	TGGAAAAAAACTTTTATAAACAATATTCTATAAAGATTATAAAGAATTCTTGATTTTTGA	

164 **Table S9.** Yeast growth conditions in this study (related to **Materials and Methods**). When required, the
165 analogous liquid medium was made with the same composition, excluding the 2% agar.
166

Growth Conditions	Composition	Used in Figure(s)
BiGGY	10.0 g/L glucose	2C, 2D, 4C, S8B, S8C
	10.0 g/L glycine	
	1.0 g/L yeast extract	
	3.0 g/L Na2SO4	
	2.5 g/L bismuth ammonium citrate	
	2 % Agar	
Homemade YNB – Inorganic Sulfates (SD-Met+Glu)	200 ul/L of Vitamin Stock (5000x)	2E
	200 ul/L of Trace Elements Stock (10 000x)	
	10 ml/L of Salt Stock (100x)	
	2.0 g/L '-met-cys+ura' drop-out (+histidine, +leucine, +lysine, +uracil)	
	1 g/L L-glutamic acid	
	2 % Glucose/Galactose	
	2 % agar	
Homemade YNB + Inorganic Sulfates (SD-Met+Glu)	200 ul/L of Vitamin Stock (5000x)	2E
	200 ul/L of Trace Elements Stock (10 000x)	
	10 ml/L of Salt Stock (100x)	
	2.0 g/L '-met-cys+ura' drop-out (+histidine, +leucine, +lysine, +uracil)	
	1 g/L L-glutamic acid	
	2 % Glucose/Galactose	
	2 % agar	
SC-Leu+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	S2
	2.0 g/L SC drop-out (-leucine)	
	1 g/L L-glutamic acid	
	2 % Galactose	
	2 % Agar	
SC-Met-Cys+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	4C, S2
	2.0 g/L SC drop-out (-methionine, -cysteine)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
SC-Met-Cys+Gal	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	S2
	2.0 g/L SC drop-out (-methionine, -cysteine)	
	1 g/L L-glutamic acid	
	2 % Galactose	
	2 % Agar	
SD-Met+EtOH	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1C, 4E, S3B, S4
	2.0 g/L '-met-cys' drop-out (+histidine, +leucine, +lysine, +uracil)	
	1 g/L L-glutamic acid	
	3 % Ethanol	
	2 % Agar	
SD-Met+Gal	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1C, 3D, S3B, S4
	2.0 g/L '-met-cys' drop-out (+histidine, +leucine, +lysine, +uracil)	
	1 g/L L-glutamic acid	
	2 % Galactose	
	2 % Agar	
SD-Met+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1B, 1C, 1E, 1F, 1G, 2B, 4B, 4C, 4D, 4E, S1, S3A, S3B, S4, S8A
	2.0 g/L '-met-cys' drop-out (+histidine, +leucine, +lysine, +uracil)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
SD-Ura+EtOH	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1C, S3B, S4
	2.0 g/L '+met-cys-ura' drop-out (+histidine, +leucine, +lysine, +methionine)	
	1 g/L L-glutamic acid	
	3 % Ethanol	
	2 % Agar	
SD-Ura+Gal	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1C, S3B, S4
	2.0 g/L '+met-cys-ura' drop-out (+histidine, +leucine, +lysine, +methionine)	
	1 g/L L-glutamic acid	
	2 % Galactose	
	2 % Agar	

SD-Ura+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1C, S3B, S4
	2.0 g/L '+met-cys-ura' drop-out (+histidine, +leucine, +lysine, +methionine)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
SD-Leu+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1C, S3B
	2.0 g/L '+met-cys-leu' drop-out (+histidine, +uracil, +lysine, +methionine)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
SD+Met+EtOH	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	4E
	2.0 g/L '+met-cys' drop-out (+histidine, +leucine, +lysine, +methionine)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
	2 % Agar	
SD+Met+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	1C, 2E, 4D
	2.0 g/L '+met-cys' drop-out (+histidine, +leucine, +lysine, +uracil, +methionine)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
SD-Met+Bi+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	S5A, S5B, S8B, S8C
	2.0 g/L '-met-cys' drop-out (+histidine, +leucine, +lysine, +uracil)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
	2.5 g/L bismuth ammonium citrate	
YPDA	20 g/L Peptone	2B, 4C, S8A
	10 g/L Yeast Extract	
	2 % Galactose	
	120 mg/L Adenine	
	2 % Agar	
SC-Met-Cys+(Fe)EDTA+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	4C
	2.0 g/L SC drop-out (-methionine, -cysteine)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
	0.048M Fe-EDTA	
SD-Met+(Fe)EDTA+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	4B, S10
	2.0 g/L '-met-cys' drop-out (+histidine, +leucine, +lysine, +uracil)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
	0 - 0.136M Fe-EDTA	
SD+His+Leu+Ura+Glu	1.7 g/L Yeast Nitrogen Base without amino acids and ammonium sulfate	S1
	2.0 g/L drop-out (+histidine, +leucine, +uracil)	
	1 g/L L-glutamic acid	
	2 % Glucose	
	2 % Agar	
GNA	50 g/L Glucose	S1
	30 g/L Difco Nutrient Broth	
	10 g/L Yeast Extract	
	2 % agar	
SPO	10 g/L Glucose	S1
	1.25 g/L Yeast Extract	
	10 g/L Potassium Acetate	
	10 ml/L (100mM lysine)	
	3 mL/L (100mM histidine)	
	2 % agar	

Homemade Yeast Nitrogen Base without Inorganic Sulfate	
Component	Concentration
	Vitamins
Biotin	2µg/L
Calcium Pantothenate	400µg/L
Folic Acid	2µg/L
Inositol	2,000µg/L
Niacin	400µg/L
p-Aminobenzoic Acid	200µg/L
Pyridoxine Hydrochloride	400µg/L
Riboflavin	200µg/L
Thiamine Hydrochloride	400µg/L
	Trace Elements
Boric Acid	500µg/L
Cupric chloride	37.74µg/L
Potassium Iodide	100µg/L
Ferric Chloride	200µg/L
Manganese Chloride	333.34µg/L
Sodium Molybdate	200µg/L
Zinc Sulfate	400µg/L
	Salts
Monopotassium Phosphate	1.0g/L
Magnesium Chloride	0.395g/L
Sodium Chloride	0.1g/L
Calcium Chloride	0.1g/L

169
170

Homemade Yeast Nitrogen Base with Inorganic Sulfate	
Component	**Concentration**
	Vitamins
Biotin	2µg/L
Calcium Pantothenate	400µg/L
Folic Acid	2µg/L
Inositol	2,000µg/L
Niacin	400µg/L
p-Aminobenzoic Acid	200µg/L
Pyridoxine Hydrochloride	400µg/L
Riboflavin	200µg/L
Thiamine Hydrochloride	400µg/L
	Trace Elements
Boric Acid	500µg/L
Copper Sulfate	40µg/L
Potassium Iodide	100µg/L
Ferric Chloride	200µg/L
Manganese Sulfate	400µg/L
Sodium Molybdate	200µg/L
Zinc Sulfate	400µg/L
	Salts
Monopotassium Phosphate	1.0g/L
Magnesium Sulfate	0.5g/L
Sodium Chloride	0.1g/L
Calcium Chloride	0.1g/L

172 **SUPPLEMENTARY REFERENCES**
173
Alberti, S., Gitler, A.D., and Lindquist, S. (2007). A suite of Gateway cloning vectors for

high-throughput genetic analysis in Saccharomyces cerevisiae. Yeast *24*, 913-919.

10.1002/yea.1502.

Brewster, J.L., Pachl, P., McKellar, J.L.O., Selmer, M., Squire, C.J., and Patrick, W.M.

(2021). Structures and kinetics of Thermotoga maritima MetY reveal new insights into

the predominant sulfurylation enzyme of bacterial methionine biosynthesis. J Biol Chem

296, 100797. 10.1016/j.jbc.2021.100797.

Butcher, R.A., Bhullar, B.S., Perlstein, E.O., Marsischky, G., LaBaer, J., and Schreiber,

S.L. (2006). Microarray-based method for monitoring yeast overexpression strains

reveals small-molecule targets in TOR pathway. Nat Chem Biol *2*, 103-109.

10.1038/nchembio762.

Douglas, A.C., Smith, A.M., Sharifpoor, S., Yan, Z., Durbic, T., Heisler, L.E., Lee, A.Y.,

Ryan, O., Gottert, H., Surendra, A., et al. (2012). Functional analysis with a barcoder

yeast gene overexpression system. G3 (Bethesda) *2*, 1279-1289.

10.1534/g3.112.003400.

Giaever, G., Chu, A.M., Ni, L., Connelly, C., Riles, L., Veronneau, S., Dow, S., Lucau-

Danila, A., Anderson, K., Andre, B., et al. (2002). Functional profiling of the

Saccharomyces cerevisiae genome. Nature *418*, 387-391. 10.1038/nature00935.

Kredel, S., Oswald, F., Nienhaus, K., Deuschle, K., Rocker, C., Wolff, M., Heilker, R.,

Nienhaus, G.U., and Wiedenmann, J. (2009). mRuby, a bright monomeric red

fluorescent protein for labeling of subcellular structures. PLoS One *4*, e4391.

10.1371/journal.pone.0004391.

Lu, H., Li, F., Sanchez, B.J., Zhu, Z., Li, G., Domenzain, I., Marcisauskas, S., Anton,

P.M., Lappa, D., Lieven, C., et al. (2019). A consensus S. cerevisiae metabolic model

Yeast8 and its ecosystem for comprehensively probing cellular metabolism. Nat

Commun *10*, 3586. 10.1038/s41467-019-11581-3.

Winston, F., Dollard, C., and Ricupero-Hovasse, S.L. (1995). Construction of a set of

convenient Saccharomyces cerevisiae strains that are isogenic to S288C. Yeast *11*, 53-

55. 10.1002/yea.320110107.

Winzeler, E.A., Shoemaker, D.D., Astromoff, A., Liang, H., Anderson, K., Andre, B.,

Bangham, R., Benito, R., Boeke, J.D., Bussey, H., et al. (1999). Functional

characterization of the S. cerevisiae genome by gene deletion and parallel analysis.

Science *285*, 901-906. 10.1126/science.285.5429.901.