1 **Title: Excessive self-grooming of *Shank3* mutant mice is associated with gene dysregulation and**

2 **imbalance between the striosome and matrix compartments in the striatum**

Author names and affiliations, including postal codes:

Ferhat Allain-Thibeault* (1,2,3), Biton Anne* (4), Verpy Elisabeth* (1), Forget Benoit (1), de

Chaumont Fabrice (1), Mueller Florian (5), Le Sourd Anne-Marie (1), Coqueran Sabrina (1), Julien

Schmitt (6), Rochefort Christelle (6), Rondi-Reig Laure (6), Leboucher Aziliz (1), Boland Anne (7),

Fin Bertrand (7), Deleuze Jean François (7,8), Boeckers Tobias M. (9,10), Ey Elodie[#] (1,11),

Bourgeron Thomas[#] (1)

* Equally contributed to the work

[#] Co senior-authors

1. Génétique Humaine et Fonctions Cognitives, Institut Pasteur, UMR3571 CNRS, Université de

Paris, Paris, France

2. Department of Neuroscience, Columbia University Medical Center, New York, NY, 10027

3. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, 10027

4. Centre de Bioinformatique, Biostatistique et Biologie Intégrative (C3BI, USR 3756 Institut Pasteur

and CNRS), Paris, France

5. Unité Imagerie et Modélisation (UMR 3691 Institut Pasteur and CNRS), Paris, France

6. Sorbonne Université, CNRS, INSERM, Institut de Biologie Paris Seine (IBPS), Neurosciences Paris

Seine (NPS), Cerebellum Navigation and Memory Team (CeZaMe), F-75005 Paris, France

7. Université Paris-Saclay, CEA, Centre National de Recherche en Génomique Humaine (CNRGH),

91057, Evry, France

8. Centre d'étude du polymorphisme humain, Paris, France

9. Institute of Anatomy and Cell Biology, Ulm University, 89081 Ulm, Germany

10. Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), 89081 Ulm, Germany

11. Université de Strasbourg, CNRS, INSERM, Institut de Génétique et de Biologie

Moléculaire et Cellulaire - UMR 7104, Illkirch-Graffenstaden, France

Corresponding author email address:

thomas.bourgeron@pasteur.fr

ferhat.allainthibeault@gmail.com

Abstract

Autism spectrum disorders (ASD) are characterised by atypical social communication and stereotyped behaviours. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are detected in 1-2% of patients with ASD and intellectual disability (ID), but the mechanisms underpinning the symptoms remain largely unknown. Here, we characterized the behaviour of *Shank3* mutant mice deleted for exon 11 (*Shank3*$^{\Delta11/\Delta11}$) from three to twelve months of age. We observed decreased locomotor activity, increased stereotyped self-grooming and atypical socio-sexual interaction compared to wild-type littermates. We then used RNAseq on four brain regions of the same animals to identify differentially expressed genes (DEG). DEGs were identified mainly in the striatum and were associated with synaptic transmission (e.g. *Grm2, Dlgap1*), G-protein-signalling pathways (e.g. *Gnal*, *Prkcg1, and Camk2g*), as well as excitation/inhibition balance (e.g. *Gad2*). Downregulated and upregulated genes were respectively enriched in the gene clusters of medium-sized spiny neurons expressing the dopamine 1 (D1-MSN) and the dopamine 2 receptor (D2-MSN). Moreover, expression of DEGs reported as striosome markers within the striatum (*Cnr1*, *Gnal1*, *Gad2*, *and Drd4*) were positively correlated with excessive self-grooming. Finally, we showed that the striosome compartment of *Shank3*$^{\Delta11/\Delta11}$ mice was enlarged and displayed higher expression of GAD65 compared to wild-type mice. Altogether, these results shed light on a possible role of the striosomes/matrix imbalance in excessive self-grooming in *Shank3*$^{\Delta11/\Delta11}$ mice. Such striatal alterations could be present in a subgroup of patients with ASD and ID and could open the way to new therapeutic approaches.

Introduction

Autism spectrum disorders (ASD) are neurodevelopmental conditions characterised by atypical social communication and interactions associated with stereotyped behaviours and restricted interests (1). More than 200 genes have been robustly associated with ASD pointing at biological pathways such as chromatin remodelling and synaptic function (2). Among these genes, *SHANK3* (SH3 and multiple ankyrin repeat domain 3) codes for a scaffolding protein located at the postsynaptic density (PSD) of glutamatergic synapses where it interacts with other scaffolding proteins, cytoskeletal proteins, glutamate receptors and signalling molecules (3).

Heterozygous *de novo* mutations deleting the *SHANK3* gene on chromosome 22q13.3 or affecting its coding region are associated with Phelan-McDermid syndrome (4) and observed in 1-2% of patients with both ASD and intellectual disability (ID), making this gene a major cause of neurodevelopmental disorders (NDD) (5,6). Remarkably, a relatively high proportion of patients (>60%) carrying *SHANK3* mutations display regression (i.e., substantial loss of language and social skills) during adolescence and adulthood (7,8). However, the mechanisms of this worsening of the phenotype remain unknown.

Through the use of multiple intragenic promoters and alternative splicing, several SHANK3 protein isoforms are differentially expressed according to developmental stage or brain region (9). Numerous *Shank3* mutant mice have been generated [for reviews see (3,10)]. Most of them bear deletion of specific exons still allowing the expression of some minor isoforms (9,11). The behavioural deficits reported for *Shank3* mutant mice depend on the strains and experimental conditions. However, the presence of stereotyped-like behaviours, such as excessive self-grooming, was reported in all *Shank3* mutant mice and is reminiscent of the stereotyped behaviours observed in patients with ASD (3,12,13). Stereotyped behaviours are frequently associated with anomalies of the striatum (14), a region that coordinates multiple aspects of cognition, including action planning, decision-making, motivation, reinforcement, and reward perception as well as motor control. Several studies identified anomalies in the striatum of *Shank3* mutant mice at the anatomical (e.g., larger volume), circuit (e.g., disruption of indirect pathway of basal ganglia) and cellular (e.g., decreased striatal expression of GluN2B, a NMDA

receptor subunit) levels (15–20). The biological pathways underpinning the observed abnormalities as

well as the biomarkers predictive of phenotypic severity remain however largely unknown.

In the present work, we first conducted a thorough longitudinal behavioural characterization of

$Shank3^{\Delta11/\Delta11}$ mice, carrying a deletion of exon 11 of $Shank3$ (which still allow the expression of minor

protein isoforms) (21), at three, eight and twelve months of age, focusing on social interactions and

stereotyped behaviours. We then compared gene expression in four brain structures - cortex,

hippocampus, cerebellum and striatum - in $Shank3^{+/+}$ and $Shank3^{\Delta11/\Delta11}$ male littermates characterised

in the behavioural study. Compared to the other brain regions, the striatum appeared to be highly

sensitive to the loss of the major SHANK3 protein isoforms. We also identified a correlation between

the level of self-grooming and an increased striatal expression of several genes, including $Gad2$

encoding GAD65, which catalyses the transformation of glutamate into GABA. Finally, using

immunofluorescence on the striatum sections at 12 months, we found that the expression of GAD65 is

increased in the patch/striosome compartment of the $Shank3^{\Delta11/\Delta11}$ mice compared to wild-type

littermates and that the surface of this compartment is also increased, relative to the matrix compartment.

Our results suggest that signalling imbalance between the striosome and matrix compartments of the

striatum causes excessive self-grooming in $Shank3^{\Delta11/\Delta11}$ mice.

Material & methods

<u>Animals and cohorts</u>

$Shank3^{\Delta11/\Delta11}$ mice were generated by Genoway (Lyon, FRANCE) using 129S1/SVImJ ES cells. The

$Shank3$ mutation, deletion of exon 11, was then transferred onto a C57BL/6J background with more

than 15 backcrosses (21). In this model, short protein isoforms are still expressed (21). Food and water

were provided *ad libitum*. The experimenters were blind to the genotype of the tested animals for data

collection and analyses. We bred $Shank3^{\Delta11/+}$ males and females to obtain $Shank3^{+/+}$, $Shank3^{+/\Delta11}$ and

$Shank3^{\Delta11/\Delta11}$ littermates. For behavioural tests, we used three cohorts (including both males and

females) that we analysed separately. See **Supplementary Methods** for details on the cohorts.

<u>Ethical statement</u>

All procedures were validated by the ethical committee of the Institut Pasteur (CEEA n°89) and of the

Sorbonne Université (CEEA n°5) and were conducted under the approval of the Ministère de

l'Enseignement Supérieur, de la Recherche et de l'Innovation under the reference APAFIS#7706-20

161 12317252460 v2 and reference APAFIS#3234-2015121619347680 v7, respectively.

<u>Behavioural tests</u>

Dark/light test, Elevated Plus-maze test, locomotion and exploratory test in the open field, rotarod,

occupant/new-comer social test with ultrasonic vocalisation recording, male behaviour in presence of

an oestrous female, three-chambered social test and starmaze test were used to investigate the mice's

behaviours. The protocols of these tests are detailed in the "behavioural tests" section of the

Supplementary Methods.

<u>Observation of stereotyped behaviour</u>

Mice were individually placed in a new test cage (Plexiglas, 50 x 25 x 30 cm; 100 lux; clean sawdust

bedding) in a soundproof chamber. After 10 min habituation, we recorded their behaviour for 10 min

(camera Logitech C920). We manually scored the time spent self-grooming and digging (The Observer,

Noldus Information Technology). When scoring self-grooming, we did not take into account the

scratching behaviour. The total duration and the average duration of each behavioural category were

calculated.

<u>Behavioural statistics</u>

All group data are represented as mean ± standard error of the mean (s.e.m.), as well as the individual

points. All statistics were performed with R software (22). In all behavioural tests, $Shank3^{+/\Delta11}$ mice

were not significantly different from $Shank3^{+/+}$ unless otherwise specified (see **Table S1**). Given the

limited sample size and the non-normality of the data, comparisons between genotypes were performed

using non-parametric Mann-Whitney U-tests. We used the non-parametric Friedman test to examine the

effect of age within each genotype. Between age points, post-hoc tests were performed using paired

Wilcoxon signed-rank tests. To analyse the variability within groups, the variances of the raw data were

compared using Ansari-Bradley tests. If required, a Bonferroni correction for multiple testing was

performed. Differences between groups were considered significant when $p < 0.05$.

<u>Transcriptome analysis</u>

Tissue collection, total RNA extraction and sequencing, mapping and reference genome, calling variants

from the RNAseq data and differential expression analysis are detailed in the "Transcriptome analysis"

section of the **Supplementary Methods.**

<u>Gene set and protein-protein interaction network analysis</u>

Gene set over-representation analyses were performed based on Ensemble of Gene Set Enrichment

Analyses (EGSEA) without considering the genes around the *Shank3* gene. Protein-protein interaction

(PPI) networks were generated using the DEGs for the four brain structures independently. A

randomised protein interaction network was created using all DEGs to determine the probability of

finding a network. Software, packages and databases are detailed in "Gene set and protein-protein

interaction network analysis" section of the **Supplementary Methods.**

<u>Quantitative RT-PCR (q-RT-PCR) and RNA and protein fluorescent detection on striatum sections</u>

We selected several genes of interest by cross checking the DEGs, ORA (gene ontology – GO –, Kyoto

encyclopaedia of genes and genomes – KEGG– and Pathway), and data from the literature. To validate

these genes, we used q-RT-PCR, either with the droplet digital Polymerase Chain Reaction (dd-PCR)

or with the real time quantitative PCR (q-PCR) technology. These procedures are detailed in the

"Quantitative RT-PCR" section of the **Supplementary Methods.** All statistical analyses were

performed using R software (22). The comparison between genotypes was performed using Mann-

Whitney U-tests. Single-molecule Fluorescent In Situ Hybridization (smFISH) experiment to detect

Drd1 and *Drd2* RNAs and immunofluorescence experiment and image analysis to quantify GAD65

immunoreactivity in the dorsal striatum are detailed in the ""Single-molecule Fluorescent In Situ

Hybridization and immunofluorescence experiments on brain sections" and "Quantitative analysis of

GAD65 immunoreactivity" section of the **Supplementary Methods.**

Results

<u>Hypoactivity, atypical social behaviour and excessive self-grooming in $Shank3^{\Delta11/\Delta11}$ mice</u>

At three months of age, $Shank3^{\Delta11/\Delta11}$ mice displayed typical body weight, normal motor coordination,

typical anxiety-like behaviour levels and intact working memory, compared to $Shank3^{+/+}$ littermates

(**Fig. S1**). In the open field, $Shank3^{\Delta11/\Delta11}$ males travelled significantly shorter distances in comparison

Figure 1: Hypoactivity, reduced exploration and increased stereotyped behaviours in *Shank3*^Δ11/Δ11^ mice at three, eight and twelve months of age. (A) Total distance travelled during 30-min free exploration of an open field in wild-type (green), heterozygous (blue) and *Shank3*^Δ11/Δ11^ (orange) for males (left panel) and females (right panel). (B) Total time spent digging in fresh bedding during 10 min observation in a new test cage, after 10 min habituation. (C) Total time spent in contact during male/female and female/female social interaction. (D) Total time spent self-grooming during 10 min observation in a new test cage, after 10 min habituation. (E) Z-score for the time spent self-grooming. (F) Number of individuals displaying different levels of severity in self-grooming. "Dead" corresponds to animals that had to be euthanized due to severe self-inflicted injuries. A-D: Black points represent mean with standard error of the mean (s.e.m). Mann–Whitney U test with Bonferroni correction for multiple testing (2 tests): *: corrected p-value <0.05; **: corrected p-value <0.01. Red lines represent statistics at 3 months of age; black lines represent statistics at 8 and 12 months of age.

with *Shank3*^+/+^ littermates (Cohort 1: W=130, p=0.007; **Fig. 1A,** left panel; confirmed in Cohort 2 and

Cohort 3: **Fig. S2**). *Shank3*^Δ11/Δ11^ males also spent less time digging in the bedding in comparison with

Shank3^+/+^ mice (Cohort 1: W=123, p=0.027; **Fig. 1B;** Cohort 2: W=173, p<0.001; **Fig. S2**). A similar

trend was present, but did not reach significance levels in females for activity and digging behaviour

(Cohort 1: W=92, p=0.07; **Fig. 1A, B**). Remarkably, in stressful swimming conditions, *Shank3*^Δ11/Δ11^

mice swam at a significantly higher speed compared to wild-type mice in the Morris water maze with

visible platform (Cohort 3: p<0.01 over all training) and in the star maze (p<0.01), without any

significant difference to find the direct path in the star maze (**Fig. S2**).

Impairments in social interactions and communication were not always observed in the different

Shank3 mouse models (10). Here, mice of all genotypes and both sexes displayed a typical preference

for a same-sex conspecific in the three-chambered test (**Fig. S3**). No significant difference was observed

in the latency for the first contact, in the total time spent in contact with same sex conspecific, in the

time spent in the different social behaviours and in the emission of the ultrasonic vocalizations (USVs) (**Fig. S3**). In contrast, we observed that $Shank3^{\Delta11/\Delta11}$ males spent significantly more time in contact with an oestrous female compared to $Shank3^{+/+}$ males (W=32, p=0.034; **Fig. 1C**), especially for oro-oral contact (W=22, p=0.005; **Fig. S3**). We also observed a reduction of time keeping the female in the visual field for $Shank3^{\Delta11/\Delta11}$ males compared to $Shank3^{+/+}$ males (W=133, p=0.006; **Fig. S3**). These altered behaviours were not accompanied with atypical vocal behaviour (**Fig. S3**).

At 3 months of age, $Shank3^{\Delta11/\Delta11}$ male mice already displayed a significant increase in the time spent self-grooming in comparison with $Shank3^{+/+}$ males (Cohort 1: W=21, p=0.002; Cohort 2: W=43, p=0.022). Similar trends did not reach significance in female mice (**Fig. 1, Fig. S2**). Interestingly, a higher group heterogeneity was observed in $Shank3^{\Delta11/\Delta11}$ mice in comparison with $Shank3^{+/+}$ mice (Cohort 1: males: Ansari Bradley test AB=111, p=0.036; females: AB=82, p=0.009; **Fig. 1D**). Furthermore, a subsample of $Shank3^{\Delta11/\Delta11}$ males (5 individuals over 12) and females (6 individuals over 12) presented hair removal that evolved into self-injuries.

Shank3 mutant mice display excessive self-grooming behaviour with age

Considering the frequent regression of patients carrying _SHANK3_ mutations, we tested whether a phenotypic deterioration occurred in $Shank3^{\Delta11/\Delta11}$ mice. For that purpose, we tested exploratory activity, social communication and stereotyped behaviours in mice at eight and twelve months of age. The reduction of distance travelled in $Shank3^{\Delta11/\Delta11}$ mice compared to $Shank3^{+/+}$ mice was not present at older age; only $Shank3^{+/+}$ and $Shank3^{+/\Delta11}$ mice reduced their activity with increasing age (**Fig. 1A**). Genotype-related differences in digging behaviour persisted with increasing age (8 months: W= 140, p<0.01; 12 months; W=107; p=0.01; **Fig. 1B**).

During social interactions, we observed a decrease of the time spent in contact over age, in particular for $Shank3^{+/+}$ males (Friedman test: F=7.384, p=0.025), $Shank3^{+/\Delta11}$ males (F=12.316, p=0.002) and $Shank3^{+/+}$ females (F=7.400, p=0.025) and a tendency for $Shank3^{\Delta11/\Delta11}$ males without reaching significance (**Fig. 1C & S3**). In female-female interactions, no behavioural differences between genotypes were observed at eight and twelve months of age, similarly to tests conducted at three months of age (**Fig. S3**). At eight months of age, the duration of oro-oral contact in $Shank3^{\Delta11/\Delta11}$ males

with an oestrous female was still significantly increased, similarly to what we observed at three months

(W=29, p=0.025), but these differences were no longer significant at twelve months of age (**Fig. S3**).

Nevertheless, at this age, we detected an increase of the "approach then escape" behavioural sequence

displayed by *Shank3*^Δ11/Δ11 males toward a C57BL/6J female in comparison with *Shank3*⁺/⁺ males

(W=15.5, p=0.005, **Fig. S3**).

At eight and twelve months of age, *Shank3*^Δ11/Δ11 males still displayed excessive self-grooming

in comparison with *Shank3*⁺/⁺ mice (eight months: W=27, p=0.009; twelve months: W=37, p=0.041;

Fig. 1D). The same was true for *Shank3*^Δ11/Δ11 females who also groomed themselves significantly more

than *Shank3*⁺/⁺ females (eight months: W=11, p=0.002; twelve months: W=20, p=0.014; **Fig. 1C**).

Interestingly, higher inter-individual variability was still present in *Shank3*^Δ11/Δ11 mice compared to

Shank3⁺/⁺ mice (at eight months for males: AB=118, p=0.002; and females: AB=75, p=0.044; at twelve

months for males: AB=102, p<0.001; **Fig. 1E**). With age, an increasing number of *Shank3*^Δ11/Δ11 mice

were outliers, performing self-grooming during a period exceeding two standard deviations (thresholds

for males: 3 months: >80.8s; 8 months: >231s; 12 months: >208.4s; thresholds for females: 3 months:

>78.3s; 8 months: >188.8s; 12 months: >171.8 s) or had to be euthanized because of severe self-inflicted

injuries. Nevertheless, for 4 out of 12 *Shank3*^Δ11/Δ11 males and 3 out of 12 *Shank3*^Δ11/Δ11 females, we did

not observe excessive self-grooming at any of the three time points (**Fig. 1E, F**).

Differential expression of genes related to synaptic function, signalling and cytoskeleton dynamics in the striatum of $Shank3^{\Delta11/\Delta11}$ mice

To investigate the impact of the loss of the major SHANK3 isoforms on the transcriptome, we performed RNAseq on four brain structures (whole cortex, striatum, hippocampus, and cerebellum) of seven $Shank3^{+/+}$ and eight $Shank3^{\Delta11/\Delta11}$ male mice after their behavioural characterization at twelve months of age. DEGs between $Shank3^{+/+}$ and $Shank3^{\Delta11/\Delta11}$ mice (FDR < 0.05) across and within structures (**Table S2**) were enriched in genes located in the vicinity of the $Shank3$ locus on chromosome 15, most likely due to the residual genomic region from the 129S1/SvImJ ES cells used to generate the $Shank3^{\Delta11/\Delta11}$ mice (**Fig. S4**) (21). We therefore excluded these genes from the analyses (more details in **Supplementary methods**). We also performed quantitative RT-PCR (q-RT-PCR) to confirm up- or downregulation of 30 DEGs (Spearman correlation R=0.809 P<0.0001; **Supplementary Fig. S5**). As expected, $Shank3$ expression levels in all four brain structures were significantly reduced in $Shank3^{\Delta11/\Delta11}$ mice in comparison with $Shank3^{+/+}$ mice (FDR < 0.001; **Fig. S6**). Notably, the $Shank3$ sequence reads still present in the $Shank3^{\Delta11/\Delta11}$ mice were aligned to exons downstream of exon 11 (**Table S2**). No significant changes in the expression of $Shank1$ and $Shank2$ were found (**Fig. S6**, **Table S2**).

We examined the DEGs (FDR < 0.05) within each of the four brain structures. The largest number of DEGs was detected for the striatum with 140 DEGs. The hippocampus displayed 26 DEGs, the cortex 13 and the cerebellum 16 (**Fig. 2A**). After removing the genes and pseudogenes located around the $Shank3$ locus, $Shank3$ was the only gene differentially expressed in the four brain structures (**Fig. 2A**). The DEGs in the striatum displayed an increased dispersion compared to the other structures, indicating an increased variability of response between animals (**Fig.S7**). Altogether, these results suggest that, among the four structures studied, the transcriptome of the striatum is by far the most impacted by the loss of the major isoforms of SHANK3.

We investigated the functions of the proteins encoded by the DEGs in the striatum using over-representation analysis (ORA) and Protein-Protein Interaction (PPI) analysis. The ORA (**Table S3**) revealed enrichment in DEGs associated with, among others, two pathways of interest: synaptic transmission (negative fold-change in $Shank3^{\Delta11/\Delta11}$ mice in comparison with $Shank3^{+/+}$ mice), and G-

protein activity (positive and negative fold-change in $Shank3^{\Delta11/\Delta11}$ mice in comparison with $Shank3^{+/+}$

mice according to the type of G-protein pathway) (**Fig. 2b** and **Table S3**). PPI analysis of the striatal

DEGs highlighted one network of 15 proteins (p-value from randomized network: <0.01; **Fig. 2C**), some

of them directly interacting with SHANK3 in the PSD, such as the metabotropic glutamate receptor 2

(encoded by *Grm2*, highly decreased) and Disks large-associated protein 1, a scaffolding protein

(encoded by *Dlgap1*, also decreased) interacting with AMPA and NMDA receptors. In addition to these

synaptic proteins, the network contains proteins involved in signal transduction (e.g. *Prkcg1*, *Camk2g*)

and cytoskeleton dynamics (e.g. *Slk*). Other small networks identified by the PPI analysis contain

proteins involved in glutamate transportation (*Slc17a7*) and decarboxylation (*Gad2*), and cytoskeleton

dynamics (*Actbl2*).

<u>Cellular expression pattern of the DEGs expressed in the striatum of $Shank3^{\Delta11/\Delta11}$ mice</u>

The inhibitory medium-sized spiny neurons (MSN) constitute the major type of striatal neuronal

population. They receive glutamatergic inputs and are the target of dopamine innervation from cortex,

thalamus, amygdala and substantia nigra pars compacta (SNpc) (23). Most of these MSN express either

dopamine 1 receptor DRD1 (D1-MSN, belonging to the direct pathway), dopamine 2 receptor DRD2

(D2-MSN, belonging to the indirect pathway), or both receptors for a small fraction of MSN. In order

to identify the striatal cell types which are the most impacted by SHANK3 deficiency, we compared the

expression pattern of the DEGs with marker genes of striatal cell types reported in two single-cell RNA

sequencing studies (24,25). We observed that DEGs under-expressed in $Shank3^{\Delta11/\Delta11}$ compared to

$Shank3^{+/+}$ mice were enriched in the D1-MSN clusters while overexpressed genes were enriched in the

D2-MSN clusters (**Fig. 3A, Table S2 & S4**). This could reflect a modified proportion of D1-MSN

(decrease) and D2-MSN (increase) in the striatum of the $Shank3^{\Delta11/\Delta11}$ mice. However, many genes

commonly considered as specific to D1-MSN (*Drd1*, *Isl1*, *Sfxn1* and *Tac1*) or D2-MSN (*Drd2*, *Adora2a*,

Penk, *Gpr6*, *Sp9*) were not found to be differentially expressed in the $Shank3^{\Delta11/\Delta11}$ animals (**Table S2**).

Moreover, using Single Molecule RNA Fluorescence In Situ Hybridization with *Drd1* and *Drd2* RNA

probes, we observed no significant difference in the number of D1 and D2 striatal neurons between the

$Shank3^{\Delta11/\Delta11}$ and $Shank3^{+/+}$ mice (**Fig. S8**). Altogether, these results indicate that the deletion of exon

Figure 3: Altered transcriptome of D1- and D2-MSN and higher expression of GAD65 in the striosomal compartment in the striatum of *Shank3*$^{\Delta 11/\Delta 11}$ mice.

(A) Enrichment of RNAseq DEGs in the striatum of *Shank3*$^{\Delta 11/\Delta 11}$ mice compared to cell-specific gene clusters suggested by Gocke et al. (2016) and Montalban et al. (2020). Genes under-expressed in *Shank3*$^{\Delta 11/\Delta 11}$ striatum are enriched in D1-MSN clusters while genes over-expressed in *Shank3*$^{\Delta 11/\Delta 11}$ striatum are enriched in D2-MSN clusters. Ratio genes correspond to the proportion of DEGs enriched into the cluster. Over-representation analyses were performed with Fisher's hypergeometric tests and p-values were adjusted for multiple testing with the Benjamini-Hochberg procedure within each category of DEGs. (B) Comparison of the log of p-values for the differential expression of selected genes in *Shank3*$^{+/+}$ vs *Shank3*$^{\Delta 11/\Delta 11}$ mice and the log of p-values for low self-grooming vs high self-grooming. (C-D) Confocal images of striatum coronal sections of one-year old *Shank3*$^{+/+}$ male mice. (C) GAD65 immunoreactivity (red) is enriched in striatal microzones identified as striosomes by immunostaining for μ-opioid receptor (MOR, in green), a canonical marker of striosomes. (D) Increased GAD65 immunoreactivity in striosomes (arrows) and subcallosal streak (two-headed arrows)

← compared to surrounding matrix is observed with two different antibodies: the rabbit polyclonal antibody (Invitrogen PA5-77983 in red) used in C, E, F and a mouse monoclonal antibody (Millipore MAB351, in green). (E-F) Distribution and quantification of GAD65 immunoreactivity in dorsal striatum sections of a *Shank3*$^{+/+}$ (E) and a *Shank3*$^{\Delta 11/\Delta 11}$ (F) mouse. The arrows point to striosomes and the arrowheads to subcallosal streak. The unlabelled myelinated fibres are coloured in red. The regions of higher expression of GAD65 (striosomes) are coloured in green, and the regions of lower expression of GAD65 (matrix) are uncoloured. (G, H, I) Comparison of GAD65 immunoreactivity in the striosome and matrix compartments of the dorsal striatum in 5 *Shank3*$^{+/+}$ (green) and 9 *Shank3*$^{\Delta 11/\Delta 11}$ (orange) one-year old male mice. (G) Relative surface of the GAD65-enriched striosome compartment (surface of striosomes / surface of (striosomes + matrix)). (H) Relative GAD65 labelling intensity in the striosome compartment compared to the matrix compartment (GAD65 labelling in striosomes / GAD65 labelling in matrix). (I) Overall GAD65 immunoreactivity in the striosomal compartment (relative surface (G) multiplied by the relative GAD65 labelling intensity (H)). Data, generated from analysis of 9 to 14 images per animal, are presented as box-plots (median, first, and third quartiles). Scale bars: 50 μm (C), 200 μm (D), 500 μm (E, F).

11 of *Shank3* leads to differential transcriptional alterations in the D1- and D2-MSN, without major

difference in the number of these types of neurons throughout the whole striatum.

<u>Genes with expression that correlate with excessive self-grooming</u>

Using RT-PCR, we investigated the correlation between the mRNA level in the striatum of DEGs

identified by RNAseq analysis and other genes of interest (*Shank1*, *Shank2*, *and Drds*) and the level of

self-grooming observed in seven 12 months-old *Shank3*$^{\Delta 11/\Delta 11}$ mice. We found that excessive grooming

was associated with higher levels of transcripts for four of the DEGs identified by RNAseq or RT-PCR:

Cnr1, *Gnal*, *Gad2,* and *Drd4* (**Fig. 3B; FigS9**). *Cnr1* encodes the cannabinoid receptor 1; *Gnal* encodes

a stimulatory G-alpha subunit of a heterotrimeric G-protein (Gαolf); *Gad2* codes for GAD65, one of the

two enzymes that convert glutamate into γ-aminobutyric acid (GABA), and *Drd4* encodes the dopamine

receptor 4. Interestingly, these proteins have been reported to be heterogeneously distributed in the

dorsal striatum, being more abundant in the spatial microzones known as striosomes or patches than in

the much larger matrix surrounding them (26–29). Given the function of GAD65, i.e. converting

glutamate into GABA, an inhibitory transmitter, we studied its striatal distribution in the dorsal striatum,

using immunofluorescence on brain sections (**Fig. 3C-I**). We found that GAD65 is enriched in

striosomes, identified by co-labelling of μ-opioid receptor (MOR), the most widely used marker for

striosomes of the rostral striatum (30) (**Fig. 3C**). GAD65 can thus be considered as a marker of the

striosome compartment in mice. We also confirmed the specificity of GAD65 labelling using a second

independent GAD65 antibody (**Fig. 3D**). Both the relative surface occupied by the GAD65-enriched

striosomes in the dorsal striatum and the GAD65 labelling intensity of these patches relative to matrix

were increased in the *Shank3*$^{\Delta11/\Delta11}$ mice compared to the *Shank3*$^{+/+}$ mice (p = 0.0016 and p = 0.0055

respectively) (**Fig. 3E-H**)), leading to an increased GAD65 immunoreactivity in the striosome

compartment vs the matrix compartment in *Shank3*$^{\Delta11/\Delta11}$ animals (p=0.0003, **Fig. 3I**).

Discussion

The present study highlighted robust behavioural deficits in *Shank3*$^{\Delta11/\Delta11}$ mutant mice, more specifically

a reduced activity in the open field, atypical social behaviour in males interacting with oestrous females,

and increased self-grooming compared to wild-type littermates. Interestingly, the majority of

Shank3$^{\Delta11/\Delta11}$ mutant mice displayed excessive self-grooming, with a tendency to a worsening of the

phenotype with increasing age for mice showing an early emergence of excessive grooming. At 12

months of age, the striatum had the most impacted transcriptomic profile compared to other brain

regions. Further molecular characterizations suggest that the excessive self-grooming phenotype might

be related to signalling imbalances between the striosome and the matrix compartments in the dorsal

striatum.

Behavioural profiling of the Shank3*$^{\Delta11/\Delta11}$ *mutant mice

Atypical social interaction: Results from the literature have highlighted either reduced duration of social

contact and reduced number of USVs emitted [e.g., *Shank3b* (31); *Shank3*$^{\Delta ex4-22}$ (32)] or no significant

difference in socio-sexual behaviours [e.g., *Shank3*^{Q321R} (33)]. Here, we observed limited difference in

social interaction with an increase in the time spent in contact for *Shank3*$^{\Delta11/\Delta11}$ males interacting with

an oestrous female (observed at 3 and 8 months of age), and more specifically in nose-to-nose contacts,

as already highlighted in previous studies [*Shank3*$\Delta^{ex14-16}$cKO (33); *Shank3*$^{\Delta11/\Delta11}$ females (34)]. Testing

the specificity of this type of contact would help to understand its significance in mouse social

communication. Further experiments should be designed to test whether this specific behaviour is

associated with a deficit in social reward processing [as suggested for *Shank2*$^{\Delta6-7/\Delta6-7}$ mice (35)]. Indeed,

several studies have shown that SHANK3 controls maturation of social circuits in the ventral tegmental area (VTA) (36,37) and synaptic strength of D2-MSN in the striatum (16,20).

Excessive grooming: Excessive self-grooming has been observed in the majority of *Shank3* mutant mice over their lifetime and therefore represents the most robust behavioural phenotype in all models. Extreme self-grooming is not due to increased skin sensitivity since previous studies reported that rescuing normal tactile reactivity in *Shank3b*$^{+/-}$ does not improve over-grooming behaviour (38). In our study, we also observed an important inter-individual variability in this trait with 30% of the mutant mice that did not seem to display excessive grooming even after one year. Increased self-grooming might reflect a response to stressful conditions, which might vary from one individual to another (difference in susceptibility and/or exposure to stressful conditions). Such increased reactivity to stressful conditions is reminiscent of what is observed in patients and deserves further systematic testing in the different *Shank3* mutant models (39).

From transcriptome to imbalance of striatum compartments and excessive grooming

Abnormal striatal transcriptome: The massive impact of the *Shank3* mutation on the transcriptome profile of the striatum suggests that this region could be the centrepiece of the behavioural deficits observed in *Shank3* mutant mice. Several studies have already reported functional alterations of the striatum and more specifically of the MSN in the diverse *Shank3* mutant strains (12,15,16,20,36,37). Our report of an alteration of the striatal transcriptome could be due to a direct role of the SHANK3 protein through gene transcription. Indeed, previous studies have reported the presence of SHANK3 in the nucleus, especially the isoform SHANK3b, which is disrupted in our model. Alternatively, the abnormal striatal transcriptome profile could be the consequence of the alteration of the glutamatergic synapses that interfere with signal transduction, such as G protein (40) and mTOR (41) signalling, and then with gene transcription or mRNA stability. We found that several transcripts encoding proteins involved in glutamatergic synaptic transmission were under-expressed in *Shank3*$^{\Delta 11/\Delta 11}$ mice. For example, we observed a decrease of *Grm2*, encoding mGluR2, a G-protein-coupled glutamate receptor which interacts with the proline-rich domain of SHANK3 and which has been found decreased in a

valproate-induced rat model of autism (42). We also observed significant under-expression of *Dlgap1*,

coding for DAP1/GKAP. This protein connects SHANK3 to the scaffolding protein PSD-95 (43) which

recruits AMPAR and NMDAR. In contrast, we did not observe diminution of transcript levels for several

proteins (GluA, GluN or mGluR5) previously shown as decreased in the synaptosomes or PSD fractions

of *Shank3* mutant mice compared to wild type. A possible explanation is that the absence of SHANK3

decreases the translation or the recruitment of these proteins at the synapse or increases their degradation

without affecting gene transcription and mRNA degradation.

A link between abnormal gene transcription, striatum organisation and excessive grooming: We

observed that expression of *Gad2*, *Cnr1*, *Gnal* and *Drd4* was positively correlated with excessive self-

grooming. Remarkably, three of these genes have been previously associated with self-grooming, but

not always in the same direction. In contrast to our study, *Cnr1* deletions (as well as CB1R antagonists)

have been reported to increase self-grooming behaviour (44–47). The overexpression of *Cnr1* in

Shank3$^{\Delta11/\Delta11}$ mice might therefore result from a compensatory upregulation in response to the alterations

of the endocannabinoid system which has been previously reported in SHANK3-deficient mice (20,48),

in valproic acid induced rat models of ASD and also in some individuals with ASD (49). Consistent

with our findings, *Gnal* haploinsufficiency is associated with a reduction of self-grooming behaviour in

a mouse model of dystonia (50). Links between *Shank3* and *Gad2* were previously reported with an

increase of GAD65 immunoreactivity in the *Shank3*ß$^{-/-}$ mice (44) and a decrease of *Gad2* transcript in

the striatum of a *Shank3*-overexpressing model (51). To our knowledge, there is no direct evidence in

the literature for an association between *Drd4* expression and grooming behaviour. It would thus be

interesting in future studies to investigate this potential link.

Interestingly, these four genes have also been reported to be enriched in striosomes (26–29). These

microzones of the dorsal striatum contain early born MSN and are embedded into the much larger

surrounding matrix including late-born MSN (52). The two compartments, initially distinguished by

differential expression of numerous molecular markers [for a review see (53)], also differ in their input

and output connectivity and electrophysiological characteristics (54), such as D1-MSN excitability (55),

rates of dopamine release (56), and response to chronic stress (57). Signalling imbalances between the

Figure 4: Proposed model linking increased activity in the striosomal pathway and imbalance between the direct and indirect pathways in the _Shank3_$^{\Delta 11/\Delta 11}$ mice. Increased activity of the striosomal pathway in _Shank3_$^{\Delta 11/\Delta 11}$ mice, due to both larger size and increased GAD65 activity of the striosomal compartment (light/dark orange) enhances GABAergic inhibition of the dopamine-producing neurons of SNpc (green). This inhibition of the SNpc neurons (light green) results in an imbalance between the direct and indirect pathways by decreasing both the activation of D1-MSN (D1) and the inhibition of D2-MSN (D2). GPI, internal globus pallidus; GPE, external globus pallidus; SNpr, substantia nigra pars reticulata; SNpc, substantia nigra pars compacta; STN, subthalamic nucleus.

striosome and matrix compartments have been implicated in several movement disorders [Huntington's disease, Parkinson's disease, dyskinesia, dystonia, for a review see (53)], psychostimulant-induced motor stereotypies (58,59), and more recently in anxiety disorders (60). Corticostriatal path targeting striosomes also control decision-making under cost-benefit conflict (61,62) and habit formation (63). Our finding that four striosome markers had increased expression associated with excessive self-grooming in _Shank3_$^{\Delta 11/\Delta 11}$ mice, prompted us to explore the compartmental architecture of the striatum in these mice. This investigation was also motivated by previous studies that have shown that enhanced

striosomal activation is highly correlated with increased repetitive behaviours, including self-grooming

in both non-human primates and rodents [for review see (14)]. Moreover, studies by Kuo and Liu (64,65)

have recently suggested that aberrant striatal compartmentation may be involved in ASD.

We first confirmed that GAD65, which had been reported to be a striosome marker in primates,

but not in rats (27) was indeed a striosome marker in the mouse. We then found that both the surface

and GAD65 labelling intensity of the striosome compartment relative to the matrix one are increased in

$Shank3^{\Delta11/\Delta11}$ mice compared to $Shank3^{+/+}$ mice. Whereas the majority of GABA in brain is produced

by the cytosolic isoform of glutamate decarboxylase (GAD67, encoded by $Gad1$), GAD65, which is

anchored to the membrane of synaptic vesicles, can supply GABA in situations of high demand (66,67),

such as stress or fear, and for fine tuning of inhibitory transmission. It is thus possible that an alteration

of striosomes in $Shank3^{\Delta11/\Delta11}$ mice is responsible for excessive self-grooming observed in these mice.

A model for striosome/matrix imbalance as a possible cause of excessive grooming in the Shank3 mutant

mice: It is well established that the striatum and the dopamine-containing nigrostriatal tract control self-

grooming behaviour (14), and striosomal MSN are the predominant striatal population innervating

dopamine neurons of the nigrostriatal tract (68). An imbalance between the striosome and the matrix

compartments may then explain, via a modification of dopamine signalling, the excessive grooming of

$Shank3^{\Delta11/\Delta11}$ mice (**Fig. 4**). Indeed, in contrast to MSN located in the matrix of the striatum, striosomal

MSN projects directly to the dopamine-producing neurons in the SNpc (53,69). Then, SNpc dopamine

neurons project back to the entire dorsal striatum. An increased activity of GAD65 in the striosomes, in

a situation of stress for example, would lead to an inhibition of dopamine release by the SNpc and thus

to a reduced dopamine modulation of both the direct and indirect pathways in the dorsal striatum.

Because dopamine activation has opposite effects on D1- and D2-MSNs by increasing and decreasing

cell excitability, respectively - a decreased dopamine release by SNpc is expected to result in

diminishing D1-MSN activity and boosting MSN-D2 activity. The differential enrichment in up- and

down-regulated DEGs in D2-MSN and D1-MSN could be related to previous observation of a distinct

impairment of LTD in D2-MSNs, but not in D1-MSN (20) in $Shank3$ mutant mice. Importantly, since

striosome cells are generated before matrix cells during development (52) and SHANK3 seems to be

involved in the early stages of neuronal development (15,70,71), the enlargement of the total relative

surface of striosomes observed in *Shank3*^{Δ11/Δ11} mice may be due to a default in the compartmentation

of the striatum during early development.

Conclusions and perspectives

Altogether, the present study precisely describes the effects over time of the *Shank3*^{Δ11} mutation on

mouse social and repetitive behaviours and suggests a critical role of the striatum in excessive self-

grooming, probably through imbalances in two intertwined systems, the striosome/matrix compartments

and the direct/indirect pathways. Although still preliminary, our results suggest that SHANK3

deficiency and, more generally, ASD could integrate the growing list of neurological conditions

implicating an imbalance in the striosome/matrix compartmentation of the dorsal striatum. Future

studies are necessary to understand the mechanisms at the origin of the striosome/matrix imbalance and

consider the possibility of its reversibility.

Conflict of interest statement

The authors have declared that no competing interests exist.

Detailed author contributions:

ATF, ABi, EV, EE and TB conceived the experiments. ATF and EE designed and acquired

behavioural data from cohorts 1 and 2. ATF extracted and dissected brains, extracted RNA from

cohorts 1 and 2. ATF and FM designed and ATF acquired data from smFISH. ATF performed analysis

and statistics of behaviour, PPI and smFISH. ABo, BFi and JFD performed the RNA sequencing. ABi

and ATF performed the analyses of ORA. ATF, BFo and AL performed and analysed the results of

quantitative RT-PCR. EV performed the immunofluorescence experiments, with the help of SC for

brain collection. FdC performed quantitative image analysis. SC and AMLS genotyped the mice. JS,

CR and LRR conducted the experiments on cohort 3. TMB generated the mouse model. ATF, ABi,

EV, BF, FdC, EE and TB wrote the manuscript.

Acknowledgments

The authors thank the members of the Génétique Humaine et Fonctions Cognitives lab for helpful discussions, especially Jean-Pierre Bourgeois for his advice and expertise, Thomas Rolland for his assistance with data analysis and Nathalie Lemière for her assistance with sample collection. The authors thank Jean-Antoine Girault, Véronique Brault, and Sonia Garel for their advice and expertise. This research was supported by Institut Pasteur, the Bettencourt-Schueller Foundation, CNRS, Université de Paris, the Conny-Maeva Charitable Foundation, the Cognacq Jay Foundation, the Fondation de France, the Fondation pour la Recherche Médicale (FRM), the GIS "Autisme et Troubles du Neuro-Développement", the Roger de Spoelberch Foundation, the Eranet-Neuron (ALTRUISM) project, the Laboratory of Excellence GENMED (Medical Genomics) (grant no. ANR-10-LABX-0013 managed by the National Research Agency (ANR) part of the Investment for the Future program), the BioPsy Labex (ANR-10-LABX-BioPsy, ANR-11-IDEX-0004-02), AIMS-2-TRIALS which received support from the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement No 777394 and the Inception program (Investissement d'Avenir grant ANR-16-CONV-0005). We gratefully acknowledge the UtechS Photonic Bioimaging (Imagopole), C2RT, Institut Pasteur, supported by the French National Research Agency (France BioImaging; ANR-10-INSB-04; Investments for the future). The views expressed here are the responsibility of the author(s) only. The EU Commission takes no responsibility for any use made of the information set out.

Bibliography

1. Association American Psychiatric. Diagnostic and Statistical Manual of Mental Disorders [Internet]. American Psychiatric Association; 2013. Available from: http://psychiatryonline.org/doi/book/10.1176/appi.books.9780890425596 https://psychiatryonline.org/doi/book/10.1176/appi.books.9780890425596

2. Leblond CS, Le T-L, Malesys S, Cliquet F, Tabet A-C, Delorme R, et al. Operative list of genes associated with autism and neurodevelopmental disorders based on database review. Mol Cell Neurosci. 2021 Jun;113:103623.

3.   Monteiro P, Feng G. SHANK proteins: Roles at the synapse and in autism spectrum disorder.

Nat Rev Neurosci. 2017 Mar;18(3):147–57.

4.   Phelan K, McDermid HE. The 22q13.3 deletion syndrome (Phelan-McDermid syndrome). Mol

Syndromol. 2012;2(3–5):186–201.

5.   Durand CM, Betancur C, Boeckers TM, Bockmann J, Chaste P, Fauchereau F, et al. Mutations

in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism

spectrum disorders. Nat Genet. 2007 Jan;39(1):25–7.

6.   Leblond CS, Nava C, Polge A, Gauthier J, Huguet G, Lumbroso S, et al. Meta-analysis of

SHANK Mutations in Autism Spectrum Disorders: A Gradient of Severity in Cognitive

Impairments. PLoS Genet. 2014;10(9).

7.   Kohlenberg TM, Trelles MP, McLarney B, Betancur C, Thurm A, Kolevzon A. Psychiatric

illness and regression in individuals with Phelan-McDermid syndrome. J Neurodev Disord

[Internet]. 2020 Feb;12(1). Available from: https://pubmed.ncbi.nlm.nih.gov/32050889/

8.   Rubeis SD, Siper PM, Durkin A, Weissman J, Muratet F, Halpern D, et al. Delineation of the

genetic and clinical spectrum of Phelan-McDermid syndrome caused by SHANK3 point

mutations. Mol Autism. 2018 Dec;9(1):31.

9.   Wang X, Xu Q, Bey AL, Lee Y, Jiang YH. Transcriptional and functional complexity of Shank3

provides a molecular framework to understand the phenotypic heterogeneity of SHANK3

causing autism and Shank3 mutant mice. Mol Autism. 2014;5(1):30.

10. Ferhat A-T, Halbedl S, Schmeisser MJ, Kas MJ, Bourgeron T, Ey E. Behavioural Phenotypes

and Neural Circuit Dysfunctions in Mouse Models of Autism Spectrum Disorder. Adv Anat

Embryol Cell Biol. 2017;224:85–101.

11. Drapeau E, Riad M, Kajiwara Y, Buxbaum JD. Behavioral Phenotyping of an Improved Mouse

Model of Phelan–McDermid Syndrome with a Complete Deletion of the Shank3 Gene. bioRxiv.

2018 Jan;5(3):ENEURO.0046-18.2018.

12. Peça J, Feliciano C, Ting JT, Wang W, Wells MF, Venkatraman TN, et al. Shank3 mutant mice

display autistic-like behaviours and striatal dysfunction. Nature. 2011 Apr;472(7344):437–42.

13. Wang X, McCoy PA, Rodriguiz RM, Pan Y, Je HS, Roberts AC, et al. Synaptic dysfunction and

abnormal behaviors in mice lacking major isoforms of Shank3. Hum Mol Genet. 2011

Aug;20(15):3093–108.

14. Kalueff AV, Stewart AM, Song C, Berridge KC, Graybiel AM, Fentress JC. Neurobiology of

rodent self-grooming and its value for translational neuroscience. Nat Rev Neurosci. 2016

Jan;17(1):45–59.

15. Peixoto RT, Wang W, Croney DM, Kozorovitskiy Y, Sabatini BL. Early hyperactivity and

precocious maturation of corticostriatal circuits in Shank3B â '/â ' mice. Nat Neurosci. 2016;

16. Bey AL, Wang X, Yan H, Kim N, Passman RL, Yang Y, et al. Brain region-specific disruption

of Shank3 in mice reveals a dissociation for cortical and striatal circuits in autism-related

behaviors. Transl Psychiatry. 2018 Dec;8(1):94.

17. Jin C, Lee Y, Kang H, Jeong K, Park J, Zhang Y, et al. Increased ribosomal protein levels and

protein synthesis in the striatal synaptosome of Shank3-overexpressing transgenic mice. Mol

Brain. 2021 Feb 23;14(1):39.

18. Reim D, Distler U, Halbedl S, Verpelli C, Sala C, Bockmann J, et al. Proteomic Analysis of

Post-synaptic Density Fractions from Shank3 Mutant Mice Reveals Brain Region Specific

Changes Relevant to Autism Spectrum Disorder. Front Mol Neurosci. 2017 Feb;10:26.

19. Schoen M, Asoglu H, Bauer HF, Müller H-P, Abaei A, Sauer AK, et al. Shank3 Transgenic and

Prenatal Zinc-Deficient Autism Mouse Models Show Convergent and Individual Alterations of

Brain Structures in MRI. Front Neural Circuits. 2019;13:6.

20. Wang W, Li C, Chen Q, Goes M-SS van der, Hawrot J, Yao AY, et al. Striatopallidal

dysfunction underlies repetitive behavior in Shank3-deficient model of autism. J Clin Invest.

2017 May;127(5):1978–90.

21. Schmeisser MJ, Ey E, Wegener S, Bockmann J, Stempel AV, Kuebler A, et al. Autistic-like

behaviours and hyperactivity in mice lacking ProSAP1/Shank2. Nature. 2012

Apr;486(7402):256–60.

22. R Core Team (R Foundation for Statistical Computing V. R: A language and environment for

statistical computing. 2020; Available from: https://www.r-project.org/

23. Park J, Coddington LT, Dudman JT. Basal Ganglia Circuits for Action Specification. Annu Rev

Neurosci. 2020 Jul 8;43:485–507.

24. Gokce O, Stanley GM, Treutlein B, Neff NF, Camp JG, Malenka RC, et al. Cellular Taxonomy

of the Mouse Striatum as Revealed by Single-Cell RNA-Seq. Cell Rep. 2016 Jul;16(4):1126–37.

25. Montalban E, Giralt A, Taing L, Nakamura Y, Martin C, Pins B de, et al. Translational profiling

of mouse dopaminoceptive neurons reveals a role of PGE2 in dorsal striatum [Internet]. 2020

Sep [cited 2021 Nov 24] p. 2020.09.02.279240. Available from:

https://www.biorxiv.org/content/10.1101/2020.09.02.279240v1

26. Davis MI, Crittenden JR, Feng AY, Kupferschmidt DA, Naydenov A, Stella N, et al. The

cannabinoid-1 receptor is abundantly expressed in striatal striosomes and striosome-dendron

bouquets of the substantia nigra. PloS One. 2018;13(2):e0191436.

27. Lévesque M, Wallman M-J, Parent A. Striosomes are enriched in glutamic acid decarboxylase in

primates. Neurosci Res. 2004 Sep;50(1):29–35.

28.   Rivera A, Cuéllar B, Girón FJ, Grandy DK, de la Calle A, Moratalla R. Dopamine D4 receptors

are heterogeneously distributed in the striosomes/matrix compartments of the striatum. J

Neurochem. 2002 Jan;80(2):219–29.

29.   Sako W, Morigaki R, Nagahiro S, Kaji R, Goto S. Olfactory type G-protein alpha subunit in

striosome-matrix dopamine systems in adult mice. Neuroscience. 2010 Oct 13;170(2):497–502.

30.   Miyamoto Y, Katayama S, Shigematsu N, Nishi A, Fukuda T. Striosome-based map of the

mouse striatum that is conformable to both cortical afferent topography and uneven distributions

of dopamine D1 and D2 receptor-expressing cells. Brain Struct Funct. 2018 Dec;223(9):4275–

91.

31.   Pagani M, Bertero A, Liska A, Galbusera A, Sabbioni M, Barsotti N, et al. Deletion of Autism

Risk Gene Shank3 Disrupts Prefrontal Connectivity. J Neurosci Off J Soc Neurosci. 2019 Jul

3;39(27):5299–310.

32.   Wang X, Bey AL, Katz BM, Badea A, Kim N, David LK, et al. Altered mGluR5-Homer

scaffolds and corticostriatal connectivity in a Shank3 complete knockout model of autism. Nat

Commun. 2016 May;7:11459.

33.   Yoo Y-E, Yoo T, Lee S, Lee J, Kim D, Han H-M, et al. Shank3 Mice Carrying the Human

Q321R Mutation Display Enhanced Self-Grooming, Abnormal Electroencephalogram Patterns,

and Suppressed Neuronal Excitability and Seizure Susceptibility. Front Mol Neurosci.

2019;12:155.

34.   de Chaumont F, Lemière N, Coqueran S, Bourgeron T, Ey E. LMT USV Toolbox, a Novel

Methodological Approach to Place Mouse Ultrasonic Vocalizations in Their Behavioral

Contexts-A Study in Female and Male C57BL/6J Mice and in Shank3 Mutant Females. Front

Behav Neurosci. 2021;15:735920.

35.   Ey E, Torquet N, Chaumont F de, Lévi-Strauss J, Ferhat A-T, Sourd A-ML, et al. Shank2

Mutant Mice Display Hyperactivity Insensitive to Methylphenidate and Reduced Flexibility in

Social Motivation, but Normal Social Recognition. Front Mol Neurosci [Internet]. 2018 Oct;11.

Available from: /pmc/articles/PMC6180161/ /pmc/articles/PMC6180161/?report=abstract

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6180161/

https://www.frontiersin.org/article/10.3389/fnmol.2018.00365/full

36.   Bariselli S, Tzanoulinou S, Glangetas C, Prévost-Solié C, Pucci L, Viguié J, et al. SHANK3

controls maturation of social reward circuits in the VTA. Nat Neurosci. 2016 Jul;19(7):926–34.

37.   Bariselli S, Bellone C. VTA DA neuron excitatory synapses in Shank3 Δex4–9mouse line.

Synapse. 2017;71(6):1–5.

38.   Orefice LL, Mosko JR, Morency DT, Wells MF, Tasnim A, Mozeika SM, et al. Targeting

Peripheral Somatosensory Neurons to Improve Tactile-Related Phenotypes in ASD Models.

Cell. 2019 Aug 8;178(4):867-886.e24.

39.   Lin R, Learman LN, Bangash MA, Melnikova T, Leyder E, Reddy SC, et al. Homer1a regulates

Shank3 expression and underlies behavioral vulnerability to stress in a model of Phelan-

McDermid syndrome. Cell Rep. 2021 Nov 16;37(7):110014.

40.   Qin L, Ma K, Wang Z-J, Hu Z, Matas E, Wei J, et al. Social deficits in Shank3-deficient mouse

models of autism are rescued by histone deacetylase (HDAC) inhibition. Nat Neurosci [Internet].

2018 Mar; Available from: http://www.ncbi.nlm.nih.gov/pubmed/29531362

http://www.nature.com/articles/s41593-018-0110-8

41.   Moutin E, Sakkaki S, Compan V, Bouquier N, Giona F, Areias J, et al. Restoring glutamate

receptosome dynamics at synapses rescues autism-like deficits in Shank3-deficient mice. Mol

Psychiatry. 2021 Jul 30;1–14.

42.   Chen Y-W, Lin H-C, Ng M-C, Hsiao Y-H, Wang C-C, Gean P-W, et al. Activation of mGluR2/3

underlies the effects of N-acetylcystein on amygdala-associated autism-like phenotypes in a

valproate-induced rat model of autism. Front Behav Neurosci. 2014 Jun;8:219.

43.   Boeckers TM, Inter C, Smalla KH, Kreutz MR, Bockmann J, Seidenbecher C, et al. Proline-rich

synapse-associated proteins ProSAP1 and ProSAP2 interact with synaptic proteins of the

SAPAP/GKAP family. Biochem Biophys Res Commun. 1999;264(1):247–52.

44.   Gogolla N, Takesian AE, Feng G, Fagiolini M, Hensch TK. Sensory integration in mouse insular

cortex reflects GABA circuit maturation. Neuron. 2014 Aug 20;83(4):894–905.

45.   Litvin Y, Phan A, Hill MN, Pfaff DW, McEwen BS. CB1 receptor signaling regulates social

anxiety and memory. Genes Brain Behav. 2013;12(5):479–89.

46.   Tallett AJ, Blundell JE, Rodgers RJ. Endogenous opioids and cannabinoids: system interactions

in the regulation of appetite, grooming and scratching. Physiol Behav. 2008 Jun 9;94(3):422–31.

47.   Terzian AL, Drago F, Wotjak C, Micale V. The Dopamine and Cannabinoid Interaction in the

Modulation of Emotions and Cognition: Assessing the Role of Cannabinoid CB1 Receptor in

Neurons Expressing Dopamine D1 Receptors. Front Behav Neurosci. 2011;5:49.

48.   Folkes OM, Báldi R, Kondev V, Marcus DJ, Hartley ND, Turner BD, et al. An

endocannabinoid-regulated basolateral amygdala-nucleus accumbens circuit modulates

sociability. J Clin Invest. 2020 Apr 1;130(4):1728–42.

49.   Zou M, Liu Y, Xie S, Wang L, Li D, Li L, et al. Alterations of the endocannabinoid system and

its therapeutic potential in autism spectrum disorder. Open Biol. 2021 Feb;11(2):200306.

50.   Pelosi A, Menardy F, Popa D, Girault J-A, Hervé D. Heterozygous Gnal Mice Are a Novel

Animal Model with Which to Study Dystonia Pathophysiology. J Neurosci. 2017 Jun

28;37(26):6253–67.

51.   Lee B, Zhang Y, Kim Y, Kim S, Lee Y, Han K. Age-dependent decrease of GAD65/67 mRNAs

but normal densities of GABAergic interneurons in the brain regions of Shank3 -overexpressing

manic mouse model. Neurosci Lett. 2017 May;649:48–54.

52.   Hagimoto K, Takami S, Murakami F, Tanabe Y. Distinct migratory behaviors of striosome and

matrix cells underlying the mosaic formation in the developing striatum. J Comp Neurol. 2017

Mar 1;525(4):794–817.

53.   Crittenden J, Graybiel A. Basal Ganglia Disorders Associated with Imbalances in the Striatal

Striosome and Matrix Compartments. Front Neuroanat. 2011;5:59.

54.   Prager EM, Plotkin JL. Compartmental function and modulation of the striatum. J Neurosci Res.

2019 Dec;97(12):1503–14.

55.   Prager EM, Dorman DB, Hobel ZB, Malgady JM, Blackwell KT, Plotkin JL. Dopamine

Oppositely Modulates State Transitions in Striosome and Matrix Direct Pathway Striatal Spiny

Neurons. Neuron. 2020 Dec 23;108(6):1091-1102.e5.

56.   Jaquins-Gerstl A, Nesbitt KM, Michael AC. In vivo evidence for the unique kinetics of evoked

dopamine release in the patch and matrix compartments of the striatum. Anal Bioanal Chem.

2021 Nov;413(27):6703–13.

57.   Friedman A, Homma D, Bloem B, Gibb LG, Amemori K-I, Hu D, et al. Chronic Stress Alters

Striosome-Circuit Dynamics, Leading to Aberrant Decision-Making. Cell. 2017 Nov

16;171(5):1191-1205.e28.

58.   Canales JJ, Graybiel AM. A measure of striatal function predicts motor stereotypy. Nat

Neurosci. 2000 Apr;3(4):377–83.

59.   Murray RC, Logan MC, Horner KA. Striatal patch compartment lesions reduce stereotypy

following repeated cocaine administration. Brain Res. 2015 Aug 27;1618:286–98.

60.   Karunakaran KB, Amemori S, Balakrishnan N, Ganapathiraju MK, Amemori K. Generalized

and social anxiety disorder interactomes show distinctive overlaps with striosome and matrix

interactomes. Sci Rep. 2021 Sep 15;11(1):18392.

61.   Friedman A, Homma D, Gibb LG, Amemori K, Rubin SJ, Hood AS, et al. A Corticostriatal Path

Targeting Striosomes Controls Decision-Making under Conflict. Cell. 2015 Jun 4;161(6):1320–

33.

62.   Friedman A, Hueske E, Drammis SM, Toro Arana SE, Nelson ED, Carter CW, et al. Striosomes

Mediate Value-Based Learning Vulnerable in Age and a Huntington's Disease Model. Cell.

2020 Nov 12;183(4):918-934.e49.

63.   Nadel JA, Pawelko SS, Scott JR, McLaughlin R, Fox M, Ghanem M, et al. Optogenetic

stimulation of striatal patches modifies habit formation and inhibits dopamine release. Sci Rep.

2021 Oct 6;11(1):19847.

64.   Kuo H-Y, Liu F-C. Valproic acid induces aberrant development of striatal compartments and

corticostriatal pathways in a mouse model of autism spectrum disorder. FASEB J.

2017;31(10):4458–71.

65.   Kuo H-Y, Liu F-C. Pathological alterations in striatal compartments in the human brain of

autism spectrum disorder. Mol Brain. 2020 Dec;13(1):83.

66.   Esclapez M, Tillakaratne NJ, Kaufman DL, Tobin AJ, Houser CR. Comparative localization of

two forms of glutamic acid decarboxylase and their mRNAs in rat brain supports the concept of

functional differences between the forms. J Neurosci Off J Soc Neurosci. 1994 Mar;14(3 Pt

2):1834–55.

67.   Kaufman DL, Houser CR, Tobin AJ. Two forms of the gamma-aminobutyric acid synthetic

enzyme glutamate decarboxylase have distinct intraneuronal distributions and cofactor

interactions. J Neurochem. 1991 Feb;56(2):720–3.

68.   Watabe-Uchida M, Zhu L, Ogawa SK, Vamanrao A, Uchida N. Whole-brain mapping of direct

inputs to midbrain dopamine neurons. Neuron. 2012 Jun 7;74(5):858–73.

69.   McGregor MM, McKinsey GL, Girasole AE, Bair-Marshall CJ, Rubenstein JLR, Nelson AB.

Functionally Distinct Connectivity of Developmentally Targeted Striosome Neurons. Cell Rep.

2019 Nov 5;29(6):1419-1428.e5.

70.   Jaramillo TC, Xuan Z, Reimers JM, Escamilla CO, Liu S, Powell CM. Early Restoration of

Shank3 Expression in Shank3 Knock-Out Mice Prevents Core ASD-Like Behavioral

Phenotypes. eNeuro. 2020 Jun;7(3):ENEURO.0332-19.2020.

71.   Huang G, Chen S, Chen X, Zheng J, Xu Z, Doostparast Torshizi A, et al. Uncovering the

Functional Link Between SHANK3 Deletions and Deficiency in Neurodevelopment Using

iPSC-Derived Human Neurons. Front Neuroanat. 2019 Mar 13;13:23.

Shank3$^{+/+}$ *Shank3*$^{\Delta 11/\Delta 11}$

excitatory glutamate   dopamine activation

inhibitory GABA   dopamine inhibition