1 **FoodMicrobionet v4: a large, integrated, open and transparent database for food bacterial**
2 **communities.**
3

4 Eugenio Parente, Teresa Zotta, Annamaria Ricciardi
5
6 Scuola di Scienze Agrarie, Forestali, Alimentari ed Ambientali, Università degli Studi della
7 Basilicata, Potenza, Italy
8
9
10 **Corresponding author**
11
12 Prof. Eugenio Parente
13
14 Scuola di Scienze Agrarie, Forestali, Alimentari ed Ambientali,
15 Università degli Studi della Basilicata,
16 Viale dell'Ateneo Lucano 10, 85100 Potenza, Italy
17
18 Email: eugenio.parente@unibas.it
19 Tel.: +390971205561
20
21

FoodMicrobionet v4: a large, integrated, open and transparent database for food bacterial communities.

Eugenio Parente, Teresa Zotta, Annamaria Ricciardi

Abstract

With the availability of high-throughput sequencing techniques our knowledge of the structure and dynamics of food microbial communities has made a quantum leap. This knowledge is however dispersed in a large number of papers and hard data are only partly available through powerful on-line databases and tools such as QIITA, MGnify and the Integrated Microbial Next Generation Sequencing platform, whose annotation is, however, not optimized for foods.

Here, we present the 4[th] iteration of FoodMicrobionet, a database of the composition of bacterial microbial communities of foods and food environments. With 180 studies and 10,151 samples belonging to 8 major food groups FoodMicrobionet 4.1.2 is arguably the largest and best annotated database on food bacterial communities. This version includes 1,684 environmental samples and 8,467 food samples, belonging to 16 L1 categories and 196 L6 categories of the EFSA FoodEx2 classification and is approximately 4 times larger than previous version (3.1, https://doi.org/10.1016/j.ijfoodmicro.2019.108249). Both FoodMicrobionet and related data and software conform to FAIR (Findable, Accessible, Interoperable, Reusable/Reproducible) criteria for scientific data and software and are freely available on public repositories (GitHub, Mendeley data).

Even if FoodMicrobionet does not have the sophistication of QIITA, IMNGS and MGnify, we

feel that this iteration, due to its size and diversity, provides a valuable asset for both the

scientific community and industrial and regulatory stakeholders.

Key words: amplicon targeted high-throughput sequencing; 16S metagenomics; bacterial

microbiota; database

1. Introduction.

In less than 15 years, next generation sequencing has added several layers of complexity to

the study of food microbiomes. At the time of writing this article (January 2022) searches

with the key words "microbiome" OR "microbiota" in Scopus and in Web of Science

retrieved 118,416 and 127,424 hits, respectively. The number of papers using amplicon

targeted next generation sequencing to study the structure of food bacterial communities

has been rising exponentially since the first pioneering publications (Humblot and Guyot,

2009; Roh et al., 2010) (see Figure 1).

The evolution of -omic approaches to the study of the food microbiome and the

methodological challenges in wet laboratory methods, sequencing and bioinformatic and

statistical approaches have been reviewed multiple times (Bokulich et al., 2020; De Filippis

et al., 2018; Pollock et al., 2018). While amplicon targeted approaches are still prevalent,

with the decreasing cost of sequencing and the increase in power of sequencing platforms

and of computational tools the number of studies using metagenomic or multi-omics

approaches is rapidly increasing (Yap et al., 2021).

Figure 1. Cumulative number of papers using metataxonomic or metagenomic approaches

for the study of food microbial communities. The data derive from a manually curated

search on Web of Science[TM].

Shotgun metagenomics offers a significantly higher resolution (down to the strain and

perhaps to single nucleotide polymorphism variants (Hildebrand, 2021) and paves the way

to accurate source tracking for contamination (De Filippis et al., 2020) and to a new

paradigm in food safety (Kovac, 2019). However, amplicon targeted approaches still have

the power to describe, in a sensitive and cost-effective way, the structure of food microbial

communities down to the genus level or below (Callahan et al., 2016a).

More than 640 papers describing the food microbiota have been published since 2009.

Exploitation of this wealth of information for metastudies or for the development of

descriptive or predictive tools requires FAIR (findable, accessible, interoperable,

reusable/reproducible: Lamprecht et al., 2020; Wilkinson et al., 2016) data and software.

Three large on line repositories on microbiome data have appeared in recent years: IMNGS

(Lagkouvardos et al., 2016), QIITA (Gonzalez et al., 2018) and MGnify (Mitchell et al., 2019).

The Integrated Microbial Next Generation Sequencing platform directly accesses 16S rRNA

gene targeted next generation sequencing data on NCBI Short Read Archive and provides

powerful tools for searching for taxa or sequences or for performing analysis of user

deposited 16S sequences (Lagkouvardos et al., 2016).

MGnify hosts metagenomic, metatranscriptomics and metataxonomic datasets which are

either contributed by users or obtained from public repositories and offers powerful

platforms for data analysis but does not allow the integration of data from different studies

(Mitchell et al., 2019).

QIITA offers a powerful suite of tools for the analysis of sequences of public and private

datasets, and allows the search and integration of data from different studies (Gonzalez et

al., 2018).

At the time of writing of this paper (January 2022) MGnify included 3,696 public studies, and

325,323 public samples, but only 83 studies/2,805 samples on food biomes

(https://www.ebi.ac.uk/metagenomics/) while QIITA included 620 public studies and

303,313 public samples but only a very limited number of public studies on food biomes

(https://qiita.ucsd.edu/stats/). The data and interfaces for all these tools respond well to

FAIR principles for scientific data (Wilkinson et al., 2016) but, unfortunately, the structure of

the metadata for studies and samples is not optimized for foods.

Some years ago, we created a database for metataxonomic studies on food bacterial

communities, FoodMicrobionet (Parente et al., 2019, 2016), whose main strength is the

annotation system for studies and samples, based on the hierarchical classification of foods

developed by the European Food Safety Authority, FoodEx 2(EFSA, 2015). Here we describe

the structure and new features of the latest version of the database.

2. Materials and methods.

The procedure used to add data to FoodMicrobionet has not changed significantly since the

last version (Parente et al., 2019) and it is represented schematically in Figure 2.

Figure 2. Schematic workflow showing how raw sequences and their metadata and

additional information from the scientific literature are used to assemble data for

FoodMicrobionet.

With the exception of 33 studies (out of 180), which were originally obtained by research

partners as abundance tables for taxa and sample metadata tables (Parente et al., 2016), all

studies in FoodMicrobionet are obtained by reprocessing raw sequence data from NCBI

Short Read Archive (SRA), and by using metadata from SRA and from the papers.

Processing of sequences is carried out in R (R Core Team, 2021) using a modified version of

the Bioconductor pipeline for amplicon targeted sequence analysis, using DADA2 for

Amplicon Sequence Variant (ASV) inference and SILVA v138.1 for taxonomic assignment

(Callahan et al., 2016a; Callahan et al., 2016b). The analysis results in the production of

phyloseq (McMurdie and Holmes, 2013) objects which are further processed by R scripts

and imported in Microsoft Excel for further manual editing of study and sample metadata.

Finally, a R script is used to process Excel tables and for quality control checks. All scripts

used for processing are publicly available in the FoodMicrobionet GitHub repository

(https://github.com/ep142/FoodMicrobionet).

The structure of the database is schematically shown in Figure 3.

Studies, samples and taxa have all a rich metadata structure. The structure of the tables is

described in detail on GitHub

(https://github.com/ep142/FoodMicrobionet/blob/master/FoodMicrobionet_tablespecs.m

d), on Mendeley data (https://data.mendeley.com/datasets/8fwwjpm79y/5) and in

supplementary material.

Recent additions compared to version 3 (Parente et al., 2019) include four fields which

complete information for bioinformatic processing and one field for geolocation for studies,

geolocation information for samples, and two further reference tables (primers and the

FoodEx2 Exposure Hierarchy classification (EFSA, 2015).

Figure 3. Schematic representation of the structure of FoodMicrobionet tables (Study,

Primers, Samples, FoodEx2, Taxa, Edges) and their relationship with phyloseq (McMurdie

and Holmes, 2013) objects and external databases.

FoodMicrobionet is available either as a R list, which allows experienced programmers to

run their searches and analyses in the most flexible and sensitive way and as an interactive

Shiny app (Parente et al., 2019) which requires minimum installation and configuration and

allows users to perform searches using a large number of criteria, to perform aggregation of

samples and taxa, to rapidly reach external resources using hyperlinks, to export data in a

variety of formats, and to obtain and save graphs and tables (Parente et al., 2019).

Use cases and example workflows have been illustrated in a previous work (Parente et al.,

2019) and we have tried to demonstrate this approach in a series of proof-of-concept

metastudies (Parente et al., 2020, 2021; Zotta et al., 2021). The code for generating graphs

and statistical analyses is fully reproducible and reusable

(https://data.mendeley.com/datasets/8fwwjpm79y/4;

https://github.com/ep142/MAN_in_cheese) and allows to reproduce the figures and tables

when a new version of FoodMicrobionet is published.

3. Results and discussion.

3.1 FoodMicrobionet facts and figures.

FoodMicrobionet v 4.1.2 has grown significantly since its last iteration (Parente et al., 2019;

Figure 4): the number of studies has grown from 44 to 180, and the number of samples from

2,237 to 10,151.

The growth of FoodMicrobionet reflects the growth of published studies and the

distribution of target gene regions reflects the shift in technology in Next Generation

Sequencing, from the defunct Roche 454 platform (with most studies targeting the V1-V3

region) to Illumina platforms, with V4 and V3-V4 regions of the 16S rRNA gene as the most

frequent targets (Figure 4, Supplementary Table 1).

The number of reads for sample after processing is also saved in the database

(Supplementary Figure 1): 70% samples have >10^4 reads after processing, but a significant

number of samples in studies targeting the V4 region has >5×10^4 reads. This makes the

detection of rare components of the microbiota possible.

Figure 4. Cumulative distribution of studies in FoodMicrobionet 4.1.2, by year and by 16S

rRNA gene target region.

The variety of food groups and food types in FoodMicrobionet is also very large: samples in

FMBN belong to 16 major food groups (L1 level of FoodEx2 exposure classification;

Supplementary Table 2). Approximately 80% of samples belong to the first three categories

(Milk and dairy products, Meat and meat products, Vegetables and vegetable products).

Samples in FMBN are further classified using levels L4 and L6 of the FoodEx2 exposure

classification, and additional fields (which allow to identify raw products, intermediates or

finished products, the level of thermal treatment and the occurrence of spoilage and/or

fermentation) allow a finer classification. Samples in FMBN belong to 109 L4 food groups

and 197 L6 food groups. There are 163 individual food types, and, combining further

information on samples (nature, heat treatment, spoilage/fermentation), there are 316

combinations (Parente et al., 2019). This number and variety of samples makes

FoodMicrobionet the largest database of metataxonomic data on food bacterial

communities, with significantly more samples compared to QIITA and MGnify.

FoodMicrobionet stores samples with 51 countries, but the 90% of samples are from 14

countries only (Supplementary Figure 2). This does not reflect the distribution of samples

and studies in published studies but rather, the distribution of those which are available in

NCBI SRA.

There are currently 9,098 taxa in the taxa table of FoodMicrobionet. Taxa have a unique

numeric and text identifier and may represent ASV identified at the species (4,497, 49.4% of

taxa) or genus level (3,259, 35.8%) or above using the DADA2 assignTaxnomy() and

assignSpecies() functions. The % of the taxa and sequences identified at the genus level or

below varies by study, depending on the quality and length of sequences and on the gene

target. Length of reads (in bp) in FoodMicrobionet studies varies between 150 and 610 bp

(median 422). Although one may question the value of taxonomic assignment at the species

level, especially for the shortest reads (Callahan et al., 2016a; Edgar, 2018; Johnson et al.,

2019; Meola et al., 2019; Pollock et al., 2018), due to the detailed information provided for

both studies and samples (gene target and region, the number of issues observed during

processing of raw sequences) and to the possibility of accessing to the processed sequences

(ASV) in phyloseq objects, users of FoodMicrobionet can make informed decision on how

and when taxonomic units should be combined at a level higher than the species, an

operation which is easily performed with the ShinyFMBN app (Parente et al., 2019).

FoodMicrobionet is fully connected to external databases: external links (as dynamically

built Uniform Resource Locators) in the study, sample and taxa tables allow rapid access to

external taxonomic databases (NCBI taxonomy, the List of Prokaryotic Names with Standing

in Nomenclature and the Florilege database, Falentin et al., 2017

http://migale.jouy.inra.fr/Florilege/#&about) and to the scientific literature (via DOI). Using

NCBI SRA Study accession number it is possible to access fine grained data on ASV sequence

and abundance stored in the phyloseq objects obtained from processing the raw sequences

(these are not public but available on request).

Is FoodMicrobionet FAIR?

FoodMicrobionet data and software are free (both are under MIT licence

https://opensource.org/licenses/MIT), open and highly reusable and support analysis

protocols which are reproducible. We have done our best (given the time and resource

constraints) to conform as closely as possible to criteria for FAIR (Findable, Accessible,

Interoperable, Reusable/Reproducible) data and software sharing (Lamprecht et al., 2020;

Wilkinson et al., 2016).

Both the database and the software are Findable (using searches on Google, Mendeley data,

or GitHub, for example) and deposited on permanent repositories (Mendeley data, GitHub)

with unique identifiers. Even if data within FoodMicrobionet may not be directly findable by

automated machine searches (and insofar they are not machine operable), since the

database is not available on line except in the form of R lists or tab delimited text files, the

richness of contextual metadata for all the main tables (studies, samples, taxa) makes it

possible to devise precisely targeted searches.

FoodMicrobionet is Accessible through the above-mentioned repositories and through our

website, and are confident that enough metadata are provided in these repositories to

easily reach the resources. In terms of user accessibility (which is not a criterion in FAIR

principles), we have done our best to make the resource accessible to both expert (the

database in the form of a R list can be used for fine-tuned searches and analyses using R

scripts) and moderately expert users. The latter can, with a minimum effort, download and

install the R Shiny app, ShinyFMBN (Parente et al., 2019), which, once launched in R, allows

easy access via an interactive and intuitive interface. A detailed manual for the app is

available on Mendeley data (https://data.mendeley.com/datasets/8fwwjpm79y/4).

Although the app could be easily deployed on RStudio Shiny apps server

(https://www.shinyapps.io), we feel that this would make the use unnecessarily slow.

FoodMicrobionet is fully Interoperable. The sample classification is based on a robust

hierarchical classification, FoodEx2, rather than on arbitrary keywords and dynamic links are

created in the studies, samples and taxa tables to reach a number of other databases.

Conversely, other databases like Florilege might quite simply create new accessions using

FoodMicrobionet and metadata in FMBN can, in principle, be used to populate QIITA and

MGnify, by linking studies and samples via the SRA accession numbers.

Data and products of search results are highly Reusable, for the same reasons. In addition,

the objects exported by the app are in formats which are compatible with metataxonomic

analysis pipelines (like phyloseq and ShinyPhyloseq: (McMurdie and Holmes, 2015, 2013);

MicrobiomeAnalyst: (Chong et al., 2020; Dhariwal et al., 2017); CoNet: (Faust and Raes,

2016); graph visualization and analysis software like Cytoscape and iGraph: (Csardi et al.,

2006; Shannon et al., 2003), microbial association network inference tools: (Kurtz et al.,

2015; Peschel et al., 2021).

5. Conclusions.

Even if FoodMicrobionet does not have the sophistication of QIITA, IMNGS and MGnify, we

feel that this iteration, due to its size and diversity, provides a significant resource for both

the scientific community and industrial and regulatory stakeholders. Scientists can access

and use a variety of stand-alone or online software tools and ShinyFMBN to compare their

own results with literature results, carry out metastudies to answer a variety of scientific

questions, build reproducible analysis workflows, get quantitative data on the

ecology/distribution of bacteria of interest, use the database as an entry point for further

searches in other databases. The size of this version, which includes >9×10^5 taxon/sample

relationships, might even allow the machine learning approaches to predict contamination

patterns of food. The ability to rapidly retrieve information on prevalence/abundance of

taxon in different foods and on the structure of microbial communities in different food

types may be useful to both the industry and regulatory agencies. Information on the

distribution of beneficial genera and species may find use for regulatory purposes (for

example to facilitate studies on the distribution of beneficial microorganisms to evaluate

their inclusion in the Qualified Presumption of Safety). The fine-grained data on the

structure of microbial communities for a large variety of raw materials, foods, food

environments may be useful for both process and product development purposes to

identify spoilage or contamination patterns, or for the design of microbiome-based starters.

We are committed to keep adding data to FoodMicrobionet, but the openness and

transparency of its software and documentation technically allows any interested party to

create new versions of the database or to significantly improve its structure and

functionality.

CRediT author statement

Eugenio Parente: Conceptualization, Methodology, Software, Writing- Original draft

preparation. **Annamaria Ricciardi** Data curation, Writing - Reviewing and Editing. **Teresa**

Zotta: Data curation, Writing - Reviewing and Editing.

Acknowledgements.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

Bokulich, N.A., Ziemski, M., Robeson, M.S., Kaehler, B.D., 2020. Measuring the microbiome:

    Best practices for developing and benchmarking microbiomics methods. Comput. Struct.

    Biotechnol. J. 18, 4048–4062. https://doi.org/10.1016/j.csbj.2020.11.049

Callahan, B. J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016a.

    DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Met. 13,

    581–583. https://doi.org/10.1038/nmeth.3869

Callahan, B. J., Sankaran, K., Fukuyama, J.A., McMurdie, P.J., Holmes, S.P., 2016b.

    Bioconductor workflow for microbiome data analysis: from raw reads to community

    analyses. F1000 Res. 5, 1492. https://doi.org/10.12688/f1000research.8986.2

Chong, J., Liu, P., Zhou, G., Xia, J., 2020. Using MicrobiomeAnalyst for comprehensive

    statistical, functional, and meta-analysis of microbiome data. Nat. Protoc. 1–23.

    https://doi.org/10.1038/s41596-019-0264-1

Csardi, G., Nepusz, T., 2006. The igraph software package for complex network research,

    InterJournal, Complex Systems 1695. https://igraph.org

De Filippis, F., Parente, E., Ercolini, D., 2018. Recent past, present, and future of the

    food microbiome. Annu. Rev. Food Sci. Technol. 9, 589–608.

    https://doi.org/10.1146/annurev-food-030117-012312

De Filippis, F., Valentino, V., Alvarez-Ordóñez, A., Cotter, P.D., Ercolini, D., 2020.

Environmental microbiome mapping as a strategy to improve quality and safety in the

food industry. Curr. Opin. Food Sci. 38, 168–176.

https://doi.org/10.1016/j.cofs.2020.11.012

Dhariwal, A., Chong, J., Habib, S., King, I.L., Agellon, L.B., Xia, J., 2017. MicrobiomeAnalyst: a

web-based tool for comprehensive statistical, visual and meta-analysis of microbiome

data. Nucl. Ac. Res. 45, W180–W188. https://doi.org/10.1093/nar/gkx295

Edgar, R.C., 2018. Accuracy of taxonomy prediction for 16S rRNA and fungal ITS sequences.

PeerJ 6, e4652. https://doi.org/10.7717/peerj.4652

E.F.S.A., 2015. The food classification and description system FoodEx 2 (revision 2). EFSA

Supporting Publications 1–90. https://doi.org/10.2903/sp.efsa.2015.en-804

Falentin, H., Chaix, E., Dérozier, S., Weber, M., Buchin, S., Dridi, B., Deutsch, S.-M., Valence-

Bertel, F., Casaregola, S., Renault, P., Champomier-Verges, M.-C., Thierry, A., Zagorec, M.,

Irlinger, F., Delbes, C., Aubin, S., Bessières, P., Loux, V., Bossy, R., Dibie, J., Sicard, D.,

Nédellec, C. (2017, October). Florilege: a database gathering microbial phenotypes of

food interest. In 4th International Conference on Microbial Diversity 2017, Bari, ITA

(2017-10-24 - 2017-10-26). http://migale.jouy.inra.fr/Florilege/#&about

Faust, K., Raes, J., 2016. CoNet app: inference of biological association networks using

Cytoscape. F1000 Res. 5, 1519–14. https://doi.org/10.12688/f1000research.9050.2

Gonzalez, A., Navas-Molina, J.A., Kosciolek, T., McDonald, D., Vázquez-Baeza, Y., Ackermann,

G., DeReus, J., Janssen, S., Swafford, A.D., Orchanian, S.B., Sanders, J.G., Shorenstein, J.,

Holste, H., Petrus, S., Robbins-Pianka, A., Brislawn, C.J., Wang, M., Rideout, J.R., Bolyen,

E., Dillon, M., Caporaso, J.G., Dorrestein, P.C., Knight, R., 2018. QIITA: rapid, web-enabled

microbiome meta-analysis. Nat. Met. 15, 1–6. https://doi.org/10.1038/s41592-018-0141-

9

Hildebrand, F., 2021. Ultra-resolution metagenomics: when enough Is not enough.

mSystems 6, e00881-21. https://doi.org/10.1128/msystems.00881-21

Humblot, C., Guyot, J.P., 2009. Pyrosequencing of Tagged 16S rRNA g   ene a   mplicons

for r   apid d   eciphering of the m   icrobiomes of f   ermented f   oods s   uch as p

earl m   illet s   lurries. Appl. Environ. Microbiol. 75, 4354–4361.

https://doi.org/10.1128/aem.00451-09

Johnson, J.S., Spakowicz, D.J., Hong, B.-Y., Petersen, L.M., Demkowicz, P., Chen, L., Leopold,

S.R., Hanson, B.M., Agresta, H.O., Gerstein, M., Sodergren, E., Weinstock, G.M., 2019.

Evaluation of 16S rRNA gene sequencing for species and strain-level microbiome analysis.

Nat. Commun. 10, 5029. https://doi.org/10.1038/s41467-019-13036-1

Kovac, J., 2019. Precision food safety: a paradigm shift in detection and control of foodborne

pathogens. mSystems 4, e00164-19. https://doi.org/10.1128/msystems.00164-19

Kurtz, Z.D., Müller, C.L., Miraldi, E.R., Littman, D.R., Blaser, M.J., Bonneau, R.A., 2015. Sparse

and compositionally robust inference of microbial ecological networks. PLoS Comput.

Biol. 11, e1004226. https://doi.org/10.1371/journal.pcbi.1004226

Lagkouvardos, I., Joseph, D., Kapfhammer, M., Giritli, S., Horn, M., Haller, D., Clavel, T., 2016.

IMNGS: A comprehensive open resource of processed 16S rRNA microbial profiles for

ecology and diversity studies. Sci. Rep. 6, 33721. https://doi.org/10.1038/srep33721

Lamprecht, A.-L., Garcia, L., Kuzak, M., Martinez, C., Arcila, R., Pico, E.M.D., Angel, V.D.D.,

Sandt, S. van de, Ison, J., Martinez, P.A., McQuilton, P., Valencia, A., Harrow, J.,

Psomopoulos, F., Gelpi, J.L., Hong, N.C., Goble, C., Capella-Gutierrez, S., 2020. Towards

FAIR principles for research software. Lect. Notes Comput. Sc. 3, 37–59.

https://doi.org/10.3233/ds-190026

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. PLoS One 8, e61217.

https://doi.org/10.1371/journal.pone.0061217

McMurdie, P.J., Holmes, S., 2015. Shiny-phyloseq: Web application for interactive

microbiome analysis with provenance tracking. Bioinformatics 31, 282–283.

https://doi.org/10.1093/bioinformatics/btu616

Meola, M., Rifa, E., Shani, N., Delbès, C., Berthoud, H., Chassard, C., 2019. DAIRYdb: a

manually curated reference database for improved taxonomy annotation of 16S rRNA

gene sequences from dairy products. BMC Genom. 20, 560.

https://doi.org/10.1186/s12864-019-5914-8

Mitchell, A.L., Almeida, A., Beracochea, M., Boland, M., Burgin, J., Cochrane, G., Crusoe,

M.R., Kale, V., Potter, S.C., Richardson, L.J., Sakharova, E., Scheremetjew, M.,

Korobeynikov, A., Shlemov, A., Kunyavskaya, O., Lapidus, A., Finn, R.D., 2019. MGnify: the

microbiome analysis resource in 2020. Nucl. Ac. Res. 48, D570–D578.

https://doi.org/10.1093/nar/gkz1035

Parente, E., Cocolin, L., De Filippis, F., Zotta, T., Ferrocino, I., O'sullivan, O., Neviani, E.,

Angelis, M.D., Cotter, P.D., Ercolini, D., 2016. FoodMicrobionet: A database for the

visualisation and exploration of food bacterial communities based on network analysis.

Int. J. Food Microbiol. 219, 28–37. https://doi.org/10.1016/j.ijfoodmicro.2015.12.001

Parente, E., De Filippis, F., Ercolini, D., Ricciardi, A., Zotta, T., 2019. Advancing integration of

data on food microbiome studies: FoodMicrobionet 3.1, a major upgrade of the

FoodMicrobionet database. Int. J. Food Microbiol. 305, 108249.

https://doi.org/10.1016/j.ijfoodmicro.2019.108249

Parente, E., Ricciardi, A., Zotta, T., 2020. The microbiota of dairy milk: a review. Int. Dairy J.

107, 104714. https://doi.org/10.1016/j.idairyj.2020.104714

Parente, E., Zotta, T., Ricciardi, A., 2021. Microbial association networks in cheese: a meta-

analysis. BiorXiv 2021.07.21.453196. https://doi.org/10.1101/2021.07.21.453196

Peschel, S., Müller, C.L., Mutius, E. von, Boulesteix, A.-L., Depner, M., 2021. NetCoMi:

network construction and comparison for microbiome data in R. Brief. Bioinform.

https://doi.org/10.1093/bib/bbaa290

Pollock, J., Glendinning, L., Wisedchanwet, T., Watson, M., 2018. The madness of

microbiome: attempting to find consensus "best practice" for 16S microbiome studies.

Appl. Environ. Microbiol. 84, 3225. https://doi.org/10.1128/aem.02627-17

R Core Team (2021). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Roh, S.W., Kim, K.-H., Nam, Y.-D., Chang, H.-W., Park, E.-J., Bae, J.-W., 2010. Investigation of

archaeal and bacterial diversity in fermented seafood using barcoded pyrosequencing.

ISME J. 4, 1–16. https://doi.org/10.1038/ismej.2009.83

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,

Schwikowski, B., Ideker, T., 2003. Cytoscape: a software environment for integrated

models of biomolecular interaction networks. Genome Res. 13, 2498–2504.

https://doi.org/10.1101/gr.1239303

Wilkinson, M.D., Dumontier, M., Aalbersberg, Ij.J., Appleton, G., Axton, M., Baak, A.,

Blomberg, N., Boiten, J.-W., Santos, L.B. da S., Bourne, P.E., Bouwman, J., Brookes, A.J.,

Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C.T., Finkers, R., Gonzalez-

Beltran, A., Gray, A.J.G., Groth, P., Goble, C., Grethe, J.S., Heringa, J., Hoen, P.A.C. 't,

Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S.J., Martone, M.E., Mons, A., Packer, A.L.,

Persson, B., Rocca-Serra, P., Roos, M., Schaik, R. van, Sansone, S.-A., Schultes, E.,

Sengstag, T., Slater, T., Strawn, G., Swertz, M.A., Thompson, M., Lei, J. van der, Mulligen,

E. van, Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., Mons, B.,

2016. The FAIR Guiding Principles for scientific data management and stewardship. Sci.

Data 3, 160018. https://doi.org/10.1038/sdata.2016.18

Yap, M., Ercolini, D., Álvarez-Ordóñez, A., O'Toole, P.W., O'Sullivan, O., Cotter, P.D., 2021.

Next-generation food research: use of meta-omic approaches for characterizing

microbial communities along the food chain. Annu. Rev. Food Sci. Technol. 13, 1–24.

https://doi.org/10.1146/annurev-food-052720-010751

Zotta, T., Ricciardi, A., Condelli, N., Parente, E., 2021. Metataxonomic and metagenomic

approaches for the study of undefined strain starters for cheese manufacture. Crit. Rev.

Food Sci. 1–15. https://doi.org/10.1080/10408398.2020.1870927