Hyperglycemia in type 2 diabetes: physiological and clinical implications of a brain centered

model

Akanksha Ojha,

Indian Institute of Science Education and Research, Pune, 411008, India

akanksha.ojha@students.iiserpune.ac.in

Milind Watve, *

Independent Researcher,

E-1-8, Girija Shankar Vihar,

Karve Nagar, Pune 411052, India

milind.watve@gmail.com

* for correspondence

Hyperglycemia in type 2 diabetes: physiological and clinical implications of a brain centered

model.

Abstract:

The hyperglycemia of type 2 diabetes has been classically believed to be a result of insulin

resistance and relative insulin deficiency. However, the unexpected outcomes of insulin

receptor knockout and insulin suppression experiments and the unexplained patterns in

clinical data have exposed the limitations of the peripheral regulation models. Absence of

fasting hyperglycemia by insulin receptor knockouts or insulin suppression, evidence for

hyperinsulinemia preceding insulin resistance, the perplexing hyperinsulinemic

normoglycemic state, reduced glucose transport to the brain preceding hyperglycemia,

absent or poor correlation between fasting glucose and insulin, very strong positive

correlation between indices of insulin resistance and β cell function in population data are

some of the anomalous findings which glucose homeostasis models have not addressed so

far. On the other hand the increasing evidence of central neuronal mechanisms of glucose

regulation necessitates incorporation of brain glucose regulation in the models. Although

models considering brain glucose regulation have been suggested, their multiple

physiological and clinical implications have not been thoroughly examined. Based on

updated evidence, we propose a refined model of glucose regulation that considers brain

glucose levels as the ultimate target of homeostasis and combines central and peripheral

mechanisms of regulation. In this model, considering reduced rate of blood to brain

transmission of glucose and insulin as primary pathology explains most of the patterns

including apparent anomalies with or without including insulin resistance. A brain centered

model accounts for the limited and inconsistent success of glucose normalization in

effectively reducing diabetic complications and mortality. A serious attempt to reevaluate

alternative models is required since the brain centered model implies radical changes in the

target and the course of diabetes treatment.

Key words: Glucose homeostasis, hyperglycemia, type 2 diabetes, steady state

Presently, any metabolic disorder with consistent raised plasma glucose and insulin

resistance, which cannot be categorized into either type 1 diabetes or gestational diabetes

is defined as type 2 diabetesmellitus (T2DM) (DeFronzo et al., 2015). It is the most common

form of diabetes, accounting for almost 90% of all cases.The currently perceived cause of

hyperglycemia in T2DM is insulin resistance followed by failure of compensatory insulin

response. Historically the role of brain in glucose homeostasis was revealed by experiments

in which damage to the fourth cerebral ventricle was shown to impair homeostatic control

(Bernard, 1859). However, at a later stage, pancreatic extracts were shown to lower glucose

and eventually insulin was identified as the active principle (Banting et al., 1922). It is

necessary to note that the differentiation of type 1 and type 2 diabetes was not made at

that time. The success of insulin was so spectacular that all other mechanisms of glucose

regulation were almost forgotten for about a century (Lundqvist et al., 2019). The distinction

between type 1 and type 2 diabetes was to become clear half a century later and it was

realized that unlike in type 1, insulin deficiency was not the primary cause of hyperglycemia

in type 2 diabetes. However, by this time the thinking in the field was so much insulin

centered that the inability of normal or increased levels of insulin to regulate plasma

glucose was termed insulin resistance, without carefully evaluating alternative hypotheses.

Further,the insulin resistance hypothesis is criticized for having a circular logic which makes

it un-falsifiable (Diwekar-Joshi&Watve, 2020). Insulin resistance is said to be responsible for

the inability of normal or raised levels of insulin to regulate glucose. However, insulin

resistance itself is measured by the inability of insulin to regulate glucose. There is no

independent measure of insulin resistance that can be used to test the causal role of insulin

resistance. Impairment of insulin signaling in experiments using (i) tissue specific insulin

receptor knockouts (Blüher et al., 2002; Kadowaki, 2000) (ii) insulin degrading enzyme

knockouts (Costes & Butler, 2014; Maianti et al., 2014) (iii) pharmaceutical insulin

suppression by diazoxide and octreotide (Giustina et al., 1991; Lamberts & Hofland, 2019;

Leahy et al., 1994; Matsuda et al., 2002)(iv) insulin gene suppression by RNAi (Mehran et al.,

2012b) (v) alteration in insulin gene dosage (Templeman et al., 2015) does not consistently

alter fasting glucose although it alters the post meal glucose curve. In epidemiological data,

fasting insulin and fasting glucose are poorly correlated while post meal insulin and glucose

are strongly correlated, unlike what would be expected by any of the models. With

converging multiple lines of evidence Diwekar-Joshi and Watve (2020) raised doubts on

whether insulin resistance and failure of compensatory hyperinsulinemia is a necessary and

sufficient explanation of fasting hyperglycemia.

Many mathematical models have been constructed based on the assumption of insulin

resistance. A common set of assumptions is shared by most of the models that fasting

glucose is the steady state achieved by a balance between glucose uptake by tissues and

glucose production by liver. One of the foundational assumptions is that both the above

processes are regulated by insulin signaling in the steady state (Matthews et al., 1985).

Therefore insulin signaling has been central to these models. Since recent research has

exposed the inadequacies of this model (Diwekar-Joshi & Watve, 2020) there is a need to

hunt for better alternative models.

An alternative to the classical model which assumes brain glucose level as the target of

homeostasis has been suggested and mathematical models incorporating this concept have

been constructed (Gaohua & Kimura, 2009; Watve, 2013). These models have not been

explored sufficiently towards a comparative evaluation with peripheral models with respect

to their predictions and their physiological as well as clinical implications. Here we refine the

brain centered model further incorporating recent evidence, evaluate its performance and

explore the physiological and clinical implications.

<u>What a glucose homeostasis model needs to explain:</u>

Most models of glucose homeostasis have a limited goal of explaining altered fasting steady

state glucose level and the alterations in the glucose curve in the pre-diabetic and diabetic

state. However, now we have a large number of experimental results that are potentially

anomalous. A model prediction matrix, in which each model is tested for matching its

prediction with multiple known empirical patterns, is an appropriate and robust approach to

compare alternative models. Such an approach has been used to test alternative hypothesis

in a number of other contexts (Thakar et al 2003, Diwekar and Watve 2016, Vibishan and

Watve 2020, Shinde et al 2021) and it can be the right approach to test alternative models

of glucose homeostasis in T2DM. A model of choice would be one which can predict all or

most of the known physiological patterns in type 2 diabetes so that no or minimal anomalies

are left. We take a few steps towards a model prediction matrix, although a lot more

exploration of many other models is required before constructing such a matrix.

Following is the list of known empirical patterns that a model needs to be compatible with.

1. The model should allow stable steady state glucose on fasting: Fasting glucose is

assumed to be a steady state with adequate evidence (Lerner & Porte, 1972;

Matthews et al., 1985; R. C. Turner et al., 1979). Achieving a steady state is one of

the primary objectives of every homeostasis model. Many of the key concepts of

classical models are also based on achieving a steady state in the fasting condition

(Matthews et al., 1985; R. C. Turner et al., 1979).

2. The model should also allow stable steady state insulin on fasting. Normally this is

not a problem for any model, however many models have a problem in explaining

the altered steady state insulin level in a prediabetic state as explained below.

3. Hyperinsulinemic normoglycemic condition in the prediabetic state: A prediabetic

state is often accompanied by high level of fasting insulin (FI) but normal fasting

glucose (FG) in the plasma. This is not easy to achieve in a model. The classical

qualitative explanation of this state is that it is a state of insulin resistance with

compensatory rise in FI such that FG remains normal. For this to happen, there is a

need to estimate the level of insulin resistance and regulate insulin secretion

accordingly. It is possible to hypothesize a mechanism of sensing insulin resistance

and conveying this information to the pancreas. Such a mechanism can be glucose

dependent or glucose independent. Currently no glucose independent mechanism of

accurately assessing the insulin resistance and conveying it to β cells is known. A

common assumption is that insulin resistance reduces glucose uptake thereby

increasing FG. The rise in FG stimulates insulin production and the rise in insulin

levels normalizes glucose again. However, this mechanism fails to achieve a steady

state hyperinsulinemic normoglycemic condition (Diwekar-Joshi & Watve, 2020).

Plasma insulin has a short half life and after glucose levels are normalized, it is

unlikely to stay at a higher level unless there is a glucose independent trigger for

insulin secretion. Explaining a hyperinsulinemic, normoglycemic steady state is a

tricky challenge that any model needs to meet.

4. Hyperinsulinemia precedes insulin resistance and T2DM: Although hyperinsulinemia is believed to be a compensatory response to insulin resistance, a number of studies show that hyperinsulinemia precedes obesity and insulin resistance (Corkey &Shirihai, 2012; Mehran et al., 2012a; Pories & Dohm, 2020; Shanik et al., 2008; Weyer et al., 2000; Wiebe et al., 2021). This raises two independent questions. One is how and why hyperinsulinemia precedes insulin resistance. The second,more perplexing question is that if hyperinsulinemia is not a compensatory response to insulin resistance, the failure of compensation explanation becomes redundant. This needs to be replaced by a more coherent explanation of hyperglycemia.

5. The model should explain the following features of post meal glucose curve and its alterations in prediabetic and diabetic states, which cumulatively are called impaired glucose tolerance curve.
   - Increased height of peaks
   - Delayed return to normal
   - Increase in the time difference in glucose and insulin peaks in the diabetic state.

6. The normal fasting glucose, impaired glucose tolerance (NFG-IGT) state: It is common to find that in a prediabetic state the glucose tolerance curve is altered with normal level of fasting glucose. A good model needs to explain the NFG-IGT phenomenon.

7. Stress hyperglycemia: Why and how stress causes hyperglycemia only in some individuals (Dungan et al., 2009) needs to be explained by a model with realistic and testable assumption.

8. Hyperglycemia after intensive exercise: If in a fasting state FG is a resultant of the rate of liver glucose production and glucose disposal, exercise that increases glucose disposal should decrease FG. However, in some studies plasma glucose remains unchanged or is increased after exercise (Coggan, 1991; Marliss & Vranic, 2002). A model needs to incorporate this phenomenon.

9. An important challenge is to explain why insulin receptor knockouts specific to muscle, fat cell or liver results into normal fasting glucose but altered post glucose load curves. Similarly, insulin degrading enzyme knockouts do not alter FG. Also insulin suppression by agents such as octreotide or diazoxide fail to alter FG (Diwekar-Joshi & Watve, 2020). A model should explain this observation. Diwekar-

Joshi and Watve (2020) further differentiated between the consequential steady state (CSS) and targeted steady state (TSS) models and demonstrated that the results of the insulin receptor knock out and insulin suppression experiments can be explained by a TSS but not by a CSS model. In effect, to be able to account for these results a model needs to be a TSS model.

10. Reduced glucose transport to brain, particularly in obese or prediabetic individuals: It is known over a long time that in type 2 diabetes the rate of glucose transport from blood to brain is slowed down substantially (Gjedde & Crone, 1981). In rodent models, the glut-1 expression in brain capillaries is shown to be reduced (Cornford et al., 1995; Matthews et al., 1985; Pardridge et al., 1990). This has been viewed as a response to hyperglycemia. However, Hwang et al (2017) show that subnormal transport is evident in obese and presumably prediabetic state, even before hyperglycemia appears. A model needs to account for this pattern as a causal or consequential phenomenon.

11. The T2DM specific islet amyloid deposition which often accompanies islet deterioration: Islet amyloid deposition is frequently associated with type 2 diabetes but not observed in type 1. The causes of islet amyloid deposition should be compatible with the glucose homeostasis model (Höppener et al., 1999; Porte & Kahn, 1989; Watve, 2014).

12. The pattern of B cell deterioration: The deterioration of β cells appears to follow a peculiar pattern in T2DM, in which a substantial proportion of β cells survive lifelong (Bacha et al., 2013; Butler et al., 2003; Clark et al., 2001; Porte & Kahn, 2001; Watve, 2014). If dysfunction and damage to the β cell population is assumed to be an essential prerequisite of hyperglycemia in a model, the models need to account for the peculiar population dynamics of β cells.

13. SGLT2 inhibitors increase liver glucose production as well as ketogenesis: Over the last decade SGLT2 inhibitors, which allow greater clearance of glucose through urine, have offered a novel means of combating hyperglycemia. Lowering of plasma glucose by SGLT2 inhibitors is accompanied by increased liver glucose production as well as increased ketogenesis (Limenta et al., 2019; Mistry & Eschler, 2021; Op den Kamp et al., 2021; Pfützner et al., 2017). A glucose homeostasis model needs to account for this phenomenon.

14. The Somogyi phenomenon i.e., hyperglycemia following infusion of insulin is seen in at least some cases of T2DM but not in healthy individuals. This can happen without being overtly hypoglycemic (Campbell, 1976; Reyhanoglu & Rehman, 2021). This phenomenon needs to be incorporated by a good model of glucose homoeostasis.

15. In cases of brain injury hyperglycemia is common (Shi et al., 2016). In bacterial meningitis hyperglycemia is often observed accompanied by low CSF glucose level (Krishnan et al., 2016; Schut et al., 2009). A model should be able to account for these phenomena.

16. In population data one sees good correlation between post meal glucose and insulin but poor correlation between FG and FI (Diwekar-Joshi & Watve, 2020). This population pattern should be explained by a comprehensive model.

17. Although FG and FI are poorly correlated in prediabetic state, HOMA-β and HOMA-IR have strong positive correlation in population data (Diwekar-Joshi & Watve, 2020). A model should predict such a pattern in a population.

18. Failure of glucose normalization to reduce the frequency of complications and mortality: Unlike T1DM, in T2DM tight regulation of plasma glucose has failed to show reduction in mortality consistently across multiple large scale clinical trials (Diabetes Prevention Program Research Group, 2015; Ferrannini & DeFronzo, 2015; King et al., 1999; Klein, 2010; Lee et al., 2021; Schwartz & Meinert, 2004; R. Turner et al., 1998; The NICE-SUGAR Study Investigators, 2009, UK Prospective Diabetes Study (UKPDS) Group, 1998). If hyperglycemia is the primary pathology of type 2 diabetes, preventing or correcting it should have reduced the frequency of complications and mortality considerably.

19. Reversal of hyperglycemia by FGF 21 in all models of rodent diabetes: Independent of the cause of hyperglycemia, a single injection of FGF 21 was able to achieve long term normalization of plasma glucose (Laeger et al., 2017). This is a potential anomaly that a model needs to resolve.

While most models intend to explain 1, 2 and 5 of the above, they either fail to explain or do not address the others. Our expectation from a satisfactory model is that it should inherently predict these outcomes or should be compatible with them. If an additional explanation that does not contradict the model, accounts for the phenomenon, the model is compatible. Compatibility with empirical findings cannot be taken as a proof or validation of

the model, but if a model directly contradicts one or more of the empirical findings, it

becomes serious grounds for rejecting the model. Later when we describe model results, we

will use the numbers in the above list in square brackets to indicate that the model

accounted for this pattern.

<u>Peripheral models of glucose homoeostasis</u>:

There is a rich history of development of mathematical models of glucose homeostasis and

the origin and progression of type 1 and type 2 diabetes (reviewed by Ajmera et al., 2013;

Mari et al., 2020). The focus of the field is so much on peripheral mechanism that the review

by Mari et al (2020) does not even cite the models incorporating the role of brain. Ajmera et

al (2013) briefly mention the Gouhua et al (2009) model but do not elaborate on its

potential implications. A class of models attempts to capture glucose homeostasis at the

systems levels whereas other models look at individual components such as insulin

dependent glucose uptake, liver glucose production, β cell dysfunction, glucose stimulated

insulin secretion and incretin effects at greater details. Nevertheless, these models together

have not accounted for many of the patterns listed above. The central assumption of these

models is more or less invariant and revolves around insulin resistance and the failure of

insulin response to adequately compensate for it (Mari et al., 2020). Diwekar-Joshi and

Watve (2020) claimed that any model with this set of assumptions is not compatible with

empirical patterns 3,4,6,9, 16 and 17 of the above. Whether any variation of these models

can do so has not been adequately explored. Furthermore, all the peripheral regulation

models are CSS models in that the fasting steady state is a consequence of the rate of liver

glucose production and a concentration dependent rate of glucose uptake. In these models

a change in either or both the rates inevitably alters the steady state. This contrasts the TSS

models in which alteration in these rates alters the time required to reach a steady state but

does not alter the steady state glucose level (Diwekar-Joshi & Watve, 2020). An attempt to

develop a TSS model with only peripheral mechanisms has not been made to the best of our

knowledge.

We do not intend to review glucose homeostasis models here. However, we observe from

published reviews of glucose homeostasis models (Ajmera et al., 2013; Mari et al., 2020)that

most models do not address the apparently anomalous empirical findings particularly 3, 4,

6, 9, 10, 11, 12 ,13 and 15-18 among the ones listed above. A detailed exploration in to

whether some variations of these models can address the apparently anomalous patterns is

beyond the scope of this paper but we remain open to this possibility.

<u>Brain centered models of glucose homeostasis</u>**:**

It is quite well known that a number of neuronal mechanisms in the brain are involved in

energy homeostasis. Nevertheless, for some reason, they do not occupy a central stage in

the mainstream thinking in T2DM and glucose regulation models. A number of recent

publications highlight the role of brain in different ways (Lam, 2005; Osundiji et al., 2012;

Perrin et al., 2004; Watve, 2013, Lundqvist et al 2019). Peters (2004), Gaohua and Kimura

(2009) and Watve (2013) proposed that the primary target of glucose homeostasis is to

regulate glucose levels in the brain. Plasma glucose levels are only a means to achieve

required supply of glucose mainly to the brain. Since transport of glucose to the brain is

more restricted, when adequate supply of glucose to the brain in ensured, it is likely that the

supply to other organs already gets ensured.

In the Gaohua and Kimura (2009) as well as in the Watve (2013) model the rate of glucose

transport across the BBB is assumed to be an adaptation to hyperglycemia. The finding that

glucose transport is reduced to an intermediate level in obese and prediabetic individuals

(Büsing et al., 2013; Hwang et al., 2017) suggests that the deficiency in transport precedes

hyperglycemia rather than following it. Therefore, it is more likely to be causal than

consequential. Glucose deficiency in the brain is known to induce liver glucose production

and suppresses insulin release through autonomic control (Boland et al., 2017). Sympathetic

tone has also been shown to be higher in T2DM (Thackeray et al., 2012). Therefore glucose

deficiency in the brain owing to altered vascular function is likely to be primary which results

in altered plasma glucose dynamics mediated by autonomic inputs. Our model incorporates

this possibility.

Impaired vascular function is known to be central to diabetic complications. The classical

thinking has been that hyperglycemia causes the types of vasculopathies typical of T2DM.

However, it has not been ruled out that vasculopathies are not primary. There is

considerable evidence that microvascular alterations precede T2DM and are good

predictors of it (Muris et al., 2012; Nguyen et al., 2007; Stehouwer, 2018; Zaleska-Żmijewska

et al., 2017). Sedentary life style and deficiency of many specific types of physical activities

and behaviors alter the expression of growth factors and endocrine mechanisms involved in

angiogenesis. It has been demonstrated repeatedly that many growth factors and

angiogenic factors are responsive to behaviors, physical activity and exercise (Aloe et al.,

1994; Cao et al., 2010; Chodari et al., 2016; Lakshmanan, 1986; Nexø et al., 1984; NEXØ et

al., 1981; Tirassa et al., 2003). Deficiency of these behaviors is likely to lead to a primary

deficiency of angiogenic mechanisms(Watve, 2013). We consider the possibility that altered

vascular function in the brain is the primary reason why glucose transport from blood to

brain is reduced. This creates chronic glucose deficiency in the brain to which the brain

reacts by influencing multiple mechanisms of glucose regulation. The predominant

mechanism is autonomic signaling. It is well known that sympathetic and parasympathetic

tones are altered in T2DM (Thackeray et al., 2012). We consider the possibility that altered

autonomic signaling is causal rather than consequential to diabetes, based on studies

showing that changes in autonomic function precede T2DM (Carnethon et al., 2003).

The brain glucose uptake is insulin independent (Gray et al., 2014; Hasselbalch et al., 1999).

Nevertheless insulin has many other functions in the brain and insulin signaling is known to

be important in memory, cognition, decision making and behavior (Kerns 2001, Shemesh et

al., 2012; Strachan, 2003). Brain is rich in insulin receptors and activation of certain cognitive

functions in the brain is a likely cause of increased brain glucose utilization by insulin

stimulation (Bingham et al., 2002; Rebelos et al., 2021). We assume therefore that brain has

a fundamental requirement for insulin, independent of glucose metabolism and therefore

mechanisms to ensure the required insulin levels in the brain are as much critical as

ensuring minimum brain-glucose level. This is also ensured by autonomic mechanisms.

Autonomic inputs are known to regulate β cell population and as well as insulin release from

β cells (Thorens, 2015). Therefore, our model also incorporates brain control over insulin

production.

Ensuring minimum supply of glucose as well as insulin to the brain is crucial during fasting

when the plasma levels are low. The post meal levels of glucose and insulin in the plasma

are much higher and at this time, the brain need not actively regulate the plasma levels of

both. This can be taken care of by the peripheral mechanisms.

<u>Objectives of the model</u>:

Our attempt is to construct a model whose predicted outcomes match with all or most of

the observed phenomena listed above, qualitatively. We intend to construct a model in

which different hypotheses for hyperglycemia can be used to make differential testable

predictions. This approach can allow us to evaluate comparatively which causal factors

individually or in combination can give us the set of predictions that match with the patterns

listed above.

Out of the parameters required for the model, only some have empirical estimates available

(table 1). In the absence of realistic estimates of all parameters, we do not intend to make a

model making quantitative predictions. When a sufficiently large number of parameters can

be optimized, it is not difficult to fit the data quantitatively. Therefore we prefer qualitative

predictions over quantitative ones. We test whether the model is able to predict the pattern

observed empirically under some set of parameters. The ability to predict a given pattern is

not a proof of the validity of the model, but the inability to predict an observed

phenomenon at any set of parameters in a realistic range is a strong reason to call the

model either inadequate or wrong. If an additional consideration compatible with the model

is able to explain a pattern not explained by the main model, the model can be called

inadequate but not falsified. However, if the model outcome directly contradicts a

consistent and reproducible empirical finding, it can be considered falsifying evidence. We

emphasize the need to evaluate our model in comparison with classical models on these

lines.

<u>Assumptions of the model</u>**:**

A unique assumption of our model based on the analysis by Diwekar-Joshi and Watve (2020)

is that the mechanism of regulation of glucose and insulin is different in the steady state and

post meal state. In the steady state the central mechanisms are more important whereas in

post meal state mainly the peripheral mechanisms are at work. Insulin induced glucose

uptake and insulin dependent inhibition of liver glucose production happen only above a

threshold glucose and insulin level respectively. The presence of such thresholds and their

context dependent variability has been known for a long time (Chen et al., 1993; Henquin et

al., 2006; Sorensen 1985). Below the threshold, other mechanisms regulate the levels

(Sorensen 1985). We assume in the model that the thresholds are under neuro-endocrine

control and fine-tuned by the brain. The thresholds found in isolated cell cultures are slightly

lower than the normal fasting blood sugar and it is difficult to estimate thresholds in vivo

(Chen et al., 1993; Henquin et al., 2006, 2015). In presence of autonomic signals, we assume

the thresholds to be above the steady state target levels. In conditions under which the

fasting levels need to increase, we assume the thresholds to increase proportionately. This

may happen with peripheral or central mechanisms. These assumptions about the threshold

make the model a TSS model.

The steady state glucose is decided by a balance between basal level of liver glucose

production which is regulated by neuronal mechanisms and insulin independent glucose

uptake by tissues.The steady state insulin level is decided by basal rate of glucose

independent insulin production, which is regulated by neuronal mechanisms against the

insulin degradation rate. Glucose is transported from plasma to brain by facilitated diffusion

proportionate to capillary surface area in the brain and glut-1 expression. Steady state brain

glucose is determined by glucose transport to brain and utilization by brain tissue. Brain

needs a target level of glucose and the brain ensures the minimum by directly regulating

liver glucose production by neuronal mechanisms.

Insulin has cognitive and other functions in the brain (Begg & Woods, 2013; Kern et al.,

2001; Rohner-Jeanrenaud & Jeanrenaud, 1983; Shemesh et al., 2012; M W J Strachan,

2005), for which the brain requires a target level of insulin, which it ensures by autonomic

regulation of β cell number and basal insulin secretion independent of glucose. For ensuring

the brain target level for insulin, a minimum plasma insulin level needs to be maintained.

Maintenance of this level is assumed to be independent of glucose stimulated insulin

secretion.

The rate of transport of glucose and insulin from blood to brain is not constant. Both are

affected by the capillary density and blood flow. But they also depend upon specific

transport mechanisms. Glucose is transported across the blood brain barrier mainly through

the specific transporter glut-1. The expression of glut-1 in brain and other tissues is variable

and dependent on multiple dietary, endocrine and growth factor related mechanisms

(Boado et al., 1994; Ge et al., 2011; Liu et al., 2018; Schüler et al., 2018).Glucose transport

from blood to brain is diminished in obesity and prediabetes (Hwang et al., 2017). Insulin

transport to brain is also reduced in obesity (Begg, 2015). The ratio of CSF to plasma insulin

is inversely proportionate to obesity and insulin resistance (Gray & Barrett, 2018). Moreover

inducing endothelial dysfunction and reducing glucose transport experimentally by

endothelial specific deletion of HIF 1 α resulted into hyperglycemia (Huang et al 2012).

Based on such multiple lines of evidence we incorporate the possibility of diminished

glucose and insulin transport to brain being the primary pathology of T2DM and

hyperglycemia only a consequence.

The model:

Since we assume that the steady state and post meal mechanisms of glucose regulations are

not identical, we model and analyze the two states in two phases of the model. The

variables and parameters used in the model and the range of parameter used are in table 1.

sr no	Variables and parameters	meaning	units	Value/range used in simulations
A. Input parameters				
1	K_1	Insulin independent glucose uptake maximum	mg/dL/min	20-50
2	K_{1m}	Half saturation constant for insulin independent glucose uptake	mg/dl	50
3	K_2	Maximum insulin induced suppression of PG, including glucose disposal and suppression of L.	per μU/mL	0.001 to 0.05
4	K_{2m}	Half saturation constant for insulin induced suppression of PG	μU/mL	15
5	K_5	Maximum glucose dependent insulin secretion	μU/mL/min	2-20
6	K_{5m}	Half saturation constant for glucose induced insulin secretion	mg/dL	50
7	K_8	Maximum rate of glucose transport to brain	Mg/dL/min	3 - 8
8	K_{8m}	Half saturation constant for glucose transport to brain	mg/dL	50-100
9	K_9	Maximum rate of Insulin transport	μU/mL/min	0.6 - 1

#	Symbol	Description	Units	Value
		to brain		390
10	K_{9m}	Half saturation constant for insulin transport to brain	µU/mL	0.3-1 391
11	K_{10}	Brain glucose utilization rate	unitless	0.08 to 0.12 392
12	d_1	Insulin degradation rate	unitless	0.1 to 0.2
13	BGt	Target BG	mg/dL	25 393
14	BIt	Target BI	µU/mL	1-5 394
15	I_s	Insulin sensitivity as compared to normal which is assumed 1.	unitless	0-1 395
16	L_{max}	Maximum capacity of liver glucose production	Mg/dL/min	30 396
17	K_{4max}	Maximum capacity of glucose independent insulin secretion	µU/mL/min	397 10 398
B. Derived variables				
18	L	Rate of liver glucose production	mg/dL/min	Eq. 6 or iterati399
19	K_4	Rate of glucose independent insulin secretion	µU/mL/min	Eq 7 and 8 or 400 iterative
C. Outcome variables				401
20	PG	Plasma glucose	Mg/dL	dynamic 402
21	PI	Plasma insulin	µU/mL	dynamic
22	$\underline{PG}$	Steady state plasma glucose	mg/dL	Eq. 5 403
23	$\underline{PI}$	Steady state plasma insulin	µU/mL	Eq. 7 404
24	BG	Brain glucose	mg/dL	dynamic 405
25	BI	Brain insulin concentration	µU/mL	dynamic 406
26	PGT	glucose threshold above which GSIS begins	mg/dL	5 to 10 mg/dL 407 above $\underline{PG}$ 408
27	PIT	Insulin threshold above which insulin induced glucose disposal begins	µU/mL	2-5 µU/mL 409 above $\underline{PI}$ 410

Table 1: all variables and parameters considered in the model.

A: Modeling steady state glucose:

As per the assumption, below a threshold *PI*, insulin induced glucose uptake and insulin

induced inhibition of liver glucose production is negligible,

1. Plasma Glucose:

$$\frac{dPG}{dt} = \begin{cases} L - \frac{K_1 * PG}{K_{1m} + PG}, \quad + Gt & (\underline{PI} < PIT) \\ L - \frac{K_1 * PG}{K_{1m} + PG} + Gt - \left(I_S * \frac{K_2 * PI}{K_{2m} + PI}\right) * PG, & (PI > PIT) \end{cases}$$ Eq. 1

2. Plasma Insulin:

$$\frac{dPI}{dt} = \begin{cases} K_4 - (d1 * PI), & (\underline{PG} < PGT) \\ K_4 - d1 * PI + \frac{K_5 * PG}{K_{5m} + PG} * (PG - PGT1), & (\underline{PG} > PGT) \end{cases}$$ Eq. 2

3. Brain glucose:

$$\frac{dBG}{dt} = \frac{K_8 * (PG - B)}{K_{8m} + (PG - B)} - K_{10} * BG$$ Eq. 3

Since brain forms only about 3 % of total body mass, we assume the reduction in plasma

glucose diffused to the brain is a negligible fraction of total plasma glucose. Therefore,

that term is not included in the plasma glucose dynamics. Similarly, insulin diffusion to

brain is not represented in the plasma insulin dynamics. Nevertheless, the rate of

plasma to brain diffusion is crucial in determining the brain glucose and insulin

dynamics.

4. Brain insulin:

$$\frac{dBI}{dt} = \frac{K_9 * (PI - BI)}{K_{9m} + (PI - B)} - d_1 * BI$$ Eq. 4

At steady state (SS) all differential terms become zero and steady state levels can be

calculated using equilibrium solutions.

As per the assumption of the model the brain needs a certain level of glucose and that is

the target of homeostasis *BGt.* In order to ensure *BGt* at given K_8, K_{10} and K_{8m}, the SS

plasma glucose *PG* needs to be

$$\underline{PG} = \frac{BG_t * K_{10} * K_{8m}}{K_8 - (BGt * K_{10})} + BGt$$ Eq. 5

We assume that the brain ensures this plasma glucose level by regulating liver glucose

production L. To maintain the desired plasma glucose level at steady state, the liver

glucose production should be:

$$L = \frac{K_1 * \underline{PG}}{K_{1m} + \underline{PG}}$$ Eq. 6

The brain can either ensure this by sending a calculated signal to the liver through the

autonomic system. Alternatively, this can be ensured by the brain by increasing

sympathetic inputs when $BG < BGt$ and parasympathetic when it is above target (Antuna-

Puente et al., 2009; D'Alessio et al., 2001; Kiba, 2004). The required L can be achieved

iteratively by $L_{(t+1)} = L_{(t)} + K_s.(BGt - BG)$

Similarly, to ensure required brain insulin levels, plasma insulin should be

$$\underline{PI} = \frac{d_1 * BIt * K_{9m}}{K_9 - (d_1 * BIt)} + BIt$$ Eq. 7

To ensure this much plasma insulin level, K4 should be

$$K_4 = d_1.\underline{PI}$$ Eq. 8

Similar to glucose, the brain can ensure the target level of insulin in the brain by

neuronally modulating K_4 in a calculated or iterative manner. Specific autonomic inputs

are known to regulate β cell number as well as insulin release from β cells (Begg &

Woods, 2013; Boland et al., 2017; Kiba, 2004).

We also assume that both L and K_4 have a fixed upper limit as L_{max} and K_{4max} when the

maximum capacity of liver and pancreas is reached. L and K_4 cannot exceed this limit

even if neuronal inputs become more intense. This happens when K_8 declines below a

threshold such that,

$$K_8 < \frac{K_1 - L_{max}(BGt * K_{10} * K_{8m})}{L_{max} * K_{1m} - BGt(K_1 - L_{max})} - (BGt * K_{10})$$ Eq. 9

There is a potential conflict here. Sympathetic tone is known to increase L (Nelles et al.,

1996; Nonogaki, 2000) whereas parasympathetic signal is required for proliferation of β

cells. Simultaneously sympathetic signal is known to inhibit insulin release from β cells

(Gilon & Henquin, 2001; Miller et al., 1976). If the brain is deficient in glucose as well as

insulin, both sympathetic and parasympathetic tones will be simultaneously higher (Watve

et al 2014). This is incorporated into the model by updating K_4 iteratively according to both

BG and BI levels simultaneously,

$$K_{4(t+1)} = K_{4(t)} + K_p(BIt - BI) - K_s(BGt - BG)$$ Eq. 10 By

this consideration if sympathetic stimulation of liver glucose production is adequate to

restore required BG, $BGt - BG$ will tend to zero and there will be no interference in insulin

release by β cells. However, in case the rise in liver glucose production is inadequate to

ensure required BG, β cell function may be apparently affected.

B. Simulating glucose tolerance curve:

While for the steady state, equilibrium solutions can be algebraically derived, the post

glucose load curve and its properties can only be obtained by simulations. Simulations were

run using the same set of equations and giving a positive ephimeral Gt to simulate food

intake.

In all the simulations used, insulin resistance can be simulated by altering Is, β cell

dysfunction by reducing K_4 and/or K_5 and vascular defect slowing down transport of glucose

and insulin to brain and other organs by altering K_8, K_9 and K_1. Mental stress is assumed to

increase K_{10}. By differentially altering these parameters the model can separately and

collectively examine the effects of insulin resistance, β cell dysfunction and reduced rates of

diffusion across BBB on the glucose tolerance curve.

T2DM- causal analysis: We ask the question which minimal set of changes can give rise to

stable increase in fasting glucose, changes in the glucose tolerance curve characteristic of

type 2 diabetes along with other patterns listed above. The putative causal factors in are

examined individually and in combination. The factors include insulin resistance (Decreased

I_s), subnormal β cell response (Reduced K_4 and/or K_5), reduced insulin independent glucose

uptake (decrease in K_1), reduced blood to brain transmission of glucose (K_8) and insulin (K_9)

and stress related increase in brain glucose consumption (K_{10}). Subnormal or defective

vasculature is expected to decrease K_1, K_8 and K_9 proportionately, however the decrease in

the three parameters may not be proportionate if the vasculature in different parts of the

body is differentially affected. Also glucose transporters and there expressions can alter in

tissue specific manner. Therefore we allow proportionate as well as differential decrease in

the three parameters.

Results:

<u>A: Steady state solutions</u>: We observed that the steady state solutions and the autonomic

iteration simulations match well in the end result except when the limits of L and K_4 are

reached. However, it took a long time to reach the desired level by iterative approach,

particularly when the desired level was substantially different from the starting level. In real

life major changes in vascular function or glucose transporter levels K_8, K_9, K_1 happen

gradually. Therefore the desired level can be attained by autonomic fine tuning over time.

Since in the iterative simulations, a long time was required to reach SS, we used the steady

state solutions for quicker results.

It can be seen that the steady state levels of PG in the model is a stable steady state since

substituting $PG < \underline{PG}$ leads to a positive value and $PG > \underline{PG}$ leads to a negative value of $\frac{dPG}{dt}$.

The same applies to plasma insulin, brain glucose and brain insulin. This ensures a stable

steady state of glucose-insulin in the plasma as well as brain on fasting [1,2]. From equations

5,6,7, and 8 we see that, FG does not change by altering Is or FI. This is compatible with the

empirical findings that insulin receptor knockouts and insulin suppression experiments fail

to increase fasting plasma glucose [9]. This result is unique to our model and is due to

segregating fasting and post meal glucose regulation mechanisms. This is the only

explanation offered so far, patterns [9], [16] and [17].

The steady state solution of our model shows that FG can increase as a result of increase in

K_{10} or decrease in K_8. That is if the glucose consumption in brain increases and/or the rate of

glucose transport from blood to brain decreases. Both the effects are interdependent and

the shape of the glucose response is decided by the interaction between K_8 and K_{10}.

Increased K_{10} increases FG marginally and almost linearly when K_8 is large. When K_8

decreases even a marginal rise in K_{10} can induce disproportionately greater rise in FG (fig 1).

This means that stress induced hyperglycemia is unlikely to be seen in healthy individuals

while it is more likely in individuals with reduced vascular transport [7].

Figure 1: The same input different output (SIDO) phenomenon:

A. A conceptual diagram with arbitrary units: A saturation relationship in glucose and

insulin transport has important consequences that can account for many phenomena

observed in prediabetic and diabetic stages. For example here for achieving an

increase in Y by 6 units with a higher Ymax, X needs to increase only by 6 units but at

a lower Ymax for the same target increase in Y 25 units of increment in X is needed.

This non-linearity explains many phenomena including prediabetic hyperinsulinemia

and stress induced diabetes in our model of glucose homeostasis.

B. A simulation using the steady state model: It can be seen that as K_8 decreases, for the

same change in brain requirement K_{10}, a non-linear escalated increase in plasma

glucose is required.

C. *A simulation result assuming a correlated decrease in K_8 and K_9 which is expected if*

the underlying cause is reduced brain vasculature. It is possible that while glucose

shows a marginal increase, insulin increases substantially, since the parameters of

glucose and insulin transport curves are different. This is a potential explanation for

hyperinsulinemic normoglycemic state.

540	The origin of hyperinsulinemic normoglycemic condition can be explained by the
541	difference in the parameters of the saturation curves of transport dynamics of
542	glucose and insulin. If the insulin diffusion is assumed to be nearer to saturation and
543	glucose diffusion is sufficiently away from saturation, at the same level of vascular
544	function deficiency, insulin will increase disproportionately more than glucose
545	(Figure 1). This is a possible cause of hyperinsulinemic normoglycemic state [3], and
546	the reason why hyperinsulinemia precedes hyperglycemia [4] which requires neither
547	insulin resistance nor compensatory hyperinsulinemic response. When this happens
548	the HOMA-IR and HOMA β increase although there is no change in true insulin
549	resistance and compensatory insulin response. This gives a false impression of insulin
550	resistance as defined popularly.
551	Further as the transport saturation constant K_8 and K_9 continue to decrease
552	proportionately, FG increases monotonically but FI shows a non-monotonic curve in
553	which FI increases with moderate decrease in transport rates but decreases after a
554	threshold decrease in transport rates. This non-monotonicity is predicted by the
555	iterative autonomic inputs model. When the required L approaches or exceeds L_{max},
556	the steady state BG remains smaller than BGt. The resultant increase in the
557	sympathetic signal inhibits insulin release from β cells. This leads to reduced insulin
558	secretion and insulin levels drop substantially. Thus, without bringing in additional
559	factors the model explains the early phase rise in fasting insulin as well as the later
560	phase decline in the course of T2DM [4] (Figure 2).

 Fingure2: Effect of correlated decrease in K_8 and K_9 on fasting plasma glucose and insulin.

 Note that for a small to moderate decrease FI increases rapidly with marginal change in FG.

 However after a threshold decrease in the transport rates FI declines sharply whereas FG

 increases rapidly. This pattern matches with the course of clinical T2DM without involving

 insulin resistance, compensatory hyperinsulinemia and β cell exhaustion. Simulation specific

 other parameters were K_{1m} =50, K_1=30, K_{2m}=15, K_2=0.01, K_{5m}=0.01, K_5=0.5, d_1=0.15, I_s=1,

 BGt=25, BIt=5, K_{8m}=80, K_8=5, K_{9m}=0.3, K_9=0.8, K_{10}=0.1(Blue line) and 0.12(Red line).

This decline in the insulin response does not require β cell dysfunction as a causal

mechanism but there are more complex possible effects on β cells. Simultaneous activation

of sympathetic and parasympathetic inputs to β cells is implicated in β cell amyloidogenesis

(Watve et al., 2014). Parasympathetic stimulation is known to increase β cell number.

However, sympathetic signaling suppresses insulin release resulting in increased retention

time of insulin along with amylin which beyond a threshold retardation results in to

spontaneous polymerization of amyloid protein leading to poisoning of β cells and

amyloidogenesis (Watve et al., 2014). This is an alternative causal interpretation of β cell

dysfunction and decline in its population. This mechanism has a built-in negative feedback

loop giving rise to a steady state β cell population. This result is compatible with the finding

in which the β cell population remains subnormal lifelong [11,12]. This contrasts the

dynamics expected by the classical thinking in which β cells are destroyed by gluco-

lipotoxicity or oxidative stress. This mechanism has a built-in positive feedback cycle. As a

part of the β cell population is destroyed, the insulin secretion is decreased which would

increase glucose further accelerating gluco-lipotoxicity and oxidative stress and thereby β

cell loss. Such a positive feedback vicious cycle can only stop with complete destruction of

the β cell population. This prediction of the classical model does not match the finding of

sustained presence of subnormal β cell population in T2DM.

By the classical model, prolonged sustained exercise in the fasting state should result into

lower plasma glucose. However, experimental increase in glucose uptake resulting from

physical activity during fasting does not necessarily result into a decrease in *FG*. This is

mainly because as *FG* starts decreasing, *BG* decreases consequently. *BG* being lower than

the brain target induces sympathetic mediated increase in *L* and the change in *FG* by

physical activity is more or less compensated. This may happen through the agency of

glucagon as well, since a strong link between autonomic signals and glucagon secretion is

known. Therefore, at times plasma glucose may actually increase after intensive exercise.

Further, if exercise involves heightened activation of cerebellar mechanisms of coordination

K_{10} can also increase resulting into higher rather than lower *FG* in response to exercise [8].

A high dose of exogenous insulin results into increased plasma glucose after a time lag, is

known as Somogyi phenomenon. The phenomenon is conditional and not seen in every

diabetic. It is said to involve the counter-regulatory response on facing central

hypoglycorrhachia. However, often this response is seen without peripheral hypoglycemia.

This is best explained by reduced K_8 that gives rise to decreases *BG/FG* ratio. With smaller

K_8, at apparently normal or increased plasma glucose, brain glucose can still be lower than

the target which gives a sympathetic signal to increase *L*. Simulations using the autonomic

iteration model shows this phenomenon quite well at low K_8. At the same level of insulin

infusion, at lower K_8 there is more intense hyperglycemic response (Figure 3) [14]. A model

without accommodating changes in K_8 does not predict Somogyi phenomenon without an

obvious hypoglycemic state.

At transmission 90%

At transmission 70%

Figure 3:Effect of high dose insulin on peripheral glucose under two rates of glucose and

insulin transport across the blood brain barrier (Correlated decrease in K_8 and K_9). A

paradoxical rise in blood sugar following insulin administration is more prominent when the

transport rate is substantially lower than normal. Other parameters (K_{1m} =50, K_1=30, K_{2m}=15,

K_2=0.08, K_{5m}=0.01, K_5=15, d_1=0.15, I_s=1, BGt=25, BIt=4.5, K_{8m}=80, K_8=5, K_{9m}=0.3, K_9=0.8,

K_{10}=0.1)

The response is restricted to a narrow set of conditions. For intense hyperglycemic response

to insulin administration, it is necessary that insulin sensitivity is good, *BG* is close to *BGt*

before insulin transfusion, and K_8 is subnormal.

Potentially one can visualize two distinct possible classes of reasons why blood sugar

increases in response to brain injury. In the first, there is some impairment in the glucose

regulation mechanism. In the second, the requirement for glucose is increased during the

wound healing process and hyperglycemia is a mechanism of the body to meet the demand.

It is known that infections such as meningitis, stroke or hemorrhage leads to transient

hyperglycemia. Such hyperglycemia is often accompanied by lower *BG* levels (Schut et al.,

2009; van Veen et al., 2016)[15]. In such cases, hypoglycemic treatments can be

counterproductive and result into delayed or derailed repair process. This is a likely reason

why in patients under intensive care, strict regulation of plasma glucose increased mortality

instead of the expected decrease (The NICE-SUGAR Study Investigators, 2009).

When hyperglycemia is a process of meeting increased glucose demand or compensating

subnormal transport, removal of plasma glucose by any means is expected to increase *L*.

This is observed to happen when SGLT2 inhibitors decrease the urinary threshold and drive

out plasma glucose. If increasing *L* is not sufficient to restore the required glucose level,

there is a shift to ketogenesis since the brain can use keto acids as alternative source of

energy.Therefore, increased *L* with or without increased ketogenesis is expected following

SGLT2 inhibitors [13].

<u>B: Glucose tolerance curve</u>: Unlike the steady state predictions, the patterns of post feeding

glucose curve by our model are not qualitatively different from classical models. The area

under the curve, height of the peak, time required to return to steady state and the time

difference between glucose and insulin peaks are increased by decreased I_s, K_4 or K_8

individually or in combination. Decrease in I_s or K_4 secretion does not increase *FG* but

changes the shape of the glucose tolerance curve. Decreased K_8 may alter both

simultaneously. The altered curve shows the three typical features namely taller peak,

delayed return to steady state and longer gap between glucose peak and insulin peak [5].

This result is not unique to our model and classical models also show the three features. The

classical model results into simultaneous and proportionate alterations in the fasting as well

as post meal glucose by changing I_s or K_4 and therefore fails to explain the NFG-IGT state. In

our model, decreasing I_s or K_1 without a change in K_8 results into a NFG-IGT state [6].

C: Population simulations: For the steady state as well as post glucose load dynamics we

give population distributions to K_1, K_5, K_8, K_9, I_S and also incorporate normally distributed

error in glucose and insulin measurements in fasting and post meal sampling. We also

incorporate correlated changes in K_1, K_8 and K_9 which are expected as a result of hypo-

vascularization in the brain. These simulations are run to observe whether we get the

anomalous correlations in fasting versus post meal state and in the HOMA indices as

observed empirically (Chawla et al., 2018, Diwekar-Joshi and Watve 2020).

By classical models the regression correlation parameters of glucose-insulin relationship are

not different in fasting state versus post glucose load although the range of variables is

different as shown previously by Diwekar-Joshi and Watve (2020). Also, if we assume

HOMA-IR to faithfully reflect insulin resistance and HOMA β to faithfully represent β cell

response, then there is no reason why the two indices should be correlated.

The assumption behind our model that there are different mechanisms at work under

fasting versus post glucose load condition is necessary to explain the large difference in the

regression correlation parameters in fasting versus post meal levels. If the same set of

mechanisms in the fasting and post meal conditions are operational, whatever the model

used, it is imperative that fasting correlation regression parameters are stronger or similar

to post meal parameters. Simulations with our model are able to give poor *FG-FI* correlation

along with strong post meal glucose insulin correlation under multiple conditions [16]

(Figure 4). When the variance in K_8 and K_9 is small but that in K_1, K_5, Gt and I_s individually or

in combination is large, the fasting correlations are weak and post meal correlations strong;

the post meal glucose-insulin regression slope is substantially greater than *FG-FI* slope. Also,

whenever *FG-FI* correlation is weak, the indices HOMA-IR and HOMA β are strongly

correlated similar to the epidemiological data [17]. The difference between fasting and post

meal regression correlation patterns is not predicted by the classical models and is unique

to our model which assumes different set of regulatory mechanisms in the fasting and post
meal state.

Figure 4: Plasma glucose and insulin correlation in the fasting steady state (A) and post meal
(B) in simulations. The empirical finding that there is a strong positive correlation in the post
meal data (R^2=0.57) but poor correlation in fasting steady state (R^2=0.014) was possible
under a variety of conditions in our model. Depicted here the simulation results in which K_1
was given a wider population distribution mean (S.D.) K_1 = 35(7)and K_5 = 10(0.00001), K_8 =
5(0.000005), and K_9 = 0.8(0.000008) had a narrow distribution. Other parameters (K_{1m} =50,
K_{mum}=15, K_2=0.05, d_1=0.15, I_s=1, BGt=25, BI=4.5, K_{8m}=80, K_{10}=0.1).

<u>D: Effect of glucose normalization on arresting diabetic complications and mortality</u>: By
classical thinking, chronic hyperglycemia is responsible for the diabetic complications and
preventing hyperglycemia should arrest complications. In contrast the thought behind our
model is that vascular problems are primary which alter the rate of glucose insulin transport
to the brain and hyperglycemia is an offshoot symptom that may not be causal to diabetic
complications which can arise directly from the vasculopathy. Therefore, controlling glucose
may not have any effect on diabetic complications [18]. On the other hand forcefully
reducing plasma sugar without addressing vascular problems can create more severe
glucose deficiency in the brain and other organs, thereby turning counterproductive.

Because of the saturating dynamics of transport, a curvilinear relationship is expected

between plasma glucose and brain glucose in such a way that moderate reduction in

hyperglycemia will change brain glucose availability marginally whereas tight glucose

regulation can have disproportionately larger effect (Figure 5). Therefore, tight glucose

regulation may increase mortality and other adverse outcomes. This prediction is

compatible with some of the tight versus moderate control clinical trials including ACCORD,

NICE sugar trial and UGDP [18].

Figure 5: Effect of glucose lowering on brain glucose availability. For example, a reduction in

FG from 400 mg/dl to 150 mg/dl corresponds to a decrease in BG of 1.68 mg/dl, but further

reduction from 150 to 70 mg/dl reduces BG by 2.41 mg/dl. Therefore a moderate sugar

control may not affect brain glucose supply drastically but tight sugar control is expected to

affect it more seriously.

Discussion:

The main inference from our model, stated most conservatively, is that a brain centered

model can potentially explain most of the anomalies faced by peripheral models and

therefore needs greater attention. If supported well, by exploring its testable predictions

suggested below and possibly more, it has a potential to bring in a radical change in the

fundamental view as well as clinical practice of T2DM.

<u>Testability of assumptions and additional predictions</u>:

The assumption that the thresholds *PGT* and *PIT* can be modified by autonomous signaling

needs to be tested empirically. Although currently the thresholds are known to be flexible,

information about the conditions and mechanisms of change are poorly known.

The assumption of our model that reduced transport of glucose and insulin to brain is the

primary pathology of T2DM leading secondarily to hyperglycemia makes more predictions

that can be tested experimentally or epidemiologically. Experimentally specifically blocking

glut1 receptors in the brain should lead to hyperglycemia. Conversely infusion of glucose

directly to the brain should reduce peripheral hyperglycemia in the short run. This is already

suggested by some experiments (Ono et al., 1983; Osundiji et al., 2012). It is also

demonstrated that inducing primary endothelial dysfunction and reduced glucose transport

to brain by endothelial deletion of hypoxia inducible factor HIF-1α results in hyperglycemia

(Huang et al 2012). More careful research in this direction to reveal the cause effect

relationship between vascular defects, brain glucose levels and plasma glucose levels will be

enlightening. Epidemiologically hypoglycemia is shown to associate with dementia and

other neuronal problems (Lipska and Montori 2013, Meneilli and Tessier 2016, Rhee 2017,

Yaffe et al 2013), tight glycemic control led to higher mortality as compared to moderate

control in many of the clinical trials (Klein 2010, Schwartz & Meinert, 2004, The NICE-SUGAR

Study Investigators, 2009) which demands investigations into the causal pathways. The

question whether tight glycemic control leads to subtle neuronal changes in the long run as

expected by our model needs careful investigation. The difference between fasting and post

meal regression correlation parameters between glucose and insulin is an important

epidemiological line of evidence we have used. Chawla et al (2017) and Diwekar-Joshi and

Watve (2020) showed this pattern across four different data sets. How generalized the

pattern is can be easily tested in multiple populations.

On the modeling front it is necessary to undertake comparative evaluation of the different

models with respect to the battery of prediction that we listed here. Perhaps a few more

predictions may be added. However, at present many of the models and their possible

modifications are not explored sufficiently to see whether they can explain the currently

unexplained patterns under certain set of conditions. A model prediction matrix would be

an appropriate approach for such a comparative evaluation, but we may have to wait till all

alternative models are explored sufficiently elaborately on which of the empirical patterns

they predict, which ones they are compatible with and which ones they contradict. We have
shown here that the brain centered model predicts or is compatible with all the patterns
listed in the introduction and does not contradict any.

<u>Possible causes of T2DM</u>:

The classically believed causal factors namely insulin resistance and β cell dysfunction are
not compatible with many of the empirical findings as shown by Diwekar-Joshi and Watve
(2020). In our model change in insulin resistance and β cell dysfunction were neither
necessary nor sufficient to account for all the patterns. Nevertheless they were helpful in
accounting for the altered glucose tolerance curve, although other factors could also
account for it independent of insulin resistance. Incorporating insulin resistance in our
model and assuming it to work only in the post feeding state, patterns 1,2,5,6,9,16 and 17
could be explained but not others. In short, our model does not rule out insulin resistance as
a phenomenon, but implies that it may not be central to T2DM. Primary vasculature
impairment reducing glucose and insulin transport in mutually correlated or uncorrelated
manner could explain all patterns and therefore makes the most parsimonious causal
hypothesis. B cell dysfunction is an inevitable effect of higher degree of vascular dysfunction
and therefore may not be needed as an independent causal factor.

Being open to alternative possibilities is an important virtue of science and it is particularly
important when the mainstream thinking has limited success and mounting anomalous
findings. Prevention of T2DM on a global scale has largely failed and treatment has limited
and inconsistent success in arresting mortality and morbidities associated with T2DM
(Brown et al., 2004; DeFronzo, 2010; Rosengren et al., 2008, Lee et al 2021). Therefore,
exploring alternative possible interpretations is a need of the time.

Triggered by multiple anomalies in the theory of glucose homeostasis and the origins of
diabetic hyperglycemia, we have articulated here an alternative paradigm which potentially
resolves in a logically and mathematically consistent manner many of the anomalous
findings. Being mathematically and logically sound and compatible with evidence is not a
proof of a theory but certainly reflects on its potential to develop into a new alternative
paradigm. For a large field such as T2DM multiple efforts would be needed to evaluate
competing paradigms. We have suggested a few more testable predictions that can help in

this task. There can be more possible ways of testing them which should come to light and

used to reach robust conclusions which have the potential to change the clinical course of

prevention as well as treatment of an important global health problem.

If our proposed causal inference that vascular dysfunction is primary to T2DM is supported

by more careful investigations, it explains the failure of glucose normalizing treatment in

arresting diabetic complications. Simultaneously it suggests alternative lines of treatments

which should focus on normalizing vascular and neuronal function rather than focusing on

glucose normalization. The deficiencies of stimuli normally required for growth factors and

other angiogenic and neuroprotective factors can be forecasted as the best bet for the new

approach. However, rigorous efforts are needed to strengthen the evidence base for

selecting the right one amongst the alternative paradigms.

References:

Ajmera, I., Swat, M., Laibe, C., Le Novère, N., & Chelliah, V. (2013). The impact of

mathematical modeling on the understanding of diabetes and related complications.

CPT: Pharmacometrics & Systems Pharmacology, *2*(7), 54.

https://doi.org/10.1038/psp.2013.30

Aloe, L., Bracci-Laudiero, L., Alleva, E., Lambiase, A., Micera, A., & Tirassa, P. (1994).

Emotional stress induced by parachute jumping enhances blood nerve growth factor

levels and the distribution of nerve growth factor receptors in lymphocytes.

Proceedings of the National Academy of Sciences, *91*(22), 10440–10444.

https://doi.org/10.1073/pnas.91.22.10440

Antuna-Puente, B., Disse, E., Lavoie, M.-E., Rabasa-Lhoret, R., Laville, M., & Bastard, J.-P.

(2009). Evaluation of the SIisOGTT index in a healthy population with normal glucose

tolerance. *Diabetes & Metabolism*, *35*(3), 236–237.

https://doi.org/10.1016/j.diabet.2009.02.004

Bacha, F., Gungor, N., Lee, S., & Arslanian, S. A. (2013). Progressive deterioration of β-cell

function in obese youth with type 2 diabetes. *Pediatric Diabetes*, *14*(2), 106–111.

https://doi.org/10.1111/j.1399-5448.2012.00915.x

805 Banting, F. G., Best, C. H., Collip, J. B., Campbell, W. R., & Fletcher, A. A. (1922). Pancreatic

806   Extracts in the Treatment of Diabetes Mellitus. *Canadian Medical Association Journal*,

807   *12*(3), 141–146. https://www.ncbi.nlm.nih.gov/pubmed/20314060

808 Begg, D. P. (2015). *Insulin Transport into the Brain and Cerebrospinal Fluid* (pp. 229–248).

809   https://doi.org/10.1016/bs.vh.2014.12.007

810 Begg, D. P., & Woods, S. C. (2013). Interactions between the central nervous system and

811   pancreatic islet secretions: a historical perspective. *Advances in Physiology Education*,

812   *37*(1), 53–60. https://doi.org/10.1152/advan.00167.2012

813 Bernard, C. (1859). *Leçons sur les phénomènes de la vie. Cours de physiologies generale du*

814   *Museum d'Histoir Naturelle*. Paris: Librairie Delagrave.

815 Bingham, E. M., Hopkins, D., Smith, D., Pernet, A., Hallett, W., Reed, L., Marsden, P. K., &

816   Amiel, S. A. (2002). The Role of Insulin in Human Brain Glucose Metabolism. *Diabetes*,

817   *51*(12), 3384–3390. https://doi.org/10.2337/diabetes.51.12.3384

818 Blüher, M., Michael, M. D., Peroni, O. D., Ueki, K., Carter, N., Kahn, B. B., & Kahn, C. R.

819   (2002). Adipose tissue selective insulin receptor knockout protects against obesity and

820   obesity-related glucose intolerance. *Developmental Cell*, *3*(1), 25–38.

821   https://doi.org/10.1016/S1534-5807(02)00199-5

822 Boado, R. J., Black, K. L., & Pardridge, W. M. (1994). Gene expression of GLUT3 and GLUT1

823   glucose transporters in human brain tumors. *Molecular Brain Research*, *27*(1), 51–57.

824   https://doi.org/10.1016/0169-328X(94)90183-X

825 Boland, B. B., Rhodes, C. J., & Grimsby, J. S. (2017). The dynamic plasticity of insulin

826   production in β-cells. *Molecular Metabolism*, *6*(9), 958–973.

827   https://doi.org/10.1016/j.molmet.2017.04.010

828 Brown, J. B., Nichols, G. A., & Perry, A. (2004). The Burden of Treatment Failure in Type 2

829   Diabetes. *Diabetes Care*, *27*(7), 1535–1540. https://doi.org/10.2337/diacare.27.7.1535

830 Büsing, K. A., Schönberg, S. O., Brade, J., & Wasser, K. (2013). Impact of blood glucose,

831   diabetes, insulin, and obesity on standardized uptake values in tumors and healthy

832   organs on 18F-FDG PET/CT. *Nuclear Medicine and Biology*, *40*(2), 206–213.

833 https://doi.org/10.1016/j.nucmedbio.2012.10.014

834 Butler, A. E., Janson, J., Bonner-Weir, S., Ritzel, R., Rizza, R. A., & Butler, P. C. (2003). -Cell

835   Deficit and Increased -Cell Apoptosis in Humans With Type 2 Diabetes. *Diabetes*, *52*(1),

836   102–110. https://doi.org/10.2337/diabetes.52.1.102

837 Campbell, I. W. (1976). The somogyi phenomenon. A short review. *Acta Diabetologica*

838   *Latina*, *13*(1–2), 68–73. https://doi.org/10.1007/BF02591583

839 Cao, L., Liu, X., Lin, E.-J. D., Wang, C., Choi, E. Y., Riban, V., Lin, B., & During, M. J. (2010).

840   Environmental and Genetic Activation of a Brain-Adipocyte BDNF/Leptin Axis Causes

841   Cancer Remission and Inhibition. *Cell*, *142*(1), 52–64.

842   https://doi.org/10.1016/j.cell.2010.05.029

843 Carmina, E., Stanczyk, F. Z., & Lobo, R. A. (2019). Evaluation of Hormonal Status. In *Yen and*

844   *Jaffe's Reproductive Endocrinology* (pp. 887-915.e4). Elsevier.

845   https://doi.org/10.1016/B978-0-323-47912-7.00034-2

846 Carnethon, M. R., Jacobs, D. R., Sidney, S., & Liu, K. (2003). Influence of Autonomic Nervous

847   System Dysfunction on the Development of Type 2 Diabetes: The CARDIA study.

848   *Diabetes Care*, *26*(11), 3035–3041. https://doi.org/10.2337/diacare.26.11.3035

849 Chawla, S., Pund, A., B., V., Kulkarni, S., Diwekar-Joshi, M., & Watve, M. (2018). Inferring

850   causal pathways among three or more variables from steady-state correlations in a

851   homeostatic system. *PLOS ONE*, *13*(10), e0204755.

852   https://doi.org/10.1371/journal.pone.0204755

853 Chen, N.-G., Tassava, T. M., & Romsos, D. R. (1993). Threshold for Glucose-Stimulated

854   Insulin Secretion in Pancreatic Islets of Genetically Obese (ob/ob) Mice is Abnormally

855   Low. *The Journal of Nutrition*, *123*(9), 1567–1574.

856   https://doi.org/10.1093/jn/123.9.1567

857 Chodari, L., Mohammadi, M., Ghorbanzadeh, V., Dariushnejad, H., & Mohaddes, G. (2016).

858   Testosterone and Voluntary Exercise Promote Angiogenesis in Hearts of Rats with

859   Diabetes by Enhancing Expression of VEGF-A and SDF-1a. *Canadian Journal of Diabetes*,

860   *40*(5), 436–441. https://doi.org/10.1016/j.jcjd.2016.03.004

Clark, A., Jones, L., de Koning, E., Hansen, B. C., & Matthews, D. R. (2001). Decreased insulin

secretion in type 2 diabetes: a problem of cellular mass or function? *Diabetes*,

50(Supplement 1), S169–S171. https://doi.org/10.2337/diabetes.50.2007.S169

Coggan, A. R. (1991). Plasma Glucose Metabolism During Exercise in Humans. *Sports*

Medicine, *11*(2), 102–124. https://doi.org/10.2165/00007256-199111020-00003

Corkey, B. E., & Shirihai, O. (2012). Metabolic master regulators: sharing information among

multiple systems. *Trends in Endocrinology & Metabolism*, *23*(12), 594–601.

https://doi.org/10.1016/j.tem.2012.07.006

Cornford, E. M., Hyman, S., Cornford, M. E., & Clare-Salzler, M. (1995). Down-regulation of

blood-brain glucose transport in the hyperglycemic nonobese diabetic mouse.

Neurochemical Research, *20*(7), 869–873. https://doi.org/10.1007/BF00969700

Costes, S., & Butler, P. C. (2014). Insulin-degrading enzyme inhibition, a novel therapy for

type 2 diabetes? In *Cell Metabolism* (Vol. 20, Issue 2).

https://doi.org/10.1016/j.cmet.2014.07.016

D'Alessio, D. A., Kieffer, T. J., Taborsky, G. J., & Havel, P. J. (2001). Activation of the

Parasympathetic Nervous System Is Necessary for Normal Meal-Induced Insulin

Secretion in Rhesus Macaques1. *The Journal of Clinical Endocrinology & Metabolism*,

86(3), 1253–1259. https://doi.org/10.1210/jcem.86.3.7367

DeFronzo, R. A. (2010). Overview of Newer Agents: Where Treatment Is Going. *The*

American Journal of Medicine, *123*(3), S38–S48.

https://doi.org/10.1016/j.amjmed.2009.12.008

DeFronzo, R. A., Ferrannini, E., Groop, L., Henry, R. R., Herman, W. H., Holst, J. J., Hu, F. B.,

Kahn, C. R., Raz, I., Shulman, G. I., Simonson, D. C., Testa, M. A., & Weiss, R. (2015).

Type 2 diabetes mellitus. *Nature Reviews Disease Primers*, *1*(1), 15019.

https://doi.org/10.1038/nrdp.2015.19

Diabetes Prevention Program Research Group. (2015). Long-term effects of lifestyle

intervention or metformin on diabetes development and microvascular complications

over 15-year follow-up: the Diabetes Prevention Program Outcomes Study. *The Lancet*

Diabetes & Endocrinology, *3*(11), 866–875. https://doi.org/10.1016/S2213-

8587(15)00291-0

Diwekar-Joshi, M., & Watve, M. (2020). Driver versus navigator causation in biology: The
case of insulin and fasting glucose. *PeerJ*, *8*. https://doi.org/10.7717/peerj.10396

Drouin, R., Lavoie, C., Bourque, J., Ducros, F., Poisson, D., & Chiasson, J.-L. (1998). Increased
hepatic glucose production response to glucagon in trained subjects. *American Journal*
of Physiology-Endocrinology and Metabolism, *274*(1), E23–E28.
https://doi.org/10.1152/ajpendo.1998.274.1.E23

Dungan, K. M., Braithwaite, S. S., & Preiser, J.-C. (2009). Stress hyperglycaemia. *The Lancet*,
373(9677), 1798–1807. https://doi.org/10.1016/S0140-6736(09)60553-5

Ferrannini, E., & DeFronzo, R. A. (2015). Impact of glucose-lowering drugs on cardiovascular
disease in type 2 diabetes. *European Heart Journal*, *36*(34).
https://doi.org/10.1093/eurheartj/ehv239

Gaohua, L., & Kimura, H. (2009). A mathematical model of brain glucose homeostasis.
Theoretical Biology and Medical Modelling, *6*(1), 26. https://doi.org/10.1186/1742-
4682-6-26

Ge, X., Chen, C., Hui, X., Wang, Y., Lam, K. S. L., & Xu, A. (2011). Fibroblast Growth Factor 21
Induces Glucose Transporter-1 Expression through Activation of the Serum Response
Factor/Ets-Like Protein-1 in Adipocytes. *Journal of Biological Chemistry*, *286*(40),
34533–34541. https://doi.org/10.1074/jbc.M111.248591

Gilon, P., & Henquin, J.-C. (2001). Mechanisms and Physiological Significance of the
Cholinergic Control of Pancreatic β-Cell Function. *Endocrine Reviews*, *22*(5), 565–604.
https://doi.org/10.1210/edrv.22.5.0440

Giustina, A., Girelli, A., Buffoli, M. G., Cimino, A., Legati, F., Valentini, U., & Giustina, G.
(1991). Low-dose octreotide is able to cause a maximal inhibition of the glycemic
responses to a mixed meal in obese type 2 diabetic patients treated with insulin.
Diabetes Research and Clinical Practice, *14*(1). https://doi.org/10.1016/0168-
8227(91)90052-F

Gjedde, A., & Crone, C. (1981). Blood-Brain Glucose Transfer: Repression in Chronic

Hyperglycemia. *Science*, *214*(4519), 456–457. https://doi.org/10.1126/science.7027439

Gray, S. M., & Barrett, E. J. (2018). Insulin transport into the brain. *American Journal of*

Physiology-Cell Physiology, *315*(2), C125–C136.

https://doi.org/10.1152/ajpcell.00240.2017

Gray, S. M., Meijer, R. I., & Barrett, E. J. (2014). Insulin regulates brain function, but how

does it get there? *Diabetes*, *63*(12), 3992–3997. https://doi.org/10.2337/db14-0340

Gruetter, R., Novotny, E. J., Boulware, S. D., Rothman, D. L., Mason, G. F., Shulman, G. I.,

Shulman, R. G., & Tamborlane, W. V. (1992). Direct measurement of brain glucose

concentrations in humans by 13C NMR spectroscopy. *Proceedings of the National*

Academy of Sciences, *89*(3), 1109–1112. https://doi.org/10.1073/pnas.89.3.1109

Hasselbalch, S. G., Knudsen, G. M., Videbaek, C., Pinborg, L. H., Schmidt, J. F., Holm, S., &

Paulson, O. B. (1999). No effect of insulin on glucose blood-brain barrier transport and

cerebral metabolism in humans. *Diabetes*, *48*(10), 1915–1921.

https://doi.org/10.2337/diabetes.48.10.1915

Henquin, J.-C., Dufrane, D., & Nenquin, M. (2006). Nutrient Control of Insulin Secretion in

Isolated Normal Human Islets. *Diabetes*, *55*(12), 3470–3477.

https://doi.org/10.2337/db06-0868

Höppener, J. W. M., Oosterwijk, C., Nieuwenhuis, M. G., Posthuma, G., Thijssen, J. H. H.,

Vroom, T. M., Ahrén, B., & Lips, C. J. M. (1999). Extensive islet amyloid formation is

induced by development of Type II diabetes mellitus and contributes to its progression:

pathogenesis of diabetes in a mouse model. *Diabetologia*, *42*(4), 427–434.

https://doi.org/10.1007/s001250051175

Huang Y, Lei L, Liu D, Jovin I, Russell R, Johnson RS, Di Lorenzo A, Giordano FJ. Normal

glucose uptake in the brain and heart requires an endothelial cell-specific HIF-1α-

dependent function. Proc Natl Acad Sci U S A. 2012 Oct 23;109(43):17478-83. doi:

10.1073/pnas.1209281109. Epub 2012 Oct 9. PMID: 23047702; PMCID: PMC3491491.

Hwang, J. J., Jiang, L., Hamza, M., Sanchez Rangel, E., Dai, F., Belfort-DeAguiar, R., Parikh, L.,

Koo, B. B., Rothman, D. L., Mason, G., & Sherwin, R. S. (2017). Blunted rise in brain

glucose levels during hyperglycemia in adults with obesity and T2DM. *JCI Insight*, *2*(20).

https://doi.org/10.1172/jci.insight.95913

Sorensen, John Thomas (1985). A physiologic model of glucose metabolism in man and its
use to design and assess improved insulin therapies for diabetes. Thesis (Sc. D.)
Massachusetts Institute of Technology, Dept. of Chemical Engineering.

Kadowaki, T. (2000). Insights into insulin resistance and type 2 diabetes from knockout
mouse models. *Journal of Clinical Investigation*, *106*(4), 459–465.
http://www.ncbi.nlm.nih.gov/pmc/articles/PMC380257/

Kern, W., Peters, A., Fruehwald-Schultes, B., Deininger, E., Born, J., & Fehm, H. L. (2001).
Improving Influence of Insulin on Cognitive Functions in Humans. *Neuroendocrinology*,
74(4), 270–280. https://doi.org/10.1159/000054694

Kiba, T. (2004). Relationships Between the Autonomic Nervous System and the Pancreas
Including Regulation of Regeneration and Apoptosis. *Pancreas*, *29*(2), e51–e58.
https://doi.org/10.1097/00006676-200408000-00019

King, P., Peacock, I., & Donnelly, R. (1999). The UK prospective diabetes study (UKPDS):
clinical and therapeutic implications for type 2 diabetes. *British Journal of Clinical*
Pharmacology, *48*(5), 643–648. https://doi.org/10.1046/j.1365-2125.1999.00092.x

Klein, R. (2010). Intensive treatment of hyperglycaemia: ACCORD. *The Lancet*, *376*(9739),
391–392. https://doi.org/10.1016/S0140-6736(10)61028-8

Krishnan, S., Chang, A. C., Stoltz, B. M., & Prasadarao, N. V. (2016). *Escherichia coli* K1
Modulates Peroxisome Proliferator–Activated Receptor γ and Glucose Transporter 1 at
the Blood-Brain Barrier in Neonatal Meningitis. *Journal of Infectious Diseases*, *214*(7),
1092–1104. https://doi.org/10.1093/infdis/jiw306

Laeger, T., Baumeier, C., Wilhelmi, I., Würfel, J., Kamitz, A., & Schürmann, A. (2017). FGF21
improves glucose homeostasis in an obese diabetes-prone mouse model independent
of body fat changes. *Diabetologia*, *60*(11), 2274–2284. https://doi.org/10.1007/s00125-
017-4389-x

Lakshmanan, J. (1986). Aggressive behavior in adult male mice elevates serum nerve growth
factor levels. *American Journal of Physiology-Endocrinology and Metabolism*, *250*(4),
E386–E392. https://doi.org/10.1152/ajpendo.1986.250.4.E386

Lam, T. K. T. (2005). Regulation of Blood Glucose by Hypothalamic Pyruvate Metabolism.

Science, *309*(5736), 943–947. https://doi.org/10.1126/science.1112085

Lamberts, S. W. J., & Hofland, L. J. (2019). Octreotide, 40 years later. In *European Journal of*

Endocrinology (Vol. 181, Issue 5). https://doi.org/10.1530/EJE-19-0074

Leahy, J. L., Bumbalo, L. M., & Chen, C. (1994). Diazoxide causes recovery of β-cell glucose

responsiveness in 90% pancreatectomized diabetic rats. *Diabetes*, *43*(2).

https://doi.org/10.2337/diabetes.43.2.173

Lee, C. G., Heckman-Stoddard, B., Dabelea, D., Gadde, K. M., Ehrmann, D., Ford, L., Prorok,

P., Boyko, E. J., Pi-Sunyer, X., Wallia, A., Knowler, W. C., Crandall, J. P., & Temprosa, M.

(2021). Effect of Metformin and Lifestyle Interventions on Mortality in the Diabetes

Prevention Program and Diabetes Prevention Program Outcomes Study. *Diabetes Care*,

44(12), 2775–2782. https://doi.org/10.2337/dc21-1046

Lerner, R. L., & Porte, D. (1972). Acute and steady-state insulin responses to glucose in

nonobese diabetic subjects. *Journal of Clinical Investigation*, *51*(7), 1624–1631.

https://doi.org/10.1172/JCI106963

Limenta, M., Ho, C. S. C., Poh, J. W. W., Goh, S.-Y., & Toh, D. S. L. (2019). Adverse Drug

Reaction Profile of SGLT2 Inhibitor-Associated Diabetic Ketosis/Ketoacidosis in

Singapore and their Precipitating Factors. *Clinical Drug Investigation*, *39*(7), 683–690.

https://doi.org/10.1007/s40261-019-00794-5

Lipska KJ, Montori VM. (2013) Glucose control in older adults with diabetes mellitus--more

harm than good? JAMA Intern Med. 22;173(14):1306-7. doi:

10.1001/jamainternmed.2013.6189. PMID: 23753259

Liu, M., Cao, H., Hou, Y., Sun, G., Li, D., & Wang, W. (2018). Liver Plays a Major Role in FGF-

21 Mediated Glucose Homeostasis. *Cellular Physiology and Biochemistry*, *45*(4), 1423–

1433. https://doi.org/10.1159/000487568

Lundqvist, M. H., Almby, K., Abrahamsson, N., & Eriksson, J. W. (2019). Is the Brain a Key

Player in Glucose Regulation and Development of Type 2 Diabetes? *Frontiers in*

Physiology, *10*. https://doi.org/10.3389/fphys.2019.00457

Maianti, J. P., McFedries, A., Foda, Z. H., Kleiner, R. E., Du, X. Q., Leissring, M. A., Tang, W.-J.,

Charron, M. J., Seeliger, M. A., Saghatelian, A., & Liu, D. R. (2014). Anti-diabetic activity

of insulin-degrading enzyme inhibitors mediated by multiple hormones. *Nature*,

511(7507), 94–98. https://doi.org/10.1038/nature13297

Mari, A., Tura, A., Grespan, E., & Bizzotto, R. (2020a). Mathematical Modeling for the

Physiological and Clinical Investigation of Glucose Homeostasis and Diabetes   . In

Frontiers in Physiology   (Vol. 11, p. 1548).

https://www.frontiersin.org/article/10.3389/fphys.2020.575789

Mari, A., Tura, A., Grespan, E., & Bizzotto, R. (2020b). Mathematical Modeling for the

Physiological and Clinical Investigation of Glucose Homeostasis and Diabetes. *Frontiers*

in Physiology, *11*. https://doi.org/10.3389/fphys.2020.575789

Marliss, E. B., & Vranic, M. (2002). Intense Exercise Has Unique Effects on Both Insulin

Release and Its Roles in Glucoregulation: Implications for Diabetes. *Diabetes*,

51(Supplement 1), S271–S283. https://doi.org/10.2337/diabetes.51.2007.S271

Matsuda, M., Kawasaki, F., Mikami, Y., Takeuchi, Y., Saito, M., Eto, M., & Kaku, K. (2002).

Rescue of β-cell exhaustion by diazoxide after the development of diabetes mellitus in

rats with streptozotocin-induced diabetes. *European Journal of Pharmacology*, *453*(1).

https://doi.org/10.1016/S0014-2999(02)02389-0

Matthews, D. R., Hosker, J. P., Rudenski, A. S., Naylor, B. A., Treacher, D. F., & Turner, R. C.

(1985). Homeostasis model assessment: insulin resistance and β-cell function from

fasting plasma glucose and insulin concentrations in man. *Diabetologia*, *28*(7).

https://doi.org/10.1007/BF00280883

Mehran, A. E., Templeman, N. M., Brigidi, G. S., Lim, G. E., Chu, K.-Y., Hu, X., Botezelli, J. D.,

Asadi, A., Hoffman, B. G., Kieffer, T. J., Bamji, S. X., Clee, S. M., & Johnson, J. D. (2012a).

Hyperinsulinemia Drives Diet-Induced Obesity Independently of Brain Insulin

Production. *Cell Metabolism*, *16*(6), 723–737.

https://doi.org/10.1016/j.cmet.2012.10.019

Mehran, A. E., Templeman, N. M., Brigidi, G. S., Lim, G. E., Chu, K. Y., Hu, X., Botezelli, J. D.,

Asadi, A., Hoffman, B. G., Kieffer, T. J., Bamji, S. X., Clee, S. M., & Johnson, J. D. (2012b).

Hyperinsulinemia drives diet-induced obesity independently of brain insulin

production. *Cell Metabolism*, *16*(6). https://doi.org/10.1016/j.cmet.2012.10.019

Meneilly GS, Tessier DM. (2016) Diabetes, Dementia and Hypoglycemia. Can J Diabetes.

2016 Feb;40(1):73-6. doi: 10.1016/j.jcjd.2015.09.006. PMID: 26778684.

Miller, R., Waid, T., & Joyce, M. (1976). Direct neural inhibition of insulin secretion in

response to systemic hypoglycemia. *American Journal of Physiology-Legacy Content*,

230(4), 1090–1094. https://doi.org/10.1152/ajplegacy.1976.230.4.1090

Mistry, S., & Eschler, D. C. (2021). Euglycemic Diabetic Ketoacidosis Caused by SGLT2

Inhibitors and a Ketogenic Diet: A Case Series and Review of Literature. *AACE Clinical*

Case Reports, *7*(1), 17–19. https://doi.org/10.1016/j.aace.2020.11.009

Muris, D. M. J., Houben, A. J. H. M., Schram, M. T., & Stehouwer, C. D. A. (2012).

Microvascular Dysfunction Is Associated With a Higher Incidence of Type 2 Diabetes

Mellitus. *Arteriosclerosis, Thrombosis, and Vascular Biology*, *32*(12), 3082–3094.

https://doi.org/10.1161/ATVBAHA.112.300291

Nelles, E., Butzler, C., Jung, D., Temme, A., Gabriel, H. D., Dahl, U., Traub, O., Stumpel, F.,

Jungermann, K., Zielasek, J., Toyka, K. V, Dermietzel, R., & Willecke, K. (1996). Defective

propagation of signals generated by sympathetic nerve stimulation in the liver of

connexin32-deficient mice. *Proceedings of the National Academy of Sciences*, *93*(18),

9565–9570. https://doi.org/10.1073/pnas.93.18.9565

NEXØ, E., HOLLENBERG, M. D., & BING, J. (1981). Aggressive behaviour in mice provokes a

marked increase in both plasma epidermal growth factor and renin. *Acta Physiologica*

Scandinavica, *111*(3), 367–371. https://doi.org/10.1111/j.1748-1716.1981.tb06749.x

Nexø, E., Olsen, P. S., & Poulsen, K. (1984). Exocrine and endocrine secretion of renin and

epidermal growth factor from the mouse submandibular glands. *Regulatory Peptides*,

8(4), 327–334. https://doi.org/10.1016/0167-0115(84)90043-0

Nguyen, T. T., Wang, J. J., & Wong, T. Y. (2007). Retinal Vascular Changes in Pre-Diabetes

and Prehypertension. *Diabetes Care*, *30*(10), 2708–2715. https://doi.org/10.2337/dc07-

0732

Nonogaki, K. (2000). New insights into sympathetic regulation of glucose and fat metabolism. *Diabetologia*, *43*(5), 533–549. https://doi.org/10.1007/s001250051341

ONO, T., STEFFENS, A., & SASAKI, K. (1983). Influence of peripheral and intracerebroventricular glucose and insulin infusions on peripheral and cerebrospinal fluid glucose and insulin levels. *Physiology & Behavior*, *30*(2), 301–306. https://doi.org/10.1016/0031-9384(83)90023-9

Op den Kamp, Y. J. M., de Ligt, M., Dautzenberg, B., Kornips, E., Esterline, R., Hesselink, M. K. C., Hoeks, J., Schrauwen-Hinderling, V. B., Havekes, B., Oscarsson, J., Phielix, E., & Schrauwen, P. (2021). Effects of the SGLT2 Inhibitor Dapagliflozin on Energy Metabolism in Patients With Type 2 Diabetes: A Randomized, Double-Blind Crossover Trial. *Diabetes Care*, *44*(6), 1334–1343. https://doi.org/10.2337/dc20-2887

Osundiji, M. A., Lam, D. D., Shaw, J., Yueh, C.-Y., Markkula, S. P., Hurst, P., Colliva, C., Roda, A., Heisler, L. K., & Evans, M. L. (2012). Brain Glucose Sensors Play a Significant Role in the Regulation of Pancreatic Glucose-Stimulated Insulin Secretion. *Diabetes*, *61*(2), 321–328. https://doi.org/10.2337/db11-1050

Pardridge, W. M., Triguero, D., & Farrell, C. R. (1990). Downregulation of Blood-Brain Barrier Glucose Transporter in Experimental Diabetes. *Diabetes*, *39*(9), 1040–1044. https://doi.org/10.2337/diab.39.9.1040

Perrin, C., Knauf, C., & Burcelin, R. (2004). Intracerebroventricular Infusion of Glucose, Insulin, and the Adenosine Monophosphate-Activated Kinase Activator, 5-Aminoimidazole-4-Carboxamide-1-β- <scp>d</scp> -Ribofuranoside, Controls Muscle Glycogen Synthesis. *Endocrinology*, *145*(9), 4025–4033. https://doi.org/10.1210/en.2004-0270

Pfützner, A., Klonoff, D., Heinemann, L., Ejskjaer, N., & Pickup, J. (2017). Euglycemic ketosis in patients with type 2 diabetes on SGLT2-inhibitor therapy—an emerging problem and solutions offered by diabetes technology. *Endocrine*, *56*(1), 212–216. https://doi.org/10.1007/s12020-017-1264-y

Pories, W. J., & Dohm, G. L. (2020). Letter to the Editor Re: Sera of Obese Type 2 Diabetic Patients Undergoing Metabolic Surgery Instead of Conventional Treatment Exert

Beneficial Effects on B Cell Survival and Function: Results of a Randomized Clinical Study. *Obesity Surgery*, *30*(9), 3603–3604. https://doi.org/10.1007/s11695-020-04665-3

Porte, D., & Kahn, S. E. (1989). Hyperproinsulinemia and Amyloid in NIDDM: Clues to Etiology of Islet -Cell Dysfunction? *Diabetes*, *38*(11), 1333–1336. https://doi.org/10.2337/diab.38.11.1333

Porte, D., & Kahn, S. E. (2001). B-cell dysfunction and failure in type 2 diabetes: potential mechanisms. *Diabetes*, *50*(Supplement 1), S160–S163. https://doi.org/10.2337/diabetes.50.2007.S160

Rebelos, E., Rinne, J. O., Nuutila, P., & Ekblad, L. L. (2021). Brain Glucose Metabolism in Health, Obesity, and Cognitive Decline—Does Insulin Have Anything to Do with It? A Narrative Review. *Journal of Clinical Medicine*, *10*(7), 1532. https://doi.org/10.3390/jcm10071532

Reyhanoglu, G., & Rehman, A. (2021). *Somogyi Phenomenon*.

Rohner-Jeanrenaud, F., & Jeanrenaud, B. (1983). [The central nervous system-endocrine pancreas axis]. *Annales d'endocrinologie*, *44*(4), 217–227.

Rhee SY. (2017) Hypoglycemia and Dementia. *Endocrinol Metab (Seoul)*.32(2):195-199. doi:10.3803/EnM.2017.32.2.195

Rosengren, A., Jing, X., Eliasson, L., & Renström, E. (2008). Why Treatment Fails in Type 2 Diabetes. *PLoS Medicine*, *5*(10), e215. https://doi.org/10.1371/journal.pmed.0050215

Schüler, R., Seebeck, N., Osterhoff, M. A., Witte, V., Flöel, A., Busjahn, A., Jais, A., Brüning, J. C., Frahnow, T., Kabisch, S., Pivovarova, O., Hornemann, S., Kruse, M., & Pfeiffer, A. F. H. (2018). VEGF and GLUT1 are highly heritable, inversely correlated and affected by dietary fat intake: Consequences for cognitive function in humans. *Molecular Metabolism*, *11*, 129–136. https://doi.org/10.1016/j.molmet.2018.02.004

Schut, E. S., Westendorp, W. F., de Gans, J., Kruyt, N. D., Spanjaard, L., Reitsma, J. B., & van de Beek, D. (2009). Hyperglycemia in bacterial meningitis: a prospective cohort study. *BMC Infectious Diseases*, *9*(1), 57. https://doi.org/10.1186/1471-2334-9-57

Schwartz, T. B., & Meinert, C. L. (2004). The UGDP Controversy: thirty-four years of
contentious ambiguity laid to rest. *Perspectives in Biology and Medicine*, *47*(4), 564–
574. https://doi.org/10.1353/pbm.2004.0071

Shanik, M. H., Xu, Y., Skrha, J., Dankner, R., Zick, Y., & Roth, J. (2008). Insulin Resistance and
Hyperinsulinemia: Is hyperinsulinemia the cart or the horse? *Diabetes Care*,
31(Supplement 2), S262–S268. https://doi.org/10.2337/dc08-s264

Shemesh, E., Rudich, A., Harman-Boehm, I., & Cukierman-Yaffe, T. (2012). Effect of
Intranasal Insulin on Cognitive Function: A Systematic Review. *The Journal of Clinical
Endocrinology & Metabolism*, *97*(2), 366–376. https://doi.org/10.1210/jc.2011-1802

Shi, J., Dong, B., Mao, Y., Guan, W., Cao, J., Zhu, R., & Wang, S. (2016). Review: Traumatic
brain injury and hyperglycemia, a potentially modifiable risk factor. *Oncotarget*, *7*(43),
71052–71061. https://doi.org/10.18632/oncotarget.11958

Stehouwer, C. D. A. (2018). Microvascular Dysfunction and Hyperglycemia: A Vicious Cycle
With Widespread Consequences. *Diabetes*, *67*(9), 1729–1741.
https://doi.org/10.2337/dbi17-0044

Strachan, M W J. (2005). Insulin and cognitive function in humans: experimental data and
therapeutic considerations. *Biochemical Society Transactions*, *33*(Pt 5), 1037–1040.
https://doi.org/10.1042/BST20051037

Strachan, Mark W J. (2003). Insulin and cognitive function. *The Lancet*, *362*(9392), 1253.
https://doi.org/10.1016/S0140-6736(03)14615-6

Templeman, N. M., Clee, S. M., & Johnson, J. D. (2015). Suppression of hyperinsulinaemia in
growing female mice provides long-term protection against obesity. *Diabetologia*,
58(10). https://doi.org/10.1007/s00125-015-3676-7

Thackeray, J. T., Beanlands, R. S., & Dasilva, J. N. (2012). Altered sympathetic nervous
system signaling in the diabetic heart: emerging targets for molecular imaging.
American Journal of Nuclear Medicine and Molecular Imaging, *2*(3), 314–334.

The NICE-SUGAR Study Investigators. (2009). Intensive versus Conventional Glucose Control
in Critically Ill Patients. *New England Journal of Medicine*, *360*(13), 1283–1297.

https://doi.org/10.1056/NEJMoa0810625

Thorens, B. (2015). GLUT2, glucose sensing and glucose homeostasis. *Diabetologia*, *58*(2),

221–232. https://doi.org/10.1007/s00125-014-3451-1

Tirassa, P., Triaca, V., Amendola, T., Fiore, M., & Aloe, L. (2003). EGF and NGF injected into

the brain of old mice enhance BDNF and ChAT in proliferating subventricular zone.

Journal of Neuroscience Research, *72*(5), 557–564. https://doi.org/10.1002/jnr.10614

Turner, R. C., Holman, R. R., Matthews, D., Hockaday, T. D. R., & Peto, J. (1979). Insulin

deficiency and insulin resistance interaction in diabetes: Estimation of their relative

contribution by feedback analysis from basal plasma insulin and glucose

concentrations. *Metabolism*, *28*(11), 1086–1096. https://doi.org/10.1016/0026-

0495(79)90146-X

Turner, R., Matthews, D., Neil, a., & Mcelroy, H. (1998). Tight blood pressure control and

risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38.

British Medical Journaledical, *317*, 703–713. https://doi.org/10.1136/bmj.317.7160.703

UK Prospective Diabetes Study (UKPDS) Group. (1998). *Intensive blood-glucose control with*

SU or insulin compared with conventional treatment (UKPDS 33) - Lancet 1998.

352(Ukpds 33), 837–853.

van Veen, K. E. B., Brouwer, M. C., van der Ende, A., & van de Beek, D. (2016). Bacterial

meningitis in diabetes patients: a population-based prospective study. *Scientific*

Reports, *6*(1), 36996. https://doi.org/10.1038/srep36996

Watve, M. (2012). Doves, Diplomats, and Diabetes. In *Doves, Diplomats, and Diabetes*.

https://doi.org/10.1007/978-1-4614-4409-1

Watve, M. (2013). *Doves, Diplomats, and Diabetes*. Springer New York.

https://doi.org/10.1007/978-1-4614-4409-1

Watve, M., Bodas, A., & Diwekar, M. (2014). Altered autonomic inputs as a cause of

pancreatic β-cell amyloid. *Medical Hypotheses*, *82*(1), 49–53.

https://doi.org/10.1016/j.mehy.2013.11.002

Watve, M., & Diwekar-Joshi, M. (2019). Does insulin signalling decide glucose levels in the

fasting steady state? *BioRxiv*, 553016. https://doi.org/10.1101/553016

Weyer, C., Hanson, R. L., Tataranni, P. A., Bogardus, C., & Pratley, R. E. (2000). A high fasting
plasma insulin concentration predicts type 2 diabetes independent of insulin
resistance: evidence for a pathogenic role of relative hyperinsulinemia. *Diabetes*,
49(12), 2094–2101. https://doi.org/10.2337/diabetes.49.12.2094

Wiebe, N., Ye, F., Crumley, E. T., Bello, A., Stenvinkel, P., & Tonelli, M. (2021). Temporal
Associations Among Body Mass Index, Fasting Insulin, and Systemic Inflammation.
JAMA Network Open, *4*(3), e211263.
https://doi.org/10.1001/jamanetworkopen.2021.1263

Yaffe K, Falvey CM, Hamilton N, Harris TB, Simonsick EM, Strotmeyer ES, Shorr RI, Metti A,
Schwartz AV (2013) Health ABC Study. Association between hypoglycemia and
dementia in a biracial cohort of older adults with diabetes mellitus. JAMA Intern Med.
2013 Jul 22;173(14):1300-6. doi: 10.1001/jamainternmed.2013.6176. PMID: 23753199;
PMCID: PMC4041621

Zaleska-Żmijewska, A., Piątkiewicz, P., Śmigielska, B., Sokołowska-Oracz, A., Wawrzyniak, Z.
M., Romaniuk, D., Szaflik, J., & Szaflik, J. P. (2017). Retinal Photoreceptors and
Microvascular Changes in Prediabetes Measured with Adaptive Optics (rtx1[TM]): A Case-
Control Study. *Journal of Diabetes Research*, *2017*, 1–9.
https://doi.org/10.1155/2017/4174292

1193

1194