Midbrain somatostatin-expressing cells control pain-suppression during defensive states

Nanci Winke[1,2§✉], Frank Aby[3§], Daniel Jercog[1,2§], Thomas Bienvenu[1,2,4], Coline Riffault[1,2], Rabia Bouali-Benazzouz[3], Juliette Viellard[1,2,3], Delphine Girard[1,2], Zoé Grivet[3], Marc Landry[3], Laia Castell[5], Emmanuel Valjent[5], Stephane Valerio[1,2*], Pascal Fossat[3*] and Cyril Herry[1,2*✉]

[1]Univ. Bordeaux, Neurocentre Magendie, U1215, 146 Rue Léo-Saignat, 33077 Bordeaux, France

[2]INSERM, Neurocentre Magendie, U1215, 146 Rue Léo-Saignat, 33077 Bordeaux, France.

[3]Interdisciplinary Institute for Neuroscience (IINS, CNRS UMR 5297), University of Bordeaux, Bordeaux Cedex, France.

[4]Centre de Référence Régional des Pathologies Anxieuses et de la Dépression, Pôle de Psychiatrie Générale et Universitaire, Centre Hospitalier Charles Perrens, Bordeaux, France.

[5]Université de Montpellier, Institut de génomique fonctionnelle (IGF), CNRS, Inserm, 141 rue de la Cardonille, 34094 Montpellier, France.

§ shared authorship
* shared seniority
✉ corresponding authors: nanci.winke@inserm.fr; cyril.herry@inserm.fr

Abstract

Upon threatening situations, animals exhibit a broad range of behavioral and autonomic responses. As such, a crucial adaptive response is the inhibition of pain to facilitate relevant defensive behaviors that promote survival. Whereas the structures and mechanisms involved in fear and pain behavior are well documented, little is known about the precise neuronal mechanisms mediating the emotional regulation of endogenous pain-suppression. Here we used a combination of behavioral, anatomical, optogenetic, and electrophysiological approaches to investigate the role of that somatostatin-expressing cells in the ventrolateral periaqueductal gray matter (SST$^+$ vlPAG cells) in the control of analgesia induced during defensive states. Our data indicate that the optogenetic inhibition of SST$^+$ vlPAG cells

33 promotes analgesia irrespective of animal's defensive state. In contrast, the optogenetic

34 activation of long-range SST[+] vlPAG cells that project to the rostral ventromedial medulla

35 (RVM) abolishes the analgesia mediated by fear behavior. Together these results identify a

36 novel circuit mechanism composed of long-range SST[+] vlPAG cells projecting to the RVM

37 that regulate analgesia elicited during defensive states.

38

Introduction

39

40 During threatening situations, the inhibition of pain is an essential adaptive response that can be

41 crucial to promote self-preservation[1–4]. The inhibition of pain responses to a stimulus that would

42 typically elicit pain is defined as analgesia[5]. In addition, previous studies have described that a

43 fearful event can reduce pain sensitivity, a phenomenon referred to as fear-conditioned analgesia

44 (FCA)[6]. In particular, the pharmacological inactivation of the basolateral amygdala abolishes such

45 fear-modulated analgesia, and is associated with an increase in the immediate-early gene c-Fos

46 expression in the central amygdala and periaqueductal gray matter (PAG)[7]. The PAG is a midbrain

47 structure receiving many cortical and subcortical inputs from brain structures involved in fear

48 processing[8–11], projecting to the spinal cord primarily through a brainstem relay in the rostral

49 ventromedial medulla (RVM)[12], a key structure involved in nociceptive transmission[13,14]. Indeed,

50 the ventrolateral part of the PAG (vlPAG) is crucial for the descending control of pain[13,25] in the

51 dorsal horn of the spinal cord (DH)[12], and several reports indicate that electrical stimulation of the

52 vlPAG selectively inhibits responses to noxious stimuli in a variety of pain test conditions[12].

53 Moreover, analgesia induced upon vlPAG stimulation is opioid-dependent[15–17] and involves direct

54 descending projections to the RVM and the DH[18–20,26]. Furthermore, previous research indicate that

55 vlPAG activation induces analgesia through the recruitment of local GABAergic cells[21], which is

consistent with our knowledge of the role of the GABAergic system in the descending control of

pain[22,23]. Thus, the vlPAG is ideally located to allow the emotional regulation of pain behavior, yet

the underlying neuronal circuits and mechanisms are largely unknown. Here we investigated the

precise neural circuits within the vlPAG that mediates the suppression of pain behavior during

threatening situations. We developed a unique behavioral paradigm where a cued-elicited defensive

state promotes changes in pain sensitivity. Combining behavioral, anatomical, optogenetic, and

electrophysiological approaches, we identified a novel midbrain to brainstem circuit mechanism

composed of somatostatin positive (SST$^+$) vlPAG cells projecting to the RVM that is critically

involved in the suppression of pain behavior during conditioned fear states.

Results

Cued-fear conditioning elicits analgesia

To evaluate the contribution of specific vlPAG cells populations in the fear modulation of pain

behavior, we developed a novel cued fear-conditioned analgesia (CFCA) paradigm, consisting of

a pain sensitivity assay during the presentation of threat-predicting cues (**Figure 1a**). In this

paradigm, an initially neutral stimulus (the conditioned stimulus, CS$^+$) becomes associated with a

footshock (the unconditioned stimulus, US), whereas another neutral stimulus serves as a control

(CS$^-$). After conditioning, re-exposure to the CS$^+$ elicited freezing behavior, a behavioral readout

of associative learned fear[37] (**Figure 1b and Supplementary Figure 1a**). Following the fear

retrieval session, mice were submitted to the pain sensitivity assay. Mice were placed on a Hot-

Plate (HP) apparatus in which the floor plate temperature progressively increases while either CS$^-$

or CS$^+$ were concomitantly presented. Plate temperature increased, and CSs were continuously

played until mice displayed a nociceptive (NC) response reflecting pain behavior (**Figure 1c**),

defined by either jumping or hindpaw licking (**Figure 1d**). The detection of a NC response

terminated the test session (**see Methods**).

We observed a significant delay in the NC response performance for CS^+ compared to CS^-

presentations, reflecting that pain-suppression was selectively promoted by the threat-predicting

CS (**Figure 1e and Supplementary Figure 1b**). This increased NC latency during CS^+

presentation could alternatively be explained by a competition between freezing and pain-related

responses. However, freezing probability at the time of the NC response for CS^+ was remarkably

low and comparable to CS^- levels, suggesting that CS^+ increased NC response latency was not due

to competing freezing responses (**Figure 1f, see Methods**). Instead, these results suggest that fear

induced by CS^+ presentations drives the increase in NC latency. Importantly, such states induced

by our CFCA paradigm reflected fear associative processes (**Supplementary Figure 2**). To

demonstrate a causal link between cued-fear and analgesia, mice were submitted to an extinction

procedure to devaluate the CS^+-US contingency and abolish the negative emotional state during

CS^+ presentations (**Figure 1g**). In these conditions, CS^+ presentation after extinction had no effects

on NC response latency (**Figure 1h**). Furthermore, the absence of auditory cues or the presence of

auditory cues not associated with a threat had comparable NC response latencies as those observed

during CS^- presentations (**Figure 1i**). Finally, previous studies indicate that fear can lead to

vasoconstriction, which could result in a redirection of the blood flow to the skeletal musculature,

decreasing the temperature in the extremities of the body[24]. However, monitoring the back and tail

temperature of mice submitted to the HP test did not reveal any difference between the CSs

presentations, suggesting that increased latency was not due to fear-induced vasoconstriction

(**Supplementary Figure 1c-e**). Overall, our results indicate that the presence of a conditioned

threat-predicting cue can selectively promote analgesia.

Photostimulation of SST⁺ vlPAG cells alters pain and fear behavior

To identify the cell types in the vlPAG mediating the analgesic effect in the CFCA task, we focused on GABAergic interneurons, which activity is suggested to be inhibited during analgesia through μ-opioid receptor-dependent mechanisms[12,27]. Using single-molecule fluorescent in situ hybridization (smFISH, **see Methods**), we confirmed that the majority of vlPAG SST mRNAs were expressed in inhibitory neurons (**Supplementary Figure 3**). As such, since somatostatin-expressing interneurons (SST) are the most abundant interneurons class in the vlPAG[28], we focused on this cell population. Thus, to evaluate the contribution of SST⁺ vlPAG cells in CFCA, SST-Cre mice were injected in the vlPAG with a Cre-dependent AAV expressing the Channelrhodopsin (ChR2), Archaerhodopsin (ArchT), or GFP while optic fibers were bilaterally implanted targeting this area (**Figure 2a**). Next, mice were submitted to the CFCA task while photostimulating SST⁺ cells in the vlPAG (**Figure 2b**; **see Methods**). After fear retrieval (**Figure 2c, e**; **Supplementary Figure 4a-d**), the optogenetic activation of SST⁺ vlPAG cells during the HP test had no effect on CS⁻ presentation, whereas the same manipulation performed during CS⁺ blocked the increase in NC response latency compared to GFP controls (**Figure 2d and Supplementary Figure 5a**). In contrast, optogenetic inhibition of SST⁺ vlPAG cells increased NC response latency for both CS⁻ and CS⁺ conditions compared to GFP controls (**Figure 2f and Supplementary Figure 5b**). Importantly, these changes in NC response latencies were not due to motor or aversive effects of the optogenetic manipulations, since photoactivation of SST⁺ vlPAG cells did not induce changes in distance travelled in an open field nor place aversion in a real-time place preference test (**Supplementary Figure 5c-f**). In addition, we did not observe changes in fear learning or expression when the footshock was replaced or paired with the SST⁺ vlPAG cells photoactivation during the fear conditioning procedure (**Supplementary Figure 6**).

Altogether, these experiments confirmed that SST$^+$ vlPAG activation did not promote pain

responses and instead suggested that the SST$^+$ vlPAG cells are key elements of an endogenous

pain-suppression circuit. Additionally, the optogenetic effect observed during CFCA was not due

to alteration of somatostatin levels in our homozygous SST-Cre mice[29], as we observed the same

effect in heterozygous SST-Cre mice (**Supplementary Figure 7a-d**). Finally, in contrast with

STT$^+$ vlPAG cells, the optogenetic inhibition of another important class of vlPAG inhibitory cells

expressing the vasoactive intestinal peptide (VIP$^+$) did not change the NC response latencies during

CS$^-$ or CS$^+$ presentations (**Supplementary Figure 7e-h**).

Because our CFCA paradigm is dependent on fear associative processes (**Figure 1 and**

Supplementary Figure 1-2), it is possible that the optogenetic manipulation of SST$^+$ vlPAG cells

may have altered the expression of fear behavior. To control for this possibility, we optogenetically

activated or inhibited SST$^+$ vlPAG cells during a fear retrieval session following auditory fear

conditioning (**Figure 3a, b**). Our data indicate that whereas the optogenetic inhibition of SST$^+$

vlPAG cells had no effect on freezing behavior relative to GFP controls, their optogenetic

activation reduced freezing levels (**Figure 3c, d**). Altogether, our data show that the local SST$^+$

vlPAG network can modulate both fear expression and the analgesia promoted during defensive

states.

Photostimulation of SST$^+$ vlPAG cells regulates spinal cord nociceptive transmission

Consistent with our observations, previous studies found that the vlPAG is involved in fear-

conditioned analgesia[7]. It is conceivable that the SST$^+$ vlPAG cells are indeed part of a circuit

involved in the fear regulation of endogenous pain-suppression. To evaluate the effect of the

manipulation of SST$^+$ vlPAG cells on analgesia independently of the expression of fear behavior,

we evaluated the impact of SST$^+$ vlPAG cells manipulation on the control of nociceptive

transmission directly at the level of the spinal cord DH. Therefore, we performed extracellular local

field potential and single-unit recordings in the spinal cord DH during optogenetic stimulation of

SST$^+$ vlPAG cells in anesthetized mice (**Figure 4a**). Consistent with previous studies[30], nociceptive

stimulation induces early and late neuronal responses at the level of the spinal cord DH (A-

comprising non-nociceptive Aβ and nociceptive Aδ and C- fiber mediated, respectively; **Figure**

4a). While optogenetic activation of SST$^+$ vlPAG cells during paw suprathreshold electrical

stimulation increased the magnitude of late nociceptive field potentials, the optogenetic inhibition

reduced it (**Figure 4b, c**). To determine if this effect was specific to the nociceptive network, we

recorded wide dynamic range (WDR) cells in the DH, which receive both tactile (Aβ) and

nociceptive information (Aδ and C)[30]. We observed that SST$^+$ vlPAG cells photoactivation

increases both fast nociceptive (Aδ-fiber) and slow nociceptive (C-fiber) responses without

affecting Aβ (**Figure 4d**). Moreover, the optogenetic inhibition of SST$^+$ vlPAG cells specifically

inhibited nociceptive responses while Aβ responses remained unaffected (**Figure 4e**). In addition,

we performed subliminal electrical stimulations that did not elicit nociceptive neuronal responses.

In this condition, we observed that the removal of analgesia by the activation of SST$^+$ vlPAG cells

leads to WDR responses at latencies corresponding to nociceptive Aδ and C-fibers (**Figure 4f**).

Finally, the optogenetic activation of SST$^+$ vlPAG cells had no effect on purely tactile fast latency

spinal neurons (**Supplementary Figure 8**). Consistent with our optogenetic manipulations during

the CFCA task (**Figure 2**), these data suggest that under analgesia promoted by isoflurane,

activation and inhibition of SST$^+$ vlPAG cells increased or decreased nociceptive transmission by

specifically promoting pronociceptive and antinociceptive responses in the spinal cord DH,

respectively.

SST⁺ vlPAG-RVM-spinal cord pathway activation abolishes analgesia induced during defensive states.

We next evaluated whether SST⁺ vlPAG cells mediate their analgesia modulation by contacting WDR cells directly in the spinal cord or through the rostro ventromedial medulla (RVM), a structure known to receive direct projections from vlPAG and mediate pain modulation[18,19] (**Figure 5a**). Anatomical tracing revealed large labelling of SST⁺ fibers in RVM (**Figure 5b**) and only sparse labelling within the spinal cord (SC; data not shown). By combining fluorogold retrolabelling from the SC and GFP anterograde labelling from SST⁺ vlPAG cells (**Figure 5c**), we observed RVM cells projecting to the DH with close apposition of SST⁺ putative boutons (**Figure 5d**), suggesting that SST⁺ vlPAG cells project to the RVM and contact DH-projecting RVM cells. Optogenetic activation of SST descending fibers above the SC has no effect on WDR neural activity upon paw electrical stimulation, suggesting that SST⁺ vlPAG long-range projections to SC did not play a major role in endogenous pain-suppression (**Figure 5e**). In contrast, photoactivation of SST⁺ vlPAG inputs to the RVM increased both nociceptive (Aδ and C-fiber) responses (**Figure 5f, Supplementary Figure 9**). Altogether, these data revealed that the SST⁺ vlPAG-RVM-DH pathway mediates endogenous pain-suppression.

We next addressed if the SST⁺ vlPAG-RVM-DH pathway was specific in the mediation of endogenous pain-suppression induced by defensive states. Thus, SST-Cre mice were injected in the vlPAG with a Cre-dependent AAV expressing the ChR2 or GFP, and a single optic fiber was implanted above the RVM (**Figure 6a**). While photoactivation of the SST⁺ vlPAG-RVM pathway promoted nociception (**Figure 5f**), it did not impacted fear expression as freezing levels remained unchanged (**Figure 6b**). Consistently, activation of the SST⁺ vlPAG-RVM pathway during CFCA had no impact on CS⁻ presentation, whereas the same manipulation performed during CS⁺ blocked

196   the increase in NC response latency compared to GFP controls (**Figure 6c, d**). Overall, these data

197   suggest the existence of two populations of SST[+] vlPAG cells mediating fear expression and

198   analgesia induced by defensive states. (**Figure 6e**). Altogether, our results indicate that the SST[+]

199   vlPAG-RVM-DH pathway is critically involved in the analgesia induced during defensive states.

Discussion

In this study, we established that SST[+] vlPAG cells control the analgesia during defensive states.

Endogenous pain-suppression was mediated by long-range SST[+] vlPAG cells projecting to the

RVM, which in turn transmitted the pain signal to the DH of the spinal cord (**Figures 2 and 5**).

Finally, our data indicated that SST[+] vlPAG cells can regulate both fear expression and the

analgesia promoted during defensive states, and that the expression of these two behaviors can be

segregated by precisely targeting the SST[+] vlPAG-RVM pathway (**Figure 6**). Together these

results identify a novel midbrain to brainstem circuit mechanism composed of long-range SST[+]

vlPAG cells projecting to the RVM that regulate analgesia elicited during defensive states.

Our observations of the involvement of vlPAG in our cued-based fear conditioned analgesia

paradigm are consistent with previous studies in contextual-based FCA involving this region[7]. In

such FCA studies[6], animals undergo contextual fear conditioning followed by pain behavior testing

in the same conditioning context, limiting a temporally precise elicitation of aversive states.

Instead, our behavioral paradigm goes beyond these limitations by promoting defensive states

through a cued conditioned stimulus, advancing toward temporal-specific manipulations such as

optogenetics, and allowing us to study the phenomena with a within-subject experimental design

by subsequent CS[+/-] presentations. Nevertheless, since the circuits involved in contextual and cued

fear may differ[31,32], this might also imply different circuits involved in the emotional modulation

of pain-suppression behavior. To our knowledge, this is the first demonstration of a cued fear-

conditioned analgesia model in rodents. Addressing possible differences in the circuitry involved

in contextual and cued fear-induced analgesia will require further investigation.

Our observation that the photoactivation of SST[+] vlPAG cells not only abolished the analgesia

during defensive states but also decreased freezing expression has far-reaching functional

consequences. Two independent previous studies found that long-range inhibitory inputs from the

central medial amygdala contact inhibitory cells within the vlPAG, implicated in different roles:

the modulation of fear behavior[21] and nociceptive transmission[33]. Notably, both studies describe

the involvement of vlPAG inhibitory cells in a disinhibitory circuit mechanism of long-range

projecting excitatory cells as the underlying neural substrates responsible for such modulations[21,33].

We observed that SST[+] is the most abundant type of inhibitory cells in the vlPAG (**Supplementary**

Figure 3). Thus, it is conceivable that the SST[+] vlPAG cells are indeed part of a circuit involved

in the emotional regulation of endogenous pain-suppression. Notably, global manipulation of SST[+]

vlPAG cells cannot disentangle their role in analgesia and fear behavior at the local level (**Figure**

2 and 3). Because the studies mentioned above exclusively addressed the contribution of vlPAG

circuits in either pain behavior[33] or fear behavior[21], these studies could have been oblivious to

alternative behavioral readout interpretations. Here we demonstrated that SST[+] vlPAG cells

projecting to the RVM are involved in the modulation of analgesia induced during defensive states

without impacting the behavioral expression of fear (**Figure 6**). Thus, our results suggest that fear

and analgesia signals rely on overlapping neuronal circuits and that the discrimination between

these two states could be mediated within the vlPAG network. Altogether, our data is consistent

with the existence of different populations of SST[+] cells within the vlPAG responsible for the

mediation of fear behavior and analgesia information, defined by their output connectivity.

The lateral inhibition model of analgesia within the PAG relies on the activation of PAG

excitatory cells projecting to the RVM. More precisely, in this model, analgesia is thought to occur

through an opioid-dependent inhibition of GABAergic cells, ultimately disinhibiting excitatory

cells projecting to the RVM[12,19,20]. Our data indicate that in addition to this disinhibitory

mechanism, analgesia is also mediated by the inhibition of a direct long-range SST+ neurons

projection onto RVM cells projecting to the DH (**Figure 6**). Our results align with the parallel

inhibition-excitation model, where inhibitory and excitatory cells form two distinct, parallel

descending pathways for pain modulation[35]. Indeed, previous research demonstrated the presence

of an inhibitory pathway projecting throughout the PAG–RVM-spinal cord dorsal horn neuraxis[36].

Our results complement this study by suggesting that this previously proposed inhibitory pathway

is mediated by GABAergic SST+ vlPAG cells and has a functional role in mediating analgesia.

We identified a novel midbrain to brainstem circuit mechanism composed of SST+ vlPAG cells

that project to the RVM, which regulates analgesia during defensive states. Substantial progress

has been made that outlined the midbrain circuitry in endogenous pain-suppression pathway[12,23,37].

However, only more recently, high cortical areas have also been shown to be involved in

analgesia[7,38]. Previous reports describe connections between CeL and vlPAG inhibitory neurons in

the modulation of pain behavior[21,34]. Future work is required to elucidate as to whether the exact

circuit that regulates analgesia are mediated by different negative emotional states such as fear,

depression or stress.

Recent human reports reveal strong comorbidity between mood and anxiety disorders with and

pain-related disorders[39], suggesting that the deregulation of the circuits mediating the interplay

between emotional and pain behavior could be at the core of such psychiatric conditions.

Comprehensive knowledge about the circuits mediating the emotional regulation of pain behavior

is urgently needed. We identified an SST+ cell type- and pathway- specific involvement in the

mechanisms that mediate the emotion-dependent regulation of pain behavior. Thus, these findings

may empower the development of more effective analgesic therapies by the improvement of further

refined pharmacological targets.

Figure Legends

Figure 1. Fear modulation of pain behavior. a. Schematic of the setup and the CFCA paradigm.

The paradigm consists of a discriminative auditory fear conditioning test followed by a pain

sensitivity assay (for details, see Methods). **b.** Average freezing values for CS^+ was higher than

CS^- or baseline (BL) periods during retrieval (****, $P < 0.0001$, one-way RM ANOVA, n = 12

mice). **c.** Example of the infrared digital thermographic camera view of a mouse during the FC HP

test. After 1 min context acclimatization, plate temperature steadily increases at 6 ºC per min (HP

start). CSs started 130 s after the HP start and continued until mice displayed an NC response

(licking the hindpaw or jumping). **d.** Example of assessed hindpaw licking NC response. **e.** Latency

of NC response on the FC HP test increased during CS^+ compared to CS^- (***, $P = 0.0005$,

Wilcoxon matched-pairs signed rank test, n = 12 mice). **f. Left**: dynamics of freezing probability

at CS onset and NC response for both CS^- and CS^+ (freezing periods for each individual mouse are

displayed by rows in grey; error bars display S.E.M; bin size = 0.5 s). **Right**: average freezing

values during 10 s after CS onset and before NC response for CS^+ were higher than CS^- only at CS

onset (***, $P_{onset} = 0.0015$, paired t-test, n = 12 mice). **g.** Mean freezing values during fear

extinction, 24 CS^+ were presented across two separate extinction sessions (joined by dashed lines).

Mice acquired the CS^+-US association (1st CS^+ block vs. BL/CS^- ****, $P < 0.0001$, Friedman test,

n = 10 mice), followed by a rapid extinction (5th & 6th block of CS^+ vs BL/CS- ns, $P > 0.05$, one-

way RM ANOVA). **h.** After extinction, there was no difference in the NC response latency for

both CSs (ns, P > 0.05, paired t-test n = 10 mice). **i.** Average NC response latencies for the different

HP tests. *No tone* - mice submitted to the HP test without conditioning nor tone presentation. *Tone*

- mice submitted to the HP test paired with unconditioned tone presentation. *CS$^-$/CS$^+$* - mice were

submitted to the CFCA protocol. (No tone/Tone/CS$^-$ vs. CS$^+$ **, P < 0.0001, Krustal-Wallis test).

For exact P values and test statistics on this and all subsequent figures, see Supplementary Table

1.

Figure 2. SST$^+$ cells in vlPAG mediate the fear modulation of pain behavior. a. Sst-IRES-Cre

mice were bilaterally injected with viruses that expressed ChR2 (right, upper), ArchT (right,

bottom), or GFP (control) in SST$^+$ vlPAG cells. **b**. Schematics of the stimulation protocol used for

photoactivation of the SST$^+$ vlPAG cells: blue light delivered at 2 Hz with 5 ms pulse duration

during CSs presentation until NC response. For photoinhibition experiments, green light was

delivered continuously. **c, e.** Average freezing values during retrieval for ChR2- (**c**) and ArchT-

infected mice (**e**) and their respective GFP controls. The opsin and respective control groups were

pooled together because no difference was found in the conditioning level (see Supplementary

Figure 4). Average freezing values during CS$^+$ was higher than CS$^-$ or baseline (BL) periods (****,

P < 0.001, one-way RM ANOVA, (**c**) n = 16 mice and (**e**) n = 23 mice). **d.** Photoactivation of SST$^+$

vlPAG cells abolished the analgesic effect induced by fear (*, P = 0.0112, opsin x CS - two-way

RM ANOVA, n = 7 GFP, n = 9 ChR2). The NC response latency for the CS$^+$ was significantly

different between the ChR2 and GFP groups (*, P = 0.0344, Bonferroni post-hoc test). For the

ChR2 group, the NC response latency during CS$^+$ was equivalent to the CS$^-$ (ns, P = 0.3876,

Bonferroni post-hoc test). On the contrary, the NC response latency between the CSs differed for

the GFP group (***, P = 0.0003, Bonferroni post-hoc test). **f.** Photoinhibition of SST$^+$ vlPAG cells

increased the analgesic effect for the ArchT group when compared to the GFP (*, P = 0.0037, opsin

x CS - two-way RM ANOVA, n = 12 GFP, n = 11 ArchT). The NC response latency for the CS^-

and CS^+ was significantly different between the ArchT and GFP group (CS^-: ****, P < 0.0001,

Bonferroni post-hoc test; CS^+: *, P = 0.0265, Bonferroni post-hoc test). For the GFP group, the

latency of NC response was higher for the CS^+ trials when compared to the CS^- trials (***, P =

0.0003, Bonferroni post-hoc test), yet this was not the case for the ArchT group (ns, P > 0.999,

Bonferroni post-hoc test).

Figure 3. Activation of SST^+ vlPAG cells reduced fear expression a. Protocol for optogenetic

manipulation during fear retrieval. Day 1 and 2 were done as described previously for the CFCA

paradigm. During retrieval, there were 12 CS^+ presentations divided into three blocks. **b.** The

stimulation protocol used for photoactivation of the SST^+ vlPAG cells. The optogenetic

manipulation was done during the 2nd block of the CS^+ presentation (analogously to CFCA

manipulations, **Figure 2b**). **c.** Light inhibition of SST^+ vlPAG cells did not modulate freezing levels

(ArchT ns, P = 0.9396, opsin x CS two-way RM ANOVA, n = 6, GFP, n = 8). **d.** Photoactivation

of the SST^+ vlPAG cells had no effect on the GFP group, but it transiently decreased the freezing

levels for the ChR2 group (***, P < 0.0001, opsin x CS two-way RM ANOVA, n = 9 GFP, n = 8

ChR2).

Figure 4. Manipulation of SST^+ vlPAG cells alters spinal cord-related pain signals. a.

Schematics of *in vivo* anesthetized experiments. Extracellular local field potential and single-unit

recordings in the spinal cord DH were performed during optogenetic stimulation of SST^+ vlPAG

cells with concomitant noxious electrical paw stimulation. **b.** Example of average nociceptive field

potentials in the lumbar SC before (OFF), during (ON), and after (OFF) photoactivation of SST^+

vlPAG cells (top). Photoactivation induces a significant increase in the nociceptive fields (bottom;

, P = 0.0001, one-way RM ANOVA, n = 8 mice). **c. Analogous to panel b, but for photoinhibition

of SST$^+$ vlPAG cells. Photoinhibition induces a significant decrease in the nociceptive fields

(bottom; *, P = 0.0275, one-way RM ANOVA, n = 5 mice). **d.** Example of WDR single-unit

activity before and during photoactivation of SST$^+$ vlPAG cells (**top**). Photoactivation of SST$^+$

vlPAG cells induces a significant and global increase in WDR response to both Aδ- and C-

mediated nociceptive fibers (**bottom**; **, $P_{Aδ\ fiber}$ = 0.0022, paired t-test, n = 11 cells; **, $P_{C\ fiber}$ <

0.0033, paired t-test, n = 11 cells). **e.** Analogous to panel d, but for photoinhibition of SST$^+$ vlPAG

cells. Photoinhibition induces a significant and specific inhibition of WDR response to Aδ- and C-

fibers (**bottom**; **,** $P_{Aδ\ fiber}$ = 0.0241, ****, $P_{C\ fiber}$ < 0.0001, n = 14 cells**,** Wilcoxon matched-pairs

signed rank test). **f.** Example traces of single-unit recordings of WDR cells with subthreshold

electrical stimulation accompanied by photoactivation of SST$^+$ vlPAG cells (**top**). The

photoactivation of SST$^+$ vlPAG cells elicits WDR response to A- and C- mediated peripheral fibers

(**bottom**; *, $P_{Aδ\ fiber}$ = 0.0156, n = 10 cells; **, $P_{C\ fiber}$ = 0.0020, n = 10 cells, Wilcoxon matched-

pairs signed rank test).

Figure 5. SST$^+$ vlPAG-RVM-DH pathway activation removes analgesia. a. Schematics of

possible SST$^+$ vlPAG circuits mediating analgesia by long-range projections directly to the SC or,

alternatively, by projecting to the SC via the rostral ventromedial medulla (RVM). **b.** Sst-IRES-

Cre mice were injected with an anterograde AAV Cre-dependent GFP virus in the vlPAG (left).

Example of SST fibers labelling at the level of the RVM (middle)**.** Higher magnification (right)

reveals putative axonic buttons in the RVM (examples indicated by white arrows). **c.** Sst-IRES-

Cre mice were injected concomitantly with anterograde AAV Cre-dependent GFP virus in the

vlPAG and retrograde fluorogold in the lumbar DH of the SC. **d.** Fluorogold positive cells (red)

cross SST$^+$ vlPAG fibers in the RVM (green). Higher magnification (**right**) shows close contacts

between the putative SST$^+$ button and fluorogold$^+$ cells or fibers (white arrows). **e.** Sst-IRES-Cre

mice were injected in the vlPAG with AAV Cre-dependent ChR2 virus and optic fiber placed above

the lumbar SC. Single-unit recordings of WDR cells while photoactivation of the SST$^+$ vlPAG SC

fibers and electrically stimulating the paw in anesthetized mice (**left**). Photoactivation of vlPAG

projections to SC did not affect nociceptive transmission (**right**; ns, P = 0.3683, paired t-test, n =

11 cells). **f.** Same experimental design as in panel e, except that the optic fiber was placed above

the RVM (left). Photoactivation of SST$^+$ vlPAG inputs to the RVM induces a significant increase

in WDR response to both Aδ- and C-mediated nociceptive fibers stimulation (**middle**; *, $P_{Aδ\ fiber}$ =

0.0346, Paired t-test, n = 10 cells; **, $P_{C\ fiber}$ = 0.0020, Wilcoxon matched-pairs signed rank test, n

= 10 cells). Removal of analgesia by activation of SST$^+$ vlPAG-RVM induces a significant increase

in WDR response to C-fibers responses (right; **, P = 0.0078, Wilcoxon matched-pairs signed

rank test, n = 9 cells).

Figure 6. SST$^+$ vlPAG-RVM-SC pathway activation abolishes analgesia induced by CFCA.

a. Sst-IRES-Cre mice were bilaterally injected with an AAV expressing ChR2 or GFP in the

vlPAG, and optic fibers were implanted above the RVM (left). Representative example of SST$^+$

vlPAG terminals in the RVM (right). **b.** Photoactivation of SST$^+$ vlPAG-RVM pathway did not

modulate freezing levels (ns, P = 0.6443, opsin x CS two-way RM ANOVA; CS$^+$ 2nd block, GFP

vs. ChR2 ns, P > 0.9999, Bonferroni post-hoc test, n = 7 GFP, n = 6 ChR2). **c.** Average freezing

values during retrieval for pooled ChR2 and GFP-infected mice (groups were pooled together since

no difference was found in the conditioning level). Average freezing values during CS$^+$ were higher

than CS$^-$ or baseline (BL) periods (****, P < 0.0001, one-way RM ANOVA, n = 14 mice). **d.**

Photoactivation of SST$^+$ vlPAG-RVM pathway abolished the analgesic effect modulated by fear

(**, P = 0.0063, opsin x CS - two-way RM ANOVA, n = 7 GFP, n = 7 ChR2). For the ChR2 group,

the NC response latency during CS^+ was equivalent to the CS^- (ns, $P > 0.9999$, Bonferroni post-

hoc test). On the contrary, the NC response latency between the CSs differed for the GFP group

(****, $P < 0.0001$, Bonferroni post-hoc test). **e**. Working model schematics for SST^+ control

analgesia induced during defensive states. Activation of SST^+ cells in the vlPAG is associated with

the inhibition of fear-induced freezing behavior. Selective activation of RVM projecting SST^+

vlPAG cells is associated with an inhibition of analgesia induced during defensive states.

Methods

Subject details

We used either male C57BL6/J mice (Janvier), heterozygous or homozygous Sst-IRES-Cre

mice (Jackson laboratory), or heterozygous VIP-IRES-Cre mice (Jackson Laboratory) age 8-14

weeks that were individually housed under a 12 h light-dark cycle and provided with food and

water ad libitum. All procedures were performed in accordance with standard ethical guidelines

(European Communities Directive 86/60-EEC) and were approved by the committee on Animal

Health and Care of Institut National de la Santé et de la Recherche Médicale and the French

Ministry of Agriculture and Forestry (agreement #A3312001).

Behavioral apparatus

Cued fear-conditioned analgesia *task* was performed in three different contexts (**Figure 1a**).

Context A was used for *Habituation* and *Retrieval* and consisted of a plexiglass cylinder (25 x 24

cm diameter) with a grey, smooth plastic floor and house lights (16 lux). Context B was used for

Fear conditioning and consisted of a square plexiglass (25 x 40 cm) with a grid floor connected to

a shocker (Coulbourn Instruments) and brighter house lights (40 lux). A total of 5 scrambled

footshocks of 1 s duration and intensity of 0.8 mA were delivered via the grid floor and served as

the unconditioned stimulus (US). Context A and B were cleaned, respectively, with 70% ethanol

or 1 % acetic acid between different mice. Both contexts contained an infrared beam detection that

automatically scored freezing periods, determined by no change in infrared beam crossing for at

least 2 s.

Context C was used for the *FC Hot Plate (HP) test*. A steady increase in temperature was

controlled by the Incremental Hot/Cold Plate Analgesia Meter (IITC) device. To restrain the mice's

movement, the HP apparatus had a testing surface (plate) enclosed in a square plexiglass surface

(20.3 x 10 x 20.5 cm) and house lights (100 lux). The mice's temperature and surroundings were

recorded with an infrared digital thermographic camera (Testo 885) placed ~ 50 cm above the

testing surface. The thermal camera had a spatial resolution of 320×240 pixels, a sampling rate of

25 Hz, and thermal sensitivity of 0.03 °C at 30 °C. The testing surface of context C was cleaned

with water between different mice. All three contexts were enclosed in an acoustic foam isolated

box with speakers mounted on the top of each compartment. The auditory conditioned stimulus

(CS) consists of either 7.5 kHz or white-noise 50 ms pips at 1 Hz repeated 27 times, 2 ms rise and

fall, and 80 dB sound pressure level.

Open field task was performed in a square plexiglass arena (36 x 36 x 25cm). A LED mounted

on the top-right side of the arena signalled the start and end of different epochs for offline analyses.

A video camera recorded from above the arena at 30 fps for offline video-tracking purposes.

Real-Time Place Preference (RTPP) task was performed in a shuttlebox consisting of a

plexiglass box (40 x 10 x 30 cm) with a floor grid, where a small plastic hurdle (1 cm height)

divided the arena into two equal compartments while infrared beams detection automatically

monitored the mice shuttling between compartments (Imetronic). A video camera recorded from

above the arena at 30 fps for offline video-tracking purposes.

For both the open field and RTPP task, a free user video-tracking software (idTracker: Tracking

individuals in a group by automatic identification of unmarked animals) together with in-house

codes in Matlab (The MathWorks, Inc., Natick, MA, USA) were used to analyze each condition.

Behavioral paradigm

Cued fear-conditioned analgesia (CFCA). C57BL6/J mice (n = 12) were habituated to the

context and tones (day 1). Four white-noise (CS⁻) and four 7.5 kHz (CS⁺) tones were presented

sequentially and without US reinforcement. In Fear conditioning (day 2), five CS⁻ and five CS⁺

were presented in an intermingled fashion. The CS⁺ presentations were paired with a mild

footshock (US) at tone offset, whereas the CS⁻ was never reinforced. The Fear retrieval session

(Day 3) was done 24h after conditioning and in the same context as the habituation session. As in

habituation, four CS⁻ and four CS⁺ were presented sequentially and without US reinforcement

(**Figure 1a**).

Since the focus of this study was the emotional modulation of pain sensitivity, it was compulsory

to evaluate the associative fear levels before measuring its impact on pain sensitivity. Two indices

were computed: the discrimination index (DI), to assess the discrimination between CS⁻ and CS⁺,

and the conditioning index (CI), which indicated the level of freezing to the tone predicting the US.

These indexes were calculated as follows: $DI = \frac{(Freezing\ to\ CS+)-(Freezing\ to\ CS-)}{(Freezing\ to\ CS+)+(Freezing\ to\ CS-)}$ and $CI = DI \times$

(Freezing to CS⁺). Based on preliminary data, mice were only submitted to the FC HP test if DI ≥

0.4 & CI ≥ 0.3. Mice that did not fit the criteria were conditioned a second time. There was a

minimum interval of 2 h between the retrieval and the FC HP test.

The FC HP test consisted of two trials, one where the CS⁺ was presented and another where the

CS⁻ was presented. The two trials were counterbalanced within the group. The testing surface was

set at 30 ºC. After a 60 s acclimatization period, its temperature gradually increased at 6 ºC per

minute (HP start). Tone presentation (CS$^+$ or CS$^-$) started 130 s after the HP start. Temperature

increase and tone presentation terminated concomitantly with the display of a nociception response.

Valid nociception readout responses included jumping or licking the hindpaw. The effect of the

negative emotional modulation on pain sensitivity was assessed by comparing the latency (or

temperature) of the NC response between the two HP trials. There was a minimum of 30 min

interval between the two trials for each mouse, during which mice returned to their home cage.

Freezing during the HP test was assessed by automatically analyzing thermographic camera

videos using custom scripts (available at: github.com/djercog/TestoFreez). Briefly, individual

video frames were binarized to define a mask for the mouse within each frame. Consecutive

binarized frames were then subtracted, and the number of non-zero pixels on the resultant image

determined if the mouse was considered immobile when below a certain threshold (defined to

approximate manual freezing estimation on random videos). Immobilization episodes of at least

0.5 s were considered as freezing periods. In contrast, non-immobilization episodes shorter than

0.1 s were merged together with the neighbouring detected freezing episodes.

Extinction training. On days 3 and 4, mice (n =10) were submitted to an extinction training

protocol established by Courtin et al. 2014[40]. Briefly, there were 4 CS$^-$ and 12 CS$^+$, presented in a

non-reinforced manner (**Figure 1g**). Mice were considered as having successfully extinguished the

fear expression if the level of freezing in the last 4 CS$^+$ was not statistically different from the CS$^-$

. After fear extinction, mice were submitted to the FC HP test, using the same parameters as in the

CFCA paradigm.

Stability training. The first three days consisted of the classical CFCA paradigm. On day 4,

mice (n = 10) repeated the protocol applied on day 3 (**Supplementary Figure 2e-i)**. All mice that

passed the fear conditioning criteria (DI $\geq$ 0.4 & CI $\geq$ 0.3) on day 3 were kept for the following

day, independent of their level of Conditioning on day 4.

Vasoconstriction assay. To determine if vasoconstriction was a confounding factor in the FC

HP test outcomes, we used a modified version of the CFCA paradigm (**Figure 1f)**. During the HP

test (n = 13), the temperature was not gradually increased but maintained at 30 ºC while CSs were

presented. The tones (CS^+ or CS^-) were presented for 120 s, and mice were kept for an additional

50 s before the HP trial terminated. Offline, with the infrared videos, mice's back and tail were

measured at 30 s intervals. For each measuring point, the temperature of three spots of the two

body part was averaged to create the body temperature for the back and tail for a given time point

and mouse.

Basal nociception assay (No tone). Naïve mice (n = 10) were submitted to the HP test (**Figure**

1i and Supplementary Figure 2a-d). Each mouse underwent two identical trials. The HP test was

similar to the one described in the CFCA paradigm, except no tones were presented. There was a

minimum of 30 min interval between the two trials, during which mice returned to a restful state

in their home cage. Since the two trials were not significantly different, for simplification, in panel

i from Figure 1, the two trials were pooled together.

Tone-specific nociception assay. Naïve mice (n = 12) were submitted to the HP test without

auditory fear conditioning (**Figure 1i and Supplementary Figure 2a-d)**. Each animal underwent

two trials, one trial with a 7.5 kHz tone presentation and another with a WN tone presentation. The

tone presentation was counterbalanced. The HP test parameters were similar to the one described

for the CFCA, except that both tones were unconditioned. Since the two trials were not significantly

different, for simplification, in panel i from Figure 1, the two trials were pooled together.

Open Field assay. This test was used to determine the effect of optogenetic stimulation on

locomotion. Therefore, only mice from optogenetic experiments were submitted to this assay. Mice

could freely move during the entire test. The locomotion assay had a total duration of 9 min and

was divided into 3 min epochs. The first and third epochs were OFF periods in which no optical

stimulation occurred. During the second epoch, mice received photostimulation (ON period). The

optogenetic stimulation effect was analyzed by comparing the overall distance travelled between

the OFF and ON epochs (**Supplementary Figure 5c, d)**. Mice injected with GFP were used to test

the effect of heat and light on the stimulation itself. Mouse tracking data was extracted from video

recordings using idTracker[41] and imported to Matlab for further analysis using custom scripts to

calculate the total distance travelled during stimulation.

Real-time Place Preference assay. Only mice from optogenetic experiments were submitted to

this assay. During the entire duration of the assay, mice could freely shuttle between the two

compartments. Under a closed-loop stimulation, mice received photo-stimulation upon entry into

one of the two compartments. The stimulated compartment was counterbalanced within the group.

The optical stimulation effect was assessed by comparing the time spent in the stimulated

compartment between the ChR2 and GFP groups (**Supplementary Figure 5e, f)**. Permanence time

in the stimulated compartment was automatically assessed by apparatus software (Imetronic).

Virus injections and optogenetics

For optogenetic manipulation of SST$^+$ cells in the vlPAG, 0.15-0.2 μL of either ChR2 (AAV5-

EF1a-DIO-hChR2(H134R)-EYFP, titer: 3.2×10^{12} - Vector Core, University of North Carolina),

ArchT (AAV9-CAG-FLEX-ArchT-GFP, titer: 4.7×10^{12} - Vector Core, University of North

Carolina) or GFP (AAV5-FLEX-GFP, titer: 4.5×10^{12} - Vector Core, University of North Carolina)

were bilaterally injected into the vlPAG of 8/9 weeks old Sst-IRES-Cre mice from glass pipettes

(tip diameter 20-30 μm) at the following coordinates relative to bregma: - 4.4 mm AP; ± 1.5 mm

ML; -2.45 mm DV from dura, with a 20 degrees angle. Injection coordinates for manipulating VIP$^+$

vlPAG cells were the following: - 4.4 mm AP; ± 1.35 mm ML; -2.5 mm DV from dura, with a 20

degrees angle.

At two weeks after the injections, mice were bilaterally implanted with custom-built optic fibers (diameter: 200 μm; numerical aperture: 0.39; Thorlabs) above the vlPAG at the following coordinates relative to bregma: i) Sst-IRES-Cre mice: - 4.4 mm AP; ± 1.0 mm ML; -1.8 mm DV from dura, with a 10 degrees angle; ii) VIP-IRES-Cre mice: - 4.4 mm AP; ± 0.8 mm ML; -2.0 mm DV from dura, with a 10 degrees angle. For RVM manipulations, mice were implanted at the following coordinates relative to bregma: -5.8 mm AP; 0.0 mm ML; -5.2 DV from the dura. All implants were secured using three stainless steel screws and Super-Bond cement (Sun Medical). During surgery, long- and short-lasting analgesic agents were injected (Metacam, Boehringer; Lurocaïne, Vetoquinol). After surgery, mice were allowed to recover for at least five days. Afterward, mice were handled daily to familiarize themselves with being restrained for the optic fibers connection. Behavioral experiments were performed at least four weeks after viral injections. Only mice with correct placement of optic fibers and virus expression were included in the analyses.

For optogenetic excitation, light stimulation consisted of blue light (473 nm, ~8-10 mW at fiber tip) delivered with 2 Hz frequency and 5 ms pulse duration. In contrast, light stimulation consisted of green light (532 nm, ~8-10 mW at fiber tip) delivered continuously for optogenetic inhibition. For optogenetic manipulations during the CFCA paradigm of either SST^+ or VIP^+ vlPAG cells, the light was delivered during the FC HP test and paired with the tone presentation.

Two sets of experiments were performed sequentially for stimulation during the fear conditioning. First, the footshock US was replaced by optical stimulation. Then, the US became the optical stimulation combined with the footshock (**Supplementary Figure 6**). For both conditions, the optogenetic stimulation started 5 s before and lasted until 5 s after CS^+ offset. The same mice were used for both experiments. For the manipulations during the fear retrieval, there

545   were 12 CS$^+$ presentations divided into blocks of 4 CS$^+$. The optogenetic stimulation was paired

546   with the second block of CS$^+$.

Anatomical tracing

Fluorogold injection. An incision between one to two cm was made slightly caudal to the peak

of the dorsal hump to expose the lumbar spinal region. The vertebra of interest was identified.

Then, a small incision was made between the tendons and the vertebral column on either side. The

vertebra was then secured using spinal adaptor clamps, and all tissue was removed from the surface

of the bone. Pulled borosilicate glass capillaries (Ringcaps, disposable capillary pipettes with ring

mark, DURAN, Hirschmann Laborgeräte, Germany) were inserted between 2 vertebrae and

allowed to microinject 0.5 μL of fluorogold 2% in the dorsal horn of the spinal cord on both sides.

555   *In vivo* electrophysiology

Mice were anesthetized with isoflurane 4% for induction, then 1.5% maintenance. The

experiment was started as soon as there was no longer any reflex. The colorectal temperature was

kept at 37 °C with a heating blanket. Two metal clamps were used to set the animal spine in a

stereotactic frame (M2E, France) for stability during electrophysiological recordings. Then, a

laminectomy was performed at T13-L1 to expose the lumbar part of the spinal cord. The dura mater

was carefully removed. Custom-made optical fibers were placed 1mm above the dorsal spinal cord

for optogenetic manipulations. C-fiber-evoked field potentials were recorded in the deep lamina of

the DH (at a depth range of 250 and 500 μm) with borosilicate glass capillaries (2 MΩ, filled with

NaCl 684 mM; Harvard Apparatus, Cambridge, MA, USA). Field potentials were recorded with

an ISODAM-amplifier (low filter: 0.1Hz to high filter: 0.1 kHz; World Precision Instruments,

USA) in response to electrical stimulation of the ipsilateral paw. Single unit recordings of WDR

DH cells were made with the same borosilicate glass capillaries mentioned above and placed in the

dorsal part of the spinal cord. The criterion for selecting a cell was the presence of an A-fiber-evoked response (0-80 ms) followed by a C-fiber-evoked response (80-150 ms) to electrical stimulation of the ipsilateral sciatic nerve.

The paw of the mice was stimulated by trains (every 30 s) of electrical stimulation at two times the threshold for C-fibers were performed before, during, and after optogenetic stimulations with an optic fiber placed above the recording site. Subthreshold stimulations were performed below the threshold for C-fiber and A-fiber, respectively.

Histology analyses

Mice were administered a lethal dose of Exagon and underwent transcardial perfusions via the left ventricle with 4% w/v paraformaldehyde (PFA) in 0.1 M PB. Following dissection, brains were post-fixed for 24 h at 4°C in 4% PFA. Brain sections of 80 μm-thick were cut on a vibratome, mounted on gelatin-coated microscope slides, and dried. To verify correct viral injections and optic fiber location, serial 80 μm-thick slices containing the regions of interest were mounted in VectaShield (Vector Laboratories) and imaged using an epifluorescence system (Leica DM 5000) fitted with a 10-x dry objective. The location and the extent of the injections/infections were visually controlled. Only infections targeting the vlPAG and optic fibers terminating, depending on the experiment, above the vlPAG or RVM were included in the analyses.

Statistics

All statistics details are contained in Supplementary Table 1. Box-whisker plots indicate median, interquartile range, and 5th - 95th percentiles of the distribution. All statistics are indicated where used. Statistical analyses were performed with Prism software (GraphPad). We tested the normality of all data with the Kolmogorov–Smirnov normality test in Prism software (GraphPad), and used nonparametric statistical tests for nonnormally distributed datasets, as indicated where

used. No statistical methods were used to predetermine sample sizes, but sample sizes were based

on previous studies in our laboratory. Blinding for the opsin was done for optogenetics

experiments. Animals used for the CFCA paradigm that did not satisfy the DI and CI criteria after

two conditioning sessions were discarded from the study. Mice with an incorrect injection of the

opsins or misplaced location of the optic fibers were discarded. For *in vivo* electrophysiology, field

potentials were measured as the area above the curve in the C-fiber range (80-300 ms). Absolute

values were used to compare values in each condition (OFF vs. ON optogenetic manipulation). In

single-unit recordings, the number of A- and C-fiber induced spikes of WDR cells were measured

after each electrical stimulation, during, and after optogenetic manipulation. An average of the four

stimuli for each WDR recorded was used for statistical analysis. Outliers were removed using the

Tukey method.

No other mice were excluded. Significance levels are indicated as follows: *$p<0.05$, **$p<0.01$,

***$p<0.001$, **** $p<0.0001$.

Data and script availability

605   Data supporting the findings of this study and data analysis codes are available from the

606   corresponding author on request.

References

1.   Bolles, R. C. & Fanselow, M. S. A perceptual-defensive-recuperative model of fear and
pain. 291–323 (1980).

2.   Rhudy, J. L. & Meagher, M. W. Fear and anxiety: divergent effects on human pain
thresholds. *Pain* **84,** 65–75 (2000).

3.   Beckham, J. C. *et al.* Chronic posttraumatic stress disorder and chronic pain in vietnam
combat veterans. *J. Psychosom. Res.* **43,** 379–389 (1997).

4.   Flor, H. & Grüsser, S. M. Conditioned stress-induced analgesia in humans. *Eur. J. Pain* **3,**

616   317–324 (1999).

5. International Association for the Study of Pain. IASP Terminology. *Iasp* (2017).

6. Butler, R. K. & Finn, D. P. Stress-induced analgesia. *Prog. Neurobiol.* **88,** 184–202 (2009).

7. Rea, K., Roche, M. & Finn, D. P. Modulation of conditioned fear, fear-conditioned analgesia, and brain regional C-fos expression following administration of muscimol into the rat basolateral amygdala. *J. Pain* **12,** 712–721 (2011).

8. Floyd, N. S., Price, J. L., Ferry, A. T., Keay, K. a & Bandler, R. Orbitomedial prefrontal cortical projections to distint longitudinal columns of the periaqueductal gray in the rat. *J. Comp. Neurol.* **422,** 556–578 (2000).

9. LeDoux, J. E., Iwata, J., Cicchetti, P. & Reis, D. J. Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. *J. Neurosci.* **8,** 2517–29 (1988).

10. Vertes, R. P. Differential Projections of the Infralimbic and Prelimbic Cortex in the Rat. *Synapse* **51,** 32–58 (2004).

11. Carrive, P. & Morgan, M. M. *Periaqueductal Gray - Chapter 10. The Human Nervous System* (Elsevier, 2012). doi:10.1016/B978-0-12-374236-0.10010-0

12. Basbaum, A. I. & Fields, H. L. Endogenous Pain Control Systems: Brainstem Spinal Pathways and Endorphin Circuitry. *Annu. Rev. Neurosci.* **7,** 309–338 (1984).

13. Millan, M. J. Descending control of pain. *Prog. Neurobiol.* **66,** 355–474 (2002).

14. Fields, H. L., Heinricher, M. M. & Mason, P. Neurotransmitters in nociceptive modulatory circuits. *Annu. Rev. Neurosci.* **14,** 219–245 (1991).

15. Castilho, V. M., Macedo, C. E. & Brandão, M. L. Role of benzodiazepine and serotonergic mechanisms in conditioned freezing and antinociception using electrical stimulation of the dorsal periaqueductal gray as unconditioned stimulus in rats. *Psychopharmacology (Berl).* **165,** 77–85 (2002).

16. Helmstetter, F. J. & Landeira-Fernandez, J. Conditional hypoalgesia is attenuated by Naltrexone applied to the periaqueductal gray. *Brain Res.* **537,** 88–92 (1990).

17. Bellgowan, P. S. F. & Helmstetter, F. J. The role of mu and kappa opioid receptors within the periaqueductal gray in the expression of conditional hypoalgesia. 83–89 (1998).

18. Jacquet, Y. F. The NMDA receptor: central role in pain inhibition in rat periaqueductal gray. *Eur. J. Pharmacol.* **154,** 271–276 (1988).

19. Moreau, J. L. & Fields, H. L. Evidence for GABA involvement in midbrain control of medullary neurons that modulate nociceptive transmission. *Brain Res.* **397,** 37–46 (1986).

20. Osborne, P. B., Vaughan, C. W., Wilson, H. I. & Christie, M. J. Opioid inhibition of rat periaqueductal grey neurones with identified projections to rostral ventromedial medulla in

vitro. *J. Physiol.* **490,** 383–389 (1996).

21.   Tovote, P., Esposito, M. S., Botta, P., Chaudun, F. & Jonathan, P. Midbrain circuits for
defensive behaviour. *Nature, Press* (2016). doi:10.1038/nature17996

22.   Fields, H. L. State-dependent opioid control of pain. *Nat. Rev. Neurosci.* **5,** 565–75 (2004).

23.   Lau, B. K. & Vaughan, C. W. Descending modulation of pain: the GABA disinhibition
hypothesis of analgesia. *Curr. Opin. Neurobiol.* **29,** 159–164 (2014).

24.   Vianna, D. M. L. & Carrive, P. Changes in cutaneous and body temperature during and after
conditioned fear to context in the rat. *Eur. J. Neurosci.* **21,** 2505–2512 (2005).

25.   Samineni, V. K. *et al.* Divergent Modulation of Nociception by Glutamatergic and
GABAergic Neuronal Subpopulations in the Periaqueductal Gray. *eNeuro* **4,** (2017).

26.   Harris, J. A. Descending antinociceptive mechanisms in the brainstem: their role in the
animal's defensive system. *J. Physiol. Paris* **90,** 15–25 (1996).

27.   Fields, H. State-dependent opioid control of pain. *Nat. Rev. Neurosci.* **5,** 565–75 (2004).

28.   Smith, G. S. T. *et al.* Distribution of messenger RNAs encoding enkephalin, substance P,
somatostatin, galanin, vasoactive intestinal polypeptide, neuropeptide Y, and calcitonin
gene-related peptide in the midbrain periaqueductal grey in the rat. *J. Comp. Neurol.* **350,**
23–40 (1994).

29.   Viollet, C. *et al.* Somatostatin-IRES-Cre Mice: Between Knockout and Wild-Type? **8,** 1–8
(2017).

31.   Gasser, H. S. The classification of nerve fibers. *Ohio J. Sci.* (1941).

32.   Tovote, P., Fadok, J. P. & Lüthi, A. Neuronal circuits for fear and anxiety. *Nat. Rev.*
Neurosci. **16,** 317–331 (2015).

33.   Maren, S., Phan, K. L. & Liberzon, I. The contextual brain: implications for fear
conditioning, extinction and psychopathology. *Nat. Rev. Neurosci.* **14,** 417–428 (2013).

34.   Yin, W. *et al.* A Central Amygdala–Ventrolateral Periaqueductal Gray Matter Pathway for
Pain in a Mouse Model of Depression-like Behavior. *Anesthesiology* **132,** 1175–1196
(2020).

35.   Cleary, D. R., Neubert, M. J. & Heinricher, M. M. Are opioid-sensitive neurons in the rostral
ventromedial medulla inhibitory interneurons? *Neuroscience* **151,** 564–571 (2008).

36.   Morgan, M. M., Whittier, K. L., Hegarty, D. M. & Aicher, S. A. Periaqueductal gray neurons
project to spinally projecting GABAergic neurons in the rostral ventromedial medulla. *Pain*
140, 376–386 (2008).

37.   Ossipov, M. H., Dussor, G. O. & Porreca, F. Review series Central modulation of pain. **120,**
(2010).

38. Butler, R. K. *et al.* Molecular and electrophysiological changes in the prefrontal cortex-amygdala-dorsal periaqueductal grey pathway during persistent pain state and fear-conditioned analgesia. *Physiol. Behav.* **104,** 1075–1081 (2011).

39. McWilliams, L. A., Cox, B. J. & Enns, M. W. Mood and anxiety disorders associated with chronic pain: an examination in a nationally representative sample. *Pain* **106,** 127–133 (2003).

40. Courtin, J. *et al.* Prefrontal parvalbumin interneurons shape neuronal activity to drive fear expression. *Nature* **505,** 92–6 (2014).

41. Pérez-Escudero, A., Vicente-Page, J., Hinz, R. C., Arganda, S. & de Polavieja, G. G. idTracker: tracking individuals in a group by automatic identification of unmarked animals. *Nat. Methods* **11,** 743–748 (2014).

Acknowledgments

We thank Joshua Johansen, David Finn, and Nadine Gogolla for critical and stimulating discussions; K. Deisseroth and E. Boyden for generously sharing material, S. Laumond, J. Tessaire, and the technical staff of the housing and experimental animal facility of the Neurocentre Magendie. We thank Claire Francioni for her technical support. Microscopy was performed in the Bordeaux Imaging Center of the CNRS-INSERM and Bordeaux University, member of France BioImaging. This work was supported by grants from the French National Research Agency (ANR-FEARLESSPAIN, ANR-10-EQPX-08 OPTOPATH), the Fondation pour la Recherche Médicale (FRM-PhD grant to NW 2020-2021) and the French Society for the Study and Treatment of Pain (SFETD – grant to NW 2022).

Author Contributions

N.W performed behavioral experiments and optogenetic experiments on freely moving animals.

F.A, J.V, performed electrophysiological and optogenetic experiments on anesthetized animals.

E.V, L.C, N.W, F.A, Z.G, T.B, C.R, R.B, and D.G performed histology. N.W, M.L, S.V. P.F, and

C.H. designed the experiments. N.W, F.A, D.J, and E.V analyzed the data, N.W, D.J. and C.H

wrote the paper.

Supplementary Figures

Supplementary Figure 1. Cued fear-conditioned analgesia behavior. a. During habituation, the

freezing levels for the context (BL) and the two tones were low but significantly different (****, P

= 0.0003, Friedman test, n = 12 mice). **b.** Mean temperature at which the NC response was observed

in the HP test. The temperature of NC response was higher during CS^+ trials when compared to the

CS^- trials (***, P = 0.0005, Wilcoxon matched-pairs signed rank test, n = 12 mice). **c.** Schematic

representation of how the back and tail temperature of the mice was measured. Every 30-sec three-

point temperature was taken for each of the two body parts. The temperature of the mice back and

tail were measured by the infrared digital thermographic camera and analyzed offline (see

Methods). The average temperature of the mice back (**d**) and tail (**e**) while the CS^+ or the CS^- were

presented. There were no differences in body temperature for the different CSs trials (ns, P > 0.05,

mixed-effects model, n = 13 mice). Vertical dashed lines correspond to the average time of NC

response for the CS^+ (red) and CS^- (blue) during the standard CFCA protocol.

Supplementary Figure 2. CFCA depends on associative processes. a. Top, protocol for the *No*

Tone: naïve mice were submitted to two HP trials without conditioning nor tone presentation.

Bottom, protocol for the *Tone*: naïve mice were submitted to two HP trials paired with an

unconditioned tone presentation. **b, c.** Mean latency and temperature of NC response for the two

tests mentioned above and the CFCA. The transient cued-fear induced analgesia (CS^+) compared

to the basal nociception, the tone, and the CS^- trials (***, P = 0.0001, Krustal-Wallis test). Trials

between the same type of test (*No tone* or *Tone*) were not significantly different (ns, P > 0.05,

Bonferroni post-hoc test, see statistical table for all comparisons) **d.** After fear extinction, there was

no difference in the mean temperature response between the two CSs (ns, P = 0.3276 paired t-test,

n = 10 mice). **e.** Protocol for stability training (see methods). Mice were submitted to two rounds

of the CFCA paradigm. During retrieval (**f, g**), average freezing values during CS^+ was higher than

CS^- or baseline (BL) periods (****, P < 0.0001, one-way RM ANOVA, n = 10 mice). **h, i.** Mean

latency and temperature of NC response during CS^- and CS^+ trials for FC HP test 1 (**h**) and test 2

(**i**). CS^+ presentation increased the latency and temperature of NC response in FC HP test 1 and test

2 (*, **, and ***, P < 0.05, paired t-test; for details, see supplementary table 1).

Supplementary Figure 3. Two distinct neuronal populations in SST- Cre mice within the

PAG. a. Representative picture of single-molecular fluorescent in situ hybridization for Sst

mRNAs in the PAG. Scale bar, 400 μm. **b, c.** Single-molecular fluorescent in situ hybridization for

Sst (green) and Cre (red) mRNAs in the vlPAG (left). Histograms showing the co-expression of

Sst/Cre as percentage of Sst-expressing cells (green) and as percentage of Cre-expressing cells

(red) in the dlPAG and vlPAG (right). Scale bar, 20 μm. **d.** Single-molecular fluorescent in situ

hybridization for Sst (red) and Slc32a1 (green) within the dlPAG (upper panel) and vlPAG (bottom

panel). Scale bar, 20 μm. **e.** Quantification of colocalization within the dlPAG (upper panel) and

vlPAG (bottom panel) of Sst^+ and $Slc32a1^+$. In the dlPAG approximately 19% of $Slc32a1^+$ cells

are Sst^+ and 17% of Sst^+ cells are $Slc32a1^+$. On the contrary, in the vlPAG approximately 95% of

$Slc32a1^+$ cells are Sst^+ and 61% of Sst^+ cells are $Slc32a1^+$. **f.** Single-molecular fluorescent in situ

hybridization for Sst (red) and Slc17a6 (green) within the dlPAG (upper panel) and vlPAG (bottom

panel). Scale bar, 20 μm. **g.** Quantification of colocalization within the dlPAG (upper panel) and

vlPAG (bottom panel) of Sst^+ and $Slc17a6^+$. In the dlPAG approximately 91% of $Slc17a6^+$ cells

are Sst^+ and 38% of Sst^+ cells are $Slc17a6^+$. On the contrary, in the vlPAG approximately 34% of

$Slc17a6^+$ cells are Sst^+ and 33% of Sst^+ cells are $Slc17a6^+$.White arrows indicate colocalization.

Supplementary Figure 4. Comparable fear levels prior to the FC HP test. After retrieval, fear

conditioning levels between the opsins and their respective GFP group was tested to ensure

equivalent fear levels. The discrimination index (see methods) between the GFP and ChR2 (**a**, ns,

P = 0.6052 unpaired t-test, n = 16 mice) or ArchT (**b,** ns, P = 0.4575, unpaired t-test, n = 23 mice)

were not significantly different. All the mice discriminated equally between CSs. The conditioning

index (see methods) between the GFP and ChR2 (**c,** ns, P = 0.2333, unpaired t-test n = 16 mice) or

ArchT (**d,** ns, P = 0.1320, unpaired t-test, n = 23 mice) were also not significantly different. All

mice displayed a similar high freezing level to the CS^+.

Supplementary Figure 5. Optogenetic manipulation of SST^+ vlPAG cells does not affect

locomotion or produce aversion. a. Photoactivation of SST^+ vlPAG cells abolished the analgesic

effect induced during defensive states (**, P = 0.0035, opsin x CS - two-way RM ANOVA, n = 7

GFP, n = 9 ChR2). The mean temperature of NC response for the CS^+ was significantly different

between the ChR2 and GFP groups (*, P = 0.0322, Bonferroni post-hoc test). For the ChR2 group,

the mean temperature of NC response during CS^+ was equivalent to the CS^- (ns, P = 0.1783,

Bonferroni post-hoc test). On the contrary, the mean temperature of NC response between the CSs

differed for the GFP group (****, P < 0.0001, Bonferroni post-hoc test). **b.** Photoinhibition of SST^+

vlPAG cells increased the analgesic effect for the ArchT group when compared to the GFP (**, P

= 0.0035, opsin x CS - two-way RM ANOVA, n = 12 GFP, n = 11 ArchT). The mean temperature

of NC response for the CS^- was significantly different between the ArchT and GFP groups (****,

P < 0.0001, Bonferroni post-hoc test). For the GFP group, the mean temperature of NC response

was higher for the CS^+ trials when compared to the CS^- trials (***, P = 0.0003, Bonferroni post-

hoc test), yet this was not the case for the ArchT group (ns, P > 0.999, Bonferroni post-hoc test).

c. The photoactivation of SST$^+$ vlPAG cells was performed during the ON epoch (blue shaded

area), and the average distance travelled during the ON epoch was not different from the OFF

epochs when comparing the two opsins (ns, P = 0.5121, opsin x light - two-way RM ANOVA, n =

7 GFP, n = 9 ChR2). **d.** The photoinhibition of SST$^+$ vlPAG cells was performed during the ON

epoch (yellow shaded area), and the average distance travelled during the ON epoch was not

different from the OFF epochs when comparing the two opsins (ns, P =0.4047, opsin x light - two-

way RM ANOVA, n = 12 GFP, n = 11 ArchT). **e.** Real-time place-preference location plot from a

representative animal while submitted to optogenetic activation of SST$^+$ vlPAG cells in the left

compartment throughout the 15-min session. **f.** There was no difference between ChR2- expressing

mice and the control group in the time spent on the stimulated and non-stimulated compartments

(ns, P = 0.1812, Mann Whitney test).

Supplementary Figure 6. SST$^+$ vlPAG cells photoactivation during conditioning does not

impact fear learning. a. Protocol for optogenetic manipulations during fear conditioning. On Day

2, mice received 5 CS-US associations. The US was either optogenetic stimulation alone or

optogenetic stimulation plus footshock. **b.** Average freezing levels during the CS-US association

of optogenetic stimulation alone. There was no difference in the overall freezing levels for the CS$^-$

and CS$^+$ between ChR2 and GFP (ns, P = 0.4804, opsin x CS - two-way RM ANOVA, n = 6 GFP,

n = 7 ChR2). **c.** During retrieval, there was no effect on freezing upon activation of SST$^+$ vlPAG

cells as an US (ns, P = 0.4804, opsin x CS effect - two-way RM ANOVA, n = 6 GFP, n = 7 ChR2).

d. Average freezing levels during the CS-US association of optogenetic stimulation plus footshock.

There was no difference in the overall freezing levels for the CS$^+$ and CS$^-$ between ChR2 and GFP

(ns, P = 0.8663, opsin x CS effect - two-way RM ANOVA, n = 6 GFP, n = 7 ChR2). **e.** During

retrieval, there was no difference in the fear expression by the activation of the SST$^+$ vlPAG cells

(ns, P = 0.8967, opsin x CS effect - two-way RM ANOVA, n = 6 GFP, n = 7 ChR2).

Supplementary Figure 7. Optogenetic effect observed during CFCA is not due to alteration

of somatostatin levels nor mediated by VIP$^+$ vlPAG cells. a. Sst-IRES-Cre heterozygotic mice

received a bilateral injection of opsins in the vlPAG, and optic fibers were implanted above the

vlPAG. **b.** A representative example of expression patterns of ChR2 within SST$^+$ vlPAG cells. **c.**

Average freezing values during retrieval. ChR2 and GFP groups were pooled together because no

differences were found in the conditioning level (data not shown). The average freezing values

during CS$^+$ was higher than CS$^-$ or baseline (BL) periods (****, P < 0.0001, Friedman test, n = 13

mice). **d.** Optogenetic activation of SST$^+$ vlPAG cells abolished the analgesic effect induced by

defensive states (**, P = 0.0048, opsin x CS - two-way RM ANOVA, n = 6 GFP, n = 7 ChR2).

The mean latency of NC response for the CS$^+$ was significantly different between the ChR2 and

GFP groups (*, P = 0.0300, Bonferroni post-hoc test, n = 6 GFP, n = 7 ChR2). For the ChR2 group,

the latency of NC response during CS$^+$ was equivalent to the CS$^-$ (ns, P = 0.3342, Bonferroni post-

hoc test). On the contrary, the latency of NC response between the CSs was different for the GFP

group (****, P = 0.0001, Bonferroni post-hoc test). **e.** VIP-IRES-Cre mice received a bilateral

injection of opsins in the vlPAG, and optic fibers were implanted above the vlPAG. **f.**

Representative example of the expression pattern of ArchT within VIP$^+$ vlPAG cells. **g.** ArchT and

GFP groups were pooled together because no differences were found in the conditioning level (data

not shown). The average freezing values during CS$^+$ was higher than CS$^-$ or BL periods (****, P <

0.0001, one-way RM ANOVA, n = 10 mice). **h.** Optogenetic inhibition of VIP$^+$ cells did not change

the analgesic effect for the ArchT group when compared to the GFP (ns, P = 0.5455, opsin x CS

effect - two-way RM ANOVA, n = 5 GFP, n = 5 ArchT). The latency of NC response was higher

for the CS$^+$ trials compared to the CS$^-$ trials for both GFP and ArchT (***, P < 0.01, Bonferroni

post-hoc test, n = 5 GFP, n = 5 ArchT).

Supplementary Figure 8. Optogenetic activation of SST$^+$ vlPAG cells did not affect tactile

sensitivity. a. Perievent stimulus time histogram indicating the spiking latency of a purely tactile

neuron recorded in the spinal cord in response to gentle brushing of the skin. **b.** Representative

spiking activity latency of a purely tactile neuron recorded in the spinal cord in response to gentle

brushing of the skin with or without activation of ChR2 expressing SST$^+$ vlPAG cells. **c.** The

activation of ChR2 expressing SST$^+$ vlPAG cells of tactile neuron recorded in the spinal cord in

response to gentle brushing of the skin had no effect on the average firing activity (Paired t-test, P

= 0.6929, n = 5 cells).

Supplementary Figure 9. Optogenetic manipulation of GFP expressing vlPAG cells did not

affect analgesia. a. Single-unit recordings of WDR cells in the lumbar spinal cord during

optogenetic manipulation of GFP expressing SST$^+$ vlPAG inputs to the RVM. **b**. Photostimulation

of SST$^+$ vlPAG inputs to the RVM has no effect.

Winke et al., Figure 1

Winke et al., Figure 2

Winke et al., Figure 3

Winke et al, Figure 4

Winke et al., Figure 5

Winke et al., Figure 6