A quantitative, genome-wide analysis in *Drosophila* reveals transposable elements' influence on gene expression is species-specific.

Marie Fablet[1,2], Judit Salces-Ortiz[1,6], Angelo Jacquet[1,7], Bianca F. Menezes[1,8], Corentin Dechaud[3], Philippe Veber[1], Camille Noûs[4], Rita Rebollo[5], Cristina Vieira[1]

[1] Laboratoire de Biométrie et Biologie Evolutive, Université de Lyon; Université Lyon 1; CNRS; UMR 5558, Villeurbanne, France.

[2] Institut Universitaire de France (IUF)

[3] Institut de Génomique Fonctionnelle de Lyon, Univ Lyon, CNRS UMR 5242, Ecole Normale Supérieure de Lyon, Université Claude Bernard Lyon 1, 46 allée d'Italie, F-69364 Lyon, France

[4] Laboratoire Cogitamus

[5] Univ Lyon, INRAE, INSA-Lyon, BF2I, UMR 203, 69621 Villeurbanne, France.

[6] Current address: Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra) P° Marítimo de la Barceloneta, 37-49 08003 Barcelona, Spain

[7] Current address: Symbiotron; FR3728 Biodiversité, Eau, Environnement, Ville, Santé; Université Claude Bernard Lyon 1; Villeurbanne 69622, France

[8] Current address: Federal Institute of Rio de Janeiro (IFRJ), Pinheiral – RJ, Brazil

Corresponding authors: marie.fablet@univ-lyon1.fr; cristina.vieira@univ-lyon1.fr

Abstract

Transposable elements (TEs) are parasitic DNA sequences that are able to move and multiply along the chromosomes of all genomes. They are controlled by the host through the targeting of silencing epigenetic marks, which may affect the chromatin structure of neighboring sequences, including genes. In this study, we used transcriptomic and epigenomic high-throughput data produced from ovarian samples of *Drosophila melanogaster* and *Drosophila simulans* wild-type strains, in order to finely quantify the influence of TE insertions on gene RNA levels and histone marks (H3K4me3 and H3K9me3). We find general, negative impacts of TE insertions on gene RNA levels and H3K4me3 enrichment, and global, positive effects on H3K9me3 enrichment. However, we also uncover that TE insertions within exons are associated with a reduction in H3K9me3, which we propose is associated with the removal of TE sequences from the transcript through splicing. Overall, we find that gene RNA and histone mark levels show contrasted dependencies on TE insertions regarding the species. The fold-decrease in TE-carrying gene proportions between extreme expression classes is 2.6 in *D. melanogaster* and 1.8 in *D. simulans,* while it is strongly contrasted between extreme classes of H3K4me3 enrichment: 4.4 in *D. melanogaster* and 19.2 in *D. simulans.* This provides a new light on the considerable natural variability provided by TEs, which may be associated with contrasted adaptive and evolutionary potentials.

Introduction

Transposable elements (TEs) are parasitic DNA sequences that are able to move and multiply along the chromosomes of all genomes (Wells and Feschotte, 2020). As such, they are source of mutations and genome instability if uncontrolled (Biémont and Vieira, 2006; Malone and Hannon, 2009; Senti and Brennecke, 2010). Control generally consists in the targeting of silencing epigenetic marks to TE copies, which may spread to neighboring sequences and impact gene expression. In this regard, few attempts were made to finely analyze and quantify TEs' influence at the whole genome scale (Cridland et al., 2015; Hollister and Gaut, 2009; Lee and Karpen, 2017; Uzunović et al., 2019). However, TE contents, activities

and control pathways vary in nature (Akkouche et al., 2013, 2012; Baduel et al., 2021; Mérel et al., 2021,

2020). The present study therefore sheds new light on this question integrating natural variability both at the

intra- and inter-specific levels in a *Drosophila* system. Specifically, we assessed the transcriptomic and

epigenetic status of genes in the ovaries of wild-type strains of *D. melanogaster* and *D. simulans* in relation

to TE content.

Previous studies focusing on a few TE families in wild-type strains of *D. simulans* and *D. melanogaster*

uncovered between-species differences in histone mark landscapes (Rebollo et al., 2012a). Lee and Karpen

(Lee and Karpen, 2017) previously provided an analysis on the repressive histone mark H3K9me2 (Histone

3 Lysine 9 dimethylation) around TEs from two Drosophila Genetic Reference Panel (DGRP) strains

(*D. melanogaster*), and concluded to pervasive epigenetic effects of TEs. Here, we focused on H3K9me3

(Histone 3 Lysine 9 trimethylation, repressive) and H3K4me3 (Histone 3 Lysine 4 trimethylation,

permissive) marks, which proved to be particularly relevant in TE epigenetic dynamics (Ozata et al., 2019),

and we produced genome-wide data from four wild-type strains of *D. melanogaster* and four wild-type

strains of *D. simulans*, which allows us to provide a quantitative assessment of the contribution of TEs to

gene expression, in a comparative genomics perspective (Fig. 1).

Previous research already described the effects of TE insertions on gene expression using collections of

strains of *D. melanogaster* (Cridland et al., 2015; Everett et al., 2020; Osada et al., 2017; Zhang et al., 2020).

The novelties of ours is 1) to bring together TE insertion polymorphism, RNA-seq, ChIP-seq on two histone

marks, and small RNA-seq data on the exact same strains, 2) to focus on ovaries, which is the organ where

strong selective pressures act on TE control (and not on whole bodies or mix of tissues), 3) to provide a new,

quantitative approach based on linear models, and 4) to reach a larger evolutionary scale thanks to the

inclusion of data not only from *D. melanogaster* strains but also from its sister species *D. simulans*. We

investigated the impacts of TEs on gene expression at the whole genome scale using eight previously

characterized, wild-type strains of *D. melanogaster* and *D. simulans* (Mohamed et al., 2020). These strains

derived from samples collected in France and Brazil, two strains per location and per species. Using the

Oxford Nanopore long read sequencing technology, we previously produced high quality genome assemblies

at the chromosome resolution for each of the strains (Mohamed et al., 2020).

Henceforth, we reveal that TE insertions within genes (exons and introns) are associated with stronger

negative effects on gene RNA levels and H3K4me3 enrichment compared to TE insertions upstream genes.

On the contrary, downstream TE insertions have positive impacts on RNA levels in *D. simulans* only. The

proportion of genes carrying TE insertions decreases when RNA levels and H3K4me3 increase, and

increases with H3K9me3 enrichment, a silencing histone mark known to target TE sequences (Le Thomas et

al., 2013). We also uncover a novel observation regarding TE insertions within exons: they are associated

with a reduction in H3K9me3 enrichment. Considering that H3K9me3 is involved in the definition of splice

sites (Bieberstein et al., 2016; Saint-André et al., 2011), we speculate that TE insertions within exons are

not purged by natural selection provided they are skipped from the corresponding transcripts, which may be

achieved through reductions in H3K9me3 marks. Across strains, we find that the more genes display

different RNA amounts the more likely they have different numbers of TE insertions. In addition, genes that

have more TE insertions tend to be enriched in H3K9me3 marks. As a whole, these results participate in the

accurate, quantitative understanding of TEs' impacts in genomes. Moreover, our major finding in this

research work is that, across genes, RNA and histone mark levels show contrasted dependencies on TE

insertions regarding the species, *D. melanogaster* or *D. simulans*. This provides a new light on the

considerable natural variability resulting from TEs, which may be associated with contrasted adaptive and

evolutionary potentials, all the more sensible in a rapidly changing environment (Baduel et al., 2021; Fablet

and Vieira, 2011; Mérel et al., 2021).

Figure 1. Graphic summary of the study.

Eight wild-type strains from *D. melanogaster* and *D. simulans* were included in the study. The present

datasets are RNA-seq and ChIP-seq for H3K4me3 and H3K9me3 marks, and were prepared from ovarian

samples. They were analyzed in parallel with already published data produced from the same drosophila

strains: ovarian piRNA repertoires and genome assemblies based on Oxford Nanopore long read sequencing

(Mohamed et al., 2020). For RNA-seq and ChIP-seq, TE-derived reads were analyzed at the TE family level,

and gene-derived reads were analyzed in relation to TE insertions inside or near genes (therefore restricted to

the TE insertions included within the gray bubbles).

Results

TE expression and epigenetic targeting in Drosophila ovaries

We first considered TE-derived RNA-seq reads from all our samples, which we analyzed at the TE-family

level (Fig. 1). TEs account for 0.6% to 1.2%, and 0.5% to 0.7%, of read counts corresponding to annotated

sequences (genes and TEs) within the ovarian transcriptomes of our strains of *D. melanogaster* and

D. simulans, respectively (Fig. 2A). We find significant positive correlations between per TE family RNA

counts and family sequence occupancy (total number of bp spanned by each TE family along the genome)

(Spearman correlations, rho = 0.20 to 0.31, and 0.28 to 0.35, in *D. melanogaster* and *D. simulans*,

respectively; Supplemental_Fig_S2A). We also find significant positive correlations between per TE family

RNA counts and piRNA counts (Spearman correlations, rho = 0.65 to 0.70, and 0.64 to 0.68, in

D. melanogaster and *D. simulans*, respectively; Supplemental_Fig_S2B), as already described for wild-type

strains (Lerat et al., 2017).

We assessed the contribution of histone mark enrichment to TE RNA amounts considering the following

linear model on log-transformed normalized read counts: RNA ~ H3K4me3 + H3K9me3 + input. These

models led to adjusted r^2 as high as 0.48 to 0.64, and 0.45 to 0.60, in *D. melanogaster* and *D. simulans*

strains, respectively, suggesting that this approach captures a significant portion of TE RNA amount

variation. We find that TE RNA amounts are positively correlated with H3K4me3 and negatively correlated

with H3K9me3 amounts (Fig. 2B), as expected considering that H3K4me3 is an activating mark while

H3K9me3 is a silencing one. We used a similar approach to analyze piRNA amounts, and considered the

following linear model on log-transformed read counts: piRNA ~ H3K4me3 + H3K9me3 + input. We

obtained even higher adjusted r^2 values, from 0.71 to 0.75, and 0.64 to 0.68, in *D. melanogaster* and

D. simulans strains, respectively.   We find that TE-derived piRNA amounts are positively correlated both

with permissive H3K4me3 and repressive H3K9me9 enrichment (Fig. 2C).

TE insertions within or near genes

In the following sections, we focus on gene-derived reads from all our samples, which we analyzed in

relation with the presence of TE insertions within or near genes. Based on gene annotations, we

distinguished the different functional regions of genes: exons, introns, upstream, or downstream sequences

(5 kb flanking regions). In the present study, we considered a set of 17,417 annotated genes for

D. melanogaster, and 15,251 for *D. simulans* (see Material and Methods). We quantified the number of TE

insertions within genes (Fig. 2D), and found that they account for ~10% of the total number of TE insertions

per genome. The majority of genes (93%) are devoid of TE insertions within gene bodies, and 6% have

exactly one TE insertion (Fig. 2E, Supplemental_Fig_S1). Regarding flanking regions, 92% to 94% of genes

are devoid of TE insertions upstream (over 5 kb), and 6% to 8% display exactly one insertion. The

proportions are the same regarding the downstream flanking regions.

Figure 2. (*A*) Proportions of TE read counts in RNA-seq data over read counts corresponding to genes and

TEs. For each strain, two biological replicates are shown. (*B*) Contributions of H3K4me3 and H3K9me3

enrichment to TE-derived RNA read counts (according to the model RNA ~ H3K4me3 + H3K9me3 + input

calculated on log transformed read count numbers, at the TE family level). Plain-colored bars: p-values

< 0.01, light-colored bars: p-values in [0.01; 0.05], empty bars: p-values > 0.05. (*C*) Contributions of

H3K4me3 and H3K9me3 enrichment to TE-derived piRNA read counts (according to the model piRNA ~

H3K4me3 + H3K9me3 + input calculated on log transformed read count numbers, at the TE family level).

Plain-colored bars: p-values < 0.01, light-colored bars: p-values in [0.01; 0.05], empty bars: p-values > 0.05.

(*D*) Numbers of TE insertions per functional region per strain. Upstream and downstream regions are 5 kb

sequences directly flanking transcription units 5' and 3', respectively. (*E*) Numbers of genes for each value of

TE insertion numbers. dmgoth101 and dsgoth31 are shown as examples; all strains can be found in

Supplemental_Fig_S1. Error bars are standard errors.

TEs contribute to RNA level variability across genes within genomes

One of the objectives of the present work was to quantify the association between TE insertions and the

differences in expression levels that we observe across the set of genes of a genome. We focused on the

subset of genes that we found expressed in the ovaries (see Material and Methods). They account for 8,028

to 8,209 genes depending on the strains, and 6,149 to 6,431 genes, in *D. melanogaster* and *D. simulans*,

respectively. We quantified TE insertion contribution to gene RNA levels using the following linear models

built on log-transformed TPM (Transcript Per Million): TPM ~ exon + intron + upstream + downstream,

where these variables correspond to the numbers of TE insertions in exons, introns, 5 kb upstream, and 5 kb

downstream regions. We found that the most pervasive effect is a negative correlation between gene RNA

amounts and the number of TE insertions in introns (Fig. 3A). TE insertions that are located upstream of

genes are not correlated to gene transcript amounts on average. On the contrary, TE insertions that are

located downstream of genes are associated with increased gene transcript amounts, however only in

D. simulans strains (Fig. 3A; Wilcoxon test comparing downstream effect sizes across *D. melanogaster* and

D. simulans: p-value = 0.029).

Then, considering each genome individually, we defined 10 TPM classes. Using generalized linear models to

analyze the TE presence / absence variable against the TPM class factor, we could reject independence for all

D. melanogaster strains and two *D. simulans* strains (Supplemental_Table_S1). The dependency nicely fits

to negative linear correlations (Fig. 4A). This indicates that the more the genes are transcribed, the lower the

probability that they carry TE insertions. Depending on the strain, we observe a 2.2 to 2.9-fold decrease in

the proportions of such genes between the two extreme expression classes in *D. melanogaster*, and 1.5 to 2.2

in *D. simulans*. These negative correlations still hold when considering only introns or exons, but not in the

case of upstream or downstream flanking regions (Supplemental_Fig_S3).

One could object that the differences we observe between *D. melanogaster* and *D. simulans* are due to the

gene sets being different between species rather than to the intrinsic specific differences. We therefore

performed the same analyses for the subset of 4,498 genes that have 1:1 ortholog in the other species and that

are expressed in ovaries. Similarly, in all strains but dssj9, we uncover significant dependency between

proportion of TE-carrying genes and bin of expression (generalized linear model, Supplemental_Table S1).

However, this leads to patterns that are now highly similar between *D. melanogaster* and *D. simulans* (Fig.

4B). We speculate that this is due to the specificity of 1:1 orthologous genes because they are enriched in

conserved genes that are the most likely to evolve under purifying selection. In this respect, we found that

the contrast between *D. melanogaster* and *D. simulans* strains is even stronger when we consider all but 1:1

orthologous genes: an average 3.9-fold decrease between extreme expression classes in *D. melanogaster vs*

an average 1.8-fold decrease in *D. simulans* (Fig. 4C).

TEs contribute to histone enrichment variability across genes within genomes

We used a similar approach to analyze H3K4me3 and H3K9me3 enrichment (*i.e.* we aligned ChIP-seq reads

against whole gene sequences and computed corresponding read counts). As for RNA levels, TE insertions

within introns are significantly negatively correlated with H3K4me3 amounts (Fig. 3B). We could not detect

consistent, significant impacts from insertions in flanking regions.

Considering H3K9me3 amounts, the impacts of TE insertions are stronger. Using the model H3K9me3 ~

input + exon + intron + upstream + downstream, we detect significant, positive contributions of TE

insertions in introns, upstream, and downstream (Fig. 3C). These results are in agreement with TEs being the preferential targets for H3K9me3 deposition, which then spreads to neighboring regions (Le Thomas et al., 2013; Rebollo et al., 2011). However, we also find that TE insertions within exons have a clear, negative impact on gene H3K9me3 amounts (Fig. 3C).

Then, we classified genes according to their histone mark enrichment levels, and computed the proportions of TE-carrying genes for each class (Fig. 4D,G). Again, we could clearly reject independence between the proportion of TE-carrying genes and histone enrichment class (generalized linear model, Supplemental_Table_S1). We find significant, negative linear correlations between these proportions and H3K4me3 enrichment levels, which are stronger for *D. simulans* strains compared to *D. melanogaster*. The reduction in the proportion of TE-carrying genes is 3.1 to 5.7-fold, and 17.1 to 20.5-fold between the extreme histone classes, in *D. melanogaster* and *D. simulans*, respectively (Wilcoxon test contrasting both species, p-value = 0.029). When considering detailed functional regions, we found that this effect holds true for exons and introns, but not for flanking regions (Supplemental Fig S4).

Regarding H3K9me3, we reject independence (generalized linear model, Supplemental_Table_S1) and found significant, strong positive correlations between the proportion of TE-carrying genes and histone levels in *D. simulans* genomes: the proportion increases 8.3 to 11.7-fold between the extreme histone classes. On the contrary, no linear correlations are found in *D. melanogaster* genomes. However, we detected strong, positive correlations between the proportions of genes carrying TE insertions within flanking regions (both upstream and downstream) and H3K9me3 amounts, consistently across all eight genomes (Supplemental Fig S5).

We analyzed separately 1:1 orthologous genes, and found that *D. melanogaster vs D. simulans* differences disappear for H3K4me3 enrichment (Fig. 4E), while they are more contrasted when considering all but 1:1 orthologous genes (Fig. 4F), suggesting once more that differences between *D. melanogaster* and *D. simulans* are held of the fraction of the least conserved genes. On the contrary, patterns for H3K9me3 enrichment are consistent across sets of genes (Fig. 4G-I)

Figure 3. (*A*) Contribution of TE insertion numbers (within exons, introns, or in the 5 kb upstream or downstream sequences) to gene-derived RNA read counts. (*B*) Contribution of TE insertion numbers (within exons, introns, or in the 5 kb upstream or downstream sequences) to gene H3K4me3 enrichment. (*C*) Contribution of TE insertion numbers (within exons, introns, or in the 5 kb upstream or downstream sequences) to gene H3K9me3 enrichment. Plain-colored bars: p-values < 0.01, light-colored bars: p-values in [0.01; 0.05], empty bars: p-values > 0.05. Error bars are standard errors.

Figure 4. (*A-C*) Proportions of genes carrying at least one TE insertion according to gene expression classes. We plotted 0.95 confidence intervals for the proportion of TE-carrying genes and we fitted a linear regression line with mean TPM per class (Pearson correlation coefficients and the corresponding p-values are indicated). (*D-F*) Proportions of genes carrying at least one TE insertion according to gene H3K4me3 enrichment classes. (*G-I*) Proportions of genes carrying at least one TE insertion according to gene H3K9me3 enrichment classes. We plotted 0. 95 confidence intervals for the proportion of TE-carrying genes and we fitted a linear regression line with mean histone enrichment per class (Pearson correlation coefficients and the corresponding p-values are indicated). *A, D, G* correspond to the complete sets of genes that are expressed in ovaries; *B, E, H* to 1:1 orthologous genes; and *C, F, I* to all but 1:1 orthologous genes.

Private and shared TE insertions are associated with similar patterns

It has already been described that TE impacts on genes differ across young (*i.e.* polymorphic) and old (*i.e.* fixed) TE copies; this is due to the pool of old TE insertions having been purged from deleterious insertions by natural selection (Hollister and Gaut, 2009). Indeed, Uzunovic *et al.* (Uzunović et al., 2019) showed in the plant *Capsella* that young TE insertions had a negative effect on gene expression while old insertions were more likely to increase expression. In this view, we performed the same analyses as above but on TE insertions that are unique to one genome ("private") and those that are shared by all four strains of the species ("common"). The majority of the TE insertions (>80%) are common, leading to differences in statistical power between both sets of insertions. Despite this bias, the observed patterns are highly similar (Supplemental_Table_S2). Since our approach is gene-centered, it is very likely that our primary set of TE insertions is already biased: when deleterious, insertions within or near genes have such a negative impact that we are not able to catch them from natural samples. Therefore, our primary set of TE insertions may already correspond to copies that have passed the filter of natural selection. Alternatively, we cannot exclude that they may also lie in particular chromatin environments where there is retention bias (Sultana et al., 2017), and that the associations detected here are due to these particular chromatin features.

TE insertions are associated with variability in expression in histone enrichment between orthologous genes

Next, we used our experimental dataset to infer the contribution of TE insertions at the inter-genomic level, *i.e.* we compared expression levels of the same genes across genomes. We first considered *D. melanogaster* and *D. simulans* separately. For each gene that displays variation in TE insertion numbers across strains, we computed the mean difference of gene expression (TPMs) between high and low TE insertion numbers (scaled by gene mean). We performed the same approach on histone enrichment. Our assumption was that a global effect of TE insertions would shift the distribution of the mean difference away from 0. However, this is not what we observed for RNA levels (0 departure t test; *D. melanogaster*: mean difference = 0.009, p-

value = 0.644); *D. simulans*: mean difference = -0.015, p-value = 0.507) nor for H3K4me3 enrichment (0

departure t test; *D. melanogaster*: mean difference = 0.002, p-value = 0.445); *D. simulans*: mean difference

= -5e-5, p-value = 0.948). On the contrary, we find a clear increase in H3K9me3 enrichment associated with

high TE insertion numbers (0 departure t test; *D. melanogaster*: mean difference = 0.004, p-value = 0.002;

D. simulans: mean difference = 0.005, p-value = 3e-5). The signal for H3K9me3 enrichment is the strongest

for TE insertions in introns and upstream regions (Supplemental Fig S6).

We also took the opportunity to consider 1:1 orthologous genes so as to include all eight strains

(*D. melanogaster* and *D. simulans*) in the same analysis. Computation strategies were the same as above and

revealed a decrease in RNA levels for high TE insertion numbers (mean difference = 0.077, pvalue = 0.024)

but no significant differences in the H3K4me3 (mean difference = 0.023, pvalue = 0.601) nor H3K9me3

enrichment (mean difference = 0.016, p-value = 0.149). However, such an analysis considering all strains

from both species at once has to be considered cautiously because gene sequences differ across species (GC

content, length, etc.), which may interfere with mapping and read counting, and was not accounted for in this

work.

That being said, it is not unlikely that the direction of the effect of TEs on gene expression, if any, may vary

across TE insertions. Therefore, we decided to perform an unpolarized analysis as well —*ie* blind to the

direction of the effect. In this respect, we computed pairwise comparisons and we calculated the coefficient

of variation (CV, square root of the variance divided by the mean) of TPM for genes that are transcribed in

our ovarian samples. We defined 10 classes based on CV values, and computed the proportion of genes that

displayed different numbers of TE insertions within each class. The general trend is positive correlation

between these proportions and the mean CVs of each class, and the adjustment is generally better using the

square root of mean CV. This indicates that the more the genes show differences in expression levels, the

greater the probability that they have different numbers of TE insertions (Fig. 5C, Supplemental_Fig_S7,

Supplemental_Fig_S8). The effect appears to be stronger in *D. melanogaster* compared to *D. simulans*.

Figure 5. Variability in gene expression and histone enrichment according to TE insertion numbers

across strains. (*A*) Mean expression difference (scaled by gene average) between strains with high and low

TE insertion numbers; mean histone enrichment difference (log-transformed, scaled by gene average)

between strains with high and low TE insertion numbers. Analyses are performed separately for both species.

Significance levels correspond to t tests comparing observed mean to 0. (*B*) Same analyses across all eight

strains considering 1:1 orthologous genes. Significance levels correspond to t tests comparing observed mean

to 0: 0 *** 0.001 ** 0.01 * 0.05 · 0.1. (*C*) For each pair of strains, proportions of orthologous genes carrying

different numbers of TE insertions according to differences in gene RNA levels (computed based on

variation coefficient). We plotted 0.95 confidence intervals for the proportion of genes carrying different

numbers of TEs and we fitted a linear regression line with square root of mean CV per class (Pearson

correlation coefficients and the corresponding p-values are indicated).

Discussion

The common-held view is that, as parasites that are fought against by the genomes, TEs have a general negative impact on gene expression (Cridland et al., 2015; Lee, 2015; Lee and Karpen, 2017). Our present findings are in agreement with this idea. However, the originality of this research work is to provide an unprecedented quantitative view, which allows to precisely decipher TE impacts, integrating intra- and inter-species data from wild-type strains of two closely related drosophila species. In addition, this study combines genomic, transcriptomic, and epigenetic high-throughput sequencing data, all produced from the same strains and tissues, which allows to draw actual connections between the different nucleic repertoires of the cell. Moreover, the chosen tissue is ovary, which is known to be the place of the strongest epigenetic control of TE sequences, through the piRNA pathway (Malone and Hannon, 2009; Senti and Brennecke, 2010). This is therefore where we are to expect the strongest impacts of TEs on genes.

Expression and epigenetic marks of TE sequences

Our results uncover a lower contribution of TEs to the *D. simulans* transcriptome as compared to *D. melanogaster*. This is in agreement with the previously described lowest contribution of TEs in the genomes of *D. simulans*, in terms of sequence occupancy and copy numbers (Mohamed et al., 2020; Vieira et al., 1999). In both species, we found that H3K9me3 marks on TE sequences are associated with a decrease in TE-derived RNA amounts, and the opposite for H3K4me3 marks. On the contrary, we observed that both histone marks are positively correlated with TE-derived piRNA amounts, which is congruent with the piRNA-targeted deposition of H3K9me3 marks at transcriptionally active TE copies (Czech et al., 2018). However, one should note that these results reflect average behaviors at the TE family level, but TE copies may differ from one another within TE families.

TE impact on gene expression and histone marks

Consistent with an expected general negative impact of TEs on gene expression, our analyses reveal that TE insertions are associated with a general decrease in gene RNA levels and H3K4me3 enrichment, except for downstream insertions in *D. simulans*. This last observation may be related to TEs donating enhancers, as was already described on several instances (Moschetti et al., 2020; Sundaram et al., 2014; Villanueva-Cañas et al., 2019) or disrupting inhibitory sequences. Globally, we found that the impacts of TE insertions on RNA levels and histone enrichment are stronger for TE insertions within gene bodies (exons and introns) compared to TE insertions in flanking regions. This had already been observed by Cridland *et al.* (Cridland et al., 2015), who found a stronger correlation with extreme gene expression for TEs inserted into genes compared to TEs upstream of genes.

Considering that TEs are a preferential target for H3K9me3 deposition (Le Thomas et al., 2013), we expected a positive effect of TE insertion numbers on H3K9me3 enrichment. This is what we clearly observed for TE insertions within introns and flanking regions. However, we found a so far undescribed negative influence of TE insertions in exons on H3K9me3 enrichment. This finding may be explained by the interaction between H3K9me3 and the splicing machinery: it was shown in mammals that H3K9me3 favors the inclusion of alternative exons while demethylation is associated with exon skipping (Bieberstein et al., 2016; Saint-André et al., 2011). We propose that the TE insertions in exons that are maintained and not purged by natural selection, are those that are spliced out from transcripts, so that the coding sequence of the gene is not impaired. This specific splicing would be possible for TE insertions that are associated with a reduction in H3K9 methylation, which allows the splicing out of the TE. Alternatively, we cannot rule out the possibility that exonic regions that are resistant to formation of H3K9me3 are also more likely to have TE insertions, which would result in a negative association of TE insertions in exons and H3K9me3 enrichment.

As a conclusion to the intra-genomic analysis, we have to note that there are many parameters other than the numbers of TE insertions that differ across the genes (the family and length of the TEs, gene sequence composition, presence of transcription factor binding sites, etc. (Hill et al., 2021; Wittkopp and Kalay, 2011)) and yet we were able to capture statistical signal from the numbers of TE insertions. This suggests of a widespread, strong influence of TEs on gene expression. Nevertheless, we have to acknowledge that our

study uncovers here the impacts of TEs on gene expression through epigenetic modifications; however, this is not the only possible mechanism for such an influence. TEs may also disrupt functional elements, especially for those inside genes, or add transcription factor binding sites (Horváth et al., 2017; Rebollo et al., 2012b; Ullastres et al., 2021). Moreover, we have to note that TE insertions may accumulate in specific chromatin environments due to insertional preference or different levels of selection in these environments (Sultana et al., 2017). In such cases, this behavior would prevent us from disentangling accurately the impact of the TE insertion in itself.

In addition to contributing to variability across the genes of a genome, TEs also participate in variability across genomes through insertion polymorphism. Indeed, our analysis across genomes reveals that genes with the highest varibility in TE insertion numbers also display the highest variability in RNA levels, and this is clearer in *D. melanogaster* compared to *D. simulans*. However, the analysis combining data from *D. melanogaster* and *D. simulans* uncovers an average negative effect of TE insertion numbers on RNA levels. Similarly, more TE insertions are associated with higher H3K9me3 enrichment.

TE impacts on genes are different between *D. melanogaster* and *D. simulans*

Our analysis separating 1:1 ortholous genes from the others reveals that the impacts of TEs on genes are very similar between *D. melanogaster* and *D. simulans* for this particular set of genes, which corresponds to the most conserved and constrained. However, differences between species appear when considering all but 1:1 orthologous genes expressed in the ovaries. For these more rapidly evolving set of genes, we find that the epigenetic impacts of TEs are more pronounced in *D. simulans*. Indeed, we found that the probability that a gene carries at least one TE insertion is more strongly correlated with its histone enrichment in *D. simulans* compared to *D. melanogaster* (Fig. 6). Similarly, although using a different, more indirect genome-wide approach, Lee and Karpen (Lee and Karpen, 2017) also came to the conclusion that, at the same time, *D. simulans* displays a lower TE content and stronger epigenetic effects of TE control, compared to *D. melanogaster*. Again, it sustains the idea that *D. simulans* epigenetic machinery is less efficient since a strong pattern in H3K9me3 and/or H3K4me3 results in a weak pattern at the RNA levels, or in other words,

since weak histone patterns in *D. melanogaster* are enough to produce a strong RNA response. Alternatively, upstream of the regulatory effectors, it may suggest that the targeting of TE sequences for histone mark deposition is more efficient in *D. simulans*, maybe because the colonization of the *D. simulans* genome by TEs started more recently (as suggested by our previous results (Mohamed et al., 2020)). We expect that the strength of the epigenetic control of TEs should decrease after the colonization, as the equilibrium between the genome and its TEs is reached. In this respect, we may propose that, rather than a species-specific influence of TEs on gene expression, we uncovered a contrasted influence related to the time point in TE dynamics where the considered species lies. Another explanation for the differences that we highlight regarding the influence of TEs on genes between *D. melanogaster* and *D. simulans* is that the proportions of the different TE families are different (Mohamed et al., 2020). Depending on the TE class, family, length distribution, etc., we expect that the effects of TE insertions and their epigenetic effects vary. However, it seems difficult to use the present dataset to tackle this question because splitting the TE repertoire in too many categories would result in a loss of statistical power.

In any case, these contrasted impacts of TE insertions on genes through epigenetic marks across the species provide an additional demonstration of the considerable natural variability due to TEs. We predict that this leads to contrasted adaptive and evolutionary potentials, all the more sensible in a rapidly changing environment (Baduel et al., 2021; Fablet and Vieira, 2011; Mérel et al., 2021).

Figure 6. Graphic summary. The probability that a gene carries at least one TE insertion is more strongly correlated with its histone enrichment in *D. simulans* compared to *D. melanogaster*. On the contrary, this probability is more strongly correlated with gene RNA levels in *D. melanogaster*.

Methods

Drosophila strains

The strains under study in the present work were previously described in Mohamed et al. (Mohamed et al., 2020). The eight samples of *D. melanogaster* and *D. simulans* wild-type strains were collected using fruit baits in France (Gotheron, 44°56'0"N 04°53'30"E - "goth" strains) and Brazil (Saõ Jose do Rio Preto 20°41'04.3"S 49°21'26.1"W – "sj" strains) in June 2014. Two isofemale lines per species and geographical origin were established directly from gravid females from the field (French *D. melanogaster*: dmgoth63, dmgoth101; Brazilian *D. melanogaster*: dmsj23, dmsj7; French *D. simulans*: dsgoth613, dsgoth31; Brazilian *D. simulans*: dssj27, dssj9). Brothers and sisters were then mated for 30 generations to obtain inbred strains with very low intra-line genetic variability. Strains were kept at 24°C in standard laboratory conditions on cornmeal–sugar–yeast–agar medium.

Genome annotation

Genome assemblies were produced in (Mohamed et al., 2020) and have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB50024 (https://www.ebi.ac.uk/ena/browser/view/PRJEB50024).

We used the genome assemblies produced for each of the eight strains using the Oxford Nanopore sequencing technology, and previously published (Mohamed et al., 2020). Throughout the present analysis, we kept scaffolds corresponding to complete chromosomes 2L, 2R, 3L, 3R, 4, and X.

TE annotation: We used RepeatMasker 4.1.0 (http://repeatmasker.org/) -species Drosophila in order to identify TE sequences in the assemblies, followed by OneCodeToFindThemAll (Bailly-Bechet et al., 2014) with default parameters, in order to parse RepeatMasker results.

Gene annotation: For *D. melanogaster*, we retrieved dmel-all-exon-r6.16.fasta, dmel-all-intron-r6.16.fasta, and dmel-all-gene-r6.16.fasta, which are available at ftp.flybase.net/genomes/Drosophila_melanogaster/dmel_r6.16_FB2017_03/fasta/. For *D. simulans*, we retrieved dsim-all-exon-r2.02.fasta, dsim-all-intron-r2.02.fasta, and dsim-all-gene-r2.02.fasta, which are

available at ftp.flybase.net/genomes/Drosophila_simulans/dsim_r2.02_FB2017_04/fasta/. These different sequence sets were aligned to our assemblies using BLAT (Kent, 2002) and pslReps from the BLAT suite, in order to keep the best hit (-singleHit). We filtered out genes with 0 hit and genes with multiple hits. For each gene, we also checked that the positions of the hits for each of the internal regions (*i.e.* exon, intron) were included in the positions of the complete gene hit. We excluded regions that did not fulfill these criteria. We also excluded hits that were not on the same chromosome as the corresponding gene on the reference genome. Then, we used the GenomicRanges R package (version 1.38.0) (Lawrence et al., 2013) and the subsetByOverlaps function to cross gene and TE annotations.

1:1 orthologs: We retrieved ortholog information from FlyBase (ftp://ftp.flybase.net/releases/current/precomputed_files/orthologs/ dmel_orthologs_in_drosophila_species_fb_2022_01.tsv.gz) and kept only those genes for which there was a 1 to 1 correspondence between *D. melanogaster* and *D. simulans*.

TE genomic sequence occupancy (bp) was computed using OneCodeToFindThemAll (Bailly-Bechet et al., 2014). Data corresponding to the *DNAREP1* transposon (also known as *INE-1*) were removed because it is a highly abundant element displaying mainly fixed insertions in the *melanogaster* complex of species (Chakraborty et al., 2021).

In order to determine which TE insertions were common (shared) to the four strains of a species or unique (private) to one strain, we performed pairwise comparisons of TE gff using Liftoff (Shumate and Salzberg, 2020). Private insertions to one strain are those that appear in the unmapped output of the three other strains. Common insertions are those that are found in all pairwise comparisons.

Small RNA extraction, sequencing, and analyses

Small RNA extraction, sequencing, and analyses had already been performed and described in Mohamed et al. (Mohamed et al., 2020). Sequence files had been deposited in NCBI SRA under the accession number PRJNA644327.

RNA-seq preparation

RNA was extracted from ovaries of 30 3-to-5 day-old females. Two replicates per strain were produced. RNA extraction was carried out using RNeasy Plus (Qiagen) kit following manufacturer's instructions. After DNAse treatment (Ambion), quality control was performed using an Agilent Bioanalyzer. Libraries were constructed from mRNA using the Illumina TruSeq RNA Sample Prep Kit following manufacturer's recommendations. Libraries were sequenced on Illumina HiSeq 3000 with paired-end 150 nt reads.

RNA-seq analysis

TE read counts were computed at the family level using the TEcount module of TEtools (Lerat et al., 2017) and the list of TE sequences available at ftp://pbil.univ-lyon1.fr/pub/datasets/Roy2019/. In agreement with other research works (Chakraborty et al., 2021), we removed the *DNAREP1* family from the analyses because it is a highly abundant element displaying mainly fixed insertions in the *melanogaster* complex of species (it is also known as *INE-1*, and corresponds to DNAREP1_DM, DNAREP1_DSim, and DNAREP2_DSim entries from our TE sequence list).

Genome and gene annotations: The genomes of *D. melanogaster* and *D. simulans* were downloaded from flybase (http://flybase.org/) along with GTF gene annotations. *D. melanogaster*: *f*tp://ftp.flybase.net/genomes/Drosophila_melanogaster/dmel_r6.16_FB2017_03/fasta/dmel-all-chromosome-r6.16.fasta.gz; ftp://ftp.flybase.net/genomes/Drosophila_melanogaster/dmel_r6.16_FB2017_03/gtf/dmel-all-r6.16.gtf.gz. *D. simulans*: ftp://ftp.flybase.net/releases/FB2017_03/dsim_r2.02/fasta/dsim-all-chromosome-r2.02.fasta.gz; ftp://ftp.flybase.net/releases/FB2017_03/dsim_r2.02/gtf/dsim-all-r2.02.gtf.gz.

Reads mapping: The alignment of all read samples on their corresponding reference genome was performed with Hisat2 version 2.1.0 (Kim et al., 2019) using the following command line options: hisat2-build $Genome $GenomeIndex to build an index of the genome; hisat2 -p 7 -k 2 --dta -x $GenomeIndex -1 $Sample_1.fq.gz -2 $Sample_2.fq.gz -S $Sample.sam to align each sample on the reference genome. The --dta option is used as recommended in the New tuxedo pipeline to perform an assembly of the transcripts (Pertea et al., 2016). The -k 2 option allows to report a maximum of two alignments per sequencing read. As we wanted to exclude potentially expressed TEs from the assembly, we discarded multimapped reads using

the following command line: samtools view -h -@ 7 -f 0x2 $Sample.bam |awk '{if(substr($1, 0, 1)=="@" ||

$5==60){print $0}}' >$Sample.sam. The option -f 0x2 extracts the properly aligned paired reads from the

bam. The awk command also keeps the header section of the SAM file as well as the reads with a

MAPQ=60. In Hisat2, a MAPQ=60 means that the read has been uniquely aligned regardless of the

mismatches / indels number. The filtered SAM file is finally converted to BAM and sorted: samtools view -h

-@ 7 -b -S $Sample.sam >$Sample_filtered.bam && samtools sort -@ 7 $Sample_filtered.bam sorted.bam

&& mv sorted.bam $Sample_filtered.bam.

Transcripts assembly: We used StringTie (Pertea et al., 2015) to assemble the transcripts using the genomic

coordinates of the aligned reads, for each sample: stringtie -p 7 -o $Sample_assemb.gtf -j 2

$Sample_filtered.bam && echo "$Sample$_assemb.gtf" >> mergelist.txt. The -j option specifies that at least

two reads are needed to support an exon-exon junction and create a new transcript. This is thus more

stringent than the default value of 1 used by StringTie. One assembly per sample is created; we used the

following command line to merge these assemblies: stringtie --merge -o merged.gtf mergelist.txt.

Transcript expression quantification: Stringtie was also used to quantify transcript expression in each sample:

stringtie -e -B -p 7 -G merged.gtf -o $Sample/$Sample.gtf $Sample_filtered.bam. -e option is used to

estimate the expression of transcripts of the GTF file given to the -G option. -B option is used to create an

output needed to run ballgown (Frazee et al., 2015) in the next step.

Count   matrix   generation: We   used   the   prepDE.py

(https://github.com/gpertea/stringtie/blob/master/prepDE.py) script from StringTie (Pertea et al., 2015) to

generate the count matrix from the output of the previous step.

Gene expression level quantification: We used the ballgown R package (Frazee et al., 2015) to estimate the

TPM (Transcript Per Million) expression of each gene or transcript. Data are loaded with ballgown from the

output of StringTie -e -B. Transcripts with low expression are filtered out as recommended in the new tuxedo

procedure (Pertea et al., 2016) using: subset(data,"rowMeans(texpr(data))>0.5", genomesubset=TRUE).

FPKM were then calculated using texpr function from ballgown. TPM where finally computed from this

table by normalizing the sum of each sample to 1e6.

Comparison to gene annotation: Genes and transcripts reconstructed by StringTie were compared to the

published gene annotation of the genome using gffcompare

(https://ccb.jhu.edu/software/stringtie/gffcompare.shtml). Each new transcript was assigned a class code

depending on its similarity to a reference transcript, as described in gffcompare documentation.

Quantification of the effects of TE insertions on gene transcript levels: considering only genes expressed in

ovaries, we computed mean TPMs across replicates and used the following linear models after log

transformation: TPM ~ exon + intron + upstream + downstream, where "exon", "intron" "upstream", and

"downstream" are the numbers of TE insertions in exons, introns, 5 kb upstream sequences, and 5 kb

downstream sequences, respectively. Size effects for each of these factors were then recorded. In parallel, for

each strain, we defined 10 classes of genes based on mean TPM values, and computed the proportion of

genes carrying at least one TE insertion in each of the 10 classes.

For the intergenomic analysis, we considered all pairs of strains within species, and for each pair of strains

and each gene, we computed the difference in TE insertion numbers on one hand and the coefficient of

variation (CV, square root of variance divided by mean) for TPM on the other hand. We then defined 10

classes of genes based on CV values and, for each class, we computed the proportions of genes carrying

different numbers of TEs.

ChIP-seq preparation

Chromatin immunoprecipitation was performed using 50 ovary pairs dissected from three to five day old

females. Ovaries were re-suspended in A1 buffer containing 60mM KCl, 15mM NaCl, 15mM Hepes, 0,5%

Triton and 10mM Sodium butyrate. Formaldehyde (Sigma) was added to a final concentration of 1.8% for

secondary cross-linking for 10 min at room temperature. Formaldehyde was quenched using glycine (0.125

M). Cross-linked cells were washed and pelleted twice with buffer A1, once with cell lysis buffer (140mM

NaCl, 15mM Hepes, 1mM EDTA, 0.5mM EGTA, 1% Triton X100, 0.1% Sodium deoxycholate, 10mM

sodium butyrate) , followed by lysis in buffer containing ,140mM NaCl, 15mM Hepes, 1mM EDTA,

0.5mMEGTA, 1% Triton X100, 0.5% SDS, 0.5% N-Laurosylsarcosine, 0.1% sodium deoxycholate, 10mM

sodium butyrate for 120 min at 4°C. Lysates were sonicated in Bioruptor sonicator to reach a fragment size

window of 200-600 bp.

Chromatin was incubated overnight at 4°C with the following antibodies: for H3K9me3 ChIP using α-

H3K9me3 (actif motif #39161, 3μg/IP) and for H3K4me3 using α-H3K4me3 (millipore #07-473, 3μg/IP)

antibodies. The Magna ChIP™ A/G Chromatin Immunoprecipitation Kit (cat# 17-10085) was used

following manufacturer's instructions. Final DNA recovery was perfomed by classic phenol/chloroform

DNA precipitation method using MAxtract high density tubes to maximize DNA recovery.

DNA fragments were then sequenced on an Illumina HiSeq 4000 apparatus, with paired-end 100 nt reads.

Due to technical issues, only one replicate could be used for dsgoth31 input.

ChIP-seq quality check: Validation of H3K4me3 enrichment around promoters.

Raw reads were trimmed using trim_galore (https://zenodo.org/record/5127899#.YbnMs73MLDc) with

default parameters along with --paired, --clip_R1 9, --clip_R2 9, and --max_n 0. Mapping was

performed using Bowtie2 (Langmead and Salzberg, 2012) with --sensitive-local against

the *D. melanogaster* r6.16 and *D. simulans* r2.02 genomes. Samtools was used to convert SAM to

coordinated sorted BAM files, while sambamba (Tarasov et al., 2015) was used to filter for uniquely

mapping reads and to remove duplicates (sambamba view -h -t 2 -f bam -F "[XS] == null and not

unmapped and not duplicate"). For *D. melanogaster* datasets, we filtered available blacklisted regions

(Amemiya et al., 2019) with bedtools. Finally coverage files containing reads per genome coverage

(RPGC) were obtained with DeepTools (Ramírez et al., 2016) bamCoverage with --extendReads, --

effectiveGenomeSize 129789873 for *D. melanogaster* available from the Deeptools suite, and --

effectiveGenomeSize 121102921 computed with unique-kmers.py from khmer (https://github.com/dib-

lab/khmer). Promoter regions were obtained with gencode_regions

(https://github.com/saketkc/gencode_regions) and along with coverage files were used in DeepTools

computeMatrix and plotProfile to build the average coverage of H3K4me3 and H3K9me3 around

transcription start sites in both species. The corresponding profiles looked as expected

(Supplemental_Fig_S9).

ChIP-seq analysis

For each of the immunoprecipitated samples (H3K4me3, H3K9me3, input), TE read counts were computed at the family level using the TEcount module of TEtools (Lerat et al., 2017) and the list of TE sequences available at ftp://pbil.univ-lyon1.fr/pub/datasets/Roy2019/.

ChIP-seq counts were normalized across samples of the same species using the counts(normalize=T) function of DESeq2 1.26.0 (Love et al., 2014). This was done independently for each of the immunoprecipitated samples (H3K4me3, H3K9me3, input). We then performed a log-transformation using the rlogTransformation function of DESeq2, and subsequently considered mean values across replicates. We only kept genes expressed in ovaries. We chose to work on log-transformed values because log-transformation of count variables makes them fit normal assumption and thus makes them suitable for linear models. In addition, a ratio becomes a difference when log-transformed, which ensure the strict equivalence with the classical normalization approach consisting in dividing histone counts with input counts: log([H3Kime3 counts] / [input counts]) = log(H3Kime3 counts) – log(input counts).

In order to quantify the effects of TE insertions on histone marks enrichment, we used the following linear models on log transformed read counts: histone mark (either H3K4me3 or H3K4me3) ~ input + exon + intron + upstream + downstream, where "exon", "intron", "upstream", and "downstream" are the numbers of TE insertions in exons, introns, 5 kb upstream sequences, and 5 kb downstream sequences, respectively. Size effects for each of these three factors were then recorded.

In parallel, for each strain, we defined 10 classes of genes based on histone mark enrichment, built upon the values of {log transformed histone mark read counts (either H3K4me3 or H3K9me3) – log transformed input read counts}, calculated for each gene. We computed the proportion of genes carrying at least one TE insertion in each class.

For the inter-genomic analysis, for each pair of strains and each ovary-expressed gene, we considered the model H3Kime3 (i stands for 4 or 9) ~ input + strain, and retrieved the absolute value of the effect of the strain factor. We defined 10 classes of genes based on the values of this variable (square root transformed), and computed the proportion of genes that carried different numbers of TE insertions within each class.

Data access

587 The RNA-seq data generated in this study have been submitted to the NCBI BioProject database

588 (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number PRJNA795668. The ChIP-seq

589 data generated in this study have been submitted to the NCBI BioProject database under accession

590 number PRJNA796157. TE and gene annotations have been deposited to Zenodo doi:

591 10.5281/zenodo.5844405.

Acknowledgments

We thank Francois Sabot, Matthieu Boulesteix, and Vincent Mérel for useful discussions and

technical help. We thank Gladys Mialdea, Justine Picarle, Sonia Janillon, and Nelly Burlet for

technical help. This work was performed using the computing facilities of the CC LBBE/PRABI.

Sequencing was performed by the GenomEast platform, a member of the "France Génomique"

consortium (ANR-10-INBS-0009). This work was supported by Fondation pour la Recherche

Médicale (grant DEP20131128536) and Agence Nationale de la Recherche (grant ExHyb ANR-14-

CE19-0016-01). The authors declare no competing interests.

602 References

Akkouche A, Grentzinger T, Fablet M, Armenise C, Burlet N, Braman V, Chambeyron S, Vieira C. 2013. Maternally deposited germline piRNAs silence the tirant retrotransposon in somatic cells. *EMBO Rep* **14**:458–464. doi:10.1038/embor.2013.38

Akkouche A, Rebollo R, Burlet N, Esnault C, Martinez S, Viginier B, Terzian C, Vieira C, Fablet M. 2012. tirant, a newly discovered active endogenous retrovirus in Drosophila simulans. *J Virol* **86**:3675–3681. doi:10.1128/JVI.07146-11

Amemiya HM, Kundaje A, Boyle AP. 2019. The ENCODE Blacklist: Identification of Problematic Regions of the Genome. *Sci Rep* **9**. doi:10.1038/s41598-019-45839-z

Baduel P, Leduque B, Ignace A, Gy I, Gil JJ, Loudet O, Colot V, Quadrana L. 2021. Genetic and environmental modulation of transposition shapes the evolutionary potential of Arabidopsis thaliana. *Genome Biol* **22**. doi:10.1186/s13059-021-02348-5

Bailly-Bechet M, Haudry A, Lerat E. 2014. "One code to find them all": a perl tool to conveniently parse RepeatMasker output files. *Mob DNA* **5**:13. doi:10.1186/1759-8753-5-13

Bieberstein NI, Kozáková E, Huranová M, Thakur PK, Krchňáková Z, Krausová M, Carrillo Oesterreich F, Staněk D. 2016. TALE-directed local modulation of H3K9 methylation shapes exon recognition. *Sci Rep* **6**:29961. doi:10.1038/srep29961

Biémont C, Vieira C. 2006. Genetics: junk DNA as an evolutionary force. *Nature* **443**:521–524. doi:10.1038/443521a

Chakraborty M, Chang C-H, Khost DE, Vedanayagam J, Adrion JR, Liao Y, Montooth KL, Meiklejohn CD, Larracuente AM, Emerson JJ. 2021. Evolution of genome structure in the Drosophila simulans species complex. *Genome Res* **31**:380–396. doi:10.1101/gr.263442.120

Cridland JM, Thornton KR, Long AD. 2015. Gene expression variation in Drosophila melanogaster due to rare transposable element insertion alleles of large effect. *Genetics* **199**:85–93. doi:10.1534/genetics.114.170837

Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, Hannon GJ. 2018. piRNA-Guided Genome Defense: From Biogenesis to Silencing. *Annu Rev Genet* **52**:131–157. doi:10.1146/annurev-genet-120417-031441

Everett LJ, Huang W, Zhou S, Carbone MA, Lyman RF, Arya GH, Geisz MS, Ma J, Morgante F, St Armour G, Turlapati L, Anholt RRH, Mackay TFC. 2020. Gene expression networks in the Drosophila Genetic Reference Panel. *Genome Res* **30**:485–496. doi:10.1101/gr.257592.119

Fablet M, Vieira C. 2011. Evolvability, epigenetics and transposable elements. *BioMol Concepts* **2**:333–341.

Frazee AC, Pertea G, Jaffe AE, Langmead B, Salzberg SL, Leek JT. 2015. Ballgown bridges the gap between transcriptome assembly and expression analysis. *Nat Biotechnol* **33**:243–246. doi:10.1038/nbt.3172

Hill MS, Vande Zande P, Wittkopp PJ. 2021. Molecular and evolutionary processes generating variation in gene expression. *Nat Rev Genet* **22**:203–215. doi:10.1038/s41576-020-00304-w

Hollister JD, Gaut BS. 2009. Epigenetic silencing of transposable elements: a trade-off between reduced transposition and deleterious effects on neighboring gene expression. *Genome Res* **19**:1419–1428. doi:10.1101/gr.091678.109

Horváth V, Merenciano M, González J. 2017. Revisiting the Relationship between Transposable Elements and the Eukaryotic Stress Response. *Trends Genet* **33**:832–841. doi:10.1016/j.tig.2017.08.007

Kent WJ. 2002. BLAT--the BLAST-like alignment tool. *Genome Res* **12**:656–664. doi:10.1101/gr.229202

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**:907–915. doi:10.1038/s41587-019-0201-4

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:357–359. doi:10.1038/nmeth.1923

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ. 2013. Software for computing and annotating genomic ranges. *PLoS Comput Biol* **9**:e1003118. doi:10.1371/journal.pcbi.1003118

Le Thomas A, Rogers AK, Webster A, Marinov GK, Liao SE, Perkins EM, Hur JK, Aravin AA, Tóth KF. 2013. Piwi induces piRNA-guided transcriptional silencing and establishment of a repressive chromatin state. *Genes Dev* **27**:390–399. doi:10.1101/gad.209841.112

Lee YCG. 2015. The Role of piRNA-Mediated Epigenetic Silencing in the Population Dynamics of Transposable Elements in Drosophila melanogaster. *PLoS Genet* **11**:e1005269. doi:10.1371/journal.pgen.1005269

Lee YCG, Karpen GH. 2017. Pervasive epigenetic effects of Drosophila euchromatic transposable elements impact their evolution. *eLife* **6**. doi:10.7554/eLife.25762

Lerat E, Fablet M, Modolo L, Lopez-Maestre H, Vieira C. 2017. TEtools facilitates big data expression analysis of transposable elements and reveals an antagonism between their activity and that of piRNA genes. *Nucleic Acids Res* **45**:e17. doi:10.1093/nar/gkw953

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**:550. doi:10.1186/s13059-014-0550-8

Malone CD, Hannon GJ. 2009. Small RNAs as guardians of the genome. *Cell* **136**:656–668. doi:10.1016/j.cell.2009.01.045

Mérel V, Boulesteix M, Fablet M, Vieira C. 2020. Transposable elements in Drosophila. *Mob DNA* **11**:23. doi:10.1186/s13100-020-00213-z

Mérel V, Gibert P, Buch I, Rodriguez Rada V, Estoup A, Gautier M, Fablet M, Boulesteix M, Vieira C. 2021. The Worldwide Invasion of Drosophila suzukii Is Accompanied by a Large Increase of Transposable Element Load and a Small Number of Putatively Adaptive Insertions. *Mol Biol Evol* **38**. doi:10.1093/molbev/msab155

Mohamed M, Dang NT-M, Ogyama Y, Burlet N, Mugat B, Boulesteix M, Mérel V, Veber P, Salces-Ortiz J, Severac D, Pélisson A, Vieira C, Sabot F, Fablet M, Chambeyron S. 2020. A Transposon Story: From TE Content to TE Dynamic Invasion of Drosophila Genomes Using the Single-Molecule Sequencing Technology from Oxford Nanopore. *Cells* **9**. doi:10.3390/cells9081776

Moschetti R, Palazzo A, Lorusso P, Viggiano L, Marsano RM. 2020. "What You Need, Baby, I Got It": Transposable Elements as Suppliers of Cis-Operating Sequences in Drosophila. *Biology* **9**:E25. doi:10.3390/biology9020025

Osada N, Miyagi R, Takahashi A. 2017. Cis- and Trans-regulatory Effects on Gene Expression in a Natural Population of Drosophila melanogaster. *Genetics* **206**:2139–2148. doi:10.1534/genetics.117.201459

Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD. 2019. PIWI-interacting RNAs: small RNAs with big functions. *Nat Rev Genet* **20**:89–108. doi:10.1038/s41576-018-0073-3

Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc* **11**:1650–1667. doi:10.1038/nprot.2016.095

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* **33**:290–295. doi:10.1038/nbt.3122

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**:W160–W165. doi:10.1093/nar/gkw257

Rebollo R, Horard B, Begeot F, Delattre M, Gilson E, Vieira C. 2012a. A snapshot of histone modifications within transposable elements in Drosophila wild type strains. *PloS One* **7**:e44253. doi:10.1371/journal.pone.0044253

Rebollo R, Karimi MM, Bilenky M, Gagnier L, Miceli-Royer K, Zhang Y, Goyal P, Keane TM, Jones S, Hirst M, Lorincz MC, Mager DL. 2011. Retrotransposon-induced heterochromatin spreading in the mouse revealed by insertional polymorphisms. *PLoS Genet* **7**:e1002301. doi:10.1371/journal.pgen.1002301

Rebollo R, Romanish MT, Mager DL. 2012b. Transposable elements: an abundant and natural source of regulatory sequences for host genes. *Annu Rev Genet* **46**:21–42. doi:10.1146/annurev-genet-110711-155621

Saint-André V, Batsché E, Rachez C, Muchardt C. 2011. Histone H3 lysine 9 trimethylation and HP1γ favor inclusion of alternative exons. *Nat Struct Mol Biol* **18**:337–344. doi:10.1038/nsmb.1995

Senti K-A, Brennecke J. 2010. The piRNA pathway: a fly's perspective on the guardian of the genome. *Trends Genet TIG* **26**:499–509. doi:10.1016/j.tig.2010.08.007

Shumate A, Salzberg SL. 2020. Liftoff: accurate mapping of gene annotations. *Bioinforma Oxf Engl* btaa1016. doi:10.1093/bioinformatics/btaa1016

Sultana T, Zamborlini A, Cristofari G, Lesage P. 2017. Integration site selection by retroviruses and transposable elements in eukaryotes. *Nat Rev Genet* **18**:292–308. doi:10.1038/nrg.2017.7

Sundaram V, Cheng Y, Ma Z, Li D, Xing X, Edge P, Snyder MP, Wang T. 2014. Widespread contribution of transposable elements to the innovation of gene regulatory networks. *Genome Res* **24**:1963–1976. doi:10.1101/gr.168872.113

Tarasov A, Villeva AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast processing of NGS alignment formats. *Bioinforma Oxf Engl* **31**. doi:10.1093/bioinformatics/btv098

Ullastres A, Merenciano M, González J. 2021. Regulatory regions in natural transposable element insertions drive interindividual differences in response to immune challenges in Drosophila. *Genome Biol* **22**:265. doi:10.1186/s13059-021-02471-3

Uzunović J, Josephs EB, Stinchcombe JR, Wright SI. 2019. Transposable Elements Are Important Contributors to Standing Variation in Gene Expression in Capsella Grandiflora. *Mol Biol Evol* **36**:1734–1745. doi:10.1093/molbev/msz098

Vieira C, Lepetit D, Dumont S, Biémont C. 1999. Wake up of transposable elements following Drosophila simulans worldwide colonization. *Mol Biol Evol* **16**:1251–1255.

Villanueva-Cañas JL, Horvath V, Aguilera L, González J. 2019. Diverse families of transposable elements affect the transcriptional regulation of stress-response genes in Drosophila melanogaster. *Nucleic Acids Res* **47**:6842–6857. doi:10.1093/nar/gkz490

Wells JN, Feschotte C. 2020. A Field Guide to Eukaryotic Transposable Elements. *Annu Rev Genet*. doi:10.1146/annurev-genet-040620-022145

Wittkopp PJ, Kalay G. 2011. Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nat Rev Genet* **13**:59–69. doi:10.1038/nrg3095

Zhang X, Zhao M, McCarty DR, Lisch D. 2020. Transposable elements employ distinct integration strategies with respect to transcriptional landscapes in eukaryotic genomes. *Nucleic Acids Res* **48**:6685–6698. doi:10.1093/nar/gkaa370