A quantitative, genome-wide analysis in *Drosophila* reveals transposable elements' influence on gene expression is species-specific

Marie Fablet[1,2], Judit Salces-Ortiz[1,5], Angelo Jacquet[1,6], Bianca F. Menezes[1,7], Corentin Dechaud[3], Philippe Veber[1], Rita Rebollo[4], Cristina Vieira[1]

[1] Laboratoire de Biométrie et Biologie Evolutive, Université de Lyon; Université Lyon 1; CNRS; UMR 5558, Villeurbanne, France.

[2] Institut Universitaire de France (IUF)

[3] Institut de Génomique Fonctionnelle de Lyon, Univ Lyon, CNRS UMR 5242, Ecole Normale Supérieure de Lyon, Université Claude Bernard Lyon 1, 46 allée d'Italie, F-69364 Lyon, France

[4] Univ Lyon, INRAE, INSA-Lyon, BF2I, UMR 203, 69621 Villeurbanne, France.

[5] Current address: Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra) P° Marítimo de la Barceloneta, 37-49 08003 Barcelona, Spain

[6] Current address: Symbiotron; FR3728 Biodiversité, Eau, Environnement, Ville, Santé; Université Claude Bernard Lyon 1; Villeurbanne 69622, France

[7] Current address: Federal Institute of Rio de Janeiro (IFRJ), Pinheiral – RJ, Brazil

Corresponding authors: marie.fablet@univ-lyon1.fr; cristina.vieira@univ-lyon1.fr

Abstract

Transposable elements (TEs) are parasite DNA sequences that are able to move and multiply along the chromosomes of all genomes. They are controlled by the host through the targeting of silencing epigenetic marks, which may affect the chromatin structure of neighboring sequences, including genes. In this study, we used transcriptomic and epigenomic high-throughput data produced from ovarian samples of several *Drosophila melanogaster* and *Drosophila simulans* wild-type strains, in order to finely quantify the influence of TE insertions on gene RNA levels and histone marks (H3K9me3 and H3K4me3). Our results reveal a stronger epigenetic effect of TEs on ortholog genes in *D. simulans* compared to *D. melanogaster*. At the same time, we uncover a larger contribution of TEs to gene H3K9me3 variance within genomes in *D. melanogaster*, which is evidenced by a stronger correlation of TE numbers around genes with the levels of this chromatin mark in *D. melanogaster*. Overall, this work contributes to the understanding of species-specific influence of TEs within genomes. It provides a new light on the considerable natural variability provided by TEs, which may be associated with contrasted adaptive and evolutionary potentials.

Introduction

Transposable elements (TEs) are parasite DNA sequences that are able to move and multiply along the chromosomes of all genomes (Wells and Feschotte, 2020). They are source of mutations and genome instability if uncontrolled (Biémont and Vieira, 2006; Malone and Hannon, 2009; Senti and Brennecke, 2010). Control of TEs generally consists in the targeting of particular chromatin marks to TE copies, which induce transcriptional and post-transcriptional gene silencing and may spread to neighboring sequences and impact gene expression. In this regard, few attempts were made to finely analyze and quantify TEs' influence at the whole genome scale (Cridland et al., 2015; Hollister and Gaut, 2009; Huang et al., 2016; Lee and Karpen, 2017; Uzunović et al., 2019; Wei et al., 2022). In addition, since the very beginning of TE studies, species-specific differences in TE contents, activities and control pathways have been reported in nature, and particularly between *D. melanogaster* and *D. simulans* (Akkouche et al., 2013, 2012; Fablet et al., 2014; Kofler et al., 2015b; Lee and Karpen, 2017; Mérel et al., 2020; Vieira et al., 2012, 1999). Previous research described the effects of TE insertions on gene expression using collections of strains of *D. melanogaster* (Cridland et al., 2015; Everett et al., 2020; Osada et al., 2017; Zhang et al., 2020), and other studies focusing on a few TE families in wild-type strains of *D. simulans* and *D. melanogaster* uncovered between-species differences in histone mark landscapes (Rebollo et al., 2012a). Lee and Karpen (Lee and Karpen, 2017) provided an analysis on the repressive histone mark H3K9me2 (Histone 3 Lysine 9 dimethylation) around TEs from two *Drosophila* Genetic Reference Panel (DGRP) strains (*D. melanogaster*), and concluded to pervasive epigenetic effects of TEs. However, rather than H3K9me2, it is H3K9me3 (Histone 3 Lysine 9 trimethylation) that is known to be associated with the

activity of dual-stranded piRNA clusters and the production of TE-derived silencing piRNAs (Le Thomas et al., 2013; Mohn et al., 2014; Sienski et al., 2012). H3K9me3 differs from H3K9me2 in that it is more strongly bound by Rhino, which is abundant in ovaries and leads to piRNA production through alteration of the local transcription program (Mohn et al., 2014).

Several limitations remained from the previous studies, which we propose to address in the present work. First, we connect TE insertion polymorphism, RNA-seq, ChIP-seq on two histone marks, and small RNA-seq data on the same strains. We use eight previously characterized, wild-type strains of *D. melanogaster* and *D. simulans* (Mohamed et al., 2020) that are derived from samples collected in France and Brazil, two strains per location and per species. Using the Oxford Nanopore long read sequencing technology, we previously produced high quality genome assemblies at the chromosome resolution for each strain, which provides us with the various TE insertion sites in each genome (Mohamed et al., 2020). Second, all data are produced from ovaries, *i.e.* the exact same tissue and not mix of tissues. As previously stated, Rhino is known to bind to H3K9me3 and promote the non-canonical transcription of dual-stranded piRNA clusters, in ovaries only (Mohn et al., 2014). Therefore, we expect the strongest control of TEs in this tissue and thus potentially the strongest impact on neighboring genes. In particular, we can speculate that genes located nearby TE insertions may be affected by the local production of piRNAs and hence we searched for gene-derived piRNAs, in association with increased levels of H3K9me3 deposition on gene sequences. We also studied H3K4me3 (Histone 3 Lysine 4 trimethylation), which is known to be associated with active, canonical transcription. Third, the production of genome-wide data from four wild-type strains of *D. melanogaster* and four wild-type strains of *D. simulans* brings the opportunity to statistically test for species-specific differences and provide a quantitative assessment of the contribution of TEs to gene expression, in a comparative genomics perspective (Fig. 1). In addition, the use of linear models allows to finely quantify and compare the contributions at different levels.

The original approach and subsequent analyses reveal a stronger epigenetic influence of TEs on orthologous genes in *D. simulans* compared to *D. melanogaster*, and are in agreement with the recent work published by Lee's lab (Huang et al., 2022). At the same time, we uncover a larger contribution of TEs to genome architecture in *D. melanogaster*: in particular, TE insertions contribute more to gene H3K9me3 level variance in *D. melanogaster* compared to *D. simulans*, which is evidenced by a stronger association of TEs around genes with the levels of this chromatin mark in *D. melanogaster*. Overall, this work contributes to the understanding of species-specific influence of TEs within genomes. As a whole, these results participate in the accurate, quantitative understanding of TEs' impacts on genomes, and highlight the species-specific differences in the interaction between TEs and the host genome. This provides a new light on the considerable natural variability resulting from TEs, which may be associated with contrasted adaptive and evolutionary potentials, all the more sensible in a rapidly changing environment (Baduel et al., 2021; Fablet and Vieira, 2011; Mérel et al., 2021).

Figure 1. Graphic summary of the study.

Eight wild-type strains from *D. melanogaster* and *D. simulans* were included in the study. The present datasets are RNA-seq and ChIP-seq for H3K4me3 and H3K9me3 marks, and were prepared from ovarian samples. They were analyzed in parallel with already published data produced from the same *Drosophila* strains: ovarian small RNA repertoires and genome assemblies based on Oxford Nanopore long read sequencing (Mohamed et al., 2020). For RNA-seq and ChIP-seq, TE-derived reads were analyzed at the TE family level, and gene-derived reads were analyzed in relation to TE insertions inside or near genes (therefore restricted to the TE insertions included within the gray bubbles).

Results

TE expression and epigenetic targeting in *Drosophila* ovaries

We first considered TE-derived RNA-seq reads from all samples, which we analyzed at the TE-family level (Fig. 1). As performed by other research studies (Chakraborty et al., 2021; Kofler et al., 2015b), we removed the non-autonomous *DNAREP1* helentron (also known as *INE-1*) from our analyses because it is a highly abundant element displaying mainly fixed insertions in the *melanogaster* complex of species (Thomas et al., 2014). However, a recent study revealed an expansion of this family in the *Drosophila nasuta* species group (Wei et al., 2022), indicating its activity and potential genomic impacts. We therefore performed a *DNAREP1*-dedicated analysis, apart from the other families. TEs account for 0.6% to 1.2%, and 0.5% to 0.7%, of read counts corresponding to annotated sequences (genes and TEs) within the ovarian transcriptomes of *D. melanogaster* and *D. simulans* strains, respectively (Fig. 2A), and *DNAREP1* accounts for 6% to 13%, and for 5% to 9% of the total number of TE read counts in *D. melanogaster* and *D. simulans*, respectively. This contribution is very weak with regard to the ~4,000 copies of *DNAREP1* identified by our procedure within each genome. We removed *DNAREP1* and found significant positive correlations between per TE family RNA counts and family sequence occupancy (quantified as the total number of bp spanned by

each TE family along the genome) (Spearman correlations, rho = 0.33 to 0.37, and 0.39 to 0.44, in *D. melanogaster* and *D. simulans*, respectively; Supplemental Fig. S2A). Regarding TE-derived piRNA production, it was previously described in control conditions in wild-type strains that the amounts of piRNAs were positively correlated with the amounts of RNAs, at the TE family level (Lerat et al., 2017). This remains true in the present dataset: we find significant positive correlations between per TE family RNA counts and piRNA counts (Spearman correlations, rho = 0.39 to 0.48, and 0.48 to 0.56, in *D. melanogaster* and *D. simulans*, respectively; Supplemental Fig. S2B). In both cases, correlations are significantly stronger in *D. simulans*, compared to *D. melanogaster* (Wilcoxon rank tests for *D. melanogaster vs D. simulans* comparisons; correlation coefficients between TE RNA counts and TE sequence occupancy: p-value = 0.029; correlation coefficients between TE RNA counts and TE piRNA counts: p-value = 0.029), suggesting a more efficient production of TE-derived piRNAs.

We assessed the contribution of histone mark enrichment to TE RNA amounts considering the following linear model on log-transformed normalized read counts: RNA ~ H3K4me3 + H3K9me3 + input. These models led to adjusted r^2 as high as 0.48 to 0.64 depending on the strains in *D. melanogaster* and 0.45 to 0.60 in *D. simulans*, suggesting that these models capture significant portions of TE RNA amount variation. We find that TE RNA amounts are positively correlated with H3K4me3 and negatively correlated with H3K9me3 amounts (Fig. 2B), as expected considering that H3K4me3 is an activating mark while H3K9me3 is a silencing one. We used a similar approach to analyze piRNA amounts, and considered the following linear model on log-transformed read counts: piRNA ~ H3K4me3 + H3K9me3 + input. We obtained even higher adjusted r^2 values, from 0.70 to 0.75, and 0.64 to 0.68, depending on the strains in *D. melanogaster* and *D. simulans*, respectively. We find that TE-derived piRNA amounts are positively correlated both with permissive H3K4me3 and repressive H3K9me9 levels (Fig. 2C). The tighter correlations may be due to the strong dependency of piRNA production mechanisms on chromatin marks and H3K9me3 in particular, while RNA transcription also involves other factors, such as transcription factors, which binding sites vary a lot across TE sequences.

TE insertions within or nearby genes

In the following sections, we focus on gene-derived reads from all samples, which we analyzed with regard to the presence of TE insertions within or nearby genes (Fig. 1). Based on gene annotations, we distinguished the different functional regions of genes: exons, introns, upstream, or downstream sequences (5 kb flanking regions). Exons are both UnTranslated Regions (UTRs) and Coding Sequences (CDSs). Sequences that may both behave as exons or introns depending on alternative splicing are included in "exons". In this first step, we considered a set of 17,417 annotated genes for *D. melanogaster*, and 15,251 for *D. simulans* (see Material and Methods). We quantified the number of TE insertions within genes (Fig. 2D), and found that

they account for ~16% and ~25% of the total number of TE insertions per genome in *D. melanogaster* and *D. simulans*, respectively. The majority of genes (93%) are devoid of TE insertions within gene bodies, and very few display more than one TE insertion (Fig. 2E, Supplemental Fig. S1). Among the copies of *DNAREP1* that we identified along the genomes, our analysis revealed that 1,343 to 1,374 insertions from this family are found within genes in *D. melanogaster*, and 1,075 to 1,089 insertions in *D. simulans* (Supplemental Fig. S3).

Figure 2. (*A*) Proportions of TE read counts in RNA-seq data relative to read counts corresponding to genes and TEs. For each strain, two biological replicates are shown. (*B*) Contributions of H3K4me3 and H3K9me3 enrichment to TE-derived RNA read counts (according to the model RNA ~ H3K4me3 + H3K9me3 + input calculated on log10 transformed read count numbers, at the TE family level). Colored bars: p-values < 0.05, empty bars: p-values > 0.05. Error bars are standard errors. (*C*) Contributions of H3K4me3 and H3K9me3 enrichment to TE-derived piRNA read counts (according to the model piRNA ~ H3K4me3 + H3K9me3 + input calculated on log10 transformed read count numbers, at the TE family level). Colored bars: p-values < 0.05, empty bars: p-values > 0.05. Error bars are standard errors. (*D*) Number of TE insertions per functional region per strain. Upstream and downstream regions are 5 kb sequences directly flanking transcription units 5' and 3', respectively. (*E*) Number of genes for each value of TE insertion numbers. dmgoth101 and dsgoth31 are shown as examples; all strains can be found in Supplemental Fig. S1.

TEs contribute to RNA level variability across genes within genomes

One of the novelties of the present work is to quantify the contribution of TE insertions to the variance in gene expression levels within distinct genomes. We focused on the subset of genes that we found expressed in the ovaries (see Material and Methods), *i.e.* 7,883 to 8,135 genes depending on the strains of *D. melanogaster*, and 7,653 to 8,121 genes in *D. simulans*. We quantified TE insertion contribution to gene RNA levels using the following linear models built on log-transformed TPM (Transcript Per Million): TPM ~ exon + intron + upstream + downstream, where these variables correspond to the number of TE insertions within exons, introns, 5 kb upstream, and 5 kb downstream regions, respectively. We find that TE insertions contribute significantly, albeit weakly, to gene expression variance (Fig. 3A): 1.6% to 1.9% of total variance in *D. melanogaster*; 1.2% to 1.9% in *D. simulans*. These values may look low at first sight; however, gene expression levels are known to be primarily regulated by many other factors, such as transcription factor binding, sequence composition and polymorphism, etc. This reveals that our approach is powerful enough to capture low levels of variation and that TEs are significant actors of this variability. Although total contribution to gene expression variance does not differ between species (Wilcoxon rank test, p-value = 0.685), we found significant differences when considering specific gene regions. For instance, the contribution of TE insertions within introns was higher in *D. simulans* compared to *D. melanogaster* (mean values: 0.03% *vs* 0.14%; Wilcoxon rank test, p-value = 0.029), while the contribution of TE insertions downstream genes was higher in *D. melanogaster* compared to *D. simulans* (mean values: 0.06% *vs* 0.21%; Wilcoxon rank test, p-value = 0.029).

When we computed the corresponding size effects, we observed significant, negative associations between gene expression levels and TE insertions within exons and introns, and significant, positive associations for

TE insertions around genes (Fig. 3B). The association with gene expression was stronger for
D. melanogaster compared to *D. simulans* for downstream TE insertions (Fold-change = 1.6; Wilcoxon rank
test, p-value = 0.029), and it was stronger in *D. simulans* compared to *D. melanogaster* for TE insertions
within introns (Fold-change = 6.2; Wilcoxon rank test, p-value = 0.029) and upstream TE insertions (Fold-
change = 1.9; Wilcoxon rank test, p-value = 0.029).

Nevertheless, one could argue that the species-specific differences that we observe here are due to gene sets
not being exactly the same across species. In order to correct for this bias, we focused on the subset of 6,417
genes that have 1:1 ortholog in the other species and that are expressed in ovaries. The results were very
similar regarding size effects, reinforcing our conclusions (Supplemental Fig. S4). However, we noticed that
TE contribution to gene expression variance was increased in this subset of genes: 3.2% and 2.9% on
average in *D. melanogaster* and *D. simulans*, respectively (Supplemental Fig. S4).

Collectively, our data show a weak but significant contribution of TEs to the variance in gene expression
within genomes, which varies across species and is due to negative correlations between gene RNA levels
and TE numbers in exons and introns, and positive correlations with TE numbers upstream and downstream
genes.

RNA levels

H3K4me3 levels

H3K9me3 levels

Figure 3. (*A*) Contribution of TE insertion numbers to gene expression total variance estimated using the linear model gene TPM (log) ~ exon + intron + upstream + downstream, and (*B*) corresponding size effects. (*C*) Contribution of TE insertion numbers to gene H3K4me3 total variance estimated using the linear model gene H3K4me3 level (log) ~ exon + intron + upstream + downstream, and (*D*) corresponding size effects.

214 (*E*) Contribution of TE insertion numbers to gene H3K9me3 total variance estimated using the linear model

215 gene H3K9me3 level (log) ~ exon + intron + upstream + downstream, and (*F*) corresponding size effects.

216 Significance indications above graphs in *(B, D, E)* are *D. melanogaster vs D. simulans* comparisons using

217 Wilcoxon rank tests. Colored bars: p-values < 0.05, empty bars: p-values > 0.05. Error bars are standard

218 errors.

TEs contribute to histone enrichment variability across genes within genomes

We used a similar approach to analyze H3K4me3 and H3K9me3 enrichment (*i.e.* we aligned ChIP-seq reads

against whole gene sequences and computed corresponding read counts). We found that TE insertions

contributed significantly (except in dsgoth613), albeit very weakly, to gene H3K4me3 levels variance

(0.07% to 0.10% total variance in *D. melanogaster*; 0.04% to 0.09% in *D. simulans*; Wilcoxon rank test for

D. melanogaster vs D. simulans comparison, p-value = 0.200) (Fig. 3C). When computing size effects, the

only significant and consistent result is a negative association of TE insertions within exons with gene

H3K4me3 levels, in *D. melanogaster* only (Fig. 3D).

The contribution of TE insertions to total variance is higher for H3K9me3 levels: 0.29% to 0.65% in

D. melanogaster, and 0.07% to 0.14% in *D. simulans* (Fig. 3E; Wilcoxon rank test for *D. melanogaster vs*

D. simulans comparison, p-value = 0.029). The largest contribution comes from TE insertions around genes

and within introns, while TE insertions within exons virtually do not contribute to H3K9me3 variance. The

computation of size effects reveals a consistent, positive association of TE insertions within introns,

upstream and downstream genes with H3K9me3 levels, in both species. These results are in agreement with

TEs being the preferential targets for H3K9me3 deposition, which then spreads to neighboring regions (Le

Thomas et al., 2013; Rebollo et al., 2011). Alternatively, we cannot exclude that they may also lie in

particular chromatin environments where there is retention bias (Sultana et al., 2017), and that the

associations detected here are due to these particular chromatin features. The effects are stronger in

D. melanogaster compared to *D. simulans* for TE insertions around genes (Fig. 3F; Upstream: fold-change =

1.8, Wilcoxon rank test, p-value = 0.029; Downstream: fold-change = 1.7, Wilcoxon rank test, p-value =

0.029).

When considering only the set of 1:1 orthologous genes, patterns are highly similar for size effects, except

that the association between TE insertions within introns and H3K9me3 levels is now significantly stronger

in *D. melanogaster* compared to *D. simulans.* In addition, the contribution to H3K4me3 total variance is

higher for this subset of genes compared to the total set, although it remains very low, up to 0.73% in

D. melanogaster and 0.37% in *D. simulans*. (Supplemental Fig. S4).

While the observation of concomitant negative correlations with RNA levels and positive correlations with H3K9me3 for TE insertions within introns is in agreement with a negative impact of a heterochromatic mark on gene expression, the results for TE insertions around genes appears a little bit at odds. Indeed, TE insertions upstream and downstream of genes are at the same time positively correlated with RNA levels and H3K9me3 enrichment. One hypothesis for these TE insertions could be that their positive association with RNA levels is due to the multiple transcription factor binding sites that they bring —some transcription factors such as CTCF are known to be insensitive to chromatin (Isbel et al., 2022)—, and this ends up counteracting the negative impact of H3K9me3 targeting.

Patterns are globally conserved across TE classes and ages

We next analyzed TE insertions according to TE class, *i.e.* LTR (Long Terminal Repeat) elements, LINEs (Long Interspersed Nuclear Elements), DNA transposons, and *DNAREP1*. We used the same linear models on the same sets of genes, but considering only TE insertions belonging to each particular class. TE insertion numbers vary across classes (Supplemental Fig. S3), which leads to differences in statistical power (the higher power associated with the higher number of TE insertions). Despite this, the computation of size effects on gene RNA levels, H3K4me3, and H3K9me3 levels revealed highly consistent patterns across TE classes (Supplemental Fig. S3). *DNAREP1* patterns are similar to other DNA transposons. The major difference with global patterns (Fig. 3) is a trend for a positive association of DNA transposons and *DNAREP1* insertions in exons with gene expression in *D. melanogaster* only. Differences between transposons (DNA transposons and *DNAREP1*) and retrotransposons (LTR elements and LINEs) might be related to different waves of transposition: Kofler *et al.* described that LTR insertions are mostly of recent origin in both species, while DNA and non-LTR insertions are older, and that DNA transposons showed higher activity levels in *D. simulans* (Kofler et al., 2015b). The positive association between TE insertions in exons and gene expression would be characteristics of the families with the most ancient transposition activity, and potentially domestication events.

Irrespective of TE classes, it has already been described that TEs' impacts on genes differ across young (*i.e.* polymorphic) and old (*i.e.* fixed) TE copies; this is due to the pool of old TE insertions having been purged from deleterious insertions by natural selection (Hollister and Gaut, 2009). Indeed, Uzunovic *et al.* (Uzunović et al., 2019) showed in the plant *Capsella* that young TE insertions had a negative effect on gene expression while old insertions were more likely to increase gene expression. In this view, we distinguished insertions that are unique to one genome ("private") —and therefore correspond to the most recent insertions —, and those that are shared by all four strains of the species ("common") —thus the oldest ones. The majority of the TE insertions that are considered here (71% to 78%) fall in the "common" category. This may seem at odds regarding previous knowledge and the work of Kofler *et al.* in particular, who found that >80%

TE insertions had low frequency in pool seq data (Kofler et al., 2015b). However, the majority of these insertions are intergenic while we only focus on TEs within or around genes in the present study, which explains the differences in proportions between the two studies. The difference in subset sizes between "common" and "private" categories also leads to a reduced statistical power for the set of private insertions. Despite this difference, the observed patterns are rather consistent between both sets of TEs, and very similar to the global patterns including all TEs regardless of insertion polymorphism (Fig. 3, Supplemental Fig. S5). In the "common" pool, we do not observe the positive association between TE insertions in exons and gene expression reported by (Uzunović et al., 2019), maybe because the majority of these insertions are not old enough, or at least not as old as the above-described DNA transposon pool in *D. melanogaster*. Since our approach is gene-centered (Fig. 1), it is very likely that our complete set of TE insertions is already biased: when deleterious, insertions within or near genes have such a negative impact that we are not able to catch them from natural samples. Therefore, our complete set of TE insertions may already correspond to copies that have passed the filter of natural selection, and thus does not show critical differences between "common" and "private" patterns. However, some species-specific difference appears in the private set of insertions within introns: they display stronger negative association with gene expression levels in *D. simulans* only, and stronger positive association with H3K9me3 levels in *D. melanogaster* only. We speculate that this reveals species-specific differences in the efficiency of TE control at the first stages of TE invasion.

Gene-derived small RNAs and epigenetic effects

It has been demonstrated that TEs are sources of piRNA biogenesis in the ovary through the action of Rhino that promotes non-canonical transcription (Mohn et al., 2014). We took advantage of our extensive dataset made of RNA-seq, ChIP-seq and small RNA-seq produced from the ovaries of the exact same strains to test for the impact of piRNA cluster activity on neighbouring genes. In addition, siRNAs were previously shown to be produced from piRNA clusters and participated in TE silencing in ovaries (Shpiz et al., 2014). Therefore, we searched for gene-derived piRNAs and siRNAs, which could result from the spreading of small RNA production machinery from TE insertions. We filtered small RNAs based on read length, which does not allow us to distinguish siRNAs from miRNAs in the pool of 21 nt reads. We will therefore refer to them as "21 nt RNAs". In agreement with this scenario, we found a significant positive correlation between gene-derived piRNAs and gene-derived 21 nt RNAs (Spearman correlation coefficients; *D. melanogaster*: 0.517 to 0.536; *D. simulans*: 0.526 to 0.661; all p-values < 1e-10). In addition, we found that gene-derived piRNA production was significantly positively correlated with gene H3K9me3 levels (Supplemental Fig. S6), as expected in case of spreading of the piRNA cluster transcription to nearby gene sequences (Spearman correlation coefficients; *D. melanogaster*: 0.561 to 0.586; *D. simulans*: 0.475 to 0.525; all p-values < 1e-10). Remarkably, correlations were stronger for *D. melanogaster* compared to *D. simulans* (Wilcoxon rank test,

p-value = 0.029). Gene-derived 21 nt RNA production was also significantly positively correlated with gene
H3K9me3 (Spearman correlation coefficients; *D. melanogaster*: 0.470 to 0.517; *D. simulans*: 0.437 to
0.504; all p-values < 1e-10) but the strength of the correlation was not significantly different between
species.

In addition, our expectation is that the epigenetic spreading from piRNA clusters should be stronger for more
recent TE insertions, which are expected to be potentially more harmful because recently active. Therefore,
in order to focus on these recent TE insertions, we studied genes which polymorphic TE insertions were only
"private". We found that piRNA production from these genes were more frequently higher than the third
quartile than expected (except in dsgoth613) (Supplemental Table S1). These results demonstrate that the
control of TE sequences by the piRNA pathway impacts neighboring genes through the production of gene-
derived small RNAs and the increased deposition of H3K9me3 marks.

TE insertions are associated with variability in expression and histone enrichment between ortholog genes

We used our experimental dataset to infer the contribution of TE insertions at the inter-genomic level, *i.e.* we
compared expression levels of the same genes across genomes. We first considered *D. melanogaster* and
D. simulans separately. For each gene that displays variation in TE insertion numbers across strains, we
computed the mean difference of gene expression (TPM, scaled by gene average) between the strain that had
the highest TE insertion numbers and the strain that had the lowest. When several strains had the same
numbers of TE insertions, we computed their average gene expression level. We performed the same
approach on histone enrichment. Our assumption was that a general effect of TE insertions would shift the
distribution of the mean difference away from 0. This is not what we observed for RNA levels nor for
H3K4me3 enrichment (0 departure t tests, all p-values > 0.05) (Fig. 4). However, we find an increase in
H3K9me3 enrichment associated with high TE insertion numbers, but only in *D. simulans* and for TE
insertions within introns and upstream genes (0 departure t test; within introns: mean difference = 0.003, p-
value = 0.0005; upstream: mean difference = 0.003, p-value = 0.0019). These results are congruent with
recent studies, which observed a clear association between TE insertions and heterochromatin but no
predominant negative impact on the expression of neighboring genes (Huang et al., 2022; Wei et al., 2022).

We also took the opportunity to consider 1:1 ortholog genes so as to include all eight strains
(*D. melanogaster* and *D. simulans*) in the same analysis. Computation strategies were the same as above and
revealed significant decreases in RNA levels for strains with the highest TE insertion numbers in exons
(mean difference = -0.129, p-value = 1e-10) and introns (mean difference = -0.077, p-value = 9e-5). We also
found significant increase in H3K4me3 levels as well as H3K9me3 levels for strains with the highest TE

insertion numbers in exons and introns (H3K4me3, TEs within exons: mean difference = 0.012, p-value =
0.0201; within introns: mean difference = 0.019, p-value = 1e-5; H3K9me3, TEs within exons: mean
difference = 0.037, p-value = 0.0092; within introns: mean difference = 0.028, p-value = 2e-5). However,
such an analysis including all strains from both species at once has to be considered with caution because
gene sequences differ across species (GC content, length, etc.), which may interfere with mapping and read
counting, and was not accounted for in this work.

Figure 4. Variability in gene expression and histone enrichment according to TE insertion numbers across strains. (*A*) Mean expression difference (in TPM, scaled by gene average) between strains with the highest and the lowest TE insertion numbers for each region of each gene; mean histone enrichment difference (log-transformed, scaled by gene average) between strains with the highest and the lowest TE insertion numbers. Analyses are performed separately for both species (blue: *D. melanogaster*, orange: *D. simulans*), only considering genes that show different TE insertion numbers across strains. Significance levels correspond to t tests comparing observed mean to 0. (*B*) Same analyses across all eight strains considering 1:1 ortholog genes. Significance levels correspond to t tests comparing observed mean to 0: p-value 0 *** 0.001 ** 0.01 * 0.05.

Discussion

The common-held view is that, as parasites that are fought against by the genomes, TEs have a general negative impact on gene expression (Cridland et al., 2015; Lee, 2015; Lee and Karpen, 2017). Our present findings are in agreement with this idea. However, the originality of this research work is to provide an unprecedented quantitative view, which allows to precisely decipher TE impacts, integrating data gathered from wild-type strains of two closely related *Drosophila* species. This study combines genomic, transcriptomic, and epigenetic high-throughput sequence data, all produced from ovaries, where TEs are tightly controlled by epigenetic mechanisms through the piRNA pathway (Malone and Hannon, 2009; Senti and Brennecke, 2010) and therefore where we are to expect the strongest impacts of TEs on genes.

Expression and epigenetic marks of TE sequences

Our results uncover a lower contribution of TEs to the *D. simulans* transcriptome as compared to *D. melanogaster* (0.6% vs 1.1% on average, Fig. 2A). This is in agreement with the previously described lowest contribution of TEs in the genomes of *D. simulans* in terms of sequence occupancy and copy numbers (Mohamed et al., 2020; Vieira et al., 1999). In both species, we found that H3K9me3 marks on TE sequences are associated with a decrease in TE-derived RNA amounts, and the opposite for H3K4me3 marks. On the contrary, we observed that both histone marks are positively correlated with TE-derived piRNA amounts, which is congruent with the piRNA-targeted deposition of H3K9me3 marks at transcriptionally active TE copies (Czech et al., 2018; Sienski et al., 2012). However, one should note that these results reflect average behaviors at the TE family level, and TE copies may differ from one another within TE families.

What emerges from the different analyses that we performed is a remarkable variability across TEs, as illustrated by the width of dot distributions in Fig. 4 for instance. This highlights the huge variability across

TE sequences on many aspects: class, family, length, insertion site preference, chromosome distribution, activity, transposition rate, etc. For instance, in their pool-seq analysis of *D. melanogaster* and *D. simulans,* Kofler *et al.* found that half of the TE families showed evidence of variation of activity through time and were not the same depending on the species (Kofler et al., 2015b). It is congruent with the conclusions of Wei *et al.*, working on the *Drosophila nasuta* complex of species, who emphasize that TE insertions can have multiple effects on gene expression, from no effect to silencing or over-expression (Wei et al., 2022). This also echoes the work of Malone *et al.* and Sienski *et al.,* who described different groups of TEs depending on their sensitivity to different piRNA pathways and thus different effects on neighboring genes (Malone et al., 2009; Sienski et al., 2012). In addition, it has already been suggested and demonstrated that TEs' influence on gene expression is only manifested in case of stress (Naito et al., 2009), which adds another layer of variability and difficulty to disentangle biological impacts.

Intra- and inter-genomic analyses tell distinct, although complementary stories

In the intra-genomic analysis, we gather all expressed genes from a given genome, which we compare for their TE insertions, expression level, chromatin marks, and piRNA production. These are therefore heterogeneous sets of genes, which work coordinately in living cells. In the inter-genomic analysis, we compare the same ortholog genes in different genomes. We assume that these genes differ mainly based on their TE insertions.

When TE insertions are associated with differences in gene expression or chromatin state, it is very difficult to tell apart whether these TE insertions are causative or not. Nevertheless, the inter-genomic analysis is a way to demonstrate causality because it compares versions of the same genes but displaying different numbers of TE insertions —however with the limitation of neglecting nucleotide polymorphism. This approach has already successfully been followed by others and led to the conclusion of the causative role of the TE insertions (Lee and Karpen, 2017; Rebollo et al., 2011). On the contrary, in the intra-genomic study, we draw general patterns from the analysis of the complete set of genes at once, which differ from TE insertion numbers but also from many other aspects (sequence, length, expression level, tissue-specificity, local recombination rate, etc.). The intra-genomic analysis allows to identify associations between TE insertions, gene expression and chromatin environment, and therefore brings us to draw species-specific gene landscapes.

Here, the inter-genomic analysis on the complete dataset (orthologous genes from both species, Fig. 4B) reveals that TE insertions within, but not around genes, have a negative impact on gene RNA levels, and a positive impact on both histone marks, H3K4me3 and H3K9me3. This H3K4me3 result may be related to TEs donating promoters or *cis* regulatory sequences, as was already described on several instances

(Moschetti et al., 2020; Sundaram et al., 2014; Villanueva-Cañas et al., 2019) or disrupting inhibitory
sequences. The impact on H3K9me3, however, appears to be stronger since the net result is negative on gene
RNA levels. This result corresponds to TEs being a preferential target for H3K9me3 deposition (Le Thomas
et al., 2013), which then spreads to neighboring sequences.

In addition, the inter-genomic analysis reveals stronger epigenetic impacts of TE insertions in *D. simulans*
compared to *D. melanogaster* (Fig. 4A). These results support the previous findings from Lee & Karpen,
which found higher enrichment and spread of H3K9me2 from TE insertions in *D. simulans* compared to
D. melanogaster (Lee and Karpen, 2017). These results were recently confirmed in a larger set of species
(Huang et al., 2022). They proposed that this leads to stronger selection against TE insertions close to genes
in *D. simulans* compared to *D. melanogaster*, which explains the lower total number of TE insertions and
the lower proportion of TE insertions within or nearby genes in *D. simulans*.

On the other hand, the intra-genomic analysis confirms the already described trend of TE insertions within
genes to be associated with a reduction in gene RNA levels. However, our results also reveal that TE
insertions around genes are associated with increased gene expression on average. Overall, TE insertions are
virtually not associated with particular H3K4me3 patterns, except for TE insertions in exons in
D. melanogaster, which are associated with a decrease in H3K4me3. As previously known and confirmed by
the inter-genomic analysis, TE insertions are associated with increased levels of H3K9me3. The novelty
brought by the intra-genomic analysis is that the association is particularly strong for TE insertions around
genes and not within genes, particularly in *D. melanogaster* compared to *D. simulans*. *D. melanogaster* TEs
contribute more to gene H3K9me3 level variance compared to *D. simulans*. This suggests that there is a
stronger structuration or stratification of genes according to TE insertion numbers and histone marks in this
species compared to *D. simulans*. TE insertions are more frequently found with higher H3K9me3 (and even
H3K4me3 to a lesser extent) enrichment in *D. melanogaster*.

Interpretations from inter- and intra-genomic analyses seem contradictory at first sight. However, they may
illustrate the two facets of RNA interference, *i.e.* defense *vs* regulation (Torri et al., 2022). We may speculate
that in *D. simulans*, the defense facet appears prominent while the regulation prevails in *D. melanogaster*.
Such differences in closely related species are not unexpected in the piRNA pathway, which is known to be
evolving at a particularly elevated rate (Fablet et al., 2014; Obbard et al., 2009). Again, we may speculate
that this is related —whether as a cause or a consequence cannot be told— to the different tempo of TE
activity and genome colonization between both species.

In the intra-genomic analysis, many parameters other than the numbers of TE insertions differ across the
genes (the family and length of the TEs, gene sequence composition, presence of transcription factor binding
sites, etc. (Hill et al., 2021; Wittkopp and Kalay, 2011)) and yet we were able to capture statistical signal
from the numbers of TE insertions. This suggests a widespread influence of TEs on gene expression. The
underlying mechanisms may be chromatin mark spreading, but not only. TEs may also disrupt functional

elements, especially for those inside genes, or add transcription factor binding sites ((Horváth et al., 2017;

Rebollo et al., 2012b; Ullastres et al., 2021)). Moreover, we have to note that TE insertions may

accumulate in specific chromatin environments due to insertional preference or different levels of selection

in these environments (Sultana et al., 2017).

TEs' influence on genomes is different between *D. melanogaster* and *D. simulans*

The intra- and inter-genomic analyses performed here both reveal species-specific differences, however not

at the same scale (Fig. 5). The intra-genomic analysis uncovers stronger epigenetic effects in

D. melanogaster compared to *D. simulans* and a positive association between gene expression and TE

insertions located in the flanking regions (Fig. 3). It means that genes that have many TEs in

D. melanogaster on average have higher H3K9me3 levels than genes that have many TEs in *D. simulans*.

This may be due to differences in TE insertion landscapes or to differential retention in particular chromatin

regions. This analysis therefore reveals how TE sequences may participate in the structure of the genome and

how this differs between species. This reflects more long-term and intimate interactions between the host

genome and its TEs.

The species-specific differences that we observe for TE influence on genes may be due to variability in the

efficiency of epigenetic machinery, as suggested by (Lee and Karpen, 2017; Rebollo et al., 2012a).

Alternatively, it may also reveal different tempo of TE dynamics between these species. A recent peak of

activity of TEs can be seen in *D. melanogaster*, which is much smaller in *D. simulans (M*érel et al., 2020),

indicating that the colonization of the *D. simulans* genome by TEs started more recently (as suggested by our

previous results (Mohamed et al., 2020) and others (Kofler et al., 2015a)). Such ongoing colonization would

also lead to the selection of more efficient TE control mechanisms.

These contrasted impacts of TE insertions on genes through epigenetic marks across the species provide an

additional demonstration of the considerable natural variability due to TEs. We predict that this leads to

contrasted adaptive and evolutionary potentials, all the more sensible in a rapidly changing environment

(Baduel et al., 2021; Fablet and Vieira, 2011; Mérel et al., 2021).

A Defense

B Regulation

Figure 5. (A) The defense function of the piRNA pathway is prominent in *D. simulans*: TE epigenetic effects are stronger in this species (orange) compared to *D. melanogaster* (blue). (B) The regulation function of the piRNA pathway is prominent in *D. melanogaster*: Genome architecture is more tightly associated with TE insertions in *D. melanogaster,* as suggested by the stronger positive correlation between the numbers of TE insertions and gene H3K9me3 levels in this species.

Material and Methods

Drosophila strains

The strains under study in the present work were previously described in Mohamed *et al.* (Mohamed et al., 2020). The eight samples of *D. melanogaster* and *D. simulans* wild-type strains were collected using fruit baits in France (Gotheron, 44°56'0"N 04°53'30"E - "goth" strains) and Brazil (Saõ Jose do Rio Preto 20°41'04.3"S 49°21'26.1"W – "sj" strains) in June 2014. Two isofemale lines per species and geographical origin were established directly from gravid females from the field (French *D. melanogaster*: dmgoth63, dmgoth101; Brazilian *D. melanogaster*: dmsj23, dmsj7; French *D. simulans*: dsgoth613, dsgoth31; Brazilian *D. simulans*: dssj27, dssj9). Brothers and sisters were then mated for 30 generations to obtain inbred strains with very low intra-line genetic variability. Strains were kept at 24°C in standard laboratory conditions on cornmeal–sugar–yeast–agar medium.

Genome annotation

Genome assemblies were produced in (Mohamed et al., 2020) and have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB50024 (https://www.ebi.ac.uk/ena/browser/view/PRJEB50024). Throughout the present analysis, we kept scaffolds corresponding to complete chromosomes 2L, 2R, 3L, 3R, 4, and X.

TE annotation: We used RepeatMasker 4.1.0 (http://repeatmasker.org/) -species Drosophila in order to identify TE sequences in the assemblies, followed by OneCodeToFindThemAll (Bailly-Bechet et al., 2014) with default parameters, in order to parse RepeatMasker results. We include all TE sequences in the subsequent analyses, whether they are full length or truncated.

Gene annotation: We retrieved gtf files from FlyBase : ftp.flybase.net/genomes/Drosophila_melanogaster/dmel_r6,46_FB2022_03/gft/dmel-all-r6.46.gtf.gz and ftp.flybase.net/genomes/Drosophila_simulans/dsim_r2,02_FB2017_04/gtf/dsim-all-r2,02.gtf.gz. The corresponding fasta files were also downloaded from FlyBase: ftp.flybase.net/genomes/Drosophila_melanogaster/dmel_r6,46_FB2022_03/fasta/dmel-all-chromosome-r6.46.fasta.gz and ftp.flybase.net/genomes/Drosophila_simulans/dsim_r2,02_FB2017_04/fasta/dsim-all-chromosome-r2,02.fasta.gz. We used Liftoff (Shumate and Salzberg, 2020) to lift over gene annotations from the references to our genome assemblies. We used -flank 0.2 and only kept the "gene" and "exon" terms. Then, we used the GenomicRanges R package (version 1.38.0) (Lawrence et al., 2013) and the subsetByOverlaps function to cross gene and TE annotations.

1:1   orthologs: We   retrieved   ortholog   information   from   FlyBase
(ftp://ftp.flybase.net/releases/current/precomputed_files/orthologs/
dmel_orthologs_in_drosophila_species_fb_2022_01.tsv.gz) and kept only those genes for which there
was a 1 to 1 correspondence between *D. melanogaster* and *D. simulans*.

TE genomic sequence occupancy (bp) was computed using OneCodeToFindThemAll (Bailly-Bechet et al.,
2014).

In order to determine which TE insertions were common (shared) to the four strains of a species or unique
(private) to one strain, we performed all pairwise comparisons of TE gff using Liftoff -flank 0.2 (Shumate
and Salzberg, 2020). In the output, we filtered insertions with coverage >0.80 and sequence identity >0.80.
We ran *ad hoc* bash scripts to retrieve private and common insertions for each strain with the following
rationale: Private insertions to one strain are those that appear in the unmapped outputs of all pairwise
comparisons with the three other strains. Common insertions are those that are found in all pairwise
comparisons with the three other strains.

RNA-seq preparation

RNA was extracted from ovaries of 30 three to five day-old females. Two replicates per strain were
produced. RNA extraction was carried out using RNeasy Plus (Qiagen) kit following manufacturer's
instructions. After DNAse treatment (Ambion), quality control was performed using an Agilent Bioanalyzer.
Libraries were constructed from mRNA using the Illumina TruSeq RNA Sample Prep Kit following
manufacturer's recommendations. Libraries were sequenced on Illumina HiSeq 3000 with paired-end 150 nt
reads.

RNA-seq analysis

TE read counts were computed at the family level using the TEcount module of TEtools (Lerat et al., 2017)
and the list of TE sequences available at ftp://pbil.univ-lyon1.fr/pub/datasets/Roy2019/.

Genome sequences from *D. melanogaster* and *D. simulans* were downloaded from FlyBase (dmel-all-
chromosome-r6.16.fasta and dsim-all-chromosome-r2.02.fasta) and then masked using RepeatMasker
(http://repeatmasker.org/). For each species, we then built a multifasta file of gene sequences
using bedtools getfasta (Quinlan and Hall, 2010) with gff files available from FlyBase (dmel-all-r6.16.gff
and dsim-all-r2.02.gff).

Raw reads were processed using Trimmomatic 0.39 (Bolger et al., 2014) ILLUMINACLIP:TruSeq3-
PE.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:36, then mapped to genes
using HiSat2 (Kim et al., 2019). Alignment files were converted to BAM and sorted using SAMtools (Li et
al., 2009), and TPM and effective counts were then computed using eXpress (Roberts et al., 2011).

Quantification of the associations between TE insertions and gene transcript levels: considering only genes expressed in ovaries, we computed mean TPM across replicates and used the following linear models after log transformation: TPM ~ exon + intron + upstream + downstream, where "exon", "intron" "upstream", and "downstream" are the numbers of TE insertions in exons, introns, 5 kb upstream sequences, and 5 kb downstream sequences, respectively. Size effects for each of these factors were then recorded. To compute the contribution to total variance, we divided the Sum Square of the corresponding variables by the Total Sum Square, provided by the ANOVA of the linear model.

ChIP-seq preparation

Chromatin immunoprecipitation was performed using 50 ovary pairs dissected from three to five day old females. Ovaries were re-suspended in A1 buffer containing 60mM KCl, 15mM NaCl, 15mM Hepes, 0,5% Triton and 10mM Sodium butyrate. Formaldehyde (Sigma) was added to a final concentration of 1.8% for secondary cross-linking for 10 min at room temperature. Formaldehyde was quenched using glycine (0.125 M). Cross-linked cells were washed and pelleted twice with buffer A1, once with cell lysis buffer (140mM NaCl, 15mM Hepes, 1mM EDTA, 0.5mM EGTA, 1% Triton X100, 0.1% Sodium deoxycholate, 10mM sodium butyrate), followed by lysis in buffer containing ,140mM NaCl, 15mM Hepes, 1mM EDTA, 0.5mMEGTA, 1% Triton X100, 0.5% SDS, 0.5% N-Laurosylsarcosine, 0.1% sodium deoxycholate, 10mM sodium butyrate for 120 min at 4ºC. Lysates were sonicated in Bioruptor sonicator to reach a fragment size window of 200-600 bp.

Chromatin was incubated overnight at 4ºC with the following antibodies: for H3K9me3 ChIP using α-H3K9me3 (actif motif #39161, 3μg/IP) and for H3K4me3 using α-H3K4me3 (millipore #07-473, 3μg/IP) antibodies. The Magna ChIP™ A/G Chromatin Immunoprecipitation Kit (cat# 17-10085) was used following manufacturer's instructions. Final DNA recovery was perfomed by classic phenol/chloroform DNA precipitation method using MAxtract high density tubes to maximize DNA recovery.

DNA fragments were then sequenced on an Illumina HiSeq 4000 apparatus, with paired-end 100 nt reads. Due to technical issues, only one replicate could be used for dsgoth31 input.

ChIP-seq quality check: Validation of H3K4me3 enrichment around promoters and H3K9me3 on heterochromatic regions.

Raw reads were trimmed using trim_galore (https://zenodo.org/record/5127899#.YbnMs73MLDc) with default parameters along with --paired, --clip_R1 9, --clip_R2 9, and --max_n 0. Mapping was performed using Bowtie2 (Langmead and Salzberg, 2012) with --sensitive-local against the *D. melanogaster* r6.16 and *D. simulans* r2.02 genomes. Samtools was used to convert SAM to

coordinated sorted BAM files, while sambamba (Tarasov et al., 2015) was used to filter for uniquely mapping reads and to remove duplicates (sambamba view -h -t 2 -f bam -F "[XS] == null and not unmapped and not duplicate"). For *D. melanogaster* datasets, we filtered available blacklisted regions (Amemiya et al., 2019) with bedtools. Finally coverage files containing reads per genome coverage (RPGC) were obtained with DeepTools (Ramírez et al., 2016) bamCoverage with --extendReads, --effectiveGenomeSize 129789873 for *D. melanogaster* available from the Deeptools suite, and --effectiveGenomeSize 121102921 computed with unique-kmers.py from khmer (https://github.com/dib-lab/khmer). Promoter regions were obtained with gencode_regions (https://github.com/saketkc/gencode_regions) and along with coverage files were used in DeepTools computeMatrix and plotProfile to build the average coverage of H3K4me3 and H3K9me3 around transcription start sites in both species and on chromosomes for H3K9me3. The corresponding profiles looked as expected (Supplemental Fig. S7, S8).

ChIP-seq analysis

For each of the immunoprecipitated samples (H3K4me3, H3K9me3, input), TE read counts were computed at the family level using the TEcount module of TEtools (Lerat et al., 2017) and the list of TE sequences available at ftp://pbil.univ-lyon1.fr/pub/datasets/Roy2019/.

ChIP-seq counts were normalized across samples of the same species using the counts(normalize=T) function of DESeq2 1.26.0 (Love et al., 2014). This was done independently for each of the immunoprecipitated samples (H3K4me3, H3K9me3, input). We then performed a log-transformation using the rlogTransformation function of DESeq2, and subsequently considered mean values across replicates. We only kept genes expressed in ovaries. We chose to work on log-transformed values because log-transformation of count variables makes them fit normal assumption and thus makes them suitable for linear models. In addition, a ratio becomes a difference when log-transformed, which ensure the strict equivalence with the classical normalization approach consisting in dividing histone counts with input counts: log ([H3Kime3 counts] / [input counts]) = log(H3Kime3 counts) – log(input counts).

In order to quantify the associations between TE insertions and histone marks enrichment, we used the following linear models on log transformed read counts: histone mark (either H3K4me3 or H3K4me3) ~ input + exon + intron + upstream + downstream, where "exon", "intron", "upstream", and "downstream" are the numbers of TE insertions in exons, introns, 5 kb upstream sequences, and 5 kb downstream sequences, respectively. Size effects for each of these three factors were then recorded. To compute the contribution to total variance, we divided the Sum Square of the corresponding variables by the Total Sum Square, provided by the ANOVA of the linear model.

Small RNA extraction, sequencing, and analyses

Small RNA extraction, sequencing, and analyses dedicated to TEs had already been performed and described in Mohamed *et al.* (Mohamed et al., 2020). Sequence files had been deposited in NCBI SRA under the accession number PRJNA644327.

Gene-derived small RNAs: Sequencing adapters were removed using cutadapt (Martin, 2011), and 23-30 nt reads from one hand (considered as piRNAs) and 21 nt reads from the other hand (considered as siRNAs) were extracted using PRINSEQ lite (Schmieder and Edwards, 2011), as described in (Mohamed et al., 2020). Reads were then aligned on previously masked genomes (see above, RNA-seq section) using bowtie --best (Langmead et al., 2009). Aligned reads were counted using eXpress (Roberts et al., 2011) and "tot_counts" were considered.

Data access

The RNA-seq data generated in this study have been submitted to the NCBI BioProject database (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number PRJNA795668. The ChIP-seq data generated in this study have been submitted to the NCBI BioProject database under accession number PRJNA796157. TE and gene annotations have been deposited to Zenodo doi: 10.5281/zenodo.7189887.

Count tables for TE insertions, RNA-seq and ChIP-seq data are provided as Supplemental Material.

Acknowledgments

We thank Francois Sabot, Matthieu Boulesteix, and Vincent Mérel for useful discussions and technical help. We thank Gladys Mialdea, Justine Picarle, Sonia Janillon, and Nelly Burlet for technical help. This work was performed using the computing facilities of the CC LBBE/PRABI. Sequencing was performed by the GenomEast platform, a member of the "France Génomique" consortium (ANR-10-INBS-0009). This work was supported by Fondation pour la Recherche Médicale (grant DEP20131128536) and Agence Nationale de la Recherche (grant ExHyb ANR-14-CE19-0016-01). The authors declare no competing interests.

References

Akkouche A, Grentzinger T, Fablet M, Armenise C, Burlet N, Braman V, Chambeyron S, Vieira C. 2013. Maternally deposited germline piRNAs silence the tirant retrotransposon in somatic cells. *EMBO Rep* **14**:458–464. doi:10.1038/embor.2013.38

Akkouche A, Rebollo R, Burlet N, Esnault C, Martinez S, Viginier B, Terzian C, Vieira C, Fablet M. 2012. tirant, a newly discovered active endogenous retrovirus in Drosophila simulans. *J Virol* **86**:3675–3681. doi:10.1128/JVI.07146-11

Amemiya HM, Kundaje A, Boyle AP. 2019. The ENCODE Blacklist: Identification of Problematic Regions of the Genome. *Sci Rep* **9**. doi:10.1038/s41598-019-45839-z

Baduel P, Leduque B, Ignace A, Gy I, Gil JJ, Loudet O, Colot V, Quadrana L. 2021. Genetic and environmental modulation of transposition shapes the evolutionary potential of Arabidopsis thaliana. *Genome Biol* **22**. doi:10.1186/s13059-021-02348-5

Bailly-Bechet M, Haudry A, Lerat E. 2014. "One code to find them all": a perl tool to conveniently parse RepeatMasker output files. *Mob DNA* **5**:13. doi:10.1186/1759-8753-5-13

Biémont C, Vieira C. 2006. Genetics: junk DNA as an evolutionary force. *Nature* **443**:521–524. doi:10.1038/443521a

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinforma Oxf Engl* **30**:2114–2120. doi:10.1093/bioinformatics/btu170

Chakraborty M, Chang C-H, Khost DE, Vedanayagam J, Adrion JR, Liao Y, Montooth KL, Meiklejohn CD, Larracuente AM, Emerson JJ. 2021. Evolution of genome structure in the Drosophila simulans species complex. *Genome Res* **31**:380–396. doi:10.1101/gr.263442.120

Cridland JM, Thornton KR, Long AD. 2015. Gene expression variation in Drosophila melanogaster due to rare transposable element insertion alleles of large effect. *Genetics* **199**:85–93. doi:10.1534/genetics.114.170837

Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, Hannon GJ. 2018. piRNA-Guided Genome Defense: From Biogenesis to Silencing. *Annu Rev Genet* **52**:131–157. doi:10.1146/annurev-genet-120417-031441

Everett LJ, Huang W, Zhou S, Carbone MA, Lyman RF, Arya GH, Geisz MS, Ma J, Morgante F, St Armour G, Turlapati L, Anholt RRH, Mackay TFC. 2020. Gene expression networks in the Drosophila Genetic Reference Panel. *Genome Res* **30**:485–496. doi:10.1101/gr.257592.119

Fablet M, Akkouche A, Braman V, Vieira C. 2014. Variable expression levels detected in the Drosophila effectors of piRNA biogenesis. *Gene* **537**:149–153. doi:10.1016/j.gene.2013.11.095

Fablet M, Vieira C. 2011. Evolvability, epigenetics and transposable elements. *BioMol Concepts* **2**:333–341.

Hill MS, Vande Zande P, Wittkopp PJ. 2021. Molecular and evolutionary processes generating variation in gene expression. *Nat Rev Genet* **22**:203–215. doi:10.1038/s41576-020-00304-w

Hollister JD, Gaut BS. 2009. Epigenetic silencing of transposable elements: a trade-off between reduced transposition and deleterious effects on neighboring gene expression. *Genome Res* **19**:1419–1428. doi:10.1101/gr.091678.109

Horváth V, Merenciano M, González J. 2017. Revisiting the Relationship between Transposable Elements and the Eukaryotic Stress Response. *Trends Genet* **33**:832–841. doi:10.1016/j.tig.2017.08.007

Huang S, Tao X, Yuan S, Zhang Yuhang, Li P, Beilinson HA, Zhang Ya, Yu W, Pontarotti P, Escriva H, Le Petillon Y, Liu X, Chen S, Schatz DG, Xu A. 2016. Discovery of an Active RAG Transposon Illuminates the Origins of V(D)J Recombination. *Cell* **166**:102–114. doi:10.1016/j.cell.2016.05.032

Huang Y, Shukla H, Lee YCG. 2022. Species-specific chromatin landscape determines how transposable elements shape genome evolution. *eLife* **11**:e81567. doi:10.7554/eLife.81567

Isbel L, Grand RS, Schübeler D. 2022. Generating specificity in genome regulation through transcription factor sensitivity to chromatin. *Nat Rev Genet*. doi:10.1038/s41576-022-00512-6

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**:907–915. doi:10.1038/s41587-019-0201-4

Kofler R, Hill T, Nolte V, Betancourt AJ, Schlötterer C. 2015a. The recent invasion of natural Drosophila simulans populations by the P-element. *Proc Natl Acad Sci U S A* **112**:6659–6663. doi:10.1073/pnas.1500758112

Kofler R, Nolte V, Schlötterer C. 2015b. Tempo and Mode of Transposable Element Activity in Drosophila. *PLoS Genet* **11**:e1005406. doi:10.1371/journal.pgen.1005406

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:357–359. doi:10.1038/nmeth.1923

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10**:R25. doi:10.1186/gb-2009-10-3-r25

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ. 2013. Software for computing and annotating genomic ranges. *PLoS Comput Biol* **9**:e1003118. doi:10.1371/journal.pcbi.1003118

Le Thomas A, Rogers AK, Webster A, Marinov GK, Liao SE, Perkins EM, Hur JK, Aravin AA, Tóth KF. 2013. Piwi induces piRNA-guided transcriptional silencing and establishment of a repressive chromatin state. *Genes Dev* **27**:390–399. doi:10.1101/gad.209841.112

Lee YCG. 2015. The Role of piRNA-Mediated Epigenetic Silencing in the Population Dynamics of Transposable Elements in Drosophila melanogaster. *PLoS Genet* **11**:e1005269. doi:10.1371/journal.pgen.1005269

Lee YCG, Karpen GH. 2017. Pervasive epigenetic effects of Drosophila euchromatic transposable elements impact their evolution. *eLife* **6**. doi:10.7554/eLife.25762

Lerat E, Fablet M, Modolo L, Lopez-Maestre H, Vieira C. 2017. TEtools facilitates big data expression analysis of transposable elements and reveals an antagonism between their activity and that of piRNA genes. *Nucleic Acids Res* **45**:e17. doi:10.1093/nar/gkw953

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinforma Oxf Engl* **25**:2078–2079. doi:10.1093/bioinformatics/btp352

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**:550. doi:10.1186/s13059-014-0550-8

Malone CD, Brennecke J, Dus M, Stark A, McCombie WR, Sachidanandam R, Hannon GJ. 2009. Specialized piRNA pathways act in germline and somatic tissues of the Drosophila ovary. *Cell* **137**:522–535. doi:10.1016/j.cell.2009.03.040

Malone CD, Hannon GJ. 2009. Small RNAs as guardians of the genome. *Cell* **136**:656–668. doi:10.1016/j.cell.2009.01.045

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**:10–12. doi:10.14806/ej.17.1.200

Mérel V, Boulesteix M, Fablet M, Vieira C. 2020. Transposable elements in Drosophila. *Mob DNA* **11**:23. doi:10.1186/s13100-020-00213-z

Mérel V, Gibert P, Buch I, Rodriguez Rada V, Estoup A, Gautier M, Fablet M, Boulesteix M, Vieira C. 2021. The Worldwide Invasion of Drosophila suzukii Is Accompanied by a Large Increase of Transposable Element Load and a Small Number of Putatively Adaptive Insertions. *Mol Biol Evol* **38**. doi:10.1093/molbev/msab155

Mohamed M, Dang NT-M, Ogyama Y, Burlet N, Mugat B, Boulesteix M, Mérel V, Veber P, Salces-Ortiz J, Severac D, Pélisson A, Vieira C, Sabot F, Fablet M, Chambeyron S. 2020. A Transposon Story: From TE Content to TE Dynamic Invasion of Drosophila Genomes Using the Single-Molecule Sequencing Technology from Oxford Nanopore. *Cells* **9**. doi:10.3390/cells9081776

Mohn F, Sienski G, Handler D, Brennecke J. 2014. The rhino-deadlock-cutoff complex licenses noncanonical transcription of dual-strand piRNA clusters in Drosophila. *Cell* **157**:1364–1379. doi:10.1016/j.cell.2014.04.031

Moschetti R, Palazzo A, Lorusso P, Viggiano L, Marsano RM. 2020. "What You Need, Baby, I Got It": Transposable Elements as Suppliers of Cis-Operating Sequences in Drosophila. *Biology* **9**:E25. doi:10.3390/biology9020025

Naito K, Zhang F, Tsukiyama T, Saito H, Hancock CN, Richardson AO, Okumoto Y, Tanisaka T, Wessler SR. 2009. Unexpected consequences of a sudden and massive transposon amplification on rice gene expression. *Nature* **461**:1130–1134. doi:10.1038/nature08479

Obbard DJ, Gordon KHJ, Buck AH, Jiggins FM. 2009. The evolution of RNAi as a defence against viruses and transposable elements. *Philos Trans R Soc Lond B Biol Sci* **364**:99–115. doi:10.1098/rstb.2008.0168

Osada N, Miyagi R, Takahashi A. 2017. Cis- and Trans-regulatory Effects on Gene Expression in a Natural Population of Drosophila melanogaster. *Genetics* **206**:2139–2148. doi:10.1534/genetics.117.201459

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinforma Oxf Engl* **26**:841–842. doi:10.1093/bioinformatics/btq033

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**:W160–W165. doi:10.1093/nar/gkw257

Rebollo R, Horard B, Begeot F, Delattre M, Gilson E, Vieira C. 2012a. A snapshot of histone modifications within transposable elements in Drosophila wild type strains. *PloS One* **7**:e44253. doi:10.1371/journal.pone.0044253

Rebollo R, Karimi MM, Bilenky M, Gagnier L, Miceli-Royer K, Zhang Y, Goyal P, Keane TM, Jones S, Hirst M, Lorincz MC, Mager DL. 2011. Retrotransposon-induced heterochromatin spreading in the mouse revealed by insertional polymorphisms. *PLoS Genet* **7**:e1002301. doi:10.1371/journal.pgen.1002301

Rebollo R, Romanish MT, Mager DL. 2012b. Transposable elements: an abundant and natural source of regulatory sequences for host genes. *Annu Rev Genet* **46**:21–42. doi:10.1146/annurev-genet-110711-155621

Roberts A, Trapnell C, Donaghey J, Rinn JL, Pachter L. 2011. Improving RNA-Seq expression estimates by correcting for fragment bias. *Genome Biol* **12**:R22. doi:10.1186/gb-2011-12-3-r22

Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinforma Oxf Engl* **27**:863–864. doi:10.1093/bioinformatics/btr026

Senti K-A, Brennecke J. 2010. The piRNA pathway: a fly's perspective on the guardian of the genome. *Trends Genet TIG* **26**:499–509. doi:10.1016/j.tig.2010.08.007

Shpiz S, Ryazansky S, Olovnikov I, Abramov Y, Kalmykova A. 2014. Euchromatic transposon insertions trigger production of novel Pi- and endo-siRNAs at the target sites in the drosophila germline. *PLoS Genet* **10**:e1004138. doi:10.1371/journal.pgen.1004138

Shumate A, Salzberg SL. 2020. Liftoff: accurate mapping of gene annotations. *Bioinforma Oxf Engl* btaa1016. doi:10.1093/bioinformatics/btaa1016

Sienski G, Dönertas D, Brennecke J. 2012. Transcriptional silencing of transposons by Piwi and maelstrom and its impact on chromatin state and gene expression. *Cell* **151**:964–980. doi:10.1016/j.cell.2012.10.040

Sultana T, Zamborlini A, Cristofari G, Lesage P. 2017. Integration site selection by retroviruses and transposable elements in eukaryotes. *Nat Rev Genet* **18**:292–308. doi:10.1038/nrg.2017.7

Sundaram V, Cheng Y, Ma Z, Li D, Xing X, Edge P, Snyder MP, Wang T. 2014. Widespread contribution of transposable elements to the innovation of gene regulatory networks. *Genome Res* **24**:1963–1976. doi:10.1101/gr.168872.113

Tarasov A, Villeva AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast processing of NGS alignment formats. *Bioinforma Oxf Engl* **31**. doi:10.1093/bioinformatics/btv098

Thomas J, Vadnagara K, Pritham EJ. 2014. DINE-1, the highest copy number repeats in Drosophila melanogaster are non-autonomous endonuclease-encoding rolling-circle transposable elements (Helentrons). *Mob DNA* **5**:18. doi:10.1186/1759-8753-5-18

Torri A, Jaeger J, Pradeu T, Saleh M-C. 2022. The origin of RNA interference: Adaptive or neutral evolution? *PLoS Biol* **20**:e3001715. doi:10.1371/journal.pbio.3001715

Ullastres A, Merenciano M, González J. 2021. Regulatory regions in natural transposable element insertions drive interindividual differences in response to immune challenges in Drosophila. *Genome Biol* **22**:265. doi:10.1186/s13059-021-02471-3

Uzunović J, Josephs EB, Stinchcombe JR, Wright SI. 2019. Transposable Elements Are Important Contributors to Standing Variation in Gene Expression in Capsella Grandiflora. *Mol Biol Evol* **36**:1734–1745. doi:10.1093/molbev/msz098

Vieira C, Fablet M, Lerat E, Boulesteix M, Rebollo R, Burlet N, Akkouche A, Hubert B, Mortada H, Biémont C. 2012. A comparative analysis of the amounts and dynamics of transposable elements in natural populations of Drosophila melanogaster and Drosophila simulans. *J Environ Radioact* **113**:83–86. doi:10.1016/j.jenvrad.2012.04.001

Vieira C, Lepetit D, Dumont S, Biémont C. 1999. Wake up of transposable elements following Drosophila simulans worldwide colonization. *Mol Biol Evol* **16**:1251–1255.

Villanueva-Cañas JL, Horvath V, Aguilera L, González J. 2019. Diverse families of transposable elements affect the transcriptional regulation of stress-response genes in Drosophila melanogaster. *Nucleic Acids Res* **47**:6842–6857. doi:10.1093/nar/gkz490

Wei KH-C, Mai D, Chatla K, Bachtrog D. 2022. Dynamics and Impacts of Transposable Element Proliferation in the Drosophila nasuta Species Group Radiation. *Mol Biol Evol* **39**:msac080. doi:10.1093/molbev/msac080

Wells JN, Feschotte C. 2020. A Field Guide to Eukaryotic Transposable Elements. *Annu Rev Genet*. doi:10.1146/annurev-genet-040620-022145

Wittkopp PJ, Kalay G. 2011. Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nat Rev Genet* **13**:59–69. doi:10.1038/nrg3095

Zhang X, Zhao M, McCarty DR, Lisch D. 2020. Transposable elements employ distinct integration strategies with respect to transcriptional landscapes in eukaryotic genomes. *Nucleic Acids Res* **48**:6685–6698. doi:10.1093/nar/gkaa370